g 3, out 1.22
ORNL 822

UNCLASSIFIED

APPLIED FISHERIES

ORNL 822

Health & Biology

SEP 22 1950
C

ORNL-822

ENGINEERING LIBRARY
MAY 24 1967
UNIVERSITY OF WASH.

THE EFFECT OF X IRRADIATION IN
OXYGEN AND IN HYDROGEN AT NORMAL
AND POSITIVE PRESSURES ON
CHROMOSOME ABERRATION FREQUENCY IN
TRADESCANTIA MICROSPORES

OAK RIDGE NATIONAL LABORATORY
OPERATED BY
CARBIDE AND CARBON CHEMICALS DIVISION
UNION CARBIDE AND CARBON CORPORATION

POST OFFICE BOX P
OAK RIDGE, TENNESSEE

metadc100487

LIBRARY
UNIVERSITY OF WASH
NOV 2 1967

UNCLASSIFIED

UNCLASSIFIED

Report Number: CRNL- 822 _____
Health and Biology
This document consists of 15 _____
pages.
Copy 161 of 167 Series A _____
ISSUED _____

Contract No. W-7405-Eng-26

BIOLOGY DIVISION

THE EFFECT OF X IRRADIATION IN OXYGEN AND IN HYDROGEN AT NORMAL
AND POSITIVE PRESSURES ON CHROMOSOME ABERRATION FREQUENCY IN
TRADESCANTIA MICROSPORES

Norman H. Giles, Jr. and Alvin V. Beatty

DATE ISSUED
SEP 15 1950

OAK RIDGE NATIONAL LABORATORY
operated by
CARBIDE AND CARBON CHEMICALS DIVISION
Union Carbide and Carbon Corporation
Post Office Box P
Oak Ridge, Tennessee

UNCLASSIFIED

ORNL 822
Health and Biology

Internal Distribution:

1. G. T. Felbeck (C&CCC)	16. W. B. Humes (K-25)	26. F. L. Steahly
2-3. 706-A Library	17. W. D. Lavers (Y-12)	27. M. T. Kelley
4. 706-B Library	18. A. M. Weinberg	28. D. W. Cardwell
5. Biology Library	19. E. J. Murphy	29. J. H. Gillette
6. Health Physics Library	20. F. C. VonderLage	30. A. S. Householder
7. Metallurgy Library	21. A. Hollaender	31. N. H. Giles, Jr.
8-9. Training School Library	22. J. S. Felton	32. A. V. Beatty
10-13. Central Files	23. J. A. Swartout	33. M. J. Skinner
14. C. E. Center	24. A. H. Snell	34. Central Files (O.P.)
15. C. E. Larson	25. K. Z. Morgan	
-
- 35-42. Argonne National Laboratory
 - 43. Armed Forces Special Weapons Project
 - 44-45. Atomic Energy Commission, Washington
 - 46. Battelle Memorial Institute
 - 47. Brush Beryllium Company
 - 48-55. Brookhaven National Laboratory
 - 56. Bureau of Medicine and Surgery
 - 57. Bureau of Ships
 - 58-61. Carbide and Carbon Chemicals Division (K-25)
 - 62-65. Carbide and Carbon Chemicals Division (Y-12)
 - 66. Chicago Operations Office
 - 67. Cleveland Area Office, AEC
 - 68. Columbia University (J. R. Dunning)
 - 69. Columbia University (G. Failla)
 - 70. Dow Chemical Company
 - 71. H. K. Ferguson Company
 - 72-74. General Electric, Richland
 - 75. Harshaw Chemical Corporation
 - 76. Idaho Operations Office
 - 77-78. Iowa State College
 - 79. Kansas City Operations Branch
 - 80-81. Kellogg Corporation
 - 82-85. Knolls Atomic Power Laboratory
 - 86-88. Los Alamos Scientific Laboratory
 - 89. Mallinckrodt Chemical Works
 - 90. Massachusetts Institute of Technology (A. Gaudin)
 - 91. Massachusetts Institute of Technology (A. Kaufmann)
 - 92-94. Mound Laboratory
 - 95-96. National Advisory Committee for Aeronautics
 - 97-98. National Bureau of Standards
 - 99-100. Naval Radiological Defense Laboratory
 - 101. New Brunswick Laboratory
 - 102-106. New York Operations Office
 - 107. North American Aviation, Inc.
 - 108. Patent Branch (Washington)
 - 109. RAND Corporation
 - 110. Sandia Laboratory

UNCLASSIFIED

External Distribution:

- 111. Santa Fe Operations Office
- 112. Sylvania Electric Products, Inc.
- 113-127. Technical Information Division (Oak Ridge)
- 128. USAF, Air Surgeon (Lt. Col. R. H. Blount)
- 129. USAF, Director of Armament (Captain C. I. Browne)
- 130-131. USAF, Director of Research and Development (Col. R. J. Mason, F. W. Bruner)
- 132. USAF, Eglin Air Force Base (Major A. C. Field)
- 133. USAF, Kirtland Air Force Base (Col. M. F. Cooper)
- 134. USAF, Maxwell Air Force Base (Col. F. N. Moyers)
- 135-136. USAF, NEPA Office
- 137-138. USAF, Office of Atomic Energy (Col. H. C. Donnelly, A. A. Fickel)
- 139. USAF, Offutt Air Force Base (Col. H. R. Sullivan, Jr.)
- 140. USAF, Wright-Patterson Air Force Base (Rodney Hudenberg)
- 141. U. S. Army, Atomic Energy Branch (Lt. Col. A. W. Betts)
- 142. U. S. Army, Army Field Forces (Captain James Kerr)
- 143. U. S. Army, Commanding General, Chemical Corps Technical Command (Col. J. A. MacLaughlin through Mrs. G. S. Benjamin)
- 144. U. S. Army, Chief of Ordnance (Lt. Col. A. R. Del Campo)
- 145. U. S. Army, Commanding Officer, Watertown Arsenal (Col. C. H. Deitrick)
- 146. U. S. Army, Director of Operations Research (Dr. Ellis Johnson)
- 147. U. S. Army, Office of Engineers (Allen O'Leary)
- 148. U. S. Army, Office of the Chief Signal Officer (C. T. Clayton through Maj. G. C. Hunt)
- 149. U. S. Army, Office of the Surgeon General (Col. W. S. Stone)
- 150. U. S. Geological Survey (T. N. Nolan)
- 151. USAF, Director of Plans and Operations (Col. R. L. Applegate)
- 152. U. S. Public Health Service
- 153. University of California at Los Angeles
- 154-158. University of California Radiation Laboratory
- 159-160. University of Rochester
- 161. University of Washington
- 162-163. Western Reserve University
- 164-167. Westinghouse Electric Company

3

THE EFFECT OF X IRRADIATION IN OXYGEN AND IN HYDROGEN AT NORMAL
AND POSITIVE PRESSURES ON CHROMOSOME ABERRATION FREQUENCY IN
*TRADESCANTIA MICROSPORES**

Norman H. Giles, Jr. and Alvin V. Beatty
Biology Division
Oak Ridge National Laboratory

Previous investigations by Giles and Riley (1, 2) have demonstrated a marked effect of oxygen in increasing the frequency of X-ray induced chromosomal aberrations in microspores of *Tradescantia paludosa*. In certain experiments (2), the relation between aberration frequency and percentage of oxygen present during irradiation was studied. Although there was an almost linear increase in aberration frequency between 2 and 21% oxygen (air), no significant difference was noted between 0% oxygen (pure helium) and 2% oxygen (plus 98% helium). Additional experiments have now been performed to reinvestigate the effect of low percentages of oxygen. Further, studies have been made of the effect on aberration frequency of irradiation in various percentages of oxygen under pressure. Preliminary experiments have also been carried out on the effects of irradiation in hydrogen at normal and positive pressure. The general methods utilized for X irradiation of inflorescences and subsequent cytological analysis have been described previously (2).

In the experiments designed to reinvestigate the effects of low percentages of oxygen, special attempts were made to remove as much oxygen as possible from the inflorescences before irradiation. The following procedure was utilized:

- (1) The inflorescences, after being placed in the gas-tight exposure chamber, were subjected to prolonged evacuation.
- (2) Helium which had been especially freed of any residual oxygen was then introduced into the evacuated exposure chamber. To remove any oxygen the helium was passed slowly through a coil in liquid nitrogen

and then over copper foil heated to 500° C before entering the exposure chamber.

(3) Finally, the inflorescences were kept in the dark for an hour before irradiation to facilitate the removal by respiration of any further oxygen. In addition to the exposures in helium, irradiations were performed in atmospheres containing 2, 5, and 10% oxygen (plus helium). The percentages of oxygen in these gas mixtures were accurate (on the basis of analyses furnished by the vendor, the Ohio Chemical and Surgical Equipment Co.) to $\pm 0.2\%$. All experiments were carried out at room temperature (approximately 25° C). To increase the statistical reliability of the determinations, a larger number of cells was scored than previously. The data are presented in Table 1. They are also plotted in Fig. 1, together with the averages of points at higher percentages of oxygen which were obtained in previous experiments (2). These results indicate that there is still a substantial yield of aberrations even in the complete (or nearly complete) absence of oxygen. When oxygen is present during irradiation, there is a rapid rise in aberration frequency above this base level. This increase is linear between 0 and 10% oxygen, after which the rise is apparently more gradual, and shows a definite leveling off at around 20% oxygen.

The irradiation experiments in atmospheres of increasing oxygen concentration just outlined were all performed at normal atmospheric pressure (ca 740 mm of Hg at Oak Ridge). Since the amount of dissolved oxygen is directly proportional to pressure, an experiment was designed to test for an oxygen effect in another way -- by carrying out exposures in which different sets of inflorescences were irradiated in the same oxygen-helium gas mixtures, but at pressures of 1, 2 and 3 atmospheres above normal pressure. The same exposure chamber was used as previously, and the pressure inside the chamber was measured by means of a standard gauge on the gas cylinder, the accuracy of which was checked against a mercury manometer over the

Table 1

The Effect of Irradiation in Various Percentages of Oxygen on the
Frequency of Chromosomal Aberrations in Tradescantia Microspores.

All X-Ray Exposures of 400 r at 50 r/Min.

Oxygen percentage	No. cells	No. interchanges	Interchanges per cell	Interstitial deletions	I.D. per cell
0	1000	234	0.234 ± 0.015	287	0.287 ± 0.017
2	1200	347	0.289 ± 0.016	394	0.328 ± 0.017
5	1000	427	0.427 ± 0.021	501	0.501 ± 0.022
10	600	364	0.607 ± 0.032	453	0.755 ± 0.035

UNCLASSIFIED

DRAWING #9556

Fig. 1

range of the first 2 atmospheres above normal. The data for exposures in 5 and 10% oxygen (plus helium) are presented in Table 2, together with results obtained in air and in pure (ca 99.8%) helium. The values for two separate experiments in 5% oxygen at normal pressure have been averaged in the table. During the exposure in compressed air at 3 atmospheres above normal a small leak developed in the exposure chamber, and although the pressure gauge did not drop, it is felt that the value for this exposure may not be as reliable as for the other observations.

These results show that there is a marked increase in aberration frequency accompanying increasing pressure in those experiments in which oxygen was present during irradiation. When exposures occurred in the absence of oxygen (in pure helium) however, no increase resulted, indicating that pressure alone, at least within this range, did not influence the yield of aberrations. In Fig. 2 the frequencies of chromosomal interchanges at various pressures in 5 and 10% oxygen have been plotted against the standard curve from Fig. 1, on the assumption that the amount of dissolved oxygen in the microspores is directly proportional to the pressure, and that this value determines the aberration yield at the standard dose of 400 r at 50 r/min. As can be seen from the graph, the agreement with previous results at comparable partial pressures of oxygen is very good. These results thus support earlier views that the amount of dissolved oxygen present in cells is an important factor in determining aberration frequency. They indicate that pressure alone, at least within the range up to 3 atmospheres above normal, does not influence this type of radiation-induced change.

Several different types of evidence (3) support the view of Thoday and Read (4) that the effect of oxygen in increasing chromosome aberration frequency may result from an indirect action of X radiation to decompose water with the production of hydrogen peroxide. However, although very little hydrogen peroxide appears to

Table 2

The Effect of Pressure on the Frequency of X-Ray Induced Chromosomal Aberrations in Tradescantia Microspores Irradiated in Atmospheres Containing Various percentages of Oxygen. All Exposures of 400 r at 50 r/Min.

Pressure in exposure chamber	No. cells	No. interchanges	Interchanges per cell	No. interstitial deletions	Interstitial deletions per cell
Irradiated in 10% oxygen (plus 90% helium)					
Normal atmospheric (ca 740 mm Hg)	800	474	0.59 ± 0.027	599	0.75 ± 0.031
One atmosphere above normal	550	436	0.79 ± 0.038	563	1.02 ± 0.043
Two atmospheres above normal	700	584	0.83 ± 0.035	875	1.25 ± 0.042
Three atmospheres above normal	700	584	0.83 ± 0.035	732	1.05 ± 0.039
Irradiated in pure helium (~99.8%)					
Normal atmospheric (ca 740 mm Hg)	950	229	0.24 ± 0.016	256	0.27 ± 0.017
Three atmospheres above normal	1000	223	0.22 ± 0.015	259	0.26 ± 0.016
Irradiated in air					
Normal atmospheric (ca 740 mm Hg)	1000	713	0.71 ± 0.027	780	0.78 ± 0.028
Three atmospheres above normal	950	786	0.83 ± 0.030	994	1.05 ± 0.033
Irradiated in 5% oxygen (plus 95% helium)					
Normal atmospheric (ca 740 mm Hg)	1850	903	0.49 ± 0.016	954	0.52 ± 0.017
One atmosphere above normal	950	583	0.61 ± 0.025	671	0.71 ± 0.027
Two atmospheres above normal	1000	683	0.68 ± 0.026	802	0.80 ± 0.028
Three atmospheres above normal	840	657	0.78 ± 0.031	783	0.93 ± 0.033

UNCLASSIFIED

DRAWING #9557

400r AT 50r/MIN.

○ = 5% O₂ (SUPERSCRIPTS) PRESSURES ABOVE NORMAL▲ = 10% O₂ (SUBSCRIPTS) PRESSURES ABOVE NORMAL

Fig. 2

be produced by X rays in oxygen-free water (5, 6), it will be recalled (Table 1) that there is still a substantial aberration frequency in inflorescences exposed in purified helium to 400 r. The question naturally arises as to the mechanism by which these aberrations are produced. It seems quite probable (3) that the effect of a substance such as hydrogen peroxide on aberration frequency would arise from an increased production of chromosome breaks rather than from an influence on the reunion behavior of broken ends. If this is the case, aberrations induced by X rays in the absence of oxygen might result (at least in part) from breaks produced by another substance, the OH radical, which is formed by the radiodecomposition of oxygen-free water. On the other hand, all the aberrations produced in the absence of oxygen might arise from the direct action of the radiation in ionizing the molecules of the chromosomes themselves. A preliminary attempt has been made to distinguish between these two possibilities by means of experiments designed to remove the OH radical by promoting, during X irradiation, the back reaction to form water. Allen (6) has shown that the presence of molecular hydrogen markedly accelerates this back reaction. Consequently, inflorescences were irradiated in pure hydrogen at normal atmospheric pressure and at 3 atmospheres above normal. These data are presented in Table 3. Although there is some decrease in aberration frequency compared to earlier results obtained with irradiation in helium, the differences for the most reliable observations, those of interchanges, are not significant. In this experiment it was not possible to obtain evidence that hydrogen was actually present in the cells during irradiation. However, previous experiments with oxygen (2) have indicated that under similar conditions this gas diffuses into microspores very rapidly. Thus, despite the fact that hydrogen is less soluble in water than oxygen, it seemed reasonable to conclude that a considerable amount of hydrogen would be present in the microspores during

Table 3

The Effect of Irradiation in Hydrogen on Chromosome Aberration Frequency
in Tradescantia Microspores. 400 r at 50 r/Min.

Pressure in exposure chamber	No. cells	No. interchanges	Interchanges per cell	Interstitial deletions	I.D. per cell
Normal atmospheric (<u>ca</u> 740 mm Hg)	707	149	0.21 ± 0.017	166	0.23 ± 0.018
Three atmospheres above normal	498	100	0.20 ± 0.020	124	0.25 ± 0.022

irradiation, and that this should react to remove the OH radical. Consequently the failure to find a significant decrease in aberration frequency when X radiation is performed in hydrogen may mean that all the residual aberrations induced by X rays in the absence of oxygen arise as a result of a direct effect of the radiation on the chromosomes. It should be pointed out, however, that this conclusion is based on the assumption that reactions leading to H_2O_2 production or suppression in cells from which oxygen has been removed as completely as possible are similar to those occurring in oxygen-free pure water. There is as yet little experimental evidence on this point, and it is quite possible that the complexity of the cellular environment may cause very different reactions to occur.

Summary. Further studies have been made on the effect of oxygen concentrations in the range between 0 and 10% (at normal atmospheric pressure) on the frequency of X-ray-induced chromosomal aberrations in Tradescantia microspores. There is still a substantial aberration yield even in the complete (or nearly complete) absence of oxygen. When oxygen is present in irradiated cells, however, there is a rapid, linear increase in aberration frequency between 0 and 10% oxygen, after which the rise is more gradual.

Experiments have also been performed to test the effect of pressure during irradiation. A marked increase in aberration frequency accompanying increasing pressure occurred in those experiments in which oxygen was present during irradiation. Exposures in 5 and 10% oxygen at different pressures up to 3 atmospheres above normal produced aberration frequencies similar to those obtained in comparable partial pressures of oxygen. These and previous results indicate that oxygen increases the frequency of X-ray-induced chromosome aberrations and that the amount of oxygen present during the time of exposure determines the extent of the effect. Since no pressure effect was obtained in these experiments in the absence of oxygen,

the apparent effect of pressure in the presence of oxygen is, in reality, an oxygen effect.

Experiments in which inflorescences were irradiated in pure hydrogen at normal atmospheric pressure and at 3 atmospheres above normal pressure showed no significant decrease in interchange frequency. These results are interpreted as supporting the view that aberrations produced by X rays in the absence of oxygen may all result from a direct effect of the radiation on the chromosomes.

References

1. Giles, M. H., Jr., and Riley, H. P. Proc. Nat. Acad. Sci., 1949, 35, 640.
2. Giles, M. H., Jr., and Riley, H. P. Proc. Nat. Acad. Sci., 1950, 36, 337.
3. Giles, M. H., Jr. Paper presented at Oberlin Symposium on Radiobiology, June, 1950.
4. Thoday, J. M. and Read, J. Nature, 1949, 163, 133.
5. Bonet-Maury, P. and Lefort, M. Nature, 1948, 162, 381.
6. Allen, A. O. J. Phy. and Cell. Chem., 1948, 52, 479.

UNCLASSIFIED

UNCLASSIFIED