SEARCH FOR SHORT LIVED PARTICLES PRODUCED BY 300 GeV
PROTONS IN EMULSIONS

E. Albini, W. Bozzoli, P. Capiluppi, A. Forino,
R. Gessaroli, G. Giacomelli, P. Lugaresi-Serra,
A. Quarenzi and F. Rimondi.

Istituto di Fisica, Università di Bologna, Bologna, Italy
Istituto Nazionale di Fisica Nucleare, Sezione di Bologna
Scuola di Perfezionamento in Fisica, Università di Bologna

S. Di Caporiacco, A.M. Cartacci and M. G. Dagliana

Istituto di Fisica, Università di Firenze, Firenze, Italy
Istituto Nazionale di Fisica Nucleare, Sezione di Firenze

It is proposed to continue the search for short-lived particles with lifetimes in the range of $10^{-12} - 10^{-14}$ sec., using 400 GeV protons in emulsion nuclei. As in the experiment No.419, the signal would be a neutral or charged decay close to the primary proton-nucleus interaction.

We request the exposure of one stack of Ilford G5 emulsions, of 80 plates of $5 \times 10 \times 0.06 \text{ cm}^3$, to 400 GeV protons with a density of $\approx 3 \times 10^4$ protons/cm², with a total exposure of 1.2×10^6 incident protons over an area of $5 \times 8 \text{ cm}^2$.

The experiment should be sensitive to the production of new particles if they have a mass of 1 - 10 GeV and if their production cross section is larger than 1 μb .

More details can be found in the similar experiment at 300 GeV (No.419).