Princeton Plasma Physics Laboratory

PPPL-

PPPL-

Prepared for the U.S. Department of Energy under Contract DE-AC02-09CH11466.

Princeton Plasma Physics Laboratory

Report Disclaimers

Full Legal Disclaimer

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Trademark Disclaimer

Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors.

PPPL Report Availability

Princeton Plasma Physics Laboratory:

<http://www.pppl.gov/techreports.cfm>

Office of Scientific and Technical Information (OSTI):

<http://www.osti.gov/bridge>

Related Links:

[U.S. Department of Energy](#)

[Office of Scientific and Technical Information](#)

[Fusion Links](#)

Effects of toroidal rotation shear on Toroidicity-induced Alfvén Eigenmodes in the National Spherical Torus Experiment

M. Podestà, R. E. Bell, E. D. Fredrickson, N. N. Gorelenkov, B. P. LeBlanc

Princeton Plasma Physics Laboratory, Princeton NJ 08543 - USA

W. W. Heidbrink

Department of Physics and Astronomy,

University of California Irvine, CA 92697, USA

N. A. Crocker, S. Kubota

University of California Los Angeles, CA 90095, USA

H. Yuh

Nova Photonics, Princeton, NJ 08543, USA

(Dated: August 16, 2010)

Abstract

The effects of a sheared toroidal rotation on the dynamics of bursting Toroidicity-induced Alfvén eigenmodes are investigated in neutral beam heated plasmas on the National Spherical Torus Experiment (NSTX) [M. Ono *et al.*, Nucl. Fusion **40** 557 (2000)]. The modes have a global character, extending over most of the minor radius. A toroidal rotation shear layer is measured at the location of maximum drive for the modes. Contrary to results from other devices, no clear evidence of increased damping is found. Instead, experiments with simultaneous neutral beam and radio-frequency auxiliary heating show a strong correlation between the dynamics of the modes and the instability drive. It is argued that kinetic effects involving changes in the mode drive and damping mechanisms other than rotation shear, such as continuum damping, are mostly responsible for the bursting dynamics of the modes.

PACS numbers: 52.55.Fa, 52.35.Bj, 52.35.Mw

I. INTRODUCTION

Waves in the Alfvén frequency range can interact with a fast-ion population, leading to loss or redistribution of fast particles [1][2]. In particular, the nonlinear evolution following saturation of multiple modes may lead to their strong coupling and, eventually, to the overlap of mode resonances in phase space [3][4]. As a result, extended regions of the fast ion distribution function can provide free energy to further sustain the growth of the modes. These phase-space (or *kinetic*) effects may be responsible for the variety of phenomena observed in experiments on toroidicity-induced Alfvén eigenmodes (TAEs). A single mode can split into multiple frequencies [5], or show a series of frequency *chirps*. The latter can result in a rapid increase as well as a decrease of the observed mode frequency [6]. The *stochastization* of fast ion phase space associated with chirping modes is thought to be responsible for the enhanced losses and redistribution of fast ions observed experimentally [7][8][9]. Clear signatures of non-linear TAE behavior and induced fast ion losses are commonly observed in neutral beam (NB) heated plasmas on the National Spherical Torus Experiment (NSTX) [10], where NB injection provides the main source of additional heating and momentum.

Understanding the mechanisms leading to strongly nonlinear mode dynamics, including mode-mode coupling, is of broad interest for the design of future fusion devices. Indeed, it represents the first step for designing specific tools to reduce, or even suppress, the deleterious loss of fast ions associated with bursting TAE activity. Experimental results from large aspect ratio tokamaks suggest that a sheared toroidal flow may represent a possible suppression mechanism. Sufficiently strong gradients in the toroidal rotation would cause a radial decorrelation of the TAE mode structure, thus reducing the saturation level and clamping the growth rate. On JET, an increase in the damping rate of global TAE modes with toroidal mode number $n = 1$ was observed for increasing values of the rotation shear [11]. Similar results have been reported from JT60-U for intermediate- n modes [12]. In this case, both NB and RF injection were used to affect the rotation profile and the fast ion drive of the modes. According to theory, an analog decorrelation mechanism could be provided by sheared *poloidal* flows, such as the self-generated $\mathbf{E} \times \mathbf{B}$ flow induced by the Reynold stress associated with TAE turbulence [13]. The study of the effects of poloidal rotation and turbulence on TAEs is beyond the scope of this paper, and is not considered in the following.

FIG. 1. (a) Spectrogram showing TAE activity in the range $f = 70 - 200$ kHz for discharge #135388. (b) Temporal evolution of the central electron density and temperature. (c) Waveforms of NB power and neutron rate. Note the drop of the neutron rate at $t \approx 360$ ms, corresponding to an *avalanche*, which indicates a loss of fast ions from the core plasma.

In this paper, the effects of a sheared toroidal rotation profile on the dynamics of bursting TAEs are investigated in NB-heated plasmas of NSTX. The mode dynamics includes the evolution from a quasi-stationary phase into a regime characterized by dramatic frequency and amplitude bursts - so called *avalanches* [8][14]. The primary interest for the present study is an intermediate phase, prodromic to the avalanching regime causing fast ion losses, during which relatively small amplitude and frequency changes compared to avalanches are observed. If the hypothesis that a sheared flow can directly affect the TAE dynamics is confirmed, new avenues may open up for controlling the TAE dynamics during this early phase, thus limiting the detrimental effect of avalanches on fast ion confinement. Testing this hypothesis is the primary goal of this work.

The paper is organized as follows. The NSTX plasmas studied in this work and the general observations on TAE dynamics are described in Sec. II. Section III presents the measurements of toroidal rotation and its relationship with other quantities, such as the fast ion profile. The effects of rotation shear are discussed in Sec. IV. The lack of clear experimental evidence of mode suppression by rotation shear motivates the extension of the study to the TAE drive (Sec. V). Section VI summarizes the main results and concludes the paper.

FIG. 2. Displacement associated with a $n = 3$ mode, measured by a multi-channel reflectometer at three different times.

II. EXPERIMENTAL SCENARIO

NSTX is a small aspect ratio device, with major and minor radius $R_0 = 0.9$ m and $a = 0.65$ m (aspect ratio $R_0/a \approx 1.3$). For the experiments discussed herein, the magnetic field is 5.5 kG. The plasma current is 700 – 900 kA, for a pulse duration of ~ 0.6 s. L-mode deuterium plasmas are mainly investigated. Density and temperature are $n_e \approx 3 \times 10^{19} \text{ m}^{-3}$ and $T_i \approx T_e \leq 1.5$ keV (subscripts e, i refer to electrons and ions, respectively). Electron and ion plasma parameters, including rotation, are measured by a Thomson scattering [15] and a charge-exchange recombination spectroscopy system [16]. The difference between the measured carbon velocity and the main ion velocity [17], which determines the Doppler shift of the TAE frequencies, has been estimated from other studies to be $< 5\%$ [18] and is negligible for the purposes of this study. The total plasma β (ratio of plasma to magnetic energy) is $\lesssim 15\%$. Up to 3 MW of NB power is injected from three sources with tangency radii of 69, 59 and 49 cm. The maximum acceleration voltage is 90 kV. The injected species is deuterium. The velocity of fast ions resulting from charge-exchange of the injected neutrals is up to five times the Alfvén velocity. Several resonances with the fast ion population can thus destabilize waves in the Alfvén frequency range, including TAEs, as commonly observed [7]. In addition to neutral beams, a High-Harmonic Fast Wave (HHFW) system [19] is used to heat the plasma with $P_{RF} = 2$ MW.

Figure 1 illustrates a spectrogram from Mirnov coils measuring magnetic field fluctuations close to the low-field side vacuum vessel wall. Multiple modes are visible in the frequency

range 70 – 200 kHz. The modes are first destabilized during the current ramp-up ($t \leq 200$ ms). In this initial stage they are identified as reverse-shear Alfvén eigenmodes (RSAEs) [20]. After the beginning of the current flat-top RSAEs gradually evolve into TAEs, driven unstable by the injected beam ions [9]. No direct measurement of the safety factor profile, $q(R, t)$, is available for this discharge in the time window $t = 240 - 400$ ms. A reconstruction of the average $q(R, t)$ has been performed through the Grad-Shafranov equilibrium reconstruction code LRDFIT [21] by varying the NB timing on a shot-to-shot basis. The analysis shows that the discharges under investigation have a reversed-shear q profile with $q_{min} \sim 2$ at $t \sim 300$ ms. Toroidal mode numbers are $n = 2 - 8$. The dominant modes have $n = 2 - 4$ and a frequency in the laboratory frame $f \approx 90 - 150$ kHz. The toroidal rotation frequency, causing the Doppler shift between the mode frequency in the plasma and laboratory frames, is typically 15 – 30 kHz on the magnetic axis. As the discharge evolves, the modes exhibit a quasi-stationary behavior, then their dynamics becomes more turbulent, especially when a second NB source is added. Interestingly, however, the transition starts even before the second NB source is turned on. After $t \approx 290$ ms relatively large excursions in both amplitude and frequency are measured, eventually ending up with a dramatic burst of TAE activity. The latter is dubbed *avalanche*, and may result in a drop in the neutron rate of up to 30% [8][9], see Fig. 1. Depending on the frequency and amplitude variation modes are classified as *quasi-stationary*, *bursting/chirping* or *avalanching*. Although somewhat arbitrary, this classification provides a practical tool for the following discussion. Note that only down-chirping TAE modes are observed here, contrary to other experiments from which both upwards and downwards frequency sweeps have been reported [6][22].

The mode structure, measured by a multi-channel reflectometer system [20], extends over good fraction of the minor radius (Fig. 2) up to $R \approx 140$ cm, with maximum amplitude at $R = 120 - 130$ cm. For reference, the magnetic axis and the last closed flux surface are at $R \sim 103$ cm and $R \sim 145$ cm. The mode evolution described above is representative of discharges spanning a broad range of plasma parameters with $P_{NB} \leq 3$ MW in both helium and deuterium plasmas.

The transition of the dominant $n = 3$ mode from quasi-stationary to bursting/chirping illustrated in Fig. 1 can be quantified through the auto-correlation time of magnetic fluctuations, τ_{corr} , as a function of time (Fig. 3). τ_{corr} is calculated from an exponential fit of the the auto-correlation $C(\tau)$, where τ is the time lag, over a moving window of 1 ms. A

FIG. 3. (Top) Auto-correlation time for the dominant $n = 3$ mode, calculated by windowing the filtered signal (bandpass, 105 – 130 kHz range) in sub-samples with duration 1 ms. Note the drop in the auto-correlation after ≈ 285 ms, corresponding to the onset of the bursting/chirping regime. (Bottom) Filtered signal from a Mirnov coil, showing the beginning of amplitude bursts after $t \approx 285$ ms.

clear transition is observed as the mode develops bursting/chirping features after approximately $t = 285$ ms. Typical values before that time are $\tau_{corr} \lesssim 0.4$ ms, indicating that the oscillation remains coherent over ~ 50 wave cycles. After the transition, τ_{corr} drops to ~ 0.1 ms. This can be explained by decomposing τ_{corr} as $\tau_{corr}^{-1} = \tau_{phase}^{-1} + \tau_{freq}^{-1}$, where the two terms represent the decorrelation time associated with phase and frequency variations of the mode, respectively. As the frequency undergoes a rapid change, $\tau_{corr} \approx \tau_{freq}$ is primarily determined by the typical frequency chirp rate. The transition between quasi-stationary and bursting/chirping regimes, as seen in $C(\tau)$, occurs in ≈ 2 ms. This is much shorter than the time-scales associated with changes in the background plasma profiles (e.g. density, temperature, rotation or safety factor profile), beam ions slowing down and scattering (~ 10 's of ms).

To investigate in more detail the TAE dynamics, we focus in the following on the time window 260 – 390 ms, when TAEs show an almost constant frequency separation between peaks with consecutive n 's. This suggests that the modes have a common frequency in the plasma frame, f_0^{TAE} , such that [23]

$$f_{lab,n}^{TAE} = f_0^{TAE} + n f_{Doppler}^{TAE} \quad (1)$$

where $f_{lab,n}^{TAE}$ is the frequency for the toroidal mode number n as measured in the laboratory

FIG. 4. (a,b) Fit of the peak frequency vs. toroidal mode number based on Eq. 1 for modes with $n = 2 - 6$. The two different times correspond to the beginning of weakly bursting activity and to strong bursts. The peak frequency and its spread (indicated by the vertical bar) are obtained from a gaussian fit of the FFT spectrum obtained from a moving time window of 1.25 ms. (c) Time history of the reconstructed mode frequency in the plasma frame, f_0^{TAE} , and TAE Doppler shift frequency, $f_{Doppler}^{TAE}$.

frame and $f_{Doppler}^{TAE}$ the Doppler shift caused by plasma rotation. Diamagnetic corrections to $f_{Doppler}^{TAE}$ are estimated to be $\lesssim 1$ kHz and are neglected here, as well as the differential rotation between impurity and main ions [17]. In practice, a gaussian fit of the peaks in the FFT spectra from 1.25 ms time windows is first used to determine the center frequency and frequency spread of the different modes. Then, a linear fit based on Eq. 1 provides the values of f_0^{TAE} and $f_{Doppler}^{TAE}$, see Fig. 4a-b. The results as a function of time are reported in Fig. 4c. The calculated Doppler shift is remarkably constant in time. f_0^{TAE} does not change substantially for $t \lesssim 290$ ms, corresponding to a phase with only one NB source.

Interestingly, Eq. 1 represents well the frequency behavior over time scales $\gtrsim 1$ ms. The mechanism through which the different TAEs lock on a similar frequency are not yet clear. At present, it can be excluded that the modes are harmonics of a fundamental TAE mode or that they couple to each other through a stationary low frequency mode, as was

observed in Ref. [23]. The sub-millisecond dynamics is more complex, with each mode chirping independently of the others except for the largest bursts. The detailed analysis of *fast* frequency evolution and of the differentiation in time scales, which suggests that mode-mode coupling is at play for large amplitude bursts, are deferred to a separate publication.

By knowing the TAE Doppler shift frequency and the rotation profile, the radius at which they match each other is easily calculated and can be compared, for example, with the location of maximum TAE drive. The comparison makes more meaningful the study of the possible decorrelation mechanisms, as discussed in the next Section.

III. ROTATION SHEAR EVOLUTION

Several quantities can affect the drive and evolution of TAEs. Here we focus on (i) the fast ion profile and its radial gradient and (ii) the toroidal rotation profile, f_{rot} , and its radial shear, $\propto \partial f_{rot}/\partial R$. The fast ion density is measured on NSTX by a fast ion D-alpha (FIDA) diagnostic based on active charge-exchange recombination spectroscopy [24]. For the discharge under investigation the fast ion content increases as the second NB source is turned on at 290 ms. At the same time, the gradient on the low-field side steepens up. This correlates with an increase of activity in the TAE frequency range, consistent with the picture that the observed TAEs are driven unstable by the radial gradient of the fast ion profile. The fast ion density decreases after the avalanche at ≈ 360 ms, which causes a relaxation of the gradient [8].

Since NB injection also represents the main source of torque, the fast ion profile evolution correlates with the plasma rotation in many aspects. Figure 5 shows the measured evolution of the toroidal rotation profile and its shear for the discharge illustrated in Fig. 1. The rotation profile peaks on axis and gradually increases to ~ 30 kHz after the second NB source is added. At the same time the profile broadens. The position of maximum radial shear, here defined as the radial gradient of the rotation profile, shifts outward from $R \approx 120$ cm to $R \approx 125$ cm (Fig. 5b). The shear increases in time and a broader spatial region is affected by relatively large values.

The radius at which the plasma rotation matches the estimated Doppler shift frequency of the modes can be calculated from the measured rotation profile. This is illustrated in Fig. 5b, along with the position of steepest fast ion gradient. A first observation is that

FIG. 5. (a) Temporal evolution of the profiles of toroidal rotation frequency. (b) Radial shear of the rotation frequency, calculated as the radial gradient. The magnetic axis is at $R \approx 100$ cm. The red triangles indicate the radius at which the plasma rotation frequency equals the Doppler shift frequency estimated from the modes, cf. Fig. 4. The black diamonds show the position of steepest gradient of the fast ion profile on the low-field side.

the radii where $f_{rot} = f_{Doppler}^{TAE}$ and of the steepest fast ion gradient are approximately equal and track each other in time. Second, the same position also corresponds to the location of largest rotation shear. A qualitative explanation implies that the fast ion profile resulting from NB injection determines at the same time the TAE driving term (through its gradient) and the toroidal rotation profile. This double role of the fast ion population means that the TAE modes naturally sit around the region of maximum velocity shear, thus favoring a competition between driving and decorrelating mechanisms. These observations, along with results from previous studies [11][12], indicate that effects of rotation shear on the TAE modes might be expected.

For instabilities which are convected by a background flow, an upper limit for the decorrelation (or *shearing*) rate τ_{dec}^{-1} caused by a shear layer located at $R \approx R_{mode}$ can be calculated from the toroidal velocity profile and the parallel wave number. τ_{dec} represents the minimum

FIG. 6. Correlation analysis of two reflectometer channels displaced by ~ 8 cm across the shear layer. (a) Raw signals associated with the $n = 3$ mode measured at 117 cm (black) and 125 cm (red). (b) Frequency evolution for the two channels, reconstructed from the band-pass filtered signals. (c) Cross-correlation between the two signals. (d) Cross-phase reconstructed from the time-shift between raw signals.

time required to displace two regions of the mode structure separated by ΔR_{mode} under the effects of a local differential velocity $|\partial(2\pi R f_{rot})/\partial R|$ calculated at $R = R_{mode}$:

$$\tau_{dec}^{-1} \simeq k_{\parallel} \Delta R_{mode} 2\pi R_{mode} \left| \frac{\partial f_{rot}}{\partial R} \right|_{R=R_{mode}} \quad (2)$$

where $k_{\parallel} \approx |n - m/q_{min}|/R_{mode}$ is the parallel wavenumber, R_{mode} the location of maximum mode amplitude and ΔR_{mode} the radial extension of the mode. From Fig. 1-5, typical values of $|\partial f_{rot}/\partial R|$ measured at $R = 120 - 135$ cm, where the TAEs have their maximum amplitude, result in a decorrelation time $\tau_{dec} \sim 1$ ms. Values $\Delta R_{mode} = 10 - 20$ cm, derived from reflectometer measurements of the mode structure (Fig. 2), are used. Although these values are roughly consistent with the calculated auto-correlation time for the dominant TAEs, no steep transition in the rotation (or its shear) is observed. Therefore, the decorrelation from sheared rotation alone seems not responsible for the observed mode behavior.

FIG. 7. Frequency of the $n = 3$ mode calculated from two reflectometer channels (cf. Fig. 6b) for discharge no. 135388, showing no systematic difference across the shear layer.

IV. MODE FREQUENCY, CROSS-PHASE AND EFFECTIVE GROWTH/DAMPING RATES

The NSTX multi-channel reflectometer system provides radially localized information on the mode evolution. An eventual shearing off of the mode through differential plasma rotation should therefore appear from correlation analysis of channels measuring across the shear layer. Results for the dominant $n = 3$ mode are shown in Fig. 6 for a time window corresponding to two bursts (Fig. 6a). The two channels measure at $R \approx 117$ cm and $R \approx 125$ cm. For each channel, the instantaneous frequency (in the laboratory frame) is reconstructed from the band-pass filtered signals (Fig. 6b). The mode frequency shifts down by ~ 10 kHz over $300 \mu\text{s}$ with no significant difference as a function of the minor radius (Fig. 7). The frequencies are significantly different only when the signal-to-noise is too low and the analysis fails. This can be seen from the vanishing amplitude and quantified by a sudden drop in the cross-correlation (Fig. 6a-c). Finally, the phase shift between the two channels is shown in Fig. 6d. No systematic variations of the cross-phase are observed. The invariance of frequency and cross-phase is in contrast with the picture that the mode experiences a differential rotation, which should appear as a Doppler shift contribution to $f_{lab,n}^{TAE}$ (Eq. 1) that depends on the radial position.

The analysis of the time scales of the TAE bursts complements the previous conclusions. The mode amplitude evolution during a burst provides insight into the effective rise and

FIG. 8. (a) Evolution of the maximum rotation shear. (b) Rms mode amplitude for the $n = 3$ mode in Fig. 1. (c) Amplitude rise (red, diamonds) and decay (blue, triangles) rates normalized to the mode frequency, estimated from an exponential fit of the rising/decaying part of TAE bursts. The dashed lines show the decorrelation rate from rotation shear calculated through Eq. 2.

decay time for the mode amplitude, γ_{rise} and γ_{decay} . The rising (falling) amplitude is fitted with a increasing (decreasing) exponential to estimate the corresponding time constant. Considering the possible non-linear dynamics of the modes [4] and the still finite drive and damping at play during both growth and decay phases, these quantities should not be confused with the linear growth and damping rates of the modes. They represent an experimentally-accessible estimate of the typical time scales governing the mode evolution. Only in the limit of vanishing damping during the amplitude growth and vanishing drive during its decay the two time constants approximate the linear rates, under the additional assumption of linear or weakly non-linear dynamics. The normalized ratio of rise/decay rate to the mode frequency can be inferred by dividing γ_{rise} and γ_{decay} by $\omega = 2\pi f_{lab,n}^{TAE}$. Figure 8 shows the results of this analysis by comparing the temporal evolution of the maximum rotation shear, rms mode amplitude (as measured by Mirnov coils) and the estimated γ_{rise}/ω and γ_{decay}/ω ratios. Rise and decay terms do not vary in a consistent way with the $> 50\%$ increase in the rotation gradient, the dramatic change in the mode amplitude and the overall

mode dynamics. The damping rate from Eq. 2, calculated for $n = 3$ and poloidal mode number $m = -6 \dots +6$, remains below 1%, i.e. considerably smaller than the estimated γ_{rise} and γ_{decay} . (In this case, the γ/ω ratio is calculated by using the reconstructed frequency in the plasma frame).

In general, no statistically significant trend of the effective damping and decay rates as a function of the maximum shear is observed. This is illustrated in Fig. 9 for $n = 3$ mode activity from a series of similar discharges with varying level of injected NB and RF power. (Similar results are obtained for other modes with different n 's). Since electron density and temperature span a wide range, four subsets are shown separately to group data characterized by similar values of damping on the thermal plasma. The role of other damping mechanisms, such as continuum damping, is not considered in this analysis.

V. IMPORTANCE OF TAE DRIVE VS. ROTATION SHEAR EFFECTS

The results from the previous Sections suggest an alternative interpretation for the bursting TAE behavior based on the role of the TAE drive instead of the damping through rotation shear. In fact, the condition $\gamma_{decay} \approx \gamma_{rise}$ with both terms large ($> 2\%$, according to Figs. 8c-9) reminds the requisite of many models for the appearance of bursting behavior, namely nearly equal and large drive and damping rates [25]. Although the models are usually one-dimensional and based on a bump-on-tail excitation of the modes, their results can be qualitatively generalized to a more complex case with spatial (instead of velocity) gradient as steady source of free energy, to which kinetic effects resulting from wave-particle interactions add.

Experimentally, two ways are available to modify the drive for TAEs. First, NB sources with different tangency radius can be used. For the shot discussed so far (cf. Fig. 1) the two sources with smaller tangency radius, implying more perpendicular injection, were used. In the following, the source at tangency radius of 69 cm is utilized, providing more tangential injection and the possibility of direct measurements of the q -profile through Motional Stark Effect [26]. Second, RF injection can be used to affect the evolution of the fast ion and rotation profiles [12][27], both ultimately dependent upon the beam deposition profile in NB-only discharges. RF waves modify the original beam distribution by scattering and accelerating fast ions. The effect is especially important in a spherical tokamak such as

FIG. 9. (Top) Parameter space from an ensemble of 10 discharges with different NB timing and power. The bottom panels show the amplitude decay (left column) and rise (right column) rates vs. maximum rotation shear for the subsets of density and temperature values indicated in the top panel as regions *I* to *IV*. Symbols and vertical bars indicate the average and standard deviation of the measured rates over 10 ms, corresponding to the sampling time of toroidal rotation measurements. Red symbols denote time points with up to 1.5 MW of injected RF power in addition to NB injection.

NSTX where numerous resonances are simultaneously present over the entire minor radius [28].

Three shots with different timing of NB and RF injection are compared. Apart from the auxiliary power the plasma profiles are similar, at least in the initial ≈ 350 ms of the discharge (Fig. 10). Unlike the discharge of Fig. 1, plasmas transition into H-mode. Reflectometer data is therefore not available because of the flat density profile inside $R \sim 140$ cm. The first discharge has $P_{NB} = 2$ MW with a beam notch (i.e. zero power) at 320 – 340 ms, see Figs. 11 and 12a. The injected RF power is constant at $P_{RF} = 2$ MW from $t \approx 260$ ms on. Activity in the Alfvén band is similar to what was described earlier, with RSAEs

FIG. 10. Radial profiles of (a) electron density, (b) electron temperature, (c) toroidal velocity shear and (d) q profile at $t = 310$ ms for the L-mode discharge in Fig. 1 (black) and for two H-mode discharges with (red) and without (blue) HHFW injection. Strong TAE activity with similar bursting/chirping features is measured (Fig. 11) except for no-RF discharge.

shifting into TAEs at the beginning of the current flat-top. A bursting phase precedes a phase with larger mode activity, during which avalanches eventually occur (Fig. 11). TAE activity decays during the beam notch, then increases again as the beam turns back on, consistent with the fact that the injected fast ions are the primary drive for TAEs. No change in rotation shear is observed (Fig. 12d). The second discharge is similar, but with constant P_{NB} and a notch in P_{RF} at 350 – 370 ms (Fig. 12b). TAE activity decreases during the notch, where only small bursts are observed until the RF is turned back on and the mode amplitude is restored to a level comparable to that before the notch. Again, rotation does not change substantially (Fig. 12e). In the third discharge the injected RF power is zero. RSAEs are observed during the ramp-up phase, but they decay after $t \approx 270$ ms. No TAEs are measured afterwards. These observations are explained by assuming that a threshold in the fast ion population has not been crossed [9] compared to the discharge in Fig. 1, and NB-only injection is not sufficient to destabilize TAEs. When RF is applied, acceleration of fast ions partly compensate the Coulomb drag and their distribution function is modified [28], making the drive for TAEs stronger. However, RF alone is not enough to sustain the fast ion population and TAEs decay when the NB is switched off as in Fig. 12a. All these

FIG. 11. (Top) Spectrogram of TAE activity for discharge #134934. Note the decay of TAE activity during the NB notch indicating that the fast ion population, hence the drive for the modes, is decreasing, as observed in the neutron rate. (Bottom) Waveforms of injected NB and RF power and measured neutron rate.

considerations point to the fast ion drive as primary cause for the observed TAE dynamics, with rotation shear playing only a marginal role.

VI. SUMMARY AND CONCLUSIONS

The role of a shear layer in the toroidal rotation profile on the dynamics of TAE modes has been studied in L- and H-mode discharge on NSTX. Toroidal rotation is driven by NB injection, which also represents the source term for the fast ion population. This determines a coupling between the spatial location of the shear layer and of the TAE drive. H-mode discharges with simultaneous NB and RF injection are partially successful in decoupling the two terms, but no strong modifications of the TAE behavior is observed. Results from a multi-channel reflectometer system measuring TAE fluctuations across the shear layer indicate that the modes rotate rigidly, without any evidence of spatial decorrelation on the frequency and phase shift evolution caused by the velocity gradient. This indicates that sheared rotation is much less efficient in decorrelating global, coherent modes as TAEs as compared to other turbulent instabilities, for which sheared flows provide an effective suppression mechanism [29][30]. The experimental results lead to consider the TAE drive term, rather than the decorrelation from rotation shear, as the primary cause for the observed bursting dynamics of the modes. Experiments aimed at altering the TAE drive, both in real

FIG. 12. Comparison of rms TAE activity (blue) for three discharges with different NB (black) and RF (red) power waveforms. TAE activity is stronger when both auxiliary power sources are ON, whereas NB injection alone is insufficient to sustain the modes. (a) $P_{NB} \approx P_{RF} \approx 2$ MW, with a notch in P_{NB} . (b) $P_{NB} \approx P_{RF} \approx 2$ MW, with a notch in P_{RF} . (c) $P_{NB} \approx 2$ MW and $P_{RF} = 0$. (d)-(e) Same as in Fig. 5 for discharges #134934 (with RF) and #134935 (no RF).

space (through variations of the tangency radius of the NB source) and in velocity space (by applying RF heating) are consistent with this conclusion.

Apart from the shear in rotation, it is important to point out that the rotation profile itself can strongly modify the structure of the Alfvén continuum [31]. As a consequence, the stability of TAE modes may also change with rotation, e.g. because of changes in the continuum damping. An example based on the NOVA-K code [32] is illustrated in Fig. 13 for the discharge in Fig. 1 at $t = 280$ ms, i.e. just before the beginning of the bursting/chirping phase. It is seen that f_{rot} , and not its gradient, may determine whether an unstable mode in the TAE gap feels a strong continuum damping or not. Note that in this calculation the finite rotation is included as a simple Doppler shift of the calculated gap frequencies; it is not included self-consistently in the computation of TAE dispersion relation and continuum structure. A closer comparison with numerical codes would be required for more detailed investigations. A version of NOVA-K including the physics of finite plasma rotation is being developed, and will be used in future studies. Other codes, such as the non-linear, self-consistent code M3D-K ([33] and references therein) already include toroidal

FIG. 13. (a) TAE gap for $n = 3$ calculated via the NOVA-K code at $t = 280$ ms for discharge #135388, cf. Fig. 1. The blue dashed line shows the Doppler shift frequency for $n = 3$, as determined from the measured rotation profile. The red line indicates the measured frequency (in the laboratory frame) of the $n = 3$ mode, which interacts with the continuum at $R \approx 114$ cm. (b) Same calculation, but assuming a 6% reduction in the rotation. The measured frequency would now interact with the continuum at the magnetic axis, where the mode amplitude vanishes), which implies a negligible contribution from continuum damping to the total damping.

rotation. Initial results indicate that modifications of the TAE gap structure for different rotation profiles are more important for TAE dynamics than the rotation shear [34], which is consistent with the results presented herein. Regarding stability calculations for TAEs, a correct quantitative treatment of continuum damping, including for instance kinetic effects, is still missing in the numerical codes. It would be required for quantitative comparisons of calculated mode stability against experiments. At the same time good progress has been made in the past few years to understand and model the interaction between fast ions and RF [19][28][35]. Integrating those tools in codes such as TRANSP [36], in order to make analysis of RF discharges routinely available, is still work in progress. It may provide a powerful tool to investigate TAE dynamics on a quantitative basis.

Future experiments on NSTX will further investigate the role of rotation and drive in the TAE dynamics. Improved characterization of the mode structure is expected through Beam Emission Spectroscopy [37]. A second FIDA system, more sensitive to the tangential

component of the fast ion distribution function, will provide more details on the fast ion dynamics [38]. In addition, the possibility of using magnetic braking [39] to alter the rotation profile independently of the NB torque source will be explored.

ACKNOWLEDGMENTS

Discussions with Dr. G.-Y. Fu and the support of the NSTX NB and RF groups are gratefully acknowledged. This work is supported by US-DOE grant DE-FG02-06ER54867 and by US DOE contracts no. DE-AC02-09CH11466 and DE-FG02-99ER54527.

LIST OF FIGURES

1	(a) Spectrogram showing TAE activity in the range $f = 70 - 200$ kHz for discharge #135388. (b) Temporal evolution of the central electron density and temperature. (c) Waveforms of NB power and neutron rate. Note the drop of the neutron rate at $t \approx 360$ ms, corresponding to an <i>avalanche</i> , which indicates a loss of fast ions from the core plasma.	3
2	Displacement associated with a $n = 3$ mode, measured by a multi-channel reflectometer at three different times.	4
3	(Top) Auto-correlation time for the dominant $n = 3$ mode, calculated by windowing the filtered signal (bandpass, 105 – 130 kHz range) in sub-samples with duration 1 ms. Note the drop in the auto-correlation after ≈ 285 ms, corresponding to the onset of the bursting/chirping regime. (Bottom) Filtered signal from a Mirnov coil, showing the beginning of amplitude bursts after $t \approx 285$ ms.	6
4	(a,b) Fit of the peak frequency vs. toroidal mode number based on Eq. 1 for modes with $n = 2 - 6$. The two different times correspond to the beginning of weakly bursting activity and to strong bursts. The peak frequency and its spread (indicated by the vertical bar) are obtained from a gaussian fit of the FFT spectrum obtained from a moving time window of 1.25 ms. (c) Time history of the reconstructed mode frequency in the plasma frame, f_0^{TAE} , and TAE Doppler shift frequency, $f_{Doppler}^{TAE}$	7
5	(a) Temporal evolution of the profiles of toroidal rotation frequency. (b) Radial shear of the rotation frequency, calculated as the radial gradient. The magnetic axis is at $R \approx 100$ cm. The red triangles indicate the radius at which the plasma rotation frequency equals the Doppler shift frequency estimated from the modes, cf. Fig. 4. The black diamonds show the position of steepest gradient of the fast ion profile on the low-field side.	9

6	Correlation analysis of two reflectometer channels displaced by ~ 8 cm across the shear layer. (a) Raw signals associated with the $n = 3$ mode measured at 117 cm (black) and 125 cm (red). (b) Frequency evolution for the two channels, reconstructed from the band-pass filtered signals. (c) Cross-correlation between the two signals. (d) Cross-phase reconstructed from the time-shift between raw signals.	10
7	Frequency of the $n = 3$ mode calculated from two reflectometer channels (cf. Fig. 6b) for discharge no. 135388, showing no systematic difference across the shear layer.	11
8	(a) Evolution of the maximum rotation shear. (b) Rms mode amplitude for the $n = 3$ mode in Fig. 1. (c) Amplitude rise (red, diamonds) and decay (blue, triangles) rates normalized to the mode frequency, estimated from an exponential fit of the rising/decaying part of TAE bursts. The dashed lines show the decorrelation rate from rotation shear calculated through Eq. 2. . .	12
9	(Top) Parameter space from an ensemble of 10 discharges with different NB timing and power. The bottom panels show the amplitude decay (left column) and rise (right column) rates vs. maximum rotation shear for the subsets of density and temperature values indicated in the top panel as regions <i>I</i> to <i>IV</i> . Symbols and vertical bars indicate the average and standard deviation of the measured rates over 10 ms, corresponding to the sampling time of toroidal rotation measurements. Red symbols denote time points with up to 1.5 MW of injected RF power in addition to NB injection.	14
10	Radial profiles of (a) electron density, (b) electron temperature, (c) toroidal velocity shear and (d) q profile at $t = 310$ ms for the L-mode discharge in Fig. 1 (black) and for two H-mode discharges with (red) and without (blue) HHFW injection. Strong TAE activity with similar bursting/chirping features is measured (Fig. 11) except for no-RF discharge.	15
11	(Top) Spectrogram of TAE activity for discharge #134934. Note the decay of TAE activity during the NB notch indicating that the fast ion population, hence the drive for the modes, is decreasing, as observed in the neutron rate. (Bottom) Waveforms of injected NB and RF power and measured neutron rate.	16

- 12 Comparison of rms TAE activity (blue) for three discharges with different NB (black) and RF (red) power waveforms. TAE activity is stronger when both auxiliary power sources are ON, whereas NB injection alone is insufficient to sustain the modes. (a) $P_{NB} \approx P_{RF} \approx 2$ MW, with a notch in P_{NB} . (b) $P_{NB} \approx P_{RF} \approx 2$ MW, with a notch in P_{RF} . (c) $P_{NB} \approx 2$ MW and $P_{RF} = 0$. (d)-(e) Same as in Fig. 5 for discharges #134934 (with RF) and #134935 (no RF). 17
- 13 (a) TAE gap for $n = 3$ calculated via the NOVA-K code at $t = 280$ ms for discharge #135388, cf. Fig. 1. The blue dashed line shows the Doppler shift frequency for $n = 3$, as determined from the measured rotation profile. The red line indicates the measured frequency (in the laboratory frame) of the $n = 3$ mode, which interacts with the continuum at $R \approx 114$ cm. (b) Same calculation, but assuming a 6% reduction in the rotation. The measured frequency would now interact with the continuum at the magnetic axis, where the mode amplitude vanishes), which implies a negligible contribution from continuum damping to the total damping. 18

-
- [1] A. Fasoli, C. Gormezano, H. L. Berk, B. N. Breizman, S. Briguglio, D. S. Darrow, N. N. Gorelenkov, W. W. Heidbrink, A. Jaun, S. V. Konovalov, R. Nazikian, J.-M. Noterdaeme, S. Sharapov, K. Shinoara, D. Testa, K. Tobita, Y. Todo, G. Vlad, and F. Zonca, *Nucl. Fusion* **47**, S264 (2007).
 - [2] W. W. Heidbrink, *Phys. Plasmas* **15**, 055501 (2008).
 - [3] B. V. Chirikov, *Phys. Rep.* **52**, 263 (1979).
 - [4] H. L. Berk, B. N. Breizman, and M. S. Pekker, *Phys. Plasmas* **2**, 3007 (1995).
 - [5] A. Fasoli, B. N. Breizman, D. Borba, R. F. Heeter, M. S. Pekker, and S. E. Sharapov, *Phys. Rev. Lett.* **81**, 5564 (1998).
 - [6] M. P. Gryaznevich and S. E. Sharapov, *Plasma Phys. Control. Fusion* **46**, S15 (2004).
 - [7] S. S. Medley, N. N. Gorelenkov, R. Andre, R. E. Bell, D. S. Darrow, E. D. Fredrickson, S. M. Kaye, B. P. LeBlanc, A. L. Roquemore, and NSTX team, *Nucl. Fusion* **44**, 1158 (2004).
 - [8] M. Podestà, W. W. Heidbrink, D. Liu, E. Ruskov, R. E. Bell, D. S. Darrow, E. D. Fredrickson, N. N. Gorelenkov, G. J. Kramer, B. P. LeBlanc, S. S. Medley, A. L. Roquemore, N. A. Crocker, S. Kubota, and H. Yuh, *Phys. Plasmas* **16**, 056104 (2009).
 - [9] E. D. Fredrickson, N. A. Crocker, R. E. Bell, D. S. Darrow, N. N. Gorelenkov, G. J. Kramer, S. Kubota, F. M. Levinton, D. Liu, S. S. Medley, M. Podestà, K. Tritz, R. B. White, and H. Yuh, *Phys. Plasmas* **16**, 122505 (2009).
 - [10] M. Ono, S. M. Kaye, Y.-K. M. Peng, G. Barnes, W. Blanchard, M. D. Carter, J. Chrzanowski, L. Dudek, R. Ewig, D. Gates, R. E. Hatcher, T. Jarboe, S. C. Jardin, D. Johnson, R. Kaita, M. Kalish, C. E. Kessel, H. Kugel, R. Maingi, R. Majeski, J. Manickam, B. McCormack, J. Menard, D. Mueller, B. Nelson, B. Nelson, C. Neumeyer, G. Oliaro, F. Paoletti, R. Parsells, E. Perry, N. Pomphrey, S. Ramakrishnan, R. Raman, G. Rewoldt, J. Robinson, A. L. Roquemore, P. Ryan, S. Sabbagh, D. Swain, E. J. Synakowski, M. Viola, M. Williams, J. R. Wilson, and NSTX Team, *Nucl. Fusion* **40**, 557 (2000).
 - [11] D. Testa, C. Boswell, A. Fasoli, and JET-EFDA Contributors, *Nucl. Fusion* **45**, 907 (2005).
 - [12] M. Saigusa, Y. Kusama, T. Ozeki, H. Kimura, T. Fujita, S. Moriyama, T. Fujii, M. Azumi, V. I. Afanassiev, and Y. Neyatani, *Nucl. Fusion* **37**, 1559 (1997).
 - [13] D. A. Spong, B. A. Carreras, and C. L. Hedrick, *Phys. Plasmas* **1**, 1503 (1994).

- [14] E. D. Fredrickson, N. N. Gorelenkov, R. E. Bell, J. E. Menard, A. L. Roquemore, S. Kubota, N. A. Crocker, and W. Peebles, Nucl. Fusion **46**, S926 (2006).
- [15] B. P. LeBlanc, Rev. Sci. Instr. **79**, 10E737 (2008).
- [16] R. E. Bell, Phys. Plasmas, accepted for publication(2010).
- [17] Y. B. Kim, P. H. Diamond, and R. J. Groebner, Phys. Fluids B **3**, 2050 (1991).
- [18] S. M. Kaye, W. Solomon, R. E. Bell, B. P. LeBlanc, F. Levinton, J. E. Menard, G. Rewoldt, S. Sabbagh, W. Wang, and H. Yuh, Nucl. Fusion **49**, 045010 (2009).
- [19] G. Taylor, R. E. Bell, J. C. Hosea, B. P. LeBlanc, C. K. Phillips, M. Podestà, E. J. Valeo, J. R. Wilson, J.-W. Ahn, G. Chen, D. L. Green, E. F. Jaeger, R. Maingi, P. M. Ryan, J. B. Wilgen, W. W. Heidbrink, D. Liu, P. T. Bonoli, T. Brecht, M. Choi, and R. W. Harvey, Phys. Plasmas **17**, 056114 (2010).
- [20] N. A. Crocker, E. D. Fredrickson, N. N. Gorelenkov, G. J. Kramer, D. S. Darrow, W. W. Heidbrink, S. Kubota, F. M. Levinton, H. Yuh, J. E. Menard, B. P. LeBlanc, and R. E. Bell, Phys. Plasmas **15**, 102502 (2008).
- [21] J. E. Menard, private communication.
- [22] N. P. Young, S. E. Sharapov, V. M. Nakariakov, and JET EFDA contributors, Plasma Phys. Control. Fusion **48**, 295 (2006).
- [23] E. J. Strait, W. W. Heidbrink, and A. D. Turnbull, Plasma Phys. Control. Fusion **36**, 1211 (1994).
- [24] M. Podestà, W. W. Heidbrink, R. E. Bell, and R. Feder, Rev. Sci. Instr. **78**, 10E521 (2008).
- [25] H. L. Berk, B. N. Breizman, J. Fitzpatrick, M. S. Pekker, H. V. Wong, and K. L. Wong, Phys. Plasmas **3**, 1827 (1996).
- [26] F. M. Levington, R. J. Fonck, G. M. Gammel, R. Kaita, H. W. Kugel, E. T. Powell, and D. W. Roberts, Phys. Rev. Lett. **63**, 2060 (1989).
- [27] D. Borba, G. D. Conway, S. Guenter, G. T. A. Huysmans, S. Klose, M. Maraschek, A. Mueck, I. Nunes, S. D. Pinches, F. Serra, and the ASDEX Upgrade Team, Plasma Phys. Control. Fusion **46**, 809 (2004).
- [28] D. Liu, W. W. Heidbrink, M. Podestà, R. E. Bell, E. D. Fredrickson, S. S. Medley, R. W. Harvey, and E. Ruskov, Plasma Phys. Control. Fusion **52**, 025006 (2010).
- [29] K. H. Burrell, Phys. Plasmas **4**, 1070 (1997).
- [30] P. W. Terry, Rev. Mod. Phys. **72**, 109 (2000).

- [31] B. van der Holst, A. J. C. Belien, and J. P. Goedbloed, *Phys. Plasmas* **7**, 4208 (2000).
- [32] C. Z. Cheng, *Phys. Rep.* 211 **1** (1992).
- [33] J. Lang, Y. Chen, S. E. Parker, and G. Fu, *Phys. Plasmas* **16**, 102101 (2009).
- [34] G.-Y. Fu, private communication.
- [35] M. Choi, D. Green, W. W. Heidbrink, R. W. Harvey, D. Liu, V. S. Chan, L. A. Berry, F. Jaeger, L. L. Lao, R. I. Pinsky, M. Podestà, D. N. Smithe, J. M. Park, and P. T. Bonoli, *Phys. Plasmas* **17**, 056102 (2010).
- [36] R. V. Budny, *Nucl. Fusion* **34**, 1247 (1994).
- [37] D. Smith *et al.*, *Rev. Sci. Instr.*, Proceedings of the 18th HTPD meeting, to be published(2010).
- [38] A. Bortolon, W. W. Heidbrink, and M. Podestà, *Rev. Sci. Instr.*, Proceedings of the 18th HTPD meeting, to be published(2010).
- [39] W. Zhu, S. A. Sabbagh, R. E. Bell, J. M. Bialek, M. G. Bell, B. P. LeBlanc, S. M. Kaye, F. M. Levinton, J. E. Menard, K. C. Shaing, A. C. Sontag, and H. Yuh, *Phys. Rev. Lett.* **96**, 225002 (2006).

The Princeton Plasma Physics Laboratory is operated
by Princeton University under contract
with the U.S. Department of Energy.

Information Services
Princeton Plasma Physics Laboratory
P.O. Box 451
Princeton, NJ 08543

Phone: 609-243-2245
Fax: 609-243-2751
e-mail: pppl_info@pppl.gov
Internet Address: <http://www.pppl.gov>