10
21-74

Dr. 656

LA-5409

The Kinetics of an H_2-F_2 Explosion as
Initiated by a Pulse of Fluorine Atoms

MASTER

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

Printed in the United States of America. Available from
National Technical Information Service
U. S. Department of Commerce
5285 Port Royal Road
Springfield, Virginia 22151
Price: Printed Copy \$4.00; Microfiche \$1.45

LA-5409

UC-34

ISSUED: February 1974

The Kinetics of an H_2 - F_2 Explosion as Initiated by a Pulse of Fluorine Atoms

by

John H. Sullivan
Roy C. Feber

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

GO

THE KINETICS OF AN H_2-F_2 EXPLOSION AS INITIATED
BY A PULSE OF FLUORINE ATOMS

by

John H. Sullivan and Roy C. Feber

ABSTRACT

The elementary reactions occurring in an H_2-F_2 explosion are given with each vibrational level of H_2 and HF considered as a separate reacting species. Rate coefficients for all the elementary reactions are either estimated or calculated, and the major assumptions entering into the determination of each rate coefficient are given. The effect of non-Maxwell distributions in the translational energies of some of the species is estimated. For strong pulses of F atoms, the explosion is complete before any appreciable contribution to the rate is made by thermal dissociation of F_2 or H_2 .

The chemical change that produces lasing energy in H_2-F_2 -HF systems has recently been modeled on computers.¹⁻⁵ The purpose of such computations is to gain insight into the processes that determine the amount of lasing power available and to try to reproduce gross (as compared to the detail of the model), macroscopic observations. The chemical change takes place by a series of elementary chemical reactions (such as those given in a thorough account by Cohen⁶) in which each vibrational state of H_2 and HF is necessarily considered as a separate species. However, in these models¹⁻⁵ other nonequilibrium effects in the chemistry have generally been neglected, and the transferability of thermal rate coefficients from one system to another has been implicitly accepted.

The kinetics model described here differs from previous models: (1) The rate coefficients for

are obtained for reactions taking place with a mixture of "hot" and thermal atoms. (2) The reactions of

are included and are shown to have an appreciable effect on the chemical change. (3) In an explosion initiated by a pulse of F atoms the rate of the overall reaction is shown to be fast compared with the vibrational relaxation of F_2 and H_2 . The distribution of F_2 and H_2 over vibrational states is unaffected by the rise in translational temperature. The rates of dissociation of F_2 , H_2 , and HF are negligible during the explosion. Also, the distribution of $HF(v)$ from $H + F_2 \rightarrow HF(v) + F$ is primarily from $H + F_2(0) \rightarrow HF(v) + F$. (4) The distribution of H atoms over translational energies is shown to be much different from a Maxwell distribution. When one is characterizing H-atom reactions in the H_2-F_2 explosion, this distribution over translational energies rather than the thermal distribution, should be used in future trajectory calculations. (5) The rates of V-R,T relaxation by H and F atoms of the higher vibrational states in HF are based on trajectory calculations and not on the harmonic oscillator approximation. The rates of V-R,T relaxation of $HF(v>1)$ by HF and H_2 are estimated from the rates of relaxation by F atoms as determined by trajectory calculations. (6) Qualitative assessments of the effect of rotational excitation are made when possible.

Many assumptions are made. For example, the rate coefficients for the V-R,T self-relaxation of the higher vibrational states of HF are based on at least five assumptions: (1) The V dependence is the same as that for V-R,T relaxation by F atoms. (2) All HF(v') have the same efficiency in relaxing HF(v). (3) The relaxation is described by single-quantum transitions. (4) The effect of rotation excitation on the relaxation rate is taken to be negligible. (5) The temperature dependence of the relaxation rates for the higher V states is the same as that for the v = 1 state.

The model applies to a reaction initiated by a strong pulse of F atoms, and the time dependence of the concentration of F atoms in the initiating pulse can be modeled to fit experimental data. Although oxygen is used to stabilize experimental H₂-F₂ mixtures, our model contains no reactions of H or F atoms with O₂ because the rates of these reactions are negligible compared to the rates of the chain-carrying reactions H + F₂ and F + H₂. The kinetics in the self-explosion of H₂-F₂-O₂ mixtures, however, depends strongly on reactions of atoms with O₂.

The elementary reactions and the rate coefficients are given in Table I. The table also gives free-energy terms for reactions in which the vibrational states of H₂ and HF are specified. These terms modify the usual thermochemical free-energy change and are necessary for the determination of the rates of reverse reactions. Reverse reactions are included for all reactions except (23)-(30), where the reverse reactions are strongly endothermic, and (157)-(178), where the reverse rates are low because the diatoms are not vibrationally excited by the translational environment. The relaxation of HF by He, reactions (136)-(142), is included so that the model can be used with systems moderated with inert gas. The rates of the recombination reactions (157)-(186) and of the branching reactions (187)-(195) [where a choice of v is offered in H₂(v) and HF(v)] are negligible in a fast explosion, but the reactions are included for use in studies on explosion limits.

Some rate coefficients, not directly applicable to an exploding H₂-F₂ system, are calculated as an approximation to the rates present in the experimental system, and also for the purpose of comparing and assessing different methods of obtaining rate

coefficients. The reverse reactions (-2)-(-22) take place with hot H atoms in the experimental system. Initially, however, we obtained the rates of these reactions by two methods for thermal distributions of H atoms but not for the distribution of H atoms actually present in the system. The thermal rate coefficients for (-2)-(-22) were obtained from equilibrium constants and the rates of the forward (near-thermal) reactions, where the forward rates were obtained from a combination of experimental results and trajectory calculations. The thermal rate coefficients for (-2)-(-22) were also obtained directly from trajectory calculations. Although not directly applicable to our model, the two sets of rate coefficients are necessary as a first approximation and also are needed in any attempt to assess the accuracy of various methods of calculation. Rate constants for (-2)-(-22) are presently being obtained from trajectory calculations, using for H atoms the distributions of velocity that we think are present in the experimental system.

Similarly, some quantities such as the translational distributions of F atoms are calculated to obtain a better picture of the physical system, although such distributions may not affect the rate coefficients importantly.

The relative importance of reactions involving atoms or other "intermediates" in determining the concentrations of HF(v) can be realized only when the concentrations of the atoms or intermediates are known. We have justified the use of some reactions in our model by obtaining their contributions from computer runs in which the explosion was simulated and in which the reactions were shown to significantly affect the concentrations of HF(v). These computations used earlier and different values for rate coefficients, but we consider the conclusions to be qualitatively correct. Computer runs using the present set of rate coefficients are now being carried out.

The kinetics model consists of Table I, which lists the elementary reactions and rate coefficients, and Table II, which lists the vibrational energy levels of H₂ and HF. The main body of this report comprises the "Notes" referred to in Table I and discussed at length immediately following Table II. In these Notes we attempt to justify our choice of reactions and our methods of obtaining the rate

TABLE I
RATE COEFFICIENTS USED IN SIMULATING THE EXPLOSIVE REACTION OF H₂ WITH F₂
WHEN THE EXPLOSION IS INITIATED BY A STRONG PULSE OF F ATOMS

Reaction No.	Reaction	Rate Coefficients ^a (cm ³ /mole-sec and cm ³ /mole ² -sec)	ΔE_v (cm ⁻¹)	Note ^b
(1)	F + H ₂ (0) = HF(0) + H	RK(1) = 0	0	1,2,4 5
(2)	(1)	(2) = $1.75 \times 10^{13} 10^{-340/T}$	-3961.6	
(3)	(2)	(3) = 4RK(2)	-7751.0	
(4)	(3)	(4) = 2.4RK(2)	-11,373.0	
(5)	F + H ₂ (1) = HF(1) + H	RK(5) = 2.1×10^{13}	+198.8	3,4,5
(6)	(2)	(6) = 1.09RK(5)	-3590.6	
(7)	(3)	(7) = 2.36RK(5)	-7212.6	
(8)	(4)	(8) = 4.63RK(5)	-10,671.5	
(9)	F + H ₂ (2) = HF(2) + H	RK(9) = 2.44×10^{13}	+333.4	3,4,5
(10)	(3)	(10) = 1.62 RK(9)	-3288.6	
(11)	(4)	(11) = 3.37RK(9)	-6747.4	
(12)	(5)	(12) = 6.5RK(9)	-10,046.8	
(13)	F + H ₂ (3) = HF(3) + H	RK(13) = 7.5×10^{13}	+405.3	3,4,5
(14)	(4)	(14) = RK(13)	-3053.5	
(15)	(5)	(15) = 1.57RK(13)	-6352.8	
(16)	(6)	(16) = 2.78RK(13)	-9495.7	
(17)	(7)	(17) = 0.78RK(13)	-12,484.2	
(18)	F + H ₂ (4) = HF(3) + H	RK(18) = 1.07×10^{13}	+3872.7	3,4,5
(19)	(4)	(19) = 4RK(18)	+413.8	
(20)	(5)	(20) = 7RK(18)	-2885.5	
(21)	(6)	(21) = 13RK(18)	-6028.3	
(22)	(7)	(22) = 25RK(18)	-9016.9	
(23)	H + F ₂ = HF(0) + F	RK(23) = 0	0	6,7
(24)	(1)	(24) = $3.2 \times 10^{11} 10^{-370/T}$	-3961.6	
(25)	(2)	(25) = 55RK(24)	-7751.0	
(26)	(3)	(26) = 177RK(24)	-11,373.0	
(27)	(4)	(27) = 220RK(24)	-14,831.9	
(28)	(5)	(28) = 309RK(24)	-18,131.2	
(29)	(6)	(29) = 166RK(24)	-21,274.0	
(30)	(7)	(30) = 72RK(24)	-24,262.6	
(31)	HF + HF = HF + HF	F		8
(31)	1 + 1 = 0 + 2	1.	+172	
(32)	1 + 2 = 0 + 3	0.63	+339	
(33)	1 + 3 = 0 + 4	0.4	+503	
(34)	1 + 4 = 0 + 5	0.2	+662	
(35)	1 + 5 = 0 + 6	0.12	+819	
(36)	1 + 6 = 0 + 7	0.11	+973	
(37)	2 + 2 = 1 + 3	3.	+167	
(38)	2 + 3 = 1 + 4	1.8	+331	
(39)	2 + 4 = 1 + 5	1.	+490	
(40)	2 + 5 = 1 + 6	0.54	+647	
(41)	2 + 6 = 1 + 7	0.28	+801	
(42)	3 + 3 = 2 + 4	6.	+163	
(43)	3 + 4 = 2 + 5	3.5	+322	
(44)	3 + 5 = 2 + 6	1.9	+480	
(45)	3 + 6 = 2 + 7	1.1	+633	
(46)	4 + 4 = 3 + 5	10.	+159	
(47)	4 + 5 = 3 + 6	5.7	+316	
(48)	4 + 6 = 3 + 7	3.1	+470	
(49)	5 + 5 = 4 + 6	16.	+156	
(50)	5 + 6 = 4 + 7	8.7	+310	
(51)	6 + 6 = 5 + 7	23.	+154	

Reaction No.	Reaction	Rate Coefficients ^a (cm ³ /mole-sec and cm ⁶ /mole ² -sec)	ΔE_v (cm ⁻¹)	Note ^b
	HF + H ₂ = HF + H ₂			
(52)	0 + 1 = 1 + 0	1.	+199	9
(53)	1 + 1 = 0 + 2	3.6	+38	
(54)	1 + 2 = 0 + 3	2.3	+268	
(55)	1 + 3 = 0 + 4	1.3	+494	
(56)	1 + 4 = 0 + 5	0.75	+720	
(57)	1 + 5 = 0 + 6	0.33	+947	
(58)	1 + 6 = 0 + 7	0.2	+1179	
(59)	1 + 1 = 2 + 0	1.	+371	
(60)	1 + 2 = 2 + 1	5.2	+135	
(61)	2 + 2 = 1 + 3	9.	+56	
(62)	2 + 3 = 1 + 4	4.9	+322	
(63)	2 + 4 = 1 + 5	2.6	+548	
(64)	2 + 5 = 1 + 6	1.4	+755	
(65)	2 + 6 = 1 + 7	0.63	+1007	
(66)	2 + 1 = 3 + 0	0.8	+538	
(67)	2 + 2 = 3 + 1	4.1	+302	
(68)	2 + 3 = 3 + 2	14.	+72	
(69)	3 + 3 = 2 + 4	14.	+155	
(70)	3 + 4 = 2 + 5	7.5	+380	
(71)	3 + 5 = 2 + 6	3.6	+608	
(72)	3 + 6 = 2 + 7	1.8	+840	
(73)	3 + 1 = 4 + 0	0.58	+702	
(74)	3 + 2 = 4 + 1	2.9	+465	
(75)	3 + 3 = 4 + 2	10.	+235	
(76)	3 + 4 = 4 + 3	32.	+8	
(77)	4 + 4 = 3 + 5	19.	+217	
(78)	4 + 5 = 3 + 6	9.6	+445	
(79)	4 + 6 = 3 + 7	4.5	+677	
(80)	4 + 1 = 5 + 0	0.4	+861	
(81)	4 + 2 = 5 + 1	1.9	+625	
(82)	4 + 3 = 5 + 2	6.9	+395	
(83)	4 + 4 = 5 + 3	2.2	+168	
(84)	5 + 4 = 4 + 5	42.	+58	
(85)	5 + 5 = 4 + 6	22.	+285	
(86)	5 + 6 = 4 + 7	10.5	+517	
(87)	5 + 1 = 6 + 0	0.27	+1017	
(88)	5 + 2 = 6 + 1	1.4	+781	
(89)	5 + 3 = 6 + 2	4.7	+551	
(90)	5 + 4 = 6 + 3	14.9	+324	
(91)	5 + 5 = 6 + 4	45.	+99	
(92)	6 + 5 = 5 + 6	47.	+129	
(93)	6 + 6 = 5 + 7	21.	+361	
(94)	6 + 1 = 7 + 0	0.2	+1172	
(95)	6 + 2 = 7 + 1	0.8	+935	
(96)	6 + 3 = 7 + 2	3.1	+705	
(97)	6 + 4 = 7 + 3	9.5	+479	
(98)	6 + 5 = 7 + 4	28.	+253	
(99)	6 + 6 = 7 + 5	84.	+26	
(100)	7 + 6 = 6 + 7	46.5	+206

(101)	1 = 0	RK(101) = 8.2 x 10 ⁸ T + 1.2 x 10 ¹⁷ T ⁻² + 3961.	10
(102)	2 = 1	(102) = 3RK(101)	
(103)	3 = 2	(103) = 5.8RK(101)	
(104)	4 = 3	(104) = 9.7RK(101)	
(105)	5 = 4	(105) = 13.RK(101)	
(106)	6 = 5	(106) = 17.5RK(101)	
(107)	7 = 6	(107) = 22.RK(101)	

Reaction No.	Reaction	Rate Coefficients ^a (cm ³ /mole-sec and cm ⁶ /mole ² -sec)	ΔE_v (cm ⁻¹)	Note ^b
HF(v) + M₂ = HF(v-1) + M₂ M ₂ = all H ₂				
(108)	1 = 0	RK(108) = 1.61x10 ¹¹ xTx10 ^{-42.6/T^{1/2}}	+3961.	11
(109)	2 = 1	(109) = 3RK(108)	+3789.	
(110)	3 = 2	(110) = 5.8RK(108)	+3622.	
(111)	4 = 3	(111) = 9.7RK(108)	+3459.	
(112)	5 = 4	(112) = 13RK(108)	+3299.	
(113)	6 = 5	(113) = 17.5RK(108)	+3143.	
(114)	7 = 6	(114) = 22RK(108)	+2988.	
HF(v) + F = HF(v-1) + F				
(115)	1 = 0	RK(115) = 3.99x10 ¹⁰ xTx10 ^{-40.9/T^{1/2}}	+3961.	12
(116)	2 = 1	(116) = 3RK(115)	+3789.	
(117)	3 = 2	(117) = 5.8RK(115)	+3622.	
(118)	4 = 3	(118) = 9.7RK(115)	+3459.	
(119)	5 = 4	(119) = 13RK(115)	+3299.	
(120)	6 = 5	(120) = 17.5RK(115)	+3143.	
(121)	7 = 6	(121) = 22RK(115)	+2988.	
HF(v) + H = HF(v-1) + H				
(122)	1 = 0	RK(122) = 1.6 x 10 ¹³	+3961	13
(123)	2 = 1	(123) = 3.RK(122)	+3789.	
(124)	3 = 2	(124) = 4.RK(122)	+3622.	
(125)	4 = 3	(125) = 12.RK(122)	+3459.	
(126)	5 = 4	(126) = 20.RK(122)	+3299.	
(127)	6 = 5	(127) = 30.RK(122)	+3143.	
(128)	7 = 6	(128) = 50.RK(122)	+2988.	
HF(v) + F₂ = HF(v-1) + F₂				
(129)	1 = 0	RK(129) = 5.06x10 ¹⁰ xTx10 ^{-42.6/T^{1/2}}	+3961.	14
(130)	2 = 1	(130) = 3RK(129)	+3789.	
(131)	3 = 2	(131) = 5.8RK(129)	+3622.	
(132)	4 = 3	(132) = 9.7RK(129)	+3459.	
(133)	5 = 4	(133) = 13.RK(129)	+3299.	
(134)	6 = 5	(134) = 17.5RK(129)	+3143.	
(135)	7 = 6	(135) = 22.RK(129)	+2988.	
HF(v) + He = HF(v-1) + He				
(136)	1 = 0	RK(136) = 5.41x10 ¹¹ xTx10 ^{-43.9/T^{1/2}}	+3961.	15
(137)	2 = 1	(137) = 3RK(136)	+3789.	
(138)	3 = 2	(138) = 5.8RK(136)	+3622.	
(139)	4 = 3	(139) = 9.7RK(136)	+3459.	
(140)	5 = 4	(140) = 13RK(136)	+3299.	
(141)	6 = 5	(141) = 17.5RK(136)	+3143.	
(142)	7 = 6	(142) = 22RK(136)	+2988.	
H₂(v) + M₃ = H₂(v-1) + M₃ M ₃ = all but H ₂ and H				
(143)	1 = 0	RK(143) = 5.05x10 ¹⁰ xTx10 ^{-43.6/T^{1/2}}	+4160.	16
(144)	2 = 1	(144) = 2RK(143)	+3924.	
(145)	3 = 2	(145) = 3RK(143)	+3694.	
(146)	4 = 3	(146) = 4RK(143)	+3467.	
(147)	5 = 4	(147) = 5RK(143)	+3242.	
(148)	6 = 5	(148) = 6RK(143)	+3014.	
(149)	7 = 6	(149) = 7RK(143)	+2782.	
H₂(v) + M₄ = H₂(v-1) + M₄ M ₄ = H ₂ + H				
(150)	1 = 0	RK(150) = 2.10x10 ¹¹ xTx10 ^{-43.4/T^{1/2}}	+4160.	17
(151)	2 = 1	(151) = 2RK(150)	+3924.	
(152)	3 = 2	(152) = 3RK(150)	+3694.	
(153)	4 = 3	(153) = 4RK(150)	+3467.	
(154)	5 = 4	(154) = 5RK(150)	+3242.	
(155)	6 = 5	(155) = 6RK(150)	+3014.	
(156)	7 = 6	(156) = 7RK(150)	+2782.	

Reaction No.	Reaction	Rate Coefficients ^a (cm ³ /mole-sec and cm ⁶ /mole ² -sec)	ΔE_v (cm ⁻¹)	Note ^b
(157)	2F + He → F ₂ + He	RK(157) = 4.6 × 10 ¹²	0	18
(158)	2F + F ₂ → F ₂ + F ₂	(158) = 1.21 × 10 ¹³	0	
(159)	2F + M ₂ + F ₂ + M ₂ , M ₂ = all H ₂	(159) = 0	0	
(160)	2F + M ₁ + F ₂ + M ₁ , M ₁ = all HF	(160) = RK(161)	0	
(161)	2F + F → F ₂ + F	(161) = 6 × 10 ⁻³	0	
(162)	2F + H → F ₂ + H	(162) = 0	0	
(163)	2H + M ₆ → H ₂ (0) + M ₆ M ₆ → all except H	RK(163) = 2 × 10 ¹⁴	0	19
(164)	→ 1	(164) = RK(163)	-4160.	
(165)	→ 2	(165) = RK(163)	-8084.	
(166)	→ 3	(166) = RK(163)	-11,778.	
(167)	→ 4	(167) = RK(163)	-15,245.	
(168)	→ 5	(168) = RK(163)	-18,487.	
(169)	→ 6	(169) = RK(163)	-21,501.	
(170)	→ 7	(170) = RK(163)	-24,283.	
(171)	2H + H → H ₂ (0) + H	RK(171) = 10RK(163)	0	19
(172)	→ 1	(172) = 10RK(163)	-4160.	
(173)	→ 2	(173) = 10RK(163)	-8084.	
(174)	→ 3	(174) = 10RK(163)	-11,778.	
(175)	→ 4	(175) = 10RK(163)	-15,245.	
(176)	→ 5	(176) = 10RK(163)	-18,487.	
(177)	→ 6	(177) = 10RK(163)	-21,501.	
(178)	→ 7	(178) = 10RK(163)	-24,283.	
(179)	H + F + M ₅ → HF(0) + M ₅	RK(179) = 1.25 × 10 ¹⁴	0	20
(180)	→ 1	(180) = 1.25 × 10 ¹⁴	-3962.	
(181)	→ 2 M ₅ = all species	(181) = 1.25 × 10 ¹⁴	-7751.	
(182)	→ 3	(182) = 1.25 × 10 ¹⁴	-11,373.	
(183)	→ 4	(183) = 1.25 × 10 ¹⁴	-14,831.	
(184)	→ 5	(184) = 1.25 × 10 ¹⁴	-18,131.	
(185)	→ 6	(185) = 1.25 × 10 ¹⁴	-21,274.	
(186)	→ 7	(186) = 1.25 × 10 ¹⁴	-24,262.	
(187)	HF(4) + F ₂ → HF(0) + 2F	RK(187) = 7 × 10 ⁶	+14,832.	21
(188)	5 → 0	(188) = 0	+18,131.	
(189)	6 → 0	(189) = 0	+21,274.	
(190)	7 → 0	(190) = 0	+24,263.	
(191)	H ₂ (0) + F ₂ → HF(0) + H + F	RK(191) = 0	0	21
(192)	1 → 0	(192) = 10 ⁵	+4160.	
(193)	2 → 0	(193) = 0	+8084.	
(194)	3 → 0	(194) = 0	+11,778.	
(195)	4 → 0	(195) = 0	+15,245.	

^a RK indicates values used in the kinetics model.

^b These Notes are discussed immediately following Table II.

coefficients, most of which have not been measured experimentally. The accuracy of the model depends on the reactions chosen and the accuracy of the values of the rate coefficients. It is obvious that the values for most of the rate coefficients are uncertain, but this report may have some value in that the detailed descriptions of our assumptions and methods may be useful as a basis for further work. Despite the amount of experimental and theoretical (trajectory calculation) work that has been completed, the determination of rate coefficients is unfortunately necessarily made from work that is not directly applicable. We have tried to cite all references that bear directly on the reactions used, but for the large number of references on the experimental determinations of the V-R,T relaxation of HF(v) by HF and by F we refer to the recent paper of Blair, Breshears, and Schott.⁷ All rate coefficients and relaxation times are converted to cm³/mole-sec or cm⁶/mole²-sec; the symbol "RK" is used only for the values that we use in the kinetics model.

Reverse reactions are included for all reactions except (23)-(30) and (157)-(178), and the rate constants are given by

$$RK_r(i) = RK_f(i) \times K(i), \quad (1)$$

where K(i) is an equilibrium constant calculated from JANAF tables. Equilibrium constants pertain to systems in which reactants and products are in complete thermodynamic equilibrium, but the reactions considered here involve H₂ and HF in individual vibrational levels. However, in the 1000-to-3000°K temperature range of interest here (and at lower temperatures as well), equilibrium distributions of H₂ or HF contain less than 10% of the molecules in vibrational levels v > 0. For reactions where H₂ or HF, or both, are in the 0th vibrational levels, the reverse rate constant is then given with little error by Eq. (1); see Note 4. For these cases (H₂ or HF in 0th vibrational levels) the equilibrium constant may be written

$$K(i) = 10^{-\Delta G/2.302RT},$$

where $\Delta G = \Delta G_f(\text{reactants})$ minus $\Delta G_f(\text{products})$. For

reactions involving H₂ or HF in higher vibrational states, an additional free-energy term is necessary so that

$$K(i) = 10^{-(\Delta G + \Delta E_v)/2.302RT},$$

where ΔE_v = vibrational energy in reactants minus vibrational energy in products. A value of ΔE_v for each reaction is given in Table I. The values of ΔG are only weakly dependent on temperature, and we have taken ΔG to be independent of temperature and equal to the value of 2000°K for all reactions except (157)-(162). The values of ΔG are

	Reaction No.	ΔG (cm/sec)
ΔG_1	(1)-(22)	10,700
ΔG_2	(23)-(30)	36,050
ΔG_3	(157)-(162)	12,240 (1-1.25x10 ⁻⁴ T)
ΔG_4	(163)-(170)	34,550
ΔG_5	(179)-(186)	45,000
ΔG_6	(187)-(190)	- ΔG_3
ΔG_7	(191)-(195)	$\Delta G_5 - \Delta G_4 - \Delta G_3$

The value of R in cm²/deg is 0.6950.

Table II gives the energy levels that determine the values of ΔE_v . Levels for H₂ were obtained from Wooley, Scott, and Brickwedde,⁸ and for HF from Mann et al.⁹

TABLE II
VIBRATIONAL ENERGY LEVELS FOR HF AND H₂

v	H ₂ (cm ⁻¹)	HF (cm ⁻¹)
0	0	0
1	4160.4	3961.6
2	8084.4	7751.0
3	11778.3	11373.0
4	15245.7	14831.9
5	18487.4	18131.2
6	21501.6	21274.0
7	24283.7	24262.6

NOTES

1. The rate of the thermal reaction

is taken to be the mean of the two values

$$k = 1.6 \times 10^{14} \cdot 10^{-349/T} \quad (\text{Ref. 10})$$

$$k = 0.59 \times 10^{14} \cdot 10^{-349/T} \quad (\text{Ref. 11}) .$$

The latter was obtained from a single value near room temperature combined with the temperature dependence of Ref. 10. Then for a thermal distribution of F atoms and H₂(O) molecules

$$k_a = 1.1 \times 10^{14} \cdot 10^{-349/T} \text{ cm}^3/\text{mole-sec.} \quad (2)$$

In an H₂-F₂ explosion however, the species do not have a Maxwell distribution over translation energy. The H₂(O) molecules reacting in Reaction (a) probably have a Maxwell distribution, but the F atoms that are formed by a photolytic flash, by irradiation with electrons, or by chemical reaction are initially hot (having translational energy sufficiently high that the energy that can enter into excitation in an F + H₂ collision is greater than the activation energy, 1.6 kcal). Since the fraction of translational energy that can enter into excitation is $M_{\text{H}_2}/(M_{\text{H}_2} + M_{\text{F}}) = 0.095$, the initial energy, if the atom is hot, must be ≥ 16.8 kcal. It can be shown from the kinetic theory of hot atoms¹² that in a 1:1:H₂:F₂ mixture about 20% of the F atoms from a photolytic flash (absorption centered at about 2800 Å to produce F atoms with initial translational energies of 32 kcal/mole) collide with H₂ while hot.

The major part of the F atoms entering into reactions (1)-(4) is produced by reactions (23)-(30). The relative reaction rates in (23)-(30) and the distribution of initial [as formed in (23)-(30)] translational energies of F atoms were determined experimentally near room temperature for a thermal distribution of H atoms,¹³ and are given in Table III. Although not stated in Ref. 13, the reaction is primarily with F₂ in the 0th vibrational state (see Note 6). Table III also shows the distributions

TABLE III

RELATIVE RATES $k(\text{v})$ OF PRODUCTION OF HF(v) AND THE TRANSLATIONAL ENERGY OF F ATOMS $E_T(\text{F})$ FROM THE THERMAL REACTION $\text{H} + \text{F}_2 \rightarrow \text{HF}(\text{v}) + \text{F}$ AT ROOM TEMPERATURE

v	$k(\text{v})$ (Ref. 13)	$E_T(\text{F})$ (Ref. 13) (kcal/mole)	$k(\text{v})$ (Ref. 14)	$k(\text{v})$ (Ref. 15)
0	<0.1	—	0.04	
1	0.12	33	0.09	
2	0.13	38	0.11	
3	0.25	32	0.13	0.4
4	0.35	28	0.43	0.9
5	0.78	23	0.89	1.85
6	1.00	19	1.00	1.00
7	0.40	15	0.43	
8	0.26	12	0.20	
9	<0.16	—	<0.04	

$k(\text{v})$ obtained experimentally by Jonathan et al.¹⁴ and theoretically by Wilkins.¹⁵

The translational energy in products can also be obtained from the work of Blais,¹⁶ who determined theoretically the vibrational distributions for HF(v) from $\text{H} + \text{F}_2(\text{O}) \rightarrow \text{HF}(\text{v}) + \text{F}$ at fixed relative velocities $V_R = 4.57 \times 10^5$, 5.97×10^5 , 7.09×10^5 , 8.20×10^5 , and 1.05×10^6 cm/sec. These velocities correspond to the most effective relative velocities for reaction of Maxwell distributions at roughly 400, 1000, 1500, 1800, and 2500°K. The translational energies in the F atoms, as shown in Table IV, were obtained by subtracting the energy in vibration and rotation in the products from the total energy available and then dividing by two. The energy in rotation in the products was assumed to be independent of the relative velocity of the reactants and equal to 3 kcal/mole.¹³ The energy available to products was taken to be $\Delta E_0^\circ + \frac{1}{2} m V_R^2 + RT$, where RT was taken as 2 kcal/mole. Our assumptions about the amount of energy in rotation have a negligible effect on the distribution of the translational energies of the F atoms.

Although the distributions $k(\text{v})$ from Blais' theoretical work¹⁶ for $V_R = 4.6 \times 10^5$ cm/sec and from Polanyi's experimental work¹³ at 300°K are in only moderate agreement, the average values of $E_T(\text{F})$ are in good agreement. Blais' work shows that an appreciable fraction of translational energy in the H atom is carried over into translational energy of the F atom and that many distributions of translational energies are present.

TABLE IV

RELATIVE RATES $k(v)$ OF PRODUCTION OF $\text{HF}(v)$ AND THE TRANSLATIONAL ENERGY OF F ATOMS $E_T(F)$ FROM THE REACTION $\text{H} + \text{F}_2 \rightarrow \text{HF}(v) + \text{F}$
(Data taken from Figs. 5 through 9 of Ref. 15)

		$10^4 v_R$					
		4.6	5.97	8.20	10.5		
v	$k(v)$	$E_T(F)$		$E_T(F)$		$E_T(F)$	
		(kcal/mole)	$k(v)$	(kcal/mole)	$k(v)$	(kcal/mole)	$k(v)$
0	0	50	0	50.5	0	52.5	0
1	0	44	0	45	0	47	0
2	0	39	0	40	0	41.5	0.055
3	0.006	34	0.02	35	0.075	36.5	0.177
4	0.199	29	0.21	30	0.25	30.5	0.220
5	0.265	24	0.31	25	0.375	26.5	0.309
6	0.331	19	0.36	20	0.225	22	0.166
7	0.199	16	0.10	16	0.075	17	0.072

However, the hot F atoms have little effect on the rate coefficient for $\text{F} + \text{H}_2(0) \rightarrow \text{HF}(v) + \text{H}$. Blais' work¹⁷ on this reaction indicates that the cross section increases only 40% as the relative translational energy increases from 1 to 8 kcal/mole. We increase the rate coefficient by 20% (well within the experimental error) and use

$$\text{RK}(1) + \text{RK}(2) + \text{RK}(3) + \text{RK}(4) = 1.3 \times 10^{14} \cdot 10^{-34v}/T.$$

A negligible number of super-hot atoms is formed by such reactions as

etc.

The temperature T is computed from the instantaneous heat release of all the reactions and from the sum of the instantaneous rotational and translational heat capacities. The rotational and translational temperatures are assumed to be equal. However, the rate of the self-rotational relaxation of H_2 is slow.¹⁸ A rough extrapolation of the rates reported¹⁸ for 77 to 170°K to temperatures of interest here (~1000°K) would indicate that the number of H_2 - H_2 collisions is about 100 for relaxation of the excess rotational energy by a factor of e^{-1} . The rate of rotational relaxation of H_2 by HF or F_2 would not be expected to be greater. The rotational temperature of H_2 may therefore lag the translational temperature by about 100 collisions (~10⁻⁸ sec in a

system at 1 atm). If the activation energy for (1)-(4) lies in rotation, then T should properly be a rotational temperature, and our computed values of T at any instant are too high. If the activation energy lies in translation, then T should refer to a translational temperature, and our computed values of T at any instant are too low.

2. The relative rates of production of $\text{HF}(v)$ from the reaction of thermal F atoms,

as determined by various workers, are given in Table V.

TABLE V
RELATIVE RATES OF PRODUCTION OF $\text{HF}(v)$ FROM $\text{F} + \text{H}_2(0) \rightarrow \text{HF}(v) + \text{H}$

v	Ref. 19 ^a	Ref. 20 ^a	Ref. 21 ^a	Ref. 17 ^{b,c}	Ref. 22 ^{b,d}	Ref. 23 ^b	Ref. 24 ^{b,e}
0	0	0	0	0	0	0	0
1	0.31	0.29	0.18	0.39	0.14	0.53	0.21
2	1.00	1.00	1.00	1.00	1.00	1.00	1.00
3	0.47	0.76	1.33	1.66	0.58	1.07	0.49

^aExperimental values at room temperature.

^bTheoretical values.

^cAverage ratios calculated from the ratios obtained at fixed relative translational energies $E_T = 1.0, 2.0, 4.0, 8.0$ kcal/mole. The ratios of cross sections, $\sigma(3)/\sigma(2)$, at these relative energies are 1.54, 1.61, 1.72, 1.82, respectively. The energies correspond to average energies for Maxwell distributions at 330, 660, 1300, and 2600°K. A different method of assigning HF to vibrational levels gives $\sigma(3)/\sigma(2) \approx 1.0$ independent of the relative energy.

^dCalculated for relative velocity, $v_R = 4 \times 10^3$ cm/sec, $J = 3$.

^eThermal rate constants are given for $k(v)$ in the range 100 to 1000°K. The ratio $k(3)/k(2)$ is independent of temperature; $k(1)/k(2) = 0.21$ at 1000°K, and

$$k(1) = 10^{13.000} 10^{-34v}/T$$

$$k(2) = 10^{13.000} 10^{-34v}/T$$

$$k(3) = 10^{13.000} 10^{-34v}/T$$

The results of Blais¹⁷ and Wilkins²⁴ indicate only a slight temperature dependence for $k(3)/k(2)$; the experimental ratios^{19,20} are in good agreement with two of the theoretical values.^{22,24} Coombe and Pimentel²⁵ have recently shown experimentally that there is a small temperature dependence for $k(3)/k(2)$ and $k(1)/k(2)$ in the range 200 to 400°K: $k(3)/k(2) = 0.39 \exp(+117/RT)$, $k(1)/k(2) = 0.47 \exp(-254/RT)$. We use $k(3)/k(2) = 0.60$ and $k(1)/k(2) = 0.25$, and consider both ratios to be independent of temperature. We use the experimental

temperature dependence¹⁰ for the overall rate -- and not the theoretical temperature dependence, as would be given by the weighted sum of $k(v)$ of Ref 24. The rate constants are given in Table VI.

TABLE VI
RATE CONSTANTS FOR REACTIONS (1)-(4),
 $F + H_2(0) \rightarrow HF(v) + H$

Reaction No.	v	$k(v)/k(2)$	RK^a
(1)	0	0	0
(2)	1	0.25	$1.75 \times 10^{13} \times 10^{-349}/T$
(3)	2	1.00	4 RK(2)
(4)	3	0.6	2.4 RK(2)

^aRK indicates values used in the kinetics model.

About 15 to 20% of the F atoms react while hot, and we consider their effect on the relative rates to be small

3. There are no experimental data for the rates of $F + H_2(v > 0) \rightarrow HF(v') + H$. The rates for $v = 0, 1, 2, 3$, as given in Table VII, were taken from Blais and Truhlar²² and were determined for a fixed relative translational velocity $V_R = 4 \times 10^5$ cm/sec and $J = 3$.

TABLE VII

VIBRATIONAL ENERGY DISTRIBUTIONS AND REACTION CROSS SECTIONS FOR
 $F + H_2(v) \rightarrow HF(v') + H$ AT $V_R = 4 \times 10^5$ cm/sec, $J = 3$

$v \backslash v'$	Fraction of HF with given v'								$\sigma_{R,A}^0$
	0	1	2	3	4	5	6	7	
0	0	0.08	0.58	0.34	0	0	0	0	2.7
1	0.03	0.08	0.12	0.26	0.47	0.04	0	0	7.9
2	0	0	0.08	0.13	0.27	0.47	0.05	0	12.7
3	0.02	0.02	0.02	0.08	0.14	0.22	0.39	0.11	22.3

Rate constants for $F + H_2(v)$ were calculated from the cross sections in Table VII by using $k = V_R \cdot 10^{16} \sigma_R \cdot N_A$, where N_A is Avogadro's number. The results are given in Table VIII. The value of $k(0)$ corresponds to the rate at $T \approx 1200^\circ K$ as given by Eq. (2). The rates $k(v)$ for $v = 1, 2, 3$, as given in Table VIII, are then obtained relative to

TABLE VIII

RELATIVE RATES FOR $F + H_2(v) \rightarrow HF + H$

v	σ_R ($\text{\AA}^2$)	$k(v)$ ($\text{cm}^3/\text{mole-sec}$)
0	2.7	0.65×10^{14}
1	7.9	1.90×10^{14}
2	12.7	3.05×10^{14}
3	22.3	5.36×10^{14}

the experimental rate at $1200^\circ K$ and are taken to be independent of temperature. Although the rates $k(v)$ and the relative rates for production of $HF(v')$ (as shown in Table VII) are given for fixed V_R and J , we assume these values in our calculations. Where the rate of production of $HF(v')$ (Table VII) is very low, we have added that rate to the rate for the neighboring $(v'-1)$ or $(v'+1)$ state to obtain the fractional contributions given in Table IX. The total rate for $v = 4$ as given in Table IX was taken to be equal to the total rate for $v = 3$, and the relative rates for production of $HF(v')$ were estimated from the relative rates (Table VII) in $k(2)$ and $k(3)$.

TABLE IX

RATE CONSTANTS FOR REACTIONS (5)-(22), $F + H_2(v) \rightarrow HF(v') + H$

Reaction No.	v	v'	Fraction of HF with v'	RK^a ($\text{cm}^3/\text{mole-sec}$)
(5)	1	1	0.11	2.1×10^{12}
(6)	1	2	0.12	1.09 RK(5)
(7)	1	3	0.26	2.36 RK(5)
(8)	1	4	0.51	4.63 RK(5)
(9)	2	2	0.08	2.44×10^{13}
(10)	2	3	0.13	1.62 RK(9)
(11)	2	4	0.27	3.37 RK(9)
(12)	2	5	0.52	6.5 RK(9)
(13)	3	3	0.14	7.5×10^{13}
(14)	3	4	0.14	RK(13)
(15)	3	5	0.22	1.57 RK(13)
(16)	3	6	0.39	2.78 RK(13)
(17)	3	7	0.11	0.78 RK(13)
(18)	4	3	0.02	1.07×10^{13}
(19)	4	4	0.08	4. RK(18)
(20)	4	5	0.14	7. RK(18)
(21)	4	6	0.26	13. RK(18)
(22)	4	7	0.50	25. RK(18)

^aRK indicates values used in the kinetics model.

The $H_2(v)$ molecules that react in (5)-(22) are formed by V-V transfers from $HF(v')$ or by reactions (-5)-(-22), and initially have a nonthermal distribution over rotational states. The rate of self-rotational relaxation of H_2 is low.¹⁸ As stated in Note 1, a rough extrapolation of the rates reported¹⁸ for 77 to 170°K to temperatures of interest here (~1000°K) would indicate that the number of H_2 - H_2 collisions is about 100 for relaxation of the excess rotational energy by a factor of e^{-1} . An appreciable fraction of the $H_2(v)$ molecules then collides with F atoms [$(F)/(H_2) \approx 0.01$ in an explosion] while still rotationally excited. Our rate constants for (5)-(22) were determined for $J = 3$ and are good approximations to the rates in an explosion only if the cross sections are not dependent on J . Muckerman's²⁶ trajectory calculations show a decrease of about 75% in cross section for $F + H_2(0)$ as J increases from 1 to 3. Later work by Blais and Truhlar,²⁷ however, shows little dependence of the cross section for $0 \leq J \leq 6$, and in view of current data we do not correct our rate coefficients for rotational excitation.

The importance of reactions (5)-(22) in carrying the chemical reaction can be estimated from the relative rates given in Table X. The relative concentrations of $H_2(v)$ were taken from the results of a computer-simulated explosion of H_2 - F_2 . The initial conditions for the computation were 300 torr F_2 , 300 torr H_2 , 40 torr O_2 , 300°K, 1% of the F_2 dissociated in a 4.8-nsec pulse; the rate constants used in these computations were different from the rates proposed here, but the relative rates in Table X are probably not far out of line. The values in Table X indicate that at 1000 and 2400°K the contributions of $F + H_2(v > 0)$ to the total $HF(v')$ produced by $F + H_2(v) \rightarrow HF(v') + H$ are 13 and 30%, respectively.

TABLE X

RELATIVE CONCENTRATIONS $H_2(v)/H_2(0)$ AND RELATIVE RATES $k(v) \cdot H_2(v)/k(0) \cdot H_2(0)$ AT 1000 AND 2400°K IN A SIMULATED EXPLOSION

v	1000°K		2400°K	
	$H_2(v)/H_2(0)$	$k(v) \cdot H_2(v)/k(0) \cdot H_2(0)$	$H_2(v)/H_2(0)$	$k(v) \cdot H_2(v)/k(0) \cdot H_2(0)$
0	1	1	1	1
1	0.012	0.036	0.05	0.15
2	0.002	0.01	0.016	0.08
3	0.0115	0.094	0.017	0.14
4	0.0016	0.013	0.007	0.06

4. From detailed balancing, the rates of the reverse reactions (-2)-(-22) may be obtained in terms of the rate of (-1). If $\epsilon_1, \epsilon_2, \epsilon_3, \dots$ and $h_1, h_2, h_3, \dots$ are energy differences between successive vibrational levels of HF and H_2 , respectively, then

$$k_{-2} = k_2 \frac{k_{-1}}{k_1} \exp(\epsilon_1/RT)$$

$$k_{-3} = k_3 \frac{k_{-1}}{k_1} \exp(\epsilon_2/RT)$$

.

$$k_{-5} = k_5 \frac{k_{-1}}{k_1} \exp(\epsilon_1 - h_1)/RT, \text{ etc.}$$

.

The ratio k_1/k_{-1} and the thermodynamic equilibrium constant K are defined by

$$\frac{k_1}{k_{-1}} = \frac{(HF(0)) \cdot (H)}{(H_2(0)) \cdot (F)}$$

$$K = \frac{[(HF(0)) + (HF(1)) + \dots (HF(n))] \cdot (H)}{[(H_2(0)) + (H_2(1)) + \dots (H_2(n))] \cdot (F)}$$

where the concentrations are those at thermodynamic equilibrium. The relative equilibrium concentrations $(HF(1))/(HF(0))$ and $(H_2(1))/(H_2(0))$ at 1000, 2000, and 3000°K are 0.003, 0.06, 0.15 and 0.0025, 0.05, 0.136, respectively, so that, with little error, the ratio k_1/k_{-1} may be replaced by K . Then

$$k_{-2} = k_2 \cdot K \cdot \exp(\epsilon_1/RT), \text{ etc.} \quad (3)$$

The equilibrium constant is a function of temperature $K(T) = 10^{-\Delta G/RT}$, and we take ΔG to be independent of temperature and equal to 10,600 cm^{-1} . The additional free energies that contribute to each reaction, $\epsilon_1, \epsilon_2, \epsilon_3 - h_1$, etc., are given as ΔE_v in Table I. Only some of the reverse reactions in (-2)-(-22) are sufficiently near thermoneutrality to have significantly large rate coefficients; the values of these at 1000 and 2000°K (Table XI) were calculated to determine if these rates compete significantly with other reactions.

TABLE XI
VALUES OF RATE COEFFICIENTS FOR SOME REACTIONS OF NEAR-THERMAL
DISTRIBUTIONS OF H ATOMS, $H + HF(v) \rightarrow H_2(v') + F$,
AS CALCULATED FROM THE FORWARD REACTIONS

v	v'	Reaction No.	RK ^a	
			1000°K	2000°K
2	0	(-3)	4.5×10^{11}	5.6×10^{13}
3	0	(-4)	4.2×10^{13}	4.2×10^{13}
3	1	(-7)	3.3×10^{11}	4.1×10^{13}
4	1	(-8)	9.7×10^{13}	9.7×10^{13}
4	2	(-11)	2.76×10^{11}	4.8×10^{13}
5	2	(-12)	6.2×10^{13}	1×10^{14}
6	3	(-16)	3.67×10^{13}	8.7×10^{13}
7	3	(-17)	7.5×10^{13}	2.1×10^{14}
7	4	(-22)	2.3×10^{13}	7.9×10^{13}

^aRK indicates values used in the kinetics model.

These rates are calculated from forward rates in which there is little hot-atom effect; the values in Table XI are therefore determined for a near-thermal distribution of H atoms, but in the explosive system a large fraction is very hot (see Note 6).

Wilkins²⁸ calculated rate coefficients for thermal, translational, and rotational distributions in the reaction $H + HF(v) \rightarrow H_2(v') + F$ in the temperature range 100 to 1000°K. Rate coefficients as estimated from his Fig. 1 are given in Table XII.

TABLE XII
RATE COEFFICIENTS FOR $H + HF(v) \rightarrow H_2(v') + F$

v	v'	Ref. 28		Present Work	
		1000°K	2000°K	1000°K	2000°K
		(10 ¹² cm ³ /mole-sec)		(10 ¹² cm ³ /mole-sec)	
6	0	3	3.5		
5	1	4	5		
6	2	9	10		
5	0	3	4		
5	1	6	8		
4	0	4	4		
4	1	3	4	97.	97.
3	0	5.5	7	42.	42.

Two significant differences between our values and those of Wilkins are shown in Tables XI and XII:

(1) our values are a factor of 7 to 30 larger than those of Wilkins, and (2) our rate coefficients increase markedly with the exothermicity of the reaction, and Wilkins' do not. We have obtained preliminary values of the rate coefficients for

(-2)-(-22) by trajectory calculations; these values are lower than the values in Table XI by factors of 5 to 10.

Our rate coefficients for (-2)-(-22) as calculated from Eqs. (3) are for reactions of thermal distributions of HF over J, but the HF molecules, as initially formed in reactions (2)-(30), have nonthermal distributions^{13,16,17,19,24} over J with rotational excitation as high as J = 20. There are no experimental data for the rates of rotational relaxation of HF. Polanyi and Woodall,²⁹ however, modeled HCl chemiluminescence experiments to obtain upper limits for transition probabilities $P_{J+\Delta J}^J$ as a function of J and ΔJ for both upward and downward transitions in HCl. For $\Delta J = 1$, over the range $1 < J < 20$, which accounts for an appreciable fraction of the downward transitions, $P_{J+\Delta J}^J$ for downward transitions varies between 0.015 and 0.07. If we assume the same upper limits for probabilities for transitions in HF-H₂ collisions, then [with (H)/H₂] ~ 0.01 in a typical explosion initiated by a burst of F atoms] an appreciable fraction of the HF molecules collides with H atoms while still rotationally excited. Rotational self-relaxation of HCl takes place in about seven collisions.³⁰ If we assume that rotational self-relaxation of HF is equally rapid, then [with (HF)/(H) ~ 30 at 1000°K in a typical explosion] an appreciable fraction of excited HF still collides with H atoms. Relaxation in HF-F₂ collisions would be expected to be slower than that in HF-H₂ collisions. We conclude that rotationally excited HF molecules are colliding with H atoms and that our rate coefficients may have an added uncertainty if the rates of (-2)-(-22) depend appreciably on rotational excitation.

The following reactions are not included in the model because they are endothermic:

As pointed out by Cohen,⁶ the reverse reactions, however, are exothermic and occur to some extent. But, as indicated by Wilkins' results in Table XII,

reactions with large quantum jumps are not as important as those with small quantum jumps. And the rates are probably more uncertain.

5. It is of interest to consider the relative importance of reactions (-2)-(-22) in competing for H atoms and in degrading vibrational energy in HF(v).

The rate coefficients for $H + F_2 \rightarrow HF + F$, as calculated from Eq. (4), see Note 6, are 8.6×10^{13} and 1.65×10^{14} cm³/mole-sec at 1000 and 2000°K. From a computation of a simulated explosion (initial conditions given previously), the ratios HF(3)/F₂, HF(4)/F₂, HF(5)/F₂, HF(6)/F₂, and HF(7)/F₂ at 1000 and 2000°K are 0.033, 0.025, 0.019, 0.012, 0.0055, and 0.09, 0.07, 0.05, 0.038, 0.027, respectively. These figures, when taken with the rate coefficients in Table XI indicate that in an H₂-F₂ explosion at 1000°K less than 5% of the H atoms react with HF(v), and at 2000°K less than 10% of the H atoms react with HF(v).

One of the most important reactions for V-R,T de-excitation of HF(v) is

For a given v, the reaction among (-2)-(-22) that competes with Reaction (b) in degrading vibrational energy is

The efficiency of Reaction (c) relative to that of Reaction (b) in degrading vibrational energy in an H₂-F₂ explosion was calculated by using

$$\frac{R(c)}{R(b)} = \frac{(v-v')k_c}{k_b} \frac{(H)}{(HF)}$$

for those Reactions (c) where k_c is large. The values of k_c were obtained from Table XI, the requisite values of k_b from the coefficients RK(101)-RK(107), and the concentrations (H) and (HF), where (HF) is the sum of concentrations of all vibrational levels of HF, were obtained from a computed simulated explosion (initial conditions given previously). The ratios (H)/(HF) at 1000 and 2000°K were 0.025 and 0.0113, respectively. The relative

efficiencies are given in Table XIII. If our rate coefficients for (-2)-(-22) are approximately correct, then the values in Table XIII indicate that some of these reactions are very important in degrading the vibrational energy. However, our rate coefficients in Table XI may overestimate the rates of (-2)-(-22), and these rates would not be appreciable if we used lower values of the rate coefficients from the Wilkins results shown in Table XII, or the values obtained from our trajectory calculations.

TABLE XIII

RELATIVE EFFICIENCIES R(c)/R(b) FOR DEGRADATION OF VIBRATIONAL ENERGY BY REACTIONS
 $HF(v) + HF \rightarrow HF(v-1) + HF$ (b) AND $H + HF(v) \rightarrow H_2(v') + F$ (c) IN AN H₂-F₂ EXPLOSION

R(c)/R(b)			
v	v'	1000°K	2000°K
2	0	0.01	0.02
3	0	0.6	0.14
4	1	0.8	0.2
5	2	0.4	0.15
6	3	0.2	0.1

6. The rate of $H + F_2 \rightarrow HF + F$ for thermal H atoms was taken from

$$k = 1.2 \times 10^{14} 10^{-525/T} \quad (\text{Ref. 31})$$

$$k = 2.0 \times 10^{14} 10^{-570/T} \quad (\text{Ref. 11}).$$

We use the temperature dependence of Rabideau, Hecht, and Lewis¹¹ and take the rate coefficient for thermal H atoms to be

$$k = 1.6 \times 10^{14} 10^{-570/T}.$$

The fact that the activation energy (2600 kcal = 912 cm⁻¹) is about equal to the difference in vibrational levels of F₂ (892 cm⁻¹) suggests that the rate can be attributed solely to

However, trajectory calculations have shown that

has an appreciable cross section compared to that for $H + F_2(1)$.^{14,16} The major part of the reaction near room temperature is carried by $F_2(0)$ because 98.6% of the F_2 molecules are in the 0th vibrational state. The distribution of $HF(v')$ over v' depends strongly on v in $F_2(v)$,¹⁶ and thus would change markedly with temperature as the reaction progressed if the F_2 relaxed vibrationally. However, the rate of relaxation of F_2 into upper vibrational states is slow compared to the rate of reaction — as shown by Table XIV. The rate at which the translational temperature changes, as given in Table XIV, was taken from a computation in which the initial conditions were such as to produce a rather low rate, i.e., 0.1% F_2 dissociated over a 144-nsec interval.

The rate of transfer of vibrational energy into F_2 by V-V processes can also be shown to be too slow compared to the rate of the overall reaction. An upper limit, $k < 2 \times 10^9$ cm³/mole-sec, was obtained experimentally³⁴ for the rate coefficient for either or both of the processes:

The fractional rate per microsecond is

$$\begin{aligned} + \frac{dF_2(v)}{F_2(0)dt} &< 2 \times 10^9 \cdot HF(v) \cdot 10^{-6} \\ &< 2 \times 10^3 \cdot HF(v). \end{aligned}$$

In a system with initial conditions 300 torr H_2 , 300 torr F_2 , the sum of concentrations of $HF(v)$ for $v \geq 1$ does not exceed 5×10^{-6} before lasing ceases (computation of simulated explosion). The fractional increase in $F_2(v)$ is less than 0.01/ μ sec, and this rate is far exceeded by the rate of temperature increase. The rate of V-V transfer by

might also be expected to be slow.

We take the $F_2(0):F_2(1)$ ratio to be unchanged at its 300°K value during the reaction, and the production of $HF(v')$ to be due to the reaction

TABLE XIV

VIBRATIONAL RELAXATION TIMES OF F_2 COMPARED TO RATES OF REACTION IN AN H_2-F_2 EXPLOSION

(Values for dT/dt were taken from a computation of a simulated explosion, and the initial conditions for the computation were 300 torr H_2 , 300 torr F_2 , 0.1% F_2 dissociated in a 144-nanosecond pulse)

Temp. (°K)	$\tau P(F_2-F_2)$ (Refs. 32,33) (μ sec-atm)	$dT(\text{translational})/dt$ (from simulated explosion) (deg/ μ sec)
300	11.27	1000 (at 400°K)
1000	0.71	6000
2000	0.204	~20,000
3000	0.110	~20,000

The relative rates of production of $HF(v)$ from $H + F_2 \rightarrow HF(v) + F$ are given in Table III for a reaction of thermal H atoms near room temperature. These results and the above equation for the rate coefficient for $H + F_2$ as a function of temperature are not applicable to an exploding H_2-F_2 mixture, and are given only for orientation, comparison, and for possible use in heavily moderated systems.

The applicable rates and the distributions of $HF(v)$ over v are obtained from a consideration of the translational energies of H actually present in an exploding system.

The H atoms in reactions (23)-(30) are formed in reactions (2)-(22) with high translational energy, and a fraction of them react while hot (defined as having translational energy greater than $E_a = 2600$ kcal). The distribution of initial translational energies of the H atoms depends on the relative rates in the exploding system of the reactions

The relative rate coefficients, as obtained for a fixed relative velocity equal to the arithmetical average velocity for a Maxwell distribution at 1200°K, are obtained from the rate coefficients in Table VIII and are given in Table XV. The relative concentrations of $H_2(v)$, obtained near the peak of lasing in computer simulated explosions, are also given in Table XV. From these the relative rates are calculated.

TABLE XV

RELATIVE RATES OF $F + H_2(v)$

[$k(v)/k(0)$ from Ref. 22, $H_2(0)/H_2(v)$ at $T = 1860^\circ K$ from a computed simulated explosion with initial conditions 300 torr H_2 , 300 torr F_2 , 1% F_2 dissociated in 4.8 nsec]

v	$k(v)/k(0)$	$H_2(v)/H_2(0)$	Rate $k_v H_2(v) /$ Rate $k_0 H_2(0)$
0	1.00	1.00	1.00
1	2.9	0.033	0.096
2	4.7	0.01	0.047
3	8.3	0.016	0.133

The values in Table XV were obtained for a fixed relative translational velocity (corresponding to the arithmetical average velocity of a Maxwell distribution at $1200^\circ K$) and for a ratio $H_2(v)/H_2(0)$ that is present at the peak of lasing at $1860^\circ K$. However, we assume that the relative rates in Table XV apply to translational distributions of F and $H_2(v)$ present during most of the lasing period in an exploding H_2-F_2 system. When the relative rates of production of $HF(v')$ from a given $H_2(v)$ (as shown in Table VII) are multiplied by the relative rates of reaction of each v , the relative rates of production of $HF(v')$ are obtained for all

and are given in Table XVI.

TABLE XVI

RELATIVE RATES OF PRODUCTION OF $HF(v')$ FROM
 $H_2(v) + F \rightarrow HF(v') + H$ IN A LASING SYSTEM

v/v'	0	1	2	3	4	5	6	7
0	0	0.08	0.58	0.34	0	0	0	0
1	0.003	0.007	0.011	0.025	0.045	0.004	0	0
2	0	0	0.004	0.006	0.012	0.022	0.002	0
3	0.002	0.002	0.002	0.01	0.018	0.029	0.05	0.014

The values in Table XVI indicate that about 21% of the H atoms are formed from

and the distributions of $HF(v')$ are shifted to higher v' when compared to the distribution from

The shift in v' is almost proportional to the increase in v of $H_2(v)$; if there is no strong dependence on v of the amount of rotational energy in the products, the distribution of translational energies of the H atoms from the $H_2(v > 0)$ reactions is then not greatly different from that from the $H_2(0)$ reactions. To obtain the initial translational energies of the H atoms we then consider only $F + H_2(0)$ reactions and use the relative rates as given in Table VI. The translational energies of the H atoms, as given in Table XVII, are average values obtained from Fig. 10 of Ref. 19, and are for a room-temperature experiment. The extra 2 or 3 kcal/mole that might arise from translational energy of the reactants at higher temperatures are neglected.

TABLE XVII

AVERAGE INITIAL TRANSLATIONAL ENERGIES (AITE) OF
H ATOMS IN AN H_2-F_2 EXPLOSION
AT ROOM TEMPERATURE

v	$k(v)/k(2)$	AITE of H Atoms (kcal/mole)
0	0	--
1	0.25	20
2	1.00	10
3	0.6	2

We assumed that the atoms with 2 kcal/mole energy H_{th} react with a thermal rate, and we calculated rate constants for those H atoms, H^* , that had initial energies of 10 kcal/mole, and for those, H^{**} , that had initial energies of 20 kcal/mole. We assumed H^* and H^{**} to be separate species, each with a distribution of discrete velocities. The distribution was obtained by considering that the initial energies were either 10 or 20 kcal/mole and that the energy of the atom was always reduced by 21% (the average fraction lost) in collisions with H_2 .³⁵ The discrete velocities of the H atom were then $v, v', v'' \dots$. We took these to be equal to the relative velocities in collisions with F_2 . The cross sections $\sigma_v, \sigma_{v'}, \sigma_{v''}$, etc., of H atoms with these relative velocities were obtained from Fig. 3 of Ref. 16, and were determined using a classical

distribution of J at 1000°K. The rate of rotational relaxation of heavy molecules (such as F₂) is fast enough³⁶ so that we can consider the rotational and translational temperatures of F₂ to be about equal and the F₂ to attain a rotational temperature of 1000°K during an explosion. The effective rate constant for each species, H* and H** , was calculated from

$$k = N_A [f \cdot v \cdot \sigma_v + f' \cdot v' \cdot \sigma_{v'} + f'' \cdot v'' \cdot \sigma_{v''} + \dots]$$

where f, f', f''... are the fractions of H* and H** reacting while at velocities v, v', v'', etc., and N_A is Avogadro's number. About 65% of the H* atoms and about 85% of the H** react with F₂ before being reduced to thermal (<2.6 kcal/mole) energies. The rate constants obtained for the atoms at 10 and 20 kcal/mole were, respectively,

$$k_h = 5.5 \times 10^{14} \text{ cm}^3/\text{mole-sec}$$

$$k'_h = 9.9 \times 10^{14} \text{ cm}^3/\text{mole-sec.}$$

In the program we combined all H atoms into one species and calculated an effective rate constant k_{eff} :

$$k_{th}(H_{th})(F_2) + k_h(H^*)(F_2) + k'_h(H^{**})(F_2) \\ = k_{eff}(H) \cdot (F_2).$$

The value of k_{eff} is given by

$$k_{eff} = \frac{1}{\frac{x}{k_{th}} + \frac{x_1}{k_h} + \frac{1-x-x_1}{k'_h}},$$

where x is the fraction of H atoms that reacts with a thermal rate constant, x₁ is the fraction that started as H* and reacted with rate constant k_h, and (1-x-x₁) is the fraction that started as H** and reacted with rate constant k'_h. Because 65% of the H* and 85% of the H** react while hot, the fractional contributions in Table XVII are corrected to x = (0.6 + 0.35(1.00) + 0.15(0.25))/1.85 = 0.53, x₁ = 0.65/1.85 = 0.35, and 1-x-x₁ = 0.12. When

these and the values for k_{th}, k_h, and k'_h are used in the above equation for k_{eff}, the net effect is $k_{eff} = 2 k_{th}$:

$$k_{eff} = 3.2 \times 10^{14} 10^{-570/T}. \quad (4)$$

We use

$$\sum_{i=23}^{30} RK(i) = 3.2 \times 10^{14} 10^{-570/T}.$$

The initial velocities of the 10- and 20-kcal H atoms are 9.1 x 10⁵ and 1.29 x 10⁶ cm/sec, respectively. About half the H atoms reacting in the system have velocities close to these values. To obtain the relative values of the rate coefficients among reactions (23)-(30) we used Blais' distribution (Ref. 16, Fig. 9) of HF(v) over v as calculated for V_R = 1.08 x 10⁶ cm/sec:

$$RK(23) = 0$$

$$(24) = 3.2 \times 10^{11} \cdot 10^{-570/T}$$

$$(25) = 55 RK(24)$$

$$(26) = 177 RK(24)$$

$$(27) = 220 RK(24)$$

$$(28) = 309 RK(24)$$

$$(29) = 166 RK(24)$$

$$(30) = 72 RK(24).$$

7. The rate coefficients for reactions (-23)-(-30) are taken to be zero. All the reactions are strongly endothermic and the probability is negligible that a hot F atom collides with an HF molecule with sufficiently high vibrational energy (v ≥ 6) to overcome the energy barrier. In a computed simulated explosion, the concentrations of the F₂ molecules were about a factor of 10 to 20 larger than the concentrations of HF(v ≥ 6), and were sufficiently high to moderate the F atoms before they collided with HF(v ≥ 6).

8. The rates of V-V transfer reactions among HF molecules, all measured near room temperature, are summarized in Table XVIII. The rate of 1 + 1 → 0 + 2 was measured by Bott³⁷ by monitoring the rate of appearance of the V = 2 level. The rates of V + 0 → (V - 1) + 1 were determined^{38,39,40}

from the rate of disappearance of the V level with the assumption that the major process was the V-V transfer and not the V-R,T transfer. The three

TABLE XVIII

MEASURED VALUES OF HF-HF V-V TRANSFER RATES

Reaction No.	Reaction	Temp (°K)	k (cm ³ /mole-sec)	Ref.
(31)	1 + 1 + 0 + 2	295	2.2 x 10 ¹³	37
	2 + 0 + 1 + 1	290	1.19 x 10 ¹³	38
	3 + 0 + 2 + 1	290	2.88 x 10 ¹³	39
	2 + 0 + 1 + 1	300?	1.12 x 10 ¹³	40
	3 + 0 + 2 + 1	300?	1.20 x 10 ¹³	40
	4 + 0 + 3 + 1	300?	>3.2 x 10 ¹³	40
	5 + 0 + 4 + 1	300?	>4.9 x 10 ¹³	40
(31)	1 + 1 + 2 + 0	290	2.8 x 10 ¹³	38
(31)	1 + 1 + 2 + 0	300?	2.6 x 10 ¹³	40 ^a
(32)	2 + 1 + 3 + 0	300?	6.1 x 10 ¹³	40 ^a
(33)	3 + 1 + 4 + 0	300?	>3.5 x 10 ¹⁴	40 ^a
(34)	4 + 1 + 5 + 0	300?	>1.17 x 10 ¹⁵	40 ^a

^aCalculated from the reverse reaction to obtain the rate of the exothermic reaction.

values of the rate coefficients for 1 + 1 + 2 + 0 are in good agreement. Bott and Cohen⁴¹ have shown that the rate coefficient for

decreases by about 50% as the temperature increases from 300 to 700°K and that there may be a minimum near 700°K with the rate coefficient increasing as T increases above 700°K. The work of Ahl and Cool⁴² at 300 and 350°K indicates that the rate coefficient for this reaction may decrease as T increases. For reaction (31) we consider the temperature dependence to be slight and we use

$$\text{RK}(31) = 2.5 \times 10^{13} \text{ cm}^3/\text{mole-sec}$$

at all temperatures.

The rates of (32)-(51) relative to (31) were calculated from $R(i) = F(i) \cdot R(31)$ where the factor $F(i)$ is

$$F(i) = \frac{[(\lambda+1) \cdot \ell]_i}{[(\lambda+1) \cdot \ell]_{31}} \cdot \frac{f_i}{f_{31}}$$

and where the ratio $[(\lambda+1) \cdot \ell]_i / [(\lambda+1) \cdot \ell]_{31}$ is obtained from theory⁴³

$$P_{\ell \rightarrow \ell-1}^{\lambda \rightarrow \lambda+1} = (\lambda+1) \cdot \ell \cdot P_{1 \rightarrow 0}^{0 \rightarrow 1}$$

The factor f_i/f_{31} is used on the assumption that an energy defect between products and reactants reduces the rate.⁴⁴ Straight-line plots of the logarithm of measured V-V energy transfer probability vs energy defect for the 1 → 0, 0 → 1 transition are in the literature for HF-diatomic, DCl-diatomic, HBr-diatomic, and HCl-diatomic systems,⁴⁵ for HF-diatomic systems,⁴¹ for various diatomic systems,⁴⁶ and for the HCl-hydrogen halide systems^{47,48,49} (the HCl-hydrogen halide plots from the same data). The data in Ref. 45 for the HF-diatomic system are for HF and nonpolar molecules and are perhaps not a close approximation to the effects in HF-HF V-V transfer. We considered originally that the effect of the energy defect on HF-HF V-V transfer might be approximated best by the HCl-hydrogen halide systems, and we obtained the factors f_i/f_{31} from Fig. 3 of Ref. 47. The same factors f_i/f_{31} were obtained from later data⁴² when a straight-line plot was made using the two points defined by

$$E_d = 172 \text{ cm}^{-1} \quad k = 2.5 \times 10^{13} \text{ cm}^3/\text{mole-sec}$$

$$E_d = 1055 \text{ cm}^{-1} \quad k = 2 \times 10^{12} \text{ cm}^3/\text{mole-sec.}$$

Similar, but not identical, factors would be obtained from the points for these two reactions as given by Bott and Cohen⁴¹ in their Fig. 12. The factors $F(i)$ which give the rates of (32)-(51) relative to that of (31) are given in Table XIX.

Multiquantum transitions are not included, since there are no hard experimental data, although theoretical^{50,51} work indicates that such transitions may be significant.

The rates of reactions (32), (33), and (34) as obtained from Table XIX are much lower than those measured by Airey and Smith⁴⁰ and given in Table

TABLE XIX

FACTORS RELATING RK(32)-RK(51) TO RK(31)

Reaction No.	HF + HF → HF + HF	E_d (cm ⁻¹)	$(\lambda+1) \cdot \nu$	f_i	$F(i)$
(31)	1 + 1 = 0 + 2	172	2	1.	1.
(32)	1 + 2 = 0 + 3	339	3	0.42	0.63
(33)	1 + 3 = 0 + 4	503	4	0.2	0.4
(34)	1 + 4 = 0 + 5	662	5	0.08	0.2
(35)	1 + 5 = 0 + 6	819	6	0.04	0.12
(36)	1 + 6 = 0 + 7	973	7	0.03	0.11
(37)	2 + 2 = 1 + 3	167	6	1.	3.
(38)	2 + 3 = 1 + 4	331	8	0.45	1.8
(39)	2 + 4 = 1 + 5	490	10	0.2	1.0
(40)	2 + 5 = 1 + 6	647	12	0.09	0.54
(41)	2 + 6 = 1 + 7	800	14	0.04	0.28
(42)	3 + 3 = 2 + 4	164	12	1.	6.
(43)	3 + 4 = 2 + 5	322	15	0.46	3.5
(44)	3 + 5 = 2 + 6	480	18	0.21	1.9
(45)	3 + 6 = 2 + 7	633	21	0.095	1.1
(46)	4 + 4 = 3 + 5	159	20	1.	10.
(47)	4 + 5 = 3 + 6	316	24	0.48	5.7
(48)	4 + 6 = 3 + 7	470	28	0.22	3.1
(49)	5 + 5 = 4 + 6	156	30	1.1	16.
(50)	5 + 6 = 4 + 7	310	35	0.5	8.7
(51)	6 + 6 = 5 + 7	154	42	1.1	23.

XVIII. Some of the experimentally determined rate coefficients in Table XVIII, even for reactions with large energy defects, are higher than gas-kinetic collision rates. We have assumed that the relationship between energy defect and rate holds for higher vibrational states, but this may not be correct. It then appears that the only reasonable sure value among the rates of (31)-(51) is RK(31). The others must be considered to be estimates.

9. The reported values of the rate of (52) and (-52) are given in Table XX. The measured temperature dependence of k_{52} in the range 300 to 1000°K is slight.⁴¹ We use, at all temperatures,

$$RK(52) = 1.1 \times 10^{12} \text{ cm}^3/\text{mole-sec.}$$

There are no experimental data on reactions (53)-(100). We assume that each rate constant RK(i) can be obtained from $RK(i) = F(i) \cdot RK(52)$ where

$$F(i) = \frac{[(\lambda+1) \cdot \nu]_i}{[(\lambda+1) \cdot \nu]_{52}} \cdot \frac{f_i}{f_{52}}.$$

TABLE XX

MEASURED VALUES OF THE RATE CONSTANTS
FOR REACTIONS (52) AND (-52)

Temp. (°K)	k_{-52} (cm ³ /mole-sec)	k_{52} (cm ³ /mole-sec)	Ref.
294	--	1.15×10^{12}	52
294	4.4×10^{11}	1.1×10^{12} ^a	45,52
298	4.4×10^{11}	1.1×10^{12} ^a	53
295	3.05×10^{11}	8.1×10^{11} ^a	41

^aCalculated from $k_{52} = k_{-52} \cdot 10^{1.99/1.6T}$.

The factors f_1/f_{52} were obtained from Fig. 3 of Ref. 45 where the V-V exchange probability is plotted as a function of energy defect using data for the systems HF-H₂, HF-D₂, and HF-N₂. The factors F(i) that are used in the program are given in Table XXI.

10. The V-R,T relaxation rate of

was recently measured in the range 600 to 2400°K and near 300°K by Blair, Breshears, and Schott.⁷ Their work contains a summary of the results of previous investigations. In obtaining an equation for the rate coefficient of Reaction (d), we weighted their work heavily. Our rate coefficient

$$k_a = 8.2 \times 10^8 \cdot T + 1.2 \times 10^{17} T^{-2}$$

reproduces the values of τ_P given in Fig. 7 of their paper within about 5% in the range 400 to 2400°K, and at 300°K gives a value of τ_P about 20% higher than that of Ref. 7, but a value that is still centered within values obtained by other workers. There are no data on the rates of V-R,T relaxation of HF(1) by HF in higher vibrational states. We assume that all states have the same effectiveness as HF(0). For the reaction

TABLE XXI

RELATIVE RATE CONSTANTS FOR REACTIONS (53)-(100)

Reaction No.	HF + H ₂ → HF + H ₂	E _d (cm ⁻¹)	(λ+1)·λ	f _i	F(i)
(52)	0 + 1 = 1 + 0	199	1	1.	1.
(53)	1 + 1 = 0 + 2	37	2	1.8	3.6
(54)	1 + 2 = 0 + 3	268	3	0.76	2.3
(55)	1 + 3 = 0 + 4	494	4	0.32	1.3
(56)	1 + 4 = 0 + 5	720	5	0.13	0.75
(57)	1 + 5 = 0 + 6	947	6	0.056	0.33
(58)	1 + 6 = 0 + 7	1180	7	0.028	0.2
(59)	1 + 1 = 2 + 0	370	2	0.52	1.
(60)	1 + 2 = 2 + 1	135	4	1.3	5.2
(61)	2 + 2 = 1 + 3	96	6	1.5	9.
(62)	2 + 3 = 1 + 4	322	8	0.62	4.9
(63)	2 + 4 = 1 + 5	548	10	0.26	2.6
(64)	2 + 5 = 1 + 6	755	12	0.12	1.4
(65)	2 + 6 = 1 + 7	1007	14	0.045	0.63
(66)	2 + 1 = 3 + 0	538	3	0.27	0.8
(67)	2 + 2 = 3 + 1	302	6	0.68	4.1
(68)	2 + 3 = 3 + 2	72	9	1.6	14.
(69)	3 + 3 = 2 + 4	155	12	1.2	14.
(70)	3 + 4 = 2 + 5	380	15	0.5	7.5
(71)	3 + 5 = 2 + 6	608	18	0.2	3.6
(72)	3 + 6 = 2 + 7	840	21	0.085	1.8
(73)	3 + 1 = 4 + 0	702	4	0.145	0.58
(74)	3 + 2 = 4 + 1	465	8	0.36	2.9
(75)	3 + 3 = 4 + 2	235	12	0.86	10.
(76)	3 + 4 = 4 + 3	8	16	2.	32.
(77)	4 + 4 = 3 + 5	217	20	0.94	19.
(78)	4 + 5 = 3 + 6	445	24	0.4	9.6
(79)	4 + 6 = 3 + 7	677	28	0.16	4.5
(80)	4 + 1 = 5 + 0	860	5	0.08	0.4
(81)	4 + 2 = 5 + 1	625	10	0.19	1.9
(82)	4 + 3 = 5 + 2	395	15	0.46	6.9
(83)	4 + 4 = 5 + 3	168	20	1.1	2.2
(84)	5 + 4 = 4 + 5	58	25	1.7	42.
(85)	5 + 5 = 4 + 6	285	30	0.72	22.
(86)	5 + 6 = 4 + 7	517	35	0.3	10.5
(87)	5 + 1 = 6 + 0	1017	6	0.045	0.27
(88)	5 + 2 = 6 + 1	780	12	0.11	1.4
(89)	5 + 3 = 6 + 2	550	18	0.26	4.7
(90)	5 + 4 = 6 + 3	324	24	0.62	14.9
(91)	5 + 5 = 6 + 4	99	30	1.5	45.
(92)	6 + 5 = 5 + 6	130	36	1.3	47.
(93)	6 + 6 = 5 + 7	360	42	0.52	21.
(94)	6 + 1 = 7 + 0	1170	7	0.025	0.2
(95)	6 + 2 = 7 + 1	935	14	0.06	0.8
(96)	6 + 3 = 7 + 2	705	21	0.15	3.1
(97)	6 + 4 = 7 + 3	480	28	0.34	9.5
(98)	6 + 5 = 7 + 4	253	35	0.8	28.
(99)	6 + 6 = 7 + 5	26	42	2.	84.
(100)	7 + 6 = 6 + 7	206	49	0.95	46.5

where M_i includes all vibrational states of HF, we use

$$RK(101) = k_a$$

A recent theoretical calculation of the rate of Reaction (d) is that of Berend and Thommarson.⁵² A mechanism that reproduces the temperature dependence of the relaxation rate of HF has been given by Thompson.⁵³

The rate of V-R,T relaxation of the v = 2 state by HF(0) is twice the rate of relaxation of the v = 1 state.^{37,38,54} There is evidence that the v = 3 state also relaxes with a rate proportional to v.³⁹ However, trajectory calculations⁵⁵ on the rates of relaxation of higher vibrational states in a somewhat similar reaction

indicate that if all the vibrational energy transfer is ascribed to single-quantum transfers, then the rate of reaction is proportional, roughly, to v^{1.6} (see Note 12). In lieu of any other information on the self-relaxation of higher v states in HF we considered the relaxation rate to be proportional to v^{1.6}, and we used the following rate coefficients

$$RK(102) = 3 RK(101)$$

$$RK(103) = 5.8 RK(101)$$

$$RK(104) = 9.7 RK(101)$$

$$RK(105) = 13 RK(101)$$

$$RK(106) = 17.5 RK(101)$$

$$RK(107) = 22 RK(101)$$

Blais and Thompson⁵⁶ showed that rotational excitation considerably enhances the cross section for vibrational relaxation by F atoms:

The same effect should be expected in reactions (102)-(107), where rotationally excited HF(v) would be colliding with HF.

11. There are virtually no experimental data on the rate of

Upper limits of $k = 1.1 \times 10^{10}$ cm³/mole-sec and 1.8×10^{10} cm³/mole-sec at room temperature are given, respectively, by Bott and Cohen⁴¹ and by Hancock and Green.^{45,57} Vasil'ev et al.⁵⁸ obtained a value of 9×10^8 cm³/mole-sec, but this is for a mixed process, V-V and V-R,T transfer; Hinchey⁵⁹ assumed a value 7.3×10^{10} cm³/mole-sec. Kapralova et al.⁶⁰ obtained a theoretical value of 2.7×10^9 cm³/mole-sec. There are no data at higher temperatures. For

where M_2 includes all the vibrational states of H_2 , we used for all temperatures the rate obtained by Bott and Cohen⁶¹ in the 1350-to-3000°K range for

Thus,

$$\text{RK}(108) = 1.61 \times 10^{11} \cdot T \cdot 10^{-42.0/T^{1/3}} .$$

All vibrational states of H_2 are assumed to have the same effectiveness as D_2 . The rate as given by $\text{RK}(108)$ is about a factor of 7 to 10 greater than the rates of the V-R,T process in HF-He in the range 1350 to 3000°K, and would seem to be reasonable for V-R,T transfer in H_2 in the same temperature range. At 300°K the above expression for $\text{RK}(108)$ gives 2.6×10^7 cm³/mole-sec, which is somewhat lower than the upper limit of $k < 7 \times 10^8$ cm³/mole-sec, as determined at 300°K for HF-He V-R,T relaxation;⁴⁵ it is also considerably lower than the upper limits (given above) that were obtained by Bott and Cohen⁴¹ and by Hancock and Green^{45,57} for HF- H_2 V-R,T relaxation. The correct value is very likely $2.6 \times 10^7 < \text{RK}(108) < 1.1 \times 10^{10}$ cm³/mole-sec. We take the rates of relaxation from the upper vibrational states to be

$$\text{RK}(109) = 3 \text{ RK}(108)$$

$$\text{RK}(110) = 5.8 \text{ RK}(108)$$

$$\text{RK}(111) = 9.7 \text{ RK}(108)$$

$$\text{RK}(112) = 13 \text{ RK}(108)$$

$$\text{RK}(113) = 17.5 \text{ RK}(108)$$

$$\text{RK}(114) = 22 \text{ RK}(108) .$$

Relaxation rates would probably be enhanced by the rotational excitation of $\text{HF}(v)$.

12. The rate of

was measured by Blair, Breshears, and Schott⁷ in the range 1500 to 2500°K, by Bott and Cohen,⁶² and by Solomon et al.⁶³ Blair et al. found that F atoms are about 2 to 5 times more efficient than HF in relaxing HF, while Bott and Cohen and Solomon found that they were about 20 times more efficient. We use

$$\text{RK}(115) = 3.99 \times 10^{10} \cdot T \cdot 10^{-14.9/T^{1/3}} ,$$

which reproduces within 0.5% the measured relaxation rates of Ref. 7. Although 10 to 20% of the hot F atoms collide with $\text{HF}(v)$ before being moderated, we make no correction to the thermal rate. Thompson⁵⁵ and Wilkins⁶⁴ calculated the rates theoretically. The rate coefficients are compared in Table XXII.

TABLE XXII

RATE COEFFICIENTS FOR $\text{HF}(1) + \text{F} \rightarrow \text{HF}(0) + \text{F}$
(NOT INCLUDING ATOM EXCHANGE REACTIONS) AS
DETERMINED BY TRAJECTORY CALCULATIONS

Temp. (°K)	(10 ¹² cm ³ /mole-sec)		
	$\text{RK}(115)$	$k(\text{Wilkins})^a$	$k(\text{Thompson})^b$
1000	1.29	0.7	3.5
1500	2.98	1.	4.5
2000	5.19	~1.2	6.8
2500	7.88	~1.3	--

^aAs read from Fig. 1 of Ref. 64.

^bReference 55.

There are no experimental data on relaxation of higher vibrational states of HF by F. In some of Thompson's trajectories amounts of vibrational energy corresponding to several quanta were transferred. When rate coefficients were calculated by attributing all the energy transferred to one-quantum transfers, he obtained the values given in Table XXIII.

TABLE XXIII

RATE COEFFICIENTS FOR $\text{HF}(v) + \text{F} \rightarrow \text{HF}(v-1) + \text{F}$ AS TAKEN FROM TABLE II OF REF. 55

Temp. (°K)	$(10^{12} \text{ cm}^3/\text{mole-sec})$		
	$k_{1 \rightarrow 0}$	$k_{3 \rightarrow 2}$	$k_{5 \rightarrow 4}$
1000	3.5	15.5	
1500	4.5	26.1	75.4
2000	6.8	30.3	119.

As Thompson points out, the rapid increase in k with v (greater than the theory⁴³ $k_{v \rightarrow v-1} \approx vk_{1 \rightarrow 0}$) suggests that multiquantum transfers took place. Wilkins⁶⁴ also found in his trajectory calculations that amounts of energy corresponding to several quanta were transferred out of vibration. He interpreted his calculations in two ways: (1) He partitioned the continuum states of the products into discrete quantum states and reported rate constants as a function of temperature for the individual processes

for the cases $3 \rightarrow 2$, $3 \rightarrow 1$, $6 \rightarrow 5$, $6 \rightarrow 4$, $6 \rightarrow 3$, $6 \rightarrow 2$. (2) He (as did Thompson) also attributed all the energy transferred to single-quantum transfers and calculated the rates for these processes. There is some question about the validity of partitioning the products into discrete states. We report Wilkins' results in Table XXIV as read from Figs. 1 and 6 of his paper,⁶⁴ and as calculated on the assumption of single-quantum transfers.

Blais and Thompson⁵⁶ have shown that the cross sections for relaxation by atom exchange and by inelastic scattering are about equal. To account for relaxation by both processes, the rate coefficients

TABLE XXIV

RATE COEFFICIENTS FOR $\text{HF}(v) + \text{F} \rightarrow \text{HF}(v-1) + \text{F}$ AS DETERMINED IN REF. 64

Temp. (°K)	$(10^{12} \text{ cm}^3/\text{mole-sec})$			
	$k_{1 \rightarrow 0}$	$k_{2 \rightarrow 1}$	$k_{3 \rightarrow 2}$	$k_{6 \rightarrow 5}$ ^a
1000	0.7	1.	2.	25.
1500	1.	1.5	2.8	36.
2000	~1.2	1.6	3.2	43.

^aRecalculated from the results in Fig. 6 of Ref. 64 to obtain rates on the assumption of single-quantum transfers. We took Wilkins' values for the rate coefficients k' as he reported them for the individual transitions $\text{F} + \text{HF}(v) \rightarrow \text{HF}(v') + \text{F}$ (including atom exchange reactions), and then calculated $k_{6 \rightarrow 5}$ (in which all the change in vibrational energy, previously distributed among the individual transitions, is now attributed to the $6 \rightarrow 5$ transition),

$$k_{6 \rightarrow 5} = k'_{6 \rightarrow 5} + 2k'_{6 \rightarrow 4} + 3k'_{6 \rightarrow 3} + 4k'_{6 \rightarrow 2}.$$

of Wilkins and Thompson in Table XXII, those in Table XXIII, and the values of $k_{1 \rightarrow 0}$, $k_{2 \rightarrow 1}$, $k_{3 \rightarrow 2}$ in Table XXIV should be doubled.

Thompson's results for the $1 \rightarrow 0$ transition are in good agreement with the experimental data, and his results for higher v states indicate that if the approximation of single-quantum transfers is made, then the rate coefficients increase with v at a rate greater than $k_{v \rightarrow v-1} = vk_{1 \rightarrow 0}$, predicted by theory.⁴³ From Thompson's results we take the rate coefficients for the higher vibrational states to be

$$k_v = v^{1.4} k_{1 \rightarrow 0},$$

and our rate coefficients are

$$\text{RK}(116) = 3 \text{ RK}(115)$$

$$(117) = 5.8 \text{ RK}(115)$$

$$(118) = 9.7 \text{ RK}(115)$$

$$(119) = 13. \text{ RK}(115)$$

$$(120) = 17.5 \text{ RK}(115)$$

$$(121) = 22. \text{ RK}(115)$$

If an appreciable fraction of HF collides with H atoms while the HF molecules are rotationally

excited (Note 4), a similar fraction collides with F atoms. Blais and Thompson⁵⁶ showed that rotational excitation considerably enhances the cross section for vibrational relaxation by F atoms, and our rates as written above may be low.

13. There are no experimental data for the rate of

Thompson,⁶⁵ by trajectory calculations, obtained rates for this reaction and also for relaxation from higher states. His Table II is copied here as Table XXV.

TABLE XXV

RATE COEFFICIENTS FOR H-ATOM VIBRATIONAL RELAXATION OF HF, AS CALCULATED FROM CLASSICAL TRAJECTORIES (REF. 65)

T (°K)	(10 ¹² cm ³ /mole-sec)			
	$k_{1 \rightarrow 0}$	$k_{3 \rightarrow 2}$	$k_{5 \rightarrow 4}$	$k_{7 \rightarrow 6}$
600	1.4	2.3		
1000	2.3	6.7	75.5	324.
1500	5.0	15.1	99.8	
2000	10.7	64.0	151.	348.

Wilkins²⁸ reported rate constants as a function of temperature in the range 100 to 1000°K for each of the processes

We recalculated his results for 1000°K, as read from his Fig. 3 and assuming single-quantum transfers. We used his values for $k'_{v \rightarrow v'}$, the rate coefficients for the individual transitions $\text{HF}(v) \rightarrow \text{HF}(v')$, to obtain values of the rate coefficients k_v , in which all the change in vibrational energy is attributed to single-quantum transfers,

$$k_v = k'_{v \rightarrow v-1} + 2k'_{v \rightarrow v-2} + \dots + vk'_{v \rightarrow 0}.$$

The results are in Table XXVI. The values in Table XXVI correspond roughly to values that would be obtained from Blais' potential-energy surface (used

by Thompson) for a system at a temperature slightly greater than 2000°K.

TABLE XXVI

RATE COEFFICIENTS FROM TRAJECTORY CALCULATIONS (REF. 28, T = 1000°K)

(10 ¹² cm ³ /mole-sec)					
$k_{1 \rightarrow 0}$	$k_{2 \rightarrow 1}$	$k_{3 \rightarrow 2}$	$k_{4 \rightarrow 3}$	$k_{5 \rightarrow 4}$	$k_{6 \rightarrow 5}$
8.	47.	90.	105.	190.	250.

If we assume that the probability of the $1 \rightarrow 0$ relaxation per collision is the same for H atoms as for F atoms, we can obtain approximate rates for relaxation by H atoms based on the experimental data for relaxation by F atoms [Ref. 7 and RK(115)]. For thermal H atoms the collision rate would be greater than that for F atoms by about a factor of three. By using the values of RK(115) in Table XXII we obtain the rate coefficients for thermal H atoms seen in Table XXVII.

TABLE XXVII

RATE COEFFICIENTS FOR $\text{H} + \text{HF}(1) \rightarrow \text{H} + \text{HF}(0)$ OBTAINED FROM EXPERIMENTAL VALUES FOR $\text{F} + \text{HF}(1) \rightarrow \text{F} + \text{HF}(0)$

Temp. (°K)	$k(122) = 3\text{RK}(115)$ (10 ¹² cm ³ /mole-sec)
1000	3.9
1500	9.
2500	23.

These rates are about a factor of two larger than Thompson's, and the rate for 1000°K is one-half that given by Wilkins. The velocity distribution of H atoms in an $\text{H}_2\text{-F}_2$ explosion, however, is part thermal and part peaked at 9×10^5 cm/sec (10 kcal/mole) and 1.3×10^6 cm/sec (20 kcal/mole). Trajectory calculations were made for $v_R = 9 \times 10^5$ cm/sec and rotational temperature = 2000°K to obtain $k(122) = 5.3 \times 10^{13}$ cm³/mole-sec. The rate coefficient for thermal atoms is about 1×10^{13} cm³/mole-sec (Tables XXV, XXVI, XXVII). About 53% of the H atoms are thermal and 47% are hot (Note 6). Then, by combining

the rate coefficients by using the equation in Note 6 we obtain $RK(122) = 1.6 \times 10^{13} \text{ cm}^3/\text{mole-sec.}$

Both Thompson's and Wilkins' results indicate that if we attribute all the change in vibrational energy to single-quantum transfers, the dependence of the rate on v is much stronger than $k_{v \rightarrow v-1} = vk_{1 \rightarrow 0}$. We use the following rates, as estimated from Tables XXV and XXVI

$$RK(123) = 3 RK(122)$$

$$(124) = 4 RK(122)$$

$$(125) = 12 RK(122)$$

$$(126) = 20 RK(122)$$

$$(127) = 30 RK(122)$$

$$(128) = 50 RK(122) .$$

An appreciable fraction of rotationally excited HF molecules collides with H atoms (Note 4). As with F atoms, the V-R,T relaxation rate is probably enhanced over the rates given for (122)-(128).

14. An upper limit has been reported³⁴ for the rate of relaxation of HF by F_2 at 350°K, $k < 2 \times 10^9 \text{ cm}^3/\text{mole-sec.}$; the relaxation was either V-R,T or V-V. There are no other data for

The rate of relaxation of HCl by Cl_2 molecules is reported⁶⁶ to be 1/2000th the rate of relaxation by Cl atoms. We take the chemical affinity between F_2 and HF to be slight and take the rate at all temperatures to be given by the rate expression determined at 1350 to 4000°K for Ar,⁶¹

$$RK(129) = 5.06 \times 10^{10} \cdot T \cdot 10^{-46.6/T^{1/3}} .$$

The equation for RK(129) gives $2.26 \times 10^6 \text{ cm}^3/\text{mole-sec}$ at 350°K, which can be compared to the upper limit $k < 2 \times 10^9 \text{ cm}^3/\text{mole-sec.}$ The rates of relaxation of the upper states are taken to be proportional to $v^{1.6}$. The HF(v) collide with F_2 molecules while rotationally excited, and the rates are probably greater than those we use.

15. The rate of

was measured by Bott and Cohen⁶¹ in the range 1350 to 3500°K. An upper limit of $7 \times 10^8 \text{ cm}^3/\text{mole-sec}$ at 294°K was determined by Green and Hancock.⁴⁵ We take the rate expression determined from the results of Bott and Cohen to be applicable at all temperatures and we use

$$RK(136) = 5.41 \times 10^{11} \cdot T \cdot 10^{-57.9/T}.$$

This expression gives $RK(136) = 3.2 \times 10^3 \text{ cm}^3/\text{mole-sec}$ at 294°K. The rates of relaxation from higher vibrational states are taken to be proportional to $v^{1.6}$.

Again, the rates are probably greater than those we use because of rotational excitation of HF.

16. There are no experimental data for the rates of V-R,T relaxation of H_2 by F_2 or by F atoms. There are room-temperature data for the V-R,T relaxation by HF,

Hinchen⁵⁹ estimates $k = 5 \times 10^{11} \text{ cm}^3/\text{mole-sec.}$ Lower values of upper limits for the rate coefficient are given by other workers: Bott and Cohen,⁴¹ $k = 3 \times 10^{10} \text{ cm}^3/\text{mole-sec.}$; Hancock and Green,⁶⁷ $k = 2.2 \times 10^{10} \text{ cm}^3/\text{mole-sec.}$; Osgood,⁶⁸ $k = 5 \times 10^{10} \text{ cm}^3/\text{mole-sec.}$ A comparison of the rate of this reaction (based on these upper limit values) with the rate of

in a computed simulated H_2-F_2 explosion shows that the rate of Reaction (e) is lower than the rate of self-relaxation of HF by a factor of about 300. Reaction (e) is then not important in the degradation of the vibrational energy in the system. We assumed that F_2 , F, and HF (all vibrational states) have the same efficiency for V-R,T relaxation of H_2 as does Ar, which rate was determined in the range 1500 to 2700°K as 0.24 times the rate of relaxation by H_2 .⁶⁹ For

where M_3 means all species except H_2 and H , we then use for all temperatures

$$\text{RK}(143) = 5.05 \times 10^{10} \cdot T \cdot 10^{-43.4/T^{1/3}}$$

$$= 0.24 \text{ RK}(150).$$

The rate of relaxation of H_2 in Ar was measured⁷⁰ at 300°K and found to be about 40% lower than the value obtained from the above equation. We take the rates of relaxation of the higher states to be proportional to v .

17. The rate of relaxation

was measured in the range 1100 to 2700°K.⁶⁹ The rate coefficient was

$$k_f = 2.105 \times 10^{11} \cdot T \cdot 10^{-43.4/T^{1/3}} .$$

We assume that H atoms and all vibrational states of H_2 are equally effective in relaxing $\text{H}_2(1)$ and that the above equation is valid at all temperatures of interest. We use

$$\text{RK}(150) = k_f$$

for the reaction

where M_4 represents H atoms and H_2 in all vibrational levels. The equation for $\text{RK}(150)$ gives $k = 2.08 \times 10^7 \text{ cm}^3/\text{mole-sec}$ at 300°K. A measured value at 300°K is $7.94 \times 10^6 \text{ cm}^3/\text{mole-sec}$.^{71,72} The rate of

at 300°K has been measured as $k = 1.2 \times 10^8 \text{ cm}^3/\text{mole-sec}$.⁷³ Should the rate of relaxation by H atoms be greater by a factor of five to six than that by H_2 , our calculations are not significantly affected;

the concentration of H atoms is always less than a few percent of the concentration of H_2 molecules, and the rate of V-R,T relaxation of $\text{H}_2(v)$ is low compared to the dominant rate of V-R,T relaxation in the system, that of HF . We take the rates of relaxation of the higher v states to be proportional to v .

18. For reactions (157)-(162) we obtain the recombination rate from the dissociation rate and the equilibrium constant. The rates of dissociation of F_2 by Ar and by F_2 were recently measured, and the rate of dissociation of F_2 by F was estimated in the range 1400 to 2600°K by Breshears and Bird.⁷⁴ (Their paper also contains references to previous work.) We assume the rate of dissociation in He to be the same as that in Ar . Then, using their rates,

$$\text{RK}(157) = 4.6 \times 10^{12} \text{ cm}^6/\text{mole}^2\text{-sec}$$

$$\text{RK}(158) = 1.21 \times 10^{13} \text{ cm}^6/\text{mole}^2\text{-sec}$$

$$\text{RK}(161) = 6 \times 10^{13} \text{ cm}^6/\text{mole}^2\text{-sec} .$$

The rate coefficient for (159) is taken to be zero because of the high reactivity of F with H_2 . The rate coefficient for (160) is taken to be equal to that for (161),⁷⁵ and all vibrational states of HF are taken to be equally effective:

[Reaction (160)]

$$\text{RK}(160) = \text{RK}(161) .$$

In explosions initiated by 10- to 100-nsec pulses of F atoms, the rates of both the forward and reverse reactions for (157)-(162) are negligible.

19. In explosions initiated by pulses of F atoms the rates of the forward and reverse reactions for (163)-(178) are also negligible. The forward reactions are included because they might contribute to the overall reaction in other types of explosions. Recombination rates are given by Shui and Appleton⁷⁶ and by Stepukhovich and Umanskii.⁷⁷ We assume that the rates of recombination into all vibrational levels are equal and we use, as obtained from Shui and Appleton's Fig. 2,

$$\text{RK}(163) = 2 \times 10^{16} \text{ cm}^6/\text{mole}^2\text{-sec}$$

for

where M_6 is all species except H. There is some evidence⁷⁷ that H is more efficient as a third body than is H_2 . We take

$$RK(171) = 10 \text{ RK}(163)$$

and the rates of recombination into all vibrational states to be the same. The dissociation rate of H_2 is taken to be zero; the rate of vibrational excitation is probably too low to be excited by T-R,V transfers, and the rate of dissociation from lower levels ($v \leq 6$) is low because of the high activation energy

20. The comments on the recombination and dissociation of H_2 apply to the recombination of H and F atoms into HF. We estimate

$$RK(179) = 1.25 \times 10^{14} \text{ cm}^6/\text{mole}^2\text{-sec}$$

and assume that the rates of recombination into all vibrational levels are equal. The rate of dissociation of HF is taken to be zero. M_5 is all species.

21. Analysis⁷⁸ of experimental data^{79,80} on the explosion limits of $H_2 + F_2$ has shown that the two reactions

contribute to branching. These do not contribute to a pulse-initiated explosion but are included so that the program can be used to study explosion limits of H_2 - F_2 mixtures.

ACKNOWLEDGMENTS

We wish to thank Normand C. Blais, Jon B. Cross, Donald L. Thompson, Garry Schott, Dale Breshears, Richard W. Getzinger, N. R. Greiner, and Professor John M. White of the University of Texas at Austin for helpful and illuminating discussions -- Normand C. Blais and Donald L. Thompson for permission to use their potential energy surface and trajectory program for calculations on $F + H_2 \rightarrow HF + H$.

REFERENCES

1. R. L. Kerber, G. Emanuel, and J. S. Whittier, "Computer Modeling and Parametric Study for a Pulsed $H_2 + F_2$ Laser," *Appl. Opt.* **11**, 1112-1123 (1972).
2. N. L. Rapagnani, "The Effect of Various Chain-Branching Rates on the Pulsed HF Chemical Laser," Lawrence Livermore Laboratory report UCID-16006.
3. S. N. Suchard, R. L. Kerber, G. Emanuel, and J. S. Whittier, "Effect of H_2 Pressure on Pulsed $H_2 + F_2$ Laser, Experiment and Theory," *J. Chem. Phys.* **57**, 5065 (1972).
4. R. L. Kerber, N. Cohen, and G. Emanuel, "A Kinetic Model and Computer Simulation for a Pulsed DF- CO_2 Chemical Transfer Laser," *IEEE J. Quantum Electron.* **QE-9**, 94-113 (1973).
5. R. L. Kerber and J. S. Whittier, "The Effect of Various Parameters on the Behavior of the Pulsed HF Chemical Laser," Aerospace Corp. report TOR-0059(6753-10)-1.
6. N. Cohen, "A Review of Rate Coefficients for Reactions in the H_2 - F_2 Laser System," SAMS0-TR-72-23. Aerospace Corp. report TR-0172(2779)-2.
7. L. S. Blair, W. D. Breshears, and G. L. Schott, "Vibrational Relaxation of HF in the Temperature Range 600-2400°K," submitted to *J. Chem. Phys.*
8. H. W. Wooley, R. B. Scott, and F. G. Brickwedde, "Compilation of Thermal Properties of Hydrogen in Its Various Isotopic and Ortho-Para Modifications," *J. Res.* **41**, 379-475 (1948).
9. D. E. Mann, B. A. Thrush, D. R. Lide, Jr., J. J. Ball, and N. Acquista, "Spectroscopy of Fluorine Flames. I. Hydrogen-Fluorine Flame and the Vibration-Rotation Emission Spectrum of HF," *J. Chem. Phys.* **34**, 420-431 (1961).
10. K. H. Homann, W. C. Solomon, J. Warnatz, H. Gg. Wagner, and C. Zetzsch, "Eine Methode zur Erzeugung von Fluoratomen in inerter Atmosphäre," *Bunsen-Ges. Phys. Chem.* **74**, 585-589 (1970).
11. S. W. Rabideau, H. G. Hecht, and W. B. Lewis, "A Study of the Kinetics of the Reaction Between H_2 and F_2 by EPR Methods," *J. Magn. Resonance* **6**, 384-395 (1972).
12. A. Wolfgang, "The Hot Atom Chemistry of Gas-Phase Study," *Prog. in React. Kinet.* **3**, 97-169 (1965), p. 118.
13. J. C. Polanyi and J. J. Sloan, "Energy Distribution among Reaction Products. VII. $H + F_2$," *J. Chem. Phys.* **57**, 4988-4998 (1972).
14. N. Jonathan, C. M. Melliar-Smith, and D. H. Slater, "Initial Vibrational Energy Level Populations Resulting from the Reaction $H + F_2$ as Studied by Infrared Chemiluminescence," *J. Chem. Phys.* **53**, 4396-4397 (1970).
N. Jonathan, S. Okuda, and D. Timlin, "Initial Vibrational Energy Distributions Determined by Infrared Chemiluminescence. III. Experimental

- Results and Classical Trajectory Calculations for the $H + F_2$ System," *Mol. Phys.* **24**, 1143 (1972).
15. R. L. Wilkins, "Monte Carlo Calculations of Reaction Rates and Energy Distributions among Reaction Products. III. $H + F_2 \rightarrow HF + F$ and $D + F_2 \rightarrow DF + F$," *J. Chem. Phys.* **58**, 2326-2332 (1973).
 16. N. C. Blais, "Monte Carlo Trajectories, Dynamics of the Reaction $H + F_2$," Los Alamos Scientific Laboratory report LA-4687 (July 1971).
 17. N. C. Blais, "Monte Carlo Trajectories: The Dynamics of the Reaction $F + H_2$," Los Alamos Scientific Laboratory report LA-4603 (April 1971).
 18. P. Borrell, "Relaxation Processes in Gases," *Advan. Mol. Relaxation Processes* **1**, 69-108 (1967-68).
 19. J. C. Polanyi and K. B. Woodall, "Energy Distribution among Reaction Products. VI. $F + H_2, D_2$," *J. Chem. Phys.* **57**, 1574-1586 (1972).
 20. N. Jonathan, C. M. Melliar-Smith, and D. H. Slater, "Initial Vibrational Energy Distributions Determined by Infra-red Chemiluminescence. I. The Reaction of Fluorine Atoms with Hydrogen and Methane," *Mol. Phys.* **20**, 93-102 (1971).
 21. J. H. Parker and G. C. Pimentel, "Vibrational Energy Distribution through Chemical Laser Studies. I. Fluorine Atoms plus Hydrogen or Methane," *J. Chem. Phys.* **51**, 91-96 (1969).
 22. N. C. Blais and D. G. Truhlar, "Monte Carlo Trajectories: Dynamics of the Reaction $F + D_2$ on a Semiempirical Valence-Bond Potential Energy Surface," *J. Chem. Phys.* **58**, 1090-1108 (1973).
 23. J. T. Muckerman, "Classical Dynamics of the Reaction of Fluorine Atoms with Hydrogen Molecules. II. Dependence on the Potential Energy Surface," *J. Chem. Phys.* **56**, 2997-3006 (1972).
 24. R. L. Wilkins, "Monte Carlo Calculations of Reaction Rates and Energy Distributions among Reaction Products. I. $F + H_2 \rightarrow HF + H$," *J. Chem. Phys.* **57**, 912-917 (1972).
 25. R. D. Coombe and G. C. Pimentel, "Temperature Dependence of the Vibrational Energy Distributions in the Reactions $F + H_2$ and $F + D_2$," *J. Chem. Phys.* **59**, 251-257 (1973).
 26. J. T. Muckerman, "Monte Carlo Calculations of Energy Partitioning and Isotope Effects in Reactions of Fluorine Atoms with H_2 , HD , and D_2 ," *J. Chem. Phys.* **54**, 1155-1164 (1971).
 27. Private discussion with N. Blais.
 28. R. L. Wilkins, "Monte Carlo Calculations of Reaction Rates and Energy Distribution among Reaction Products. II. $H + HF(v) \rightarrow H_2(v') + F$ and $H + HF(v) \rightarrow HF(v') + H$," *J. Chem. Phys.* **58**, 3038-3046 (1973).
 29. J. C. Polanyi and K. B. Woodall, "Mechanism of Rotational Relaxation," *J. Chem. Phys.* **56**, 1563-1572 (1972).
 30. M. A. Breazeale and H. O. Kneser, "Rotational Relaxation in Hydrogen Chloride," *J. Acoust. Soc. Amer.* **32**, 885-886 (1960).
 31. A. F. Dodonov, G. K. Lavroskaya, I. I. Morozov, R. G. Albright, V. L. Tal'roze, and A. K. Lyubimova, "Mass Spectrometric Measurement of the Rate Constant of Elementary Reactions. VI. Reactions of Hydrogen Atoms with Chlorine and Fluorine," *Kinet. i Katal.* (English translation) **11**, 677 (1970).
R. G. Albright, A. F. Dodonov, G. K. Lavroskaya, I. I. Morosov, and V. L. Tal'roze, "Mass-Spectrometric Determination of Rate Constants for H-Atom Reactions with Cl_2 and F_2 ," *J. Chem. Phys.* **50**, 3632-3633 (1969).
 32. Numerical values from $\log_{10} p_T$ ($\mu\text{sec-atm}$) = $-6.431 + 60.97 \times T^{-1/2}$ as quoted by W. D. Breshears, LASL Group L-3, and obtained from group at Boston College
 33. The values given are about 10 to 30% lower than those given in R. C. Millikan and D. R. White, "Systematics of Vibrational Relaxation," *J. Chem. Phys.* **39**, 3209-3213 (1963).
 34. S. S. Fried, J. Wilson, and R. L. Taylor, "Measurement of the Temperature Dependence of the Vibrational Relaxation Rate of HF and the Effect of SF_6 , N_2 , and F_2 as Diluents," *IEEE J. Quantum Electron.* **QE-9**, 59-64 (1973).
 35. The average logarithmic loss per collision was taken to be equal to that for T in D_2 . See R. N. Porter and S. Kunt, "Theoretical Studies of Hot-Atom Reactions. II. Yields for Exchange Reactions of Hot Tritium Atoms with Hydrogen and Deuterium Molecules: Simple Models for Nonreactive Scattering," *J. Chem. Phys.* **52**, 3240-3250 (1970).
 36. A. B. Callear and J. D. Lambert, "The Transfer of Energy between Chemical Species," in *Comprehensive Chemical Kinetics* **V. 3**, C. H. Bamford and C. F. H. Tipper, eds. (Elsevier, New York, 1969), Chapter 4, pp. 182-273.
 37. J. F. Bott, "HF Vibrational Relaxation Measurements Using the Combined Shock Tube-Laser-Induced Fluorescence Technique," *J. Chem. Phys.* **57**, 96-102 (1972).
 38. R. M. Osgood, Jr., A. Javan, and P. B. Sackett, "Measurement of Vibration-Vibration Energy Transfer Time in HF Gas," *Appl. Phys. Lett.* **20**, 469-472 (1972).
 39. R. M. Osgood, Jr., P. B. Sackett, and A. Javan, "Measurement of V-V Transfer Rate from HF $v=3$ Using Simultaneous Optical Pumping on the HF $v=2 \rightarrow 1$ and $v=1 \rightarrow 0$ Bands," *Appl. Phys. Lett.* **22**, 254-256 (1973).
 40. J. R. Airey and I. W. M. Smith, "Quenching of Infrared Chemiluminescence: Rates of Energy Transfer from HF($v \leq 5$) to CO_2 and HF, and

- from DF($v \geq 3$) to CO₂ and HF," J. Chem. Phys. 57, 1669-1676 (1972).
41. J. F. Bott and N. Cohen, "Temperature Dependence of V-V and V-R, T Energy Transfer Measurements in Mixtures Containing HF," J. Chem. Phys. 58, 4539-4549 (1973).
42. J. L. Ahl and T. A. Cool, "Vibrational Relaxation in the HF-HCl, HF-HBr, HF-HI, and HF-DF Systems," J. Chem. Phys. 58, 5540-5548 (1973).
43. R. N. Schwartz, Z. I. Slawsky, and K. F. Herzfeld, "Calculation of Vibrational Relaxation Times in Gases," J. Chem. Phys. 20, 1591-1599 (1952).
44. D. Rapp and T. Kassal, "The Theory of Vibrational Energy Transfer Between Simple Molecules in Nonreactive Collisions," Chem. Rev. 69, 61-102 (1969).
45. W. H. Green and J. K. Hancock, "Laser-Excited Vibrational Energy Transfer Studies of HF, CO, and NO," IEEE J. Quantum Electron. QE-9, 50-58 (1973).
46. P. F. Zittel and C. B. Moore, "V-T, R and V-V Relaxation in DCl Systems," J. Chem. Phys. 58, 2922-2928 (1973).
47. Hao-Lin Chen and C. B. Moore, "Vibration-Vibration Energy Transfer in Hydrogen Chloride Mixtures," J. Chem. Phys. 54, 4080-4084 (1971).
48. B. M. Hopkins and Hao-Lin Chen, "Vibrational Excitation and Relaxation of HCl ($v = 2$) State," J. Chem. Phys. 57, 3816-3821 (1972).
49. S. R. Leone and C. B. Moore, "V-V Energy Transfer in HCl with Tunable Optical Parametric Oscillator Excitation," Chem. Phys. Lett. 19, 340-344 (1973).
50. J. D. Kelley, "Vibrational Energy Transfer Processes in Collisions between Diatomic Molecules," J. Chem. Phys. 56, 6108-6117 (1972).
51. T. A. Dillon and J. C. Stephenson, "Multi-quantum Vibrational-Energy Exchange," Phys. Rev. 6A, 1460-1468 (1972).
52. G. C. Berend and R. L. Thommarson, "Vibrational Relaxation of HF and DF," J. Chem. Phys. 58, 3203 (1973).
53. D. L. Thompson, "Mechanisms for Hydrogen Halide Vibrational Relaxation," J. Chem. Phys. 57, 2589-2590 (1972).
54. M. J. Bina and C. R. Jones, "Use of Direct Overtone Excitation in the Measurement of the HF($v=2$) Deactivation Rate," Appl. Phys. Lett. 22, 44-45 (1973).
55. D. L. Thompson, "Monte Carlo Classical Trajectory Calculation of the Rates of F-Atom Vibrational Relaxation of HF and DF," J. Chem. Phys. 57, 4164-4169 (1972).
56. N. C. Blais and D. L. Thompson, "Classical Dynamics: The Role of Atom Exchange in Vibrational Relaxation of HF and DF by F Atoms," to be published in J. Chem. Phys.
57. J. K. Hancock and W. H. Green, "Vibrational Deactivation of HF($v=1$) in Pure HF and in HF-Additive Mixtures," J. Chem. Phys. 57, 4515 (1972) and "Laser-Excited Vibrational Relaxation Studies of Hydrogen Fluoride," J. Chem. Phys. 56, 2474-2475 (1972).
58. G. K. Vasil'ev, E. F. Makarov, V. G. Papin, and V. L. Tal'roze, "Spectroscopic Investigation of the Fluorescent and Radiationless Relaxation of Vibrationally Excited HF Molecules," Dokl. Phys. Chem. 191, 296-299 (1970).
59. J. J. Hinchey, "Vibrational Relaxation of Hydrogen and Deuterium Fluorides," J. Chem. Phys. 59, 233-240 (1973).
60. G. A. Kapralova, E. E. Nikitin, and A. M. Chaikin, "Nonempirical Calculation of Probability of Vibrational Transition in Hydrogen Halides," Kinet. i Katal. (English translation) 10, 793-799 (1969).
61. J. F. Bott and N. Cohen, "Shock-Tube Studies of HF Vibrational Relaxation," J. Chem. Phys. 55, 3698-3706 (1971).
62. J. F. Bott and N. Cohen, "HF Vibrational Relaxation by F Atoms," J. Chem. Phys. 55, 5124-5125 (1971).
63. W. C. Solomon, J. A. Blauer, F. C. Jaye, and J. C. Hnat, "The Vibrational Excitation of Hydrogen Fluoride Behind Incident Shock Waves," Int. J. Chem. Kinet. 3, 215-222 (1971).
64. R. L. Wilkins, "Monte Carlo Calculations of Reaction Rates and Energy Distributions among Reaction Products. IV. $F + HF(v) \rightarrow HF(v') + F$ and $F + DF(v) \rightarrow DF(v') + F$," J. Chem. Phys. 59, 698-704 (1973).
65. D. L. Thompson, "Monte Carlo Classical Trajectory Calculation of the Rates of H- and D-Atom Vibrational Relaxation of HF and DF," J. Chem. Phys. 57, 4170-4173 (1972).
66. N. C. Craig and C. B. Moore, "Vibrational Relaxation of Hydrogen Chloride by Chlorine Atoms and Chlorine Molecules," J. Phys. Chem. 75, 1622-1623 (1971).
67. As quoted in Ref. 41.
68. As quoted in Ref. 41.
69. J. H. Kiefer and R. W. Lutz, "Vibrational Relaxation of Hydrogen," J. Chem. Phys. 44, 668-672 (1966).
70. M.-M. Audibert, C. Joffrin, and J. Ducuing, "Vibrational Relaxation in Hydrogen-Rare Gases Mixtures," Chem. Phys. Lett. 19, 26-28 (1973).

71. Quoted in J. B. Calvert, "Distorted Wave Calculation of Vibrational Excitation and Transfer in H_2 and D_2 ," J. Chem. Phys. 56, 5071-5074 (1972).
72. J. Ducuing and F. De Martini, "L'Effet Raman Stimulé et la Mesure de la Durée de Vie Vibratoire de l'Hydrogène," J. Chim. Phys. 64, 209-213 (1967).
73. R. F. Heidner III and J. V. V. Kasper, "An Experimental Rate Constant for $H + H_2(v \approx 1) \rightarrow H + H_2(v \approx 0)$," Chem. Phys. Lett. 15, 179-184 (1972).
74. W. Dale Breshears and Paul F. Bird, "Density Gradient Measurements of F_2 Dissociation Rates in Shock Waves," J. Chem. Phys. 58, 5176 (1973).
75. Discussion with W. D. Breshears and G. L. Schott, LASL.
76. V. H. Shui and J. P. Appleton, "Gas-Phase Recombination of Hydrogen. A Comparison between Theory and Experiment," J. Chem. Phys. 55, 3126-3132 (1971).
77. A. D. Stepukhovich and V. M. Umanskii, "Kinetics and Mechanism of Three-Body Recombination of Atoms and Radicals," Russ. Chem. Revs. 38, 590-607 (1969).
78. J. H. Sullivan and R. C. Feber, Los Alamos Scientific Laboratory, personal communication.
79. G. A. Kapralova, E. M. Margolina, and A. M. Chaikin, "Rate Constants for Some Elementary Stages of the Reaction of Fluorine with Hydrogen," Chem. Abstr. 78, 152006e (1973).
80. G. A. Kapralova, E. M. Margolina, and A. M. Chaikin, "The Effect of Fluorine, Hydrogen, or Inert Gases on the Upper Self-Ignition Limit in the Reaction of Fluorine with Hydrogen," Kinet. i Katal. (English translation) 10, 23-27 (1969).