Ladd 1697

AECU-2916

WS 44377

NUCLEAR SCATTERING OF HIGH ENERGY NEUTRONS*
AND THE OPTICAL MODEL OF THE NUCLEUS

47761

A Thesis
Presented to the Faculty of the Graduate School
of Cornell University for the degree of
Doctor of Philosophy

APPROVED FOR RELEASE
DATE <u>7-9-54</u>
For The Atomic Energy Commission
<u>William A. J. ...</u>
Chief, Declassification Branch <i>L</i>

by

Theodore Brewster Taylor

Los Alamos Scientific Laboratory

June, 1954

I

*For subsequent publication in the Physical Review ✓

072 002

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

TABLE OF CONTENTS

	PAGE
I. Introduction	1
II. A Brief History of the Optical Model of the Nucleus.	3
III. Description of the Optical Model and the Method of Solution of the Scattering Problem. . .	7
IV. Analysis of Experimental Data Above 70 Mev . . .	19
V. Discussion of Results.	26
VI. Analysis of Experimental Data Below 70 Mev . . .	34

II

072 002

I. INTRODUCTION

The primary purpose of this paper is to attempt to explain the energy dependence of the total cross sections of atomic nuclei for high energy neutrons by using the so-called "optical model" of the nucleus. Nuclei are represented by spheres of radius R within which there exist a nuclear potential V and a coefficient of neutron absorption K , and the scattering problem is solved in essentially the same manner as that which would be used for solving the problem of the scattering of light by a sphere having a complex index of refraction. There are three main steps in the analysis: 1) We inquire whether there exists a set of nuclear radii and a set of values of V and K which may vary smoothly with the energy of the incident neutron beam, but which are the same for all nuclei, such that we obtain agreement between calculated and observed total cross sections for high energy neutrons. It is found that this can be done. 2) We try to determine within what limits the empirically derived nuclear parameters are determined. 3) The nuclear radii obtained by this analysis are compared with those obtained by other methods, and an attempt is made to explain qualitatively the values of K and V which come from the analysis.

Since a rather considerable amount of literature dealing with the optical model of the nucleus has appeared in recent years, it has appeared useful to give a brief history of the model with references to articles that have appeared on the

subject. This will be followed by a somewhat detailed description of the model and the method which was used for solving the scattering problem. Finally the three stages of the analysis outlined above will be given.

II. A BRIEF HISTORY OF THE OPTICAL MODEL OF THE NUCLEUS

Prior to 1940 the interactions of beams of neutrons with nuclei were generally considered from the point of view of the theory of the compound nucleus. Emphasis was placed primarily on what happens after an incoming particle has been absorbed by a nucleus, rather than on the details of the probability of forming a compound nucleus in the first place. Total cross sections were usually expressed in terms of a "sticking probability" $\frac{\sigma}{\sigma_0}$, which was assumed to be equal to unity for high energy particles, and to be proportional to velocity for low energy particles, but there existed no detailed theory of the sticking probability.

In 1940 Bethe⁽¹⁾ gave a theory of the sticking probability. With energies above 100 Kev in mind it was assumed that nuclear energy levels were so dense that they could be considered as a continuum. Nuclei were represented by a potential well consisting of an ordinary real potential plus an imaginary potential, the latter of which can be directly related to a coefficient for absorption of neutrons in nuclear matter. The scattering problem was then set up in terms of the one-particle Schroedinger equation

$$\nabla^2 \psi + \frac{2M}{\hbar^2} (E - V_1(r) - i V_2(r)) \psi = 0 \quad (1)$$

where M is the mass of the incident particle and E is its energy. V_1 was assumed to be much smaller than V_2 , and an expression was derived for $\frac{\sigma}{\sigma_0}$ in terms of E and V_2 . Bethe also

obtained an expression for the absorption cross section of nuclei for very high neutron energies by assuming V_2 to have the form of a square well and by using the WKB approximation to solve the scattering problem. In the limit of very large V_2 , corresponding to a "black" nucleus, both the absorption cross section and the elastic "shadow scattering" cross section reduce to πR^2 , where R is the nuclear radius.

It was pointed out by Serber ⁽²⁾ in 1947 that at neutron energies of the order of 100 Mev nuclei should appear partially transparent to the neutrons because at that energy one would expect that the inelastic scattering mean free path for a neutron traversing nuclear matter would become comparable to the nuclear radius. In a paper by Fernbach, Serber, and the present author ⁽³⁾ it was shown that total cross section measurements for a variety of elements by Cook et al ⁽⁴⁾ for about 90 Mev neutrons could be satisfactorily accounted for by using the optical model of the nucleus. It was found that very close agreement with the measurements could be obtained by assuming nuclear radii equal to $1.37 A^{1/3}$ (A is the nuclear mass number.), a real potential equal to 30 Mev, and an absorption coefficient in nuclear matter equal to $2.2 \times 10^{12} \text{ cm}^{-1}$. A subsequent comparison between calculated and observed elastic scattering angular distributions by Bratenahl et al ⁽⁵⁾ for 84 Mev neutrons also showed reasonably good agreement with the predictions of the optical model.

By 1951 total neutron cross section measurements were extended up to 280 Mev ^{(6),(7),(8)}. It was found that the

rather rapid drop observed in the cross sections above 100 Mev could be only qualitatively obtained from the optical model by letting the effective potential of nuclei drop to zero for neutron energies above about 150 Mev, but that quantitative agreement, using the nuclear radii determined by the 90 Mev data, was impossible.

A compilation of data on the energy dependence of total neutron cross sections in the low energy region (1-3 Mev) by Barschall(9) showed striking regularities in the neutron cross sections of the elements. Maxima and minima appear in the cross sections of a given element as a function of energy, and the positions of these maxima and minima appear to vary smoothly with nuclear mass number. In an article by Feshbach, Porter and Weisskopf(10) calculations of low energy neutron cross sections using the optical model were described, and it was shown that a quantitative agreement with Barschall's data could be obtained by assuming nuclear radii equal to $1.4 A^{1/3} \times 10^{-13}$ cm., a real nuclear potential of 19 Mev, and an imaginary potential of 1 Mev. Since the quantity which determines the positions of the maxima and minima in the total cross sections is the product of the neutron wave number inside the nucleus and the nuclear radius, somewhat different choices of R and V will presumably also allow reasonably good fitting of the experimental data. It is understood by the author that at the present time these calculations are being extended toward somewhat higher energies, and that the effects of changing the spatial dependence of both the real and imaginary parts of the

nuclear potential well are being investigated.

Recent total neutron cross section measurements for a variety of nuclei for neutron energies between 30 and 150 Mev have been reported by Taylor and Wood⁽¹¹⁾, Linlor and Ragent⁽¹²⁾, and Ragent⁽¹³⁾. These data show definite maxima and minima in the cross sections as a function of neutron energy rather similar to those pointed out by Barschall for low energy neutrons. The analysis of these data and some reported at higher neutron energies by Nedzel⁽¹⁴⁾ (400 Mev) and Snow et al⁽¹⁵⁾ (1.4 Bev) is the main subject of this thesis.

III. DESCRIPTION OF THE OPTICAL MODEL AND THE METHOD OF SOLUTION OF THE SCATTERING PROBLEM

The optical model of the nucleus consists of a spherically symmetrical aggregate of material which we may call "nuclear matter," which is completely described by its ability to absorb and refract incident particles. In what follows we shall assume that the incident particles are neutrons; the model which we shall discuss may be assumed to apply equally well if the incident particles are protons if their energies are high enough so that Coulomb effects may be neglected.

We should clarify what we mean by refraction and absorption. By the former is meant an interaction between the incident particle and some effective real potential within nuclear matter such that there is no net transfer of energy between the particle and any internal degrees of freedom which nuclear matter may possess. This corresponds to scattering which is coherent. By absorption is meant any process which leaves the nucleus in an energy state different from its initial one, and thus includes any incoherent scattering.

It might be expected that the potential energy of a neutron traversing nuclear matter will vary greatly from place to place, depending on whether or not it passes very close to individual nucleons; we might picture the potential qualitatively as shown below:

One of the basic assumptions in the optical model is that it is reasonable to replace the "actual" potential by an average, smooth function of r .

A particular model of a nucleus is specified by giving the functional forms of the real potential, V , and the coefficient for particle absorption, K , as a function of the distance from the center of the nucleus. In general, we may expect that the real potential and the absorption coefficient will depend upon the energy of the incident neutron. As the wave length of the incident neutron decreases it can explore more of the details of the shape of the nuclear potential, so that the average potential will appear to change. A phenomenological study of neutron-nuclear interactions could be made, in principle, by assuming various forms of $V(r,E)$ and $K(r,E)$ and then comparing calculated and observed elastic and inelastic scattering cross sections and elastic scattering angular distributions. In order to get some idea of the complexity of the problem, let us consider the scattering of 100 Mev neutrons by lead nuclei. The wave number k of a 100 Mev neutron is about $2 \times 10^{13} \text{ cm}^{-1}$. If we were to solve the problem by the method of partial waves, we should expect that the number of important phase shifts would be of the order of k times the distance at which the nuclear potential of lead drops to a negligible value. Thus, assuming a "radius" for lead of about $8 \times 10^{-13} \text{ cm}$., we see that we would have to calculate of the order of 16 phase shifts. It is clear, then, that the kind of study we have suggested could be carried out only by finding some fairly

simple approximate way of solving the scattering problem, or by using high speed computing machines. Furthermore, it is highly questionable whether there exist enough experimental data on the scattering of high energy neutrons to warrant so detailed an investigation.

It is possible to solve the scattering problem analytically in the special case that V and K are taken to be constant out to some distance R , and to vanish elsewhere; the scattering corresponds to that of scattering by a complex square well, and the phase shifts for each l may be expressed in terms of spherical Bessel functions of complex argument⁽¹⁶⁾. This is essentially what was done by Feshbach, Porter, and Weisskopf⁽¹⁰⁾. These exact calculations, however, become too laborious for neutron energies greater than a few Mev to be practicable without the aid of high speed computing machines.

There appears to be enough interesting detail in the energy variation of neutron-nuclear cross sections above about 50 Mev to warrant an analysis of the experimental data by means of a method which is only applicable for rather high energy neutrons, but which is simple enough to involve only a moderate amount of numerical calculation. The method which was chosen for this purpose is that described by Fernbach, Serber, and Taylor⁽³⁾, in which the main simplification comes from replacing the sum over discrete partial waves by an integral over a continuous impact parameter. This method should be applicable to high energy scattering even if the real and imaginary parts of the potential are not assumed to have the form of a square well;

calculations of the effect of decreasing the sharpness of the nuclear boundary, however, have been left for the future. A preliminary look at the approximate solution of the problem of scattering by a non-square well shows that a detailed analysis of the existing scattering data would probably have to be performed with the use of a high speed computing machine. We may consider the restriction to a square well as a first step in a program to test the applicability of the optical model.

The problem is that of the scattering of a neutron wave by a sphere of material characterized by a radius R , and absorption coefficient K , and an index of refraction. The index of refraction is determined by the mean potential energy, $-V$, of the incident neutron when it is inside the nucleus. It is convenient to introduce a quantity k_1 , which is the change in the wave number of a neutron as it enters the nucleus. k_1 may be related to the incident neutron kinetic energy, E , and the nuclear potential V as follows:

$$k_1 = k \left[\left(1 + \frac{V}{E} \right)^{1/2} - 1 \right] \quad (2)$$

where k , the wave number of the incident neutron, is equal to $\sqrt{2ME}/\hbar$. Equation (2) is valid only for non-relativistic neutron energies. A relativistic expression for k_1 which is valid for any neutron energies is

$$k_1 = \frac{Mc}{\hbar} \frac{\left[\frac{2V}{Mc^2} + \frac{2VE}{M^2c^4} + \frac{V^2}{M^2c^4} \right]}{\sqrt{\frac{2(E+V)}{Mc^2} + \frac{(E+V)^2}{M^2c^4}} + \sqrt{\frac{2E}{Mc^2} + \frac{E^2}{M^2c^4}}} \quad (3)$$

We now make the following assumptions:

1) $kR \gg 1$. This is analogous to the assumption made in geometrical optics that the wave length of light incident upon an obstacle is small compared to the dimensions of the obstacle; the assumption allows us to consider the classical trajectories of neutrons traversing the nucleus.

2) k_1 and $K \ll k$. This condition is required if we are to neglect the reflection of the neutron wave at the surface of the nucleus and the change in direction of the neutron propagation vector as it enters the nucleus. The order of magnitude of the error made by this assumption may be gotten from an analogy with optics, for which the reflection coefficient for normal incidence would be equal to

$$\left(\frac{\frac{k+k_1}{k} - 1}{\frac{k+k_1}{k} + 1} \right)^2 = \left(\frac{k_1}{2k+k_1} \right)^2$$

Thus, for a neutron energy of 100 Mev and a potential of 30 Mev, $k = 2 \times 10^{13} \text{ cm.}^{-1}$, $k_1 = .3 \times 10^{13} \text{ cm.}^{-1}$, and the reflection coefficient is equal to about .005. It should be noted that the neglect of reflection at the boundary is justified further if it is assumed that the potential and the absorption coefficient drop to zero in a distance of the order of 10^{-13} cm. , rather than perfectly sharply, because the values of λ for the neutrons which we shall consider are less than 10^{-13} cm. Thus, in this particular sense, our method of calculation assumes a lack of perfect sharpness in the nuclear boundary.

A qualitative description of the scattering may be given as follows: Consider the incident neutrons as plane waves striking a sphere of radius R . Part of the neutron wave will be absorbed, with the absorption cross section given by πR^2 times the probability that a neutron striking the nucleus will be inelastically scattered. The disturbance of the incident plane waves created by this absorption will give rise to "diffraction scattering," analogous to the diffraction of light by an obstacle. In the limit of completely opaque nuclei, both the absorption and the diffraction cross section will be equal to πR^2 . If the nucleus is partly transparent, however, that part of the neutron wave which is transmitted will emerge with a phase different from that of the neutron beam as a whole, with the result that the diffraction scattering will be complicated by further interference effects.

The solution to the scattering problem is straightforward. Consider the portion of the incident neutron wave which strikes the nucleus at a distance ρ from the center. It will emerge after travelling a distance $2s$, with $s^2 = R^2 - \rho^2$. The amplitude of the neutron wave after emerging is $a = \exp(-K + 2ik_1)s$, where K is defined in such a way that the intensity of the neutron wave after travelling a distance $2s$ is equal to $|a|^2$, or e^{-Ks} . Thus, on a plane behind the sphere, the wave differs from a plane wave by an amplitude $1 - a$. The absorption cross section is now calculated as

$$\begin{aligned}
\sigma_a &= 2\pi \int_0^R (1 - |a|^2) \rho d\rho = 2\pi \int_0^R (1 - e^{-2\kappa\rho}) \rho d\rho \\
&= 2\pi \int_0^R (1 - e^{-2\kappa\rho}) \rho d\rho \\
&= \pi R^2 \left\{ 1 - \frac{[1 - (1+2x)e^{-2x}]}{2x^2} \right\}, \quad (4)
\end{aligned}$$

where $x = \kappa R$. Note that σ_a approaches πR^2 as x approaches infinity.

Similarly the diffraction cross section will be given by

$$\begin{aligned}
\sigma_d &= 2\pi \int_0^R |1 - a|^2 \rho d\rho = 2\pi \int_0^R |1 - e^{(-\kappa + 2i k_1)\rho}|^2 \rho d\rho \\
&= 2\pi \int_0^R |1 - e^{(-\kappa + 2i k_1)\rho}|^2 \rho d\rho \\
&= \pi R^2 \left[1 + \frac{1}{2x^2} \left\{ 1 - (1+2x)e^{-2x} \right\} \right. \\
&\quad \left. - \frac{1}{\left(\frac{x^2}{4} + y^2\right)^2} \left(\frac{x^2}{4} - y^2 + e^{-x} \left\{ [2y\left(\frac{x^2}{4} + y^2\right) + xy] \sin 2y \right. \right. \right. \\
&\quad \left. \left. \left. - \left[\frac{x^2}{4} - y^2 + x\left(\frac{x^2}{4} + y^2\right) \right] \cos 2y \right\} \right) \right] \right], \quad (5)
\end{aligned}$$

where $x = \kappa R$ and $y = k_1 R$. σ_d also approaches πR^2 as x approaches infinity.

When x is equal to zero (5) becomes

$$\sigma_d = \pi R^2 \left[2 - \frac{1}{y^2} (2y \sin 2y + \cos 2y - 1) \right]. \quad (6)$$

When y is equal to zero (5) becomes

$$\sigma_d = \pi R^2 \left\{ 1 + \frac{1}{2x^2} [1 - (1+2x)e^{-2x}] - \frac{4}{x^2} [1 - (1+x)e^{-x}] \right\}. \quad (7)$$

The total cross section is simply the sum of the absorption and diffraction cross sections:

$$\sigma_t = \sigma_d + \sigma_a. \quad (8)$$

The angular distribution of the elastically scattered neutrons may be calculated in the same way as that used for calculating the Fraunhofer diffraction pattern for a circular obstacle. The scattered amplitude at an angle θ to the incident beam is found by adding up the contributions from spherical wavelets originating from a disk in the shadow of the sphere. In this way the scattered amplitude is easily shown to be

$$f(\theta) = k \int_0^R [1 - a] J_0(k\rho \sin \theta) \rho d\rho$$

$$= k \int_0^R [1 - e^{(-k + 2ik_1)\rho}] J_0(k\rho \sin \theta) \rho d\rho. \quad (9)$$

In the limit of a completely opaque nucleus ($KR \rightarrow \infty$), this becomes the well known expression

$$f(\theta) = \frac{kR^2}{2} \left[\frac{J_1(2kR \sin \frac{\theta}{2})}{kR \sin \frac{\theta}{2}} \right]. \quad (10)$$

The amplitude for scattering in the forward direction is easily evaluated and is found to be

$$f(0) = \frac{kR^2}{2} \left\{ 1 + \frac{(y - i\frac{x}{2})^2}{2(\frac{x^2}{4} + y^2)^2} [1 - (1 + x - 2iy) e^{-x + 2iy}] \right\}. \quad (11)$$

In order to obtain the scattering cross section in the forward direction, $f(0)$ must be broken up into its real and imaginary parts. A convenient way to do this is to go back to the expression (9) for $f(\theta)$ and substitute $\theta = 0$ before performing the integration over ρ . The integrand is then broken up into its real and imaginary parts to give

$$f(0) = \frac{kR^2}{2} - \frac{kR^2}{4(k_1 R)^2} \int_0^{2k_1 R} e^{-\frac{k}{2k_1} x} (\cos x + i \sin x) x dx.$$

The two integrals on the right hand side can easily be done by using the relations

$$\int x e^{-ax} \cos x dx = -\frac{d}{da} \int e^{-ax} \cos x dx$$

$$\int x e^{-ax} \sin x dx = -\frac{d}{da} \int e^{-ax} \sin x dx$$

where the integrals on the right hand side are given in standard tables of integrals.

The final expression for the scattering in the forward direction is given by

$$\left(\frac{d\sigma}{d\Omega} \right)_{\theta=0} = \frac{k^2 R^4}{4} \left[\left(1 - \frac{\bar{X}^2}{2y^2} \right)^2 + \frac{\bar{Y}^2}{4y^4} \right], \quad (12)$$

where

$$\bar{X} = \frac{e^{-2ya}}{(a^2+1)^2} \left[-(a^2+2ya^3-1+2ya) \cos 2y + (2ya^2+2y+2a) \sin 2y \right]$$

$$+ \frac{a^2-1}{(a^2+1)^2} \quad a = \frac{1K}{2k_1}$$

and

$$\bar{Y} = \frac{e^{-2ya}}{(a^2+1)^2} \left[-(a^2+2ya^3-1+2ya) \sin 2y - (2ya^2+2y+2a) \cos 2y \right]$$

$$+ \frac{2a}{(a^2+1)^2}$$

In the limit of $k_1 = 0$ we have

$$f(0) = \frac{kR^2}{2} \left\{ 1 - \frac{2[1 - (1+x)e^{-KR}]}{x^2} \right\} \quad (13)$$

$\frac{\sigma_a}{\pi R^2}$ is plotted as a function of KR in Fig. 1. It can be seen that $\frac{\sigma_a}{\pi R^2}$ approaches 1 for very large values of KR , and that it is greater than 0.9 for KR greater than 2. Thus it is to be expected that transparency effects will become important when the mean free path in nuclear matter is equal to about one half the nuclear radius. It should also be noted that if KR is greater than about 2, measurements of the absorption cross sections do not provide an accurate means for determining K .

In Fig. 2 we have plotted $\frac{\sigma_t}{\pi R^2}$ as a function of $k_1 R$ for various values of k_1/K . A qualitative understanding of these curves can be obtained by considering the simple problem of the scattering from a disk of radius R and thickness T . The absorption, scattering, and total cross sections are given by

$$\sigma_a = \pi R^2 (1 - |a|^2) = \pi R^2 (1 - e^{-KT})$$

$$\sigma_d = \pi R^2 |1 - a|^2 = \pi R^2 (1 - 2e^{-\frac{KT}{2}} \cos k_1 T + e^{-KT})$$

$$\sigma_t = \pi R^2 (2 - 2e^{-\frac{KT}{2}} \cos k_1 T).$$

If $k_1 T$ is very small, the phase change in the transmitted wave is small, and the total cross section is primarily determined by the value of KT . When $k_1 T$ is increased to the value $\frac{\pi}{2}$, the total cross section reaches the value $2\pi R^2$, and is independent of KT . When $k_1 T$ is further increased to the value π , there is constructive interference between the transmitted wave and the rest of the beam; the excess of σ_t over $2\pi R^2$ will be determined by the value of KT . For still higher values of $k_1 T$ a curve of $\frac{\sigma_t}{\pi R^2}$ as a function of $k_1 T$ will oscillate about the

value 2 with a period of 2π , and the relative heights of the maxima and minima will depend upon the values of KR .

The curves in Fig. 2 show these same qualitative features; since, however, the phase change for a particular ray through the nucleus is a function of the impact parameter, the first maximum occurs when $k_1 R$ is equal to 2, rather than when $k_1 T = \pi$ and, when $k_1 R$ is equal to one, $\frac{\sigma_T}{\pi R^2}$ is only approximately independent of KR , and is somewhat smaller than 2.0.

For a given value of V , large values of k_1 correspond to low energy, so that, for a given element, one expects that as neutron energies are decreased from some high value the cross section will rise to a maximum when $k_1 R$ is equal to 2, decrease again to some minimum value when the interference is destructive, and continue to go through successive maxima and minima until the neutron energy goes to zero. The cross section will oscillate about the value $2\pi R^2$ only if λ of the neutron is small compared to the nuclear radius; when the neutron energy is so low that this is no longer true, then it is to be expected that the method we have used for solving the scattering problem will generally give too small cross sections. There are two effects that tend to damp out the oscillations of σ_T about the value $2\pi R^2$ as $k_1 R$ gets very large. If k_1/K is kept constant, then KR increases as $k_1 R$ increases, and the intensity of the transmitted wave causing the interference effects decreases. Furthermore, as the phase change of the transmitted wave becomes very large, the phase change for a particular ray through the nucleus becomes a sensitive function of impact parameter, and integration over the impact parameter cancels out the interference effects.

Fig. 3 is a plot of $\frac{V_t}{\pi R^2}$ as a function of KR for various values of k_1/K . This is a convenient way of plotting equation (5) when we are interested in energies for which k_1 is rather small.

The author has been unable to find a convenient form for the integral in equation (9) for arbitrary values of the nuclear parameters and of θ . Because of the large numbers of parameters involved it has not seemed worthwhile to obtain a set of curves of $f(\theta)$ by numerical integration of equation (9). Calculations by Bratenahl et al (5) indicate that, even if the nuclear transparency is rather high, the angular distributions calculated from equation (9) are not very different from those obtained from equation (10), which is valid for completely opaque nuclei; the angular widths at half maximum are practically identical, and the only significant differences occur at very small or very large angles.

IV. ANALYSIS OF EXPERIMENTAL DATA ABOVE 70 MEV

We shall first attempt to answer question 1 in the Introduction. For the sake of generality in the analysis of the experimental data we shall assume no previous knowledge of any of the nuclear parameters, except for their orders of magnitude. Attention will first be confined only to the energy variation of the total cross sections; having found a set of nuclear parameters which will allow us to fit the total cross section data, we shall then compare predicted and observed values of a number of other quantities which have been observed experimentally. The experimental data which will be analyzed are those of Taylor and Wood⁽¹¹⁾, Linlor and Ragent⁽¹²⁾, Ragent⁽¹³⁾, and Nedzel⁽¹⁴⁾. All of these data have been obtained within the last two years by methods which permitted accurate knowledge of the incident neutron energy to within a few Mev, and for which the statistical accuracy in measurement is a few percent.

Throughout the analysis we shall make the following assumptions, besides those listed in the discussion of the method of solution of the scattering problem:

1. Both K and V , for a particular neutron energy, are the same for all nuclei. This implies nucleon densities which are independent of nuclear mass number, and after the analysis of the experimental data we must check for consistency between the nuclear radii which are determined and the assumption of constant nucleon density.

2. The effective radius of a given nucleus is independent of the energy of the bombarding neutron. It is conceivable that changes in the opacity of the nuclear matter near the edges of nuclei might have the effect of making the effective radii appear to change slightly with the incident energy. The neglect of this effect is partly justified by the result that K does not appear to vary greatly with energy over the energy range considered here.

3. Both K and V are smooth functions of the bombarding energy. A reason for expecting V to depend upon the bombarding energy is given on p. 8. K would be expected to vary with energy because it is presumably closely connected with the free n - p and n - n cross sections, both of which are energy dependent. (We assume equality of the p - p and n - n interactions.)

The analysis of the experimental data proceeds as follows:

1) We assume that the observed maximum in the lead cross section as a function of energy is the first interference maximum, corresponding to $k_1 R = 2.0$. This maximum occurs at an energy of about 85 Mev. Then, for several assumed values of $\gamma_{85} = k_1/K$ at 85 Mev, we use Fig. 2 to determine corresponding values of the radius of lead which will give the measured value of the lead cross section. Thus, for each assumed value of γ_{85} we have values of k_1 , K , and R_{pb} and, again using Fig. 2 and the measured cross sections of the other elements at 85 Mev, we can obtain a set of nuclear radii for the other elements.

2) We note that the curves in Fig. 2 are such that, for $k_1 R$ equal to 1.05, $\frac{\sigma_t}{\pi R^2}$ is essentially independent of k_1/K and is equal to 1.75. Hence, by obtaining from the experimental data the energy for which $\sigma_t = 1.75\pi R^2$ for each element, using the radii determined in step (1), we can obtain a curve of k_1 as a function of energy. Note that we obtain one curve for each assumed value of γ_{85} .

3) K is determined at each energy for which k_1 has been determined by using the measured values of the total cross sections of any one of the elements.

4) We now have obtained, for each choice of γ_{85} , a set of nuclear radii and curves of k_1 and K as a function of energy. As we shall see presently, the range of neutron energies which has been covered so far is from about 70 Mev to about 150 Mev, which includes most of the Harwell and Berkeley data. We now compute the detailed energy variation of the total cross sections for all of the nuclei for which measurements exist, and compare the results with the experimental data.

The above procedure was followed through for four different choices of γ_{85} : 1.0, 1.25, 1.5, and 2.0. In all cases smooth curves of k_1 as a function of energy and R as a function of $A^{1/3}$ were obtained. In order to give some idea of the sensitivity of the curves to the choice of γ_{85} , these curves are shown in Figs. 4 and 5. It will be noted that the least well determined parameter is K , while the nuclear radii are rather insensitive to the choice of γ_{85} .

A detailed comparison between observed and calculated cross sections for all the elements showed that the best agreement was obtained by using the set of nuclear parameters corresponding to the choice of γ_{85} of 1.25, and that 1.0 and 2.0 are ruled out if overall agreement between calculated and observed cross sections within quoted experimental errors is insisted upon.

Since there is no sharp dividing line between nuclear parameters which will and which will not allow a close fit to the experimental data over the entire energy range considered, it is difficult to say precisely how well the nuclear parameters are determined by the data. The parameters corresponding to the choice of γ_{85} of 1.25 seem to give the best fit. A general idea of the uniqueness of the derived parameters can be obtained from the following limits, within which a reasonably good fit to the data is possible: Nuclear radii, $\pm$ about 3%. k_1 at 85 Mev = $(.238 \pm .02) \times 10^{13} \text{ cm}^{-1}$. K at 85 Mev = $(.18 \pm .04) \times 10^{13} \text{ cm}^{-1}$.

The total cross sections obtained by Nedzel⁽¹³⁾ at 410 Mev were analyzed as follows: Using the nuclear radii determined by the lower energy data, a search was made for values of k_1 and K which would give agreement between calculated and observed cross sections. It was found that agreement within experimental error for all the cross sections measured (excluding that of hydrogen!) could be obtained by choosing k_1 equal to $.067 \times 10^{13} \text{ cm}^{-1}$ and K equal to $0.18 \times 10^{13} \text{ cm}^{-1}$, and that insistence

on a good fit to the experimental data determined k_1 to within about 5% and K to within about 20%. It is interesting that the value of V determined by the 410 Mev data is 10 Mev, which is only slightly smaller than V at 150 Mev.

By means of the relations (2) and (3) the values of k_1 were converted to values of V at each neutron energy. The curves of k_1 , V , and K as a function of energy were joined smoothly to the values determined at 400 Mev; the resulting curves are shown in Figs. 6 and 7. The parts of the curves extending below 70 Mev will be discussed in Chapter VI. Comparisons between the calculated and observed total cross sections are shown in Figs. 8 and 9, and, for a neutron energy of 410 Mev, in Table I. The radii of the lightest and heaviest elements included in Table I were obtained by linear extrapolation of the curve of R as a function of $A^{1/3}$ obtained from the analysis of the experimental data of Taylor and Wood.

There are a number of measurable quantities besides the total cross sections which can be used as further tests of the validity of the optical model:

- 1) Total inelastic (absorption) cross sections. Since the absorption cross sections are independent of the value of V , measurements of these provide a much more sensitive determination of K than the total cross sections do. "Poor" geometry experiments for several elements have been reported by DeJuren and Knable⁽¹⁷⁾ for 95 Mev neutrons and by DeJuren⁽⁷⁾ for 270 Mev neutrons. In both cases the neutrons in the incident beam contained a rather broad band of energies, and the use of

threshold detectors with thresholds considerably lower than the mean incident neutron energy meant that inelastic collisions involving small energy transfers were not detected. The ratios $\frac{\sigma_a}{\sigma_t}$ rather than σ_a are given by these authors. Since the total cross sections for some of the elements vary rather rapidly with energy, particularly in the region of 95 Mev, we have made a comparison between calculated values of σ_a and values of σ_a obtained by multiplying the experimental values of $\frac{\sigma_a}{\sigma_t}$ by the values of σ_t measured by the same authors, i.e., with a rather broad energy spread in the incident neutron beam. Note that the approximate constancy of K with energy implies absorption cross sections which vary little with energy in this energy range.

It was found that, by choosing $K = 2 \times 10^{13} \text{ cm.}^{-1}$, it was possible to obtain agreement between calculated and observed values of the absorption cross sections of a number of elements for 1.4 Bev incident neutrons⁽¹⁴⁾. These comparisons are also shown in Table II.

2) Angular distributions of elastically scattered neutrons. Because of its dependence on the 4th power of the nuclear radius, the elastic scattering cross section in the forward direction would appear to be the best measurement for determining nuclear radii by means of neutron-nuclear scattering. The forward scattering cross section also depends rather strongly on both K and V, however, so that this cross section becomes of particular value only if taken together with other experimental data at the same energy. Angular distributions of elastically

scattered "85" Mev neutrons have been reported and analyzed by Bratenahl et al⁽⁵⁾. Since the spread in incident neutron energies in the experiment covered an energy band within which V apparently changes very rapidly with energy, a detailed analysis of the experiments has not been made by this author; the low energy part of the incident neutron spectrum includes energies for which the applicability of the calculation we have described becomes highly questionable. As is shown in Reference (5), the angular distributions can be fitted fairly well by using either the opaque nucleus model or a somewhat transparent model compatible with the total cross section measurements. Since the different choice of constants for the transparent model which we have obtained in the present analysis will presumably affect the scattering in the forward direction more than at other angles, we include in Table III a comparison between calculated values of the cross section in the forward direction and those obtained by extrapolating the measured angular distributions to zero angle. Because of the beam energy spread just mentioned, it seems likely that the close agreement is fortuitous.

V. DISCUSSION OF RESULTS

The overall conclusion that can be drawn from the analysis just described is that virtually all of the existing experimental data on the scattering of high energy neutrons can be quantitatively explained by using a simple form of the optical model of the nucleus and a small number of adjustable nuclear parameters. We shall now discuss some of the specific results.

1) Nuclear radii. The "best fit" nuclear radii are plotted as a function of $A^{1/3}$ in Fig. 10. Although the curve is not exactly a straight line, it can be well approximated by the line $R = (0.6 + 1.23 A^{1/3}) \times 10^{-13}$ cm. Before comparing these values of nuclear radii with those obtained by other methods we shall discuss the meaning of a "nuclear radius" as it applies to the analysis described in this paper. Let us suppose, for a moment, that the density distribution of nuclear matter in nuclei is such that it is constant out to some radius, and then drops to zero in a distance which is small compared to the radius itself. Then the quantity which we are presumably "measuring" in neutron scattering experiments is the radius of a volume within which there is an interaction between an incident neutron and the nucleus. Because of the finite range of nuclear forces, we should then expect that this radius will be somewhat larger than the radius of the envelope of the scattering centers of which nuclear matter is composed. Thus we may interpret the radius we have determined in this way:

The constant term in the expression for R as a function of $A^{1/3}$, a number of the order of the range of nuclear forces, corresponds to the distance from the edge of the nucleus within which a neutron must pass in order that it be subjected to a strong interaction, while the term proportional to $A^{1/3}$ corresponds to the "true" radius of the nucleus, i.e., the radius of the envelope of the scattering centers themselves. Now, as we have said before, there are reasons for believing that nuclei do not have a sharp boundary; the density of nucleons may be a nearly constant function of radius out to some radius beyond which there is a more or less rapid fall off to zero in a distance of the order of the range of nuclear forces. The author is unable to make any quantitative statements about the effect of assuming such a density distribution; however, because it has been possible to account satisfactorily for a considerable amount of scattering data by assuming that a well defined radius exists, it is believed that it would be impossible to choose between various possible density distributions differing from the square one only near the nuclear boundary by using existing neutron scattering data at high energies alone. It seems reasonable to assume that we have determined a sort of mean value of the radius, and that other scattering data than those which we have used must be used for giving more details about the density distribution.

We see that we may justify the assumption that, for a given neutron energy, K is the same for all nuclei, if we accept the above interpretation of the nuclear radii derived from the

experiments; our interpretation implies the same nucleon density for all nuclei. We shall discuss this question more fully when we discuss the energy dependence of K.

We now compare the nuclear radii which we have obtained with those obtained by other methods.

A) Scattering of high energy electrons by nuclei. Recent experiments by Hofstadter et al⁽¹⁸⁾ on the angular distribution of high energy electrons elastically scattered by nuclei indicate that, if the protons are assumed to be uniformly distributed throughout the nucleus, the nuclear radii must be assumed to be equal to about $1.2 \times A^{1/3} \times 10^{-13}$ cm⁽¹⁹⁾. Since what is being "measured" in these experiments is the radius of the proton envelope, the results are in good agreement with those obtained from neutron scattering, provided that we assume that the "proton radius" is the same as the "neutron radius." It has been pointed out by Johnson and Teller⁽²⁰⁾ that one should expect the former to be somewhat smaller than the latter because of the higher potential energy of the protons due to electrostatic repulsion, but the author is not aware of any calculations of this effect. A detailed analysis of the electron scattering data⁽¹⁹⁾⁽²¹⁾ indicates that a best fit to the data is obtained by smearing out the proton "edge" over a distance of the order of 10^{-13} cm.

B) Comparisons of energies of mirror nuclei. Recent calculations by Cooper and Henley⁽²²⁾ show that the "proton radii" of mirror nuclei agree very well with those obtained from

electron scattering, and, therefore, with the term proportional to $A^{1/3}$ in our radius formula.

C) Lifetimes of α -radioactive nuclei. The radii of heavy nuclei determined by this method scatter about values somewhat larger than those obtained by method (A), and are compatible with those obtained from neutron scattering. We assert that this is to be expected, since the radii obtained from α -decay lifetimes are those of volumes within which there is a strong nuclear interaction. Questions concerning the "size" of α -particles introduce some uncertainty about the meaning of nuclear radii determined by this method.

2) The absorption coefficient of nuclear matter for neutrons. It is of interest to compare the empirically derived energy dependence of K with that which one might infer from the measured nucleon-nucleon cross sections. Calculations have been done by Goldberger⁽²³⁾, in which nuclear matter is considered to consist of a mixture of two non-interacting Fermi gases of neutrons and protons bound in a uniform potential well. The exclusion principle was taken into account by demanding that the incident as well as the target nucleon be outside the occupied Fermi momentum sphere after the scattering has taken place. It was found that, assuming that the nucleon-nucleon scattering is isotropic in the center of mass system and is independent of the relative momenta of the two particles for the range of momenta existing in the Fermi gas, the effective cross section $\bar{\sigma}$ of the nucleons bound in nuclear matter may be written

$$\bar{\sigma} = \sigma(E_n) \left(1 - \frac{1}{5} \frac{E_F}{E_n} \right), \quad (14)$$

where E_F is the Fermi energy, E_n is the energy of the neutron inside the nucleus, and $\sigma(E_n)$ is the free particle nucleon-nucleon cross section. The expression is valid for $E_n \geq 2E_F$. According to the well known formula,

$$E_F = \frac{\hbar^2}{2m} \left(\frac{3\pi^2 N}{V} \right)^{2/3} = 1.98 \times 10^{-24} n^{2/3} \text{ Mev.} \quad (15)$$

where n is the number of nucleons per cubic centimeter. Now, the Fermi energy is a quantity generally ascribed to systems consisting of a very large number of identical Fermions in a volume which is large compared to the average volume per particle. In our case, the nuclear radius is not orders of magnitude larger than either the inter-nucleon distances or the range of nuclear forces, and so there is bound to be some ambiguity in defining, among other things, the Fermi energy. In particular, it is not clear whether nuclear volumes should be computed using what we have called the "true" nuclear radius or the larger, effective radius determined from neutron scattering; the difference between the two will lead to considerable differences in the computed Fermi energy. It was decided to use the larger radii, since these, as we have argued before, correspond to the radius of the potential well as seen by a probe particle. We now compute the proton and neutron Fermi energies for lead. Using $R = 8.1 \times 10^{-13}$ cm., $Z = 82$, and $N = 125$, we obtain 29 Mev for the neutron Fermi energy and 22 Mev for the proton Fermi energy. (These numbers would have been 38 Mev

and 29 Mev if we had used the smaller radii equal to $1.2 A^{1/3} \times 10^{-13}$ cm.) The average Coulomb energy of a proton in lead, assuming a radius of 7.1×10^{-13} cm., and a uniform distribution of protons is 10 Mev, which is close enough to the difference between the neutron and proton Fermi energies to assume that it accounts for it.

We can now calculate K as a function of neutron energy for lead by using (14), a Fermi energy of 29 Mev, and the measured n-p and p-p total cross sections given in Fig. 11, with K given by

$$K = 3 [Z \bar{\sigma}_{n-p}(E_n) + (A-Z) \bar{\sigma}_{p-p}(E_n)] / 4\pi R^3, \quad (16)$$

where E_n is the incident neutron energy plus the value of the nuclear potential obtained from Fig. 7. We are again faced by an ambiguity in the choice of R. It is not necessarily the same value of the radius as that used in computing the Fermi energy; but, since we are assuming that K is constant out to the large radius, it seems most reasonable to use the larger radius in equation (16). The curve of K as a function of energy which is obtained in this way is compared with the empirically derived K curve in Fig. 7. The shapes of the two curves are very nearly the same, but the computed K curve lies somewhat higher. Because of the ambiguities involved in calculating both the Fermi energy and the nucleon density from which K is obtained, the difference in magnitudes of the computed and "observed" values of K is not considered to be

significant. It is interesting, however, that both curves show an essentially constant value of K over a very wide energy range, in spite of the rapid decrease of the nucleon-nucleon cross sections with energy.

Although proton-proton scattering is isotropic over the energy range considered here, the neutron-proton angular distributions are peaked in both the forward and backward directions in the center of mass system, whereas we have assumed isotropy for both types of scattering. Since small angle scattering goes with small energy transfers, it should be depressed by the exclusion principle, and we should expect a further reduction in K over that which we have calculated. Calculations by Goldberger (loc. cit.) indicate that the use of a more realistic angular distribution for neutron-proton scattering further decreases the bound neutron-proton cross section by about 10%, assuming an incident neutron energy of 90 Mev and a Fermi energy of 18 Mev. The protons in lead contribute to about one half of the value of K ; since we have used a larger value of the Fermi energy than Goldberger did, we might expect that if we take into account the anisotropy of neutron-proton scattering at 90 Mev, K will be lowered by about 10%.

3) The energy dependence of the nuclear potential V . If the nucleon-nucleon interaction becomes highly repulsive at very short distances, than one might expect that, as the probe neutron wave length becomes shorter and more details of the nucleon-nucleon interaction are explored, the effective nuclear potential will decrease. The very rapid drop in V above 70 Mev,

and the approach to a constant value at about 200 Mev is left unexplained, however. It may be possible to obtain some idea of the expected energy dependence of V by using the nucleon-nucleon forward scattering amplitudes as a function of energy, as suggested by Jastrow⁽²⁴⁾. It is interesting to note that an analysis by Mandl and Skyrme⁽²⁵⁾ of the total cross sections for neutron energies up to 150 Mev gives a "best fit" curve of V as a function of energy which is very nearly the same above 70 Mev as that shown in Fig. 6, in spite of the facts that the nuclear radii were taken to be equal to $1.37 A^{1/3} \times 10^{-13}$ cm., which are considerably smaller than the radii which we have obtained for the light elements, and considerably larger values of K were used.

VI. ANALYSIS OF EXPERIMENTAL DATA BELOW 70 MEV

Figs. 12 and 13 show the detailed energy dependence of the measured total cross sections of a number of elements for neutron energies in the vicinity of 70 Mev. It will be noted that the cross sections of the heavy elements show a very sharp increase in cross section from 60 to 70 Mev, while the cross section of iron shows no such sudden change in cross section near these energies. This interesting behavior of the total cross sections below 70 made it appear worthwhile to extend the analysis of the cross sections to energies below 70 Mev, in spite of the fact that the assumption we have made in our calculations that kR is very large compared to one can no longer apply to the light elements at these low energies.

For an incident neutron energy of 40 Mev, the values of kR for various elements are as follows: lead, 11.5; iron, 3.0; carbon, 4.9. Thus, at 40 Mev we might expect the geometrical optics approximation to hold reasonably well for elements as light as iron, but the approximation certainly becomes highly questionable for the very light elements. The neglect of reflection at the nuclear boundary is not justified at an energy as low as 40 Mev, which is comparable to the depth of the potential well; assuming that V is about 30 Mev for an incident neutron energy of 40 Mev, the coefficient of reflection for normal incidence is of the order of 1%. This is somewhat misleading, however, because the reflection coefficient is much

larger for rays striking the sphere with large impact parameters. We may again partly justify our neglecting reflection at the nuclear boundary if we assume that the nucleon density does not suddenly drop to zero near the boundary. It should be noted, however, that, if the nucleon density drops from its constant value to zero in a distance of about 10^{-13} cm., λ for a 40 Mev neutron is of the same order, so that there will be some reflection even if the nuclear boundary is not well defined. It is believed by the author that the gradual breakdown of the approximate method which has been used for solving the scattering problem as lower energies are considered will not affect the qualitative results of the analysis which is described below.

The procedure which was used in this analysis was again to search for values of V and K as a function of energy which allow a close fit to the experimental data. If we first consider the data for lead, bismuth, and uranium alone, the sudden drop in cross section from 70 Mev to 60 Mev can be accounted for by assuming that V , and therefore k_1 , or K , or both, are very much higher for neutron energies of 60 Mev than 70 Mev. A sudden increase in $k_1 R$ for lead from its value at 70 Mev lowers the cross section by moving off the first interference maximum (See Fig. 2), while a sudden increase in the value of $K R$ removes the transmitted wave and destroys most of the constructive interference, thereby also lowering the cross section. If we now consider the experimental data for iron, however, a sudden change in K between 60 and 70 Mev is ruled out, because

there appears to be no substantial change in the cross section between 60 and 70 Mev; since k_1R for iron is about 1.7 at 70 Mev, it can be seen from Fig. 2 that a sudden increase in K would lower the cross section by about the same factor as for lead. k_1R for iron, on the other hand, can change upward from its value at 70 Mev by as much as 50% without any important change in cross section. Thus the only way to extrapolate our nuclear parameters to 60 Mev to obtain agreement with experiment is to make V rise sharply, but to make a small change, if any, in K .

Next we consider the experimental cross sections near 40 and 50 Mev. The heavy element cross sections rise again as the energy is lowered from 60 to 40 Mev. This rise can be duplicated by our calculations by a further increase in V to such a large value at 40 Mev that we again get constructive interference (k_1R for lead must be equal to about 4.5 at 40 Mev). In order to make the calculated rise in cross section as large as that observed, however, it must be assumed that K drops to such a low value that k_1/K is of the order of 10. If extrapolate V and K in this way and calculate the cross sections of cadmium, copper, and iron at 50 and 40 Mev, we obtain cross sections which are 20 to 35% lower than those observed because k_1R for these elements is such that the interference is highly destructive.

The other alternative for fitting the 40 and 50 Mev cross sections of the heavy elements is to decrease the value of k_1

as the neutron energy is lowered below 60 Mev, thus moving the heavy element cross sections back up towards the first interference maximum. If this is done, it is also possible to fit the cross sections of cadmium, copper and iron by appropriate choices of K at 40 and 50 Mev which join smoothly to the curve of K as a function of energy above 70 Mev. The apparent small dip in the copper and cadmium cross sections at 40 Mev can be obtained by using a somewhat smaller value of k_1/K at 40 Mev than at 50 Mev.

Following the qualitative observations we have just made, a search was made for values of V and K which would allow a best fit to the experimental data: The results are shown in Figs. 6 and 7. The observed and calculated total cross sections are compared in Figs. 12 and 13.

There is some latitude in the possible choices of V and K which will allow a reasonably good fit to the total cross sections below 70 Mev, but the author has found it impossible to duplicate even the qualitative trends of the data without assuming that V goes through a sharp maximum at a neutron energy of about 60 Mev. Since the only data available at present for determining the nuclear parameters in the neighborhood of 60 Mev are the total neutron cross sections, we have no further check on the apparent resonance-like behavior of V at this energy. For this reason some thought has been given to possible experiments which could be done in this energy range in an attempt to find more direct evidence for the existence of the potential resonance.

Since the cross section for elastic scattering in the forward direction depends very strongly on the value of V , calculations have been made of the expected zero angle cross sections of lead and copper in the neighborhood of 60 Mev, using the values of V and K shown in Figs. 6 and 7. The results of these calculations are shown graphically in Fig. 14. When the k^2 dependence of the forward scattering cross section is taken out, it is seen that the forward cross section for lead shows an increase of almost 40% from a neutron energy of 60 to 70 Mev, whereas the forward cross section for copper is essentially constant. In addition the lead cross section drops sharply from 40 to 60 Mev, while the copper cross section rises. It is suggested that these predicted effects could be looked for experimentally by observing the small angle scattering of neutrons by lead and copper, using a technique for energy discrimination similar to that described by Taylor and Wood(11). Since the expected change in cross section with energy is very sharp near 60 Mev, it would be important to select as narrow neutron energy bands as possible. In addition, the smaller the scattering angle, the more strongly energy dependent will be the cross section; the ideal experiment would be the measurement of the scattering at small enough angles to allow a reasonably good extrapolation to zero angle.

Although, as we have pointed out, there is strong evidence that the near discontinuity in the heavy element total cross sections near 60 Mev is not due to a sudden change in the ab-

sorption coefficient, it would be worthwhile to try to check this more directly by measuring the absorption cross section of lead as a function of energy. Because of the extreme difficulty of measuring absorption cross sections directly with the required energy resolution, however, it is suggested that it might be more feasible to measure the elastic scattering angular distribution as a function of energy and obtain the total scattering cross section by integration over angle. It would appear reasonable to embark on an experimental program as difficult as this only if the small angle scattering experiments should show conclusively that nuclear scattering near 60 Mev is anomalous.

TABLE I

Comparison Between Calculated and Observed
Total Cross Sections at 410 Mev

Element	Observed Total Cross Section ⁽¹⁴⁾ (millibarns)	Calculated Total Cross Section*
D	62.0 ± 4	63
Be	231 ± 4	231
C	297 ± 3	294
O	378 ± 5	374
Al	587 ± 8	583
S	672 ± 9	672
Cl	742 ± 10	735
Fe	1073 ± 12	1070
Cu	1187 ± 14	1185
Cd	1848 ± 21	1870
Pb	2890 ± 30	2920
Th	3210 ± 40	3200
U	3230 ± 40	3260

* $k_1 = 0.067 \times 10^{13} \text{ cm.}^{-1}$

$K = 0.18 \times 10^{13} \text{ cm.}^{-1}$

$V = 10 \text{ Mev}$

40

TABLE II

Comparisons Between Measured and Observed
Absorption Cross Sections

E = 95 Mev (Ref. 16)

Element	Observed Absorption Cross Section	Calculated Absorption Cross Section
C	.22 ± .01 barns	.216 barns
Al	.42 ± .015	.406
Cu	.78 ± .03	.765
Pb	1.74 ± .04	1.70

E = 270 Mev (Ref. 17)

C	.145 ± .015	.202
Cu	.580 ± .03	.72
Pb	1.42 ± .04	1.63

E = 1.4 Bev (Ref. 14)

C	.231*	.230
Al	.446	.432
Cu	.755	.795
Sn	1.220	1.22
Pb	1.61	1.74

* The statistical accuracies quoted by the authors are 3 to 5%.

TABLE III

Calculated and Observed Elastic Scattering Cross
Sections in the Forward Direction for 84 Mev Neutrons

$$\left(\frac{d\sigma}{d\Omega}\right)_{\theta=0} \text{ (barns/ster.)}$$

<u>Element</u>	<u>Calculated</u>	<u>Observed</u> ⁽⁵⁾
Lead	60	66 ± 9
Copper	14.5	15 ± 2
Aluminum	5.0	5.2 ± 5

44

FIG. 2 $\frac{\sigma_T}{\pi R^2}$ vs. $R_1 R$ FOR VARIOUS
VALUES OF $\eta = \frac{R_1}{K}$

072 C46

45

FIG. 3 $\frac{\sigma}{\pi R^2}$ vs. KR FOR VARIOUS $\eta = \frac{k_1}{K}$

072 047

FIG. 4 μ_1 VS E FOR VARIOUS VALUES
OF η_{85}

FIG. 9 CALCULATED AND OBSERVED⁽¹⁾ TOTAL
CROSS SECTIONS VS NEUTRON ENERGY

072 055

FIG. 13 OBSERVED AND CALCULATED TOTAL CROSS SECTIONS VS. ENERGY

FIG. 14 ELASTIC SCATTERING IN THE FORWARD
DIRECTION VS. NEUTRON ENERGY

REFERENCES

- 1) H. A. Bethe, Phys. Rev. 57, 1125 (1940).
- 2) R. Serber, Phys. Rev. 72, 1114 (1947).
- 3) S. Fernbach, R. Serber, and T. B. Taylor, Phys. Rev. 75, 1352 (1949).
- 4) L. J. Cook, E. M. McMillan, J. M. Peterson, and D. C. Sewell, Phys. Rev. 75, 7 (1949).
- 5) A. Bratenahl, S. Fernbach, R. H. Hildebrand, C. E. Leith, and B. J. Moyer, Phys. Rev. 77, 597 (1950).
- 6) R. Fox, C. Leith, and L. Wouters, Phys. Rev. 80, 23 (1950).
- 7) J. DeJuren, Phys. Rev. 80, 27 (1950).
- 8) J. DeJuren and B. J. Moyer, Phys. Rev. 81, 919 (1951).
- 9) H. H. Barschall, Phys. Rev. 86, 431 (1952).
- 10) H. Feshbach, C. E. Porter, and V. F. Weisskopf, Phys. Rev. 90, 166 (1953).
- 11) A. E. Taylor and E. Wood, Phil. Mag. 44, 95 (1953).
- 12) W. I. Linlor and B. Ragent, Phys. Rev. 92, 835 (1953).
- 13) B. Ragent, UCRL-2337, Berkeley (1953).
- 14) V. A. Nedzel, Phys. Rev. 94, 174 (1954).
- 15) G. A. Snow, T. Coor, D. A. Hill, W. F. Hornyak, L. W. Smith, Bull. Am. Phys. Soc. 29, 54 (1954).
- 16) L. I. Schiff, "Quantum Mechanics," McGraw Hill Book Co. (1944).
- 17) J. DeJuren and N. Knable, Phys. Rev. 77, 606 (1950).
- 18) R. Hofstadter, H. R. Fechter, and J. A. McIntyre, Phys. Ref. 92, 978 (1953).
- 19) D. R. Yennie, D. G. Ravenhall, and R. N. Wilson, Phys. Rev. 93, 948 (1954).
- 20) M. H. Johnson and E. Teller, Phys. Rev. 93, 357 (1954).
- 21) R. Hofstadter, Invited Paper at Jan. 1954 Meeting of Am. Phys. Soc. at New York, N.Y.

- 22) L. N. Cooper and E. M. Henley, Phys. Rev. 92, 801 (1953).
- 23) M. L. Goldberger, Phys. Rev. 74, 1268 (1948).
- 24) R. Jastrow, Phys. Rev. 82, 261 (1951).
- 25) F. Mandl and T. H. R. Skyrme, Phil. Mag. 44, 1028 (1953).

I-II
1-58

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

1954 JUL 2 AM 10:22