UCRL 8364

MASTER

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

ANALYSIS OF THE BEVATRON
K⁻ BEAM BY MEANS OF AN EMULSION
STACK

BERKELEY, CALIFORNIA

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

UCRL-8364
Physics and Mathematics

UNIVERSITY OF CALIFORNIA

Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

ANALYSIS OF THE BEVATRON K^- BEAM
BY MEANS OF AN EMULSION STACK

John N. Dyer

July 3, 1958

ANALYSIS OF THE BEVATRON K^- BEAM
BY MEANS OF AN EMULSION STACK

John N. Dyer

Radiation Laboratory
University of California
Berkeley, California

July 3, 1958

Abstract

The Bevatron K^- beam separated by the velocity spectrometer has been analyzed with the aid of an emulsion stack, and particle density and distributions are given.

ANALYSIS OF THE BEVATRON K^- BEAM BY MEANS OF AN EMULSION STACK

John N. Dyer

Radiation Laboratory
University of California
Berkeley, California

July 3, 1958

We have exposed a stack consisting of 108 pellicles of K. 5 emulsion, 600 μ by 3 by 6 inches, to the K^- beam separated by the coaxial velocity spectrometer.¹ The stack was placed with its face 6 inches behind the $1 \times 1/2$ -in. aperture of collimator C-2 and oriented so that the beam entered parallel to the 6-in. axis of the assembled stack. This face was protected from x-rays by $1/4$ -in. Pb, and additional shielding of Pb, Cd, and paraffin was used for the remaining faces of the stack. The duration of the exposure was about 50 hours, during which 3.25×10^{14} protons were incident on the Bevatron target.

From our analysis of the exposed stack we are able to make the following observations:

1. From data taken at the entrance face of the stack:

(a) beam size (defined by C-2)	1×1.7 in. at face
(b) flux of near-minimum tracks	$6.7 \times 10^4 / \text{cm}^2$
(c) flux of K^- tracks	$670 / \text{cm}^2$
(d) K^- /near-minimum ratio	$\sim 1/100$

2. From the distribution of stopping K^- particles:

(a) Nearly all stopping K^- particles are contained in a volume of about 50 cm^3 , therefore the density of stopping K^- particles is about $70 / \text{cm}^3$ in this volume, and about 200 per cubic centimeter in the region of highest density.

(b) The range distribution of stopping K^- mesons is as shown in the histogram in Fig. 1 (by range, we mean the distance of penetration into the emulsion stack. This is about 1 mm shorter than the true range because of scattering in the emulsion and the divergence of entrance angles). The momentum corresponding to the true range represented by the maximum of this distribution is 430 Mev/c, with a momentum spread of about $\pm 5\%$ at the half-maximum points.

(c) The distributions of stopping K^- particles in the vertical and horizontal directions respectively are as represented in Figs. 2 and 3. The stack was situated so that the horizontal direction was normal to the pellicle surfaces; hence, the horizontal distribution is obtained by counting the total number of stopping K^- particles in each pellicle. We have in our possession only one

¹Horwitz, Murray, Ross, and Tripp, 450-Mev/c K^- and $\bar{p}$ Beams at the Northwest Target Area of the Bevatron Separated by the Coaxial Velocity Spectrometer, UCRL-8269, June 1958.

half of the original stack, so that we can establish only one-half the horizontal distribution. (The other half is at present being studied by Dr. Howard Taft at Yale University.

A schematic drawing of the geometry is shown in Fig. 4.

This work was done under the auspices of the U. S. Atomic Energy Commission.

MU-15585

Fig. 1. Range distribution of stopping K^- mesons in emulsion.

MU-15584

Fig. 2. Vertical distribution of stopping K^- mesons in emulsion.

MU-15586

Fig. 3. Horizontal distribution of stopping K^- mesons in emulsion.

Fig. 4. Geometry of beam and emulsion stack.

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, express or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission to the extent that such employee or contractor prepares, handles or distributes, or provides access to, any information pursuant to his employment or contract with the Commission.