25180

~~SECRET~~

ORNL-144
Physics-General

Contract W-35-058 eng. 71

This document consists 12
pages
Copy 82 of 150 Series A.

ISSUED: K. Way
(for Declassification)

CLASSIFICATION CANCELLED

For the Atomic Energy Commission

Charles A. Geller
Jmm

Acad - 2368

This document is
PUBLICLY RELEASABLE
Hugh Kinner/OSTI
Authorizing Official
Date: 7/25/68

✓ GRAPHS SHOWING NEUTRON CROSS SECTIONS AS FUNCTIONS OF A, Z OR N ✓

by
G. Haines and K. Way ✓

Oak Ridge National Laboratories

Date of Manuscript: September 8, 1948
Date Declassified: November 2, 1948

Date Received: 9/8/48

Date Issued: OCT 4 1948

"This document contains restricted data within the meaning of the Atomic Energy Act of 1946 and/or information affecting the National defense of the United States within the meaning of the Espionage Act, 50 U.S.C. 31 and 32 as amended. Its transmission or the revelation of its contents in any manner to an unauthorized person is prohibited and may result in severe criminal penalty."

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

DISTRIBUTION:

1. A. O. Allen	24. M. M. Mann
2. D. C. Bardwell	25. F. H. Murray
3. S. Bernstein	26. K. Z. Morgan
4. E. P. Blizard	27. E. J. Murphy
5. G. E. Boyd	28. L. C. Noderer
6. A. R. Brosi	29. M. D. Peterson
7. W. E. Cohn	30. H. Pomerance
8. E. C. Campbell	31. M. E. Rose
9. J. A. Cox	32. H. C. Schweinler
10. N. Dismuke	33. S. Siegel
11. M. A. Feldman	34. M. M. Shapiro
12. S. Freed	35. N. M. Smith
13. W. W. Good	36. A. H. Snell
14. W. B. Harrison	37. B. I. Spinrad
15. J. E. Hill	38. R. Stoughton
16. A. Hollaender	39. C. G. Shull
17. A. Householder	40. L. D. Roberts
18. J. R. Huffman	41. J. A. Swartout
19. W. E. Johnson	42. E. H. Taylor
20. K. A. Kraus	43. R. S. Thackeray
21. H. A. Levy	44. P. C. Tompkins
22. A. B. Martin	45. K. Way
23. F. T. Miles	46. A. M. Weinberg
	47. E. O. Wollan
- 48-49. Chemistry Library
50-51. Physics Library
52-53. Training School Library
54-75. Central Files
76-83. K. Way (for declassification)
- 84-91. Argonne National Laboratory
92. Armed Forces Special Weapons Project
93-94. Atomic Energy Commission, Washington
95. Battelle Memorial Institute
96-103. Brookhaven National Laboratories
104-107. Carbide & Carbon Chemicals Corporation (K-25 Area)
108-111. Carbide & Carbon Chemicals Corporation (Y-12 Area)
112. Columbia University (Dunning)
113-116. General Electric Company
117-121. Hanford Directed Operations
122. Iowa State College
123-125. Los Alamos
126-127. Monsanto Chemical Company, Dayton
128-129. National Bureau of Standards
130. Naval Radiological Defense Labs.
131. NEPA
132-133. New York Directe Operations
134. Patent Advisor, Washington
135-149. Technical Information Division, ORDO
150. UCLA Medical Research Laboratory (Warren)
151-155. University of California Radiation Laboratory
156-157. University of Rochester

~~SECRET~~

GRAPHS SHOWING NEUTRON CROSS SECTIONS AS FUNCTIONS OF A, Z, OR N

By G. Haines and K. Way

The accompanying graphs were prepared from the neutron cross section data of Tables of Neutron Cross Sections, Mon P-405, October 1947, and Mon P-405 Supplement and Errata, April 1948. A, Z, and N stand for atomic number, proton number, and neutron number respectively. References to sources of data are given on the graphs.

Activation cross sections for fast neutrons of two different energy regions are shown in Figure 1 as a function of neutron number. The dips at neutron numbers 50, 82, and 126 have been commented on by Dr. Maria Mayer, Phys. Rev., 74, 235 (1948), in connection with other evidence that nuclei with these neutron numbers have particularly tight bindings and that the 51st, 83rd, or 126th neutrons are rather weakly bound.

Figure 2 shows γ -ray yields from bombardment with 2.4-Mev neutrons which are presumably roughly proportional to inelastic scattering cross sections. The atomic number A of the most abundant isotope of the element investigated has been chosen as abscissa. This curve shows a striking resemblance to Hughes curve (GP-3750) for fast activation cross sections. Nuclei with 82 and 126 neutrons again show abnormally small cross sections, as is to be expected if formation of the compound nucleus is less probable for nuclei with these neutron numbers than for their neighbors. No anomaly appears at neutron number 50 (Zr_{90}^{40}).

"This document contains restricted data within the meaning of the Atomic Energy Act of 1946 and/or information affecting the National defense of the United States within the meaning of the Espionage Act, 50 U.S.C. 31 and 32 as amended. Its transmission or the revelation of its contents in any manner to an unauthorized person is prohibited and may result in severe criminal penalty."

~~SECRET~~

~~SECRET~~

F384

Figures 3 and 4 show total cross sections as functions of the most abundant A for neutron energies of 2.5 MeV and 14 MeV , respectively. Comparison of these two figures shows that as the neutron energy increases the cross sections decrease slightly and the curve tends to smooth out. For 90- MeV neutrons, Cook, et al., Phys. Rev., 72, 1264 (1947), have recently found cross sections very nearly proportional to the geometrical area of the nucleus. Cook's values are slightly lower than those of Amaldi.

F5

Figure 5 shows thermal neutron absorption cross sections of elements as function of charge number Z . Such cross sections are, of course, averages of isotopic cross sections for all stable nuclei with the same Z in which the cross section for each nucleus is weighted with its relative abundance. There is a general trend toward large cross sections as Z increases. This is to be expected if the cross section is determined chiefly by the nearness of an energy level of the compound nucleus to the energy state produced by neutron capture, i.e. by the difference $|E_r^{(N)} - E_B|$, where E_r is a resonance energy and E_B is the binding energy of a neutron to the target nucleus. As the atomic number increases the level density near E_B increases, the chance for a small $|E_r - E_B|$ increases and the probability of large cross sections grows correspondingly greater.

One would expect peaks and dips to occur at random on the general rise. Figure 5 shows no pronounced pattern. It is a little surprising, however, to find that two of the dips occur in the region of proton numbers 50 and 82.

These dips still appear when isotopic cross sections are plotted against Z as in Figure 6. Some of the isotopic values are due to mass spectrographic depletion measurements, others to activation data. Figure 6 shows the values of all known individual cross sections for each value of Z . Most of the peaks and dips of Figure 5 are considerably blurred, but the pronounced minima just above 50 and 80 remain. Additional measurements may show that these do not exist or that they are connected with the neutron numbers 82 and 126. F6

If the binding energies of the 51st, 83rd, or 127th neutrons are actually abnormally low, the excited state of the compound nucleus will be in a region of lower than normal-level density. The chance of finding a small value of $|E_F^{\textcircled{D}} - E_B|$ will, therefore, be smaller than usual and so one would expect a greater probability for low absorption cross sections for nuclei with 50, 82, or 126 neutrons.

Figure 7, which shows isotopic cross sections plotted against neutron number, indicates that this expectation may be fulfilled. There are not yet, however, enough data to allow any clear cut conclusions. F7

AECD - 2368

UNITED STATES ATOMIC ENERGY COMMISSION

GRAPHS SHOWING NEUTRON CROSS SECTIONS AS FUNCTIONS OF A, Z, OR N

by

G. Haines

K. Way

Oak Ridge National Laboratories

Date of Manuscript: September 8, 1948
Date Declassified: November 2, 1948

This document is for official use.
Its issuance does not constitute authority
for declassification of classified copies
of the same or similar content and title
and by the same authors.

Technical Information Branch, Oak Ridge, Tennessee
AEC, Oak Ridge, Tenn., 2-23-49--750-A1349

TECHNICAL LIBRARY
TECHNICAL INFORMATION BRANCH, ORE
ATOMIC ENERGY COMMISSION

937-1

GRAPHS SHOWING NEUTRON CROSS SECTIONS AS FUNCTIONS OF A, Z, OR N

By G. Haines and K. Way

The accompanying graphs were prepared from the neutron cross section data of Tables of Neutron Cross Sections, Mon P-405, October 1947, and Mon P-405 Supplement and Errata, April 1948. A, Z, and N stand for atomic number, proton number, and neutron number respectively. References to sources of data are given on the graphs.

Activation cross sections for fast neutrons of two different energy regions are shown in Figure 1 as a function of neutron number. The dips at neutron numbers 50, 82, and 126 have been commented on by Dr. Maria Mayer, Phys. Rev., 74, 235 (1948), in connection with other evidence that nuclei with these neutron numbers have particularly tight bindings and that the 51st, 83rd, or 126th neutrons are rather weakly bound.

Figure 2 shows γ -ray yields from bombardment with 2.4 Mev neutrons which are presumably roughly proportional to inelastic scattering cross sections. The atomic number A of the most abundant isotope of the element investigated has been chosen as abscissa. This curve shows a striking resemblance to Hughes curve (CP-3750) for fast activation cross sections. Nuclei with 82 and 126 neutrons again show abnormally small cross sections, as is to be expected if formation of the compound nucleus is less probable for nuclei with these neutron numbers than for their neighbors. No anomaly appears at neutron number 50 (Zr^{90}).

Figures 3 and 4 show total cross sections as functions of the most abundant A for neutron energies of 2.5 and 14 Mev, respectively. Comparison of these two figures shows that as the neutron energy increases the cross sections decrease slightly and the curve tends to smooth out. For 90-Mev neutrons, Cook, et al., Phys. Rev., 72, 1264 (1947), have recently found cross sections very nearly proportional to the geometrical area of the nucleus. Cook's values are slightly lower than those of Amaldi.

Figure 5 shows thermal neutron absorption cross sections of elements as function of charge number Z. Such cross sections are, of course, averages of isotopic cross sections for all stable nuclei with the same Z in which the cross section for each nucleus is weighted with its relative abundance. There is a general trend toward large cross sections as Z increases. This is to be expected if the cross section is determined chiefly by the nearness of an energy level of the compound nucleus to the energy state produced by neutron capture, i.e. by the difference $|E_r - E_B|$, where E_r is a resonance energy and E_B is the binding energy of a neutron to the target nucleus. As the atomic number increases the level density near E_B increases, the chance for a small $|E_r - E_B|$ increases and the probability of large cross sections grows correspondingly greater.

One would expect peaks and dips to occur at random on the general rise. Figure 5 shows no pronounced pattern. It is a little surprising, however, to find that two of the dips occur in the region of proton numbers 50 and 82.

These dips still appear when isotopic cross sections are plotted against Z as in Figure 6. Some of the isotopic values are due to mass spectrographic depletion measurements, others to activation data. Figure 6 shows the values of all known individual cross sections for each value of Z. Most of the peaks and dips of Figure 5 are considerably blurred, but the pronounced minima just above 50 and 80 remain. Additional measurements may show that these do not exist or that they are connected with the neutron numbers 82 and 126.

Figure 1. Activation cross sections as a function of neutron number for fast neutrons of two different energy regions.

Figure 2. γ -ray yields from bombardment with 2.4-Mev neutrons.

If the binding energies of the 51st, 83rd, or 127th neutrons are actually abnormally low, the excited state of the compound nucleus will be in a region of lower than normal-level density. The chance of finding a small value of $|E_r - E_B|$ will, therefore, be smaller than usual and so one would expect a greater probability for low absorption cross sections for nuclei with 50, 82, or 126 neutrons.

Figure 7, which shows isotopic cross sections plotted against neutron number, indicates that this expectation may be fulfilled. There are not yet, however, enough data to allow any clear cut conclusions.

Figure 3. Total cross sections as functions of the most abundant A for neutron energies of 2.5 Mev.

Figure 4. Total cross sections as functions of the most abundant A for neutron energies of 14 Mev.

Figure 5. Thermal neutron absorption cross sections of elements as functions of charge number Z.

937

Figure 6. Cross sections of isotopes for thermal neutrons for each value of Z.

Figure 7. Isotopic cross sections plotted against neutron number.

Figure Captions AECD-2368

Figure 1. Activation cross sections as a function of neutron number^Y for fast neutrons of two different energy regions.

Figure 2. γ -ray yields from bombardment with 2.4-Mev neutrons.

Figure 3. Total cross sections as functions of the most abundant A for neutron energies of 2.5 Mev.

Figure 4. Total cross sections as functions of the most abundant A for neutron energies of 14 Mev.

Figure 5. Thermal neutron absorption cross sections of elements as functions of charge number Z.

Figure 6. Cross sections of isotopes for thermal neutrons for each value of Z.

Figure 7. Isotopic cross sections plotted against neutron number.