3
SCR-103

MASTER

Sandia Corporation

REPRINT

QUADRATURE FORMULAS INVOLVING
DERIVATIVES OF THE INTEGRAND

by

Preston C. Hammer

and

Howard H. Wicke

July 1959

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

To be published in: MATHEMATICAL TABLES AND
OTHER AIDS TO COMPUTATION
Forthcoming issue, 1960.

The part of this work contributed by Professor
Hammer was supported in part by the Office of Ordnance
Research, U.S. Army Contract No. DA-11-022-ORD-2301.
The part of the work contributed by Dr. Wicke was supported
by the United States Atomic Energy Commission.

SANDIA CORPORATION REPRINT

QUADRATURE FORMULAS INVOLVING
DERIVATIVES OF THE INTEGRAND

by

Preston C. Hammer
Director, Numerical Analysis Laboratory,
University of Wisconsin
Consultant to Sandia Corporation

and

Howard H. Wicke
Systems Analysis Department
Sandia Corporation

July 1959

ABSTRACT

In this paper, the authors demonstrate the existence of quadrature formulas involving derivatives of the integrand. One such formula is, for k an odd positive integer,

$$\int_{-1}^1 f(x) dx = 2 \sum_{i=0}^{(k-1)/2} \frac{f^{(2i)}(0)}{(2i+1)!} + \sum_{j=1}^m a_j \left[f^{(k)}(x_j) - f^{(k)}(-x_j) \right] + R_{m,k}(f).$$

The authors use the theory of orthogonal polynomials to establish these formulas and provide both recurrence formulas for the polynomials and an expression for the remainder. The weight functions involved in the proof appear to be different from those treated heretofore in the literature.

QUADRATURE FORMULAS INVOLVING DERIVATIVES OF THE INTEGRAND

Introduction -- In this paper we demonstrate the existence of quadrature formulas of the following types. For k an odd positive integer,

$$(1) \quad \int_{-1}^1 f(x)dx = 2 \sum_{i=0}^{(k-1)/2} \frac{f^{(2i)}(0)}{(2i+1)!} + \sum_{j=1}^m a_j \left[f^{(k)}(x_j) - f^{(k)}(-x_j) \right] + R_{m,k}(f).$$

For k an even positive integer,

$$(2) \quad \int_{-1}^1 f(x)dx = 2 \sum_{i=0}^{(k-2)/2} \frac{f^{(2i)}(0)}{(2i+1)!} + \sum_{j=1}^m a_j \left[f^{(k)}(x_j) + f^{(k)}(-x_j) \right] + R_{m,k}(f).$$

Each of formulas (1) and (2) will be shown to exist so that they are exact for polynomials of degree at most $4m+k$ for odd k , and $4m+k-1$ for even k .

It is possible to use tables published by Hammer, Marlowe, and Stroud [2] and extended by Fishman [1] to obtain formulas of the form

$$(3) \quad \int_0^1 f(x)dx = f(1) + \sum_{j=1}^m a_j f'(x_j) + R_m(f)$$

and likewise formulas using higher derivatives. These formulas are asymmetrical and for some uses would be more appropriate than formula (1) or (2).

It is considered that formula (1) or (2) or (3) may be useful when the integrand function is represented as a variable integral for which derivatives

may be easier to compute than the integrand function itself. It is also anticipated that these formulas may be used in the numerical solution of differential equations.

Kopal [3] has devised formulas using the first derivative values. His approach leaves difficulties in establishing existence and reality of evaluation points and in some cases he has computed more than one formula of the same degree. The method we propose has direct connection with the established theory of orthogonal polynomials which gives the existence, reality, and distinctness of the evaluation points and the positiveness of the weights a_j . A linear transformation to give integration limits $-h, h$ in formula (1) or (2) results in multiplying each derivative of order n by h^{n+1} .

We have not identified the orthogonal polynomial systems with any treated in detail in the literature. However, tables of the a_j and x_j for $k = 1, 2, m = 1 (1) 10$, and $k = 3, 4, m = 1 (1) 9$, have been computed by G. W. Struble in [5], where tables for the $\mu_{m,k}^{(m)}$ of the remainder terms are also to be found. For purposes of computation we give a standard type recursion formula permitting generation of each polynomial in a sequence from its two predecessors.

Throughout the paper we assume that the integrand function has all-order derivatives appearing and that these are continuous.

Reduction of the Problems -- It is well known and readily established that

$$(4) \quad \int_0^1 \left(\int_0^x \right)^n g(x) (dx)^{n+1} = \frac{1}{n!} \int_0^1 (1-x)^n g(x) dx,$$

where $\left(\int_0^x\right)^n g(x)(dx)^n$ denotes the result of integrating $g(x)$ successively n times over $(0, x)$. Now our formulas (1) and (2), by choice of form, will hold for every odd integrand function $f(x)$ and hence we write $f(x) = f_0(x) + f_1(x)$, where $f_0(x) = \frac{f(x) + f(-x)}{2}$ and $f_1(x) = \frac{f(x) - f(-x)}{2}$. The functions f_0 and f_1 may be called the even and odd components of f , respectively. We now observe

$$(5) \quad \int_{-1}^1 f(x)dx = 2 \int_0^1 f_0(x)dx \text{ and } f_0^{(k)}(x) = \frac{f^{(k)}(x) + (-1)^k f^{(k)}(-x)}{2}.$$

Moreover,

$$(6) \quad \int_0^1 f_0(x)dx = \int_0^1 \left(\int_0^x\right)^k f_0^{(k)}(x)(dx)^{k+1} + S_k(f), \text{ where}$$

$$S_k(f) = \begin{cases} \sum_{i=0}^{(k-1)/2} \frac{f^{(2i)}(0)}{(2i+1)!}, & \text{for } k \text{ odd} \\ \sum_{i=0}^{(k-2)/2} \frac{f^{(2i)}(0)}{(2i+1)!}, & \text{for } k \text{ even.} \end{cases}$$

Hence our formulas (1) and (2) may be written, using formulas (4), (5), and (6), and dropping $2S_k(f)$ from each side, as follows:

$$(7) \quad \frac{2}{k!} \int_0^1 (1-x)^k f_0^{(k)}(x)dx = 2 \sum_{j=1}^m a_j f_0^{(k)}(x_j) + R_{m,k}(f).$$

Now we only need derive formulas (7) exact for even-degree polynomials replacing $f_0(x)$. Then, if k is odd we have with $u = x^2$

$$\frac{2}{k!} \int_0^1 (1-x)^k x P(x^2)dx = \frac{1}{k!} \int_0^1 (1-\sqrt{u})^k P(u)du = 2 \sum_{j=1}^m a_j \sqrt{u_j} P(u_j), \text{ and thus,}$$

$$(8) \quad \frac{1}{k!} \int_0^1 (1 - \sqrt{u})^k P(u) du = \sum_{j=1}^m b_j P(u_j), \text{ where } b_j = 2a_j \sqrt{u_j}.$$

On the other hand, if k is even, so that $f_0^{(k)}(x)$ is replaced by an even polynomial $P(x^2)$, then we have with $u = x^2$, $b_j = 2a_j$,

$$(9) \quad \frac{1}{k!} \int_0^1 \frac{(1 - \sqrt{u})^k}{\sqrt{u}} P(u) du = \sum_{j=1}^m b_j P(u_j).$$

Now formulas (8) and (9) are in standard form to apply the theory of orthogonal polynomials. The weight functions are positive-valued and appropriate. Hence there exists a sequence $\{P_{m,k}(u)\}$ of orthogonal polynomials for each $k = 1, 2, \dots$, the zeros of which are the squares of the proposed evaluation points x_j , and positive numbers b_j from which the a_j are determined. The zeros are real, distinct, and in the interval.

The weight functions are simple in form but we have been unable to find references in which this class has been treated. For $k = 1$ we have verified that the system of polynomials is not a classical one since $\{P'_{m,1}(u)\}$ is not orthogonal. For purposes of calculating formulas, however, we state in the next section recursion formulas which permit sequential generation of the polynomials for each $k \geq 1$.

Recursion Formula for Polynomials -- The recursion formula stated here is of the standard sort, but its form is preferable to the one given by Szegö [4], p. 41, since it involves only the coefficients of P_{n-1} and P_{n-2} to calculate the multipliers B_n and C_n . This is important when explicit formulas

for the multipliers have not been determined. We assume $\{P_n(u)\}$ is a system of polynomials with leading coefficient 1, orthogonal over the interval $[a, b]$ with respect to the weight function $w(u)$. Then we have

$$(10) \quad P_n(u) = (u + B_n)P_{n-1}(u) - C_n P_{n-2}(u), \quad n \geq 2$$

If we define

$$(11) \quad \mu_n^{(k)} = \int_a^b w(u) u^k P_n(u) du$$

we may observe $\mu_n^{(k)} = 0$, $0 \leq k < n$, and $\mu_n^{(n)} = \int_a^b w(u) u^n P_n(u) du = \int_a^b w(u) (P_n(u))^2 du$. We also define $c_{n-1}^{(j)}$ as the coefficient of u^j in $P_{n-1}(u)$. Then B_n and C_n are given by

$$(12) \quad B_n = - \left(c_{n-1}^{(n-2)} + \frac{\mu_{n-1}^{(n)}}{\mu_{n-1}^{(n-1)}} \right)$$

$$(13) \quad C_n = \frac{\mu_{n-1}^{(n-1)}}{\mu_{n-2}^{(n-2)}}.$$

For our particular problem, the $\mu_{n-2}^{(n-2)}$, $\mu_{n-1}^{(n-1)}$, and $\mu_{n-1}^{(n)}$ are linear combinations of the coefficients of P_{n-2} and P_{n-1} with rational multipliers and the coefficients of every P_n are rational. Hence, in the absence of better formulas, starting with P_0 and P_1 , we can generate the remaining polynomials, in principle. In practice, of course, the coefficients grow very rapidly in exact rational form.

Now $P_{0,k} = 1$ for every k , and, for degree one, $\{P_{1,k}(u)\}$ is given by

$$(14) \quad P_{1,2n-1}(u) = u - \frac{3}{(2n+3)(n+1)}, \quad n = 1, 2, 3, \dots,$$

and

$$(15) \quad P_{1,2n}(u) = u - \frac{1}{(2n+3)(n+1)}, \quad n = 1, 2, 3, \dots$$

Hence, with the recursion formula (10), successive higher-degree polynomials may be calculated.

The orthonormal sequence $\{Q_n(u)\}$ is determined by

$$(16) \quad P_n(u) = \sqrt{\mu_n^{(n)}} Q_n(u).$$

Remainders for the Integration Formulas -- In this section we give the remainder formulas. We will give the explicit functions to which Rolle's Theorem may be applied. The method is a variation of Markoff's for Hermite (osculating) interpolation (Szegő [4], p. 369.) As usual, we obtain the highest-degree polynomial $H(x)$ for which the formula is exact and which agrees with the integrand function or its derivatives at all evaluation points. Then we estimate $f^{(k)}(x) - H^{(k)}(x)$ and $f_0^{(k)}(x) - H_0^{(k)}(x)$, and use formula (7) to find $R_{m,k}(f)$. (Here H_0 is the even component of H .)

For k odd and $2m+1$ evaluation points $0, \pm x_1, \dots, \pm x_m$, we require $H(0) = f(0)$, $H^{(n)}(0) = f^{(n)}(0)$ for $n = 1, \dots, k$, $H^{(k)}(\pm x_j) = f^{(k)}(\pm x_j)$, $H^{(k+1)}(\pm x_j) = f^{(k+1)}(\pm x_j)$, $j = 1, \dots, m$. Then $H(x)$ is a polynomial of degree at most $4m+k$. Let $P_m(x^2)$ be the polynomial $P_{m,k}(u)$ determining the evaluation points, x , by $P_m(x_j^2) = 0$, $j = 1, \dots, m$. Then let x be any number which is none of $0, \pm x_j$. Define

$$(17) \quad F(z) = f^{(k)}(z) - H^{(k)}(z) - \frac{f^{(k)}(x) - H^{(k)}(x)}{x[P_m(x^2)]^2} z [P_m(z^2)]^2.$$

Then $F(z)$ vanishes at $x, 0, \pm x_j$ ($2m + 2$ distinct points). Also, $F'(z)$ vanishes at $\pm x_j$ and $2m + 1$ other distinct points, or $4m + 1$ points. Hence, applying Rolle's Theorem there exists a ξ_1 such that $F^{(4m+1)}(\xi_1) = 0$, and thus

$$(18) \quad f^{(k)}(x) - H^{(k)}(x) = \frac{f^{(4m+k+1)}(\xi_1)x \left[P_m(x^2) \right]^2}{(4m+1)!}.$$

From equation (18) applied to x and $-x$, and from the continuity of $f^{(4m+k+1)}(x)$, it follows that the even components satisfy

$$(19) \quad f_0^{(k)}(x) - H_0^{(k)}(x) = \frac{f^{(4m+k+1)}(\xi_2)x \left[P_m(x^2) \right]^2}{(4m+1)!}.$$

Hence, from equation (7) by the continuity of $f^{(4m+k+1)}(x)$ and the first theorem of the mean for integrals, we have, with $u = x^2$, and for some $\eta \in [-1, 1]$,

$$(20) \quad R_{m,k}(f) = \frac{f^{(4m+k+1)}(\eta)}{(4m+1)!} \int_0^1 \frac{(1-\sqrt{u})^k}{k!} \left[P_m(u) \right]^2 du, \text{ for } k \text{ odd},$$

or replacing the integral by $\mu_{m,k}^{(m)}$,

$$(21) \quad R_{m,k}(f) = \frac{f^{(4m+k+1)}(\eta)}{(4m+1)!} \mu_{m,k}^{(m)}, \text{ for } k \text{ odd}.$$

For k even, we require $H(x)$ of at most degree $4m + k - 1$ such that $H(0) = f(0)$, $H^{(n)}(0) = f^{(n)}(0)$, for $n = 1, \dots, k - 1$, $H^{(k)}(\pm x_j) = f^{(k)}(\pm x_j)$, $H^{(k+1)}(\pm x_j) = f^{(k+1)}(\pm x_j)$, for $j = 1, \dots, m$. Then, for x not equal to one of $\pm x_j$, we define

$$(22) \quad F(z) = f^{(k)}(z) - H^{(k)}(z) - \frac{f^{(k)}(x) - H^{(k)}(x) \left[P_m(z^2) \right]^2}{\left[P_m(x^2) \right]^2}.$$

Now $F'(z)$ vanishes at $4m$ distinct points, and there exists a ξ_1 such that $F^{(4m)}(\xi_1) = 0$, which implies

$$(23) \quad f^{(k)}(x) - H^{(k)}(x) = \frac{f^{(4m+k)}(\xi_1) [P_m(x^2)]^2}{(4m)!}.$$

The even components f_0 and H_0 then satisfy

$$(24) \quad f_0^{(k)}(x) - H_0^{(k)}(x) = \frac{f^{(4m+k)}(\xi_2) [P_m(x^2)]^2}{(4m)!},$$

using the same argument used in the passage from (18) to (19). Then, as before, substitution in equation (7) gives, with $u = x^2$, and for some $\eta \in [-1, 1]$

$$(25) \quad R_{m,k}(f) = \frac{f^{(4m+k)}(\eta)}{(4m)!} \int_0^1 \frac{(1 - \sqrt{u})^k}{k! \sqrt{u}} [P_m(u)]^2 du, \text{ for } k \text{ even,}$$

or, with the integral replaced by $\mu_{m,k}^{(m)}$,

$$(26) \quad R_{m,k}(f) = \frac{f^{(4m+k)}(\eta)}{(4m)!} \mu_{m,k}^{(m)}, \text{ for } k \text{ even.}$$

The number $\mu_{m,k}^{(m)}$ may be determined simply in the low-degree cases by substituting in the numerical formula integrands x^{4m+k+1} if k is odd, and x^{4m+k} if k is even.

It may be noted that our explicit choice of H is made for convenience in each case. That is, formulas (1) and (2) hold with $f = H$ and $R_{mk} = 0$, and H is a highest-degree polynomial for which this is true--i.e., no higher even power of x may be included. Moreover, in formula (6) we have

$$(27) \quad \int_0^1 H_0(x) dx = \int_0^1 \left(\int_0^x \right)^{k+1} H_0^{(k)}(x) (dx)^{k+1} + S_k(f),$$

since $f^{(j)}(0) = H^{(j)}(0)$, for appropriate values of j . Numerical tables of a_j , x , and the constant in the remainder have been calculated by G. W. Struble [5] for small values of k .

REFERENCES

1. H. Fishman, Numerical integration constants, Mathematical Tables and Other Aids to Computation, 11, 1. 1957.
2. P. C. Hammer, O. J. Marlowe, and A. H. Stroud, Numerical integration over simplexes and cones, Mathematical Tables and Other Aids to Computation, 10, 130-137. 1956.
3. Z. Kopal, Numerical Analysis, Wiley & Sons, New York, 1956.
4. G. Szegő, Orthogonal Polynomials, American Mathematics Society Colloquium Publication, 23, 1939.
5. G. W. Struble, Quadrature formulas using values of derivatives (to be submitted for publication).