CONF-731012--2

TITLE:

THE BIOLOGICAL EFFECTS OF NEGATIVE PIONS

AUTHOR(S):

M. R. Raju

MASTER

SUBMITTED TO:

Excerpta Medica, Proceedings of the XIIth International Congress of Radiology, Madrid, Spain (October 15, 1973)

By acceptance of this article for publication, the publisher recognizes the Government's (license) rights in any copyright and the Government and its authorized representatives have unrestricted right to reproduce in whole or in part said article under any copyright secured by the publisher.

The Los Alamos Scientific Laboratory requests that the publisher identify this article as work performed under the auspices of the U. S. Atomic Energy Commission.

NOTICE
This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

THE BIOLOGICAL EFFECTS OF NEGATIVE PIONS

M. R. Raju

Biophysics and Instrumentation Group, Los Alamos Scientific Laboratory
University of California, Los Alamos, New Mexico 87544 U.S.A.

INTRODUCTION

Study of negative pion biological effects relevant to radiotherapy has been limited because negative pion beams of appropriate momentum spread and intensity for therapy were not available. However, significant progress has been made using negative pion beams designed mainly for physics experiments at Berkeley and at CERN. These beams are nearly monoenergetic ($\sim 5\%$ momentum spread). The depth-dose distribution of these beams gives narrow Bragg peaks insufficient to cover most of the treatment volumes in radiotherapy. A special negative pion beam for biomedical work recently was built at the 7-GeV proton synchrotron (NIMROD) in England. This beam has a momentum spread of about 13%, closer to radiotherapeutic use. However, the dose rates, over a beam size of ~ 2 by 3 cm, are ~ 3 rads/hr at CERN and 30 to 60 rads/hr at the Berkeley and NIMROD accelerators. Nearly 15% of the dose at the peak position is due to contamination from electrons and μ mesons in all these beams. In spite of these beam limitations, some relevant radiobiological information has been obtained using systems that are relatively sensitive. These results were reviewed two years ago (Raju and Richman, 1972). I will present a brief review of the biological effects of negative pions including some of the unpublished data from England.

BIOLOGICAL EFFECTS AT THE PLATEAU

It is clear from previous presentations that, in the plateau region, most of the dose is deposited by negative pion energy loss and that a small fraction of the dose is deposited from nuclear interactions by negative pions in flight. The dose at the plateau region is lower than at the peak. Because the dose rate and relative biological effectiveness (RBE) of the beam at the

plateau region are low, very few measurements of biological effects have been measured in this region.

The biological effectiveness at the plateau was measured by looking for polyploidy induction in ascites tumor cells. It was found that for ^{60}Co gamma rays polyploidy induction was linear with dose and independent of dose rate in the 5 to 20 rads/hr region. Mice with ascites tumor cells were exposed to the pion beam (5 rads/hr) at the plateau region. The RBE relative to ^{60}Co gamma rays was found to be very nearly 1.0 (Loughman *et al.*, 1968). Nias *et al.* (1973) also found that the RBE at the plateau is nearly 1.0 for frozen HeLa cells. Winston *et al.* (1973) measured the biological effectiveness and oxygen enhancement ratio (OER) using *Vicia faba* 10-day root growth. They found the RBE of negative pions, compared to radium gamma rays, was 1.31 and the OER 2.04. The OER for radium gamma rays was 3.01. These results indicate that nuclear interactions by negative pions in flight are effective in increasing RBE and in reducing OER. The differences in biological effects at the plateau could be partly because of the biological systems and the end points used in these investigations. More work needs to be done.

BIOLOGICAL EFFECTS AT THE PEAK

RBE and OER were measured in various biological systems, and the results at the peak of the depth-dose distribution are given in Table 1. A high RBE value of 5.0 was obtained with ascites tumor cells when compared with an equivalent dose rate with ^{60}Co gamma rays. However, this system was dose-rate dependent. Because of a relatively high LET component at the peak, the D_0 for negative pions at the peak may not change appreciably with increase in dose rate, whereas for ^{60}Co gamma rays the D_0 decreases considerably. Hence, the

RBE for an acute dose rate of pions can be expected to be ~ 3.0 in comparison with an acute dose rate of 45 rads/min of ^{60}Co gamma rays. In addition, 5-day-old ascites tumor cells used in this investigation were known to be hypoxic. *Vicia faba* 10-day growth data from CERN showed a high RBE value of 4.5. This value was obtained at the back part of the Bragg peak where the LET component is expected to be higher than at the peak. Otherwise, the RBE was ~ 2.0 to 3.0 and the OER ~ 1.5 to 1.8 at the peak of the depth-dose distribution in most of the other biological systems used. The RBE values at the peak will be slightly higher and the OER values very slightly lower for a negative pion beam with relatively less contamination. However, these values change according to the width of the peak, as can be seen by comparing the data for *Vicia faba* 10-day growth from England and Berkeley. The RBE will be considerably reduced and the OER slightly higher for beams with the larger momentum spreads.

VARIATIONS IN BIOLOGICAL EFFECTIVENESS WITHIN THE PEAK REGION

Significant differences in biological effectiveness around the peak region have been detected by all investigators. Figure 1 shows the results from Berkeley using human kidney cells in culture, CERN results using bean roots, and results from England using bean roots and HeLa cells in culture. The biological effectiveness at the back side of the Bragg peak (marked as position III in Fig. 1a) was higher than at the front side (position II). This can be expected because, at the front side of the Bragg peak, a bigger fraction of the dose is delivered by relatively low-LET events due to energy loss of negative pions, whereas at the back side of the Bragg peak, a bigger fraction of the dose is delivered at a relatively higher LET by low-energy, heavy, charged particles arising out of negative pion capture. These differences in biological effectiveness will be reduced when two opposed fields are used. Mill *et al.* (1973)

also have done some fractionation experiments at the plateau and in the peak region, marked positions II and III in Fig. 1a. Their results indicate that there was practically no recovery at the back of the peak (position III) but that there was significant recovery in the front of the peak (position II) and in the plateau.

CELL-SURVIVAL CALCULATIONS

Cell survival was calculated with some success for beams of mixed LET such as fast neutrons and for negative pions using cell-survival data at various discrete LET values (Bewley, 1968; Curtis, 1971). Katz *et al.* (1971) have recently developed a model to calculate cell survival for radiations of different LET. This model gives results in fair agreement with fast-neutron experimental results (Armstrong and Chandler, 1973). Armstrong and Chandler (1973), using the physical data for negative pions obtained from Monte Carlo calculations and the model of Katz to predict cell survival, have calculated cell survival for negative pions. Figure 2 shows the theoretical calculations compared with the experimental data (Raju *et al.*, 1972) for negative pions at the peak of the depth-dose distribution when cells are exposed under oxygenated and hypoxic conditions. Although this excellent agreement could be somewhat fortuitous, these calculations may be very helpful in predicting cell survival for various beam configurations of interest for therapy.

BIOLOGICAL EFFECTS OF NEGATIVE PION BEAMS OF INTEREST FOR THERAPY

Negative pion beams of larger momentum spread (~15 to 30%) are necessary to treat large tumors. For such beams, LET differences between the plateau and peak regions are diminished because of the increase in dose contribution at

the peak by energy loss events compared to negative pion absorption processes. For such beams, the RBE is expected to be much smaller than shown in Table 1 for the narrow peak, and the OER is expected to increase somewhat. Professor Lindop and her associates in England, in their preliminary work (private communication, 1973), were unable to detect significant differences in RBE between the plateau and peak when biological systems such as thymic weight loss, spleen weight loss, and colony-forming units (bone-marrow cells) were used at the NIMROD negative pion beam. However, enhancement of RBE at the peak of the NIMROD negative pion beam was observed when *Vicia faba* 10-day root growth and frozen HeLa cells were used.

Experimental data with other mixed LET beams indicate that even a small fraction of the high-LET component is very effective in reducing the OER significantly, although the RBE does not increase accordingly (Raju *et al.*, 1972). Our recent results for various mixtures of x-rays and alpha particles (residual energy ~ 3 MeV) from plutonium also confirm this finding. These results indicate that, although RBE corrections for therapeutically useful negative pion beams may not be large, a significant reduction in OER is expected.

SUMMARY AND CONCLUSIONS

The biological effectiveness for nearly monoenergetic negative pions at the peak of the depth-dose distribution was found to be significantly higher than at the entrance. For therapeutically relevant pion beams of larger momentum spread, the differences in biological effectiveness are reduced considerably; however, a significant reduction in OER still can be expected. Considerable changes in biological effectiveness within the peak region have

been found. These changes will be reduced when opposed fields of negative pions are used. Although the current knowledge on biological effects sheds some light regarding what to expect when negative pion beams are used in therapy, more work needs to be done with intense negative pion beams before their possible application in therapy. Since the increase in biological effectiveness for therapeutically useful negative pion beams is expected to be small, the biological systems selected for pretherapeutic radiobiological work should be able to detect small changes in biological effects in addition to their relevancy to clinical situations.

ACKNOWLEDGMENTS

It is indeed a pleasure to thank Professor P. J. Lindop, Dr. R. J. Berry, Dr. D. H. Reading, and Dr. T. W. Armstrong for giving me permission to quote their unpublished results and for their comments to improve this manuscript. This work was performed under the auspices of the U. S. Atomic Energy Commission.

REFERENCES

- Armstrong, T. W., and Chandler, K. C. (1973): Oak Ridge National Laboratory Report ORNL-TM-4294 (to be published).
- Baarli, J., Bianchi, M., Sullivan, A. H., and Quintalinani, M. (1970): Internat. J. Radiat. Biol. 19, 537.
- Baarli, J., and Bianchi, M. (1972): Internat. J. Radiat. Biol. 22, 183.
- Bewley, D. K. (1968): Radiation Res. 34, 446.
- Burki, H. J., Barendsen, G. W., Raju, M. R., Amer, N. M., and Curtis, S. B. (1969): Lawrence Radiation Laboratory Report UCRL-18793.
- Curtis, S. B. (1971): Radiation Res. 46, 557.
- Feola, J. M., Richman, C., Raju, M. R., Curtis, S. B., and Lawrence, J. H. (1968): Radiation Res. 34, 70.
- Feola, J. M., Raju, M. R., Richman, C., and Lawrence, J. H. (1970): Radiation Res. 44, 637.
- Gnanapurani, M., Raju, M. R., Wolff, S., and Richman, C. (1972): Internat. J. Radiat. Biol. 21, 49.
- Katz, R., Ackerson, B., Homayoonfar, M., and Sharma, S. C. (1971): Radiation Res. 47, 402.
- Mill, A. J., Lewis, J. D., and Lindop, P. J. (1973): Private communication from Professor P. J. Lindop.
- Loughman, W. D., Feola, J. M., Raju, M. R., and Winchell, H. S. (1968): Radiation Res. 34, 56.
- Madhvanath, U. (1971): Effects of Densely Ionizing Radiations on Human Lymphocytes Cultured *in vitro*. Ph.D. thesis, University of California, Report UCRL-20680.

Nias, A. H. W., Greene, D., Major, D., Perry, D. R., and Reading, D. H. (1973):
Brit. J. Radiol. (submitted).

Raju, M. R., Amer, N. M., Gnanapurani, M., and Richman, C. (1970): Radiation
Res. 41, 135.

Raju, M. R., Gnanapurani, M., Stackler, B., Martins, B. I., Madhvanath, U.,
Howard, J., Lyman, J. T., and Mortimer, R. K. (1971): Radiation Res. 47,
635.

Raju, M. R., Gnanapurani, M., Martins, B. I., Richman, C., and Barendsen, G. W.
(1972): Brit. J. Radiol. 45, 178.

Raju, M. R., and Richman, C. (1972): Current Topics in Radiation Research
Quarterly 8, 159.

Richman, S. P., Richman, C., Raju, M. R., and Schwartz, B. (1967): Radiation
Res. Suppl. 7, 182.

Winston, B. M., Berry, R. J., and Perry, D. R. (1973): Brit. J. Radiol. 46,
541.

TABLE 1. RBE of Negative Pions at the Peak of the Depth-Dose Distribution

Biological System	RBE	OER	Reference
<u>BERKELEY</u>			
Abnormal anaphases in <i>Vicia faba</i> root meristems	2.4		Richman <i>et al.</i> (1967)
Chromatid aberrations in <i>Vicia faba</i>	3.0	1.8	Gnanapurani <i>et al.</i> (1972)
<i>Vicia faba</i> 10-day growth	3.0	1.35-1.5	Raju <i>et al.</i> (1970)
Arginine reversions in yeast	1.4	1.9	Raju <i>et al.</i> (1971)
Proliferative capacity of ascites tumor cells	3.0-5.0		Feola <i>et al.</i> (1968, 1970)
Polyploidy induction in ascites tumor cells	2.5		Loughman <i>et al.</i> (1968)
Human kidney cells (frozen)	1.6		Burki <i>et al.</i> (1969)
Human kidney cells (room temperature)	2.4	1.6	Raju <i>et al.</i> (1972)
Human lymphocytes	2.0		Madhvanath (1971)
<u>CERN</u>			
Spermatogonia	3.7		Baarli <i>et al.</i> (1971)
<i>Vicia faba</i> 10-day growth	2.5-4.5		Baarli and Bianchi (1972)
<u>ENGLAND</u>			
<i>Vicia faba</i> 10-day growth	2.5	1.7-1.8	Winston <i>et al.</i> (1973)
HeLa cells (frozen)	1.86		Nias <i>et al.</i> (1973)
HeLa cells (room temperature)	2.5		Mill <i>et al.</i> (1973)

Fig. 1. (a) Schematic depth-dose distribution showing various positions in the peak region where biological materials were exposed; (b) survival curves with T_1 cells measured at Berkeley; (c) 10-day growth of bean roots measured at CERN; (d) 10-day growth of bean roots measured at NIMROD; and (e) survival curves with HeLa cells measured at NIMROD.

Fig. 2. Survival curves of T_1 cells at the peak of the negative pion depth-dose distribution. Calculated results are by Armstrong and Chandler (1973) using the Katz model, and experimental results are by Raju *et al.* (1972).