RESEARCH MEMORANDUM

This is a working paper. It may be expanded,
modified, or withdrawn at any time.

**DO NOT
PHOTOSTAT**

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

I

RESEARCH MEMORANDUM

THE SPECIFIC HEAT OF A DEGENERATE ELECTRON
GAS AT HIGH DENSITY

Murray Gell-Mann

RM-1823-AEC

ASTIA Document Number AD 112391

11 November 1956

Assigned to _____

This is a working paper. It may be expanded, modified, or withdrawn at any time.

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

The **RAND** Corporation

1700 MAIN ST. • SANTA MONICA • CALIFORNIA

ABSTRACT

The methods of the preceding paper (RAND Research Memorandum RM-1802-AEC, "The Correlation Energy of an Electron Gas at High Density," by K. A. Brueckner and M. Gell-Mann) are generalized so that not only the ground state but also the low excited states of an electron gas can be discussed. The energy levels relative to the ground state are the same as those of a gas of independent particles where the energy of each particle (in Rydbergs) is a certain function $W(p)$ of its momentum (expressed in units of the Fermi momentum). The specific heat of the gas at low temperature is proportional to the density of single particle levels at the surface of the Fermi sea, or inversely proportional to $\left(\frac{dW}{dp}\right)_{p=1}$. This last quantity is calculated for high density (small r_s , where density is proportional to r_s^{-3}) and compared to the corresponding quantity for a free electron gas. The ratio is found to be $1 + .083 r_s [-\ln r_s - .203] + \text{higher terms in } r_s$. The expansion is exact and may be compared with the approximate result of Pines, who finds $1 + .083 r_s [-\ln r_s + 1.47] + \dots$

In the preceding paper⁽¹⁾, to be referred to as I, the energy of the ground state of an electron gas is computed at high density. Here we shall treat the energies of low excited states by the same method and thus calculate the low temperature specific heat at high density.

We may label the states by referring them to corresponding states of a free electron gas. The familiar methods of perturbation theory (suitably modified to fit the problem) will convert a given energy eigenstate Φ_n of the system of free electrons into an energy eigenstate Ψ_n of the system with Coulomb interactions among the electrons. We shall be concerned, of course, with the perturbed energy E_n associated with Ψ_n , and not with the unperturbed (purely kinetic) energy associated with Φ_n . Nevertheless, we shall refer to the n 'th state by describing the unperturbed state Φ_n .

Thus we shall speak of the ground state as that in which all one-particle momentum states inside the Fermi sphere are occupied and all those outside are unoccupied. This is, of course, a description of the unperturbed ground state Φ_0 . But when we speak of the energy E_0 of the ground state, we shall mean the perturbed energy associated with the exact eigenstate Ψ_0 of the interacting system.

Similarly, we shall specify an excited state by listing the momenta and spins $\{\underline{p}_j, s_j\}$ (with $j = 1, \dots, \mathcal{N}$) of the vacated one-particle states inside the Fermi sphere and the momenta $\{\underline{k}_j\}$ of the occupied one-particle states outside the sphere. (The momenta are expressed in units of the Fermi momentum P .) Again we are describing an unperturbed state; and again, in speaking of the energy, we shall mean the energy E of the perturbed state.

As in I, we consider a macroscopic sample of gas; the number N of electrons is very large, say $\sim 10^{23}$. We now restrict ourselves to treating those states of the gas in which the number ν of excited particles is small compared to N . At low temperatures, only such states will be important. For these states of the electron gas, the energy E (in units of the Rydberg) may be written in the form

$$E = E_0 + \sum_{j=1}^{\nu} \{W(k_j) - W(p_j)\} + O(\frac{\nu}{N}). \quad (1)$$

This can be done because the interaction energy of the excited particles among themselves is of order ν/N compared to their interaction energy with the rest of the gas. Under our assumptions the term of order ν/N can be neglected, and the specific heat at low temperatures depends only on the form of the function $W(p)$.

Thus for the computation of the specific heat at low temperatures we are justified in treating the interacting electron gas as a system of independent particles obeying Fermi-Dirac statistics and with the energy (in Rydbergs) of each given as a function of momentum (in units of P) by $W(p)$. It is well known⁽²⁾ that under these conditions the specific heat C per electron at constant volume is proportional to the density of "one-particle energy levels" at the surface of the Fermi sphere, that is, proportional to $[(dW/dp)_{p=1}]^{-1}$.

For a free electron gas, we have

$$W_F(p) = \frac{p^2 P^2}{2m} \left[\frac{e^4 m}{2\hbar^2} \right]^{-1} = p^2 / \alpha^2 r_s^2, \quad (2)$$

and

$$\left[(dW_F/dp)_{p=1} \right]^{-1} = \frac{\alpha^2 r_s^2}{2}, \quad (3)$$

where, as in I, $\frac{4}{3} \pi r_s^3$ is the volume per electron (in units of the Bohr radius cubed) and α is $(\frac{4}{9\pi})^{1/3}$. The specific heat of a free electron gas at low temperature is given by a familiar formula⁽³⁾, which we write as follows:

$$C_F(T, r_s) = m^{-1} \hbar^2 e^{-4} K^2 T \cdot \alpha^2 r_s^2, \quad (4)$$

where T is temperature and K is Boltzmann's constant.

Evidently the formula for the low temperature specific heat of an electron gas with interactions is

$$C(T, r_s) = m^{-1} \hbar^2 e^{-4} K^2 T \cdot 2 \left[\left(\frac{dW}{dp} \right)_{p=1} \right]^{-1}. \quad (5)$$

and in (4) we have the special case of free electrons, for which $W = W_F$. Our remaining task is to calculate $W(p)$ (or at least $\frac{dW}{dp}$ at $p = 1$) at high density for an electron gas with Coulomb interactions.

We may think of $W(p)$ in the following way: We consider the ground state of the gas, in which the filled one-particle states are those inside the Fermi sphere. We now examine the change in the energy (exact or perturbed energy) of the system when one electron, with momentum $\vec{p}$, is annihilated. (Clearly $p \leq 1$ here.) Then $W(p)$ is the energy lost in the annihilation, to within any additive constant. Similarly, if we start with the gas in its ground state and create an electron of momentum $\vec{k}$ ($k \geq 1$), then $W(k)$ is the energy gained in the creation. This interpretation of the quantity W is evidently consistent with equation (1).

Now in I we have treated the perturbation series for the ground state energy per particle ϵ (in Rydbergs). The fact that the terms of the series diverge is of no importance, since we have shown how to cure the divergence. In each term of the series for the ground state energy, there are sums over occupied one-particle states with momenta $\vec{p}_i$ such that $p_i \leq 1$ and unoccupied states with momenta $\vec{k}_i$

(equal, say, to $\vec{p}_i + \vec{q}$) such that $k_i > 1$. For example, see equations (9), (10), and (11) of reference 1. (We shall refer to these as I-9, I-10, etc.)

If a particle with momentum $\vec{p}$ and spin s is annihilated, the only change in the perturbation series for the ground state energy consists in amending the condition " $p_i \leq 1$ " for an occupied state by inserting the exception " $\vec{p}_i \neq \vec{p}$ " if $s_i = s$ and amending the condition " $k_i > 1$ " for an unoccupied state by allowing $\vec{k}_i$ to equal $\vec{p}$ if $s_i = s$. The number of occupied states is decreased by one and that of unoccupied ones correspondingly increased. The resulting energy change can best be discussed by means of an example.

One term in the series for the ground state energy per particle is given in equation I - 8:

$$\epsilon_a^{(2)} = - \frac{3}{8\pi^5} \int \frac{d^3 q}{q^4} \int_{p_1 < 1} d^3 p_1 \int_{p_2 < 1} d^3 p_2 \frac{1}{q^2 + \vec{q} \cdot (\vec{p}_1 + \vec{p}_2)} \quad (I-8)$$

$$|\vec{p}_1 + \vec{q}| > 1 \quad |\vec{p}_2 + \vec{q}| > 1$$

If we multiply $\epsilon_a^{(2)}$ by N , the number of electrons, we obtain the corresponding term in the total energy of the ground state, at least to order N . We may write N as $2\mathcal{V} (2\pi)^{-3} 4/3\pi$, where $\mathcal{V}$ is the volume of the gas in units of $\hbar^3 P^{-3}$ and the factor of 2 comes from the two spin states. Furthermore, we may replace the integral over p_1 , say by $\mathcal{V}^{-1} (2\pi)^3$ times a sum over p_1 . We have, then, for the part of the total ground state energy corresponding to $\epsilon_a^{(2)}$:

$$E_a^{(2)} = - 2 \cdot \frac{4}{3} \pi \cdot \frac{3}{8\pi^5} \int \frac{d^3 q}{q^4} \sum_{p_1 < 1} \int_{p_2 < 1} d^3 p_2 \frac{1}{q^2 + \vec{q} \cdot (\vec{p}_1 + \vec{p}_2)} \quad (6)$$

$$|\vec{p}_1 + \vec{q}| > 1 \quad |\vec{p}_2 + \vec{q}| > 1$$

Now let us remove from the gas a particle with spin up and momentum $\underline{p}$.

There are four contributions to the loss of energy:

- (a) For one of the spin states, the term $\underline{p}_1 = \underline{p}$ is dropped from the sum.

(The one-particle state with momentum $\underline{p}$ and spin up is no longer occupied.)

- (b) For one of the spin states, a term $\underline{p}_1 + \underline{q} = \underline{p}$ is added to the sum.

(The one-particle state with momentum $\underline{p}$ and spin up is now among the unoccupied ones.)

- (c) and (d) Corresponding contributions from the sum or integral over $\underline{p}_2$ rather than $\underline{p}_1$. These are evidently equal to (a) and (b).

We have, then, for the contribution $W_a^{(2)}$ of this process to the total energy removed, the expression

$$W_a^{(2)} = -2 \cdot 4/3\pi \cdot \frac{3}{8\pi^5} \left\{ \int \frac{d^3q}{q^4} \int d^3p_2 \frac{1}{q^2 + \underline{q} \cdot (\underline{p} + \underline{p}_2)} \right. \\ \left. \begin{array}{l} |\underline{p} + \underline{q}| > 1 \quad p_2 < 1 \\ |\underline{p}_2 + \underline{q}| > 1 \end{array} \right. \\ \left. - \int \frac{d^3q}{q^4} \int d^3p_2 \frac{1}{\underline{q} \cdot (\underline{p} + \underline{p}_2)} \right\}, \quad (7) \\ \begin{array}{l} |\underline{p} - \underline{q}| < 1 \quad p_2 < 1 \\ |\underline{p}_2 + \underline{q}| > 1 \end{array}$$

where the factor 2 in front is really the product of three factors: 2 from the two spin states as in (6); 1/2 from the selection of spin up only; and 2 from the existence of (c) and (d) in addition to (a) and (b).

By an obvious generalization of this method, we can compute the contribution to $W(p)$ corresponding to each term in ϵ that we investigated in I. The contribution $W_F(p)$ corresponding to the Fermi energy ϵ_F is already given by (2). We must take up next the term $W_X(p)$ corresponding to ϵ_X , the exchange energy. The expression (I-2) for ϵ_X comes from the more explicit formula

$$\epsilon_X = - \frac{3}{8\pi^3} \frac{1}{ar_s} \int_{p_1 < 1} d^3 p_1 \int_{p_2 < 1} d^3 p_2 \frac{1}{(\vec{p}_1 + \vec{p}_2)^2} \quad (8)$$

The method outlined above then gives at once

$$W_X(p) = - 2 \cdot 4/3\pi \cdot \frac{3}{8\pi^3} \frac{1}{ar_s} \int_{p_2 < 1} d^3 p_2 \frac{1}{(\vec{p} + \vec{p}_2)^2} \quad (9)$$

Now let us attempt to compute $\left(\frac{dW}{dp}\right)_{p=1}$ as a series of ascending powers of r_s . The leading term is of course $\left(\frac{dW_F}{dp}\right)_{p=1} = \frac{2}{a^2 r_s^2}$ and the next one should be $\left(\frac{dW_X}{dp}\right)_{p=1}$, presumably of order $\frac{1}{r_s}$. However, this comes out logarithmically divergent⁽⁴⁾. In fact, the situation encountered in I is repeated here, a series of increasing divergences, but each occurring one order earlier than in I.

Corresponding to equation (I-7), we have

$$\left(\frac{dW}{dp}\right)_{p=1} = \frac{2}{a^2 r_s^2} + \frac{1}{r_s} (\log. \text{divergence}) + \frac{1}{r_s^2} (\text{quadratic div.}) + \dots \quad (10)$$

Again, as in I, we expect that the divergent integrals, when summed, will cut themselves off at $q_{\min.} \sim r_s^{1/2}$, so that in place of (10) we will find

$$\left(\frac{dW}{dp}\right)_{p=1} = \frac{2}{a^2 r_s^2} + \frac{1}{r_s} (B \ln r_s + D) + \text{higher terms in } r_s. \quad (11)$$

In order to obtain this form, we must, as in I, sum the leading divergence in each order. Moreover, the leading divergence in each order beyond the first is supplied by precisely the same process as in I, since the cause of divergence is still the piling-up of factors $\frac{1}{q^2}$ coming from successive Coulomb interactions with the same momentum transfer. So from just the processes considered in I, we can obtain exact values of B and D. Moreover, in all orders beyond the first, we may employ the crudest approximation that preserves the leading divergence. In particular, in second order, we may ignore $W_b^{(2)}$ compared to $W_a^{(2)}$, which alone has a quadratic divergence in $(dW/dp)_{p=1}$.

The series that we must examine, then, is $W_x + W_a^{(2)} + W^{(3)} + W^{(4)} + \dots$, where these terms correspond precisely to $\epsilon_x, \epsilon_a^{(2)}, \epsilon^{(3)}, \epsilon^{(4)}$, etc., of I. We have already calculated W_x and $W_a^{(2)}$, and we may now study the derivatives of these terms at $p = 1$. For the first order term W_x we have from (9)

$$W_x = -\frac{1}{\pi^2 a r_s} \int \frac{d^3 q}{q^2} \eta(1 - |\vec{p} + \vec{q}|), \quad (12)$$

where $\eta(x)$ is unity for positive x and zero for negative x . The derivative is then

$$\left(\frac{dW_x}{dp}\right)_{p=1} = \frac{1}{\pi^2 a r_s} \int \frac{d^3 q}{q^2} \delta(1 - |\vec{n} + \vec{q}|) \vec{n} \cdot (\vec{n} + \vec{q}), \quad (13)$$

where $\vec{n}$ is a unit vector in the direction of $\vec{p}$. We may put $\vec{n} + \vec{q}$ equal to a unit vector $\vec{n}'$ pointed into the element of solid angle $d\Omega'$. Then we have

$$\left(\frac{dW_x}{dp}\right)_{p=1} = \frac{1}{\pi^2 a r_s} \int \frac{d\Omega'}{(\vec{n}' - \vec{n})^2} \vec{n} \cdot \vec{n}' = \frac{2}{\pi a r_s} \int_{-1}^1 \frac{x dx}{2(1-x)} \quad (14)$$

We have put $x = \vec{n} \cdot \vec{n}'$. Note the result is indeed logarithmically divergent.

We have treated $\left(\frac{dW}{dp}\right)_{p=1}$ without approximation. In the higher terms, however, we keep only the leading divergence. Now in differentiating formula (7) for $W_a^{(2)}$, we keep the leading divergence if we differentiate with respect to p only the limits of integration on $\vec{q}$: " $|\vec{p} + \vec{q}| > 1$ " and " $|\vec{p} - \vec{q}| < 1$." We find, in fact,

$$\left(\frac{dW_a^{(2)}}{dp}\right)_{p=1} \approx -\frac{2}{\pi^4} \int_{p_2 < 1} d^3 p_2 \int_{|\vec{p}_2 + \vec{q}| > 1} d^3 q \frac{1}{q^4} \times$$

$$\times \frac{1}{q^2 + \vec{q} \cdot (\vec{n} + \vec{p}_2)} \delta(1 - |\vec{n} + \vec{q}|) \vec{n} \cdot (\vec{n} + \vec{q}), \quad (15)$$

where the factor of two comes from the existence of two terms in (7), which make equal contributions to (15). As before, we put $\vec{n} + \vec{q} = \vec{n}'$ and obtain

$$\left(\frac{dW_a^{(2)}}{dp}\right)_{p=1} \approx -\frac{2}{\pi^4} \int \frac{d\vec{n}'}{[(\vec{n}' - \vec{n})^2]^2} \vec{n} \cdot \vec{n}' \int_{p_2 < 1} d^3 p_2 \frac{1}{(\vec{n}' - \vec{n}) \cdot (\vec{n}' + \vec{p}_2)} \cdot \quad (16)$$

$$|\vec{n}' - \vec{n} + \vec{p}_2| > 1$$

We are still interested only in the leading divergence at $\vec{n}' \approx \vec{n}$ and so we may take the limit $\vec{n}' - \vec{n} \rightarrow 0$ in the integral over $\vec{p}_2$. The limit is 2π . Thus we have

$$\left(\frac{dW_a^{(2)}}{dp}\right)_{p=1} \approx -\frac{8}{\pi^2} \int_{-1}^1 \frac{x dx}{[2(1-x)]^2} \cdot \quad (17)$$

Adding together the terms calculated so far in (3), (14) and (17), we have

$$\left(\frac{dW}{dp}\right)_{p=1} = \frac{2}{\alpha^2 r_s^2} + \frac{2}{\pi \alpha r_s} \int_{-1}^1 \frac{x dx}{2(1-x)} \left\{ 1 - \frac{4\alpha r_s}{\pi} \frac{1}{2(1-x)} + \dots \right\}. \quad (18)$$

To complete this series to the desired accuracy, we must look at formula (I-19) for $\epsilon^{(n)}$ and use it to calculate $W^{(n)}$ for $n > 2$. If we do that and compute the leading divergence in $(dW^{(n)}/dp)_{p=1}$, we find that the series in (18) becomes simply a geometric series. We have, thus, with sufficient accuracy to obtain exact values of B and D in equation (11), the result:

$$\left(\frac{dW}{dp}\right)_{p=1} \approx \frac{2}{\alpha^2 r_s^2} + \frac{2}{\pi \alpha r_s} \int_{-1}^1 \frac{x dx}{2(1-x)} \frac{1}{1 + \frac{4\alpha r_s}{\pi} \frac{1}{2(1-x)}}, \quad (19)$$

or

$$\left(\frac{dW}{dp}\right)_{p=1} = \frac{2}{\alpha^2 r_s^2} + \frac{1}{\pi \alpha r_s} \left\{ \ln \pi / \alpha r_s - 2 \right\} + \dots \quad (20)$$

At low temperature, then, the specific heat of a free electron gas is modified through Coulomb interactions by the factor

$$C/C_F = \left[1 + \frac{\alpha r_s}{2\pi} (-\ln r_s + \ln \pi / \alpha - 2) + \dots \right]^{-1}, \quad (21)$$

where the expansion is valid at high densities. Inserting numerical values, we have

$$C/C_F = \left[1 + .083 r_s (-\ln r_s - .203) + \dots \right]^{-1}. \quad (22)$$

This is to be compared with the approximate result of Pines⁽²⁾, who finds

$$C/C_F = \left[1 + .083 r_s (-\ln r_s + 1.47) + \dots \right]^{-1}. \quad (23)$$

The method given here permits the computation of higher terms in the series, and the next correction is now being calculated. Applications to the specific

heats of metals are also being studied.

The author would like to thank Dr. Richard Latter of the RAND Corporation and Dr. Keith A. Brueckner of the University of Pennsylvania for many interesting discussions.

REFERENCES

1. Gell-Mann, M. and K. A. Brueckner, The Correlation Energy of an Electron Gas at High Density, RM-1802-AEC, October 3, 1956.
2. D. Pines, Solid State Physics, 1, 367 (1955).
3. F. Seitz, Modern Theory of Solids (McGraw-Hill Book Company, Inc., New York and London, 1940), p. 150.
4. J. Bardeen, Phys. Rev., 50, 1098 (1936). That the divergences would disappear in a correct calculation was pointed out by E. P. Wigner, Trans. Faraday Soc., 34, 678 (1938).