THEORETICAL PARTICLE PHYSICS

Progress Report

Shu-Yuan Chu, Archibald W. Hendry and Don B. Lichtenberg

Physics Department, Indiana University

Bloomington, Indiana 47401

December 1, 1973 - November 30, 1974

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

Prepared for the U. S. Atomic Energy Commission under Contract No.

AT (11-1) - 2009, Task B.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

87

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

ABSTRACT

The major part of our research activity has been on the phenomenological treatment of two-body reactions. We have calculated the differential cross section and polarization for proton-proton elastic scattering, pion-nucleon elastic scattering, and the reaction $pp \rightarrow \pi^+ d$. We have considered optical models, Regge pole models in both t - and s -channels, a parton model, and a phase-band method. We have also considered the parametrization of strong-interaction resonances, taking into account how threshold effects can shift the masses and widths of resonances from their apparent values. Other topics treated were $SU(3)$ -breaking mass formulas, a bootstrap scheme for particle interactions based on an eigenvalue condition for the fine-structure constant in non-abelian gauge theories, and a geometric interpretation of gauge fields in an $SU(4)$ invariant theory of particle interactions.

This Report describes the research work performed during the past year or so which was supported by the AEC Contract No. AT(11-1) - 2009, Task B. Below, we discuss in turn the various topics in which we have been actively engaged. These topics cover a wide range of subjects. As will be seen, a considerable part of our work has involved close collaboration throughout the year with the experimental side (Task A) of this Contract, while other activities range from the practical analysis of resonances to the more esoteric areas of parton models and non-abelian gauge theories.

1. A STUDY OF PROTON-PROTON ELASTIC POLARIZATION (Hendry, with G. Abshire)

During the past year, the experimental group (Task A) of this Contract have been analyzing their large t proton-proton elastic scattering measurements, taken earlier at the Argonne ZGS with a polarized proton target and beam momenta 5.15, 7.0 and 12.33 GeV/c. Their results¹ have turned out to be exceedingly interesting, since one can observe the steady development of a succession of dip structures in the polarization as the beam momentum is increased; at 12.33 GeV/c, there seem to be three definite structures (possibly double zeros) located at $-t \approx 0.8, 2.3$ and about 5 (GeV/c)². (These results are particularly interesting to us since it is just what we had predicted before the experiment was done, on the basis of an optical approach; indeed we gave some guidance to the experimentalists as regards where to look for the dips.)

We have examined² these new pp polarization data using several theoretical models in an effort to understand the origin of the observed structure. These models included

- (1) A standard Regge model by Rarita et al. which uses $P, P', P'', \omega, \omega'$ trajectories with various kinds of ghost-killing mechanisms.
- (2) A Regge model, proposed by Booth et al., which incorporates "linear exchange degeneracy breaking."
- (3) Several Regge models with cyclic residue functions, as first suggested by Barger and Phillips.
- (4) The Chou-Yang eikonal optical model, as developed by Durand and Lipes.
- (5) An optical model developed by Chu and Hendry.

We found that it was difficult to get a good fit to the polarization data with any of the Regge models, whereas optical models were rather more successful. It strongly suggests the interesting possibility that optical ideas, which seem to work at ISR energies, may well be applicable also even at these much lower energies.

2. REGGE MODEL PREDICTION FOR THE REACTION $pp \rightarrow \pi^+ d$ (Lichtenberg, with R. M. Heinz and R. Shtokhamer)

During the year, we have begun a Regge model calculation of the differential cross section and proton polarization for the reaction $pp \rightarrow \pi^+ d$. There are as yet no data for the polarization, but with the good polarized proton targets now available, the experiment is feasible to do and may possibly be done in the coming year at Argonne by Task A. Our calculation is based on the formalism of H. Lee,³ but we have generalized the form of the Regge residues. The reason for this is that in Lee's formalism, used in a previous calculation of Barger and Michael,⁴ the predicted polarization

is zero. In our model, the polarization is in general non zero. If we restrict ourselves to the contributions from the N_α and N_γ Regge trajectories, we find that the polarization has the characteristic feature of vanishing at wrong-signature nonsense zeros of both trajectories. This qualitative feature is independent of the details of the model, and lends itself to an experimental test. These qualitative results form part of the proposal for the $pp \rightarrow \pi^+ d$ experiment at Argonne. This work is still in progress.

3. PARTON MODEL PREDICTIONS AND LARGE ANGLE HADRON-HADRON SCATTERING (Hendry)

Parton models⁵ have very simple predictions for large angle scattering, namely that the differential cross-section should behave like $d\sigma/dt \sim s^{-N} f(\theta)$, where N depends on the particular model as well as on the particular scattering process involved. This power law dependence for fixed θ is easily testable in the case of elastic scattering, where there is a reasonably plentiful supply of experimental data.

To date, we have examined the situation for elastic proton-proton and elastic π^+ -proton scattering. We find that, when $(d\sigma/dt)_\theta$ is plotted against $\ln s$, the data in both cases do not lie on straight lines, as a previous cursory examination by Landshoff and Polkinghorne⁶ seemed to indicate. The data for fixed θ in fact seems to contain a sequence of breaks.

This result is particularly important in view of the present-day great interest and activity in parton models. Structure in the large angle differential cross-section is certainly unexpected from these simple parton models, and one has to conclude that they still lack some important ingredients.

Having drawn attention to this structure, we have attempted to give an explanation for its origin. By plotting the same data in a different way, namely $(d\sigma/dt)_\theta$ versus t , we show that the observed structure occurs at or near the oscillations of Bessel functions. This leads to a simple interpretation and to the important conclusion that strong diffractive (and hence coherent parton) effects are present even at large angles, and cannot be ignored in any description of large angle scattering.

4. MASSSES AND WIDTHS OF RESONANCES (Lichtenberg)

The masses and widths of elementary particle resonances listed in the latest tables of the Particle Data Group⁷ show errors which are sometimes larger than necessary. In some cases, the reason the errors listed in the tables are so large is that different experiments to determine the masses and widths are seemingly incompatible. In fact, however, the different values sometimes result from differences in analysis rather than incompatible data. As an example, we have examined⁸ the recent precise data^{9,10} on the masses and widths of the $\Sigma^*(1380)$. In the paper of Borenstein et al.,¹⁰ the mass of the Σ^{*-} is given as 1383 ± 2 MeV, while in the paper of Baltay et al.,⁹ the mass is given as 1391 ± 1 MeV. These results do not agree within the stated errors. However, Borenstein et al. found the position of the pole of the relevant amplitude in the complex energy plane, but Baltay et al. did not. When we use the experimental results of Baltay et al. to compute this pole, we obtain a mass of 1383 ± 1 MeV, a result in complete agreement with that of Borenstein et al. In other cases, the improvement is not so marked, but it should be possible to determine approximately how much of the discrepancies between different experiments arise from experimental errors and how much from differences in

analysis.

We have also examined the effects of thresholds on the masses and widths of resonances. A resonant amplitude near threshold is commonly described phenomenologically by a Breit-Wigner form with an energy-dependent width. In this case, the true mass and width (as determined by the position of the pole of the amplitude) are smaller than the mass and width parameters used in describing the amplitude. We have found⁸ that the mass correction goes as the square of the width, while the width correction goes like the width cubed, and have obtained simple formulas to estimate the size of the corrections.

Previously, we had proposed¹¹ a new parametrization for a resonance which incorporates the threshold behavior of the amplitude and still gives the position of the pole directly. Recently Borenstein et al.¹⁰ have applied this parametrization to the $\Sigma^*(1380)$ and found it better than other parametrizations commonly used.

5. UNITARY-SYMMETRY BREAKING OF BARYON MASSES (Lichtenberg)

The recent precise measurements of the masses of the members of the baryon decimet have aroused new interest in baryon mass-breaking formulas. Borenstein et al.¹⁰ have noticed that the baryon decimet masses agree better with the Gell-Mann-Okubo mass formula if squares of masses are used in the relation rather than linear masses. Oneda and Takasugi¹² have argued that this fact means that quadratic mass formulas should be used for all baryons. We have examined¹³ the agreement of the octet and decimet mass formulas with experiment, using pole values for the masses. Our conclusion is that although the quadratic formula agrees better with experiment for the decimet than the linear formula, the linear and quadratic forms both fail about

equally for the octet. It was also found that the Okubo second-order decimet formula agrees with experiment either in mass or mass-squared. It is therefore still not clear from experiment whether a linear or quadratic mass formula should be used for baryons.

6. A PARAMETRIZATION FOR PERIPHERAL SCATTERING (Chu and Hendry)

It has become increasingly clear over the past two or three years that the concept of peripheral scattering is very important in particle collisions. Possibly the clearest example of this is the well-known cross over effect in forward differential cross-sections, which occurs (even for rather low energies) in the vicinity of $-t \approx 0.2 \text{ (GeV/c)}^2$. Simple t-channel Regge pole models expect the crossover to occur at $-t \approx 0.6 \text{ (GeV/c)}^2$ where the ρ , ω trajectories pass through zero; thus to get the crossover at the experimentally observed position, one needs to introduce absorbing cuts. (Essentially, these cuts reduce the low partial wave component of the amplitudes, hence making these amplitudes relatively more peripheral.)

Alternatively however, one can ascribe the crossover effect directly to peripheral s-channel scattering, and as pointed out for example by Harari,¹⁴ an impact parameter s-channel picture immediately gives a contribution of the form $J_0(R\sqrt{-t})$; the crossover is then associated with the first zero of the J_0 Bessel function.

In the present work, we have attempted to obtain a description of peripheral scattering which is based directly on an s-channel approach, rather than the more orthodox method of absorbed t-channel Regge poles. Actually this is a fairly well-known problem in several areas of physics, for example classical

electromagnetic theory and nuclear physics, and we have relied on this previous work to develop a formalism for peripheral scattering appropriate to high energy physics. The classical form for peripheral scattering (obtained originally long ago by Sommerfeld and subsequently used by Regge in his historic papers on the Schrödinger equation) is $P_{\alpha(s)}(-\cos\theta_s)/\sin \pi\alpha(s)$. This is of course similar in appearance to the t-channel Regge pole expression; however, there are some crucial differences. In particular, since we are in the s-channel physical region, $\alpha(s)$ is complex (being an analytic function of s), and in order that the amplitude corresponds to peripheral scattering, the real part of $\alpha(s)$ must be kR .

After examining the spinless case, we extended our consideration to the more realistic case of meson-baryon scattering, paying particular attention to analyticity and the crossing properties of the amplitudes. Comparison with forward pion-nucleon charge exchange scattering yielded reasonable agreement with experiment. Our results show that a direct s-channel approach to peripheral scattering seems quite feasible as an alternative to the usual t-channel approach with poles and absorbing cuts. (V. Barger also mentioned this possibility at the recent international London conference; some related work has been done recently by B. and F. Schrempp.¹⁵⁾

7. A PHASE SHIFT ANALYSIS OF PION-NUCLEON SCATTERING FROM 1.6 TO 5.0 GeV/c (Hendry)

In recent years, a tremendous amount of experimental data for pion-nucleon scattering has accumulated in the energy range up to 5 GeV/c. There are several reasons why it is important that we understand what is

happening in this region. For example, we will be able to determine what higher mass resonances there are, thus providing a good test of models for baryon spectra, particularly the harmonic oscillator quark model which has been so successful to date. Also, this is a crucial transitional energy region: the lower energies are dominated by a few resonances, whereas by 5 GeV/c all the important structures that we normally associate with high energies (dips, etc.) have already set in. It will be extremely useful to try to understand how this transition takes place.

A usual type of phase shift analysis however is impractical - a large number of partial waves is required, which means a hundred or more parameters. We have therefore adopted the technique of using angular momentum distributions (making the assumption that ℓ -distributions for $\delta(\ell)$, $\eta(\ell)$, or equivalently amplitude $f(\ell)$, are smooth). This in a sense is a standard method, often used in nuclear physics, being particularly applicable where no one partial wave dominates but where scattering effects are due to a large number of similar partial waves. (The idea has been used to a small extent before in high energy physics, for example by Moravcsik et al.¹⁶ in their phase band method.)

This work is in progress. We have already obtained preliminary fits to the new charge-exchange data in this region, and are at present combining this with elastic $\pi^\pm$ -proton scattering. The method looks very promising, and should it be successful, it would provide a much needed useful tool to analyze data for other processes in this important energy range.

8. BOOTSTRAP SCHEME FOR PARTICLE INTERACTIONS BASED ON AN EIGENVALUE
CONDITION FOR THE FINE-STRUCTURE CONSTANT IN NON-ABELIAN GAUGE
THEORIES (Chu)

In this work, it is assumed that the new renormalization-group equations derived by Weinberg¹⁷ allow a non-trivial bootstrap solution besides the trivial free-field solution. Specifically, we start from a Lagrangian of N massless spin- $\frac{1}{2}$ fields with interactions invariant under a gauge group $SU(N')$, $1 < N' < N$ and characterized by a single coupling constant. The bootstrap scheme assumes the existence of a non-trivial zero of the function β and that the physical coupling constant α is fixed at the non-trivial zero of β . This eigenvalue condition¹⁸ serves to determine uniquely not only α but N , N' and the nature of the fermion representation. Furthermore, with the eigenvalue condition satisfied, we can set the unrenormalized coupling constant and gauge parameter equal to zero and still have a non-trivial solution, thereby achieving a truly bootstrap situation. The gauge symmetry $SU(N')$ is then spontaneously broken to leave only a $U(1)$ gauge group of electric charge unbroken. Fermion masses within the same $SU(N')$ representation are not split. The gauge vector bosons, except the photon, acquire heavy masses related to the fermion mass differences. All the dimensionless quantities in the bootstrap scheme become calculable in principle. In particular there is a direct way of calculating α assuming perturbation methods are applicable. The asymptotic behavior of the bootstrap scheme is that of broken scale invariance, with deviations from canonical scaling characterized by small anomalous dimensions of order α . The validity of the bootstrap assumptions and the reliability of the perturbation methods

can therefore be tested in deep-inelastic lepton-hadron scattering experiments.

9. GEOMETRICAL INTERPRETATION OF THE GAUGE FIELDS IN AN $SU(4)$ INVARIANT THEORY OF PARTICLE INTERACTIONS (Chu)

The assumption that the elementary particle interactions are $SU(4)$ gauge invariant is a natural way to reconcile the success of the Cabibbo theory with the absence of the strangeness changing neutral current. We have shown that the gauge fields in an $SU(4)$ gauge invariant theory can be given a geometrical interpretation based on the formulation of the principle of general relativity using the conformal group.¹⁹ The present theory contains Einstein's theory of gravitation as a special case in the absence of other interactions and with quantum effects neglected.²⁰ The theory suggests that $SU(4)$ gauge invariance may be the only one consistent with macroscopic phenomena, and may lead to a solution of the long standing problem of constructing a renormalizable quantum field theory of gravitation. We are at present examining the detailed consequences of the present proposal, as well as its overall consistency.

The three principal investigators are spending 100% of their time during two of the summer months and 30% of their time during the academic year, on the research project.

A number of theoretical physicists from outside institutions have made short-term visits of a day or two to Indiana University during the year to discuss problems of interest to the principal investigators.

These visitors were

Dr. S. J. Chang, University of Illinois

Dr. H. Jay Melosh, University of Chicago

Dr. W. Zimmerman, New York University

Dr. J. Sullivan, University of Illinois

Dr. T. K. Kuo, Purdue University

Dr. S. P. Rosen, Purdue University

Dr. Jan Dash, Argonne National Laboratory

Dr. James Trefil, University of Virginia

Dr. E. Fischbach, Purdue University

Dr. J. Rosner, University of Minnesota

REFERENCES

1. G. W. Abshire et al., Phys. Rev. Lett 32, 1261 (1974).
2. A. W. Hendry and G. W. Abshire, to be published in Phys. Rev.
3. H. Lee, Phys. Rev. 174, 2130 (1968).
4. V. Barger and C. Michael, PRL 22, 1330 (1969).
5. For a review, see S. Brodsky, Proceedings of Conference on High Energy Collisions at Stony Brook (1973), ed. by C. Quigg.
6. P. Landshoff and J. C. Polkinghorne, Phys. Lett. 443, 293 (1973).
7. Particle Data Group: V. Chaloupka et al., Phys. Lett. 50B, 1 (1974).
8. D. B. Lichtenberg, Report No. COO-2009-80.
9. C. Baltay et al., in Baryon Resonances-73, Purdue University, Lafayette, Indiana (1973); M. Habibi, Thesis, Columbia University, Nevis-199 (1973).
10. S. R. Borenstein et al., Phys. Rev. D9, 3006 (1974).
11. D. B. Lichtenberg, in Baryon Resonances-73, C. Fowler, ed., Purdue University, Lafayette, Indiana (1973), Report No. COO-2009-52;
D. B. Lichtenberg, Lett. Nuovo Cimento 7, 727 (1973), Report No. COO-2009-53.
12. S. Oneda and E. Takasugi, University of Maryland Technical Report No. 74-103 (1974).
13. D. B. Lichtenberg, Ninth Balaton Symposium on Particle Physics, Balatonfured, Hungary, June 12-18, 1974, Report No. COO-2009-71.
14. H. Harari, Phys. Rev. Lett. 26, 1400 (1971); Phys. Lett. 35B, 239 (1971).
15. B. Schrempp and F. Schrempp, CERN Th. 1812 (1974).
16. M. Moravcsik et al., Phys. Rev. Lett 25, 771 (1970).

17. S. Weinberg, Phys. Rev. D8, 3497 (1973).
18. For systematic discussion of the eigenvalue condition in quantum electrodynamics, see S. L. Adler, Phys. Rev. D5, 3021 (1972).
19. Preliminary concepts of the present approach are contained in Sec. IV of the report, New Approach to Five-Dimensional General Relativity, Shu-Yuan Chu, Report No. C00-2009-46 (1973).
20. The present approach is related, but with different fundamental physical assumptions, to that of C. N. Yang, Integral Formalism for Gauge Fields, University of Wroclaw and State University of New York, Stony Brook, preprint (1974).

RESEARCH PAPERS SUPPORTED BY AEC CONTRACT NO. AT(11-1) - 2009, TASK B

1. Analysis of $\pi^+ p$ Elastic Scattering from 2.75 to 4.0 GeV/c at all Angles (S.-Y. Chu and A. W. Hendry), Phys. Rev. D4, 2743 (1971).
2. 5 GeV/c Pion-Nucleon Scattering at all Angles (S.-Y. Chu and A.W. Hendry), Phys. Rev. D4, 3282 (1971).
3. Duality Diagrams with Exotic t-channel Exchanges in Meson-Baryon Scattering (S.-Y. Chu and P. Chylek), Lett. Nuovo Cimento 2, 229 (1971).
4. On the Equivalence of Regge and Optical Models (A. W. Hendry), Phys. Rev. D5, 215 (1972).
5. Proton-Proton Polarization Data: Possible Evidence for Exotic Peripheral Resonances (S.-Y. Chu and A. W. Hendry), Phys. Rev. D6, 190 (1972).
6. A Unified Picture of Proton-Proton Elastic Scattering at All Angles from 3 to 24 GeV/c (T. Y. Cheng, S.-Y. Chu and A. W. Hendry), Phys. Rev. D7, 86 (1973).
7. Regge and Optical Models (A. W. Hendry), Proceedings of Argonne ZGS Workshop, Vol. 1 (1971).
8. Violation of CP Invariance in the Dyon Model (S.-Y. Chu), Phys. Rev. D7, 853 (1973).
9. Resonances and Background Contributions to Meson-Nucleon Total Cross-Sections (P. Chylek and D. B. Lichtenberg), Lett. Nuovo Cimento 5, 435 (1972).

10. Unitarity Corrections to Approximate Amplitudes in Multichannel Scattering (D. B. Lichtenberg), *Particles and Nuclei* 5, 119 (1973).
11. Are Regge and Optical Models Equivalent, by an Amplitude Rotation? (A. W. Hendry), *Phys. Rev.* D7, 910 (1973).
12. Comment on the Parameters of the 3-3 Resonance (T. Y. Cheng and D. B. Lichtenberg), *Phys. Rev.* D7, 2249 (1973).
13. A New Approach to 5-Dimensional General Relativity (S.-Y. Chu), submitted to *Phys. Rev.*
14. On the Widths and Elasticities of Meson-Baryon Resonances (D. B. Lichtenberg), in *Baryon Resonances - 73*, Earle C. Fowler, ed., Purdue University, Lafayette, Indiana, 465 (1973).
15. Parametrization of a Broad Resonance (D. B. Lichtenberg), in *Baryon Resonances - 73*, Earle C. Fowler, ed., Purdue University, Lafayette, Indiana, 23 (1973).
16. Mass and Width of a Broad Resonance (D. B. Lichtenberg), *Lett. Nuovo Cimento* 7, 727 (1973).
17. Veneziano Model and the Elasticities of Pion-Nucleon Resonances (D. B. Lichtenberg), contributed to Berkeley Conference of the Division of Particles and Fields, August (1973), to be published in *Particles and Nuclei* (1974).
18. Theoretical Models for Proton-Proton Elastic Polarization (A. W. Hendry and G. W. Abshire), accepted for publication in *Phys. Rev.* (1974).
19. Evidence for Coherent Effects in Large Angle Hadron-Hadron Collisions (A. W. Hendry), accepted for publication in *Phys. Rev.* (1974).

20. Peripheral Scattering and an S-Channel Regge Approach to Particle Scattering (S.-Y. Chu and A. W. Hendry), submitted for publication.
21. Baryon Decimet Mass Splittings (D. B. Lichtenberg), presented at the Ninth Balaton Symposium on Particle Physics, Balatonfured, June (1974); COO-2009-71.
22. Bootstrap Scheme for Particle Interactions Based on an Eigenvalue Condition for the Fine Structure Constant (S.-Y. Chu), submitted to the 17th International Conference on High Energy Physics, London, July (1974); COO-2009-78.
23. Structures in Proton-Proton Elastic Scattering and their Interpretation (A. W. Hendry), invited talk at Argonne ZGS Workshop on Polarization Phenomena, July (1974); COO-2009-86.