I

U N C L A S S I F I E D

UR-463

Health and Biology

THE UNIVERSITY OF ROCHESTER
Atomic Energy Project
P. O. Box 287, Station 3
Rochester 20, New York

* * *

Contract W-7401-eng-49 between the U. S. Atomic
Energy Commission and the University of Rochester,
administered by the Department of Radiation Biology
of the School of Medicine and Dentistry.

LOW LEVEL POLONIUM DETERMINATION OF TISSUE AND URINE

by

Stuart C. Black

Division: Radiology and Biophysics

Division Head: W. F. Bale

Section: General Biophysics

Section Head: W. F. Bale

Date Completed: 10/5/56

Date of Issue: 10/18/56

U N C L A S S I F I E D

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

TABLE OF CONTENTS

ABSTRACT.	vi
LIST OF FIGURES	vii
LIST OF TABLES	viii
I. INTRODUCTION	1
II. CHARACTERISTICS OF POLONIUM	6
III. REVIEW OF PREVIOUSLY REPORTED PROCEDURES	8
A. Procedure of Scott and Stannard (HNO_3 , H_2SO_4)	8
B. Procedure Used by Fink (HNO_3 , HClO_4)	10
C. Procedure Used by Comber, <u>et al</u> (HNO_3 , H_2O_2)	11
D. Procedure of Mulhaney, <u>et al</u> (HClO_4 , H_2O_2)	12
E. Procedure Used by Rundo (coppt. with Te)	13
F. Comment on the Digestion Procedures	14
IV. APPARATUS USED	15
A. Counting Equipment	15
B. Plating Apparatus	17
V. PRELIMINARY EXPERIMENTS ON THE RECOVERY OF POLONIUM	22
A. Per cent Recovery as a Function of Activity Level	22
B. Per cent Recovery as a Function of Plating Time	23
C. Absolute Recovery of Polonium	24
D. Effect of Nitric Acid on Recovery of Polonium	25

E. Effect of Undissolved Solids on Recovery of Polonium	27
F. Effect of Ferric Ion on Recovery	30
VI. RECOVERY OF SMALL AMOUNTS OF POLONIUM FROM URINE	30
A. Introduction and Survey of Previously Reported Procedures	30
B. Plating of Urinary Polonium on Silver, Nickel, and Copper	32
C. Recoveries from Urine Using Nickel Powder and Nickel Filings	33
D. Recovery from Large Volumes of Urine Using Nickel Powder	36
E. Recovery from Large Volumes of Urine by Concentration	39
F. Recovery of Polonium from the Urine of Polonium Injected Rats	41
G. Method Developed for the Recovery of Polonium from Large Volumes of Urine	45
H. Polonium in the Urine of Non-exposed Personnel	47
1. Method used.	47
2. Procedure	47
3. Discussion of the results	49
4. Calculations on expected urinary excretion of polonium	51
I. Summary and Conclusions on Section VI	56

VII.	RECOVERY OF POLONIUM FROM SOFT TISSUES AND BONE . . .	57
A.	Experiments Based on Nitric Acid Digestion . . .	57
1.	Recovery from mice injected 24 hours previously	57
2.	Recoverable polonium from non-exposed mice .	60
3.	Analysis of bats for polonium	62
4.	Recovery of added polonium from bone	63
a.	Fresh bone	63
b.	Ashed bone	64
c.	Fresh bone with fat extraction	66
5.	Recovery of added polonium from fat	69
6.	Discussion of Part A.	70
B.	Method for Analysis of Soft Tissue and Bone for Polonium	71
1.	Description of the method	71
2.	Natural polonium activity of beef bone	73
C.	Studies on Rats Killed Immediately after Radon Exposure	74
1.	Recovery of polonium from rats	74
2.	Calculations on the amount of polonium in the rats.	77
3.	Discussion	80
D.	Studies on Mice Exposed to Radon and Allowed to Live until Death	81
1.	Mice exposed to 1 millicurie-hour/liter . . .	81
2.	Calculations on the amount of polonium in the mice.	84

3. Polonium Content in the Bones of Mice Exposed to Radon	87
a. Introduction	87
b. Procedure	87
c. Discussion	88
d. Calculations	90
E. Studies on Human Autopsy Material	91
1. Introduction	91
2. Experiments with formaldehyde	92
3. Experiments with bone	96
a. Procedure	96
b. Results	96
c. Discussion	98
4. Experiments with lung tissue	99
VIII. SUMMARY AND CONCLUSIONS	101
BIBLIOGRAPHY	108
APPENDIX. Decontamination Procedure	113

ABSTRACT

Methods were developed for analysis of up to 50 grams of soft tissue and bone and liter volumes of urine for polonium content of the order of 1 dpm above background. Bone and lung tissue obtained at autopsy of a uranium miner contained approximately twenty times as much polonium as similar material from non-miners. Mice and rats exposed to radon were analyzed for polonium. Calculations gave polonium activities of the order of magnitude found by analysis of these animals. The data indicate that approximately 50% of the polonium found in mice was in the skeleton, suggesting that it was derived from radiolead stored therein as a result of exposure to radon approximately 120 days previously.

LIST OF FIGURES

<u>Figure Number</u>	<u>Page</u>
1. Decay scheme--Radon to stable lead	4
2. Plating equipment	19
3. Plating bottle and stirring rod	21
4. Apparatus for boiling off large volumes of fluids	42
5. Glass cage for polonium injected mice	59
6. Build-up and decay of radon in fat of rats sacrificed immediately after exposure	79
7. Build-up of radon (1), maintenance of level (2), and blowoff (3) in tissues of mice allowed to live after exposure	79

LIST OF TABLES

<u>Table Number</u>	<u>Page</u>
I. Recovery of polonium from 0.5N HCl as a function of radioactivity level	23
II. Recovery of polonium from 0.5N HCl as a function of plating time	23
III. The same amount of polonium as in Table II deposited on a silver foil and air-dried gently under an infra-red lamp	24
IV. The effect of nitric acid on the recovery of polonium from 0.5N HCl	25
V. The effect of various treatments to improve recovery of polonium when the silver foil has been discolored during plating	27
VI. Effect of undissolved solids on the recovery of polonium	29
VII. Plating of added polonium from acidified urine onto various metals	32
VIII. Recovery of added polonium from acidified urine and 0.5N HCl by various treatments. Volumes of approximately 200 ml	35
IX. Recovery of added polonium from acidified urine. Comparison of nickel powder and nickel filings	37
X. Recovery of added polonium from acidified urine by evaporation of large volumes	40
XI. Effect of wet digestion on the recovery of polonium from the urine of polonium injected rats	44
XII. Recovery of polonium from liter volumes of tap and distilled water, and urine from unexposed personnel	49
XIII. The recovery of polonium from mice sacrificed and analyzed 24 hours after IP injection of polonium	60

XIV.	Polonium content of non-exposed mice of the local colony .	61
XV.	The polonium content of two bats trapped in a uranium mine	63
XVI.	The recovery of added polonium from bone after digestion with nitric and nitric-sulfuric acid	65
XVII.	The recovery of added polonium from bone after various digestion procedures with and without the extraction of fat	68
XVIII.	The polonium content of beef femur samples used in these experiments	74
XIX.	The polonium content of two rats exposed to a daughter-free radon level of 125 $\mu\text{c}/\text{liter}$ for 3.5 hours, sacrificed immediately, and analyzed 246 days after exposure .	76
XX.	The polonium content of mice exposed to a daughter-free radon atmosphere of 200 $\mu\text{c}/\text{liter}$ for 5 hours, living a nearly normal life span. Analyzed 774 days after exposure	83
XXI.	Polonium content of mice which had been exposed to a daughter-free radon concentration of 200 $\mu\text{c}/\text{liter}$ for 15 hours and had died of this exposure in 90 to 100 days. For three of these, the bones were analyzed separately from the soft tissue. Analysis was approximately one month after death	89
XXII.	The change in weight of bone and lung after soaking 8 days in a 10% solution of formaldehyde in water	93
XXIII.	The polonium content of formaldehyde solutions used to preserve various polonium containing samples	95
XXIV.	The polonium content of bone from a uranium miner compared with bone from people with no history of radium or radon exposure	97
XXV.	The polonium content of lung from a uranium miner compared with lung from a person with no history of radium or radon exposure	100
XXVI.	The polonium content of various samples from presumably unexposed animals	106

I. INTRODUCTION

In recent years, increased mining and processing of uranium has subjected numerous individuals working in this field to concentrations of radon and its short-lived daughters that are potentially toxic (1).

In addition, recent work has shown that even in mines not worked for uranium or radium potentially toxic concentrations may exist (2,3).

The permissible level of radon and its daughters has been subject to frequent change and some controversy. Based on recommendations by Evans and Goodman (4), a permissible working level of 1×10^{-11} curies/liter of radon was at one time widely used in the radium dial painting industry. The last published U.S. Department of Commerce Handbook 52, "Maximum permissible amounts of radioisotopes in the human body and maximum permissible concentrations in air and water," (5) gives a permissible level of 1×10^{-11} curies/liter of radon in equilibrium with its daughters for continuous exposure. For a 40 hour work week, on the basis of their auxiliary recommendations (Appendix 3, Handbook 52), this is changed to 3×10^{-11} curies/liter. In 1954 the International Commission on Radiological Protection recommended a level of 1×10^{-10} curies/liter for radon in equilibrium with its daughters for continuous exposure or 3×10^{-10} curies/liter for a 40 hour work week (6).

A study of the data and reasoning leading to these various recommended levels suggests that the assumptions on which they were based have a substantial degree of qualitative and numerical uncertainty. The results of published experimental investigations and theoretical studies indicate that the radiation dosage to portions of the respiratory tree may exceed 0.3 rem per week at the permissible exposure level given by the International Commission [Bale (7,8), Shapiro (9), Cohn et al (10), Chamberlain and Dyson (11)]. It has also been suggested by Morgan (12) that this level is too high. Furthermore this recommended level is within a factor of two or three of that believed by Evans to have existed in the Joachimsthal and Schneeberg mines, and which he believed to be responsible for the higher than normal incidence of lung cancer among the miners working in those mines (4). It may therefore be reasonable to conclude that further investigative work in this field is worthwhile.

One important question not yet definitely answered is whether or not radon exposure of human beings in the United States and Canada has so far led to lung cancer or other disability. A method for determining some quantity directly proportional to previous summated radon exposure in humans may make possible an answer to this question. For it would then be possible to look for a positive correlation between incidence of lung cancer or other disease and the relative magnitude of radon exposure in miners who have worked in similar environments.

Monitoring of individuals for degree of radon exposure is at present difficult to do on a routine basis. Radon levels vary from time to time and from place to place in mines. Also, many mines are small and only infrequently surveyed for radon and radon daughter levels. Therefore a reliable method for determining by a urine analysis some quantity directly proportional to a miner's summated radon and radon daughter exposure would be useful in this field.

It is possible that urinary polonium excretion of exposed individuals can furnish data for estimates of the relative integrated exposure to radon plus its daughters. This is evident from the following considerations. Radon decays through several relatively short-lived intermediaries to the long-lived RaD (Pb^{210}) as shown in Figure 1. Some of the RaD formed in the body by radon exposure is deposited in bone where it decays through RaE (Bi^{210}) to RaF (Po^{210}). Some of this polonium then appears in the general circulation from which a portion is excreted in the urine. Thus the polonium excretion of individuals exposed at the same times under the same general conditions may be proportionally related to their integrated exposure.

Sultzter and Hursh (13) have shown that active uranium miners do, in fact, excrete larger amounts of polonium than unexposed individuals, and that the amount excreted is probably at least roughly proportional to the magnitude of exposure.

Figure 1. Decay scheme - radon to stable lead

However, no data are yet available which give polonium excretion as a function of time after the exposure to radon took place. In general, one would expect daily polonium excretion to decrease with time due to at least three factors. First, the radiolead store will decrease by radioactive decay; second, the radiolead will be gradually excreted; and third, the radiolead may become buried increasingly deeper in bone which is relatively inert physiologically with the result that polonium formed from it would not reach the circulation and so be available for excretion. Therefore to evaluate the usefulness of urinary polonium measurements as a monitoring method and as a relative indication of integrated previous radon exposure, it will be necessary to determine the rate of decline of polonium excretion as a function of time after a known period of exposure.

In a research program designed to provide these data, it will be essential to have reliable, quantitative methods for determining the polonium content of urine and tissues at or close to the levels resulting from the exposure of individuals to normal atmospheric concentrations of radon. This will require development of methods for quantitatively isolating on one foil, suitable for alpha counting, the polonium from a liter or more of urine and from 50 grams or more of soft tissue and bone. Progress in the development of these methods and their use in preliminary determinations of the types of samples mentioned above are the subjects of this paper.

II. CHARACTERISTICS OF POLONIUM

Polonium²¹⁰ is a naturally occurring radioactive element discovered by the Curies during investigation of the radioactivity of pitchblende. It is a member of the radium series known as RaF. It is formed by beta decay of RaE (Bi²¹⁰) and decays by alpha emission to form a stable isotope of lead. The half-life of Po²¹⁰ has been measured as $138\frac{1}{2} \pm 0.1\%$ days (14). Polonium is also a member of other radioactive series each of which has two or three isotopes of polonium. Other isotopes of polonium have also been produced artificially practically all of which are alpha emitters (15).

The atomic number of 84 places polonium in group VI B of the periodic table so it would have properties like S, Se, and Te if it were a non-metal, or properties like Tl, Pb and Bi if metallic.

It has been shown that polonium is a metal existing in two allotropic forms with 42 and 44 micro-ohm-cm specific resistivity.

The crystalline structure is simple cubic for the α form and simple rhombohedral for the β form with the transition occurring between 65° and 130° C. It melts at 254° C, has a density of 9.4 gm/cc, and is similar to lead in hardness. Its thermal coefficient is large and negative, $-300 \pm 100 \times 10^{-6}$ per °C. The valence of polonium is variable, ranging from +4 to -3, and it shows a marked tendency to form complexes; pyrogallates, acetylacetonates, oxalates, camphorates,

and tartrates, among others, have been prepared. Electrochemically polonium behaves like a noble metal (16, 17, 18, 19).

It was discovered rather early that polonium could be readily vaporized from other materials, for example it vaporized at temperatures 250° C lower than RaE (Bi) and 300° C lower than RaD (Pb).

This effect has been used to purify polonium by vaporizing it in a stream of gas from which it is deposited in a cold portion of the system (20, 21, 22).

The electrochemical behavior of polonium has also been used to purify it. Marchi (23) extracted it from solution by bubbling hydrogen through the solution while rotating a Pt electrode in it. Prestwood (24) separated 94% of the polonium in a RaD, E and F solution by stirring with a platinum electrode in a hydrogen atmosphere. Polonium in acid solution will also deposit by electrochemical displacement on copper, nickel, silver and alloys containing nickel, such as monel metal and stainless steel, Marckwald (25) observed this "plating" on silver and bismuth in 1905.

In spite of the ease of removal of polonium from acid solutions, there are several ions which will interfere with the "plating" process. These include Ca^{++} , Ca^{+} , Ag^{+} , and Fe^{+++} as well as some components of the usual digestion mixtures such as the Van Slyke-Folch, Nicleaux and Kjeldahl reagents (31). In addition, polonium which has been metabolized in the body, may possibly form complexes

which would also impede its plating. For these reasons it is necessary to develop a method for complete digestion of biological samples and for treatment of the digest to minimize the above mentioned impedances before quantitative recoveries of polonium can be expected.

However, it may be possible to plate on metal foil or powder by electrochemical displacement most if not all of the polonium content of urine without prior digestion.

III. REVIEW OF PREVIOUSLY REPORTED PROCEDURES

The following methods have been used for digestion of biological samples preliminary to assay for radioactive materials. Digestion, as used here, refers to complete oxidation of organic material and is used to insure that any radionuclide present in the tissue is placed in an ionic or inorganic state. The plating procedures depend on the electrochemical displacement by the polonium of the less noble metal of the foils.

A. Procedure Used by Scott and Stannard (26, 31).

1. Place the sample in a vessel in which it will occupy not more than 10% of the volume, then add 4 to 5 times the sample volume of concentrated nitric acid, and let stand for several hours or preferably overnight.
2. Bring to a boil and continue heating with additions of nitric acid until the digest is a clear straw color. Evaporate to minimal volume, cool and add 25 ml of water per 100 grams or less

of sample. A layer of fat indicates the need for more digestion.

10 to 15 grams of rat liver require 16-18 hours of hot digestion and 200 ml of nitric acid. For larger amounts of tissue or digests with much fat, continue digestion as follows.

3. Add 25 ml of sulfuric acid and heat to charring with the appearance of white sulfur trioxide fumes. Add up to 1 liter of nitric acid and heat. Continue the alternate sulfuric and nitric acid additions until a fat-free straw colored solution results.

4. 30% hydrogen peroxide may be added with heating until the straw colored solution results. A cat carcass will require as much as 60 hours of active digestion and 30 pounds of nitric acid. Cat and rabbit carcasses, in general, are divided into several smaller portions and digested separately.

5. Evaporate the digest to a minimal volume, add 100 ml of water and neutralize the excess acid with sodium hydroxide.

6. Add hydrochloric acid so the solution will be 0.5N when diluted to a convenient volume.

7. An aliquot containing, if possible, not less than 20 cpm nor more than approximately 100 ml in volume is taken for plating.

8. Plating is done on a silver foil by stirring the 100 ml plating volume in a 95-100° C water bath for 90 minutes.

9. This method has been used for bone and soft tissues in sizes from individual mouse organs up to cat or rabbit carcasses, and

for blood samples and urine from laboratory animals. The recoveries are reported as per cent of a dummy made by injecting into 0.5 N HCl the same amount of polonium solution used for injecting the animals. From 0.5 N HCl, recoveries averaged from $89 \pm 4\%$ for 15 cpm up to $96 \pm 4\%$ for 2×10^5 cpm. These latter percentages were based on total recoveries obtained by replating the supernatant of each plating to a total of up to 9 platings for each solution.

B. Procedure Used by Fink (27).

1. For samples less than 50 grams.
 - a. Use a Kjeldahl flask such that the sample occupies not more than 5% of the flask volume and add 3 ml of 72% perchloric acid plus 1 ml of nitric acid per gram of sample.
 - b. Heat in water bath until foaming ceases, then transfer to an air bath (asbestos covered wire gauze over a gas flame).
 - c. Continue heating until dense white fumes evolve and a clear solution results. If solution darkens, cool and add nitric acid carefully, then reheat. Continue until clear solution results.
2. For 50 to 125 grams of sample.
 - a. Add 150 ml of nitric acid and allow to stand several hours.
 - b. Heat in water bath until foaming subsides, cool, and add 150 ml of 72% perchloric acid. Reheat in water bath until foaming again subsides.

c. Add 10 ml of nitric acid carefully and glass beads and transfer to air bath.

d. Continue digestion as above.

3. Plating procedure.

a. The digests are neutralized with sodium hydroxide, then made 0.5 N in HCl and an aliquot taken for plating.

b. The polonium is plated on a silver foil by stirring 80 ml of solution which contains the aliquot and which is 0.5 N in HCl, for 90 minutes in a 90-100° C water bath.

This method was used for samples as large as 125 grams in an 800 ml Kjeldahl, with larger animals (cats or rabbits) being divided into several portions. Recoveries averaged 92% of a dummy prepared as in Part A above. It was indicated that nickel filings could be used for determining the polonium content of a liter or more of urine.

C. Procedure used by Cember et al (28).

1. Add 2 ml of nitric acid to 2 grams of tissue and allow to stand overnight.

2. Heat in water bath until digest clears, use antifoam agents if necessary.

3. Cool and add 1 ml of 30% hydrogen peroxide and reheat until a clear yellow or brown solution results.

4. Cool and add 2 ml hydrochloric acid, then heat until nitric fumes cease to come off.

This procedure was used only for 2 gram samples of tissue for analysis of sulfur-32 content. It is reviewed here merely as a digestion method which might possibly be applied to analysis of tissues for polonium since complete digestion was obtained by the method.

D. Procedure Used by Mulhaney et al (29).

1. Add 2.5 ml of 72% perchloric acid, 1.25 ml of 30% hydrogen peroxide and 0.5 ml of water per gram of material and let stand 48 hours.
2. Place on a rack heated electrically to 200° C and heat to dense white fumes with a clear solution.
3. With fat present, the solution will darken. Cool and add 1-2 ml of 30% hydrogen peroxide. Continued heating of the darkened solution may cause it to become concentrated with attendant danger of an explosion.
4. Reheat after the H_2O_2 addition until a clear solution results and perchloric acid has been destroyed by the peroxide, using more peroxide if necessary.
5. Add sufficient hydrochloric acid to make the solution 1N in HCl.
6. Plate 2 1/2 to 3 hours from 50 ml of solution by stirring rapidly enough at a temperature of 90-100° C to keep a 1 inch silver foil spinning on edge.

This procedure was used for mouse organs, whole mice and their excreta, with recoveries reported as quantitative as compared to a dummy prepared as in Part A above.

E. Procedure Used by Rundo (30).

1. Wet ash the tissue with 10 ml nitric acid and 5 ml of dilute (1:1) sulfuric acid per gram and evaporate until charring occurs.

2. Take up in 50 ml nitric acid and evaporate again. If solution is not clear, add 40 ml of $\text{HNO}_3\text{-HClO}_4$ (3:1) mixture and evaporate. Continue this until a clear water white solution which gives off white sulfur trioxide fumes is obtained.

3. Transfer residue to a 1-liter beaker and add 10 ml of a solution containing 1 mg/ml of tellurium. Add 2 grams sodium hypophosphite and boil to complete precipitation.

4. Filter the precipitate and dissolve it off the paper with 5 ml HCl saturated with bromine and wash the paper well with 0.5 N HCl.

5. Warm the dissolved precipitate and washings in a 100 ml beaker and add sufficient hydrazine hydrochloride to precipitate the Te when boiled gently.

6. Filter off the tellurium and wash well with 0.5N HCl.

7. Neutralize the filtrate plus washings with sodium hydroxide.

8. Add 4.5 ml HCl and make up to 100 ml with distilled water.

9. Plate on a clean silver foil by stirring moderately for 90 minutes at 80° C.

The method was developed to detect 1 μc of polonium in 1 liter of urine but no data were reported, though the addition of perchloric acid was reported as not necessary for digestion of urine. Recoveries from mouse carcasses, mouse kidneys, and excreta averaged 85%.

F. Comment on These Procedures.

The above do not include all procedures but have been selected as indications of the variety of procedures that have been used.

In selecting a procedure for digestion of biological materials preliminary to analysis of polonium content, the major factors to be considered include:

1. Digestion should be complete enough to destroy any organic material which might form a complex with the polonium.
2. Digestion reagents should not add ions which interfere with polonium deposition.
3. As few vessels as possible should be used.
4. The final solution should not react with the material on which the polonium is to be deposited.

The Rundo procedure was discarded immediately since it is too complicated. The procedures requiring perchloric acid were by-passed because of the possibility of explosion with loss of perhaps irreplaceable samples. It was therefore decided that the procedure of Scott and Stannard would be used initially. None of the procedures listed had been tested experimentally for recovery of polonium at levels near 1 dpm above background and so could not be applied for the analyses of the type attempted herein without such testing.

IV. APPARATUS USED IN THIS WORK

A. Counting Equipment.

Counting of the foils resulting from the analytical procedures was accomplished through the use of an alpha scintillation counter. This consisted of an NICC model 165S scale of 64 binary scaler to which a G.E. scintillation counter head was attached. The tube used was a DuMont type 6292 photomultiplier tube which was optically coupled to a plastic dish by use of a high viscosity silicone liquid.* The face of the plastic dish nearest the foil to be counted was coated with ZnS to convert the alpha ionizations to light which could be detected by the photomultiplier tube. The ZnS screen was protected by a thin plastic covering so decontamination could be accomplished by scrubbing the screen with a scouring powder. The sample table height was adjustable so the sample could be brought close to the ZnS screen for detection of the maximum possible number of particles.

To calibrate the counter set-up, a comparison was made with a Nuclear Measurements Corp. gas-flow alpha counter. The efficiency of the gas-flow counter was checked by counting the alpha particles from an aged RaD + E standard which had been calibrated by the National Bureau of Standards. The alphas come from RaF (Po^{210}) which is essentially in equilibrium with RaD + E after 3 years. Then the counting rate of a silver foil, on which polonium had been electroplated, was

*Dow Corning 200 Fluid , viscosity 10^6 cs, Dow Corning Corp., Midland, Mich.

determined. Finally the counting rate of the same polonium foil was determined in the scintillation counter, thus the efficiency was readily obtained, e.g.

1. RaD+E std #200 had an activity of 118.4 dps on 4-25-46 according to the accompanying NBS certificate

$$\text{Decay 9 yrs: } 118.4 \times 2^{-9/22} = 89.3 \text{ dps or } 5350 \text{ dpm on } 5-10-55.$$

2. In gas-flow counter 5254 counts/2 min. = 2627 cpm for RaD+E std

$$2627/5350 = 49.15\% \text{ efficiency of gas-flow counter}$$

3. Sample #5a 4904 counts/242 min. = 20.26 cpm in gas-flow counter

$$\frac{20.26}{0.4915} = 41.2 \text{ dpm on } 5-10-55$$

4. Sample #5a in scintillation counter

$$23,519 \text{ counts}/1248 \text{ min} = 18.84 \text{ cpm.}$$

This was counted previously on 4-21-55

$$1154 \text{ counts}/56 \text{ min.} = 20.6 \text{ cpm}$$

$$19 \text{ days decay: } 20.6 \times 2^{-19/138} = 18.8 \text{ cpm} \quad \text{check}$$

5. Scintillation counter efficiency = $\frac{18.84 \text{ cpm}}{41.2 \text{ dpm}} \times 100 = 45.7\%$

Once the efficiency of the scintillation counter was established, a silver foil containing an electrodeposit of uranium was counted. This provided a substandard of almost constant activity which could be used to check the performance of the counter throughout the course of this work. Uranium alphas are less energetic than polonium alphas so a decrease in counter sensitivity would result in lower efficiency.

as determined by the uranium standard than if determined by a polonium standard, however the difference would not be very great.

After a careful cleaning of all parts of the scintillation head exposed to the ZnS screen the background for this equipment as measured over a cleaned silver foil varied between 3 and 6 counts per hour, depending on the foil used.

B. Plating Apparatus.

The equipment for plating consisted of a water bath large enough to hold six plating bottles. It was heated by a 1000-watt electrical heating coil which kept the water bath at 94° C after a 30 minute warmup period. Over the water bath was a frame in which 6 stirring spindles were mounted. The stirrers were driven simultaneously by a motor, the power being transmitted by a nylon endless belt. The stirrers rotated at approximately 720 rpm. See Figure 2.

The plating bottles used were Davol brand nursing bottles modified for this purpose according to the usual procedure at this institution, as shown in Figure 3.

Figure 2. Plating equipment -- the individual stirrers are mounted in a frame constructed of $1/4"$ x $2"$ steel, the water bath is a box made of $1/2"$ plywood lined with 20 gauge copper, a $1/4"$ transite board in the bottom supports the 1000-watt electrical heater on which rests a 20 gauge copper plate with supports for positioning the plating bottles. The stirrer collets are mounted on shafts through roller bearings. The stirrers are belt driven by a variable-speed electric motor.

19

Figure 3. Plating bottle and stirring rod.

Davol brand nursing bottles with the bottom cut off. A 1/16" neoprene disk and .005" silver foil fit inside the cap which is then firmly screwed onto the bottle. The stirring rod is 1/4" glass with the bottom heated and squeezed with tongs to make two 1/2" oval flats at right angles to each other and bent to a line approximately 30° from centerline of rod.

V. PRELIMINARY EXPERIMENTS ON RECOVERY OF POLONIUM

Before proceeding to analytical work, it was necessary to determine the conditions which would give the best possible recoveries using the equipment available. For this purpose the following experiments were performed.

A. Per cent recovery as a function of activity

This experiment was necessary since it had been reported that percentage recovery decreased with decrease in activity (31).

Six plating bottles were selected and decontaminated (See Appendix). They were then assembled, with a cleaned silver foil inserted, filled with 0.5 N HCl, and stirred 2 hours in the water bath. Counts on the foils showed an average activity of 0.1 cpm.

The bottles were then partially filled with 0.5 N HCl, after insertion of new, cleaned silver foils, and stock polonium solution was added to give the activities listed in Table I. The stock polonium had been diluted with 0.5 N HCl to give approximately 1 cpm/ml.

Sufficient 0.5N HCl was added to the bottles to make the final volume 200 ml. The solutions were then stirred 2 hours at 94° C in the water bath. The results are shown in Table I.

The experimental results indicate no obvious correlation between activity and per cent recovery. They also indicate an average recovery of 93.1% for activities in the range of 2 to 160 dpm.

TABLE I

Recovery of polonium from 0.5N HCl as a function of radioactivity level

Sample	Added Po Activity	Total Count	Time min.	Cpm	Net Cpm	% Recovery
1	1.04	1053	985	1.07	0.97	93.4
2	3.12	652	237	2.75	2.65	85.1
3	7.28	662	94	7.05	6.95	95.5
4	15.61	662	45	14.71	14.61	93.7
5	32.2	747	25	29.9	29.8	92.6
6	79.3	779	10	77.9	77.8	98.2

B. Per cent Recovery as a Function of Plating Time.

Four plating bottles were prepared as in Part A above. To each were added approximately 20 cpm of polonium. Two of the solutions were stirred for 2 hours and two for 4 hours. The results are set forth in Table II.

TABLE II

Recovery of polonium from 0.5N HCl as a function of plating time

Sample	Plating Time hours	Total Count	Time min.	Cpm
1	2	1154	56	20.6
2	2	2350	114	20.6
3	4	3151	150	21.0
4	4	1479	72	20.6
Average				20.7

This indicates that no advantage ensues by using the longer stirring time under the conditions of this experiment.

C. Absolute Recovery of Polonium.

To furnish a standard for the activity of the stock polonium solution against which actual recoveries could be checked, the following experiment was performed.

On each of 3 silver foils was deposited, by the same pipette used in Part B, the same amount of solution as used in Part B. The foils were then dried gently under an infra-red lamp so that no spattering occurred. The foils were counted when dry and the results were as set forth in Table III.

TABLE III

The same amount of polonium as in Table II deposited on a silver foil and air-dried gently under an infra-red lamp

Foil	Total Count	Time min.	Cpm
1	1225	54	22.7
2	1532	78	19.7
3	4544	229	19.9
		Average	20.7

These results indicate that the recovery of polonium from 200 ml of 0.5N HCl, when stirred rapidly for 2 hrs at 94° C over a silver foil can approach 100%.

D. Effect of HNO_3 on Recovery of Polonium

During the final plating of a digest, assuming no interfering ions are present, one of the most important factors affecting recovery is attack on the foil by components of the plating solution. Nitric acid has been held chiefly responsible for this effect (31).

1. To determine the relation between recovery and normality of HNO_3 content, the following experiment was designed. Solutions were made up to a total volume of 200 ml which were 0.5N in HCl, had approximately 85 cpm of polonium added and which had various nitric acid content. They were plated on silver foils by stirring rapidly for 2 hours at 94°C , air dried, and counted with the results as set forth in Table IV.

TABLE IV

The effect of nitric acid on the recovery of polonium from 0.5N HCl

Sample	Added Activity	Normality of HNO_3	Total Count	Time Min.	Net Cpm	% Recovery
1	85.2 cpm	0.04	745	8	93.0	109
2	" "	0.08	675	8	84.4	99
3	" "	0.16	645	8	80.5	94.5
4	" "	0.24	692	9	76.8	90.1
5	" "	0.36	944	13	72.5	85.1

These results show a definite decrease in recovery with increasing nitric acid content. Thus indicating that for best results, the

plating solution should be less than 0.16N in HNO_3 (2 ml concentrated HNO_3 in 200 ml of solution). This effect is probably due to the formation of dilute aqua regia in the plating solution which reacts readily with the silver foil.

The discoloration ranged from just perceptible on Foil #1 to moderately heavy on Foil #5.

2. The manner in which the nitric acid caused low recoveries was studied experimentally as follows.

The low recovery could be due to the discolored deposit actually overlying the polonium and thus acting as an absorber or it could be due to the deposit occluding the surface of the foil so that no silver was exposed to the solution. If the deposit was a silver salt, it could be dissolved in "hypo." Both a solution of sodium thiosulfate in water and a commercial acid fix were used.

Also the plating solutions which produced such a deposit were saved, boiled down, and as much as possible of the nitric acid boiled off, by repeated additions of concentrated HCl with heating. The results of these experiments are set forth in Table V.

Since the activity of a foil decreased when the deposit was removed, the deposit probably was not overlying the polonium.

Since more careful elimination of HNO_3 led to recovery of a large portion of the originally added activity, it is highly probable that the deposit mechanically interferes with the plating

of polonium.

Because of these results, it was decided to save all solutions until it could be determined whether or not the foils were discolored.

TABLE V

The effect of various treatments to improve recovery of polonium when the silver foil has been discolored during plating

Sample	Treatment	Net Cpm		% Change
		Before	After	
1	Foil soaked in neutral $\text{Na}_2\text{S}_2\text{O}_3$	7.6	6.6	- 13
2	Foil soaked in acid fix	19.33	13.9	- 28
3	Solutions replated on new foil after boiling down	12.41	32.8	+ 264
4	with added conc. HCl to drive off more nitric acid	5.63	31.8	+ 565

E. Effect of Undissolved Solids on Recovery of Polonium

In the course of routine digestions of various materials, it is sometimes impossible to get all the solids into solution in the 200 to 220 ml volume of 0.5 N HCl which the plating bottles will hold. A digest of 40 to 50 grams of bone will contain such undissolved solids.

To check the effect of these solids the following experiments were performed. Three dog femurs, weighing approximately 40 g

each were digested by the method described on page 58, two having added polonium, and plated according to the usual procedure.

Six samples of beef pelvic bone, weighing between 40 and 50 grams, were similarly treated, four had polonium added. Three of the latter samples were filtered through Whatman #41 filter paper (long stem funnel) and three filtered through a sintered glass disk of coarse porosity. The filtrates were plated. Some samples containing only the digest reagents with added polonium were run in parallel with the bone samples. The results of these experiments are shown in Table VI.

These results show that recovery is poor when undissolved solids are present in the plating solution and that filtering the solids does not increase the recovery appreciably. They also indicate that the filtering procedure did not remove any activity, (cf. samples 9, 10, 15, and 16 which were also filtered). Two possibilities for the low recoveries are that the solids might shield the foil or they may complex with the polonium. The data on filtering apparently supports the latter possibility. As a result of these data, it was decided that all solids should be dissolved before plating the polonium either by increasing the normality of the solution or by increasing the volume by dividing the digest between two or more plating bottles. The latter procedure, of course, would not be desirable for samples of low activity.

TABLE VI

Effect of undissolved solids on the recovery of polonium

Sample	Composition	Treatment	Net Cpm	% Recovery
1	Dog femur	Plated with undissolved solids	10.96	23
2	" "	" " " "	0	0
3	Digest reagents	containing added polonium	48.5	103
4	" "	" " "	50.1	106
5	Dog femur	No added activity	0.16	--
6	Digest reagents	No added activity	0	--
7	" "	" " "	0.03	--
8	Beef pelvis	" " "	0.294	--
9	Digest reagents	Added activity and filtered	37.8	98.8
10	" "	" " "	38.0	99.4
11	Beef pelvis	Undissolved solids filtered	10.2	26.0
12	" "	with Whatman #41 paper	16.2	42.0
13	Digest reagents	No added activity	0.02	--
14	Beef pelvis	No added activity	0.201	--
15	Digest reagents	Added activity and filtered	37.8	105
16	" "	" " "	37.6	104
17	Beef pelvis	Undissolved solids filtered	2.24	6.2
18	" "	with sintered glass disk, of coarse porosity	2.33	6.4

F. Effect of Ferric Ion on Polonium Recovery.

The effect of ferric ion on the recovery of polonium has been well established (31) so no experiments were performed on this effect. However, 100 mg or more of ascorbic acid were added routinely to all solutions just prior to plating, the amount depending on the amount of tissue originally digested, e.g. 100 mg for 50 grams or less of bone or soft tissue and 200 mg for any larger amount of bone or soft tissue up to and including a whole rat.

VI. RECOVERY OF SMALL AMOUNTS OF POLONIUM FROM URINE

A. Introduction and Survey of Previously Reported Procedures.

A satisfactory method for determining small amounts of polonium in urine should be relatively easy to use, give closely reproducible results, and be useful for large volumes. It would be advantageous to handle large volumes in order to increase the amount of polonium recovered, and thus reduce counting errors.

Several methods are used routinely for low-level polonium analysis of urine. Three of these methods are the following.

1. Nitric-perchloric acid digestion and concentration followed by plating on silver at 90-100° C, as described by Sultzer and Hursh (13) is used by the Public Health Service in Utah.

2. At Mound Laboratory approximately 100 ml of urine is acidified in a waxed paper container, a nickel foil is inserted and the solution stirred at 50° C. Deposition is on both sides of the foil. The report includes data on the effect of storage, efficiency of various metal foils, plating temperature, plating time and volume (32).

3. At the University of Rochester Atomic Energy Project, 100 ml of urine is acidified with 10 ml HCl in a waxed paper cup and stirred 2 hours at room temperature with both sides of a clean copper foil exposed to the solution (33).

The first method appears useful and was used by Sultz and Hursh in the analysis of miners' urine as well as by the author and Dr. Jacob Shapiro of this institution in the analysis of 37 additional samples of miners' urine with a maximum volume of about 250 ml of urine. The latter two methods are interesting but can handle only small volumes of urine (100 to 150 ml), with the activity deposited on both sides of the foil.

The following experiments were designed to determine, if possible, the best method for analysis of urine for polonium content at levels close to those occurring naturally.

Pooled urine of Project personnel was used in the following experiments.

B. Plating of Polonium on Silver, Nickel, and Copper

Silver, copper, and nickel foils were used. The foils were inserted in the plating bottles and 200 ml of acidified urine (0.5N in HCl) were placed in each flask. Approximately 30 cpm of polonium were added to each bottle along with 100 mg of ascorbic acid. The bottles were placed in the water bath at 94° C and stirred rapidly for 2 hours. The results are shown in Table VII.

TABLE VII

Plating of added polonium from acidified urine onto various metals

Sample	Foil	Total Count	Time min.	Cpm	Net Cpm	% Recovery	Mean $\pm$ Std. Deviation
1	Silver	1691	59	28.7	28.6	92.2	
2	"	1009	34	29.7	29.6	95.4	93.6 $\pm$ 1.9
3	"	4760	164	29.0	28.9	93.2	
4	Copper	1034	49	21.1	21.0	67.7	69.6 $\pm$ 8.4
5	"	2057	85	24.2	24.1	77.7	
6	"	755	38	19.8	19.7	63.5	
7	Nickel	1074	41	26.2	26.1	96.6	94.7 $\pm$ 3.3
8	"	2146	87	24.7	24.6	91.0	
9	"	1103	42	26.2	26.1	96.6	

The recoveries from copper averaged 23% lower than from nickel or silver and had greater variation. Perhaps this was due to the discoloration of the copper foils, an effect which did not occur with the other foils.

More samples were not run since these results confirm more extensive data already available; the few samples were run merely as a check.

C. Recoveries from Urine with Nickel Filings and Nickel Powder

These experiments were designed to develop a method for using nickel powder to recover polonium from large volumes of urine. It has been demonstrated that larger volumes require longer plating times for a given area of foil exposed to the solution (27). On the other hand, the plating time can be shortened by increasing the surface area of the foil, which can be done most readily by using a powder.

These experiments were performed with digested urine, undigested urine and 0.5N HCl as the polonium containing solutions.

The urine was digested in the following manner.

1. Approximately 30 cpm of polonium were added to 200 ml of acidified urine in a 250 ml Ehrlenmeyer.
2. 25 ml HNO_3 were added and solution boiled down to 10 ml on a hotplate.
3. 20 ml of a 1:2 mixture of nitric and 72% perchloric acids were added and boiled down until the white fumes from perchloric acid appeared.
4. A few drops of HCl were added down the side of the flask. If only a transient color change occurred in the solution and no

brown fumes appeared, the digestion was complete and all the nitric destroyed.

5. The solution was made alkaline, since the acidity was unknown, using a saturated NaOH solution with bromothymol blue indicator.

6. Reacidified with HCl to approximately 0.5N.

Three of these digests were plated on silver foil directly. To three others were added 1 gram of nickel filings (through 20 mesh onto a 160 mesh screen).

To two flasks containing 200 ml of 0.5N HCl were added a known amount of polonium and 1 gram of nickel filings.

To four flasks containing 200 ml of acidified urine were added a known amount of polonium and 1 gram of nickel filings.

Treatment proceeded as follows.

- (1) All solutions containing nickel filings were placed in a 94° C water bath and stirred rapidly for 2 hrs.
2. The flasks were removed from the water bath and the solution decanted.
3. The filings were rinsed in 25 ml of distilled water which was also decanted.
4. The filings were then dissolved in concentrated HCl by heating on a hotplate and boiling down to a few milliliters. This required 6 to 8 hours.
5. Sufficient distilled water was added to each solution, assuming it to be constant boiling HCl, to make it 0.5 N in HCl.
6. The solutions were brought to boiling and

transferred to prepared plating bottles with 3 rinses of 0.25N HCl sufficient to make the final volume about 200 ml. 7. 100 mg ascorbic acid were added and stirred rapidly for 2 hours in a 94° C water bath.

The results of these experiments are shown in Table VIII.

TABLE VIII

Recovery of added polonium from acidified urine and 0.5N HCl by various treatments. Volumes of approximately 200 ml.

Sample	Composition	Treatment	Total Count	Time Min.	Net CPM	% Recovery
1	Digested urine	Plated on silver	5899	207	28.4	96.5
2	"	"	26,301	977	26.8	91.2
3	"	"	1444	50	28.8	98.2
4	"	Plated on nickel	2403	95	25.2	85.8
5	"	filings, then on	1834	77	23.8	80.7
6	"	silver foil	1872	77	24.3	82.5
7	Acidified urine	"	1946	70	27.7	95.2
8	"	"	1581	58	27.2	93.4
9	"	"	1434	54	26.5	91.2
10	"	"	3189	120	26.5	91.2
11	0.5N HCl	"	2286	88	25.8	90.0
12	"	"	3420	121	28.1	98.0

These results show that recovery of added inorganic polonium is higher from digested urine when plated directly on silver than when nickel filings are used as an intermediary. They also show good recovery from undigested urine when the activity is first plated on nickel filings, the filings dissolved, and the activity replated on silver.

On the basis of these results, it was decided to attempt recoveries from large volumes of urine.

D. Recoveries from Large Volumes of Urine Using Nickel Powder

1. The first experiment in this series was performed to determine the recovery with nickel filings as used in Part B above and with nickel powder* which exposed a much greater area of metal to the solution.

a. 1-liter of acidified urine was placed in each of four 1-liter beakers.

b. A known amount of polonium was added to each beaker.

c. To two of the solutions were added 2 grams of nickel filings, and to the other two were added 2 grams of nickel powder.

d. Placed in a 94° C water bath and stirred rapidly for 2 hours.

e. Urine decanted and 50 ml HCl added to dissolve the nickel. The addition of 10 ml of nitric acid apparently speeded up the solution process which then required 5 to 6 hours.

f. Boiled down to approximately 10 ml of solution and 50 ml distilled water added.

g. Transferred to prepared plating bottles with 3 rinses of 0.25 N HCl to make final volume about 200 ml.

* Nickel Metal; powder; Baker and Adamson Chemicals, Gen. Chem. Div., Allied Chem. and Dye Corp, New York, N. Y.

h. 100 mg ascorbic acid added, placed in water bath, and stirred rapidly for 2 hours at 94° C.

The results are shown in Table IX.

TABLE IX

Recovery of added polonium from acidified urine.
Comparison of nickel powder and nickel filings.

Experiment	Sample	Composition	Nickel	Total Count	Time Min.	Net Cpm	% Recovery
1	1	1-liter urine	powder	2050	119	17.1	59.9
	2	" "	"	2636	130	20.1	70.5
	3	" "	filings	1328	46	28.7	100.
	4	" "	"	1105	46	23.9	83.7*
2 (powder filtered)	1	225-ml urine	powder	2847	101	28.1	100.2
	2	" "	"	1215	43	28.2	100.4
	3	" "	filings	3206	117	27.3	97.5
	4	" "	"	1311	45	29.0	103.7
	5	225 ml 0.5N HCl	powder	1025	39	26.2	95.2
	6	" "	"	2920	113	25.7	93.7

*Foil discolored

2. Since the results with nickel powder were poor, it was thought that possibly some of the fine powder was lost during decantation. To check on this possibility smaller volumes were used and filtration attempted instead of decantation.

The procedure was the same as in 1. above except that 225 ml. of urine or 0.5N HCl was used for each sample, and 1 gram of nickel filings or nickel powder was used in each sample.

After plating the added polonium on the nickel, the metal was filtered off by using Reeve Angel #201 filter paper under suction and was washed once with 25 ml of distilled water. Some powder did collect on the paper so it was placed in the flask containing the remaining powder.

50 ml of aqua regia (3 volumes HCl plus 1 volume HNO_3) were added to the flask by pouring it through the Büchner funnel to rinse out any remaining powder. The flasks were heated to boiling and the nickel and paper went into solution in 2 to 3 hours. The boiling was continued until no brown fumes were given off when a milliliter of HCl was added. The results of this experiment are also shown in Table IX. These results indicate that in analysis of urine for added polonium, the recovery will be quantitative when the procedure involving initial plating of the polonium on nickel powder, followed by filtration, solution of the powder in aqua regia and replating on silver, is used.

The appearance of a discolored foil indicates that all the nitric acid is not destroyed. For critical analysis, perchloric acid should be used to eliminate any remaining nitric acid, followed by neutralization of the solution and reacidification with HCl.

The method as developed is described in section G below and was applied to the analysis of urine from non-exposed personnel as shown in section H.

E. Recoveries from Large Quantities of Urine by Concentration

These experiments were undertaken to test another method for recovery of polonium from large volumes of fluid.

1. Twenty-four hour urines were collected from six donors in 2-quart Mason jars.

a. Sufficient nitric acid was added to make the solutions 10% HNO_3 by volume.

b. A known amount of polonium was added to each jar.

c. A portion of the urine was poured into a 400 ml beaker and boiled off, additions being made to the beakers from the collecting jars as the level became low.

d. When the urine was concentrated to about 10 ml, 10 ml of HCl were added and boiled off to drive off any remaining nitric acid.

e. The solutions were neutralized with saturated NaOH solution to the bromo thymol blue end-point and reacidified to 0.5N with HCl .

f. Transferred hot to plating bottles with 3 rinses of 0.5N HCl to make 200 ml final volume.

g. 100 mg of ascorbic acid were added and solutions stirred rapidly in a 94°C water bath for 2 hours.

After plating, the solutions were poured back into their respective beakers. When the recoveries were found to be poor, steps f. and g. above were repeated using new foils. The results of this experiment are shown in Table X.

TABLE X

Recovery of added polonium from acidified urine by evaporation of large volumes

Sample	Urine vol. ml.	Net Cpm	% Recovery
1	1630	42.4	79.1
2	1240	40.5	75.0
3	945	46.2	85.3
4	900	29.5	54.4
5	1320	38.8	71.9
6	1450	38.1	70.6
Average			72.8 ± 18.2

The recovery is not as good as with the method using the nickel powder. It may be that the final solution was either too concentrated, since two platings on two separate foils had to be made from each sample, or that some of the activity remained on the sides of the beaker. This method would not be suitable for low activities in either case.

2. To speed up the boiling off process, some 2-liter bottles having standard ground-glass tapered necks were obtained. Standard 34/45 ground-glass connectors were modified by cutting off some of the free end and affixing a short length (3") of 9 mm inside diameter glass tubing (See Figure 4).

Three of the bottles were filled with 1N HCl, a known amount of polonium added, and the modified connectors inserted. They were then inverted over 250 ml beakers and the height adjusted so about one inch of liquid ran into the beakers. The beakers were on asbestos covered wire gauze and were heated by gas flame. As the solution boiled away, more dropped into the beaker from the 2-liter bottle so the level was maintained until all the solution boiled away. More than half was boiled off in 7 hours.

The last 50 ml were transferred to a plating bottle with 3 rinses of 0.5N HCl sufficient to make the final volume 200 ml. They were stirred rapidly for 2 hours in a 94° C water bath. The average recovery was 93.1%.

F. Recovery of Polonium from Urine of Polonium Injected Rats

1. One important point not covered by the above experiments was the necessity of digesting the urine. It is possible that polonium which has been metabolized and excreted by the kidneys may be complexed with some constituents of urine. Thus recoveries from

Figure 4. Apparatus for boiling off large volumes of fluid.

undigested urine, to which polonium has been added, may be good but not quantitative for determining polonium which has been excreted.

In order to decide this issue, a quantity of urine was secured from rats which had been injected intravenously with polonium solution several weeks previously in another experiment. An amount of concentrated HCl equal to 0.1 times the volume of urine was added to prevent deposition of the polonium on the flask.

2. The experiment was performed in two parts. In one part the urine was plated as is and in the other the urine was digested then plated.

a. Five ml of rat urine were added to 125 ml of 0.5N HCl in each of five prepared plating flasks and the solutions stirred rapidly for 2 hours in a 94° C water bath. The solutions were discarded and the foils counted when dry.

b. Five ml of rat urine were added to 50 ml of 0.5N HCl in each of five 250 ml beakers and digested with nitric and perchloric acids (procedure listed in Part A.1. above). After neutralizing with NaOH pellets and reacidifying to 0.5N in HCl, the solutions were transferred to prepared plating bottles with 3 rinses of 0.5N HCl sufficient to make 125 ml final volume. They were then placed in a 94° C water bath and stirred 2 hours. The results of this experiment are shown in Table XI.

TABLE XI

Effect of wet digestion on the recovery of polonium from the urine of polonium injected rats

Sample	Treatment	Total Count	Time Min.	Net Cpm	Mean Recovery $\pm$ Std. Deviation
1	Undigested	1681	216	7.69	
2	"	7864	1041	7.45	
3	"	1193	163	7.23	7.35 $\pm$ 0.47 cpm
4	"	28,996	3819	7.49	
5	"	377	54	6.89	
6	Wet digested	1287	138	9.22	
7	"	7976	1024	7.66	
8	"	1123	118	9.44	8.89 $\pm$ 1.05 cpm
9	"	1502	160	9.29	
10	"	1092	122	8.86	

The difference in recovery by the two treatments is statistically significant with $P = 0.02$.

This shows that only 83% as much activity was recovered from undigested urine as compared to digested urine.

This indicates that procedures where the urine is merely acidified and stirred with a copper foil (recovery about 70% of added polonium) may only pick up about $70 \times 0.83 = 58\%$ of the polonium present.

G. Method Developed for Recovery of Polonium from Large Volumes of Urine.

From the results of the above experiments, the following method appears to be suitable for the recovery of added inorganic polonium from liter quantities of urine.

1. Acidify the urine by adding 50 ml of concentrated HCl per liter.
2. Add nickel powder in the amount of 2 grams per liter.
3. Add 100 mg of ascorbic acid and stir rapidly for 2 hours at a temperature of 90 - 100° C.
4. Filter the powder through a medium speed paper under suction.
5. Place the filter paper back in the original container and pour aqua regia, 30 ml per 2 grams of powder, through the funnel over the paper and powder and wash once with 25 ml of 0.5N HCl.
6. Boil the solution 2 to 3 hours to dissolve the paper and powder, then add 50 ml of a 2:1 mixture, V/V, of 70% perchloric and concentrated nitric acids.
7. Boil down until dense white fumes appear, then add 1 ml of concentrated HCl cautiously. If no brown fumes evolve then all the nitric has been destroyed.
8. Make alkaline with saturated NaOH solution using phenolphthalein as an indicator then add 5% of the volume of

concentrated HCl.

9. Cover the container and reflux gently for 2 hours to remove any polonium which may have deposited on the glass during the alkaline stage.

10. Transfer to a plating bottle fitted with a cleaned silver foil using three rinses of 0.5N HCl so total volume does not exceed 200 ml.

11. Add 100 mg of ascorbic acid.

12. Place in water bath at 90 - 100° C and stir rapidly for 2 hours.

With constant attention, 4 samples can be run in 9 to 10 hours.

If the urine is collected in a suitable container, e.g. a two-quart Mason jar, then the whole procedure up to the transfer to the plating bottle may be carried out in that container.

Urine containing excreted polonium was not available so this method could not be tested against other methods for recovery of metabolized polonium. It is possible that recovery of metabolized polonium from rat urine was improved by wet digesting prior to plating because such urine contains some food and fecal material, and that human urine which can be collected free of foreign matter may not require digestion. Experimental information of the latter point can be accomplished when urine from uranium

miners, or perhaps dogs exposed to uranium mine atmosphere becomes available.

H. Polonium in the Urine of Non-exposed Personnel.

1. Method. The method used was that using nickel powder for the initial removal of activity from the urine described in Part G above.

2. The detailed procedure was as follows:

a. 4 1-liter beakers were decontaminated by the method set forth in Appendix 1.

b. 1-liter of distilled water, 50 ml of HCl and 2 grams nickel powder placed in each of 3 beakers. Beakers placed in 94° C water bath and stirred rapidly for 2 hours with 200 mg ascorbic acid added at the start.

c. Filtered through #201 Reeve Angel paper under suction.

d. Filter paper placed back in the beaker and 30 ml of aqua regia poured over the nickel powder and paper through the Büchner funnel, rinsing down the sides, followed by rinsing with 25 ml of 0.5N HCl.

e. Warmed on hotplates until paper and nickel powder were dissolved.

f. 50 ml of a 2:1 perchloric-nitric mixture added and boiled down to the white fuming stage.

g. Made alkaline with saturated NaOH solution using

phenolphthalein indicator.

h. Reacidified with concentrated HCl to approximately half normal.

i. Transferred to plating bottles, over previously counted foils, with 3 rinses of 0.25N HCl sufficient to give 200 ml final volume.

j. 100 mg ascorbic acid added and stirred rapidly in a 94°C water bath for 2 hours.

The three foils when counted again gave blank values for this experiment.

The four 1-liter beakers were then run with hot tap water through the above procedure.

Finally, liter volumes of acidified urine were placed in the beakers and run through the above procedure. This urine was collected by placing 2-liter flasks in the mens' lavatory. When two liters of urine accumulated, 100 ml of concentrated HCl were added. Three samples were run for normal polonium content and one was run with added polonium. They were plated on the same foils as the tap water samples. This was done since available data indicated that no activity was added by the reagents (See Table XXIII) and that an initial run would remove 90% or more of any activity on the glassware, leaving negligible activity to be picked up by a second run. See Table XII for the results of this experiment.

TABLE XII
Recovery of polonium from liter volumes of tap and distilled water,
and urine from unexposed personnel

Sample	Composition	Total Count	Time Min.	Cpm	Foil cpm	Net Cpm	Remarks
1	Distilled water	493	3271	0.150	0.073	0.077	Average
2	"log activity"	271	2330	0.115	0.065	0.055	0.063 $\pm$ 0.013
3	""	188	1397	0.134	0.076	0.058	above foil counts
4	Tap water	407	3816	0.106			
5	" "	62	652	0.095			
6	" "	145	1053	0.138			
7	" "	138	1465	0.094			
8	Spiked urine	666	30	22.2	0.20	22.0	90% recovery
9	Urine	207	957	0.216	0.158	0.058	
10	" "	307	1394	0.220	0.201	0.019	Average
11	" "	1013	4379	0.231	0.157	0.074	0.05 $\pm$ 0.03

3. Discussion of These Results.

Some activity was added to the foils when tap or distilled water was run through the urine procedure. Since the reagents contain no measurable activity in the quantities used, (see page 94 and Table XXIII) it was probably due to the nickel powder used.

From the data presented, the activity averaged 0.063 μc per two grams of nickel powder though this was not explored further to definitely fix the value. However, this activity was subtracted from the urine cpm to determine the net cpm listed.

There was essentially no difference in the polonium content of the liter volumes of tap and distilled water runs detectable by the method used. This indicates that negligible polonium activity is present in one liter samples of tap water. Hursh (34) has found an average of 4×10^{-14} grams Ra per liter in public water supplies, water in equilibrium with the normal atmospheric radon of 10^{-13} curies/liter may contain 2.5×10^{-4} curies/liter of radon, and leaching from the soil might add some polonium to water so some difference in polonium content between tap and distilled water would not be unexpected.

The average value of the polonium content of urine shown in Table XII is not firmly established due to the relatively great range of the data. Many more samples, or samples of larger volume, would have to be run before the average polonium content could be fixed with any statistical validity. The variability among the samples shown in Table XII might be expected among samples from various individuals, but pooled urine as used here would be expected to yield more uniform values among various samples.

4. Calculations on Expected Urinary Polonium Excretion

There are two major sources which may contribute to the polonium activity of the urine. They are respectively the body store of radiolead which decays to polonium and the polonium obtained from ingested food and water and from inhaled air. Inhalation of polonium at most would probably contribute only a small fraction of the total polonium excretion. The body radiolead store is built up by decay of radon daughters filtered from the air by the lungs, by decay of the radon present in body tissues as a result of equilibration with atmospheric radon, and by decay of radon formed by the body radium store, with a certain portion of the radiolead so formed being stored chiefly in the bones. A fraction of the polonium derived from this stored radiolead will be excreted in the urine. Ingestion of food and water would contribute to body polonium in a manner similar to oral administration, with the resultant urinary excretion of a portion of the polonium so supplied. The following calculations are based on the foregoing considerations.

a. Stored radiolead from radon daughters filtered from the air by the lungs.

Assume: (1) an atmospheric radon concentration of

10^{-13} curies/liter (35)

(2) daughters in equilibrium with the radon

(3) lung filters out and retains 25% of the

inhaled daughters (9)

(4) Average minute volume of 14 liters over each 24 hours (NBS Handbook 52)

(5) 30% of the lead is absorbed and stored in the body (36)

(6) the effective half-life of radiolead is 880 days (13).

10^{-13} curies/liter $\times 2.22 \times 10^{12}$ dpm/curie $\times 14$ liters/min = 3.11 dpm/min of each daughter

$3.11 \text{ dpm} \times 0.25 = 0.778 \text{ dpm/min}$ for each retained daughter

Using the respective half-lives of the daughters and the following equation:

$$N = \frac{1}{\lambda} \frac{dN}{dt} \text{ where } \frac{dN}{dt} = 0.778,$$

the atoms per minute deposited in the lungs is found to be 55.6

$$55.6 \text{ atoms/min} \times 0.3 = 16.7 \text{ atoms/min stored in the body.}$$

Equating the atoms/min stored with the atoms per minute eliminated, through use of the effective decay constant which is

$$0.695/880 \text{ days} = 7.88 \times 10^{-4}/\text{day} = 5.47 \times 10^{-7}$$

per minute gives:

$$16.7 \text{ atoms/min} = N \times 5.47 \times 10^{-7} / \text{min}$$

from which $N = 3.04 \times 10^7 = \text{atoms of radiolead}$
stored at equilibrium.

b. Stored radiolead by equilibration of body tissues with
atmospheric radon

Assume: (1) 70 kg man with specific gravity of 1, i.e.

70 liters total volume,

(2) 30% retention of radiolead formed by radon decay,

(3) an average distribution coefficient of 0.89 for
man*,

(4) an atmospheric radon concentration of 10^{-13} curies/
liter

$$0.89 \times 10^{-13} \text{ curies/liter} \times 2.22 \times 10^{12} \text{ dpm/curie}$$

$$\times 70 \text{ liters} \times 0.3 = 4.15 \text{ atoms/min deposited}$$

$$\frac{4.15 \text{ atoms/min}}{5.47 \times 10^{-7} \text{ min}} = 7.59 \times 10^6 \text{ atoms of radiolead}$$

stored at equilibrium

c. Stored Pb^{210} by decay of body radium content.

Assume: (1) Average human contains 1.5×10^{-10} grams radium

(38, 39, 40)

(2) 70% of the radon formed is lost in the breath

(41, 42, 43)

*Using values for the distribution coefficients of various rat tissues as found by Nussbaum and Hursh (37) and the per cent of various tissues present in a "standard man" (6), the value of 0.89 was computed.

- (3) 100% retention of the Pb^{210} from radon decay,
since it is formed in the bones.

$$1.5 \times 10^{-10} \text{ curies} \times 0.3 \times 2.22 \times 10^{12} \text{ dpm/curie} =$$

99.9 dpm of retained radon

99.9 atoms of Pb^{210} deposited per minute

$$\frac{99.9}{5.47 \times 10^{-7}} = 1.8 \times 10^8 \text{ atoms of } \text{Pb}^{210} \text{ stored}$$

Therefore, from parts a., b., and c. there could be stored

$$0.304 \times 10^8 + .076 \times 10^8 + 1.8 \times 10^8 = 2.18 \times 10^8 \text{ atoms of } \text{Pb}^{210}$$

$$2.18 \times 10^8 \text{ atoms} \times 6.95 \times 10^{-8} \text{ min.}^{-1} = 15.1 \text{ dpm of } \text{Pb}^{210}$$

The maximum polonium activity from these three sources would
be 15.1 dpm but is probably much less.

Assuming urinary excretion of 0.13% of body content per day

$$\text{gives } 15.1 \times 1.3 \times 10^{-3} = .0197 \text{ dpm per day excreted } \text{Po}^{210}.$$

d. Urinary polonium from food and water

Assume: (1) water contains $.04 \times 10^{-12}$ g Ra per liter (34),

1.5 liters/day consumption (6)

(2) food contains 1×10^{-15} g Ra per gram (44),

1000 grams/day consumption

(3) Po^{210} in equilibrium with the radium

(4) Urinary excretion of 2% of intake (45)

$$.04 \times 10^{-12} \text{ g/liter} \times 1.5 \text{ liters} + 10^{-15} \text{ gm/gm} \times 1000 \text{ gm} =$$

$$1.06 \times 10^{-12} \text{ g Ra}$$

$1.06 \times 2.22 = 2.35$ dpm radium -- also of Po^{210} level no assigned
 $2.35 \times .02 = .047$ dpm Po^{210} in urine.

e. Total calculated urinary excretion

The total urinary excretion would be the sum of the above calculated values, $0.02 + 0.047 = .067$ dpm, or .03 cpm assuming 45% counting efficiency. This is the maximum possible value under the assumptions used. Actually 0.05 ± 0.03 cpm/liter was measured. The agreement of the calculations with the measured value is good considering the uncertainty of the assumptions required for the calculations. The difference may be even greater if the data on the recovery of polonium from digested and undigested rat urine applies to human urine (Table XI).

Some of the factors which might affect the agreement between analyzed and calculated polonium content include:

- (1) One of the persons contributing to the pooled urine may have a greater than average body radium content,
- (2) Some of the assumptions could be modified if better data were available,
- (3) Some of the 70% of the non-absorbed Pb^{210} may contribute activity to the urine,
- (4) The 0.05 ± 0.03 cpm average recovery is less than twice average of background counts (normal range 0.04 to 0.09 cpm) so some statistical uncertainty is present, thus requiring many

samples or larger volumes, to eliminate this effect

(5) The possibility of contamination cannot be eliminated with certainty but is believed unlikely, based on previous experience with the equipment and procedures used.

I. Summary and Conclusions on Section VI.

Determinations were made on the recovery of added inorganic polonium from 200 ml portions of acidified human urine by deposition on silver, nickel, and copper foils through stirring triplicate samples for 2 hours at 94° C. Recoveries exceeded 90% using silver and nickel foils, but recoveries from copper foils ranged from 64 to 78% probably because these foils were attacked by the acid. These data confirm more extensive data previously reported.

Recovery of polonium from the urine of polonium injected rats when acidified and stirred over a silver foil at 94° C for 2 hours was only 83% of the values obtained when the urine was wet digested before plating.

Recovery of added inorganic polonium from liter quantities of urine was better than 90% when the urine was acidified,

stirred 2 hours at 94° C with 2 grams of nickel powder, the powder filtered off, dissolved in aqua regia and the polonium replated on silver foil. In another procedure for recovery of polonium from large volumes, polonium was added to 2 liters of 1N HCl.

The 2-liter bottle was arranged to feed the solution into a 250 ml beaker so that a constant level of fluid was maintained in the beaker during evaporation. The 50 ml of solution remaining after evaporation were diluted to 200 ml with 0.5N HCl then stirred over a silver foil for 2 hours at 94° C. The average recovery was 93%.

The analysis of liter quantities of pooled urine from presumably unexposed personnel by the nickel powder procedure indicated an average activity of $0.11 \pm .07$ dpm per liter.

Calculations were made for expected urinary excretion using values commonly accepted for the radium content of non-exposed humans, and food and water, and for the radon concentration of atmospheric air, and using reasonable assumptions for the amount of stored lead from these sources. These indicated that a maximum of 0.066 dpm per day would be excreted in the urine, a value within the range of that actually found.

VII. RECOVERY OF POLONIUM FROM SOFT TISSUES AND BONE

A. Experiments Based on Nitric Acid Digestion

1. Recovery from mice injected with polonium 24 hours previously.

Six white female mice of the local colony were obtained. Five of the mice were injected intraperitoneally with a known

amount of polonium and placed in cages made up as shown in Figure 5.

The sixth mouse was sacrificed with ether and injected with same amount of polonium as the other five. This was called the control since it contained only non-metabolized Po.

After 24 hours the five live mice were sacrificed by introducing ether into the cages. 50 ml of nitric acid were poured over all six mice in the beakers, after removing only the water supply and covering watch glass. Thus the mice, excreta, and small amount of remaining food were analyzed simultaneously. After cold digestion overnight the beakers were placed in an oil bath at 110°C and boiled to a moist paste. 50 ml more of nitric acid were added to each beaker and again taken to a moist paste. 50 ml of 1N HCl were added to each and boiled to dryness. This was permissible since oil bath temperature was always kept below 150°C , the temperature at which losses of polonium by volatilization begin to occur. An additional 50 ml of 1N HCl were added and again taken to dryness to get rid of any nitric left in the digest. The residue was taken up in 50 ml of 1N HCl, heated to boiling and transferred to plating bottles with 3 rinses of 1N HCl each also brought to a boil. The final plating volume was about 120 ml. 200 mg of ascorbic acid were added to each solution. They were then

Figure 5. Glass cage for polonium injected mice

stirred rapidly in a water bath at 94° C. The recoveries are listed in Table XIII.

TABLE XIII

The recovery of polonium from mice sacrificed and analyzed 24 hours after IP injection of polonium

Sample	Total Counts	Time Min.	Cpm	% Recovery*
1	1181	60	19.7	100
2	15,481	867	17.8	99
3	2,361	153	15.42	86
4	3,528	210	16.8	93.7
5	2,685	161	16.7	93.2
6	13,216	915	14.48	80.7**
Average (Less #6)				94.4

*Corrected for background and decay.

**Foamed over during digestion with loss of material. This mouse was sacrificed at the start of the experiment and injected IP with the polonium solution.

The method appeared successful for recovery of inorganic polonium injected IP in mice even though a small amount of undigested fat remained in the digest.

2. Recoverable polonium from non-exposed mice.

Six female white mice of the local colony were sacrificed

with ether and digested according to the procedure in VII.A.1. above. During this period, backgrounds were run on the counter using a silver foil which had been cleaned by scrubbing with an abrasive powder. Four background runs were made during the week and the average rate is shown in Table XIV.

TABLE XIV

Polonium content of non-exposed mice of the local colony

Sample	CCI	Total Count	Time Min.	Cpm	Net Cpm
Bkg		1054	9,131	0.113	0.113
1	1.88	315	1,103	0.286	0.173
2	8.88	128	465	0.275	0.162
3	88.00	368	1,435	0.265	0.152
4	4.48	1079	4,426	0.246	0.133
5		268	1,302	0.207	0.094
6		347	1,479	0.234	0.121
U-Std II	12058		2009		Counter efficiency 45.2%

Mean polonium content per mouse $0.139 \pm .046$ cpm

These results indicate an average of 0.307 dpm per 25 gram mouse. Reagents add no detectable activity in the quantities used for mouse digests as can be seen by referring to Tables XXII and XXIII.

3. Analysis of bats for polonium.

With the experience gained from the above experiments with nitric acid digestion, the analysis of two bats was attempted. These bats had been trapped in a uranium mine in eastern Utah and were supplied through the courtesy of Dr. Mitchell R. Zavon of the Division of Special Health Services, U.S. Public Health Service, Cincinnati, Ohio. Since it is probable that the bats had been exposed to a higher than average radon concentration they should contain appreciable quantities of analyzable polonium.

Six samples could be run simultaneously with the available equipment so along with the 2 bats were run two mice, a reagent blank, and a reagent sample to which a known amount of polonium was added.

The digestion method was as described in A.1. above. The bats had been preserved in a formaldehyde solution which was divided evenly between the two flasks containing the bats. The results are shown in Table XV.

With 45% counting efficiency and correcting for background and reagent blank, these results indicate that one bat had a total body polonium content of 14.66 μmc and the other 63.63 μmc . Since more than one year elapsed between death and analysis, these figures probably represent the radiolead content of the bats.

TABLE XV

The polonium content of two bats trapped in a uranium mine

Sample	Composition	Total Count	Time Min.	Cpm	Net Cpm	Remarks
Bkgd		111	981	0.113	0.113	
U.-std		9990	5	1998		45% efficiency
1	Digest reagents	519	4360	0.119	0	
2	Spiked reagents	1355	24	56.46	56.35	103% recovery
3	Unexposed mouse	423	1477	0.286	0.173	Avg. of 0.134
4	" "	321	1406	0.208	0.095	cpm per mouse
5	Bat	14829	999	14.84	14.73	
6	"	2042	32	63.81	63.70	

4. Recoveries of added polonium from bone.

a. The initial work on recovery of added polonium from bone is described in Section V.E. above. As shown in Table VI, when 40 to 50 grams of bone are digested some precipitate occurs which is difficult to get in to the normal 200 to 220 ml volume of plating solution. To get good recoveries, 2 or 3 plating bottles must be used. This is no real disadvantage when the sample contains appreciable activity but at very low activities it would be better to have all the activity on one foil. In addition to the precipitate effect, there was the effect of undigested fat. It has been reported by Smith et al that fat contains less than 5%

of the polonium injected. It was necessary to check this fact so experiments were set up using both digestion of bone and fat and digestion of bone with extraction of the fat.

b. Recovery from ashed bone.

Recovery of polonium added to bone when large volumes are required for complete solution of the bone digest was determined as follows.

Ten grams of ash from human bone, ashed in a muffle furnace, were added to each of six flasks, this is equivalent to 20 - 30 grams of fresh bone. A known amount of polonium was added to each flask. Two of the samples were digested with nitric acid according to the procedure developed above. These two samples went into solution in two plating bottles each. To the other four samples, 5 ml of sulfuric were added after the preliminary nitric digestion. When the digests boiled almost dry, 5 ml of concentrated HCl were added and boiled off, then 5 ml of distilled water were added and boiled off. After neutralizing and reacidifying with HCl, it was found that 3 plating bottles were required to get all solids into solution. Two of the samples were plated for 2 hours and the other two plated for 3 hours. The results of this experiment are shown in Table XVI.

TABLE XVI
The recovery of added polonium from bone after digestion with nitric
or nitric-sulfuric acid

Sample	Digest Reagents	Plating Time Hrs.	Total Count	Time Min.	Net Cpm	% Recovery
1a	HNO ₃	2	1452	19	76.32	
1b	"	"	1737	34	50.99	106
2a	"	"	1459	22	66.22	
2b	"	"	7360	140	52.47	99.3
3a	HNO ₃ -H ₂ SO ₄	"	1089	49	22.12	
3b	"	"	1222	35	34.81	
3c	"	"	1841	49	37.47	88.2
4a	"	"	1161	36	32.31	
4b	"	"	2268	75	30.14	
4c	"	"	3003	109	27.45	84.0
5a	"	"	870	26	33.36	
5b	"	"	1486	36	41.18	
5c	"	"	1207	42	28.64	96.4
6a	"	"	1162	33	35.11	
6b	"	"	2462	71	34.57	
6c	"	"	1753	44	39.74	102

These results indicate that recovery of added polonium is quantitative from 10 grams of bone ash even though large volumes are required for complete solution of the digest. The addition of sulfuric acid to aid digestion caused a larger amount of precipitate which required up to 600 ml of solution to dissolve it and a longer plating time for good recoveries.

c. Recovery of added polonium from fresh bone.

Using fresh pig and beef femurs, a series of experiments were performed to check the recovery of added polonium with and without fat extraction, with various treatments prior to fat extraction and with different digestion methods. In outline form, the procedures were as follows:

- (1) Add the nitric acid, heat, add water, extract the fat, continue the digestion of both the extract and residue.
- (2) Add nitric acid, boil to moist paste, add water, extract fat, continue the digestion of both portions.
- (3) Extract the fat before any other treatment, then digest both portions.
- (4) Digest the total sample without extracting the fat.

Some of the digestions, indicated by HNO_3 in Table XVII, were done with the nitric acid method described on page 58. These had a high percentage of discolored foils so some were redigested by using a 2:1 V/V mixture of 72% perchloric and concentrated nitric acids, indicated by HNO_3 , HClO_4 in the table. Many of the samples were digested using the perchloric-nitric acid method as described on page 71 and are indicated by HClO_4 in the table.

Fat extraction was accomplished by adding 50 ml of petroleum ether to the sample in the beaker, warming the mixture to the boiling point of the ether and then shaking it gently. After standing until the two layers separated, the solvent layer was pipetted off carefully. This was repeated twice more with 50 ml portions of petroleum ether for a total volume of 150 ml of extract.

The results of this series of experiments are shown in Table XVII.

These experiments indicate that less than 0.2% of added polonium will be removed from 20 grams of fresh bone when the fat is extracted by petroleum ether after preliminary digestion of the bone with concentrated nitric acid.

Recoveries of added polonium from 20 grams of fresh bone average 95% when the nitric-perchloric digestion method is used whether or not fat extraction is used.

The nitric acid digestion as described on page 58 proved unsuitable for the bone digestions as many discolored foils were obtained. This indicates that boiling the digests down with added concentrated HCl did not remove all the nitric acid from this type of sample.

All the samples from No. 21 to No. 32 inclusive required four hours plating time to obtain the per cent recoveries

TABLE XVII

The recovery of added polonium from bone after various digestion procedures with and without the extraction of fat

Sample	Material	Weight Grams	Procedure*	%Recovery		Remarks
				Bone	Fat	
1	Pig femur	50	(1) HNO ₃ , HC10 ₄	77.2	--	extract spilled
2	" "	51	" " "	88.4	2.6	
3	" "	51	(2) HNO ₃ , HC10 ₄	90.7	0.08	
4	" "	49	" " "	91.5	0.20	
5	" "	10	" " "	94.2	0.14	
6	" "	11	" " "	96.1	0.0	
7	" "	10.6	" " "	94.1	0.0	
8	" "	11.1	" " "	85.3	0.18	
9	Beef femur	21.0	(2) HC10 ₄	101	0.15	
10	" "	20.6	" "	90.5	0.03	
11	" "	19.9	" "	90.1	0.09	
12	" "	21.2	" "	95.5	0.10	
13	" "	20.2	" "	93.5	0.03	
14	" "	20.3	" "	106	0.08	
15	Pig femur	10.9	(3) HNO ₃	76.8	1.5	
16	" "	10.0	" "	78.3	1.5	
17	" "	9.8	(3) HNO ₃ , HC10 ₄	50.4		
18	" "	10.2	" " "	50.9	1.4	All extracts
19	" "	10.1	" " "	53.4		combined
20	" "	9.7	(4) HC10 ₄	86.1		No extracts on
21	Beef femur	21.0	" "	99.1		20 through 32
22	" "	19.5	" "	103		
23	" "	20.4	" "	84.6		
24	" "	20.9	" "	96.0		
25	" "	20.6	" "	101		
26	" "	21.3	" "	92.6		
27	" "	19.5	" "	102		
28	" "	19.2	" "	97.1		
29	" "	19.0	" "	98.3		
30	" "	20.6	" "	82.7		
31	" "	20.4	" "	98.9		
32	" "	20.3	" "	85.6		

Mean recovery 95 ± 11% (less 1,2,15-19)

* This column refers to the treatment before extraction and the digestion procedure as outlined on page 66.

listed. The bone samples of 10 grams or less were plated for only two hours with comparable recoveries.

5. Recovery of added polonium from fat.

To validate the results in the preceding section, it was necessary to determine the percentage recovery of added polonium from fat. The following procedure was used for this purpose.

A known amount of polonium was added to 20 gram samples of "Pure Lard" (commercial hydrogenated animal fats) in 250 ml flasks. Twenty milliliters of concentrated sulfuric acid and 5 ml of concentrated nitric acid were added. This procedure produced a violent reaction which subsided when the flasks were cooled in running water. The digests were heated until they turned black when a few milliliters of nitric acid were added cautiously. This was continued until 300 ml of nitric acid had been used. The solutions still had a dark color indicating incomplete digestion. Five ml of 30% hydrogen peroxide had no apparent effect so 50 ml of a 2:1 perchloric-nitric mixture were added. The solutions were then boiled until they darkened when 10 ml of concentrated nitric acid were added cautiously. This was continued until a light, straw colored fluid resulted which gave off dense, white fumes when heating was continued. The solutions were made alkaline using saturated NaOH solution and bromothymol blue indicator then reacidified to 0.5N with concentrated HCl. A 100 mg tablet of ascorbic acid was added and solutions

transferred with 3 rinses of 0.5N HCl to a prepared plating bottle and stirred rapidly for 2 hours in a 94° C water bath. The recoveries were 93.4, 100, 99.8, 81.2, and 81.8% for an average of 91.2%.

This good recovery of added polonium from lard indicates that the recoveries of less than 0.2% in fat, extracted from samples to which polonium had been added before extraction of the fat, are reliable.

6. Discussion of Part A, Section VII.

Use of the nitric acid digestion method was successful for young mice, 25 g average weight, and bats. With samples which contain more fat, analysis would be more difficult.

Analysis of 25 gram mice of the local colony, presumably non-exposed, resulted in a whole body activity of 0.306 μc of polonium per average mouse for 7 mice.

Recovery of polonium injected intraperitoneally in 25 gram mice 24 hours prior to analysis was quantitative when their excreta was analyzed along with each mouse.

Two bats were analyzed for whole body polonium and activities of 14.66 and 63.63 μc respectively were detected. Higher polonium activities would be expected in these animals whose natural habitat was a uranium mine where the atmospheric radon concentration may be relatively high.

In general, recoveries of added polonium from bone or fat using nitric acid or $\text{HNO}_3\text{-H}_2\text{SO}_4$, as used by Scott and Stannard (26), were unsuccessful with the modified procedure described.

B. Method Developed for Analysis of Tissue and Bone for Polonium

Content

1. As a result of experiments reported above the digestion method considered most satisfactory was as follows:

a. Select a digestion vessel such that the sample weight is 10 to 20% of the vessel volume, e.g. a 250 ml flask for a 25 gram mouse or a 1000 ml beaker for a 200 gram rat.

b. Add 2 ml of concentrated nitric acid per gram of material up to 50 grams and 1 ml per gram for over 50 grams, e.g. 50 ml per 25 gram mouse or 200 ml per 200 gram rat.

c. After digesting at room temperature several hours, preferably overnight, heat gently, cool when foaming becomes excessive, until a moist paste results.

d. If more than 1 ml of fat is present at this stage, extract with three additions of petroleum ether, 50 ml per addition, swirl the mixture gently, then carefully pipette off the solvent layer after each addition.

e. Add a 2:1 mixture V/V of 70% perchloric acid and concentrated nitric acid to the digests in the same amount as the initial addition of nitric acid in b. above. For bone add 3 ml

of the mixture per gram.

f. Boil down until dense white fumes appear. If foam-
ing starts, with charring of the digest at the surface of the
solution, cool and add very cautiously a few ml of concentrated
nitric acid. Repeat until further heating produces the dense white
fumes.

g. Add 1 ml of concentrated HCl slowly, if no brown fumes
evolve and the solution undergoes only a transient color change
then all the nitric has been destroyed.

h. Make alkaline using NaOH pellets and/or a saturated
NaOH solution with bromothymol blue indicator.

i. Add sufficient concentrated HCl to make the solution
0.5N.

j. Cover the vessels and boil gently for 2 hours or more.

k. Transfer to a prepared plating bottle with 3 rinses of
0.5N HCl and add sufficient 0.5N HCl to make the final volume
approximately 200 ml.

l. Add 100 mg of ascorbic acid per 50 grams or less of
digested tissue and 200 mg for larger quantities including rats
of up to 200 grams.

m. Stir rapidly (600 rpm or more) in a 90 to 100°C water
bath for 2 hours, except for more than 10 grams of bone, when
4 hours are required.

This method has been used for recovering added inorganic polonium from up to 50 grams of bone and soft tissue and from mice and rats with recoveries of 90% or more. A modification of the method is required for rats of 130 grams or larger. These digests will not go into solution in the 200 ml plating volume so they must be divided among two plating bottles, with the result that the polonium is plated on two foils, and they must be stirred for four hours as is required for more than 10 grams of bone.

2. Check on the above method by determination of polonium activity of beef bone.

To check the above method the natural polonium activity of two 30 gram samples of beef femur was determined using that method. Six samples were run. They included the two bone samples, two reagent blanks, and two reagent samples to which was added a known amount of polonium. All samples were treated exactly alike, i.e. when something was added to the bone digest, the same was added to the other four samples.

The results are set forth in Table XVIII.

These results indicate a polonium activity of 0.016 μ mc per gram above reagent blank for the beef femur investigated.

TABLE XVIII

The polonium content of beef femur samples used in these experiments

Sample	Composition	Total Count	Time Min.	Cpm	Net Cpm	% Recovery
1	Digest reagents	159	979	0.162	--	--
2	" "	24	135	0.178	--	--
3	Reagents + Po	1012	13	77.84	77.67	97.6
4	" " "	1126	15	75.07	74.90	94.1
5	30.8g beef femur	625	944	0.662	0.492	
6	31.2g " "	303	457	0.666	0.496	
Average activity of beef femur 0.016 $\mu\text{c}/\text{gram}$						

The average recovery of 95.8% of added polonium, also above reagent blank, shows that the nitric-perchloric method is a satisfactory digestion procedure leading to quantitative recoveries of added polonium.

C. Studies on Rats Killed Immediately after Radon Exposure

1. Recovery of polonium from rats.

Two rats which had been exposed to a daughter-free radon atmosphere 8 months previously, in a experiment of Dr. Donald Morken of this institution, were obtained. They had been exposed for 3 1/2 hours to a daughter-free radon concentration of 125 $\mu\text{c}/\text{liter}$, and were sacrificed immediately on termination of the exposure and gross

autopsy performed on them. They were then wrapped in paper
towel and stored in a cold room until removal for the present
experiment.

Four more rats were obtained from the local colony and
killed with ether. These were analyzed along with the exposed
rats, two were used as controls and the other two had a known
amount of Po added. A reagent blank and a reagent with a known
amount of polonium added were also run in this series.

All samples were digested using the nitric-perchloric
method described above. The fat was extracted from all four of
the unexposed rats and the two exposed rats. The fat was digested
by the nitric-perchloric method also.

The four digests of unexposed rats were divided among two
plating bottles each and were stirred for four hours. The ex-
posed rat digests, after neutralization and reacidification, were
made up to a volume of 500 ml in a volumetric flask using 0.5N HCl.
Only 5 ml of this volume (1%) were added to 200 ml of 0.5N HCl in
a plating bottle and stirred for two hours.

The results of this experiment are shown in Table XIX.

TABLE XIX

The polonium content of two rats exposed to a daughter-free radon level of 125 $\mu\text{c}/\text{liter}$ for 3.5 hours, sacrificed immediately, and analyzed 246 days after exposure.

Sample	Composition	Rat Wt.	Cpm	Net Cpm	Fat		Remarks
					Net Cpm	%	
U-Std.				1996			45% Counter Eff.
Bkgd.				0.08			
1	Rat + Po	142.4	79.51	79.35	0.066	.08	101% recovery
2	" " "	164.6	71.47	71.31	0.021	.03	91.3% recovery
3	Control rat	150.4	0.92	0.76	0.067		
4	" "	137.7	0.80	0.64	0.026		
5	Digest reagents		0.077	0			
6	Reagents + Po		67.1	67.02			86% recovery*
7	Exposed Rat	187	39.7	39.62	0.111	.002	1% of digest used
8	" "	202	40.4	40.32	**	---	" " "

For the whole rat the activities are $8.8 \times 10^3 \text{ dpm}/187 \text{ g}$
and $8.95 \times 10^3 \text{ dpm}/202 \text{ g}$

* This sample foamed over during digestion.

** The extract was contaminated by foam from the spiked reagents.

These results show that 96% of the added activity was recovered. This probably should also apply to the rats exposed to radon. In confirmation of earlier results on fat extraction from spiked fresh bone, less than 0.1% of the activity in the sample was

present in the petroleum ether extract, even in the case of the rat exposed to radon.

This result is considered significant since it would be expected that a high percentage of the polonium resulting from radon decay would be in the fat. These rats were killed immediately after exposure so a good percentage of the radon in the fat should have decayed in situ.

The reagent blank showed no measurable activity above background.

2. Calculations on the amount of polonium deposited due to Rn exposure.

a. Rats exposed 3.5 hrs. to daughter-free radon concentration of 1.25×10^{-4} curies/liter on 7-25-55 and analyzed on 3-27-56.

b. A rat has 7.08% fat, 45.5% muscle, 4.95% blood, and 42.5% of organs and bone as determined by Caster et al (46).

c. The distribution coefficients are 4.83, 0.15, 0.4, and 0.3 for fat, muscle, blood, and organs respectively (37). The average for body less fat can be calculated from these data as 0.23.

d. Assume:

- (1) Radon build-up in the body is of the form $x = y^2$
- (2) No radon leakage after exposure.
- (3) Radon build-up is 85% of maximum in the fat in 3.5

hours (37)

(4) Radon build-up is maximum in the body, less fat,
in 1 hour (37)

(5) Specific gravity of rat tissues is 1.

e. The activity vs. time curve is shown in Figure 6.

f. Build-up of radon -- area under the curve $x = y^2$ is $2/3$.

Body:

(1) During build up: $1.25 \times 10^8 \mu\text{c/liter} \times 2.22 \text{ dpm}/\mu\text{c}$
 $\times 10^{-3} \text{ liter/g} \times 0.23 \times 60 \text{ min.} \times 2/3 = 2.55 \times 10^6 \text{ dis/g}$

(2) Remainder of exposure: $2.77 \times 10^8 \text{ dpm/l} \times 10^{-3} \text{ liter/gram}$
 $\times 0.23 \times 150 \text{ min} = 9.57 \times 10^6 \text{ dis/gram}$

Fat: $2.77 \times 10^8 \text{ dpm/liter} \times 10^{-3} \text{ liter/gram} \times 4.83 \times 0.85 \times 2/3$
 $\times 210 \text{ min} = 1.58 \times 10^8 \text{ dis/gram}$

g. Decay of radon

Body: $2.77 \times 10^8 \text{ dpm/liter} \times 10^{-3} \text{ liter/g} \times 0.23 \times 7.96 \times 10^3 \text{ min} =$
 $5.07 \times 10^8 \text{ dis/g}$

Fat: $2.77 \times 10^8 \text{ dpm/liter} \times 10^{-3} \text{ liter/g} \times 4.83 \times 0.85 \times 7.96 \times 10^3 \text{ min.} =$
 $9 \times 10^9 \text{ dis/g}$

where $7.96 \times 10^3 \text{ min}$ is the average life ($\frac{1}{\lambda}$) of radon

Each of the disintegrations ultimately deposits one atom
of Pb^{210} .

h. Total Pb^{210} deposited

$(.0255 + .0957 + 5.07) \times 10^8 \times 0.9292 = 4.82 \times 10^8 \text{ atoms/}$
gram total weight from body less fat

Figure 6. Build-up and decay of radon in fat of rats sacrificed immediately after exposure.

Figure 7. Build-up of radon (1), maintenance of level (2), and blowoff (3) in tissues of mice allowed to live after exposure.

$$(1.58 + 90) \times 10^8 \times 0.0708 = 6.48 \times 10^8 \text{ atoms/gram body}$$

weight from fat

$$(6.48 + 4.82) \times 10^8 \times 187 \text{ g} = 2.11 \times 10^{11} \text{ atoms Pb}^{210}$$

$$11.3 \times 10^8 \times 202 \text{ g} = 2.28 \times 10^{11} \text{ atoms Pb}^{210}$$

1. Calculated amount of polonium

From 7-25-55 to 3-27-56 is 246 days between exposure

and analysis.

At equilibrium polonium would have the same activity as

the lead but at 246 days the activity would be $N/N_0 = 1 - e^{-\lambda t} = 0.709$ of maximum. Therefore the Po activity is:

$$2.11 \times 10^{11} \text{ atoms} \times 6.95 \times 10^8 \text{ min}^{-1} \times 0.709 = 1.04 \times 10^4 \text{ dpm (187g)}$$

$$2.28 \times 10^{11} \times 6.95 \times 10^8 \times 0.709 = 1.12 \times 10^4 \text{ dpm (202 g)}$$

The actual recoveries were: 8.8×10^3 dpm for the 187 gram rat and 8.95×10^3 dpm for the 202 gram rat. The calculations give results in fair agreement considering the uncertainty of the assumptions that had to be made.

3. Discussion on Section C.

The recovery of added polonium from rats and reagents using the nitric-perchloric digestion method was approximately 96%.

Two rats exposed 3.5 hours to a daughter-free radon concentration of 1.25×10^{-4} curies/liter and sacrificed immediately after exposure were analyzed for polonium by the nitric-perchloric method.

The analysis showed 8.8×10^3 dpm for a 187 gram rat and

8.95×10^3 dpm for a 202 gram rat. Calculations on the amount of activity to be expected gave results close to those actually found.

The calculations of the amount of polonium present were made using values for the distribution coefficient of various tissues as determined at this institution by Nussbaum and Hursh (37).

Thus these results partially confirm those values, the major part of the difference in the amount recovered probably being due to leakage of radon from the carcass after death.

D. Studies on Mice Exposed to Radon and Allowed to Live until Death.

Two groups of white mice of the local colony were available. One group of ten mice had been exposed to 1 mc-hr/liter of daughter-

free radon and one group of nine mice had been exposed to 3 mc-hr/liter. These mice had been exposed by Dr. D. A. Morken, of this institution for another purpose.

1. Polonium content of mice exposed to 1 mc-hr/liter of radon

These mice had been exposed to a daughter-free radon atmosphere on 3-11-54 and were analyzed in May 1956. The males had been exposed to 1.53×10^{-4} curies/liter of radon for 6.5 hours and the females to 2×10^{-4} curies/liter for 5 hrs. These animals were allowed to die naturally. They had survived for as long a time as the control animals of the same experiment.

This experiment was set up so that a mouse, a digestion

flask, a plating bottle, and a foil were a unit. Since no measurable alpha activity is added by the reagents (cf. Table XXIII) the only source of extraneous activity is the glassware. A digestion procedure and subsequent plating should remove better than 90% of any activity in the glassware. Therefore, the quantities of reagents necessary for a nitric-perchloric method digestion of a 25 gram mouse were added to the digestion flasks and the digestion carried out as if tissue were present. After plating of the reagent digests the foils were counted and these counts served as blanks for the mouse digests. The counts on the foils were in the same range as those on foils which were merely cleaned and then counted. This indicates both the efficiency of the decontamination procedure, and the negligible activity present in the quantities of reagents used.

The mice were weighed, assigned a unit of apparatus and digested by the nitric-perchloric method. The weights listed in the table are the weights after death and not the weight at exposure time. A non-exposed control mouse was run along with three similar non-exposed mice to which a known amount of polonium were added, approximately 8 cpm to one mouse and 28 cpm to the other two.

The results of this experiment are shown in Table XX.

TABLE XX

The polonium content of mice exposed to a daughter-free radon atmosphere of 200 pc/liter for 5 hours, living a nearly normal life span. Analyzed 774 days after exposure.

Sample	Composition	Sex	Wt. gm.	Cpm Digest	Blank	Net Cpm	Remarks
U-Std						1944	45% Counter Eff.
1	Exposed mouse	M	15.7	7.57	0.072	7.50	
2	" "	F	16.9	11.83	0.04	11.79	
3	" "	M	19.6	9.43	0.05	9.38	
4	" "	F	16.3	11.56	0.06	11.50	
5	" "	M	19.1	9.28	0.06	9.22	
6	Control mouse	M	18.3	0.45	0.13	0.32	
7	Exposed mouse	M	14.3	9.38	0.09	9.29	
8	" "	F	13.7	9.81	0.07	9.74	
9	" "	F	8.7	10.51	0.09	10.42	
10	" "	F	17.6	5.61	0.07	5.54	
11	" "	M	16.6	11.98	0.05	11.93	
12	Mouse + Po	F	37.9	7.83	0.05	7.78	93.2% Recovery
13	" "	F	27.9	27.22	0.09	27.13	93.1% "
14	" "	F	24.0	26.65	0.07	26.58	91.3% "

The average recovery from the males was 20.24 dpm above control mouse and reagent blank.

The average recovery from the females was 21.06 dpm above control mouse and reagent blank.

2. Calculations on expected amount of polonium due to radon exposure.

a. Assumptions:

- (1) Mouse has 7% fat with distribution coefficient of 4.85 and maximum radon content is attained in 4 hrs. [Using published values for the rat (46)].
- (2) Average distribution coefficient for remainder of body is 0.23 with maximum radon content attained in 1 hour.
- (3) Radon in the body is blown off after exposure at the same rate as it is built up at the start of exposure.
- (4) Lung deposited daughters are negligible under the conditions of this experiment.
- (5) 30% of the Pb^{210} deposited in the body is retained (36).
- (6) The effective half-life of Pb^{210} is 880 days (13).

b. There are three parts to deal with, see Figure 7.

- (1) Build-up of Rn.
- (2) Maintenance of Rn level during exposure.
- (3) Blowing off the Rn after exposure.

c. Exposure: Male mice exposed to 1.53×10^{-4} curies/liter for 6.5 hours on 3-11-54 analyzed on 5-11-56.

d. Calculations for fat.

- (1) $(0.07 \times 25 \text{ grams}) \times 10^{-3} \text{ liters/gram} \times 1.53 \times 10^{-4} \text{ curies/liter} \times 2.22 \times 10^{12} \text{ dpm/curie} \times 4.85 \times 240 \text{ min} \times 2/3 =$

4.59×10^8 atoms Pb^{210} deposited. $2/3$ is the area under the curve $x = y^2$, which closely approximates the build-up curve.

(2) $1.75 \text{ grams} \times 10^{-3} \text{ liters/gram} \times 3.4 \times 10^8 \text{ dpm/liter} \times 4.85 \text{ min} = 4.3 \times 10^8$ atoms of Pb^{210} deposited

(3) During blowing-off period, the amount of Pb^{210} deposited is same as in (1).

e. Calculations for body less fat.

Similar calculations using 0.23 distribution coefficient as calculated previously for rats with 1 hour saturation time give:

(1) 6.54×10^6 atoms

(2) 5.99×10^8 atoms

(3) 6.54×10^6 atoms

f. Total Pb^{210} deposited is 1.96×10^9 atoms

30% retention gives 5.88×10^8 atoms retained in body.

In 774 days from exposure to analysis there will be:

$$\frac{N}{N^0} = e^{0.693t/T} = 2^{774/880} = 1.84, (5.88/1.84) \times 10^8 =$$

3.19×10^8 atoms left. These will have an activity of

$$3.19 \times 10^8 \times 6.95 \times 10^{-8} / \text{min} = 22.2 \text{ dpm of lead.}$$

The polonium activity, assuming 100% retention, should be

the same since 774 days elapsed between exposure and analysis.

g. Similar calculations for the female mice exposed to

2×10^{-4} curies/liter of daughter-free radon for 5

hours give 22.8 dpm of lead. The polonium activity

(at 100% retention) should be equal to the lead activ-

ity as the lead was deposited 774 days previously.

h. Analytical results compared to calculated amount.

$$\frac{20.24 \text{ dpm recovered} \times 100}{22.2 \text{ dpm calculated}} = 91.2\% \text{ for males}$$

$$\frac{21.06 \text{ dpm} \times 100}{22.8} = 92.3\% \text{ for females}$$

This agreement is probably fortuitous. If the polonium has an effective half-life of 40 days as it does after intravenous injection, then it can be calculated that only 6.4 dpm would be recovered. This in turn may be inaccurate since the majority of the polonium is probably formed within the bone and so may have a longer effective half-life; cf. Section 3.b. below. On the other hand, radon decay products could have been ingested by licking the fur. Also, since mice tend to huddle, the local radon concentration after exposure could have been elevated for some time while the mice were blowing off their radon store. Since this occurred in the cages, daughter products deposited on dust could have been retained in the lungs, thus adding some radiolead to the body stores. There is some uncertainty in the rate of

blow-off, the rate chosen was used since symmetry seemed most logical though Hollcroft and Lorenz (47) have found that elimination of body radon is practically complete in one hour after IV and IP injection of radon solution. Finally equilibration of body tissues with atmospheric radon may be more rapid in the mouse than in the rat.

3. Polonium recovery from bones of radon exposed mice.

a. Introduction.

There were available for this experiment 9 mice which had been exposed to a daughter-free radon concentration of 0.2 mc/liter for 15 hours and which had died of this exposure 90 to 100 days later. These mice were divided into 3 groups, one to be analyzed in toto and the other two to be analyzed after separation of the bones from the other tissues by two different methods. Analysis was performed approximately one month after death.

b. Experimental procedure.

Three of the mice were analyzed for whole body polonium content by the nitric-perchloric method. Three more of the mice were boiled in water to separate the bones from the other tissues and a third group of three mice was digested with a solution of 3% KOH in ethylene glycol to separate the bones from the other tissue as described by Crowell et al (48).

After separation of the bones, the solutions were filtered through a Whatman #41 filter under suction. The bones were picked

off of the filter paper with tweezers and analyzed separately.

The filter paper was added to the filtrate and this combination analyzed.

The separation by boiling was unsuccessful, a black tarry residue full of hairs being deposited on the paper which made it impossible to pick out the bones. The portions were therefore recombined and then analyzed for whole body content.

All digestions were done by use of the nitric-perchloric method with all glassware being organized and control runs (blanks) made as in D.I. above. The results of this experiment are shown in Table XXI.

c. Discussion. The method for separating soft tissues from bone was fairly successful. The best separation showed only 4.85% of body weight was bone. This is low since many very small bones could not be picked out of the residue on the filter paper and because some bone probably went into solution. In the first separation, many of the long bones apparently contained marrow because of reddish material visible through the bone shaft. The other separations yielded bones having a bleached appearance and in the last one, a spiked mouse, no bone was separable so this sample was discarded. The various separations were in the alcoholic KOH solution for progressively longer times.

The digestion of the ethylene glycol filtrate was difficult

TABLE XXI

Polonium content of mice which had been exposed to a daughter-free radon concentration of 200 $\mu\text{C}/\text{liter}$ for 15 hours and had died of this exposure in 90 to 100 days. For three of these, the bones were analyzed separately from the soft tissue. Analysis was approximately one month after death.

Sample	Composition	Wt. gram	Sex	Digest	Cpm Blank	Net Cpm*	Remarks
1	Exposed mouse	24.1	M	62.23	0.08	62.01	137.8 dpm
2	"	15.7	F	54.85	0.05	54.66	121.4 "
3	"	15.6	M	111.45	0.05	111.26	247.2 "
4	"	14.6	F	62.97	0.05	62.78	139.5 "
5	"	19.4	M	58.10	0.07	58.89	130.8 "
6	"	15.1	F	74.88	0.07	74.67	165.9 "
7	Mouse + Po	23.5	F	25.18	0.05	25.04	96.2% recovery
8	"	24.3	F	24.27	0.06	24.07	93.1% "
9	"	25.0	F	25.71	0.08	25.49	99.6% "
10	"	22.8	F	23.98	0.06	23.78	92.9% "
11a	Exposed mouse soft tissues	16.65	F	18.72	0.07	18.58	103.1 dpm
12a	"	27.1	M	36.75	0.06	36.62	113.9 "
13a	"	19.17	M	37.48	0.07	37.34	151.3 "
14a	Mouse + Po soft tissues	26.2	F	---	allowed to go dry, sample blew up		
11b	Exposed mouse bone	0.853	F	27.93	0.06	27.80	60% of total
12b	"	0.510	M	14.79	0.09	14.63	28.6% of total
13b	"	0.827		30.89	0.06	30.76	45% of total
14b	Mouse + Po bones	0.782	F	0.673	0.06	0.539	2.1% of total

*Net Cpm is the cpm minus blank counts and the cpm obtained as an average when six non-exposed mice of similar age were analyzed previously (Table XIV). For the separated tissues 1/2 of the 0.138 cpm was subtracted from each fraction.

due to the readiness of the sample to foam when nitric acid was added.

Since not more than 2% of the Po added to the control mouse appeared in the bone fraction after treatment with 3% KOH in ethylene glycol and filtration to separate the bones, it is possible that 50% or more of the polonium formed as a result of exposure of mice to radon is contained in the bones of these mice. This fact, if true, indicates a greater retention of polonium when formed in the bones by decay of Bi^{210} from Pb^{210} , as compared to retention after oral or intravenous administration, with a consequent longer effective half-life. Due to this, the urinary excretion may be less than the 0.13% of body content per day as shown by Stannard et al (45, 49, 50).

d. Calculations. The calculations are similar to those set forth in 2. above. For the various groups of exposed mice, the calculations yield 118 ± 2 dpm of Pb^{210} . The recovery for the 9 mice averaged 145.6 dpm with a range of 103 to 247 dpm of polonium. Thus the calculations give a maximum value which is only 80% of that determined experimentally. If one takes into account the time between exposure and analysis, the polonium would have built up to only 62.8 dpm, assuming no biological elimination, or only 43% of the activity detected. On the other hand, if one assumes the 40 day effective half-life found after

parenteral administration, the equilibrium activity would be only 34.1 dpm and the 120 days between exposure and analysis would permit a build-up of polonium to only 88% of this, which in turn is but 23% of the activity recovered. These considerations accentuate the uncertainties in the assumptions used in the calculations.

Obviously more data are needed on the retention of radon, the retention and distribution of polonium formed from decay of radiolead in the mice, the per cent retention of radiolead formed by radon in the body, and the rate of build-up of radon in mouse tissues.

E. Studies on Human Autopsy Material

1. Introduction.

There was available for analysis a one-gallon jar containing some lung and a piece of sternum with a portion of the ribs attached.* This material was obtained at autopsy from a miner who had died of lung cancer more than one year prior to analysis. It was preserved in formaldehyde. Since all the material was in the same jar, there was a possibility that the formaldehyde had leached some activity from parts of the sample and distributed it to other parts. There was also the possibility that the

* This material was obtained through the courtesy of Dr. James S. Arnold, Radiobiology Laboratory, University of Utah College of Medicine, Salt Lake City, Utah

preserved tissue differed in weight from the fresh tissue.

Finally, before any statements could be made relative to the polonium content of the miner's samples, the content of similar tissues from non-miners should be known.

This study was therefore performed in three parts, (1) experiments on formaldehyde, (2) experiments on lung tissue, and (3) on rib and sternum.

2. Experiments with formaldehyde.

a. Effect of preservation on tissue weight.

Some human lung and ribs obtained at local autopsy were used for this experiment. The material had been placed in an enamel bowl and covered with paper towels. It was weighed within two hours of removal by placing in a tared 800 ml beaker. The material was covered with a 10% solution of formaldehyde in water and allowed to soak for 8 days. The formaldehyde solution was then poured off and the tissues re-weighed with the results shown in Table XXII.

From these results it is seen that there is no change in bone weight, but there is a 7% increase in lung weight after soaking 8 days in a 10% formaldehyde solution.

b. Leaching of polonium from lung by formaldehyde.

Two formaldehyde solutions were used for the experiment. One was obtained from the miner's autopsy sample. The other was

TABLE XXII

The change in weight of bone and lung after soaking 8 days in a 10% solution of formaldehyde in water

	Weight - grams			
	Fresh		After Soaking	
	Bone	Lung	Bone	Lung
Sample + tare	37.5	442.0	37.6	443.2
Tare	<u>4.2</u>	<u>240.6</u>	<u>4.3</u>	<u>227.6</u>
Sample	33.3	201.4	33.3	215.6

obtained by soaking the lungs of a rabbit in a 10% formaldehyde solution for 36 hours. The lungs were from a rabbit which had received polonium by intratracheal injection of a polonium solution and had been sacrificed 48 hours after the injection.

The rabbit lungs and 25 ml of the formaldehyde solution in which they had been soaked were digested separately using the nitric-perchloric method. After the refluxing stage, the digests were transferred to 200 ml volumetric flasks with three 20 ml rinses of 0.5N HCl and made up to volume with 0.5N HCl. The plating bottles were filled with 200 ml of 0.5N HCl and 2 ml (1%) of the digest solutions were added to each plating bottle.

The formaldehyde solution from the miner's autopsy sample was also digested by the nitric-perchloric method and analyzed

according to the usual procedure. Four 25 ml samples were run along with two samples of 25 ml of distilled water. One of the formaldehyde samples and both of the water samples contained added polonium. The foils for each were cleaned and counted (1st Cpm in Table XXIII). Then each of the digestion flasks was run through the digestion and plating procedures using only the reagents. The foils were counted again after this procedure (blank counts). The formaldehyde and water digests were then plated on the foils. The results of this experiment are shown in Table XXIII below.

From these results the following conclusions can be drawn:

- (1) The recovery of added polonium from formaldehyde solution is quantitative.
- (2) The reagents contained no polonium detectable by the method used.
- (3) The preservative solution from the miner's autopsy sample contained no polonium detectable by the method used.
- (4) Only 0.23% of the activity present in the rabbit's lungs was removed by soaking them for 36 hours in a 10% solution of formaldehyde in water.

The above conclusions suggest that no polonium was leached from active portions of the miner's tissues by the preservative solution and further suggest that no polonium was transferred to

TABLE XXIII

The polonium content of formaldehyde solutions used to preserve various polonium containing samples

Sample	Composition	Sample Cpm	Folios		Sample Cpm	Remarks
			1st Cpm	2nd Cpm		
1	HCHO from rabbit lung	15.2		0.10	15.1	1510 cpm in total sample
2	Rabbit lung	6653		0.09	6653	6653×10^2 cpm in whole lung
3	Distilled water + Po	10.58	0.066	0.076	10.50	199.1% recovery
4	" "	11.0	0.053	0.091	10.91	103%
5	HCHO from miner's sample + Po	10.52	0.066	0.070	10.45	98.7%
6	HCHO from miner's sample	0.092	0.095	0.052	0	
7	" "	0.063	0.064	0.059	0	
8	" "	0.096	0.045	0.095	0	

less active portions of the tissue by the preservative used.

3. Experiments with bone.

a. Experimental procedure

The miner's autopsy sample of bone consisted of a piece of sternum with portions of two ribs attached. The sample weighed 58.45 grams when removed from the preservative solution. It was cut approximately in half by cutting down the middle of the sternum. One of the halves was saved for future analyses and the other cut in half by cutting down the middle of the attached rib. Two local autopsy samples consisting of rib and sternum were obtained from two men with no history of exposure to radioactive materials. They had died of heart disease and brain tumor respectively. The fresh samples weighed 118 grams and 84 grams. Each sample was divided into approximately 40 gram portions.

The bone samples were digested, using the nitric-perchloric method described in Section VII B, page 71. All were stirred for four hours to obtain maximum recovery.

The results of this experiment are shown in Table XXIV.

b. Results.

These results show a distinct difference between the polonium content of bones from a uranium miner and similar bones from 3 presumably unexposed humans.

TABLE XXIV

The polonium content of bone from a uranium miner compared with bone from people with no history of radium or radon exposure

Sample	Composition	Weight grams	Cpm	Net Cpm	Remarks
1	Rib and sternum } #A16270	42.4	1.49	1.37	.0323 $\mu\text{c}/\text{gram}$
2	" " " }	44.5	1.48	1.36	.0306 " "
3	" " " } #A16268	46.6	3.63	3.51	.0753 " "
4	" " " }	33.8	0.525	0.403	.0119 " "
5	" " ")	37.5	0.651	0.529	.0141 " "
6	Digest reagents		0.122	0.	
7	Reagents + Po		8.59	8.47	98.3% recovery
8	Ribs } #A16316	9.3	0.118	0	
9	" }	11.5	0.164	0.042	.0036 $\mu\text{c}/\text{gram}$
10	Miner's rib + sternum	14.32	11.0	10.88	0.760 " "
11	" " "	11.69	6.11	5.99	0.512 " "

The following statements can be made from the above results:

(1). In the normal human bone samples there may exist a 7:1 difference in the polonium activity of 40 gram samples. This variability has been reported for radium in bones (51, 52).

(2). The average from #A16270 was $0.0314 \mu\text{c/g}$ and for #A16268 $0.0377 \mu\text{c/g}$.

(3). The average polonium content of the miner's bone was $0.648 \mu\text{c/g}$, an activity 18 times the activity of the normal bone.

c. Discussion.

(1). A partial check on these results can be made by comparing the expected urinary excretion with that determined for uranium miners by Sultzer and Hursh (13).

Assuming: (a). 7 kg of bone in a 70 kg man (6),

(b). Activity found in the bone represents the average activity of the whole skeleton,

(c). 50% of the body polonium content is in the skeleton,

(d). 0.1% urinary excretion

Then $0.648 \mu\text{c/gram} \times 7 \times 10^3 \text{ grams} = 4.53 \times 10^{-3} \mu\text{c in bone}$

$2 \times 4.53 \times 10^{-3} \mu\text{c} = 9.06 \times 10^{-3} \mu\text{c in whole body} = 20.1 \times 10^3 \text{ dpm}$

Therefore $1 \times 10^{-3} \times 20.1 \times 10^3 = 20.1 \text{ dpm per 24-hour urine.}$

The calculated whole body content more nearly reflects the radiolead content at death rather than the polonium content at

that time due to the long interval between death and bone analysis, but even with a factor of 3 correction (40 day effective against 138 day radioactive half-life) the calculated excretion of 6.7 dpm ($3 \mu\text{c}$) is within the range of values reported.

(2) For normal bone one can check against the radium content of unexposed persons as determined by Hursh and Gates (38).

Assuming: a. 7 kg of bone in a 70 kg man

b. 85% of the body polonium is due to radium.--

(according to the calculations of Section

IV H above).

3×10^{-14} curies/gram $\times 7 \times 10^3$ grams = 2.1×10^{-10} curies Po in skeleton

2.1×10^{-10} curies $\times 0.85 = 1.8 \times 10^{-10}$ curies Ra in skeleton

The average value of the cases analyzed by Hursh and Gates was 1.5×10^{-10} grams of radium whole body content.

4. Experiments with lung tissue.

For polonium analysis of the miner's lung, approximately

50 gram samples were taken from the outer portion of the lobes.

For parallel analysis, a peripheral portion of the lung taken

at local autopsy from a person with no history of exposure to

radioactive material, was used. He had died of a heart condition.

TABLE XXV

The polonium content of lung from a uranium miner compared with lung from a person with no history of radium or radon exposure

Sample	Composition	Weight Grams	Cpm	Blank	Net Cpm	Remarks
1	Digest reagents + Po		8.59	.06	8.53	99% recovery
2	" " "		8.75	.06	8.69	101% recovery
3	Normal lung + Po	49.9	9.14	.07	9.07	100% recovery
4	Normal lung	49.5	0.195	.04	0.155	
5	" "	52.0	0.380	.07	0.310	
6	" "	53.6	0.215	.04	0.175	
7	Miner's lung + Po	48.1	14.5	.07	14.43	10.6 cpm added
8	Miner's lung	49.1	5.35	.08	5.27	
9	" "	51.2	4.51	.06	4.45	
10	" "	50.4	6.56	.11	6.45	
11	" "	51.9	5.75	.26	5.49	

A known amount of polonium was added to one of the miner's samples and to one of the "normal" lung samples.

These samples were digested and plated by the nitric-perchloric method described previously with the results shown in Table XXV.

These results show an average of $0.0057 \mu\text{c}/\text{gram}$ with a range of $.0031$ to $.0092 \mu\text{c}/\text{g}$ for unexposed lung compared to an average of $0.1 \mu\text{c}/\text{gram}$ with a range of 0.080 to $0.128 \mu\text{c}/\text{gram}$ for the miner's lung. The average values indicate that miner's lung has 20 times as much activity as unexposed lung, which is about the same ratio as for miner's bone and bone from unexposed persons.

The column headed "blank counts" represents counts obtained on the foils when the digestion reagents, without added tissue, were run through the digestion and plating procedure.

VIII. SUMMARY AND CONCLUSIONS

Methods were developed for the analysis of biological samples for polonium when the polonium levels of such samples were at or near those occurring naturally. The method for soft tissue and bone permits the isolation of the polonium from 50 grams of soft tissue or bone on a single silver foil suitable for alpha counting. Methods for urine allow the isolation of polonium on a single foil from one liter of urine.

Samples of rib and sternum obtained at autopsy from a miner who had died of lung cancer more than one year previously were analyzed for polonium. The average polonium content was 0.65 μc per gram. Due to the long interval between death and tissue analysis, the polonium is due mostly to the radiolead content of this tissue. This miner, from the Colorado Plateau, had worked in uranium mines in that area. Comparable bone samples from local autopsies with no history of radium or radon exposure showed polonium concentrations nearly twenty times lower.

Lung samples from this uranium miner showed a polonium content ranging from 0.08 to 0.128 μc per gram wet weight. Lung obtained at local autopsy from unexposed persons was again nearly twenty times lower in polonium content.

The method for soft tissue and bone was a modification of one developed by Fink and was used in the analysis of mice and rats, up to 50 grams of dog, pig, beef and human bone, and up to 50 grams of human lung. Analysis of these tissues for polonium activities in the range of 0.1 to 100 μc yielded recoveries of 90% or better for added inorganic polonium. The method included wet digestion with nitric and perchloric acid followed by plating of the polonium on silver foil. The "plating" is actually an electrochemical displacement process wherein the polonium replaces silver on the silver foil.

Liter quantities of urine were analyzed for added polonium by initial plating on nickel powder followed by solution of the powder in aqua regia and replating on silver foil.

As a result of the experiments performed by the methods described in the course of this work, the following conclusions can be drawn.

1. Recovery of added inorganic polonium in the activity range of 2 to 200 dpm from 0.5N HCl is not dependent on the activity level of the polonium.

2. Concentrations of nitric acid of 0.16N or greater in the plating solution decrease the per cent recovery appreciably. The reduced recovery is apparently due to a reaction of the silver foil with the nitric acid which results in a layer of material over the foil which shields the foil and prevents the electrochemical replacement process between the polonium and the silver foil.

3. The presence of undissolved solids in the plating solution, e.g., from digestion of 20 grams or more of bone, reduces the recovery of added polonium appreciably. Filtering off the solids does not increase the per cent recovery even though the filtering process itself does not remove any activity from a filtered solution which initially contained no undissolved solids.

4. Recovery of added inorganic polonium from triplicate samples of acidified human urine when stirred at 720 rpm for 2 hours at 94° C over various foils exceeds 90% for nickel and silver and ranges from 61% to 77% for copper.

5. Recovery of added polonium from one liter of acidified human urine exceeds 90% if the urine is first stirred for 2 hours at 94° C with 2 grams of nickel powder with subsequent filtration, solution of the nickel in aqua regia and a replating on silver foil.

6. Recovery of added polonium from two-liter volumes of 0.5N HCl also exceeds 90% when the fluid is evaporated to 50 ml in a 250 ml beaker using a constant feed method to maintain the fluid level in the beaker during evaporation.

7. The average polonium content of one-liter volumes of pooled urine from presumably non-exposed humans is of the order of 0.05 μ c or less.

8. Fat carefully extracted from tissue digests by use of petroleum ether, with most of the solvent layer being pipetted off, contains less than 1% of the polonium in the digest. Recovery of added polonium from 20 gram samples of lard is 90% or better.

9. Recovery of added polonium is better than 90% from bone, with or without fat extraction, mice, rats, and human lung tissue.

The following specific results were obtained from the experiments performed during the course of this work, using the methods developed therein.

1. Two rats exposed to a known amount of radon and sacrificed immediately after exposure were analyzed for polonium. The analysis indicated activities of 44.3 and 47 dpm/gram.

2. Ten mice exposed to 0.2 mc/liter of radon for 5 hours had lived nearly a normal life span before their death occurred. When analyzed for whole body polonium content, an average activity of 21 dpm per mouse was found.

3. Nine mice exposed to 0.2 mc/liter of radon for 15 hours who died in 90 to 100 days as a result of this exposure were analyzed for polonium content. The analysis indicated that the average whole body content of these mice was 145 dpm.

4. In three of the above mice, the skeleton was analyzed separately by a preliminary treatment with 3% KOH in ethylene glycol which digested the soft tissues, leaving most of the bones. An average of 50% of the whole body polonium content was found in the skeleton compared with 2% for a non-exposed mouse to which polonium had been added.

5. Two bats of the species Corynorhinus Rafinesquei which were trapped in a uranium mine in eastern Utah were analyzed for whole body polonium content. One bat contained 14.66 and the

other 63.63 μc . Since more than one year elapsed between death and analysis, the activity probably represents the radiolead content of these animals.

6. Recovery of polonium from the pooled urine of polonium injected rats was 20% greater after wet digestion and plating than after merely acidifying and plating. Five samples were used in each procedure.

7. The polonium content of various control samples, from presumably unexposed animals, as determined by the methods used in the course of this work is summarized in Table XXVI.

TABLE XXVI

The polonium content of various samples from presumably unexposed animals

Description of sample	Activity 10^{-14} curies/gram
Mice 3-6 months old (7 mice), whole body	0.55
One mouse 2 years old, whole body	1.70
Rats 6 to 8 months old (2 rats), whole body	0.48
Beef femur (62 grams total weight)	1.60
Beef pelvis (113 grams total weight)	0.43
Dog femur (40 grams total weight)	0.40
Human rib (20.8 grams)	0.36
Human sternum (with some attached rib - 205 grams)	3.40
Human lung (155 grams)	0.57

Calculations were made to determine the polonium content to be expected in mice and rats exposed to radon. The calculations gave expected polonium activities close to those measured experimentally though the agreement was perhaps fortuitous due to the uncertainty in the assumptions that were required. Similarly, calculations of the urinary polonium excretion gave a value within the order of magnitude of the activity found in a very limited number of pooled urine samples. Thus these results indicate that it may be possible to determine the relative integrated radon exposure of individuals by analyzing their urine for polonium content.

 

[illegible]

 

BIBLIOGRAPHY

1. Miller, S. E., D. A. Holaday, and H. N. Doyle, Health protection of uranium miners and millers. Arch. Ind. Health, 14: 48-55 (1956).
2. Jacoe, P. W. Occurrence of radon in non-uranium mines in Colorado. Arch. Ind. Health, 8: 118-24 (1953).
3. Harris, S. J., Radon in New York State mines. Arch. Ind. Health, 10: 54 (1954).
4. Evans, R. D. and C. Goodman, Determination of the thoron content of air and its bearing on lung cancer hazards in industry. J. Ind. Hyg. Toxicol., 22: 89-99 (1940).
5. Maximum permissible amounts of radioisotopes in the human body and maximum permissible concentrations in air and water. Handbook 52, National Bureau of Standards, U.S. Dept. of Commerce, 1953.
6. Recommendations of the International Commission on Radiological Protection. Brit. J. Radiol., Supplement No. 6, 1955.
7. Bale, W. F. Memorandum to the files, Div. of Biology and Medicine, A.E.C., 3-14-51.
8. Bale, W. F. Memorandum to the files, Div. of Biology and Medicine, A.E.C., 7-17-51.
9. Shapiro, J. An evaluation of the pulmonary radiation dosage from radon and its daughter products. U of R A.E.P. Report, UR-298, 4-9-54.
10. Cohn, S. H., R. K. Skow, and J. K. Gong. Rn inhalation studies in rats. Arch. Ind. Health, 7: 508-15 (1953).
11. Chamberlain, A.C. and E. D. Dyson. Dose to trachea and bronchi from the decay products of radon and thoron. Brit. J. Radiol., 29: 317 (1956).
12. Morgan, K. Z. Maximum permissible concentration of radon in the air. Unpublished memorandum dtd 10-27-54.
13. Sultzzer, M, and J. B. Hursh. Polonium in the urine of miners exposed to radon. Arch. Ind. Hyg. and Occupational Med., 9: 89-100 (1954).

14. Beamer, W. H. and W. E. Easton, Half-life of Po-210. AECD-3156, Los Alamos Scientific Lab., Los Alamos, N. Mex. (1946).
15. Karraker, D. G., A. Ghiorso, and D. H. Templeton, Alpha decay energies of Po isotopes. UCRL-1091, Univ. of Calif. Radiation Labs., 1-17-51.
16. Beamer, W. H. and C. R. Maxwell, Physical properties and crystal structure of Po. MDDC-721, Los Alamos Sci. Labs., Los Alamos, N. Mex., (1946).
17. Power, W. H., A modified Joliot apparatus for the study of electrodeposition of Po. Mound Lab., Miamisburg, Ohio, Report MLM-909, 10-28-53.
18. Wagner, John J., An investigation of the alpha-ray spectrum of Po. A.E.C. Report No. NP-1642 (1949).
19. Moyer, H. V., ed., Polonium. Mound Laboratories Report TID-5221, Miamisburg, O., 1955.
20. Curie, I. and M. Lecoq, A new gaseous compound of Po. Comptes Rendu, 192: 143-45 (1931).
21. Hoffer, M., Vaporization investigations in vacuum on Po, RaD, and RaE. Sitz. Akad. Wiss. Wien, 144, Abt IIa: 383-86 (1935).
22. Rona, E., Vaporization experiments with Po. Sitz. Akad. Wiss. Wien, 141, Abt IIa: 533-37 (1932).
23. Marchi, L.E., A new method for the electrochemical extraction of Po. Ziv Compt Rend Acad Science USSR, 25: 743 (1939).
24. Prestwood, R. J. Po extraction using a Pt electrode and hydrogen. AECD-2839, Los Alamos Sci. Labs., Los Alamos, N. Mex., (1944).
25. Marckwald, W. Uber das Radiotellur. Ber. deut. chem. Ges., 38: 591 (1905).
26. Scott, R. G. and J. N. Stannard. Modified procedure for analysis of Po²¹⁰ in biological materials. U of R AEP Report, UR-235, 3-31-53.

27. Fink, R. M., ed., Biological studies with polonium, radium and plutonium, National Nuclear Energy Series, McGraw-Hill, New York, 1951.
28. Cember, H., J. A. Watson, and T. B. Grucchi, Procedure for digestion and radioassay of animal tissue. Nucleonics, 12 (8): 40-41 (1954).
29. Mulhaney, T. J., W. P. Norris, and W. E. Kisielleski, Digestion and radioassay. pp. 103-9, Argonne Nat. Lab. Quarterly Rpt., ANL-4333, Chicago, Ill. (1949).
30. Rundo, J., The analysis of biological samples for Po. AERE HP/R-627, Harwell, England, 1950.
31. Smith, F. A., R. J. Della Rosa, and L. J. Casarett, Analytical and autoradiographic methods for polonium-210. U. of R. AEP Report, UR-305, 11-4-55.
32. Robbins, M.C., The determination of polonium in urine. Los Alamos Scientific Lab., Univ. of Calif., Report LA-1904, April, 1955.
33. Whipple, G. Hoyt, Personal communication.
34. Hursh, J. B., The Ra content of public water supplies. U. of R. A.E.P. Report, UR-257, 5-4-53.
35. Hultqvist, B., Studies on naturally occurring ionizing radiations. Kgl. Svenska Vetenskapsakad. Handl., 6: (3) (1956).
36. Scott, K. G., H. Foreman, and J. Crowley, Tracer metabolism studies. UCRL-1282, Univ. of Calif. Radiation Labs., 5-8-51.
27. Mussbaum, E. and J. B. Hursh, Radon solubility in rat tissues. Unpublished material presented at Radiation Research Society meeting, Chicago, Ill., May 17-19, 1956.
38. Hursh, J. B. and A. A. Gates. Body Ra content of individuals with no known occupational exposure. Nucleonics 7 (1): 46-59 (1950).

39. Sievert, R. M., Measurement of low level radioactivity, particularly the gamma radiation, from living subjects. Proc., Intern. Conf. on Peaceful Uses of Atomic Energy, v. 13: 187-95, United Nations, N. Y., 1956.
40. Stehney, A. F., and H. F. Lucas, Jr., Intake and retention of radium at natural levels. Argonne Natl. Lab., Lemont, Ill., Report ANL-5378, 1955.
41. Conference at Leeds -- Measurement of body radioactivity. Nature, 178: 172-75 (1956)
42. Norris, W. P., T. W. Speckman, and P. F. Gustafson, Studies of the metabolism of radium in man. Am. J. Roentg., 73: 785-801 (1955).
43. Stehney, A. F., W. P. Norris, H. F. Lucas, Jr., and W. H. Johnston, A method for measuring the rate of elimination of radon in breath. Am. J. Roentg., 73: 774-84 (1955).
44. Shandley, P. D., The radium content of common foods. U. of R. AEP Report, UR-255, 4-22-53.
45. Stannard, J. N., The distribution and excretion of orally administered Po in the rat. U. of R. AEP Report, UR-299, 9-17-54.
46. Caster, W. O., J. Poncelet, A. B. Simon, and W. D. Armstrong, Tissue weights of the rat. Proc. Soc. Exp. Biol. Med., 91: 122 (1956).
47. Hollcroft, J. W. and E. Lorenz, Retention of radon by the mouse. Nucleonics 2 (3): 63-71 (1949).
48. Crowell, C. D., Jr., H. C. Hodge, and W. R. Line, Removal of organic material from teeth. J. Dental Res., 14: 251 (1954).
49. Stannard, J. N. and R. C. Baxter, The effects of a maintained body burden of polonium in the rat. II. Plan of long term experiments, distribution, excretion, and retention data. U. of R. AEP Report, UR-376, 1-4-55.

50. Thomas, R. G. and J. N. Stannard. The metabolism of Po administered by intratracheal injection in the rat. U. of R. AEP Report, UR-430, 3-20-56.
51. Aub, J. C., R. D. Evans, L. H. Hempelmann and H. S. Martland, The late effects of internally deposited radioactive materials in man. Medicine, 31: 221-329 (1952).
52. Looney, W. B., Late skeletal roentgenographic, histopathological, autoradiographic and radiochemical findings following radium deposition. Am. J. Roentg., 75: 559-72 (1956).

APPENDIX

Decontamination Procedure.

At the start of the research, it was realized that a good decontamination procedure would have to be developed or else new glassware would have to be provided for each sample.

Fink stated that soaking in 3N HCl was an efficient procedure. Also scrubbing with an abrasive powder is a good method.

It was decided that a combination of these two processes with plating should be an acceptable procedure so the following steps were used to decontaminate previously used materials.

1. Foils -- These are merely scrubbed with an abrasive powder (such as Bab-O) until the surface is clean and bright. Counts on these foils range from 0.04 to 0.09 cpm after cleaning.
2. Glassware -- Glassware is washed in warm, soapy water, scouring powder is sprinkled onto the damp surface and then scrubbed thoroughly with a bottle brush. Then rinse under hot tap water while still scrubbing to remove any residue from the glass. Rinse the glassware three times with distilled water, fill with 2N HCl, add 1 or more cleaned silver foils and stir 2 hours or more at 94° C.

This procedure removes polonium by scrubbing, by hot treatment with 2N HCl and by plating the polonium removed by the HCl onto a silver foil.

To check the efficiency, a plating bottle to which had been added polonium which when plated amounted to 6653 cpm was decontaminated. After decontamination the plating bottle was filled with 0.1N HCl over a cleaned Ag foil and the solution stirred 2 hours at 94°C. The count on the foil was 1211 counts/4020 min = 0.301 cpm, net cpm = 0.201. Therefore only $\frac{0.201 \times 100}{6653} = 3.3 \times 10^{-3}\%$ of the original activity was platable. Assuming that the plating before decontamination removed 98% of the activity then 147 cpm should have been left in the plating bottle or solution, therefore $\frac{0.201 \times 100}{147} = 0.13\%$ of this activity was left after decontamination.

As another check, some Ehrlenmeyer flasks and plating bottles had been used with spiked bone sample digests. Approximately 80 cpm of polonium had been added to the bone. After plating the bone digests, the flasks and bottles were decontaminated according to the above method. This glassware was used in the formaldehyde study in Section E.2 above. Table XXIV shows that no detectable activity was added to silver foils when distilled water was put through the nitric-perchloric digestion and plating procedure, the normal procedure for reagent blanks.

ACKNOWLEDGMENTS

This research was suggested by Dr. William F. Bale. The author is grateful to Dr. Bale for his guidance and encouragement during the course of this work and the preparation of this thesis. His insight on many phases of the problem and critical analysis were most valuable.

The author is grateful to Drs. J. Newell Stannard and Frank A. Smith for helpful suggestions and illuminating discussions on the polonium phases of this work. Appreciation is also extended to Dr. John B. Hursh for his many valuable suggestions.

The author is grateful to Dr. Donald A. Morken who made available the radon exposed animals used in part of this work.

The author expresses his appreciation to Dr. James S. Arnold, Radiobiology Laboratory, University of Utah College of Medicine, Salt Lake City, Utah, for furnishing the miner's autopsy samples; to Dr. Mitchell R. Zavon, Division of Special Health Services, Occupational Health Field Headquarters, U.S. Public Health Service, Cincinnati, Ohio, for furnishing the bats; and to Dr. William Hawkins, Strong Memorial Hospital, Rochester, N. Y., for the autopsy material from non-miners.

Ideas and aid were extended by personnel of the Atomic Energy Project too numerous to mention individually, but their efforts are much appreciated nevertheless.