1
DASA 1858

EFFECTS OF OVERPRESSURE ON THE EAR - A REVIEW

Frederic G. Hirsch, M. D.

**Technical Progress Report
on
Contract No. DA-49-146-XZ-372**

**This work, an aspect of investigations dealing with
the Biological Effects of Blast from Bombs, was
supported by the Defense Atomic Support Agency of
the Department of Defense.**

**(Reproduction in whole or in part is permitted for any
purpose of the United States Government.)**

**Lovelace Foundation for Medical Education and Research
Albuquerque, New Mexico**

November 1966

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

FOREWORD

This report, which is primarily a review of published material concerning the effects of blast on the human and the mammalian ear, also incorporates some previously unpublished data concerning the impact of age on the vulnerability of the tympanic membrane to blast overpressures, and some further material on intraspecies scaling.

An attempt has been made to correlate such quantitative data as can be found in the literature, but this has not been found possible of accomplishment in a completely satisfactory way. By rescaling some previously published material in view of available data, and incorporating some unpublished material a value for the threshold of eardrum rupture in man is estimated to be 5 psi, and the "short"-rising overpressure required for rupture of human eardrums is 15 psi.

This information will be of use to those whose task it is to assess hazards and identify populations at risk, as well as those responsible for providing safe working environments and similar problems in the field of environmental health.

ABSTRACT

Information regarding blast effects on the ear has been reviewed in an attempt to gather quantitative information available for animals and man for help in the establishment of relationships between various levels of overpressure and the incidence of eardrum failure, the degree of damage to the middle and inner ear and other identifiable sequelae referable to cochlear or vestibular functions.

ACKNOWLEDGMENTS

The author wishes to acknowledge the considerable assistance in the preparation of this report which was rendered by the following people: to Dr. Donald R. Richmond of the Lovelace Foundation, for permission to include some of his unpublished data; to Mr. Roy H. Reider, of the Los Alamos Scientific Laboratory, for providing data on some recent industrial explosions; to Mr. I. Gerald Bowen, of the Lovelace Foundation, and Mr. Luke J. Vortman of the Sandia Corporation, for their help in pressure-distance scaling calculations; to Dr. Thomas L. Chiffelle, of the Lovelace Foundation, for assistance in preparing and interpreting histological preparations of tympanic membranes; to Dr. Thomas Shipman, of the Los Alamos Scientific Laboratory, and Dr. Donald E. Kilgore Jr., of the Lovelace Clinic, for making clinical material available; to Mr. Robert A. Smith, of the Lovelace Foundation, for the preparation of illustrative material; to Mr. Walter Scheurle, of the Lovelace Foundation, for the translation of material; to Mrs. Laura M. Carrasco and Mrs. Ruth P. Lloyd for secretarial and editorial assistance; to the Defense Atomic Support Agency for contract support; to Colonel Gerrit L. Hekhuis, USAF, MC, Defense Atomic Support Agency, Department of Defense and Lt. Colonel Edmund L. Fountain, USA, VC, Defense Atomic Support Agency, Department of Defense, for their interest and support; and finally to Dr. Clayton S. White, President-Director of the Lovelace Foundation, whose very great personal interest and valuable suggestions contributed very much to whatever merit resides in this report.

EFFECTS OF OVERPRESSURE ON THE EAR - A REVIEW

1. INTRODUCTION

For many years physicians have been aware that the ears of humans and other mammals have an especial vulnerability to injury when exposed to overpressures generated by explosions and the muzzle blast of guns. By the year 1900 there appeared in the medical literature thirteen articles on the subject, beginning with the report of Greene in 1872.¹ The most comprehensive treatment was that of Castex in 1893.² Since that time, well over one hundred papers have been published on blast injury of the ear, their number increasing during periods when a major armed conflict was being waged and diminishing during peaceful times; although, whenever major explosions have occurred, renewed interest has followed.^{3, 4, 5}

A review of that which has been written concerning blast injury of the ear reveals that, with some important exceptions, the reports deal mostly with clinical considerations, and that quantitative data relating overpressures to incidence or degree of injury are lacking. These exceptions date back quite a few years, however, to the work of Zalewski in 1906.⁶ This study, which will be dealt with more fully later on, was for almost forty years the only one of its kind until the work of Zuckerman^{7, 8} and his colleagues during World War II was done. Also about this time, Perlman⁹⁻¹¹ published his studies which have contributed so much to an understanding of the response of the otic structures to blast waves.

In the recent past, the Comparative Environmental Biology Department of the Lovelace Foundation for Medical Education and Research has had an opportunity to obtain some quantitative data on a number of mammalian species during the full-scale weapons tests, its shock tube studies, and some of its other activities.¹²⁻²¹ Recently, studies of the vulnerability of the ear to blast injury have been resumed in England by Golden and Clare.²²

It would appear useful and timely to prepare a review of what is known about otic blast trauma in view of recent advances in explosives technology, the development of new types of ordnance which have higher muzzle velocities and consequently higher muzzle blast overpressures, and for purposes of Civil Defense planning. This communication represents such an undertaking.

2.0 PHYSICAL, ANATOMICAL AND PHYSIOLOGICAL CONSIDERATIONS

The vulnerability of the ear to blast overpressures is readily appreciated when one considers that a blast wave is physically the same phenomenon as a sound wave. It is generated when air molecules are rapidly and violently compressed such as occurs when a solid or liquid substance is suddenly changed into a gas. In an explosion, this change in physical state occurs in times of 1×10^{-5} seconds or less, depending on the chemical nature of the substance. The resultant compression can be of many hundreds or thousands of pounds to the square inch at the source. Near the source, the molecules are forced outward for some distance before commencing to oscillate. Once oscillation has occurred, a wave is propagated radially. As it is propagated through the air, the peak overpressure and its velocity fall off at a rate which is faster than an inverse square of the distance relationship, so that at distances of several hundreds of feet the overpressure will have diminished to a magnitude of several atmospheres. One definition of a shock pulse can be given as a sound wave of great initial condensation and great initial velocity.¹¹ It follows, therefore, that an organ system which is built especially for the reception of sound will receive blast overpressures equally well.

The ear has evolved as an organ system for the transduction of sound waves into nerve impulses which are electrical in nature. It has developed an ability to respond to a limited band of frequencies which for the human may be regarded as lying between 20 Hz and 20,000 Hz. It has developed a high order of sensitivity. It responds to signals which have an energy level as low as 10^{-16} Watts/cm², or whose pressure is about one five billionths of an atmosphere, or whose force causes an excursion of the eardrum - a distance which is less than the diameter of a single hydrogen molecule.^{9-11, 23} The drumhead cannot respond faithfully to pulses which have periods of less than 0.3 milliseconds, but it attempts to do so by making a single large excursion which corresponds to that occasioned by a sound wave of high audible frequency.⁹⁻¹¹ It is this excursion which mediates the trauma to the ear.

The mammalian ear is divided anatomically and physiologically into the external, middle, and inner ears. Each of these plays a different role in the overall function of the auditory apparatus.

The external ear is a sound-gathering device which has two main functions: the first is that of amplification; and the second is as a direction-sensing device. From the standpoint of blast trauma, both functions are affected albeit in different ways.

Golden and Clare²² have found in their studies that, when a shock pulse is generated in the air at a distance from an ear, there is an increase of about 0.2 percent in the overpressure at the site of the tympanic membrane over that which is measured in the air at the same distance from the source of the pulse. FIGURE 1 shows their experimental arrangement and a typical

VARIATION OF PRESSURE IN AIR AND AT DRUMSITE

(After Golden & Clare)

FIGURE 1.

A diagram of the experimental arrangement of Golden and Clare ²² showing the measured pressures in air and at the site of the tympanic membrane.

set of values for the respective overpressures.

When an ear has been significantly injured in such a way that a hearing loss results, there also occurs an interference with stereognosis, which is the ability of an animal to locate the origin of sounds in its environment.¹¹ Part of this ability depends on differential loudness in one ear as opposed to the other, and the rest depends on the reception of sounds at slightly different times in each ear, so that a phase difference between the two ears is occasioned. This phase difference is mediated and interpreted by the nervous system in a complex way, the details of which are not germane to the present consideration. Suffice it to say that when there is injury to an ear, both loudness and phase are affected, since an altered impedance has been introduced into one side of the system.

Before leaving the consideration of the external ear, something should be said about the influence of ceruminous, or wax, plugs on the response of an ear to blast waves. These are not always present in the external auditory canal, but their incidence is quite high. Most often a plug which has a diameter sufficient to occlude the cross section of the canal will not be sufficiently long to fill the length of it, so that an air space exists between it and the eardrum. Less frequently, one end of a plug will be impacted against the drum.

When a ceruminous plug of the first sort is present, it will diminish the gain function of the canal and will act as an attenuator of a blast wave because, being both compressible and displaceable, it absorbs some of the energy of the shock pulse. That plugs of this sort are effective in reducing trauma to the ear by blast has been remarked by several of the authors who have written on the subject.^{9,24-26}

Sometimes, however, the presence of a ceruminous plug will behave in a manner which makes an ear injury worse. If the deep end of the plug is very close to the eardrum, or actually is in contact with it, and if the displacement of it by the shock pulse is sufficient, then the plug behaves as a ramrod causing not only an extensive rupture of the drum but also a displacement of the ear ossicles. Colledge²⁴ has remarked about this, and experience with dogs at the Nevada Test Site was confirmatory.¹²

In the latter case, a plug was placed in one ear of each experimental animal. It was made of cotton and liquid rubber which solidified after exposure to the air. These plugs completely filled the external auditory canals. The relatively "slow"-rising, but very "long"-duration overpressure, measured near the exposure locations inside underground shelters, apparently acted for sufficient time to push the plugs into the middle ear causing obliteration of the eardrums and dislocation of the ossicles.

Experience with exposure of animals to high-explosive detonations

on the one hand, and overpressures generated in shock tubes on the other, has shown that for relatively "low" overpressures there is little displacement of a ceruminous plug when the duration of the overpressure is "short" as in the case of some high-explosive detonations. When the duration of an overpressure of similar magnitude is relatively "long," as in the case of some of the shock tube-generated pulses, the plugs are frequently displaced more deeply into the canal. So whether or not a ceruminous plug in the external auditory canal protects the drum or makes matters worse depends on a variety of things: the duration of the overpressure, whether there is a space between the plug and the eardrum, and the character of the plug; i. e., its density, mass, and composition.

The eardrum, or tympanic membrane, separates the external ear from the middle ear. It is this structure which has engaged most of the attention of the clinicians who have concerned themselves with blast injuries of the ear. This is due to the fact that it is frequently ruptured. It is not, however, the most important lesion associated with blast trauma. Indeed, as will be subsequently developed, it is probably better for the drum to give way to the overpressure than for it to remain intact.

The drum and its associated ossicles act in concert to transfer acoustical energy from the external ear to the inner ear, which is the site of the actual transduction of the mechanical energy to the electrical energy that constitutes a nerve impulse. This system functions as an impedance matching device, and as such contributes an amount of gain to the signals received at the tympanic membrane. It should be emphasized that frequency, wave form, and relative amplitude are not altered by middle ear structures; but that there is a concentration of sound pressure on the oval window.^{23,27,28,29}

The adult human tympanic membrane has an area of about 70 sq. mm., which is some 22 times greater than the area of the oval window. This difference in area causes an increase in sound pressure at the oval window and serves to make up for the relative difference between the inertia of the air and that of a fluid - in this case, the endolymph of the inner ear. It is the motion of the endolymph which actually stimulates the sensing organ in the inner ear.

The natural resonant frequency of the middle ear structures is between 1200 and 1700 Hz; however, the ligaments and muscles attached to the ossicular chain serve to highly dampen the system.

There are two dampers in the middle ear, the most important of which is composed of the stapedius muscle and its ligaments. It is this one which acts to limit the vibration of the stapes when intense signals are being received. The other is the tensor tympani muscle and its ligaments, which serve to limit the vibration of the eardrum. It is of interest to note that, due to the manner of construction of the ligaments, the limitation of amplitude excursion is largely effective when positive pulses

are placed on the eardrum and less on the negative excursions.

These musculo-tendinous dampers are controlled by an involuntary reflex arc. Its stimulus is an excessive amplitude of vibration striking the otic structures which generates feedback. The feedback, in turn, activates the reflex arc in an analogous manner to an electronic automatic volume control circuit. It requires about 5-10 milliseconds for the reflex to become operative and effectual. This is, of course, too long a time for the bracing action of the stapedius and tensor tympani muscles to provide any protection against any but a "slow"-rising, relatively weak blast wave. It does provide some help, however, when a "low"-intensity blast wave assaults the ear under conditions where high ambient noise levels have existed prior to its arrival. Perlman,¹¹ Ireland,³⁰ and other authors have noticed that rupture of the drum is more common when an explosion occurs in a quiet environment than it is in a noisy one. It seems that the middle ear is quite a different target under these two circumstances.

Because of the manner in which the malleus is linked to the incus, inward displacement is greater than outward displacement. The incudo-malleolar joint is so constructed that when the malleus is subjected to an inward force the two bones move forward as one. However, when the excursion of the bones is outward, a separation of the malleus from the incus occurs, thus allowing a greater movement of the malleus vis-a-vis the incus.¹¹

This feature of the incudo-malleolar joint serves as a protective device to the inner ear. When one recalls that a blast wave is characterized by a "sharp"-rising, positive pressure peak of "short" duration which is followed by a "longer," negative phase, it becomes apparent that a longer excursion of the ossicular chain can result from the negative phase of the blast wave than from the positive one, depending on how "fast" an equalizing pressure can enter the middle ear via the eustachian tube. It also follows that the displacement of the endolymph can be of a "longer" duration during the negative phase, thus prolonging the stimulation of the hearing organ. Perlman¹¹ feels that in all probability the length of time that an overload is applied to this organ, the Organ of Corti, is of importance in the degree of damage that ensues. By permitting a longer excursion of the malleus, which is attached to the drum, without requiring the same excursion of the intermediate bone (the incus) the movement of the stapes, which transmits the motion of the endolymph, is diminished, thus affording some degree of protection. He is also of the opinion that if the excursion of the drum and ossicular system to the negative phase could be reduced further or even eliminated, a greater degree of protection to the inner ear would result. It is this sort of thing that occurs when the drum gives way in face of the positive overpressure of a blast wave. The drum having lost its integrity, there results a reduced ability for the conductive system to respond to the negative phase, so that the overload and its duration as applied to the inner ear is less - how much less is dependent on the size of the perforation. It is this that was meant when it was pointed out earlier that rupture of the drum can be regarded as more helpful than catastrophic. However, since the maximum overpressure and its rise time govern the nature of the ensuing negative phase, the former must be regarded as of over-riding importance.

3.0 QUANTITATIVE CONSIDERATIONS OF EARDRUM RUPTURE

There are a number of factors, the interplay of which makes the vulnerability of the tympanic membrane to rupture a variable thing. These may be divided into physical and anatomico-physiological factors.

It has been established that there is a direct relationship between the percentage of ruptured eardrums and the maximum overpressure.³¹ This has also been shown to be the case by the experiences of White et al.¹⁴⁻²⁰ during the several full-scale weapons tests, and by studies on experimental animals subjected to high-explosive detonations and shock tube-generated overpressures.¹⁵

Although extensive systematic studies remain yet to be done, enough data are at hand to show that there are additional parameters which are probably of some importance. These are the duration of the overpressure, its rate of rise, and perhaps the duration and magnitude of the negative phase which follows the positive wave.⁹⁻²⁰

A higher percentage of ruptures will be associated with a given peak pressure if its rise time is "fast," than will be the case when it is relatively "slow." For example, it was found during the full-scale weapons tests in Nevada that 50 percent of dogs' eardrums ruptured when exposed to a "slow"-rising overpressure of 31.2 psi; whereas, a "fast"-rising pulse generated in a shock tube needed only to reach a level of 12 psi in order to produce the same percentage of ruptures.^{15, 20}

In the opinion of Perlman,⁹ if the negative pressure which follows after the overpressure of a blast wave is relatively large and prolonged, a drum which has not been ruptured by the positive pressure may be ruptured during the negative phase, if its outward excursion is sufficiently large.

The vulnerability of an eardrum varies also with its age and whether or not it has been subjected to previous trauma or disease.^{6, 32}

Many years ago Zalewski⁶ found that when a static overpressure was applied to the tympanic membranes of fresh cadavers, the pressure required to cause rupture varied over a range of from 5.4 to 44.1 psi; however, that there was an age dependence was shown by the fact that whereas an average of 33 psi was required to produce ruptures in cadavers from the first decade of life, the figure was only 20 psi for the older age groups.

The reason for this variation of vulnerability with age has not been carefully studied, but the assumption has been that it can be explained by regarding the young membrane as having more elasticity than an older one.^{9, 32} Some current studies are under way in the Lovelace Foundation Department of Pathology in which histological preparations are being made of tympanic membranes removed from both quite young and quite old

animals and humans who come to autopsy. So far, only a few such preparations have been made and studied, but a very definite impression has emerged that the number and organization of the elastic fibers in the submucosal connective tissue in the material obtained from the young drum is quite different when compared with the old ones. In the young drum they are more abundant and organized; whereas, in the old drums they are more sparse and less well organized. It will require the evaluation of much more material than has yet become available, however, before a definite statement can be made.

The thickness and elasticity of a tympanic membrane can also be influenced by the aftereffects of previous disease or injury. Anything which leaves a scar in its wake will, of course, lessen the drum's resistance to blast. Conversely, those pathologic conditions, usually of a chronic nature, which result in a thickened membrane, will cause the drum to become more resistant according to Perlman.¹⁰

It has been known since Zalewski's work⁶ work was published in 1905 and from that of Blake et al.⁸ that the eardrums of the various mammalian species manifest differing degrees of vulnerability to rupture by overpressures. Zalewski found that in the case of the dog, the range of pressures required for rupture was between 9.1 and 22.8 psi with a mean of 14.9 psi. This was in contrast with the values obtained for the human cadavers he studied where the range was between 5.4 and 44.1 psi with a mean value for all ages of 22.9 psi.

An opportunity for interspecies comparison of this point has been afforded over the years to the Comparative Environmental Biology Department of the Lovelace Foundation and, although a comprehensive, systematic study has not yet been made, there are enough data at hand to make a few statements possible.

Exposure of young dogs to nuclear blast at the Nevada Test Site indicated that the maximal pressures for rupture of the eardrum ranged from 4.1 to 85.8 psi, with 31.2 psi being the statistically determined pressure required for 50 percent failure of the membrane.^{15,20} The highest pressure without rupture was noted to be 66.6 psi. The differences between these values and those of Zalewski⁶ possibly can be accounted for by recalling that his experiments were performed using a static overloading of the drum; whereas, the overpressures in the nuclear tests were the result of the detonation of an explosive.

FIGURE 2 illustrates what has been found in regard to species differences. Included are some data on monkeys derived from the studies of Blake, Douglas, Krohn and Zuckerman⁸ and on swine from Richmond.³³ It appears that the ear of the monkey is the most blast resistant of the species studied, and that the ear of the goat is the most vulnerable.

It is not possible at this time to attribute the difference in vulnerability to any single feature of the ears of these different species, such as

A log-normal plot of overpressure versus percentage of tympanic membrane ruptures for various species. It can be seen that the 50-percent value is least for the goat (8.5 psi) and greatest for the monkey (22 psi).

the structure of the tympanic membrane, or the anatomy of the external ear, since no intensive comparative studies have been made. That there are wide variations in the anatomy of the ear among the mammals is obvious. Except for differences in the thickness of the stratified columnar epithelial layer, there has not been found any marked variation in the histology of the tympanic membranes from a number of species which have been examined. The middle connective tissue layer seems to be the same for all of the species examined. It is probable that some of the biophysical parameters, such as resonant frequency, elasticity, etc., might well show that the drums of the various species of mammals are very different in their dynamics, and they are different targets more because of this, than because of differences in histologic structure.

The mammal of chief interest is, of course, the human species; consequently, it is desirable to know where it fits into the scheme of things. Reference to the work of Zalewski has already been made.⁶ This is the most systematic study which has been made, as well as the first. His experiments dealt with the resistance of cadaver ears to "slowly" rising overpressures. These he created by pumping air into the external auditory canal which had been sealed to prevent air escaping from the external auditory meatus. This is different than that of a "fast"-rising overpressure, such as a blast wave. He found that 11 percent of the cadaver eardrums ruptured at an overpressure which was less than 1 atmosphere (14.7 psi). At overpressures between 1 and 2 atmospheres 66 percent ruptured, and at pressures greater than 2 atmospheres the remaining 23 percent were ruptured.

Blake, Douglas, Krohn, and Zuckerman⁸ also studied the problem using cadavers. Instead of using a "slowly"-rising overpressure, they generated a shock wave by the bursting-diaphragm technique such as is used on a larger scale in some shock tubes. They found that the pressure required to rupture 50 percent of the drums was about 1 atmosphere or 14.7 psi.

They also studied the incidence of ruptured tympanic membranes amongst air raid casualties. They found that the lowest pressure resulting in rupture of the eardrum was between 2 and 4 psi. In their sample, however, 37 percent of people exposed to blast pressures in excess of 100 psi showed no damage to their tympanic membranes. They concluded that the lower limit of pressure required to rupture 50 percent of eardrums was 15 psi and that the upper limit was 50 psi.

Zuckerman⁷ in a later report concluded that over a range from 6 psi to 70 psi eardrum rupture may occur, but that the likelihood of such an event was much greater at the higher levels than at the lower.

Data such as we have just regarded leave something to be desired in the way of precision to those who have an interest in assessing casualty potentials. These people like to have numbers to use, such as the threshold pressure below which damage is not to be expected and a pressure value at

which 50 percent of an exposed population can be expected to manifest the effect. It was regarded as a useful and interesting exercise to see if such numbers could not be culled from the literature. Accordingly, a literature search was undertaken with the result that a number of germane publications were found. Most of these were reports of the clinical experiences with blast injury to the ear, and no effort was made to equate overpressures with either the incidence or extent of damage. In some, however, reference was made to a particular type of ordnance, a particular type and weight of explosive, or a measured overpressure. By equating these things with other available data dealing with explosions, scaling values for various explosives, and the known characteristics of various types of ordnance, it was possible to make some approximations which when assembled seemed orderly and reasonable. In doing this the help of people knowledgeable in the field of explosion phenomenology, such as Mr. I. G. Bowen of the Lovelace Foundation for Medical Education and Research, Albuquerque, New Mexico; Mr. L. J. Vortman of Sandia Corporation, Albuquerque, New Mexico, and Mr. R. H. Reider of the Los Alamos Scientific Laboratory, Los Alamos, New Mexico, was needed. They all gave substantial assistance which is gratefully acknowledged.

The difficulties and uncertainties surrounding an approach of this kind were discussed by Zuckerman in 1941.⁷ The peak overpressure and its scaling produced by any given explosive or type of ordnance can be very well determined for any given set of conditions, but since it is very difficult to determine exactly which set of conditions apply to a given case, some estimation is required. So, recognizing the inherent deficiencies of this sort of an approach, the following data are presented.

In the matter of the least overpressure which can be associated with eardrum rupture, Zalewski⁶ found this number to be 5.4 psi. Perlman¹⁰ found that a pressure of 3.9 psi was the lowest that produced rupture in his studies. Machle,³⁴ on the other hand, studied forty-five gunnery instructors who had frequent exposures to "fast"-rising overpressures of 4 psi and encountered failure of a drum at 6 psi. Shilling³⁵ found no case of rupture of the tympanic membrane to pressures less than 7 psi, but his subjects were subjected to a "slow"-rising overpressure. Corey³⁶ found 7 psi to be the rupture threshold in his study. It appears from these data that a "fast"-rising pressure pulse of about 5 psi is sufficient to rupture some human tympanic membranes and that this value can be regarded as the threshold.

When it comes to estimating the pressure required to rupture 50 percent of human drums, one is unable to find any report in the literature presenting data which permit making a categorical statement. However, there are some reports which do permit some estimation. Henry³⁷ on 292 men who sustained blast injuries to their ears. In these there were 152 perforations (or 52 percent). All these patients were injured when they were about 50 feet from an explosion of a land mine, 500-lb. aerial bomb, or a 3-in. mortar shell. Calculations indicate that the overpressure to which these people were subjected was probably about 17 psi.³⁸ Vadala³⁹ reports on his experience with 75 men who sustained ear injury as a result of exposure to the muzzle blast of a 37-mm. antiaircraft gun. From the studies

of Murray and Reid⁴⁰ this gun has a muzzle blast overpressure of about 6 psi. In Vadala's material he encountered 6 cases of tympanic membrane rupture, or an incidence of 8 percent. Reider⁴¹ at Los Alamos has some data from two recent industrial explosions which indicate that 85 percent of 22 exposed drums ruptured in one explosion where the peak overpressure was 30 psi. In the other explosion, where the overpressure reached 40 psi, all of the 6 exposed eardrums were ruptured.

If one plots these percentages versus their associated overpressure as a log-normal plot, one obtains a straight line best fit which is like those that have been obtained on experimental animals.¹⁴⁻²⁰ FIGURE 3 shows this plot compared with one prepared by White et al. for dogs exposed to "fast"-rising overpressures.¹⁵ It would appear from this that the probable "fast"-rising overpressure required for rupture of 50 percent of human drums is about 15 psi.

FIGURE 3.

A log-normal plot of the data on humans as derived from the references as shown. This is compared with the data on dogs as presented by White in CEX 65.4.¹⁵

4.0 HEARING LOSS CAUSED BY BLAST TRAUMA

One important result of blast injury to the ear is the amount of damage to the Organ of Corti in the inner ear caused by an excessive motion of the endolymph. Such damage can result in varying degrees of temporary or permanent deafness. That the hearing loss is related to whether or not the eardrum has ruptured has been remarked about in many of the reports which have been published in the literature. For example, Bourgeois⁴² states categorically that when the eardrum ruptures the injury to the inner ear is less, and the consequent deafness is less grave in that it is less apt to be permanent. Concurrence with this opinion has been expressed by Henry,³⁷ Colledge,²⁴ Silcox and Schenck,³¹ and others.

When a blast wave has caused the eardrum to rupture, the associated hearing loss will be of a mixed type; that is to say, there will be a loss of acuity for low tones as well as for high tones. Ordinarily the low-tone loss will be of the order of 10 to 30 db; whereas, the high-tone loss will be more in the nature of 40 to 80 db. This will be true providing there has been no dislocation of the ossicles associated with the rupture, because when this happens, there results a permanent and severe conductive deafness. From clinical reports it is apparently unusual for the ossicles to be displaced by a "fast"-rising, "short" duration overpressure unless the overpressure is very high. Ireland³⁰ found no incidence of ossicular fracture or dislocation in 317 cases of blast injury of the ear all due to high-explosive detonations; however, Barrow and Rhoads⁴³ encountered several in their study of 32 cases of blast injury of the ear. Henry³⁷ remarks in his study of 292 cases that dislocation of the ossicles was uncommon.

Henry also pointed out that he was unable to correlate the intensity of the immediate hearing loss with the size of the perforation, and other authors are found to be in agreement. Among the published reports studied, no one was found who disagreed with this opinion.

Boemer⁴⁴ in his study of 310 cases of acoustic trauma found that 78 percent of the cases of temporary deafness had not suffered a perforation of the drum and makes the point that these were the more severely deafened.

A number of studies of traumatic deafness, such as those of Murray and Reid,⁴⁰ Machle,³⁴ and Bunch³² have emphasized that there is an accumulative effect on hearing when multiple acoustical insults are sustained by an individual. It is also apparent that the time interval between incidents is of importance in that when they occur with intervals shorter than the recovery period, the permanent hearing loss is more profound. Murray and Reid,⁴⁰ Silcox and Schenck,³¹ and Perlman⁹ have all pointed out that there is a marked individual variation in susceptibility to acoustic trauma. It appears that susceptibility to inner ear damage as evidenced by high-tone hearing loss and the time required for recovery from

an acoustic insult go hand in hand; that is to say, the more susceptible inner ears require longer for recovery.⁴⁰

There are three types of hearing loss due to blast injury. The first of these is a neurosensory type which involves audition of high tones, usually considered as those above 4000 Hz. This can be either temporary or permanent and can result whether or not there has been incurred any damage to the middle ear structures. If there has been only a single traumatic episode, or if the intervals between episodes have been long enough to permit recovery to the inner ear structures, then the high-tone loss is temporary. On the other hand if multiple traumata of the inner ear are experienced without adequate recovery intervals, the loss will be permanent.

A case in point is one taken from the comprehensive article on traumatic deafness written by C. C. Bunch³² in 1937. It concerns a naval gunnery officer who had, in the course of his duties, experienced periods of time when his ears were exposed to muzzle blast overpressures from naval ordnance. He never suffered rupture of his eardrums so far as is known. The audiogram depicted in FIGURE 4 was taken on him at the age of 36 years. It shows a loss of auditory acuity beginning at 200 Hz and is most profound at 4000 Hz. This type of audiogram is regarded by otologists as characteristic of cochlear damage and also of senile degeneration of the inner ear. When one regards the age of this patient, it hardly seems likely that the hearing loss can be on the latter basis.

The second type of deafness associated with blast injury is that which is obtained when a rupture of the eardrum has occurred. Dr. D. E. Kilgore of the Lovelace Clinic has made available data on a 44-year old physicist who was involved in a hydrogen-oxygen explosion and as a result was exposed to a "fast"-rising overpressure estimated by Reider⁴¹ to be near 30 psi. He suffered bilateral tympanic membrane ruptures which did not become infected and healed completely within several months. FIGURE 5 is an audiogram showing the auditory acuity for his right ear. The record for his left was virtually identical. The solid line is post-rupture, and the broken line is the record made following healing of the perforations. The mixed type of deafness is clearly apparent, as is the degree of recovery.

The third type of hearing loss is that which results when the middle ear structures are irreversibly damaged. This is illustrated by a case, also taken from Bunch,³² concerning a soldier who was in a trench in World War I when a shell exploded less than thirty feet away from him. He suffered obliteration of his eardrums and dislocation of the middle ear ossicles as well. His audiogram, which is shown in FIGURE 6, was taken some twenty years after his injury which had occurred at the age of 25 years. The depressed acuity for all frequencies is characteristic for conductive deafness, and the fact that there has been no restoration in twenty years emphasizes the permanent nature of his disability.

FIGURE 4.

The audiogram of a 36-year old Naval Gunnery Officer who had suffered repeated noise exposures, but had never had a rupture of an eardrum. The high-tone loss cannot be attributed to aging since he is yet a young man. Solid lines represent the right ear; dotted lines represent the left ear.

FIGURE 5.

The audiogram of a 44-year old physicist with bilateral eardrum ruptures. The solid line is post-rupture. The dotted line is following healing of the perforation. Data for right ear only.

FIGURE 6.

Frequency in Hz

The audiogram of a soldier who received severe blast injury to his ears at the age of 25. This record was obtained when he was 41 years old, some sixteen years after his injury. The blast injury obliterated the structures of his middle ears. Solid lines represent the right ear; dotted lines represent the left.

5.0 PREVENTION OF BLAST INJURY OF THE EAR

A search for a satisfactory method of preventing acoustic trauma has been conducted by many people for many years. The approach used by most of them has been to recommend the use of some type of ear plug which will attenuate a blast wave (or loud sound) and dissipate some of its energy.

All of the devices which have been designed will do these things and thus give their wearers substantial degrees of protection.^{40, 44-46} They all have associated with their use, however, some degree of loss of auditory acuity which can be quite disadvantageous under certain circumstances.

As with all protective devices, such as "hard hats," safety shoes, safety glasses, and others, resistance to their use is perversely encountered in some of the population at risk.

Some otologists have regarded a simple cotton plug, constituting a protective device, as satisfactory as a more elaborate ear plug. Others, and they seem to be in the majority, decry the use of cotton plugs and point out with some merit the reasons for their objections.^{45, 46} After reviewing a substantial amount of the pertinent literature, one is forced to conclude that the proper and faithful use of an ear defender in a traumatically noisy environment, or where a real hazard of blast injury exists, is wise and worthwhile.^{32, 34, 40, 47}

The placement of gun crew members should certainly be made with the threat of muzzle blast in mind. Wherever the muzzle blast over-pressure approaches 4 psi, the crew members should be protected by ear defenders.

6.0 DISCUSSION

From what has been said, it is easy to appreciate why the ear is especially vulnerable to blast overpressures. Indeed, it is remarkable that a structure possessed of delicacy and sensitivity such as the mammalian ear can present as much resistance to damage by blast as it does. This must be due in part to the mitigative effect that rupture of the tympanic membrane frequently enters into the picture; although as has been pointed out, there are other important individual variables such as age, the previous health history of the ears, and the poorly understood differences in the susceptibility of individuals to acoustic trauma.

An entirely different set of variables are those associated with any given explosion - such as the maximum overpressure, the nature of its rise time, and its duration. To these must be added all of those things which collectively can be called "geometric positioning."

Where an individual happens to be when an explosion occurs, how he is oriented with respect to the direction of propagation of the blast wave, and whether environmental conditions are such as to afford shielding or reflections of the overpressure will cause marked differences in the magnitude of exposure in different individuals who may be quite close to one another at the time.

For all of these reasons, it is very difficult to derive quantitative data about the vulnerability of human ears from the study of what has occurred to an exposed population. Further, insofar as experimental studies of inner ear trauma are concerned, it has not been possible to equate overpressure versus damage due to the extraordinarily formidable difficulties associated with investigations of inner ear function. This has been pointed out by Wever and Bray⁴⁸ and more recently by Vartanyan and Maruseva.⁴⁹ The latter also emphasize that the frequency sensitivity range of the ear of an animal are highly species specific and that the mechanism of cochlear function and neural organization in the brain vary. For these reasons, extrapolations of data derived from one species to another are precarious.

Insofar as the middle ear structures are concerned, it can probably be safely said that the threshold pressure for damage to middle ear structures is about 5 psi and that overpressure near 15 psi will cause the rupture of 50 percent of the eardrums exposed to them. However, except when the ossicles have been displaced, damage to the middle ear is of secondary importance as compared with inner ear trauma, especially insofar as long-term and permanent disability are concerned. Thus, these numbers lose some of their significance and utility insofar as they can be applied helpfully to casualty assessment and identifying conditions of hazard. One can assume that obvious middle ear trauma is always associated with some significant degree of inner ear damage, but the converse is not a valid assumption.³² It has been shown by many, although perhaps no better than by Bunch,³² that inner ear damage can very often occur in the absence of demonstrable injury to the middle ear.

Compared with the other types of blast trauma, such as that to the lung and to those organs affected by the presence of air emboli in their vascular elements, injury to the ears is distinctly a matter of secondary importance. Even though the body of information concerning otic blast trauma may leave something to be desired concerning some of its quantitative details, enough can be discerned from reading that which has been published on the subject to enable a fairly satisfactory understanding of what happens and how it happens. Enough certainly is at hand to appropriately guide those who are concerned with environmental health and safety insofar as ears are concerned.

7.0 SUMMARY

The literature concerning blast injury of the ear has been reviewed. In general, there is apparently little in the way of quantitative data to be found. An effort has been made to arrive at some quantitative estimates relating blast overpressure to ear injury. It appears that the threshold for rupture of the human eardrum by a "fast"-rising overpressure is about 5 psi and that the overpressure required for rupture of 50 percent of eardrums is near 15 psi. It has not been found possible to equate overpressure quantitatively with damage to the inner ear. Some of the anatomical and physiological factors involved in otic trauma due to blast have been discussed, and the utility of ear defenders has been considered.

REFERENCES

1. Greene, J. O., "Cases of Injury to the Ear from External Violence," Trans. Amer. Otol. Soc. 5: 88, 1872.
2. Castex, A., "Effects of Dynamite on the Ear," J. Laryng. 7: 307, 1893.
3. Rusca, F., "Experimentelle Untersuchungen ueber die traumatische Druckwirkung der Explosionen," Zeitschr. Chir. 132: 315-374, 1915.
4. McReynolds, G. S., F. R. Guilford and G. R. Chase, "Blast Injuries to the Ear," Arch. Otolaryng. 50: 1-8, 1949.
5. Hofer, I. and O. Mauthner, Verletzungen des Ohres bei katastrophalen Explosionen, Joseph Sofar, Wien, 1913. (Reviewed by A. Tweedie in J. Laryngol. 28: 618, 1913.)
6. Zalewski, T., "Experimentelle Untersuchungen ueber die Resistenzfaehigkeit des Trommelfells," Zeitschr. f. Ohren. 52: 109-128, 1906.
7. Zuckerman, S., "Rupture of the Ear Drums by Blast," Ministry of Home Security Report B.P.C. 148/W.S. 13, Oxford, England, 1941.
8. Blake, P. M., J. W. B. Douglas, P. L. Krohn and S. Zuckerman, "Rupture of Eardrums by Blast," Medical Research Council Report B.P.C. 43/179/W.S. 21, Oxford, England (undated).
9. Perlman, H. B., "Acoustic Trauma in Man," Arch. Otolaryng. 34: 429-452, 1941.
10. Perlman, H. B., "Reaction of the Human Conduction Mechanism to Blast," Laryngoscope 55: 437-442, 1945.
11. Perlman, H. B., "The Effect of Explosions on the Acoustic Apparatus," Trans. Amer. Acad. Ophthal. Otolaryng. 47: 442-453, 1942.
12. Roberts, J. E., C. S. White and T. L. Chiffelle, "Effects of Overpressure in Group Shelters on Animals and Dummies," USAEC Civil Effects Test Group Report, WT-798, Office of Technical Services, Department of Commerce, Washington, D. C., September 1953.

13. Richmond, D. R., R. V. Taborelli, F. Sherping, M. B. Wetherbe, R. T. Sanchez, V. C. Goldizen and C. S. White, "Shock Tube Studies of the Effects of Sharp-Rising, Long-Duration Overpressures on Biological Systems," USAEC Technical Report, TID-6056, Office of Technical Services, Department of Commerce, Washington, D. C., March 10, 1959.
14. Richmond, D. R., R. V. Taborelli, I. G. Bowen, T. L. Chiffelle, F. G. Hirsch, B. B. Longwell, J. G. Riley, C. S. White, F. Sherping, V. C. Goldizen, J. D. Ward, M. B. Wetherbe, V. R. Clare, M. L. Kuhn and R. T. Sanchez, "Blast Biology — A Study of the Primary and Tertiary Effects of Blast in Open Underground Protective Shelters," USAEC Civil Effects Test Group Report, WT-1467, Office of Technical Services, Department of Commerce, Washington, D. C., June 30, 1959.
15. White, C. S., I. G. Bowen and D. R. Richmond, "Biological Tolerance to Air Blast and Related Biomedical Criteria," USAEC Civil Effects Test Operations Report, CEX-65.4, Office of Technical Services, Department of Commerce, Washington, D. C., October 18, 1965.
16. White, C. S. and D. R. Richmond, "Blast Biology," USAEC Technical Report, TID-5764, Office of Technical Services, Department of Commerce, Washington, D. C., September 18, 1959.
17. White, C. S., "Biological Blast Effects," USAEC Technical Report, TID-5564, Office of Technical Services, Department of Commerce, Washington, D. C., September 1959.
18. White, C. S., I. G. Bowen, D. R. Richmond and R. L. Corsbie, "Comparative Nuclear Effects of Biomedical Interest," USAEC Civil Effects Test Operations Report, CEX-58.8, Office of Technical Services, Department of Commerce, Washington, D. C., January 12, 1961.
19. White, C. S., "Biological Effects of Blast," Technical Progress Report, DASA-1271, Defense Atomic Support Agency, Department of Defense, Washington, D. C., December 1961.
20. White, C. S., T. L. Chiffelle, D. R. Richmond, W. H. Lockyear, I. G. Bowen, V. C. Goldizen, H. W. Merideth, D. E. Kilgore, B. B. Longwell, J. T. Parker, F. Sherping and M. E. Cribb, "Biological Effects of Pressure Phenomena Occurring Inside Protective Shelters Following Nuclear Detonation," USAEC Civil Effects Test Group Report, WT-1179, Office of Technical Services, Department of Commerce, Washington, D. C., October 28, 1957.
21. Pratt, D. E., T. L. Chiffelle, E. G. Damon and D. R. Richmond, "Threshold Lung Injury in Goats from a 500-Ton Explosion (Program 4 — Project 4.4)," Tab "D," Biomedical Program — 500-Ton Explosion — Final Report, Technical Progress Report, DASA-1656, Defense Atomic Support Agency, Department of Defense, Washington, D. C., June 1, 1965.

22. Golden, P. M. and R. Clare, "The Hazards to the Human Ear from Shock Waves Produced by High Energy Electrical Discharge," UK Atomic Energy Authority Weapons Group, Atomic Weapons Research Establishment, Report AWRE-E-1, Aldermaston, Berkshire, England, August 1965.
23. Bekesy, G., "Ueber die mechanischen-akustischen Vorgaenge beim Hoeren," Acta Otolaryng. 27: 281-296, 1939.
24. Colledge, L., "Discussion on Injuries of the Ear," J. Laryng. 55: 549-570, 1940.
25. Eldredge, D. H., "The Effects of Blast Phenomena on Man," CHABA Report No. 3, Armed Forces - National Research Council, Committee on Hearing and Bio-Acoustics, Washington, D. C., June 1955.
26. Wilson, J. G., "The Ear in Modern Warfare," Brit. Med. J. 1: 589, 1917.
27. Steudel, V., "Ueber Empfindung und Messung der Lautstaerke," Hochfrequenztechnik. Electroakustik 41: 116-128, 1933.
28. Guelke, R. and J. A. Keen, "A Study of the Movements of the Auditory Ossicles under Stroboscopic Illumination," J. Physiol. 116: 175-188, 1952.
29. Kessel, J., "Ueber das Durschschneiden des Steigbuegelmuskels beim Menschen," Arch. Ohrenh. 11: 100, 1876.
30. Ireland, P. E., "Traumatic Perforations of Tympanic Membrane Due to Blast Injury," Canad. Med. Assn. J. 54: 256-258, 1946.
31. Silcox, L. E. and H. P. Schenck, "Blast Injury of the Ears," Arch. Otolaryng. 39: 413-420, 1944.
32. Bunch, C. C., "Nerve Deafness of Known Pathology and Etiology," Laryngoscope 47: 615-691, 1937.
33. Richmond, D. R., Lovelace Foundation for Medical Education and Research, Albuquerque, New Mexico, Unpublished Data.
34. Machle, W., "Effects of Gun Blast on Hearing," Arch. Otolaryng. 42: 164-168, 1945.
35. Shilling, C. W., "Aerotitis Media," Arch. Otolaryng. 42: 169-173, 1945.
36. Corey, E. L., "Medical Aspects of Blast," U. S. Nav. Med. Bull. 46: 623-652, 1946.

37. Henry, G. A., "Blast Injuries of the Ear," Laryngoscope 55: 663-672, 1945.
38. Vortman, L. J., Sandia Corporation, Albuquerque, New Mexico, Personal Communication.
39. Vadala, A. J., "Effects of Gun Explosions on the Ear and Hearing Mechanism," Milit. Surg. 66: 810-822, 1930.
40. Murray, N. E. and G. Reid, "Temporary Deafness Due to Gun-fire," Med. J. Aust. 1: 611-617, 1946.
41. Reider, R. H., Los Alamos Scientific Laboratory, Los Alamos, New Mexico, personal communication concerning data from two industrial explosions.
42. Bourgeois, H., "L'Organe de l'audition pendant le guerre," Arch. de Med. et de Pharm. Mil. 65: 822, 1917.
43. Barrows, D. W. and H. T. Rhoads, "Blast Concussion Injuries," J.A.M.A. 125: 900-902, 1944.
44. Boemer, L. C., "Acoustic Trauma," Ann. Otol. 54: 513-517, 1945.
45. Cox, L. K. and J. W. Geller, "Anticoncussion Earplugs; Prevention of Aural Complications from Gunnery Concussion," U. S. Nav. Med. Bull. 43: 139-142, 1944.
46. Wells, W. A., "Danger to Hearing Apparatus in Modern Warfare," Trans. Amer. Laryng. Rhinol. Otol. Soc. 47: 215-223, 1941.
47. Craig, D. H., "Injury to Tympanic Membrane Caused by Explosions," Lancet 2: 40-41, 1940.
48. Wever, E. G. and C. W. Bray, "Auditory Nerve Impulses," Science 71: 215, 1930.
49. Vartanyan, I. A. and A. M. Maruseva, "Cochlear Electrical Reactions to Clicks in Rats," Fiziol. Zh. SSSR imeni I. M. Sechenov 51: 1037. Fed. Proc. Trans. Suppl. 25: T543, 1966.

ADDITIONAL REFERENCES

- Alexander, A. B., "Rupture of the Drumhead as a Wartime Injury," Brit. Med. J. 2:195-197, 1941.
- Barr, T., "Enquiry into the Effects of Loud Sounds upon the Hearing of Boilermakers and Others who Work Amid Noisy Surroundings," Proc. Glasgow Philosophical Soc. 17:223, 1886.
- Barr, T., "Injurious Effects of Loud Sounds upon the Hearing," Brit. Med. J. 2:675, 1890.
- Beck, K., "Lärmarbeit und Ohr," Ztschr. f. Hals-, Nasen- u. Ohrenh. 18:387-391, 1927-1928.
- Blake, C. J., "Influence of the Use of the Telephone upon Hearing Power," Arch. Otol. 17:240, 1888.
- Blegvad, V. R., "The Influence of their Calling upon Telephone Operators Particularly with Regard to the Hearing," J. Laryng. 24:172, 1909.
- Boot, G. W., "Caisson Worker's Deafness," Ann. Otol. 22:1121, 1913.
- Braislin, W. C., "Two Cases of Injury to the Ear by Lightning," Brooklyn Med. J. 20:133, 1906.
- Bruhl, G., "Contribution to the Pathology of the Ear," Arch. Otol. 36:91, 1907.
- Bruner, G., "An Interesting Case of Protracted Deafness Caused by the Report of a Musket," Arch. Otol. 9:333, 1880.
- Bryant, W. S., "Prevalence of Ear Injuries and Diseases in the French Army," J. Laryng. 32:338, 1917.
- Cantor, J. J., "Rupture of Tympanic Membrane Due to Blast," Ann. Otol. 54:554-567, 1945.
- Capart, A., "Diseases and Accidents of Telephone Operators," Laryngoscope 22:656, 1912.
- Coates, G. M., M. V. Miller, "War Injuries of Ear, Nose and Throat," Med. Clin. N. Amer. 25:1653-1676, 1941.
- Coggin, D., "A Curious Case of Deafness," Boston Med. Surg. J. 116:9, 1887.
- Colledge, L., "Injuries to the Ear in Modern Warfare," J. Laryng. 36:283, 1921.

- Collins, E. G., "Aural Trauma Caused by Gunfire," J. Laryng. 62:358-390, 1948.
- Connal, J. G., "Rupture of Membrani Tympani with a Clonic Spasm of the Facial Muscles in a Diver," Brit. Med. J. 2:719, 1898.
- Cott, G., "Telephone Deafness," Buffalo Med. J. 73:317, 1917-18.
- Crowe, S. J., S. R. Guild, L. M. Polvgt, "Observations on the Pathology of High Tone Deafness," Bull. Johns Hopkins Hosp. 54:315-379, 1934.
- Culler, E., G. Finch, "Effect upon Cochlear Function of Intense Tonal Stimulation," Am. J. Physiol. 113:32, 1935.
- Davis, H., A. J. Derbyshire, E. H. Kemp, M. H. Lurie and M. Upton, "Experimental Simulation of Deafness," Science 81:101, 1935.
- Dickie, J. K. M., "Experiences of an Otologist in France," J. Laryng. 36:514, 1921.
- Dore, G., "La sordita del cacciatori," Valsalva 9:27-30, 1933.
- Drooker, J. C., "Wartime Experience with Skull Injury and Blast and Their Effect upon the Ear," Arch. Otolaryng. 58:546-574, 1953.
- Duane, A., "Middle and Internal Ear Disease Produced by the Concussion due to Gunfiring," Arch. Otol. 29:62, 1900.
- Egeblad, K., "Electrical Injury to the Organ of Hearing," Psychol. Abs. 6:474, 1932.
- Faulder, T. J., "Injuries to the Ear in Modern Warfare," J. Laryng. 36:277, 1921.
- Fitz-Hugh, G. S., "Traumatic Perforation of Ear Drum," Virginia Med. Monthly 73:504-505, 1946.
- Fromm, B., O. Nylen and Y. Zotterman, "Electric Stimulation of the Cochlea," J. Physiol. 80:3, 1933-34.
- Got, A., "War Deafness with Lesions of the Internal Ear," J. Laryng. 31:374, 1916.
- Grant, D., "Varieties of Nerve Deafness," J. Laryng. 17:169, 1902.
- Grove, W. E., "Otologic Observations in Trauma of the Head," Arch. Otolaryng. 8:249, 1928.

- Guild, S. R., "Otopathies of War, with Especial Reference to Air Waves," Int. Assn. Med. Mus. Bull. 7:225, 1918.
- Guild, S. R., "War Deafness and Its Prevention," Ann. Otol. 50:70-76, 1941.
- Habermann, J., "Hardness of Hearing in Boilermakers," Arch. Otol. 20:171, 1891.
- Heiman, T., "A Blow on the Ear Followed by Death in a Week," Arch. Otol. 20:17, 1891.
- Hendricks, J. E., "War Deafness in Gunnery Instructors," Ann. Otol. St. Louis 55:68-80, 1946.
- Hipskind, M. M., "War Injuries to the Auditory Apparatus, Clinical and Audiometric Observations," Laryngoscope 55:678-705, 1945.
- Holt, E. E., "Boiler Makers Deafness and Hearing in a Noisy Environment," Trans. Amer. Otol. Soc. 3:34, 1882.
- Hoople, G. D., "Otolaryngology in World War II," Ann. Otol. 55:121-127, 1946.
- Horne, J., "Gun Deafness and Its Prevention," Lancet 2:462, 1914.
- Horton, G. P., "The Effect of Prolonged Acoustic Stimulation on the Auditory Sensitivity of Guinea Pigs," J. Comp. Psychol. 18:405, 1934.
- Ingersoll, J. M., "The Otological Work in the U. S. A. General Hospital for Head Surgery," Trans. Amer. Otol. Soc. 15:68, 1919.
- Jobson, T. B., "Normal Gun Deafness," Lancet 2:566, 1917.
- Jurgens, E., "Affections of the Ear, Nose and Throat Following the Explosion of Bombs," La Presse Otolaryng. Pt. 5, 1907. Abstracted in Arch. Otol. 37:305, 1908.
- Kingery, C. N., and B. F. Pannell, "Peak Overpressure vs. Scaled Distance from TNT Bursts," Ballistics Research Lab., Aberdeen Proving Ground, Maryland BRL-MR-1518, April 1964.
- Korkis, F. B., "Blast Injuries of the Ear," J. Laryng. 66:95-103, 1952.
- Lewis, R., "Deafness Due to Concussion from Gun Firing," Arch. Otol. 29:63, 1900.

- Loeb, H. W., "Military Surgery of the Ear, Nose and Throat," Med. War Man. 8:31, 1918. Lea and Febiger, Philadelphia.
- Lindsay, J. R., H. G. Kobrack and H. B. Perlman, "Relation of the Stapedius Reflex to Hearing Sensation in Man," Arch. Otolaryng. 23:671-678, 1936.
- McMaster, G. T., "Gun Deafness - A Danger," Am. Med. 4:417, 1909.
- Mueller, R., "Effect of Artillery Practice on the Ears," Arch. Otol. 28:264, 1889.
- Nimier, "Effects Produced upon the Ear by the Report of Firearms," Arch. Otol. 19:58, 1890.
- Packard, F. R., "Traumatic Perforations of the Membrana Tympani," J. A. M. A. 34:1160, 1900.
- Pratt, D. E., T. L. Chiffelle, E. G. Damon and D. R. Richmond, "Symposium Proceedings: Operation Snowball," Defense Atomic Support Agency Data Center Special Report 34. Published by DASA Data Center, General Electric Co. TEMPO 735 State Street, Santa Barbara, California.
- Reudi, L. and H. Minder, "Acute Acoustic Trauma," Pract. Oto-rhinolaryng. 5:169, 1943.
- Richards, H., "Two Noteworthy Cases of Traumatic Rupture of the Membrana Tympani," Trans. Amer. Otol. Soc. 4:447, 1889.
- Ridout, G. B., "Gunfire Deafness in the Navy," U. S. Nav. Med. Bull. 28:736, 1930.
- Sexton, S., "Injury of the Ear Caused by the Blast of a Bursting Shell," N. Y. Med. Rec. p. 201, Feb. 19, 1887.
- Schubert, H., "Ueber die Widerstandsfahigkeit des Trommelfells gegen Ueberdruck der Aussenluft," Z. Hals, Nasen, u. Ohren. 1:357-358, 1949. Exerpta Med. Sec. II 3:945, 1950.
- Shuter, R. E., "Notes on War Injuries of the Ear," Med. J. Aust. 1917. Reviewed in J. Laryng. 33:255, 1918.
- Simmons, M. H., "Otological Experiences During the Battle of Santiago on Board the Battleship Iowa," Laryngoscope 6:277, 1899.
- Stuhlman, O. J., "The Nonlinear Transmission Characteristics of the Auditory Ossicles," J. Acoustic Soc. Am. 9:911-921, 1937.

Wever, E. G. and W. W. Bray, "Auditory Nerve Impulses," Science 71: 215, 1930.

Wilson, J. G., "Effects of High Explosives on the Ear," J. A. M. A. 71:628, 1918.

"The Ear in Modern Warfare," Brit. Med. J. 1:589, 1917. Discussion of the report by J. G. Wilson, "The Effects of High Explosives on the Ear," Brit. Med. J. 1:353-355, 1917.

DISTRIBUTION

ARMY AGENCIES

Chief of Research and Development, Life Sciences Division, Dept. of the Army, ATTN: CRDLS, Washington, D. C. 20310 (1 copy)

Chief of Research and Development, Dept. of the Army, ATTN: CRDNCB, Washington, D. C. 20310 (1 copy)

Assistant Chief of Staff for Intelligence, Dept. of the Army, ATTN: DFISS, Washington, D. C. 20310 (1 copy)

Chief of Engineers, Department of the Army, ATTN: ENGTE-E, Washington, D. C. 20315 (1 copy)

Directorate of Transportation, U. S. Army Materiel Cmd., Washington, D. C. 20315 (1 copy)

Commanding General, Medical Research and Development Cmd., Dept. of the Army, ATTN: MEDDH-N, Main Navy Building, Washington, D. C. 20360 (2 copies)

Commanding General, Medical Research and Development Cmd., Dept. of the Army, ATTN: MEDPS-CS, Main Navy Bldg., Washington, D. C. 20360 (2 copies)

Commanding General, Medical Research and Development Command, Dept. of Army, ATTN: MEDDH-RS, Main Navy Bldg., Washington, D. C. 20360 (2 copies)

U. S. Army CDC Artillery Agency, Fort Sill, Oklahoma 73503 (1 copy)

President, Aviation Test Board, U. S. Army Aviation Center, Fort Rucker, Alabama 36362 (1 copy)

Assistant Chief of Staff for Force Development, Dept. of the Army, ATTN: Directorate of CBR and Nuclear Operations, Washington, D. C. 20310 (1 copy)

Commandant, U. S. Army C&GS College, ATTN: Archives, Fort Leavenworth, Kansas 66027 (1 copy)

Commandant, U. S. Army Air Defense School, ATTN: Command & Staff Dept., Fort Bliss, Texas 79906 (1 copy)

Commanding Officer, U. S. Army CDC Armor Agency, Fort Knox, Kentucky 40120 (1 copy)

Commanding Officer, U. S. Army CDC Artillery Agency, Fort Sill, Oklahoma 73503 (1 copy)

Commanding Officer, U. S. Army CDC Infantry Agency, Fort Benning, Georgia 31905 (1 copy)

Commanding Officer, U. S. Army CDC CBR Agency, Fort McClellan, Alabama 36205 (1 copy)

Commandant, U. S. Army CBR, Weapons School, Dugway Proving Ground, Dugway, Utah 84022 (1 copy)

Commanding General, Army Medical Service School, Brooke Army Medical Center, ATTN: DCCO, Fort Sam Houston, Texas 78234 (1 copy)

Commanding General, The Engineer Center, ATTN: Asst Cmdt Engineer School, Fort Belvoir, Virginia 22060 (1 copy)

Commanding General, 9th Hospital Center, ATTN: CO, U. S. Army Nuclear Medicine Research Det., Europe, APO New York 09180 (1 copy)

Commanding Officer, U. S. Army Chemical Research and Development Lab., Edgewood Arsenal, Maryland 21040 (1 copy)

Commandant, Walter Reed Army Institute of Research, Walter Reed Army Medical Center, Washington, D. C. 20012 (5 copies)

Commanding General, U. S. Army Natick Laboratories, ATTN: Technical Library, Natick, Massachusetts 01760 (1 copy)

Commanding Officer, U. S. Army, Combat Development Cmd. , Institute of Nuclear
 Studies, Fort Bliss, Texas 79906 (1 copy)
 Commanding General, U. S. Army Combat Development Cmd. , ATTN: CDC CD-F, Fort
 Belvoir, Virginia 22060 (2 copies)
 Commanding General, U. S. Army, Combat Development Command, ATTN: Combat
 Support Group, Fort Belvoir, Va 22060 (1 copy)
 Commanding Officer, U. S. Army, Combat Development Cmd, Medical Service Agency,
 Fort Sam Houston, Texas 78234 (1 copy)
 Commanding General, U. S. Army, Combat Development Cmd, Combat Arms Group, Fort
 Leavenworth, Kansas 66027 (1 copy)
 Director Armed Forces Institute of Pathology, Walter Reed Army Medical Center,
 625 16th Street, N.W., Washington, D.C. 20012 (1 copy)
 Director, Waterways Experiment Station, P.O. Box 631, ATTN: Library, Vicksburg,
 Mississippi 39180 (1 copy)
 Director, U. S. Army Ballistic Research Laboratories, Aberdeen Proving Ground, Mary-
 land 21005 (2 copies)
 Redstone Scientific Info Center, U. S. Army Missile Cmd, ATTN: Chief, Document
 Section, Redstone Arsenal, Alabama 35808 (1 copy)
 Commanding General, Army Materiel Command, ATTN: AMCRD-BN, Washington, D.C.
 20315 (1 copy)
 Commanding Officer, U. S. Army Nuclear Defense Laboratory, ATTN: Librarian,
 Edgewood Arsenal, Maryland 21040 (1 copy)
 Commanding General, Army Materiel Command, ATTN: EOD Division, Washington,
 D.C. 20315 (1 copy)
 Office of the Secretary of the Army, Assistant Director of OCD, ATTN: Research,
 Washington, D.C. 20310 (5 copies)

NAVY AGENCIES

Chief of Naval Operations, Navy Department, ATTN: OPO3EG, Washington, D.C. 20350
 (1 copy)
 Chief of Naval Operations, Navy Department, ATTN: OP-75, Washington, D.C. 20350
 (1 copy)
 Chief of Naval Operations, Navy Department, ATTN: OP-922G2, Washington, D.C.
 20350 (1 copy)
 Chief of Naval Personnel, Tech Library, Code 11B, Navy Department, Washington, D.C.
 20370 (1 copy)
 Chief of Naval Research, Navy Department, ATTN: Code 811, Washington, D.C. 20390
 (1 copy)
 Chief, Bureau of Medicine and Surgery, Navy Department, ATTN: Code 74, Washington,
 D.C. 20390 (2 copies)
 Chief, Bureau of Medicine and Surgery, Navy Department, ATTN: Code 71, Washington,
 D.C. 20390 (1 copy)
 Chief, NAVSHIPS, Navy Department, ATTN: Code 423, Washington, D.C. 20390 (1 copy)
 Chief, NAVSHIPS, Navy Department, ATTN: Code 364, Washington, D.C. 20390 (1 copy)
 Director, U. S. Naval Research Laboratory, Washington, D.C. 20390 (1 copy)
 Commander, U. S. Naval Ordnance Laboratory, White Oak, Maryland 20910 (1 copy)
 CO, U. S. Naval Mine Defense Laboratory, Panama City, Florida 32401 (1 copy)
 CO, U. S. Naval Unit, U. S. Army Chemical School, Fort McClellan, Alabama 36205
 (1 copy)

Nuclear Medicine Division, School of Submarine Medicine, U. S. Naval Submarine Medical Center, Box 600, U. S. Naval Submarine Base, New London, Groton, Connecticut 06340 (1 copy)
 CO, U. S. Naval Hospital, ATTN: Director, REEL, NNMCMC, Bethesda, Md. 20014 (1 copy)
 CO and Director, U. S. NRDL, ATTN: Tech Info Div, San Francisco, California 94135 (4 copies)
 CO, Nuclear Weapons Training Center, Atlantic, U. S. Naval Base, ATTN: Nuclear Warfare Department, Norfolk, Virginia 23511 (1 copy)
 CO, Nuclear Weapons Training Center, Pacific, U. S. Naval Air Station, North Island, San Diego, California 92135 (1 copy)
 CO, U. S. Naval Damage Control Training Center, Naval Base, ATTN: NBC Defense Course, Philadelphia, Pennsylvania 19112 (1 copy)
 Superintendent, U. S. Naval Postgraduate School, Monterey, California 93940 (1 copy)
 U. S. Naval Schools Command, T. I., NBC Defense Dept., Box 104, Bldg 194, ATTN: Tech Library, San Francisco, California 94130 (1 copy)
 Commanding Officer, U. S. Naval Air Development Center, ATTN: NAS, Librarian, Johnsville, Pennsylvania 18974 (1 copy)
 Officer in Charge, U. S. Naval Supply Research and Development Facility, Naval Supply Center, Bayonne, New Jersey 07002 (1 copy)
 Commanding Officer, U. S. Naval Medical Research Institute, National Naval Medical Center, Technical Reference Library, Bethesda, Maryland 20014 (2 copies)
 Commandant, U. S. Marine Corps, ATTN: Code A03H, Washington, D. C. 20380 (1 copy)
 U. S. Naval School of Aviation Medicine, U. S. Naval Aviation Medical Center, ATTN: Director of Research, Pensacola, Florida 32508 (1 copy)
 Commanding Officer, U. S. Naval Weapons Laboratory, Dahlgren, Virginia 22448 (1 copy)
 David Taylor Model Basin, Code 258, Washington, D. C. 20007 (2 copies)
 Commander, Naval Facilities Engineering Command, Department of the Navy, Washington, D. C. 20390 (1 copy)
 Commander, U. S. Naval Applied Science Laboratory (Code 900), Naval Base, Brooklyn, New York 11251 (1 copy)
 U. S. Naval Ordnance Test Station, Code 4563, China Lake, California 93555 (2 copies)

AIR FORCE AGENCIES

HQ USAF (1 copy)
 HQ USAF (AFMSPA) T-8, Wash DC 20333 (1 copy)
 HQ USAF (AFMSR), T-8, Wash DC 20333 (1 copy)
 HQ USAF, Chief of Operations and Analysis, Wash DC 20330 (1 copy)
 HQ USAF (AFRSTA), Wash DC 20330 (1 copy)
 HQ USAF (AFTAC), 6801 Telegraph Road, Alexandria, Va. 22310 (1 copy)
 HQ USAF (AFRDD), Wash DC 20330 (1 copy)

AIR FORCE MAJOR COMMANDS

AFSC (SCS-7), Andrews AFB, Wash DC 20331 (1 copy)
 AFSC (SCB), Andrews AFB, Wash DC 20331 (1 copy)
 AFSC (SCTR), Andrews AFB, Wash DC 20331 (1 copy)
 ADC, Asst for Atomic Energy, ADLDC, Ent AFB, Colorado 80912 (1 copy)
 AUL, Maxwell AFB, Alabama 36112 (1 copy)
 SAC (OA), Offutt AFB, Nebraska 68113 (1 copy)
 SAC (SUP), Offutt AFB, Nebraska 68113 (1 copy)

TAC (OA), Langley AFB, Virginia 23365 (1 copy)
AU, Office of the Surgeon, Maxwell AFB, Alabama 36112 (1 copy)
ATC, Office of the Surgeon, Randolph AFB, Texas 78148 (1 copy)

AFSC ORGANIZATIONS

AFSC Scientific and Tech Liaison Office, RTD, ATTN: AFUPO, Los Angeles, California 90045 (1 copy)
HQ, RTD (RTTW), Bolling AFB, Wash DC 20332 (1 copy)
HQ, FTD (TDFBD), Wright-Patterson AFB, Ohio 45433 (1 copy)
HQ, BSD (BSR-1), Norton AFB, California 92409 (1 copy)
HQ, ASD (ASMM), Wright-Patterson AFB, Ohio 45433 (1 copy)
HQ, RTD(SEG/SENS), Wright-Patterson AFB, Ohio 45433 (1 copy)
AFAPL(APS), Wright-Patterson AFB, Ohio 45433 (1 copy)
AFAL(AVS), Wright-Patterson AFB, Ohio 45433 (1 copy)
AFML(MAS), Wright-Patterson AFB, Ohio 45433 (1 copy)
HQ SSD(SSTR), AFUPO 45, Los Angeles, California 90045 (1 copy)
RADC(EMEAM), Griffiss AFB, NY 13440 (1 copy)
AFWL(WLR), Kirtland AFB, New Mexico 87117 (1 copy)
AFRPL(RPPP), Edwards AFB, California 93523 (1 copy)
HQ ESD (ESLE), L. G. Hanscom Fld, Bedford, Mass. 01731 (1 copy)
HQ AMD (AMR), Brooks AFB Tex 78235 (1 copy)
HQ AMD (AMRB), Brooks AFB Tex 78235 (1 copy)
HQ AMD (AMRO), Brooks AFB Tex 78235 (1 copy)
AFSWC (SWLA), Kirtland AFB NMex 87117 (1 copy)
AFWL (WLIL), Kirtland AFB NMex 87117 (1 copy)
AFWL (WLRB), Kirtland AFB NMex 87117 (2 copies)
USAF SAM (SMB), Brooks AFB Tex 78235 (1 copy)
6570 AMRL, Wright-Patterson AFB, Ohio 45433 (2 copies)
6571 ARL, Holloman AFB NMex 88330 (1 copy)

OTHER AIR FORCE AGENCIES

AFCRL, ATTN: CRQST-2, L. G. Hanscom Fld, Bedford, Mass. 01731 (1 copy)
Director, USAF Project Rand, AF Liaison Officer, Rand Corp (Library), 1700 Main St., Santa Monica, California 90401 (1 copy)
Medical Service School (Rad Branch), Sheppard AFB Tex 76311 (1 copy)
HQ OAR (RROS), 1400 Wilson Blvd, Arlington, Virginia 22209 (1 copy)
AFOSR (SRGL), 1400 Wilson Blvd, Arlington, Virginia 22209 (1 copy)
AFIT (Tech Library), Wright-Patterson AFB, Ohio 45433 (1 copy)
Air Force Western Test Range (WTOSM), Vandenberg AFB, California 93437 (1 copy)

DEPARTMENT OF DEFENSE AGENCIES

DDC, Cameron Station, Alexandria, Virginia 22314 (20 copies)
Director of Defense Research and Engineering, ATTN: Tech Library, Washington, D.C. 20301 (1 copy)
Director of Defense Research and Engineering, ATTN: Chief, Medical Services Division, Office of Science, Washington, D.C. 20301 (1 copy)
Director, Defense Atomic Support Agency, ATTN: Document Library, Washington, D.C. 20301 (3 copies)

Director, Defense Atomic Support Agency, ATTN: STMD (For TTCP), Washington, D. C. 20301 (12 copies)
 Commander, Test Command, Defense Atomic Support Agency, Sandia Base, Albuquerque, New Mexico 87115 (1 copy)
 Commander, Field Command, Defense Atomic Support Agency, Sandia Base, ATTN: Surgeon, Albuquerque, New Mexico 87115 (1 copy)
 Commander, Field Command, Defense Atomic Support Agency, ATTN: FCTG, Albuquerque, New Mexico 87115 (1 copy)
 Director, Armed Forces Radiobiology Research Institute, National Naval Medical Center, Bethesda, Maryland 20014 (2 copies)
 Director, Weapons Systems Evaluation Group, Washington, D. C. 20305 (1 copy)
 U. S. Documents Officer, Office of the United States National Military Representative-SHAPE, APO, New York, New York 09055 (1 copy)
 Director, Armed Forces Institute of Pathology, Walter Reed Army Medical Center, Washington, D. C. 20012 (1 copy)
 Director, ARPA, DOD, ATTN: Tech Info Officer, the Pentagon, Washington, D. C. 20301 (1 copy)
 Director, DIA, ATTN: DIAAP-IKZ, Washington, D. C. 20301 (1 copy)

OTHER FEDERAL AGENCIES

Atomic Energy Commission, ATTN: Asst. Director for Medicine & Health Research, DBM, Washington, D. C. 20545 (1 copy)
 Atomic Energy Commission, Civil Effects Branch, Div of Biology and Medicine, Washington, D. C. 20545 (150 copies)
 Atomic Energy Commission, Div of Technical Information Extension, P. O. Box 62, Oak Ridge, Tenn. 37830 (300 copies)
 Director, National Aeronautics & Space Admin., Washington, D. C. 20546 (1 copy)
 National Aeronautics and Space Admin., ATTN: Director, Life Sciences Program, Washington, D. C. 20546 (1 copy)
 National Library of Medicine, Accessions Branch, 8600 Wisconsin Ave., Bethesda, Maryland 20014 (1 copy)
 Federal Air Surgeon, (AM-1) HQ Federal Aviation Agency, 800 Independence Ave., S. W., Washington, D. C. 20553 (1 copy)
 Public Health Service, 4th and Jefferson Drive, S. W., Washington, D. C. 20201 (1 copy)
 Public Health Service, Southwest Radiological Health Lab., ATTN: Tech Library, P. O. Box 684, Las Vegas, Nevada 89101 (1 copy)
 USPHS, Research Branch, Division of Radiological Health, 1901 Chapman Av., Rockville, Md. 20853 (1 copy)
 USPHS, S. E. Radiobiology Health Laboratory, PO Box 61, Montgomery, Alabama 36101 (1 copy)
 Lawrence Radiation Lab., U. of Calif., P. O. Box 808, ATTN: Information Integration Group, Livermore, California 94550 (1 copy)
 Argonne National Laboratory, 9700 South Cass Ave., ATTN: Medical Div., Argonne, Illinois 60440 (1 copy)
 Batelle N. W. Laboratory, P. O. Box 999, ATTN: Dr. Bair, Rickland, Washington 99352 (1 copy)
 Brookhaven National Laboratory, Tech Information Div., Documents Group, Upton, New York 14100 (1 copy)
 Los Alamos Scientific Laboratory, ATTN: Biomedical Research Group, P. O. Box 1663, Los Alamos, New Mexico 87544 (1 copy)

Lawrence Radiation Laboratory, U. of California, P. O. Box 808, ATTN: Biomedical Research Div., Livermore, California 94550 (1 copy)

NON FEDERAL AGENCIES

National Academy of Sciences, ATTN: Richard Park, 2101 Constitution Ave., Washington, D. C. 20418 (5 copies)

Eric H. Wang Civil Engineering Research Facility, Box 188, University Station, U. of New Mexico, Albuquerque, New Mexico 87106 (1 copy)

Southwest Research Institute, ATTN: Technical Library, 8500 Culebra Road, San Antonio, Texas 78228 (1 copy)

Medical College of So. Carolina, Dept of Surgery, ATTN: Dr. M. S. Rittenbury, 55 Doughty St., Charleston, So. Carolina 29403 (1 copy)

Cincinnati General Hospital, Radioisotope Lab., ATTN: Dr. E. L. Saenger, Cincinnati, Ohio 45229 (1 copy)

New York State Veterinary College, Cornell U., Dept. of Physical Biology, ATTN: Dr. Comar, Ithaca, New York 14850 (1 copy)

Lovelace Foundation for Medical Education & Research, 5200 Gibson Blvd., S. E., ATTN: Dr. C. S. White, Albuquerque, New Mexico 87108 (50 copies)

The Rand Corporation, 1700 Main St., Santa Monica, Calif. 90401 (1 copy)

U. of Rochester, School of Medicine & Dentistry, Dept of Radiation Biology, P. O. Box 287, Station 3, Rochester, New York 14620 (1 copy)

U. of Rochester, P. O. Box 287, Station 3, ATTN: Atomic Energy Project Library, Rochester, New York 14620 (1 copy)

U. of New Mexico, ATTN: Library, Albuquerque, New Mexico 87106 (1 copy)

U. of Pennsylvania Medical School, Medical Library, ATTN: Librarian for Blast Biology, Philadelphia, Penn. 19104 (1 copy)

San Francisco State College, 1600 Holloway Ave., ATTN: Dr. Curtis Newcombe, San Francisco, Calif 94132 (1 copy)

Baylor University, ATTN: Document Library, 5th and Speight, Waco, Texas 76703 (1 copy)

University of California, ATTN: Director Radiobiology Lab., Davis, California 95616 (1 copy)

Laboratory of Nuclear Medicine & Radiation Biology School of Medicine, U. of Calif., ATTN: Library, 900 Veteran Ave., Los Angeles, Calif 90024 (1 copy)

Colorado State University, ATTN: Dir Collaborative Radiological Health Laboratory, Ft. Collins, Colo 80521 (1 copy)

Mass Institute of Technology, ATTN: MIT Library, Cambridge, Mass 02139 (1 copy)

Polaroid Corp, ATTN: Tech Library, 119 Windsor St., Cambridge, Mass 02139 (1 copy)

Medical College of Virginia, ATTN: Dept of Biophysics, 523 N. 12th St., Richmond, Virginia 23219 (1 copy)

Sandia Corporation, P. O. Box 5800, ATTN: Dir of Research, Albuquerque, New Mexico 87115 (2 copies)

Security Classification

DOCUMENT CONTROL DATA - R&D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Lovelace Foundation for Medical Education and Research Albuquerque, New Mexico 87108		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED	
		2b. GROUP	
3. REPORT TITLE "Effects of Overpressure on the Ear - A Review"			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Progress			
5. AUTHOR(S) (Last name, first name, initial) Hirsch, Frederic G., M.D.			
6. REPORT DATE November 1966		7a. TOTAL NO. OF PAGES 35	7b. NO. OF REFS 49
8a. CONTRACT OR GRANT NO. DA-49-146-XZ-372		9a. ORIGINATOR'S REPORT NUMBER(S) DASA-1858	
b. PROJECT NO. A4a 03.012			
c.			
d.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
10. AVAILABILITY/LIMITATION NOTICES "Distribution of this report is Unlimited."			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY DASA	
13. ABSTRACT Information regarding blast effects on the ear has been reviewed in an attempt to gather quantitative information available for animals and man for help in the establishment of relationships between various levels of overpressure and the incidence of eardrum failure, the degree of damage to the middle and inner ear and other identifiable sequelae referable to cochlear or vestibular functions.			

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Blast Effects Ear Injury - Blast Overpressure and Eardrum failure Human ear injury Ear Injury Criteria						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year; or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.