UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

PHYSICS DIVISION QUARTERLY REPORT

May, June, July 1955

BERKELEY, CALIFORNIA

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

1
UNIVERSITY OF CALIFORNIA
Radiation Laboratory
Berkeley, California
Contract No. W-7405-eng-48

PHYSICS DIVISION QUARTERLY REPORT

May, June, July 1955

August 24, 1955

PHYSICS DIVISION QUARTERLY REPORT

May, June, July 1955

Contents

GENERAL PHYSICS RESEARCH

Physics Research: Alvarez

Progress on the $C^{14} + p \rightarrow B^9 + \alpha$ Reaction at 32 Mev	4
Elastic Scattering of 30-Mev Protons	4
K^+ Lifetime Experiment	4
"Delayed Fragments"	5
Polarization of Deuterons in the Reaction $p + p \rightarrow \pi + d$	5
Direction of Polarization of the 184-inch Cyclotron Beam	5
Liquid Hydrogen Bubble Chambers	5

Physics Research: Barkas

K-Meson Mass Differences	8
Range-Momentum Relation	8
Experiments with the Neutron Beam from the Bevatron	9
Nuclear Track Photometer	9
Angle and Energy Distributions of Charged Particles from the High-Energy Nuclear Bombardment of Various Elements	10
Interactions of 5.7-Bev Protons in G.5 Emulsion	10

Theoretical Group: Judd 11

Physics Research: Lofgren

Elastic Proton-Proton Scattering	16
Antiproton Search	16
$C^{12}(p, pn) C^{11}$ Cross Section at Bevatron Energies	18
τ -Meson Lifetime	18
Interactions of Negative K Particles at Rest	18
K-Particle Production Studies	20

Physics Research: Moyer

Neutron Multiplicities from μ^- Capture	23
Nuclear Gamma-Ray Spectroscopy	23
Proton-Proton π^0 Production	24
Beta-Ray Spectroscopy	25
Photonic Decay of Heavy Mesons and Hyperons	25
Total-Absorption Photon Spectrometer	25
Pion-Proton Cross Sections	26
Proton Compton Scattering	26

Cloud Chamber Studies: Powell

Experiments

4.5-Bev π^- -p Interactions	28
High-Energy n-p Interactions	28
5.3-Bev p-p Interactions	28
Search for the Antiproton	29

* Preceding Quarterly Reports: UCRL-3014, UCRL-2920

Equipment and Techniques

Machine Analysis	29
New Cloud Chamber Magnet	29
Hydrocarbon Bubble Chamber	30

Physics Research: Richman

Masses of the K^+ Mesons	31
τ^+ Analysis	32
Large Emulsion Stack	32
Negative Proton Experiment	32
Direct Production of Electron Pairs by High-Energy Electrons	32
Reports Issued	33

Spiral-Orbit Spectrometer: Sagane

Physics Research: Segrè

Polarization Experiments on p-p Scattering	36
Calculations on p-p Scattering Phase Shifts	36
Proton-Neutron Triple Scattering	36
Search for Negative Protons	36
Interactions of Negative K Particles in Flight	37
Interactions of Positive K Particles in Flight	37
The Mean Life of Positive K Mesons	38

ACCELERATOR OPERATION AND DEVELOPMENT

184-Inch Cyclotron	39
60-Inch Cyclotron	39
Linear Accelerator	46
Synchrotron	46

PHYSICS DIVISION QUARTERLY REPORT

May, June, July 1955

Radiation Laboratory
University of California
Berkeley, California

August 24, 1955

GENERAL PHYSICS RESEARCHPHYSICS RESEARCH

Luis W. Alvarez in Charge

PROGRESS ON THE $C^{12} + p \rightarrow B^9 + \alpha$ REACTION AT 32 Mev

Harrold B. Knowles

During May through July there were several runs at the linear accelerator at 30° in the laboratory system. The ground state of B^9 comes in very clearly and is calculated to have a differential cross section at this angle of about 1.3 millibarns/steradian. The excited level at 2.4 Mev was probably observed, and indications of one other level were obtained. A month was devoted to obtaining a differential range foil of about 0.75 mg/cm^2 of aluminum. The differential range and target thicknesses are now about the same. This was done in order to optimize the counting rate and resolution. The last run in July yielded no results because of electronic failure.

ELASTIC SCATTERING OF 30-Mev PROTONS

John Leahy

The differential cross sections for Al, Cu, Ag, and Au have been fitted with the optical model by Dr. Saxon at University of California, Los Angeles, using a complex nuclear potential, $V + iW$, with an exponential tail. The values giving the best fit are $V = 35 \text{ Mev}$, $W = 15 \text{ Mev}$, $r_0 = 1.33$, and $a = 0.50$ (both r_0 and a are in units of 10^{-13} cm), where $R = r_0 A^{1/3}$ and a is the rounding parameter for the potential well. This value of W corresponds to a mean free path in nuclear matter of $2.4 \times 10^{-13} \text{ cm}$.

 K^+ LIFETIME EXPERIMENT

Luis W. Alvarez, Frank S. Crawford, Myron L. Good, and M. Lynn Stevenson

During this period we had a successful Bevatron run on a K^+ meson lifetime experiment. We detected and identified K mesons by type, using counter techniques for the first time. Data reduction for this run is proceeding. Preliminary measurements of the mean lives of the $K_{\mu 2}$ and $K_{\pi 2}$ mesons will be obtained upon completion of the data reduction. The best uncorrected value is $1.3 \times 10^{-8} \text{ sec}$ for both types.

Experimental difficulties prevented our obtaining in this run a measurement of the τ lifetime. These difficulties are not of a fundamental nature.

Another run is being planned, and it is expected that the mean lives of the $K_{\mu 2}$, $K_{\pi 2}$, and τ will be obtained to good accuracy.

"DELAYED FIGMENTS"

Luis W. Alvarez, Frank S. Crawford, Myron L. Good, and M. Lynn Stevenson

Several cosmic ray cloud chamber photographs have been seen elsewhere apparently showing a heavy meson decaying at rest with a mean life of the order of milliseconds. A counter experiment was performed at the Bevatron to search for such "delayed figments." The result was negative; we can say that at an incident proton energy of 4.8 Bev there is less than one "delayed figment" per 200 K^+ particles at a 90° laboratory angle. The counters were gated about 5 milliseconds after the main beam struck the target. Thus if the "figments" are sufficiently short-lived, they could have been missed. A more conclusive result could be obtained at 6.1 Bev at a more forward angle. This experiment may be undertaken if time permits.

POLARIZATION OF DEUTERONS IN THE REACTION $p + p \rightarrow \pi + d$

Robert D. Tripp

The polarization of deuterons produced at a center-of-mass angle of 115° in the $p + p \rightarrow \pi + d$ reaction, using 340-Mev unpolarized protons, has been measured. The deuterons were detected in coincidence with the mesons and were analyzed by scattering from a carbon target at 24° . The observed polarization was $0.043 \pm .040$. In conjunction with previous experiments on the reaction, this experiment completes the specification of the phenomenological parameters of Gell-Mann and Watson. Specifically it permits the determination of the relative phases of the several modes of meson production which in turn are related to the p-p phase shifts at this energy. An article is being submitted to the Physical Review.

DIRECTION OF POLARIZATION OF THE 184-INCH CYCLOTRON BEAM

Hugh Bradner

Scanning has been resumed on the experiment to determine the direction of polarization of the cyclotron proton beam via scattering at low energy in helium. Approximately 200 real scatters and 20 background events have been found to date. The expected magnitude of polarization has been recomputed from the published phase shifts of p-He scattering experiments.

LIQUID HYDROGEN BUBBLE CHAMBERS

Luis W. Alvarez and J.D. Gow, and group

General

Development work has continued on the four-inch and ten-inch chambers. In order to expedite completion of the ten-inch instrument, the engineering effort on the large chamber has been temporarily reduced. A reevaluation

R. L. Garwin, private communication of results from the M. I. T. Cloud Chamber group.

Alvarez

of the proposed use of the large chamber resulted in a slight change in size. It is now planned to be 72 inches in length, 20 inches in width and 15 inches in depth. The personnel from NBS Boulder cryogenic laboratory have continued to aid our efforts on all phases of development.

A program to determine experimentally the consequences of sudden release of liquid hydrogen into a vacuum system has been initiated, and experiments are under way. These experiments will mock up the situation that might exist in case of window failure in the 10-inch chamber. They will also give data predicting requirements for safety vent systems suitable for other liquid hydrogen devices.

The window seal program has continued, with efforts pointed toward solution of the 10-inch window problem. Three different designs based on lead gaskets have been tested successfully. A low-temperature-solder metal-glass seal is under investigation. Some material evaluation work is also being done, particularly on thermal expansions, and magnetic and tensile properties of materials at low temperature.

The group from Boulder has been concentrating on the thermodynamic problems associated with bubble chamber operation. Calculations of heat input associated with the pressure-volume cycle, including the effect of compressibility of liquid hydrogen, have been made. Comparison of various expansion schemes from a thermodynamic standpoint is under way. Preliminary results show that a vapor system is adequate for the 4-inch chamber and will work with rather large heat inputs for the 10-inch chamber. For larger chambers a low-temperature expansion system is required. The compressibility of liquid hydrogen in the range of 6 to 1 atmospheres is about 0.5 % per atmosphere, is nonlinear with pressure, and is a function of temperature.

4-Inch Status

The four-inch chamber was successfully operated with pulsed Helmholtz coils, which produced a peak magnetic field of 15 kilogauss through the chamber. The thin stainless steel chamber proved difficult to operate owing to leakage at the window seal. A four-day run was made, using the UCRL 300-Mev Synchrotron x-ray beam. This run was made primarily to evaluate the usefulness of bubble chambers with such machines, and was made in cooperation with the California Institute of Technology synchrotron group. Several other runs were made to provide experimental data on the system thermodynamics. Excellent agreement between the theoretical and experimental data was found. It is planned to install a redesigned chamber of improved mechanical design in this instrument. A complete rework of the control circuits was completed, resulting in greatly improved reliability.

10-Inch Status

By the end of July all parts for the 10-inch chamber except the expansion line and expansion cylinder were completed. All parts were leak tested, and a trial assembly was finished. The illumination system has been modified so that the flash tubes are internal to the vacuum system. The electronic control racks are essentially complete. The chamber proper

Alvarez

has been pressure tested to 160 psi with liquid nitrogen without leakage. The complete system will be tested with liquid nitrogen in the first week of August.

72-Inch Status

Design work is proceeding on the magnet, magnet power supply, hydrogen storage, vacuum vessel, and window. The refrigeration system cannot be chosen until the thermodynamic analysis is complete. At present it appears that a low-temperature expansion system is essential. Various illumination schemes are under consideration. A reflective (one-window) method, using solid Venetian blinds and a curved mirror, seems promising. As the 10-inch chamber nears completion, increased engineering manpower will be available.

Bubble Chamber Data Analysis

A. Data Reduction. Alvarez has pointed out that bubble chambers will accumulate data at a rate comparable to counter experiments, and with each event defined to the accuracy of cloud chamber experiments. Clearly the usual methods of analyzing cloud chamber data, which require the order of an hour per event, will not be adequate. We plan to make rapid measurements of the coordinates and (or) slopes and (or) curvatures of the tracks on each of two stereo photographs, and record these on IBM cards. The cards will be used in an IBM 650 machine, in conjunction with data from a previously photographed grid and information on the magnetic field, to compute for us the true three-dimensional trajectory and momentum of the particle so that the coordinates, radius of curvature, and hence momentum components of the track can be determined. It appears that high-speed comparators, which have been developed for ballistics research, can do the job of coordinated measurement with useful speed and accuracy.

The computer groups at Berkeley and Livermore have assisted in discussions of the IBM computations, and have begun sample programming for 4-inch and 10-inch chamber data.

B. Apparatus. All bubble chamber runs to date have used apparatus that Wilson Powell has generously lent us: stereo camera, "leaning tower" viewer, and measuring microscope. We have now assembled a stereo camera of our own, using Rekordak components for camera housing, film advance, and vacuum backing. This camera will be used with the 4-inch and 10-inch chambers. Construction of a stereoscopic projector is nearly finished.

Two types of film reader are being designed:

(a) A projector that will enable us to measure and record bubble coordinates onto IBM cards at a rate of about 10 points per minute. This unit, with an accuracy of approximately 0.003 inch, can be obtained commercially by small modification of a commercially available film reader.

(b) A projection comparator with somewhat higher accuracy. This unit will also punch coordinates in IBM cards. It will be used as a film reader and as a basic unit for experiments with semi-automatic track following. As a first step toward automation we have built, with William Fowler, a cathode ray display to permit the coordinates of a track center to be set very rapidly and accurately.

Alvarez

PHYSICS RESEARCH

Walter H. Barkas in Charge

K-MESON MASS DIFFERENCES

Walter Barkas, Harry Heckman, and Frances M. Smith

Considerable work has now been done on tracks of K-mesons in a stack that was exposed under conditions favorable for precise mass measurements. The momentum resolution of about 1% was achieved by stringent collimation of the K-meson beam and by reducing the gas scattering to a negligible amount by substituting helium for air in the path of the particles. Grids have been carefully aligned on the stacks and the rectified K-particle ranges now are being measured by breaking the range down into straight segments, each of which is measured. Careful account is being taken of surface and edge corrections, and the emulsion shrinkage factor is accurately known to be 2.21. When the inclination of the secondary track is less than 30° , the $K_{\mu 2}$ and $K_{\pi 2}$ groups (which together with the tau mesons constitute the bulk of the particles) are readily separated. Mass measurements are carried out only on K particles identified as to type. Thus far, ranges have been measured for about 100 K particles. The τ and $K_{\mu 3}$ mesons were identified by following the secondaries to the point where they stopped. The statistical distribution of the ranges has about the calculated variance except that about 5% of the events seem not to have come directly into the stack through the collimating system. Quotation of results will be deferred until the range measurements are completed and until the anomalous events can be more thoroughly investigated.

The τ , $K_{\pi 2}$, and $K_{\mu 2}$ mesons lend themselves to another possible mass comparison by the measurement of the ranges of their secondaries. Unfortunately, no experimental range-energy relation now exists for determining the energies from the ranges, the range measurement is a difficult one, and it is essential that the emulsion density be known at the time of exposure. It is also important that the grain density of minimum tracks be high everywhere in order that large errors not be made in measuring the segments of track in individual emulsions. Thus far, the range of only one $K_{\mu 2}$ secondary has been measured in which all these conditions were met. The range was 19.81 cm in emulsion of specific gravity 3.84. This measurement was checked in various ways, and no allowance for air gaps between the pellicles was found necessary. No paper was used between the sheets and the stack was tightly clamped.

RANGE-MOMENTUM RELATION

Walter H. Barkas, Paul H. Barrett, P. Cüer, Harry H. Heckman,
Frances M. Smith and H. Ticho

In a program designed to yield accurate data on the stopping power of emulsion for high-velocity particles, the range-momentum curve is being determined up to $\beta = 0.8$ or higher. This is being done with the use of magnetic analysis to obtain an accuracy of 0.1% for the momentum of pions,

Barkas

protons, deuterons, and other nuclei. This information, of course, is needed for the study of K-meson secondaries. Ranges are measured in emulsion of accurately known density. Special precautions are taken to avoid drying the emulsion surface when it is placed in vacuum. A good exposure was obtained with the cyclotron but no results are available yet.

EXPERIMENTS WITH THE NEUTRON BEAM FROM THE BEVATRON

Paul H. Barrett

Neutrons produced by the bombardment of a polyethylene target with 6.2 -Bev protons have been used in two experiments using nuclear emulsion. The exposure of test emulsions in the neutron beam indicated an undesirable flux of photons, so a system of lead converters and magnetic fields was used to reduce the photon flux accompanying the neutrons.

In the first experiment, 1-by-3-by-10-inch stack of Ilford G.5 emulsion was exposed to the neutron beam. The analysis of this stack should give (a) the flux of neutrons, (b) the mean free path for high-energy neutrons in emulsion, and (c) a rough determination of the neutron spectrum.

The second experiment was designed to measure the absorption cross section for high-energy neutrons (4 to 5 Bev) in lead, aluminum, and copper. The emulsions from these experiments have been processed and a preliminary analysis indicates that the exposure was very satisfactory.

NUCLEAR TRACK PHOTOMETER

Fred Inman

Refinements have been carried out on the nuclear track photometer during the past quarter. The photometer was tested by measuring protons and K particles. After calibrating an emulsion by scanning 50 proton tracks and determining a depth calibration curve and convincing ourselves of the independence of ionization on this curve, we proceeded to determine the usefulness of the instrument. At 700 microns residual range, the proton integrated density was 190.7 ± 2.6 for three protons and the K integrated density was 179.3 ± 3.1 for four K's. Thus a fair separation is obtained at 700 microns. Mass determinations were carried out on 3-mm-long K tracks and with the track densities of four K's were compared to the average for three protons; a K mass of 960 ± 90 was found. A different method of approach to track measurement was undertaken. Integrated gap-length measurements were undertaken by measuring the charge put on a condenser, the charge corresponding directly to the integrated gap length. The result of these measurements for 3-mm-long K tracks was 965 ± 50 (for four K tracks compared to three proton tracks).

ANGLE AND ENERGY DISTRIBUTIONS OF CHARGED PARTICLES FROM THE HIGH-ENERGY NUCLEAR BOMBARDMENT OF VARIOUS ELEMENTS

Evan Bailey

Absolute cross sections have been determined for charged secondaries from the 195-Mev proton bombardment of carbon, aluminum, nickel, silver, and gold, and the 220-Mev alpha bombardment of aluminum and silver. The secondaries studied are protons, deuterons, tritons, He^3 , and alpha particles. A report of this work is being written.

Nuclear track emulsion has also been exposed to secondary charged particles to obtain absolute cross sections for proton, deuteron, and alpha bombardment (at 340, 190, and 380 Mev respectively) of aluminum, nickel, silver, and gold. The targets are monitored for beta and gamma activity, and their activities are compared with those of targets exposed to the external cyclotron beam. The relative cross sections for particles produced in these bombardments were measured by Robert W. Deutsch and reported in UCRL-2258.

INTERACTIONS OF 5.7-Bev PROTONS IN G.5 EMULSION

Peter C. Giles

Close examination of the tracks under high magnification (efficiency ~100 %) has revealed many previously undetected interactions with free protons and interactions presumably with peripheral neutrons. A free path of ~33 cm for interactions of all types is now found.

A detailed analysis of all types of interactions will follow the complete cataloging of these events.

The scanning is being carried out in emulsion exposed with the proton beam penetrating normally to the plane of the emulsion.

THEORETICAL GROUP

David L. Judd in Charge

A study of renormalization has been made. The technique is being applied to the calculation of a cutoff-dependent value for the fine-structure constant. (Richard Spitzer)

The small-angle elastic scattering of high-energy (300 Mev) protons from nuclei has been examined. Relativistic effects, which cause a spin orbit potential to arise from the Coulomb interaction, result in a significant effect on the polarization of the proton in the forward direction, depending on the sign of the nuclear spin orbit potential. Various aspects of the experimental results of the polarization at small angles have been correlated to Coulomb interference effects. (Warren Heckrotte)

Some reactions of K^- particles in hydrogen and deuterium were investigated (UCRL-3099). They are of interest in connection with the bubble chamber program at the Bevatron. Another project, which is being written up, was an investigation of the symmetry in isotopic spin space of the Hamiltonian operator for strongly interacting particles. It was found that charge independence and charge conservation actually imply full rotational symmetry if the charge is related in a simple way to the rotation operator in isotopic spin space. (Kenneth M. Case, Robert Karplus, and C. N. Yang)

An analysis of the effect of Coulomb forces on allowed favored Fermi transitions has been begun. (William M. MacDonald)

Conditions under which relationships between bound-state problems and scattering theory exist have been investigated. When applied to the many-body nuclear problem, the integral equations used by Brueckner (Phys. Rev. 97, 1353 (1955)) can be somewhat clarified. A report on this work is now in preparation. (Bryce DeWitt, Roger Newton, and Alfred Reifman)

On the basis of the isotopic spin assignments of Gell-Mann for the "strange" particles currently being studied at the Bevatron, and on the assumption of isotopic spin conservation in fast reactions, relations between different cross sections for K^- -nucleon interactions were derived and summarized in Report No. UCRL-3074. Similar work on the cross sections for various "associated productions" in nucleon-nucleon collisions is in progress. (Stephen Gasiorowicz)

Preliminary work has been completed on the application of various simple perturbations to an isotropic harmonic oscillator potential, with a view to obtaining a set of energy levels and wave functions capable of predicting ground-state properties of heavy nuclei within the framework of the shell model. A calculation of energy levels and wave functions of an anisotropic harmonic oscillator with a flattened bottom and including spin-orbit coupling is now in progress. This problem is being treated by first-order perturbation theory about the isotropic harmonic oscillator. Also, some thought is being given to the possibility of setting up a machine calculation using a more realistic nuclear potential, which will include the deviations from sphericity and spin-orbit energy. (Marvin Rich)

An investigation of the treatment of particles in an ionized plasma in the absence of external forces has been completed (with Stephen Gasiorowicz and Maurice Neuman). The generalization of this work to the case in which external magnetic fields are also present has been begun. In this case, the approach is to treat the magnetic field exactly, while the interactions between particles give a perturbation on the otherwise circular motion of the particles. Because of the perturbations the particles will now diffuse in coordinate and velocity space, and these effects are being investigated. (Robert J. Riddell, Jr.)

The possibility of deriving new information on μ -meson decay from muons trapped in aligned nuclei has been considered. Because the muon lifetime is short compared with the relaxation time in an ordinary lattice, and because the decay electrons would be isotropically emitted (though polarized), this possibility is found to be very unpromising. (Sidney Bludman)

A model of all the strange particles as nucleon-anti- Λ^0 composites has been investigated qualitatively. In such a scheme, the slow decay of all the strange particles would follow from the slow decay of the Λ^0 (which itself might be attributable to a weak Fermi interaction). The mass degeneracy among K mesons of apparently different angular momentum and parity, as π and θ , appears naturally, provided the strong nucleon-anti- Λ^0 interaction is static. Any more quantitative estimate of the nucleon-anti- Λ^0 interaction appears beset by extreme relativistic and recoil difficulties. These conclusions had previously been arrived at by Marshak and Levy (Nuovo Cimento 12, Suppl. 2, 253, and private communication). (Sidney Bludman)

The possibility of finding an extended invariance principle leading to a definite form of coupling between elementary particle fields is still being studied. (Sidney Bludman)

The investigation of the validity of the Tomonaga approximation for the one-nucleon-state vector by making a second iteration by the method of Dalitz and Ravenhall has been completed, with the result that the difference between the second approximation and the first is the same as the difference between the first and the zeroth. (Kent Curtis)

Study is also being undertaken on the formal scattering theory and meson production with particular application to π^0 production in proton-proton collisions. (Kent Curtis)

The exact method of obtaining all central potentials that cause a given phase shift has been used for the derivation of variational derivatives, with respect to the phase shift, of the potential, the wave function, and the phase shift of another angular momentum. The result yields a simple exact relation between the value of the potential at the origin and the first moment of the phase shift. It can, furthermore, be used as the basis of a "phase shift perturbation method." In the presence of bound states, an "intrinsic" S-matrix suggests itself naturally as an "unperturbed" starting point, and the "intrinsic" potential can be found explicitly. The perturbation method near the "intrinsic" quantities was also generalized to the possible presence of a tensor force. A paper has been prepared. (Roger G. Newton)

An attempt to include conservation of angular momentum in statistical
Judd

theory of nuclear events is being made. (D. H. Holland)

Work has been done with the formalism of Low (Phys. Rev. 97, 1392 (1955)) in which the Dyson S-matrix is used as the starting point for developing an exact, although nonlinear, integral equation for the meson-nucleon scattering problem. An integral equation has been derived for the amplitude for scattering plus production of one additional meson. One would like to attempt a solution of this equation in the nonrelativistic limit of the γ_5 theory in a manner similar to that used by Chew to get the P-wave phase shifts from the Low equation. Comparison of results with proposed experiments with pion beams in the medium-energy range, where the cutoff theory might be valid, would be of interest. Such a reaction would be

$$\pi^- + p \rightarrow \pi^- + \pi^+ + n.$$

Also being considered is another approach to multiple meson production by putting angular momentum conservation into the statistical theory of Lepore and Neuman (Report No. UCRL-2649). (Saul Barshay)

An attempt was made to explain the observed difference between Fermi and Gamow-Teller nucleon beta decay coupling constants, using the philosophy that they are really the same in some fundamental sense. We argue that it cannot be that the couplings to the bare nucleon are equal, since the mesic corrections are infinite, differently for each coupling. Thus if the renormalized couplings are taken to be equal, the observed difference between the couplings can only be a result of the finite energy-momentum transfer at the beta decay vertex (a minute correction) and many-nucleon "cooperative" effects. The difficulty is that only the former effect operates in neutron beta decay, but here the ratio of coupling constants (as determined by angular correlation) is the same order as in nucleus beta decay. (Charles Goebel and William MacDonald)

Investigations on the consequences of causality are continuing. In particular the Kramers-Kronig relations are being applied to the pion-proton phase shifts at low energy with the aims (a) to determine the energy dependence of the S-wave phase shifts; (b) to then make a best fit of the data, in order to determine parameters of practical importance, such as the Panofsky number $\delta_1 - \delta_3/k$, and of theoretical importance, such as the threshold S-wave coupling constant g_s . (Charles Goebel and Robert Karplus)

High-energy pion-proton experiments continue to be studied, with especial reference to the possible role of meson-meson interaction. (Charles Goebel)

An article on the effect of Coulomb interference in proton-proton scattering was completed. Further calculations were made of this effect in connection with the phase-shift analysis of the Berkeley experiments being done by Ypsilantis and Stapp. In collaboration with Warren Heckrotte further estimates were made of various possible sources of error in the experiment with the spiral-orbit spectrometer measuring the energy spectrum of the electron from μ -meson decay. It appears that the change of magnetic field shape with field setting due to saturation is serious, and probably sufficient to account for the discrepancy between the previous values of ρ obtained by Sagane, Dudziak, and Vedder and of other experiments. (Alper Garren)

Judd

A Hamiltonian form of the wave equations for integral spin particles was obtained (UCRL-3067). (Kenneth M. Case)

A lower limit is found for the momentum derivative of the scattering phase shifts of a relativistic neutral two-particle system when the interaction is of finite range. (Charles Goebel, Robert Karplus, and Malvin Ruderman of the Physics Department, University of California) (UCRL 3034)

Within the framework of the higher dimensional Dirac equation a classification of the observed elementary particles has been constructed. In the course of the work a general expression for the charge operator for Baryons and Bosons suggested itself. Its consequences on decay schemes are currently being investigated. (Maurice Neuman)

A paper on the generalized effective range theory of scattering, begun last quarter, is almost completed. To be included are upper and lower limits on core radii in the 1S_0 state, and their dependence upon the type of external attractive well used. A second paper on a method of deducing properties of the force directly from the scattering, as reported in the last quarter, is in progress. Work on higher approximations in this method is continuing. (Robert Raphael)

During the past quarter, work has been completed on an investigation of polarization effects in nuclear scattering. This work was divided into several parts. First a phase-shift analysis of the polarization data for n-p scattering was carried out at Los Alamos. Secondly, since the energies involved approach values for which relativity will play a role, a relativistic formulation of the polarization effects was developed and the associated corrections were investigated. In addition, a formalism suitable for treating spin-one projectiles against spin-zero targets was developed, and, finally, some approximate calculations for the polarization of deuterons were carried out. In these, the first and second Born approximations were used, together with determination of corrections due to the presence of the d-state in the deuteron wave functions. (Henry Stapp)

A study is being made of some of the implications of charge independence within the context of the theories of Gell-Mann and Pais for the strange new particles. (David Feldman)

A paper (UCRL-3056) on the mathematical justification of the perturbation theory of differential operators has been completed. The eigenvalue problems given by

$$\frac{d^2 u}{dx^2} + \left\{ \lambda - q(x) - \epsilon p(x) \right\} u = 0, \quad 0 < x < \infty$$

and

$$\nabla^2 u + \left\{ \lambda - q(x, y, z) - \epsilon p(x, y, z) \right\} u = 0$$

in ordinary three-dimensional space are considered, where ϵ is a small real quantity. The expansions for the eigenvalues and eigenfunctions which are given by formal perturbation theory are justified as asymptotic series, valid

for a finite number of terms as $\epsilon \rightarrow 0$. The approximations are established up to second order by placing certain restrictions on the function p . (John Killeen)

While it is a matter of convention to say that the neutron and proton represent eigenstates of the isotopic spin with values -1 and $+1$, respectively, the corresponding assignments for antineutrons and antiprotons are not free once this choice is made. The correct choice may be inferred from experiment, for if these particles can be produced by gamma rays (or electrons) one must choose isotopic spin values -1 and $+1$ for antineutron and antiproton respectively. This ambiguity may also be taken to mean that there are two types of particles in nature; one in which the antiparticle has the same value of the isotopic spin, another type where this assignment is just the opposite. A possible assignment would be neutron and proton to the first type and the cascade particle to the second. (Joseph V. Lepore)

PHYSICS RESEARCH

Edward J. Lofgren in charge

ELASTIC PROTON-PROTON SCATTERING

Bruce Cork and William A. Wenzel

Measurement of the differential cross section for elastic proton-proton scattering as a function of angle at Bevatron energies is in progress. The internal beam, monitored by an electrostatic induction electrode, strikes a polyethylene target in the upstream end of a straight section. Protons scattered at angles from 10° to 45° (laboratory system) emerge through a thin (0.090-inch) aluminum window on the outside radius of the Bevatron. These are detected by a two-counter telescope in coincidence with a similar telescope on the inside radius of the Bevatron, which detects the larger-angle recoil. Fourfold coincidences with a time resolution of 6×10^{-9} second and good geometrical resolution (10^{-4} steradian) permit separation of the elastic peak from a rather large inelastic background.

Experimental results are as follows:

<u>T</u> (Bev)	<u>θ</u> c. m.	<u>$\frac{d\sigma}{d\omega}$</u> (mb/ster)
0.92	49 ⁰	3.3
	72	0.76
	92	0.45
2.24	44	1.5
	58	0.58
	70	0.33
	93	0.19
3.49	34	1.7
	49	0.26
	64	0.07
	78	0.06
4.40	36	0.61
	53	0.12
	69	0.045

An internal counter is being arranged so that measurements can be made at smaller angles and at higher energies.

ANTIPROTON SEARCH

Joseph J. Murray, Nahmin Horwitz, Bruce Cork, William A. Wenzel,
John M. Brabant, and Roger W. Wallace

A counter telescope has been used to search for antiprotons in the beam from the magnetic analyzing system devised and constructed by Segrè, Chamberlain, Wiegand, Goldhaber, Chupp, and Ypsilantis. The telescope, shown in Fig. 1, consisted of scintillation counters to define the beam, three

Lofgren

Fig. 1. Telescope schematic.

Cerenkov counters with radiators made of different materials, an absorber, and a deflecting magnet. With all scintillators and the polystyrene counter in coincidence, the water and lucite counters in anticoincidence, and the deflecting magnet set for negative particles, the telescope should respond to negative protons with $\beta = 0.76 \pm 2\%$ and should reject positive protons, K and π mesons, and electrons of any β . This system was followed by a large lead-glass Cerenkov counter in which annihilation of stopped negative protons could produce pulses much larger than those from minimum particles.

In a preliminary test using positive protons, the telescope exhibited the expected β resolution and had an efficiency of 3% to 4%. This response curve is shown in Fig. 2.

Approximately two hours were devoted to antiproton search. During this time roughly 4×10^5 π^- mesons were incident on the front aperture of the telescope, and one signature was obtained. The lead-glass Cerenkov counter pulse associated with the signature, however, was only twice the height of pulses produced by π^- mesons.

$C^{12}(p, pn)C^{11}$ CROSS SECTION AT BEVATRON ENERGIES

Nahmin Horwitz and Joseph J. Murray

The work described in the last report has been continued. The integrator-flyback circuit has been employed successfully to obtain pulse-height distributions for correlation of photomultiplier output with absolute number of protons passing through the target.

It still remains to complete a high-accuracy measurement of the linearity characteristic of the photomultiplier. Apart from the uncertainty arising from this problem, the $C^{12}(p, pn)C^{11}$ cross section appears to be essentially flat from 340 Mev to 4 Bev. In a later run measurements will be repeated at low energy and extended to 6 Bev.

τ -MESON LIFETIME

Luis W. Alvarez and Sulamith Goldhaber

The lifetime of the τ meson has been measured by a comparison of relative numbers at two different flight distances from the target. Nuclear emulsions were used as detectors to determine the relative numbers of τ mesons in a given momentum interval (350 Mev/c). The resulting mean life value is $1.0^{+0.7}_{-0.3} \times 10^{-8}$ sec.

This work is reported in detail in UCRL-3041 and will appear in Nuovo Cimento.

INTERACTIONS OF NEGATIVE K PARTICLES AT REST¹

Warren W. Chupp and Sulamith Goldhaber

K⁻ mesons produced in a copper target by 6.2 Bev are accepted into

¹ This work is done in conjunction with Francis H. Webb and Gerson Goldhaber. Lofgren

Fig. 2. Response of telescope for positive protons.

the strongfocusing-lens analyzing magnet at 90° for the proton beam.² Stacks of Ilford G.5 pellicles have been exposed to the focused K^- beam of momenta 300 Mev/c to 410 Mev/c.

K^- mesons are selected without bias by picking all tracks of grain density between two and three times minimum at the incident edge of the stack. A preliminary report on the study of 22 K^- stoppings has been sent to the Pisa Conference³ and an abstract to the Mexico City Meeting.⁴

A total of 130 K^- stoppings have been observed to date. The analysis of these events according to the reactions

$$K^- + \text{"nucleon"} \rightarrow \Sigma + \pi$$

is in progress. A prong distribution for 131 stars is given in Fig. 3. The visible energy release in most of the stars accounts for only a small part of the rest energy.

K-PARTICLE PRODUCTION STUDIES

Warren W. Chupp and Sulamith Goldhaber

Experimental evidence at present indicates that the masses of the various K mesons are all equal (within some few electron masses) and equal to the τ -meson mass ($965.5 m_e$). The experimental data on lifetimes also indicate that there is no distinction between the various K mesons. If the τ meson is distinct from any of the other types of K mesons, then it may be possible to observe a difference in production at different bombarding energies. Studies using nuclear emulsions are under way in which the ratio K/τ will be measured at different proton bombarding energies.

The experimental arrangement used for the emulsion exposures consisted of:

(a) A collimator 0.5 inch high by 6 inches wide, placed at 90° to the target and 30 inches away (just outside the thin window in the west straight section of the Bevatron);

(b) A 90° wedge-sector magnetic field. The field gap is 3 inches, and the intensity used was 12,000 gauss.

This arrangement gives good dispersion (5 Mev/c per inch in the focal plane) and permits the exposure of large emulsion stacks at several different momenta simultaneously.

² Kerth, Stork, Birge, Haddock, and Whitehead, Bull. Am. Phys. Soc. 30, No. 3, 41 (1955)

³ Chupp, Goldhaber, Goldhaber, and Webb, "Interaction of Negative K Particles at Rest", University of California Radiation Laboratory Report No. UCRL-3044, June 14, 1955

⁴ Bull. Am. Phys. Soc. 30, No. 5, 13 (1955)

MU-10036

Fig. 3. Prong distribution of K^- capture stars.

The proper time of flight for K mesons of 350 Mev/c momentum is approximately 1.6×10^{-8} sec.

Several exposures have been made using this arrangement. They are described as follows:

1. K^+ mesons of 250 Mev/c, 350 Mev/c, and 450 Mev/c momenta produced in copper by 6.2-Bev protons.
2. K^+ mesons of the same three momenta produced by 4.2-Bev protons.
3. K^+ mesons of the same three momenta produced by 2.6-Bev protons. (Emulsions were also placed along a copper wedge sufficiently thick to stop 330-Mev/c π^+ during this run. This was done to facilitate the comparison between K^- and K^+ meson exposures, which necessitates reversing the magnetic field.)
4. K^- mesons of 250 Mev/c and 350 Mev/c produced by 6.2-Bev protons (again the copper wedge was used to stop 330-Mev/c π^-).

The frontal face of the target was covered by several thin aluminum foils, which serve to monitor the integrated proton beam for each exposure. The induced Na^{24} activity was counted to determine the relative proton flux at each exposure so that comparison of K mesons at different energies is possible.

The stacks at present are being processed.

Edwin L. Iloff and Gerson Goldhaber of Prof. Emilio Segrè's research group have assisted greatly throughout the course of this program.

PHYSICS RESEARCH

Burton J. Moyer in Charge

NEUTRON MULTIPLICITIES FROM μ^- CAPTURE

George P. Millburn, Donald A. Hicks, Robert V. Pyle, and John Ise, Jr.

An experiment is planned to measure the neutrons emitted from various nuclei following capture of μ^- mesons. The neutron detector will be the large liquid scintillator tank (30 inches diameter by 30 inches) with a detection efficiency of about 60%. A series of four plastic scintillation counters will be used; two in coincidence will define a beam of cosmic rays that passes through the absorber, a third will be used to identify a minimum-ionizing particle, and a fourth in anticoincidence will indicate those particles that do not stop in the absorber. The pulses from the large tank will be displayed on a Dumont 517 oscilloscope and photographed along with the pulses from counters 3 and 4. The measured μ^-/μ^+ ratio will be used to determine the number of μ^- mesons stopping the absorber, and eight inches of lead will serve to remove the electronic component of cosmic rays.

The experiment is being done to measure the excitation energy given to the nucleus. If the μ^- is captured by a proton ($\mu^- + p \rightarrow n + \nu$ is the most likely interaction), then the momentum distribution and effective nucleon masses in the nucleus will determine the amount of energy transferred.

Previous measurements have been made with low-efficiency ($\sim 6\%$) neutron detectors and could give only $\bar{\nu}$, the average number of neutrons following μ^- capture. The present technique should also give the second moment of the multiplicity spectrum, $\nu^2 - \bar{\nu}^2$.

Counting rates, exclusive of any background, are expected to vary from 150 to 20 neutrons observed per week for lead and aluminum, respectively. The background is expected to be 1 or 2 neutrons per week.

NUCLEAR GAMMA-RAY SPECTROSCOPY

Charles N. Waddell, Harlan C. Shaw, and Harold E. Adelson

In an effort to test the hypothesis of a $T = 1$ origin for the 15.09-Mev level in C^{12} , an experiment was conducted at the 184-inch cyclotron to measure the yield of gamma rays when a carbon target was bombarded by alphas and deuterons. Isotopic spin selection rules would inhibit the production of a $T = 1$ state in C^{12} , as all the nuclei are even-even and hence have a $T = 0$ ground state. To permit a variety of bombarding energies to be used, the targets were viewed at 90° in the laboratory system by a 3-by-3-inch NaI(Tl) crystal. The pulse-height distribution was measured with a ten-channel pulse-height analyzer.

Very low beam intensities had to be used because of the ten-second resolving time of the pulse-height analyzer; consequently it was impossible to monitor the beam with our thermocouple arrangement. Therefore we measured the relative intensities of gamma rays at 90° from the 4.43- and 15.09-Mev levels. The 4.43-Mev level is $T = 0$ and hence its production will

not be inhibited by isotopic spin selection rules.

Another problem inherent in the alpha bombardments arises because normally there is a large contamination of deuterons in the alpha beam unless a sufficient amount of time has elapsed since a deuteron run. Because we could use only very low-intensity beams, we were able to take advantage of the different e/m values for alphas and deuterons to separate them by adjusting the arc timing to reduce beam intensity. We were able to test the efficacy of this technique in a bombardment of Be^9 , as with the timing set for alphas the gamma-ray spectrum was the characteristic 4.43- and 15.09-Mev gamma rays from C^{12} , whereas with the timing set for deuterons there was no trace of the C^{12} gammas. This was also the first time we have observed the 15.09-Mev level in the $\text{Be}^9(\alpha, n)\text{C}^{12*}$ reaction, as a previous run using a pair spectrometer did not give an observable yield,¹ although Rasmussen et al.² reported this reaction.

When 175-Mev alphas are incident upon a carbon target of sufficient thickness to stop the beam, there is a production of gamma rays of ~ 15 Mev which can be attributed to the 15.09-Mev level. However, when the target is pushed in to a radius where 42-Mev alphas are incident on the target, there is no observable yield of high-energy gamma rays. It seems probable that the gamma rays formed by high-energy alphas are due to the breakup of the alpha particle and subsequent excitation of the carbon nucleus by the emitted protons and neutrons. Previous experiments have shown that both neutrons and protons incident upon carbon will produce this level. Similarly, high-energy deuterons produce the level weakly, but time did not permit the reduction of energy as for the alpha experiment.

PROTON-PROTON π^0 PRODUCTION

Robert K. Squire and Robert J. Cence

The experiment to determine the cross section for π^0 production from protons on protons has almost been completed.

The cross section for the reaction $(p + p \rightarrow \pi^0 + p + p)$ is expected to be dependent upon the maximum meson momentum, η_0 , as

$$\sigma = a_1 \eta_0^2 + a_2 \eta_0^6 + a_3 \eta_0^8$$

The experimental results indicate that the term a_2 is zero, or very small, and that the ratio a_3/a_1 is at least as large as 20:1, although the best final results have not yet been determined. The quantity $a_3 \eta_0^8$ is the cross section for mesons to be produced in the p state with respect to the two-nucleon system and for the two nucleons to be left in a p state with respect to each other. There seems to be a strong preference for this over the other cases (a_1 and a_2) where the meson is emitted in the S state.

¹ D. Cohen, B. J. Moyer, H. Shaw, and C. N. Waddell, "Emission of High-Energy Gamma Line from C^{12*} ," Phys. Rev. 96, 714 (1954).

² V. K. Rasmussen, J. R. Rees, M. B. Sampson, and N. S. Wall, "A Search for 15-Mev Gamma Radiation," Phys. Rev. 96, 812 (1954).

BETA-RAY SPECTROSCOPY

Roger W. Wallace and Jack A. Welch

During the first part of June a study of $\text{Al}^{27}(\text{p}, \text{n})\text{Si}^{27}$ was made on the Linac. The results indicated

- (a) That the apparent half life was too long and varied with positron energy. This was later traced to a mixture with $\text{Al}^{27}(\text{p}, \text{n})\text{Si}^{26}$, formed when the absorber failed to move properly.
- (b) That an annoying long-lived (hours) positron activity was developed. This was blamed on brass parts in the target holder, which have now been replaced with lucite. The last part of June an exact repeat of the $\text{Al}^{27}(\text{p}, \text{n})\text{Si}^{27}$ run in June 1954 was made (i. e., stationary absorber, same target). This showed that indeed the present detector is more efficient, and that the above-mentioned difficulties were cleared up.

A good (1% statistics) Kurie plot was obtained for a 10-mil (137 kev) target. This showed a very bent line, which is probably due to poor resolution.

The first part of July was taken up with several mechanical changes:

- (a) an integral faraday cup to subtend a larger solid angle,
- (b) rearrangement of collimating slits,
- (c) reworking of absorber charger.

PHOTONIC DECAY OF HEAVY MESONS AND HYPERONS

John E. Osher and Sherwood I. Parker

The gamma-ray telescope used in this experiment was calibrated by folding theoretical curves for pair production and energy splitting with an experimental determination of electron counting efficiency. The latter was obtained as a function of geometry, energy, and remaining converter thickness during a short run at the synchrotron.

Because of the mass involved in the plunging slit system, a 3-inch-diameter probe was found to be necessary. This required a relocation of the experimental area and allowed a redesign of the slit system to take advantage of a near-by aluminum-walled well for locating further collimation and the gamma-ray telescope.

TOTAL-ABSORPTION PHOTON SPECTROMETER

John M. Brabant, Roger W. Wallace, and Robert J. Cence

The spectrometer that was previously assembled, tested, and calibrated from 40 to 320 Mev has been calibrated up to 1.5 Bev. This was accomplished by using electrons obtained from the materialization of π^0 -decay gamma rays obtained from the Bevatron and analyzed magnetically. The response of the device is very nearly linear over the entire range of calibration. More careful evaluation above 1 Bev seems necessary, both to extend this range and to investigate any possible departure from linearity of response in the higher-Moyer

energy region.

The spectrum of photons from π^0 decay at 6° to the incident beam has been obtained for incident protons of 6.2 Bev energy. A run to investigate the spectra at 90° and approximately 120° is scheduled for August 10-11. This should also provide additional information on the desirability of increasing the speed of the associated electronics. Additional mobility of operation has been achieved by a complete rearrangement of equipment presently used.

The spectrometer is also being used in connection with antiproton search by Edward J. Lofgren and his group. It is hoped that there will be a correlation between possible counter-telescope results of that group and possible evidences of 2-Bev annihilation radiation obtained simultaneously by the spectrometer.

PION-PROTON CROSS SECTIONS

Burton J. Moyer, Hoyt A. Bostick, Kenneth C. Bandtel, Roger W. Wallace, N. Fredrick Wikner, and Selig N. Kaplan

One further run in the measurement of pion-proton cross sections has been made since the last Quarterly Report. This was devoted to tying in Berkeley data with Brookhaven data by measuring a value of the cross section down near 2 Bev. The orbit actually employed delivered pions of 2.3 Bev, and at this energy the π^- -p cross section of 32 ± 4 millibarns was determined.

The large error makes it desirable to gather further data at this energy, and this is contemplated for a succeeding run.

Orbits for positive pions are being calculated in preparation for the π^+ -p measurements. Also, experimentation on counting methods to be employed here has been successfully pursued. The problem involved is the separation in detection of positive pions from protons which follow the same momentum orbit. This can be accomplished through the use of a high-pressure gas Cerenkov counter, which allows the detection of pions while protons of the same momentum are below the Cerenkov threshold. Experimental tests with this type of counting have been successful.

Preparations for exploring pion-neutron cross sections have been made through the use of light- and heavy-water attenuators. Also, it is planned to make at least one run with a liquid hydrogen attenuator, and if agreement is obtained with the C-CH₂ data, the liquid hydrogen target will probably not be further used.

PROTON COMPTON SCATTERING

Burton J. Moyer, Kenneth C. Bandtel, and Larry L. Higgins

Preliminary efforts on the Compton scattering of 140-Mev photons with protons have involved the study of background phenomena and means of reducing them to levels that allow the observation of the desired events with their

Moyer

very low cross section. The coincidence-anticoincidence counter rings described in the Quarterly Report for November to January have been employed, and satisfactory shielding against stray gamma radiation and against the effects of ambient neutrons have been achieved. The coincidence network with its timing conditions reduced the cosmic ray background to suitably low levels; however, one very serious background problem remains which is due to fast neutrons coming down the collimator tube and scattering into the counter rings. A small number of double recoil events were observed in which the neutrons produce recoil protons in both counters of a given unit after the converter, and the pulse heights are such as to make these events almost indistinguishable from the counts produced by Compton-scattered photons from the hydrogen protons. A further background which is always troublesome is the vast sea of low-energy electron-Compton-scattered photons from the hydrogen.

It has been concluded that Cerenkov counters must be introduced into the counting units following the converters in order to make them insensitive to the neutron recoil events referred to above. A given counter unit would then consist of an anticoincidence scintillation counter followed by a lead converter, and this in turn followed by a scintillation counter and a Cerenkov counter in series. As before, there will be a ring of 12 such units distributed around the target in the 90° plane.

CLOUD CHAMBER STUDIES

Wilson M. Powell in Charge

EXPERIMENTS

4.5-Bev π^- -p InteractionsWilliam B. Fowler, George Maenchen, Wilson M. Powell, George Saphir,
and Robert W. Wright

A preliminary report, UCRL-3042, was completed. Three more days of Bevatron time yielded additional events. It is anticipated that at least three more runs of three days each will be required in order to have sufficient statistics after breakdown of the data into their various categories -- i. e. elastic and different types of inelastic. The discovery of two new six-prong events in the last set of pictures has verified that the original two cases were not just the result of a statistical fluctuation but represent a real rate of very high-multiplicity meson production.

High-Energy n-p InteractionsWilliam B. Fowler, George Maenchen, Wilson M. Powell, George Saphir,
and Robert W. Wright

Interactions with protons of neutrons emitted at 0° from 6.2-Bev p-Cu collisions were studied in the 36-atmosphere diffusion cloud chamber in a three-day Bevatron run. Seventy-thousand pictures and somewhat more than 300 events showing meson production were obtained. About 10% of these events had five outgoing prongs. There were also two events with seven outgoing prongs. The five-prong events show at least three meson production and the 7-prong events show at least five meson production. These results are consistent with the very high-multiplicity meson production processes observed in the π^- -p experiment. The multiplicity of meson production is notably greater than the Brookhaven n-p results, where there were no five-prong events out of 185 events resulting from neutrons produced by 2.2-Bev protons on carbon. Two cases of heavy-meson production in n-p collisions were seen. One case is an identified θ^0 , and the other is an identified Λ^0 . In addition, two tau mesons, presumably produced in collisions in the wall of the cloud chamber, were seen to decay in the chamber.

5.3-Bev p-p InteractionsWilliam B. Fowler, George Maenchen, Wilson M. Powell, George Saphir,
and Robert W. Wright

Orbit plots were completed which show that it is feasible to obtain 5.3-Bev scattered protons external to the Bevatron. These protons are scattered at 11° from the 20-inch target and leave the Bevatron through the thin window, making an angle of about 11° with the wall of the straight section. They then enter the 5-foot deflecting magnet and are deflected 8° , which is sufficient to clear the yoke of the main magnet. The channel through the

shielding is parallel to the present neutron channel and is displaced downstream 2 feet. During the next quarter, a three-day run is scheduled with the 36-atmosphere hydrogen diffusion cloud chamber. Information on the 5.3 Bev p-p total cross section and on the elastic and inelastic cross sections should result.

Search for the Antiproton

John Elliott, William B. Fowler, Richard Lander, Larry Oswald,
Wilson M. Powell, and Howard White

An exposure with the magnet expansion chamber is scheduled for the next quarter in which it is hoped to see antiprotons if they are produced at 6.2 Bev. The chamber has five one-centimeter Pb plates and will be placed in a beam of negative particles emitted from the 8° target. Identification will be possible by change in ionization and change in momentum. The charge of the particles will be definitely known because of the magnetic field of 22,000 gauss.

EQUIPMENT AND TECHNIQUES

Machine Analysis

William B. Fowler, Wilson M. Powell, George Saphir, and Howard White

In order to handle the analysis of the rather large number of events that the high pressure hydrogen diffusion chamber accumulates, the IBM 650 computer at Livermore is being programed to compute energy and momentum balance and from these computations to decide whether the event is elastic or inelastic. If the event is inelastic the computer then decides whether a single neutral particle or two or more neutral particles have been emitted. Once a type of collision (elastic or single neutral case) is decided upon, the machine then seeks the solution for the category that requires the smallest weighted adjustment of the measured parameters. This should give a best-fit solution and therefore correspond closely to the actual collision. It is estimated that from 1 to 10 minutes of machine time will be required per event. The wide variation is due to varying numbers of prongs involved and the degree of uncertainty of the event, i. e., if the event is poorly measured the machine will fit more than one type of solution.

New Cloud Chamber Magnet

Cloud Chamber Group

During this quarter the new cloud chamber magnet was completed and the expansion cloud chamber was operated in it for a test run. The major improvements over the old magnet are (a) better temperature control system for the cloud chamber, (b) over-all improvement in electrical design.

Hydrocarbon Bubble Chamber

Larry Oswald and Wilson Powell

The body of the chamber was completed and the windows were received. A pressure test to 600 psi was performed and the pressure was held with no leaks. A system for expanding the chamber was designed and it is expected that during the next quarter the chamber will be expanded. The illumination and temperature regulation systems are being constructed.

PHYSICS RESEARCH

Chaim Richman in Charge

Robert W. Birge, Roy P. Haddock, Leroy T. Kerth, James R. Peterson,
Jack Sandweiss, Donald R. Stork, and Marian WhiteheadMASSES OF THE K^+ MESONS

A summary of the work on positive K mesons for this quarter appears as UCRL-3031, which will be published in Il Nuovo Cimento in the summary of the Pisa Conference on Fundamental Particles, and in UCRL-3083, which will appear as a letter to the editor of the Physical Review.

Physical Review.

Briefly, two emulsion stacks, exposed using the strong focusing spectrometer, have been scanned and the masses of the K mesons determined by the range-momentum method. The values obtained in the two stack are listed separately for τ mesons and K_L mesons. The uncertainties listed are standard deviations only. Systematic errors may change the absolute values.

Stack 16 (114 Mev)

$$K_L \quad 971 \pm 1.3 M_e$$

$$\tau \quad 978 \pm 4 M_e$$

Stack 17 (170 Mev)

$$962.9 \pm 1.9 M_e$$

$$965.4 \pm 3.8 M_e$$

In addition, work is in progress on the analysis of suitably oriented K_L secondaries in an attempt to identify individual mesons as belonging to the category of $K_{\mu 2}$, $K_{\pi 2}$, etc. Identification is established by grain counting the secondaries in a region at least 3.5 cm from the K ending. At least 5% statistics are obtained from each track. When a secondary is identified, the mass of the primary K_L meson is then compared to the average value for τ 's. To date 16 $K_{\mu 2}$'s, 9 $K_{\pi 2}$'s, 4 $K_{\mu 3}$'s and one $K_{e 3}$ have been identified. The mass differences obtained are

$$M_{\tau} - M_{K_{\mu 2}} = -2.2 \pm 9.9 M_e$$

$$M_{\tau} - M_{K_{\pi 2}} = +6.5 \pm 12.7 M_e$$

$$M_{\tau} - M_{K_{\mu 3}} = -0.2 \pm 19.8 M_e$$

$$M_{\tau} - M_{K_{e 3}} = +3 \pm 26 M_e$$

τ^+ ANALYSIS

According to the theory of Dalitz, the spin and parity of the τ meson may be determined by measuring the distribution of the decay configurations. To date approximately 60 τ 's have been analyzed and 30 more are in process. Preliminary results agree with a similar number of measurements collected by Dalitz, which indicate that the spin and parity of the τ is inconsistent with its being an alternate mode of decay of the $K_{\pi 2}$.

LARGE EMULSION STACK

A large emulsion stack of 95 pellicles 9 by 17.5 inches has been exposed to positive K particles behind the strong-focusing spectrometer.

At present over 600 K mesons have been found, including 34 τ 's and 2 $K_{\mu 3}$'s. Six long-range secondaries have been followed to their ends, of which four came from $K_{\mu 2}$'s and two from $K_{\pi 2}$'s. About 10 more tracks were identified before they left the stack. Both the $K_{\mu 2}$ and $K_{\pi 2}$ have a two-body decay scheme giving the secondary particle a unique energy. Hence a measure of the secondary range allows one to calculate the primary meson mass. In addition the primary mass of the identified particles is being measured by the range-momentum method, as for the K^+ mesons.

NEGATIVE PROTON EXPERIMENT

Calculations have been made for the relative cross section as a function of laboratory angle and energy for the production of negative protons in a complex-nucleus target at the Bevatron. An analysis is in process of the magneto-optics necessary to separate out these protons from other negative particles of different mass. A number of combinations of lens systems and target positions have been considered in order to optimize the flux and background conditions for nuclear emulsion stacks.

DIRECT PRODUCTION OF ELECTRON PAIRS BY HIGH-ENERGY ELECTRONS

Stanley L. Leonard

An investigation of the direct production of electron pairs in nuclear emulsions by 400-Mev electrons from the Stanford linear accelerator is being carried out. Direct-production events ("tridents") are found in the emulsion by a track-following technique and are readily identified. To date 16 such events have been found, and it can be tentatively concluded that the mean free path for trident production in emulsions agrees roughly with the production of the theories of Racah¹ and Bhaba² (as adjusted for lower energies by Block et al.).³

¹ G. Racah, Nuovo Cimento 14, 93 (1937)

² H. J. Bhaba, Proc. Roy. Soc. (London) A152, 559 (1935)

³ M. M. Block, D. T. King, and W. W. Wade, Phys. Rev. 96, 1627 (1954)

REPORTS ISSUED

Stanley L. Whetstone, "The Scattering of 22-Mev Positive Pions on Protons" (Thesis), University of California Radiation Laboratory Report No. UCRL-3087, July 1955.

RESEARCH WITH THE SPIRAL-ORBIT SPECTROMETER

Ryokich Sagane in Charge

Walter F. Dudziak and James Vedder

Two assumptions were made in our reduction of the published data on the positron spectrum produced by the decay of the μ -meson (i. e. using Michel's notation, our reported value $p = 0.23$). These assumptions were:

A. That the detection efficiency of our counting system for electrons does not change for an electron energy interval between 30 and 60 Mev. (In practice this is roughly known to be true).

B. That the solid angle covered by the spiral-orbit spectrometer does not change with different magnetic field settings. (This assumption is usually made for most spectrometer measurements.)

In May we tried to experimentally verify assumption A by using a synchrotron beam. The method adopted was: first, by using a uniform magnetic field, to select a certain energy interval of positrons produced at a converter; then, by varying the magnetic field, to compare the positron intensity measured with our four plastic crystals in quadruple coincidence with the intensity measured by a set of Geiger counters in triple coincidence. The test was not as successful as we wished because of the high-frequency pickup of our high-gain system that is necessary for counting relativistic electrons with thin crystals. However, we could say with $\pm 4\%$ accuracy, that the detection efficiency of our counting system was independent of the positron energy in a positron energy interval from 30 to 60 Mev. The high-frequency pickup is caused by the radiation emitted by the synchrotron rf system and is mainly introduced by the long cabling necessary to bring the signals into the counting area. Our plan is to repeat this experiment with the hope of improving this accuracy by (a) shortening the cabling necessary to bring the information into the counting area, and (b) introducing a high-Q wave trap that will filter out the 47-megacycle frequency of the synchrotron rf system.

The reduction of our published data as well as of new data obtained with our old spiral-orbit geometry was checked by the theoretical group and was found to be in good agreement within our accuracy. Certain points of the experiment were not clear to all concerned, as, for example, errors introduced due to contributions from the pole-face effect, electron scattering from the pole faces, Z-focusing effect, and contribution from the Pb shields protecting the crystals. To clarify these points as well as to check for any systematic errors that might be introduced by the spectrometer geometry and electronics, we drastically changed:

(a) The electronics. Among other elements this included using a pulse technique on one phototube dynode of each phototube. By destroying the focusing properties of the phototube with a negative pulse of 200 volts during each proton burst we could reduce the Pb shielding at our crystals without causing serious overloading of the electronics during each proton burst. Under certain conditions we could work with no shielding present.

(b) The light pipe geometry. We thus created a much larger region of overlap during which no magnetic effect could be detected that would influence the counting efficiency by varying the magnetic field.

(c) The magnetic field distribution of the spiral-orbit spectrometer.

Our first experiment with these changes was in July. It was not quite decisive because of the fluctuations that we experienced in our counting rates. However, we could show with sufficient accuracy that there was no large systematic error introduced by the Z-focusing or the other effects mentioned above. We did, however, find that we were measuring a different spectrum by using a different magnetic geometry. By eliminating one set of pole tips at a time and measuring the spectrum we reached the following conclusion. We could reproduce the same results as we always measured (i. e., our published results) with different magnetic geometries as long as a pair of large pole tips was a part of any particular magnetic geometry. (These large pole tips were incorporated into the design of the spiral-orbit spectrometer so as to serve as walls of a large vacuum chamber). If the spectrum was measured without the presence of these pole tips, a different result was obtained.

We established that the effect of scattered electrons from the pole tips did not affect our results significantly. This was accomplished by using a set of nonmagnetic dummy pole tips within the wide gap of our second magnetic geometry.

During our second experiment in early August we obtained very reliable results. Our electronic stability was such that we could reproduce the same results within counting statistics, not only during each day but also from day to day. The only difference between the July and August experiments was the use of a refrigerator unit that kept the fast electronic elements at a nearly constant temperature.

During the second experiment lithium targets one $2\frac{3}{8}$ in. (o. d.) by 3 in. and the other 1 in. (o. d.) by 6 in., were used to measure the spectrum in general. We also used a Li target $5/8$ in. o. d. and $1/4$ in. i. d. by 6 in. for measuring the maximum energy of the spectrum. These data are now being analyzed in detail. The results obtained after a rough reduction indicate the " ρ value" of Michel as ≈ 0.6 , which is in rough agreement with the values reported by other experimenters.

This means that assumption B was proved invalid for our old spiral-orbit geometry. This can only be explained by the unfamiliar sensitivity to saturation of the counting rate as measured by the spiral-orbit spectrometer when a slit system is used close to the detector. It has been well known that the counting rate is very unsensitive to the detector position and therefore to the order of saturation of the spiral-orbit spectrometer if the slit system does not exist.

PHYSICS RESEARCH

Emilio Segrè in Charge

POLARIZATION EXPERIMENTS ON p-p SCATTERING

James E. Simmons, James Foote, Richard C. Weingart,
John A. Baldwin and Rudolf Larsen

An experiment has been prepared in which one tries to observe the change of the longitudinal (parallel to the beam) spin component on scattering. This is done by changing the relative orientation of the spin and momentum of the protons in a known fashion by the use of a magnetic field. The result of this experiment will be a new independent parameter (A in Wolfenstein's rotation) useful for the determination of the p-p scattering matrix, and probably useful also for making a choice among the sets of phase shifts in p-p scattering that are now being calculated on the Maniac.

The apparatus has been designed and is in an advanced state of construction.

CALCULATIONS ON p-p SCATTERING PHASE SHIFTS

Henry P. Stapp, Tom Ypsilantis

The calculations have been actively pursued with the Maniac machine at Los Alamos. There are at present three or four possible solutions, and we are trying to further reduce them by the A experiment (see above) by correlation with $p + p \rightarrow \pi + d$ experiments and by study of the lower-energy data. A complete report will be given later.

PROTON-NEUTRON TRIPLE SCATTERING

John A. Baldwin and David L. Fischer

Triple-scattering experiments have been performed with a liquid deuterium target considered to contain free neutrons and protons. Wolfenstein's R and D parameters were measured at various angles. The results are being prepared for publication.

SEARCH FOR NEGATIVE PROTONS

Owen Chamberlain, Gerson Goldhaber, Emilio Segrè, and Clyde Wiegand*

We have two plans for searching for negative protons. The first is based on the use of photographic emulsions, as follows.

The proton beam at about 6.2 Bev strikes a copper target. The hypothetical p^- in the backward direction in the c.m. system comes out in the lab system in the forward direction with an energy of about 500 Mev.

¹ L. Wolfenstein, Phys. Rev. 96, 1654, (1954)

* This work was done in conjunction with W. W. Chupp.

The negative particles are deviated by the Bevatron field and come out of the tank in a region inside the shield where they are further analyzed for momentum by a steering magnet (14,000 gauss over 60-inch length). Immediately after having passed through the steering magnet they are focused by a set of 3 magnetic quadrupole lenses.

This whole system has been calculated on paper and carefully lined up with the help of a current-carrying wire under tension.

The momentum of the particles coming out was 1090 Mev/c, to 1% or 2% accuracy. About 200 π^- mesons per pulse per square inch were found in the image of the target source.

The p^- of a given momentum have also a definite range, and we have put a copper absorber and a stack of emulsions at the image in such a way that the p^- should stop in a known region of the stack. The total volume of the stack is 7 by 8 by 1.5 inches, and half of it is being scanned here, and half in Rome (Italy) under the supervision of Professor E. Amaldi.

A variation of this experiment has also been done: at the image focus of the quadrupole we have located a second analyzing magnet ($H=18,000$ gauss) preceded by an absorber. The particles coming to the focus of the lens all have the same momentum, but on passing the absorber the p^- and π^- undergo different momentum loss and they enter the second analyzing magnet with different momenta. Hence at least partial separation between π^- and p^- is possible. Also, in this experiment, the particles are stopped in emulsions.

In the scanning we will be looking for stars which can be ascribed to antiprotons, i. e., in which there is a low-energy entering charged particle liberating visible energies of 0.5 to 2 Bev.

The second experiment, wholly electronic, is just like the first one up to the stack of plates. The stack is replaced by two ordinary counters (in coincidence) and a Cerenkov counter sensitive to particles with $\beta = 1$ in anti-coincidence; this system is followed by suitable quadrupole lenses and at 40 feet from these counters the system is terminated by an ordinary scintillation counter and a Cerenkov counter sensitive to $\beta = 0.75 \pm 0.03$. This whole complex system should give the momentum and the velocity of the particles crossing the apparatus. Indeed, the velocity is measured by the last Cerenkov counter and by a time-of-flight method.

This second experiment is ready but has not yet been performed.

INTERACTIONS OF NEGATIVE K PARTICLES IN FLIGHT¹

Gerson Goldhaber, Ehsan Helmy, Edwin L. Illoff, Joseph E. Lannutti,
and Francis H. Webb²

To date more than 7 meters of track length has been scanned along negative K particles. The detailed analysis of the interactions is still in

¹ UCRL-3039, to be published in the Proceedings of the Pisa Conference (1955).

² This work was done in conjunction with Sulamith Goldhaber.

progress.

In contrast to the positive K particles, the negative K particles occasionally give rise to stars in which part of the rest energy of the negative K meson appears. In some of the interactions charged hyperons and π mesons are emitted. The cross section for interaction is about the geometric cross section which is to be compared with $1/3$ the geometric cross section for positive K mesons. The features mentioned above are all in agreement with conservation of "strangeness".

INTERACTIONS OF POSITIVE K PARTICLES IN FLIGHT¹

Gerson Goldhaber, Ehson Helmy, Edwin L. Iloff, and Joseph E. Lannutti²

To date more than 40 meters of track length has been scanned in nuclear emulsions on positive K particles. The detailed analysis of the interactions in flight is in progress. In the first 27 meters the events found were:

Elastic scatterings (Coulomb scattering subtracted)	9
Inelastic scatterings	16
Charge exchange	2

The striking features of these results are (a) they are in accord with the conservation of "strangeness", (b) the cross section is about $1/3$ of the geometric cross section.

THE MEAN LIFE OF POSITIVE K MESONS³

Gerson Goldhaber, Joseph E. Lannutti and Edwin L. Iloff⁴

The mean life of positive K mesons was obtained by scanning along the track of K mesons in nuclear emulsion exposed to the focused K^+ beam, and observing decays in flight.

On the basis of 19 decays in flight a mean life of $1.01^{+0.33}_{-0.21} \times 10^{-8}$ sec was obtained.

¹ W. W. Chupp, G. Goldhaber, S. Goldhaber, E. I. Iloff, J. E. Lannutti, A. Pevsner, and D. Ritson, UCRL-3021. To be published in the Proceedings of the Pisa Conference (1955).

² This work has been done in conjunction with W. W. Chupp and S. Goldhaber and also with A. Pevsner and D. Ritson of M. I. T.

³ E. L. Iloff, W. W. Chupp, G. Goldhaber, S. Goldhaber, J. E. Lannutti, A. Pevsner, and D. Ritson, UCRL-3064. To be published in Phys. Rev. Sept. 1, 1955.

⁴ This work has been done in conjunction with W. W. Chupp and S. Goldhaber and also with A. Pevsner and D. Ritson of M. I. T.

184-INCH CYCLOTRON

Robert L. Thornton in Charge

MODIFICATION

Richard Burleigh

Magnet

Tests are completed on soft-soldered and clamped joints for connecting the auxiliary coils to the terminal boxes. The shop is proceeding on the final joints.

The connections for grounding the auxiliary coil clamp bands are being installed.

Magnet shim-handling devices are being designed.

It has now been decided that pole-face windings should be installed at the time of the modification. Conductor has been ordered for these and detail design is proceeding.

Radiofrequency System

The two-pair vibrating blade system and new stators have been installed. In order to obtain full amplitude it was necessary to run the blades in helium. Preliminary radiofrequency tests indicate no major difficulties with this system.

The machining of the four new blades at Mare Island is almost complete. Nitriding and copper plating still remain to be done.

Miscellaneous

The drawings for the graphite beam scraper, the meson target, the meson cave slot shielding, and field measuring equipment are almost complete and will be issued shortly.

The iron shielding blocks for the meson cave are complete.

The shutdown date is now firmly set for September 12, 1955.

60-INCH CYCLOTRON

Joseph G. Hamilton in Charge

MODIFICATION

G. Bernard Rossi

Frequency Shifting

The 60-inch cyclotron was shut down for a two-week period starting June 1, 1955. The purpose of this shutdown was to modify the existing rf system to permit a quick and easy variation of the operating frequency. The

need for such frequency variation is apparent from Fig. 1, showing the resonance limitations for heavy-ion acceleration. The curve at 12.00 megacycles represents those ions that can be accelerated within the magnetic field range of 15,800 to 19,700 gauss. To accelerate ions whose e/m is less than 0.42 it becomes necessary to shift the frequency to some lower value. The frequency of 10.70 megacycles was chosen to encompass $^{16}_8\text{O}$, whose $e/m = 0.375$.

A simple device to accomplish this shift was developed in this laboratory and pretested on a one-quarter-scale model of the rf system. In principle, a portion of the existing dee stem was paralleled with a grounded stub line, permitting variations in electrical length of the resonant dee — dee-stem system. This extension of the dee stem is projected through the dee-stem tank by means of a vacuumtight alumina insulator capable of withstanding high rf voltages. Figure 2 is a layout of the stub sections, indicating the position of the external and internal shorting spiders. Operation of the full-scale machine has shown a close correlation to model measurements.

In operation, the low frequency is set by adjustment of the internal spider with the external spiders removed from the stubs. Variations of this setting can be made external to the vacuum by shortening the stub to raise the frequency or adding capacity to lower it. The addition of the external spiders to the stub permits higher frequency of operation over a range determined by the stub length and spider position.

With the present 30-inch stubs a low-frequency range of 10.7 to 10.82 megacycles is possible; the high-frequency range extends from 11.9 to 12.4 megacycles. The low-frequency range (used with the acceleration of heavy ions -- carbon to neon) corresponds to an energy variation of 9.5 to 9.75 Mev per nucleon; the high-frequency range (used with ions for hydrogen to fluorine) extends from 11.7 to 12.9 Mev per nucleon. Larger variations can be achieved by changes in the length of the stub-line.

The oscillator characteristics have proven fairly stable during shifts from 12.0 to 10.8 megacycles. Operation at either frequency requires only minor adjustment of the oscillator parameters, namely, a shift of the tube to dee-stem impedance match and a change in drive from dee stem to the grid of the tube.

A convenient scheme was developed to accomplish the external shorting of the stubs. A set of annular wedges and mating annular inclines contacts the inner and expands to the outer conductor with motion of four driving screws. This mechanism permits removal of the short or its motion along the stubs. Work with this unit indicates that a resetting of the energy to about 1% accuracy is feasible.

The rf model testing was done in this laboratory by Kenneth D. Jenkins and Robert Druet. The design of the full-scale stub and shorting system was accomplished under the direction of Charles Corum of our staff with the assistance of William Baker and Jack Riedel of the Radiation Laboratory Electronics Group.

HEAVY ION RESONANCE PARAMETERS

MU-9822

Fig. 1. Heavy-ion resonance parameters.

60" CYCLOTRON FREQUENCY SHIFTING SYSTEM

MU-9951

Fig. 2. 60-inch Cyclotron frequency-shifting system.

ΔR Measurements

In an effort to more fully understand the operation of the machine, some initial steps have been taken to determine ΔR orbital positions. In past reports (UCRL-2805) references have been made to the "burned foil" technique developed to determine the beam envelope. The present work is an effort to locate individual orbits within this envelope. To accomplish this, a 1/32-inch collimating slit was provided in a water-cooled internal probe target. Its position was chosen to permit only a small portion of the probe beam to continue beyond the radius for which the probe was set. A thin wire was stretched from the collimator, along the median geometric plane, to intercept a portion of the beam as it increased in radius. Figure 3 illustrates the method of irradiating the wire. The darkened spots along the wire indicate points of activation resulting from bombarding the wire. A radioautograph taken after such a bombardment is shown in Fig. 4. Forty-eight-Mev alpha particles were used with the internal collimator set at the 15-inch radius. Twelve distinct orbits can be discerned between the 15-inch and 19-inch radii. Rough measurements indicate $\Delta R_{15} = 7.5$ millimeters; $\Delta R_{19} = 6$ millimeters. Efforts are being made to increase the resolution at larger radii so that a more complete history of the particle trajectories from ion source to extraction can be determined.

OPERATION

Summary of operations as prepared by W. B. Jones for this quarter:

Alpha bombardments	429.6 hr
Deuteron bombardments	53.2 hr
Proton bombardments	200.2 hr
Carbon bombardments	30.6 hr
Oxygen bombardments	97.1 hr
Experimental bombardments	54.4 hr
Operations total	865.1 hr
Frequency-shifting system	162.0 hr
Available time	1027.1 hr
Outage time	108.9 hr
Shutdown	1040.0 hr
Holidays	32.0 hr
Total time	2208.0 hr
Operating efficiency: 84.2 %

DELTA R MONITORING DEVICE

MU-10093

Fig. 3. ΔR monitoring device

ZN-1365

Fig. 4. Radioautograph of ΔR .

LINEAR ACCELERATOR OPERATION

James D. Gow in Charge

The 32-Mev linear accelerator operated for a total of 909 hours in the period May-June-July. The average image was 91%. No major modifications were made. No nonscheduled Van de Graaff shutdowns were experienced in this period. The installation of a new oscillator cabinet to permit operation of two extra 3 W 10,000 oscillators was planned. A scheduled shutdown for this installation will be made in August. The principal research consisted of scattering and energy-level experiments. Some runs were made using the β -ray spectrometer.

SYNCHROTRON

Edwin M. McMillan in Charge

RESEARCH PROGRAM

Robert W. Kenney

The synchrotron beam intensity has improved steadily since the January shutdown and is now running at a satisfactory level.

Larry Higgins and Kenneth Bandtel have made preliminary runs on the scattering of photons by hydrogen below the meson threshold. Fast-neutron background has been difficult to shield against, and it appears that additional modifications of the counter telescope are required. Cerenkov counters will probably be used to detect fast electrons in the presence of the fast-neutron background. Some data on hydrogen have been taken but no results are available at present.

Dwight Dixon, Robert Cence, and Kenneth Bandtel have examined the possibility of observing the reaction $\gamma + D \rightarrow p + n$ at 0° proton angle, and it appears to be a difficult though feasible experiment. Poor synchrotron operation prevented the efficient gathering of additional data at wide angles.

William Imhof, Edward Knapp, and Victor Perez-Mendez have improved their statistics on the positive photopion production as a function of Z at 135° in the lab. Their data at 36 Mev and 46 Mev pion energy indicate a cross-section dependence between Z and $Z/A^{1/3}$ from which they derive a pion mean free path, in nuclear matter, of approximately $3 r_0$. Isotope effects in photopion production from B^{10} and B^{11} have been investigated and appear to be too small to be detected easily.

John D. Anderson and Robert Kenney obtained preliminary data on the total cross section for neutral photopion production as a function of A at full energy. The cross section varies as $A^{2/3}$, as expected. At photon energies near threshold, the dependence of the cross section is more nearly proportional to A . The work is continuing toward determining the π^0 mean free path in nuclear matter as a function of pion energy.

Dr. Ryokinchi Sagane, Walter Dudziak, and James Vedder calibrated the scintillation counters, used in the μ -electron decay-spectrum work, by utilizing a high-energy electron beam from the pair spectrometer magnet.

Robert Squire calibrated the efficiency of a photon counter, using electrons from the pair spectrometer magnet.

Herbert Steiner exposed some nuclear emulsions for an experiment on photofission. No data are yet available.

Dr. John Teem of Cal Tech, who visited the laboratory during the week of July 25, investigated the response of a liquid hydrogen bubble chamber which was placed directly into the synchrotron beam. The bremsstrahlung spectrum was altered by selective absorption of low-energy photons by a LiH absorber 60 inches in length placed in a magnetic field. The electron background due to Compton recoils was appreciably reduced by this means and it appears feasible to operate a liquid hydrogen bubble chamber in this manner if the synchrotron beam intensity does not exceed approximately one-half its present maximum value. This information will bear directly upon the proposed Cal Tech bubble chamber program. This work was done in cooperation with the Berkeley bubble chamber group.

OPERATION

Rudin M. Johnson

During the period of May, June, and July, the synchrotron was used entirely for physics research experiments. The beam intensity at full energy was about one-fifth of our previous maximum. Line voltage variations from the Bevatron operation still cause the synchrotron beam to be unstable. The synchrotron is being operated on owl shifts whenever possible so as to take advantage of the Bevatron off time. This is possible because we have full-time, experienced graduate student operators during the summer.

An improvement in beam intensity has been noticed when the synchrotron is running with the auxiliary experimental magnets, and this stray-field effect is being investigated as a possible method of improving the synchrotron beam intensity.

Ceramic doughnut sections to be used with the linear accelerator injector have arrived and are being machined and vacuum tested. These ceramic sections are of excellent quality.

Preliminary injection tests have been made into the 120° magnet section with the 2-Mev linear accelerator. Approximately 1 milliamperes of beam was measured at the orbit.

The mechanical and electrical design for the synchrotron linac Model II are nearly complete and parts are arriving from the shops.

The FCC has complained about rf radiation from the synchrotron resonator oscillator. Our tests indicate the quartz resonator is the source of radiation

and steps are being taken to eliminate this radiation.

Information Division
9-14-55 jm

McMillan