LA-5411-MS

Dr. 397

INFORMAL REPORT

**Sensitivity of Fusion Reactor Average
Cross Sections to Thermal Broadening of the
14-MeV Neutron Peak**

**los alamos
scientific laboratory**

of the University of California

LOS ALAMOS, NEW MEXICO 87544

UNITED STATES
ATOMIC ENERGY COMMISSION
CONTRACT W-7405-ENG. 36

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

In the interest of prompt distribution, this LAMS report was not edited by the Technical Information staff.

Printed in the United States of America. Available from
National Technical Information Service
U. S. Department of Commerce
5285 Port Royal Road
Springfield, Virginia 22151
Price: Printed Copy \$4.00; Microfiche \$0.95

LA-5411-MS
Informal Report
UC-20

ISSUED: October 1973

Sensitivity of Fusion Reactor Average Cross Sections to Thermal Broadening of the 14-MeV Neutron Peak

by

D. W. Muir

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

This work supported by the US Atomic Energy Commission's
Division of Controlled Thermonuclear Research.

MASTER

89

SENSITIVITY OF FUSION REACTOR AVERAGE CROSS SECTIONS TO THERMAL BROADENING OF THE 14-MeV NEUTRON PEAK

by

D. W. Muir

ABSTRACT

Ion motion in a deuterium-tritium thermonuclear plasma can cause significant broadening of the 14-MeV peak in the fusion neutron energy spectrum. The shape of the peak is shown to be nearly Gaussian; the full width at half maximum FW_{DT} being related to the plasma thermal energy kT by the relation $FW_{DT} = 5.59 \sqrt{kT}$, where both FW_{DT} and kT are expressed in MeV. Numerical examples related to fusion reactor technology illustrate the sensitivity of spectrum-averaged cross sections to the plasma temperature.

I. INTRODUCTION

In recent years numerous studies have explored the possible characteristics of controlled fusion reactors.¹ Most of the concepts discussed for the first generation of these reactors are based on the D-T cycle. In this cycle, the plasma consists of an equimolar mixture of deuterium and tritium, and the reaction $T(d,n)^4He$ accounts for nearly all of the energy production. An important part of these fusion reactor studies is the calculation of the transport of the neutrons released in the D-T reaction throughout the material regions adjacent to the plasma. In addition to the space-energy distribution of the transported neutrons, one needs to know the nuclear cross sections for the production of important neutron effects, such as tritium breeding, energy deposition, gamma-ray production, activation, transmutation, helium production, and others. The cross sections for the processes listed above often exhibit threshold behavior near 14.1 MeV, the nominal energy of the source neutrons, so that it is important to know the detailed shape of the energy spectrum near the peak.

As noted by Nagle et al.,² the thermal motion of the interacting ions in a fusion plasma can impart a considerable energy spread to otherwise monoenergetic reaction products. For a stationary plasma

with a Maxwellian distribution of ion velocities, Lehner³ has shown that the energy distribution of the products is nearly Gaussian. In the following, we present a somewhat more transparent derivation of this Gaussian shape, and we present numerical examples which illustrate the sensitivity of threshold neutron reaction rates to the temperature of a D-T fusion plasma.

II. THE GAUSSIAN DISTRIBUTION

The probability that an ion with mass m_1 will have a velocity in the range $(\vec{v}_1, \vec{v}_1 + d\vec{v}_1)$ and that another ion with mass m_2 will have a velocity in the range $(\vec{v}_2, \vec{v}_2 + d\vec{v}_2)$ is just the product of two independent Maxwell distributions. This probability product can be recast in terms of the relative velocity of $\vec{v}$ and the center-of-mass velocity $\vec{V}$ of the two-ion system. Somewhat surprisingly, the resulting distribution $N(\vec{V}, \vec{v})$ is also the product of two independent Maxwell distributions.⁴

$$N(\vec{V}, \vec{v}) d^3V d^3v = P(\vec{V}) d^3V \cdot p(\vec{v}) d^3v \quad (1)$$

where

$$P(\vec{V}) = \left[\frac{m_1 + m_2}{2\pi kT} \right]^{3/2} \exp \left[- \frac{(m_1 + m_2) V^2}{2kT} \right] \quad (2)$$

and

$$p(v) = \left[\left(\frac{m_1 m_2}{m_1 + m_2} \right) \frac{3}{2} \right] \exp \left[- \left(\frac{m_1 m_2}{m_1 + m_2} \right) \frac{v^2}{2kT} \right], \quad (3)$$

where V is the magnitude of $\vec{V}$, v is the magnitude of $\vec{v}$, k is Boltzmann's constant, T is the plasma temperature.

Let us introduce $\sigma(v)$, the cross section for the fusion reaction of interest. The probability distribution $R(\vec{V}, \vec{v})$ for ion pairs which actually undergo a fusion reaction is just

$$R(\vec{V}, \vec{v}) d^3V d^3v = P(V) d^3V \cdot q(v) d^3v, \quad (4)$$

where

$$q(v) = k\sigma(v)vp(v) \quad (5)$$

and where k is chosen such that $\int q(v) d^3v = 1$.

We now proceed to a discussion of those properties of the distribution $R(\vec{V}, \vec{v})$ which affect the shape of the fusion neutron energy spectrum.

The average neutron energy will be increased somewhat because of the kinetic energy E_v associated with the relative motion of the reacting ions. A calculation of the average value of E_v will require a numerical integration over the non-analytic function $q(v)$, Eq. (5). However, this contribution (typically a few times kT) is much smaller than the nuclear energy release or Q -value (17.590 MeV in the case of the D-T reaction). Thus, for plasma temperatures in the range of 10-50 keV, it is adequate to set the average contribution equal to a constant, say 50 keV, and to lump it together with the nuclear Q -value. This procedure results in a center-of-mass neutron energy of 14.07 MeV for the D-T reaction. In addition to this effect, which slightly shifts the location of the peak, there will be a broadening of the spectrum due to the distribution $q(v)$ of relative velocities. As shown in the following, this broadening is negligible in comparison with the effect of the distribution $P(V)$ of center-of-mass velocities.

In order to evaluate the effect of $P(V)$, we return to the expression for the probability distribution $R(\vec{V}, \vec{v})$ given in Eq. (4). Since the distribution of relative ion velocities $q(v)$ is isotropic, we can assume, without loss of generality, that the product neutron is emitted along the z -axis in the center-of-

mass system. Let us introduce the symbol E_0 for the center-of-mass energy of the neutron and v_0 for the associated velocity. Then

$$E_0 = \frac{1}{2} m v_0^2,$$

m being the mass of the neutron. The energy E of the neutron in the laboratory system is

$$E = \frac{1}{2} m \left[(v_0 + v_z)^2 + v_x^2 + v_y^2 \right] \\ E \approx \frac{1}{2} m \left[v_0^2 + 2v_0 v_z \right] = E_0 + m v_0 v_z. \quad (6)$$

Following the work of Bell,⁵ we have used the fact that the neutron speed v_0 is much greater than the center-of-mass speed V . From this last result, we have

$$v_z = \frac{E - E_0}{m v_0}$$

$$dv_z = \frac{dE}{m v_0}.$$

We denote by $S(E)$ the desired energy distribution. From Eq. (6), to first order E is a function only V_z , so that

$$S(E) dE = P_z(V_z) dv_z.$$

P_z is obtained by factoring $P(V)$ from Eq. (2), which gives

$$S(E) = \frac{1}{m v_0} \left(\frac{m_1 + m_2}{2\pi kT} \right)^{1/2} \exp \left[- \frac{(m_1 + m_2)(E - E_0)^2}{2m^2 v_0^2 kT} \right]$$

$$S(E) = \frac{1}{W\sqrt{\pi}} \exp \left[- \left(\frac{E - E_0}{W} \right)^2 \right] \quad (7)$$

where

$$W = \left(\frac{4mE_0 kT}{m_1 + m_2} \right)^{1/2}. \quad (8)$$

Thus, $S(E)$ is a normalized Gaussian distribution. The full width at half maximum FW is related to W by

$$FW = 2W\sqrt{\ln 2}. \quad (9)$$

For the D-T reaction $(m_1 + m_2)/m \approx 5$ and $E_0 = 14.07$ MeV. From Eqs. (8) and (9), then

$$FW_{DT}(\text{MeV}) = 1.665 W_{DT}(\text{MeV}) = 5.59 \sqrt{kT(\text{MeV})}. \quad (10)$$

The width of this distribution is much larger than kT for most cases of interest. For example, if $kT = 20$ keV, the width of the 14-MeV neutron peak is 0.79 MeV, or a factor of forty greater than kT .

III. THRESHOLD CROSS SECTIONS WITH LINEAR ENERGY DEPENDENCE

The simple analytic nature of the function $S(E)$ simplifies the task of cross-section averaging in certain cases. For example, we consider the case of a cross section which rises linearly above a threshold energy ϵ

$$\sigma(E) = \begin{cases} C(E - \epsilon), & E > \epsilon \\ 0, & E \leq \epsilon \end{cases}.$$

This shape fits the gross features of many $(n,2n)$ and $(n,3n)$ reactions, two examples of which are discussed in the following section. The average cross section for fusion-born neutrons can be written as follows

$$\sigma_{av} \equiv \int_{\epsilon}^{\infty} \sigma(E) S(E) dE = C \int_{\epsilon}^{\infty} (E - \epsilon) S(E) dE.$$

From Eq. (7), then

$$\sigma_{av} = \frac{1}{W\sqrt{\pi}} \int_{\epsilon}^{\infty} C(E - \epsilon) \exp \left[-\left(\frac{E - E_0}{W} \right)^2 \right] dE. \quad (12)$$

For convenience we introduce the parameter

$$t = \frac{\sqrt{2}}{W} (E - E_0). \quad (13)$$

We also introduce the tabulated function $G(t)$, which is the area under the normal curve of error.⁶

$$G(t) \equiv \frac{1}{\sqrt{2\pi}} \int_0^t e^{-x^2/2} dx \quad (14)$$

Integration of Eq. (12) then gives

$$\sigma_{av} = \frac{WC}{\sqrt{4\pi}} \left[e^{-t^2/2} - \sqrt{2\pi} t \left[1/2 - G(t) \right] \right]. \quad (15)$$

IV. NUMERICAL EXAMPLES

Consider the nuclear reaction $^{27}\text{Al}(n,2n)^{26m}\text{Al}$. The isomeric state of ^{26}Al decays by positron emission with a half-life of 6.4 sec. The average energy of the positron is about 1.4 MeV, and two 0.51 MeV annihilation photons are produced as well. Although the cross section for this reaction is small (≤ 1 mb),

the short range of the reaction products may lead to a degradation of the performance of the common electrical insulator, Al_2O_3 . The available data for the ^{26m}Al production cross section⁷ is adequately described below 16 MeV by the linear relation $\sigma(E) = C(E - \epsilon)$, where $C = 0.276$ mb/MeV and $\epsilon = 13.77$ MeV, and E is the neutron energy in MeV. Thus the cross section averaging formula, Eq. (15), can be used for this reaction.

Another reaction of interest in fusion reactor technology is $^{181}\text{Ta}(n,3n)^{179}\text{Ta}$. Tantalum has been considered as a possible material for a fusion reactor vacuum "first-wall," because it has a very small (n,α) cross section. This property should make tantalum very resistant to helium-induced embrittlement.¹⁷⁹ Ta production is of interest in that the half-life (600 d) is much longer than that of other activities which might be produced in tantalum structures. The cross section for the $(n,3n)$ reaction is represented in a recent evaluation⁸ as rising linearly from a threshold ϵ at 14.5 MeV. The slope C is 584 mb/MeV.

In order to illustrate the sensitivity of the average cross section to the plasma temperature and to the position of the threshold, we now calculate σ_{av} for the two reactions described above. The approximate average plasma temperatures for a variety of fusion reactor concepts are given in Table I (first column). In the second column are listed the widths of the 14-MeV peak, calculated from Eq. (10). The third column lists spectrum-averaged cross sections for the $^{27}\text{Al}(n,2n)^{26m}\text{Al}$ reaction ($\epsilon = 13.77$ MeV), and the fourth column contains similar information for the $^{181}\text{Ta}(n,3n)^{179}\text{Ta}$ reaction ($\epsilon = 14.50$ MeV). All cross sections shown were calculated using the linear approximation to $\sigma(E)$ given above and the averaging formula, Eq. (15).

Summarizing the results in Table I, we see that for the ^{27}Al case, where the threshold is 300 keV below E_0 (14.07 MeV), the average cross section varies by less than 20% for changes in kT of a factor of four. For the ^{181}Ta case, where the threshold is 430 keV above E_0 , the picture is quite different. Here a factor of ten change in σ_{av} is seen for this factor of four change in kT . It is expected that similar sensitivity will be observed for other high-threshold ($\epsilon > 14$ MeV) reactions.

TABLE I

SPECTRUM-AVERAGED CROSS SECTIONS FOR DIFFERENT PLASMA TEMPERATURES

Reactor Concept	Approximate Average Plasma Temperature (keV)	Width of 14-MeV Peak (fwhm in MeV)	σ_{av} for $^{27}\text{Al}(n,2n)^{26m}\text{Al}$ (μb)	σ_{av} for $^{181}\text{Ta}(n,3n)^{179}\text{Ta}$ (mb)
Wisconsin Tokamak A ⁹	11	0.59	87	2.5
Theta Pinch ¹⁰	12	0.61	87	3.0
Princeton Tokamak ¹	15	0.68	89	5.2
Oak Ridge Tokamak ¹¹	20	0.79	92	9.3
Wisconsin Tokamak B ⁹	28	0.94	97	16.5
Laser-Pellet ¹²	40	1.12	104	27.8
D-T Mirror ¹	400	3.53	210	238.2

ACKNOWLEDGMENTS

The author acknowledges helpful discussions of this problem with G. I. Bell, L. R. Veaser, L. Forman, and R. J. Doyas.

REFERENCES

1. R. L. Hirsch, "Fusion Power: An Assessment of the Ultimate Potential," U. S. Atomic Energy Commission report WASH-1239 (1973).
2. D. E. Nagle, W. E. Quinn, W. B. Riesenfeld, and W. Leland, "Ion Temperature in Scylla, as Determined from the Reaction D(d,p)T," Phys. Rev. Letters **3**, 318 (1959).
3. G. Lehner, "Reaction Rates and Energy Spectra for Nuclear Reactions in High Energy Plasmas," Z. Physik **232**, 174 (1970).
4. Donald D. Clayton, Principles of Stellar Evolution and Nucleosynthesis (McGraw-Hill, Inc., New York, 1968) p. 294.
5. G. I. Bell, Los Alamos Scientific Laboratory, private communication.
6. Mathematical Tables from the Handbook of Chemistry and Physics Eleventh Edition, A. A. Bennett and W. B. Ross, Eds. (Chemical Rubber Publishing Co., Cleveland, 1959), p. 212.
7. D. M. Arnold, "(n,2n) Excitation Functions of Several Low Neutron Excess Nuclei," thesis abstract, Dissertation Abstracts **26**, 3525 (1965).
8. E. H. Ottewitte, J. M. Otter, P. F. Rose, and C. L. Dunford, "An Evaluation of Ta-181 and Ta-182 for the ENDF/B Data File," Atomic International report AI-AEC-12990 (1971).
9. R. W. Conn, D. G. McAlees, and R. Neil, "Neutral Beam Heating and Energy Balance Studies in a CTR Tokamak," Trans. Am. Nucl. Soc., Winter Meeting, 1973.
10. T. A. Goultas, A. J. Hatch, R. A. Krakowski, and F. L. Ribe, "Engineering Design Study of a Reference Theta-Pinch Reactor (RTPR)," Los Alamos Scientific Laboratory report LA-5336 (1973).
11. A. P. Fraas, "Conceptual Design of the Blanket and Shield Region and Related Systems for a Full-Scale Toroidal Fusion Reactor," Oak Ridge National Laboratory report ORNL-TM-3096 (1973).
12. R. L. Morse, Los Alamos Scientific Laboratory, private communication.