LA-UR-7A-1A77
CONF-740558-2

TITLE:

THE IMPORTANCE OF NONUNIFORM DOSE-DISTRIBUTION IN AN ORGAN

AUTHOR(S):

C. R. Richmond

SUBMITTED TO:

To be published in Health Physics, Proceedings of the Second Annual Life Sciences Symposium on "Plutonium -- Health Implications for Man," held at the Los Alamos Scientific Laboratory, Los Alamos, New Mexico (May 22-24, 1974)

By acceptance of this article for publication, the publisher recognizes the Government's (license) rights in any copyright and the Government and its authorized representatives have unrestricted right to reproduce in whole or in part said article under any copyright secured by the publisher.

The Los Alamos Scientific Laboratory requests that the publisher identify this article as work performed under the auspices of the U. S. Atomic Energy Commission.

los alamos
scientific laboratory
of the University of California
LOS ALAMOS, NEW MEXICO 87544

NOTICE
This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

UNITED STATES
ATOMIC ENERGY COMMISSION
CONTRACT W-7405-ENG. 36

July

THE IMPORTANCE OF NONUNIFORM DOSE-DISTRIBUTION IN AN ORGAN

Running title: Nonuniform Dose-Distribution

C. R. Richmond

Health Division, Los Alamos Scientific Laboratory

University of California, Los Alamos, New Mexico 87544

ABSTRACT—The recent revival of interest in the "hot particle" problem, especially as regards particulate plutonium and other actinide elements in the lung, stimulated the preparation of this paper. Nonuniformity of dose-distribution has been of concern to standards-setting bodies and other groups such as the National Academy of Sciences and to health protectionists for many years. This paper reviews data from animal experiments that are used by some to implicate particulate plutonium as being especially hazardous to man. Other relevant biological data are also discussed.

The basis of the ICRP recommendations for derived radiation protection standards is the average radiation dose to an organ and not the number of particles in the organ. This dosimetric basis for radiological protection has been in effect for many years and is based on observations from humans and experimental work with animals. Recently proposed risk estimates for plutonium-induced lung cancer of 5×10^{-4} per particle cannot be substantiated on the basis of our current knowledge. In fact, careful consideration of the experimental data shows that plutonium is not more hazardous than the same amount of plutonium distributed uniformly. Further, the data suggest that the potential hazard from plutonium increases as the dispersion throughout the lung becomes more uniform.

Mr. Chairman, ladies, and gentlemen: It is my pleasure to discuss with you this morning the subject of the importance of nonuniform dose-distribution in an organ. This is indeed a timely subject but, as I will indicate later, one which is not new but has been with us for a long time. I will restrict most of my comments to plutonium in the lung, as most of us are more familiar with the subject of nonuniform distribution of dose in skeletal tissues, a topic which was covered yesterday in the paper prepared by Jack Healy and discussed by others.

Permissible values for the respiratory intake of radioactive materials are commonly calculated on the assumption of complete absorption of the radiation energy by the critical organ. Further, it is implicitly assumed that there is a uniform distribution of the energy per gram of tissue throughout the critical organ. This particular situation raises the interesting question as to the possibility of a unique hazard to the respiratory tissues for a given amount of radioactive material distributed in the form of a relatively small number of discrete radioactive particles as compared with a more homogeneous distribution. Stated in another way: for the same amount of material, is the biological harm to the lung greater or less if the energy is concentrated into very small tissue volumes as compared to the case in which the energy is absorbed by the entire organ? For alpha and some beta radiations, the distribution of energy will be nonuniform and consequently concentrated about the particles as foci, thereby producing intense radiation doses to the cells nearby. For the case of nonuniform distribution of alpha-emitting materials in the lung, the initial biological interaction is that of an extremely large energy deposition to a very small tissue volume. As I reported earlier (RICHMOND *et al.*, 1970), two problems arise immediately. The first is to determine what "radiation dose" (e.g.,

mean organ, at some distance or to some volume) is significant for the particular exposure, and the second is to assess the biological insult not only in terms of some "radiation dose" but also in terms of early and late effects on both the tissue involved and on the entire organism. We need to continually remind ourselves and others that radiation dose is a convenient means of providing a framework for relating effects observed among organs, species, and test systems, but it should not be considered more important than the effect. The ultimate misuse of "radiation dose" is the not too unfamiliar situation in which someone calculates dose to a critical tissue volume, speculates as to the biological harm, and recommends that protection standards be adjusted (always downward) because some previously accepted radiation dose limit will be exceeded according to the new model.

I have previously considered the case in which the functional lung of rodents and dogs was exposed to one or several highly radioactive alpha-emitting particles (RICHMOND *et al.*, 1970; RICHMOND *et al.*, 1974). In such a situation, the "organ mean dose" concept can be inadequate for the purposes of radiation protection in that the primary radiation dose equivalent rate may be substantially exceeded, even though a miniscule volume of the total organ is irradiated and at risk. In theory, exposures from a very few particles could easily exceed current radiation protection standards proposed by the International Commission on Radiological Protection (ICRP) and other advisory agencies. Please note that this statement does not mean that the current radiation protection guides should be revised downward (made more conservative), as they may indeed be entirely adequate or, in fact, overly conservative for the specific case of extreme non-uniform distribution of energy.

I should also like to point out at the outset that it is abundantly clear that the so-called "hot particle" problem (nonuniform dose-distribution) has not

been recently discovered but has, in fact, been a problem of recurrent interest which has troubled industrial hygienists, health physicists, and radiation biologists since the early 1940's. I shall elaborate further on this point during my discussion. I should also like to state that this problem of non-uniform dose-distribution, particularly as regards inhaled particles, has been of interest to standards-setting organizations such as the ICRP, NCRP, and their numerous committees and subcommittees who have spent much time considering the problem in detail. The problem has also been considered by several select committees formed by the National Academy of Sciences. During recent years, there have been accusations and implications that such prestigious bodies have ignored the problem or that they have provided no guidance. It has also been stated that the current standards for plutonium are a "travesty of public health." As Healy pointed out earlier in this symposium, this is simply not so.

Table 1 summarizes the general classes of nonuniformity of radiation dose as presented in ICRP publication 14 (ICRP, 1969). All three classes of non-uniformity of radiation dose have received and continue to receive considerable amount of attention on the part of the biomedical community and the various national and international standards-setting organizations. Again, I will restrict for the most part my comments to class 3, Irradiation from Radioactive Materials in Particulate Form, and I will restrict most of my comments further to the case in which the lung is the limiting organ.

Let us now look at the activity and number of particles of different geometric diameters that are required to deliver the occupational MPB2 for two nuclides of plutonium. As shown in Table 2, it would take 5.4×10^4 particles, each 1 μm in diameter, but 5.4×10^{10} particles, each 0.01 μm in diameter, to account

for a MPBB of 16 nCi. If the radiation dose is calculated by assuming that the energy from the 16 nCi is absorbed by the entire human lung, the resultant dose rate would be 1.5 rad/yr. Table 2 also indicates that only 200 $^{238}\text{PuO}_2$ particles, each 1 μm in diameter, are required to deliver 1.5 rad/yr to the human lung using the same assumptions. If we now turn to nonoccupational exposures, the number of 1- μm diameter $^{238}\text{PuO}_2$ particles required to deliver the dose limit becomes 20 or 6, depending on whose guidance we choose to follow. Please recall that we are assuming that the energy is absorbed by the entire lung mass which, in the case for humans, is approximately 1000 g. It is quite obvious that the 200 ^{238}Pu particles of the 1- μm diameter size would irradiate an extremely small fraction of the lung and place very few cells at risk. Obviously, this is not the case for the same quantity of energy when it is distributed throughout the entire mass of the organ in question.

Table 3 shows the relationship of particle size to the number of cells at risk for a static lung burden of 16 nCi of $^{239}\text{PuO}_2$. Before we discuss Table 3, I should emphasize several rather important points. First, particles in the lung are very dynamic, and there is a great deal of information to show that in general the static particle situation is quite unlikely (BAIR *et al.*, 1974). Also of great importance is the fact that uniform distribution of radioactive materials in the lung very seldom, if ever, actually occurs in nature. There is a considerable amount of information obtained from autoradiographic studies of lung tissue which strongly indicates an aggregation of inhaled materials. If we direct our attention to the 1- μm diameter particle, we can see that approximately 3.6×10^8 cells, about 0.03% of the lung, are at risk when the lung burden is 16 nCi. However, for the 0.1- μm diameter particle, about 3×10^{11} cells, or about 30%, are at risk. Note that, as the particle size increases, the number of particles decreases as does the number of cells at risk. However,

for the larger particle diameters, the cells are exposed to a larger activity per particle. This then is the heart of the so-called "hot particle" problem. For the same total quantity of activity, is it relatively more harmful for the lung to be exposed to larger alpha-emitting particles in which fewer cells are at risk but to higher dose rates, or to smaller particles in which case more cells are exposed but to lower dose rates?

I might also point out that one would need a single $^{239}\text{PuO}_2$ particle about 36 μm geometric diameter to have an activity of 0.016 μCi . Obviously, this is far in excess of the size that can be deposited and retained in the lung. A $^{239}\text{PuO}_2$ particle with a diameter of 3.6 μm has a small probability of being deposited in the alveolar portion of the lung. It would require 10^3 such particles to contain 0.016 μCi , yet only 10^{-5} of the lung would be irradiated if the particles did not move.

I would now like to discuss some of the experimental data which have been used or misused, as the case may be, as regards the "hot particle" problem. The data of Dr. Roy Albert and his colleagues at New York University, in which skin tumor response to various kinds of radiations was determined for rats, provide the basis for the dose-response portion of those models that have been derived to predict lung tumor incidence for humans exposed to plutonium. In the early experiments, these investigators tried to determine whether isolated small areas of irradiated rat skin would give the same tumor yield per unit as large-area irradiations (ALBERT *et al.*, 1961; ALBERT *et al.*, 1967a,b,c). For low LET radiation (electrons), the focal irradiation pattern was less efficient in producing skin tumors as was the large-area exposures (ALBERT *et al.*, 1967b). However, when these investigators used high LET radiation (protons), no significant differences in tumor yield were observed (BURNS *et al.*, 1972). It must

be kept in mind that these experiments are not entirely relevant to the "hot particle" problem because of the large volumes of tissues irradiated.

Table 4 shows the volume of tissue irradiated for both beta and proton exposures which involved either a large (25 cm^2) area of the skin or the same area subdivided into isolated slots and pores by using grid or sieve masking plates. If one assumes an average depth of 1 mm for the irradiated fields, the calculated tissue volumes exposed to these irradiations varied from $0.2\text{--}2500 \text{ mm}^3$. For comparison, a single ^{239}Pu particle would irradiate a volume of unit density tissue of approximately $2.7 \times 10^{-4} \text{ mm}^3$, which is considerably smaller than the volume used in the Albert studies by factors of $10^{-7}\text{--}10^{-3}$. This is an important point, as the Albert rat skin dose-response data (ALBERT *et al.*, 1967a) are the biological basis for models developed by GEESAMAN (1968) and used by others (TAMPLIN and COCHRAN, 1974) to argue for the occurrence of lung tumor probabilities of the order of $10^{-4}\text{--}10^{-3}$ for "hot particles." It is also of interest to note that the dose-response curve reported for electrons by ALBERT *et al.* (1967a) could not be repeated in other strains of rats or in mice (ALBERT *et al.*, 1961; ALBERT *et al.*, 1972).

These studies of electron radiation of varying energy and penetrating power and the relationship between the occurrence of skin tumors and atrophic hair follicles in the rat suggested the presence of target cells at a depth of about 0.3 mm which corresponded to the lower end of the resting hair follicle (ALBERT *et al.*, 1967a). The critical depth remained constant even when the skin was irradiated with the hair in the growing phase, at which time the follicles extend to a depth of 0.8 mm (BURNS *et al.*, 1973). There appeared to be a quantitative association between the existence of skin tumors and atrophic hair follicles for various kinds of radiation, various spatial distributions of dose in the skin, and for different phases of hair growth. A possible explanation for the

experimental results was that each follicle had a population of stem cells at a depth of approximately 0.3 mm below the surface that are concerned with the production of sebaceous cells in hair. Because the tumors were primarily of hair follicle origin (ALBERT *et al.*, 1969), the stem cells appeared to constitute the most sensitive, potentially oncogenic population of cells in the rat skin. Neoplastic transformation of a significant number of these target cells required large radiation doses which, in turn, killed most of the target cells and thus caused follicle atrophy.

Other experimental results from Albert's laboratory, as shown in Fig. 1, involved the production of skin tumors and chronic follicle damage in the rat as determined following graded single doses of 37-MeV alpha-particle radiation in two depth-dose patterns. In the first, the Bragg peak was placed at each of three depths in the rat skin corresponding to regions above, below, and at the tips of the resting hair follicle. In the second, the alpha beam was modified to produce a linear depth-dose pattern. Dissimilar dose-response curves for tumors and atrophic follicles were obtained for the various depth-dose patterns when the radiation dose was expressed in terms of the maximum dose to either the skin or the follicles or as the average follicle dose. Alpha irradiations extending from the skin to a depth of about 0.15 mm did not produce tumors (HEIMBACH *et al.*, 1969). However, this observation was consistent with the proposed existence of a target cell population at a depth of about 0.3 mm. Selective irradiation of the lower end of the hair follicle at a depth of about 0.3 mm by using the Bragg peak from the alpha beam did not produce tumors or atrophic hair follicles unless there was a substantial irradiation of the entire hair follicle. These results suggest that the production of chronic hair follicle injury and tumor induction depends on producing radiation damage to the entire hair

follicle and that the minimally damaged region, regardless of its location, controls the overall response.

Thus, we can summarize these extremely interesting experiments as follows:

(a) large volumes of skin were irradiated; (b) the assumed critical structure plus other volumes of tissue must be irradiated before correlations are seen in follicle damage and tumor formation; and (c) it is probably not proper to use dose-response data obtained from rat skin as input data for models designed to predict lung tumor response for human subjects. The choreography referred to yesterday by Dr. Brues applies to the latter statement above.

Another interesting experiment, reported by PASSONNEAU *et al.* (1952), involved the use of ^{90}Sr beta-particle radiation on rat skin. Table 5 shows the results of these experiments. The same quantity of activity was used to irradiate the same area of skin; however, the activity was distributed either as a uniform flat plate, in 50 beads, 20 beads, or as 10 beads. These data indicate a decrease in the tumor production efficiency as the radioactivity became concentrated in fewer point sources, irradiating a smaller total tissue volume. The beads with the most radioactivity produced the largest number of tumors per bead and the smallest number of tumors per microcurie. However, the relevant parameter would appear to be the number of tumors produced per microcurie, as the basic question as regards the "hot particle" problem is how their potential hazard compares with that of uniformly distributed radiation dose. As stated recently by TAMPLIN and COCHRAN (1974), "A fundamental question is then: is this intense but localized irradiation more or less carcinogenic than uniform irradiation?"

I have purposefully included the fifth column in Table 5 as it was not included (although it was discussed) in the original data presented by PASSONNEAU *et al.* (1952). The NRDC recently used this observation to argue that, despite the fact that the relative efficiency for tumor formation decreased as the 1500 μCi

activity was condensed into fewer particles, the number of tumors per bead actually increased. I personally think this is a red herring and is not relevant to the "hot particle" problem as actually indicated by TAMPLIN and COCHRAN (1974).

Other experiments relevant to the question of the biological effects of "hot particles" as compared with uniform dose-distribution were performed by Dr. Little and co-workers at Harvard University. These workers studied the effects of ^{210}Po absorbed onto hematite (ferric oxide) particles in Syrian golden hamsters following intratracheal instillation (LITTLE *et al.*, 1970a,b; GROSSMAN *et al.*, 1971; LITTLE *et al.*, 1973). The animals were given 15 weekly injections, each containing 3 mg of hematite and either 0, 0.01, or 0.2 μCi of ^{210}Po . Mean radiation doses calculated for the entire lung were 225 and 4500 rad, respectively, at the end of one year (LITTLE *et al.*, 1970a). The earliest and highest incidence of pulmonary tumors occurred in those hamsters receiving the highest dose of ^{210}Po bound to the hematite. In the group given 0.2 μCi of ^{210}Po on hematite, the first lung cancer appeared in an animal sacrificed 15 weeks after administration. This particular experiment established that the hamster was a useful model system for studying lung cancer production by alpha irradiation but did not compare the relative effectiveness of uniform versus nonuniform distribution. The hematite particles were such that 98% were smaller than 0.75 mm diameter. The first lung carcinoma occurred in an animal that died 7 weeks after the start of treatment.

In another experiment, four groups of 50 hamsters each were given separate intratracheal instillations twice per week for 7 weeks (GROSSMAN *et al.*, 1971). The hamsters received one of the following treatments: 3 mg hematite followed by 0.2 μCi ^{210}Po in saline; saline, then 0.2 μCi ^{210}Po alone; saline, then 0.2 μCi ^{210}Po on 3 mg hematite; saline, then 0.2 μCi ^{210}Po on 0.3 mg hematite.

The distribution of the ^{210}Po in the lung was found by LITTLE *et al.* (1973) to be very nonuniform for the hematite-bound ^{210}Po as compared with the ^{210}Po administered alone in saline (Fig. 2). The results suggested that alpha radiation might be more carcinogenic when it is uniformly distributed throughout the lung.

Additional experiments performed by the Harvard group yielded further information relevant to the importance of nonuniform dose-distribution in the induction of lung cancer by alpha radiation. When hamsters were given seven weekly injections of 0.2 μCi of ^{210}Po alone in saline, the cumulative radiation dose to the lung was about 800 rad as compared with about 2000 rad when the same amount of activity was absorbed on either 3 or 0.3 mg of hematite particles. Mean tumor induction time was shorter for the hamsters given ^{210}Po in saline, and the tumor incidence was lowest for the animals with the most nonuniform dose-distribution (LITTLE *et al.*, 1973).

The major differences among the experimental groups was in the microscopic distribution of the ^{210}Po as shown by autoradiography. Distribution throughout the lung was distinctly nonuniform for the ^{210}Po contained on hematite. Reduction of the quantity of hematite from 3-0.3 mg should have had the effect of further increasing the nonuniformity of the ^{210}Po in the lung, as there were only one-tenth as many particles administered and each of the particles contained 10 times as much activity. Preliminary results suggested that an equal amount of ^{210}Po adsorbed on 0.3 mg of hematite was even less effective for lung tumor induction than when adsorbed on a larger number of carrier particles of lower specific activity. LITTLE and co-workers (1973) tentatively concluded that these results have led us to the conclusion that "in the dose range studied alpha radiation is more carcinogenic when a lower but relatively

uniform dose is delivered to a large volume of lung tissue than when a similar amount of radioactivity is distributed nonuniformly such that the primary effect is to deliver much higher radiation doses to relatively small tissue volumes." Of course, one can summarily dismiss these experimental data by arbitrarily defining a "hot particle" as one which has a higher specific activity than the particles used by Little. In fact, this has been done recently (TAMPLIN and COCHRAN, 1974).

The work of Herman Cember and his associates is also of interest to the "hot particle" problem, even though he used beta- and beta/gamma-emitting isotopes. We should note that some of this work was reported in the open literature almost 20 years ago (CEMBER *et al.*, 1955) and that CEMBER (1964a) addressed the "hot particle" problem in some detail a decade ago. Cember has reviewed his studies of the carcinogenic properties of several radionuclides (^{144}Ce , ^{90}Sr , and ^{35}S) administered to animals in several different chemical and physical forms (Table 6). His work with ^{144}Ce employed both the chloride and fluoride forms given by intratracheal instillation. He and his co-workers injected rats intratracheally with $^{144}\text{CeF}_3$ particles. The particle size distribution was estimated to be $1.0 \pm 1.4 \mu\text{m}$. The total amount of CeF_3 injected was constant, but the radioactive content was varied so that animals in five groups received 50, 25, 15, or 5 μCi or stable cerium (CEMBER *et al.*, 1959). Acute radiation pneumonitis was seen in the three high level groups. Squamous cell carcinomas of the lungs were found in 13 of 93 rats (14%), with all but two being in the 25- or 50- μCi groups. The first tumor was seen at 48 d following a radiation dose of 5100 rad. The lowest dose in an animal with a tumor was 2400 rad, while the average dose to the lung for the tumor-bearing animals in the high level group was 21 000 rad. In a second study with $^{144}\text{CeF}_3$, CEMBER (1963) used intratracheal

injection to expose rats to activities of 0.5, 1.0, 2.0, or 4.0 μCi . The $^{144}\text{CeF}_3$ again had a particle size distribution of $1.0 \pm 1.4 \mu\text{m}$. Lung tumors were seen in all treatment levels. A total of 32 of 161 rats (20%) were affected, with the highest level group (4.0 μCi) having the highest incidence (14 out of 42, or 33%). The earliest tumor was seen 367 days after injection, and the lowest dose to the lung associated with tumor induction was 600 rad. Tumors were mostly carcinomas with squamous cell carcinomas, adenocarcinomas, undifferentiated carcinomas, and lymphomas all being found.

CEMBER and STEMMER (1964) exposed rats to $^{144}\text{CeCl}_3$ by intratracheal injection. Three groups received 10, 15, or 30 μCi of $^{144}\text{CeCl}_3$ per animal. Acute radiation pneumonitis accounted for the death of 28 out of 181 exposed animals within 30 d. Of 150 rats that lived beyond 60 d, 67 developed primary lung cancers. The earliest tumor was seen 70 d post-injection in an animal that received 25 000 rad to the lung. The animals in the 10- μCi group had a lung tumor incidence of 15%, those in the 15- μCi group had an incidence of 60%, and those in the 30- μCi group had an incidence of 73%. Most of the tumors were squamous cell carcinomas (27), but undifferentiated carcinomas (9), adenocarcinomas (5), and other tumors such as lymphosarcomas, fibrosarcomas, and hemangiosarcoma were found. The average lung dose for the tumor-bearing animals in the 10- μCi group was 14 000 rad. In the 15- μCi group, the dose was 19 600 rad and in the 30- μCi group 25 000 rad. One should interpret the ^{144}Ce experiments with caution, as CEMBER (1964b) noted that the $^{144}\text{CeCl}_3$ produced discrete focal areas of radioactivity in the lung even though the chloride form was soluble in solution.

When Cember gave 0, 4.5, 45, or 4500 μCi of $\text{Ba}^{35}\text{SO}_4$ as a single intratracheal injection to rats, no lung cancer or any other lesion suggesting that cancer might develop was observed in any of the experimental animals during a 9-month observation period (CEMBER *et al.*, 1955). When the $\text{Ba}^{35}\text{SO}_4$ was given as

10 weekly doses of 375 μCi each, 2 of the 16 rats which survived the injection regime died at 312 and 319 d later with extensive squamous cell carcinomas of the lung (CEMBER and WATSON, 1958a). The calculated radiation doses were on the order of 12 000 rad.

CEMBER and WATSON (1958b) implanted ^{90}Sr containing glass beads in the lungs of rats. The beads contained from 1.09-59.3 μCi of ^{90}Sr and were $320 \pm 110 \mu\text{m}$ in diameter. Seven of the 23 rats (30%) developed primary pulmonary neoplasia, 4 had squamous cell carcinomas, and 3 had lymphoid neoplasia. The earliest death in a tumor-bearing animal occurred at 169 d following implant. The total radiation dose in these animals, calculated for a sphere of tissue with a radius equal to the range of the beta radiation, ranged from 47 000-260 000 rad. Murine pneumonia was a problem with the experimental animals. No acute deaths were attributable to radiation effects, and no life-shortening was observed.

The experiments of Cember are of considerable interest to the problem of non-uniform dose-distribution. Cember stated that the question of the unique carcinogenic hazard associated with the high absorbed dose gradient around a single radioactive particle deposited in the lung seemed to be answered by the results of the acute $\text{Ba}^{35}\text{SO}_4$ exposures, together with the ^{144}Ce experiments. He also pointed out that the negative results of the long-term retention of several $\text{Ba}^{35}\text{SO}_4$ particulates, under conditions suitable for testing the hypothesis that such focal radiation presents a unique carcinogenic hazard to the lung, imply the absence of such a hazard associated with one or a very small number of loci. His review also emphasized that, for a given total amount of absorbed energy, low-level continuous exposure of the total lung may be more carcinogenic than the same amount of energy delivered acutely to a restricted volume of tissue.

Furthermore, CEMBER (1964a) realized that the quantitative relationships among total absorbed dose, the temporal and spatial distribution of the dose, and the probability of developing radiogenic lung cancer had not been established at that time. However, the similarity of the lung tumor dose-response curves for soluble $^{144}\text{CeCl}_3$ and insoluble $^{144}\text{CeF}_3$ suggested the absence of a "hot particle" effect. CEMBER (1964a) stated, "Should this be true, then it follows that radiation dose to the lung from inhaled radioactive dusts may be calculated, for purposes of estimating radiological risk, by assuming uniform absorption of energy throughout the lung."

In the summary of his review, CEMBER (1964a) states, "Experiments with rats have shown that radioactive substances deposited in the lung can lead to pulmonary neoplasia. Radiations from ^{35}S , $^{90}\text{Sr}/^{90}\text{Y}$, and ^{144}Ce elicited bronchogenic carcinoma and alveolar cell carcinoma in addition to several other tumor types. These experiments did not confirm the existence of a unique carcinogenic hazard due to the intense concentration of absorbed energy in the lung tissue immediately surrounding an inhaled radioactive particle."

Let us now turn to the experiments conducted at the Los Alamos Scientific Laboratory. RICHMOND *et al.* (1970) investigated the effects of $^{238}\text{PuO}_2$ particles lodged in the lung vasculature of rats following intravenous injection. The injected particles are carried by the venous blood to the right heart, from where they are transported to the lung and trapped in the capillaries and small blood vessels of the lung vasculature. These particles averaged about 180 μm in diameter and gave calculated average dose rates to the entire lung of about 3.5 rem/hr, with the alpha-particle dose rate at the surface of the particle on the order of 10^8 rad/hr. Among this exposure was a group of 6 rats which were sacrificed at 600 d. In all, approximately 70 animals were used in this experiment. Examination of the lung following these exposures indicated

the presence of a microlesion with complete degeneration of the cells close to the particle. However, the evidence indicated that this was not simply a stable type of scar tissue, as the lesion was in a dynamic state in which the collagen was renewed constantly and subsequently liquefied. Within this time period, there was no indication of effects which would be deleterious to the overall well-being of the animals. No lung cancers were observed in this study. It is noteworthy that the energy delivered to the lung, if averaged over the full lung, would be on the order of 2 000 000 rad at 600 d -- well in excess of those doses which have been shown to produce death in relatively short times, when more uniformly distributed, and considerably above the doses required to produce lung cancer. Figure 3 shows a photomicrograph of a rat lung section depicting the microlesion at 21 d following exposure. Note the normal appearing alveolar structure close to the microlesion. A more mature lesion showing the anastomosing collagen bundles comprising the fibroma at 180 d following implant is shown in Fig. 4. The development of the microlesions has been described in more detail (RICHMOND *et al.*, 1970). Similar lesions have also been produced in dog lung (RICHMOND *et al.*, 1974) but, as was the case for rats, no lung cancers were produced.

Current work is in progress at the Los Alamos Scientific Laboratory using a similar experimental design but with 10- μ m diameter ZrO microspheres loaded with PuO₂ to specific activities which correspond to respirable size particles. These experiments are directly applicable to the "hot particle" problem. Table 7 shows the plutonium content of the microspheres used in the preliminary LASL "hot particle" experiment. The exposure levels cover particle specific activities ranging from 0.07 to approximately 60 pCi of plutonium. Both ²³⁹Pu and ²³⁸Pu were used to keep the total weight fraction of plutonium very small, as shown in the last column. Table 8 shows the exposure conditions for this

same experiment. At each exposure level, shown in the first column, 70 hamsters received 2000 plutonium-containing particles. Approximately half the hamsters were sacrificed during the experiment, and the remainder were kept for their full life span. The total burden per hamster ranged from 0.14 to approximately 120 nCi. Table 8 also shows the radiation doses calculated by several dosimetric models and the number of tumors expected as calculated from a lung model proposed by COLEMAN and PEREZ (1969) which was based on Albert's rat skin data. This model is basically similar to those developed by GEESAMAN (1968) and by DEAN and LANGHAM (1969), as the dose-response function is based on the Albert rat skin data. About 1% of the lung mass of these animals was irradiated, and the median dose rate to those cells within alpha range of the microsphere was estimated to range from 20-1800 rad/d.

Since the original experiment shown in Table 8 was designed, other hamsters have been injected with larger numbers of microspheres containing lower specific activities per sphere. Figure 5 summarizes the distribution of exposure groups with respect to the number of spheres per animal (ordinate) and the specific activity of the spheres (abscissa). The solid lines are loci of constant lung burden and are labeled as nCi of plutonium per animal. To date, approximately 2000 animals have been used in these experiments. Figure 6 is a photomicrograph of an autoradiograph showing 10- μ m diameter ZrO particles containing plutonium. Figure 7 is a photomicrograph of a hamster lung section showing a single microsphere in an alveolar sac near the termination of the cuboidal epithelium in a respiratory bronchiole. Details on the production and administration of the microspheres and additional experimental detail are given in several progress reports (RICHMOND and VOELZ, 1972; RICHMOND and VOELZ, 1973; RICHMOND and SULLIVAN, 1974).

No aberrant clinical signs have been observed in any of the animals that have died or have been sacrificed to date. Blood samples have revealed no abnormalities even after long exposures, and there have been no regional lymph node effects. Small accumulations of macrophages occasionally are seen around the spheres, but the fibrous encapsulation previously described for the larger, more radioactive (about 180- μ m diameter or 550 μ Ci) ^{238}Pu microspheres (RICHMOND *et al.*, 1970, 1974) is not seen. Only two lung tumors have been observed to date for the hamsters shown in Table 8. One hamster developed a hemangiosarcoma of the lung after 9.5 months exposure to 2000 microspheres containing 0.42 pCi alpha activity. Another hamster developed a lung sarcoma after 12 months exposure at the same level (2A).

Approximately 1150 hamsters have lived their full life spans or have been sacrificed to date as part of the LASL experiments with 10- μ m diameter microspheres. About 6×10^6 spheres with specific activities above 0.07 pCi each were injected into these hamsters. The observation of three primary lung tumors suggests a tumor risk of roughly 10^{-7} per particle as a preliminary order of magnitude estimate. These results are particularly significant in view of the demonstration by LITTLE *et al.* (1970a,b, 1973) that the Syrian hamster develops lung tumors with high efficiency at short induction times following exposure to soluble ^{210}Po . It should also be noted that, because every animal in the preliminary experiment (Table 8) received 2000 microspheres, each in excess of 0.07 pCi, every animal should have developed several lung tumors if the lung tumor probability is 10^{-3} per particle as speculated by GEESAMAN (1968) and others. Only two lung tumors were observed.

The Los Alamos experiments using the 10- μ m diameter plutonium-containing microspheres are of special interest because of the few lung cancers produced, as compared with the large numbers predicted by theoretical models, and the

virtual lack of any observable biological damage. Figure 8 shows photomicrographs of lung sections obtained from a young control hamster (upper left), an older control animal in which hemosiderin-laden macrophages have accumulated in an alveolar space (upper right), a small focal reaction in which macrophages have surrounded a microsphere which has been extruded into the alveolar space (lower left), and a higher magnification showing more detail (lower right). It should be noted that much of the lung tissue contained in the field is unaffected (lower left).

Figure 9 shows a microsphere in a lung capillary with normal alveolar septal tissue in contact with the sphere. No reaction is visible, even though the sphere contained 4.3 pCi ^{238}Pu and was in the lung for 5 months. (The lung burden was 8.6 nCi.) Figure 10 shows a microsphere containing 0.42 pCi ^{239}Pu that has occluded a capillary and bulges into the alveolar space. The microsphere is still covered by vascular adventitial stroma, and no reaction was observed in this hamster sacrificed 8 months after injection.

The data of ANTONCHENKO *et al.* (1969) are of interest, although the experimental end point was average life span rather than carcinogenesis. Rats were given plutonium citrate or ammonium plutonium pentacarbonate by inhalation and compared to rats given plutonium nitrate intratracheally. Although the initial lung content was similar in each instance, the rats given plutonium nitrate received a more nonuniform distribution of dose yet lived four times longer than those rats receiving the plutonium by inhalation.

ROSENTHAL and LINDENBAUM (1969) showed that plutonium injected intravenously into CF1 mice was clearly more effective at producing bone sarcomas when given in the monomeric form as compared with the polymeric form. Mice receiving the monomeric plutonium started to die earlier and developed about twice as many bone tumors than those receiving polymeric plutonium. These data, although not

for lung tissue, are of interest because the polymeric form results in more nonuniform dose-distribution in bone as compared with the monomeric form.

Early experimental results reported by LAFUMA *et al.* (1974) using ^{238}Pu , ^{239}Pu , ^{241}Am , and ^{244}Cm in rats suggest that toxicity increases with the uniformity of the isotope in the lung. For equivalent radiation doses, ^{244}Cm was the most highly dispersed throughout the lung and the most toxic.

SANDERS (1973) reported an increase in lung tumors for rats at estimated lung doses of about 30 rad following inhalation of ^{238}Pu . A nonstatistically significant increase was also observed for rats accumulating lung doses of 9 rad. Sanders suggested that the occurrence of lung tumors at lower doses than those in the vicinity of 50 rad reported from the Russian literature (MOSKALEV, 1972) might be attributed to the greater uniformity of dose-distribution in the lung for the ^{238}Pu as compared with the ^{239}Pu . Sanders speculated that fewer target epithelial cells would be "hit" in the lung by "polymeric" ^{239}Pu than by an equal amount of "monomeric" ^{238}Pu .

In conclusion, I believe a careful consideration of the results obtained from these experimental studies demonstrates that plutonium distributed nonuniformly in the lung (particulates) is not more hazardous than the same amount of plutonium distributed uniformly. In fact, these data suggest that the potential hazard from plutonium increases as the dispersion throughout the lung becomes more uniform.

For cases of nonuniform exposure, as occurs for particulate plutonium, there appears to be a biological sparing effect resulting from the fact that fewer cells are exposed to the alpha radiations, and much of the alpha energy is wasted as compared with a more uniform distribution. Also, the collective defenses of the body, both local and abscopal, such as inhibition of transformed cells by normal cells and immune surveillance, are more efficient in the

case of nonuniform distribution. The key to the problem may well be the number of cells that interact with an alpha particle but are not killed. For the nonuniform distribution case, there are fewer of these cells which might have the potential to form a cancer, and they would be in an environment which would tend to inhibit their division and development to proceed to form a cancer.

It is quite clear that plutonium is seldom uniformly distributed throughout the body. MOSKALEV (1972) pointed out that plutonium, regardless of the compound inhaled, aggregates to form alpha-track stars in the macrophages of the inter-alveolar septae, in peribronchial connective tissue, in lymphatic tissue, and in bronchial spaces. He concluded that, even following the inhalation of soluble compounds, plutonium is distributed nonuniformly in the lung and that, in time, a large fraction of the lung is not irradiated at all. An added feature of the general nonuniformity is the presence of alpha stars located predominantly in the subpleural regions of the lung in both man and experimental animals.

The "hot particle" problem has recently been brought to the attention of several federal agencies (TAMPLIN and COCHRAN, 1974). A short critical review of the NRDC position was prepared by the United Kingdom's National Radiological Protection Board (DOLPHIN, 1974) which concludes: "It is noted that the basis of ICRP recommendations is the average radiation dose to an organ and not the number of radioactive particles in the organ. This dosimetric basis of radiological protection has been established for many years by observation of humans and experimental work with animals. A better evaluation than that offered by Tamplin and Cochran would be needed for this system to be set aside in favour of the hot particle concept. Their estimate that there is a risk of cancer being generated in cells surrounding a hot particle of 1 in 2,000 cannot be substantiated by our present knowledge."

Also of interest is a paper by ZALMANZON and CHUTKIN (1971) from the USSR which states that nonuniformity of exposure in the lung is partially compensated for by the migration of deposited particles in the respiratory organs and consequently that one can accept the entire mass of the lungs, as a first approximation, for the value of m .

REFERENCES

- ALBERT R. E., BURNS F. J. and BENNETT P., 1972, *J. Natl. Cancer Inst.* 49, 1131.
- ALBERT R. E., BURNS F. J. and HEIMBACH R. D., 1967a, *Radiation Res.* 30, 515.
- ALBERT R. E., BURNS F. J. and HEIMBACH R. D., 1967b, *Radiation Res.* 30, 525.
- ALBERT R. E., BURNS F. J. and HEIMBACH R. D., 1967c, *Radiation Res.* 30, 590.
- ALBERT R. E., NEWMAN W. and ALTSHULER B., 1961, *Radiation Res.* 15, 410.
- ALBERT R. E., PHILLIPS M. E., BENNETT P., BURNS F. and HEIMBACH R., 1969, *Cancer Res.* 29, 658.
- ANTONCHENKO G. P., KOSHURNIKOVA N. A. and LYUBCHANSKII E. R., 1969, *Radio-biologiya* 9(1), 75 [also English translation in *Radiobiology* 9(1), 97 (1969)].
- BAIR W. J., RICHMOND C. R. and WACHHOLZ B. W., 1974, *A Radiobiological Assessment of Spatial Distribution of Radiation Dose from Inhaled Plutonium*, U. S. Atomic Energy Commission Report WASH-1320 (Washington, D. C.: U. S. Government Printing Office), in press.
- BURNS F. J., ALBERT R. E., BENNETT P. and SINCLAIR I. P., 1972, *Radiation Res.* 50, 181.
- BURNS F. J., SINCLAIR I. P. and ALBERT R. E., 1973, *Proc. Amer. Assoc. Cancer Res.* 14, 88 (abstract).
- CEMBER H., 1963, *Health Phys.* 9, 539.
- CEMBER H., 1964a, in: *Progress in Experimental Tumor Research*, Vol. 4, Edited by HAMBURGER F. (New York: Hafner Publishing Co.), p. 251.
- CEMBER H., 1964b, *Health Phys.* 10, 1177.
- CEMBER H., HATCH R. F., WATSON J. A. and GRUCCI T. B., 1955, *AMA Arch. Ind. Health* 12, 628.

- CEMBER H. and STEMMER K., 1964, *Health Phys.* 10, 43.
- CEMBER H. and WATSON J. A., 1958a, *AMA Arch. Ind. Health* 17, 230.
- CEMBER H. and WATSON J. A., 1958b, *Amer. Ind. Hyg. J.* 19, 36.
- CEMBER H., WATSON J. A. and SPRITZER A. A., 1959, *AMA Arch. Ind. Health* 19, 14.
- COLEMAN J. R. and PEREZ L. J., Jr., 1969, *Consideration of a Tumor Probability Function and Micro-Dosimetry for the Lower Pulmonary Compartment*, Nuclear Utility Services Report NUS-654 (Rockville, Maryland).
- DEAN P. N. and LANGHAM W. H., 1969, *Health Phys.* 16, 79.
- DOLPHIN G. W., 1974, *A Brief Critical Review of "A Report on the Inadequacy of Existing Radiation Protection Standards Related to Internal Exposure of Man to Insoluble Particles of Plutonium and Other Alpha-Emitting Hot Particles" by A. R. Tamplin and T. B. Cochran, 14th February 1974,* National Radiological Protection Board, Harwell, Didcot, Berkshire, United Kingdom.
- GEESAMAN D. P., 1968, *An Analysis of the Carcinogenic Risk from an Insoluble Alpha-Emitting Aerosol Deposited in Deep Respiratory Tissue*, U. S. Atomic Energy Report UCRL-50387 and UCRL-50387 Addendum.
- GROSSMAN B. N., LITTLE J. B. and O'TOOLE W. F., 1971, *Radiation Res.* 47, 253 (abstract).
- HEIMBACH R. D., BURNS F. J. and ALBERT R. E., 1969, *Radiation Res.* 39, 332.
- International Commission on Radiological Protection, 1969, *Publication 14* (Oxford: Pergamon Press).
- LAFUMA J., 1974, *Les Radioelements Inhales*, Seminaire de Radiobiologie sur le thème de *La Contamination Radioactive Interne*, 18, 19 et 20 Mars 1974-Paris-organisé par la Société Française de Radioprotection.

- LITTLE J. B., GROSSMAN B. N. and O'TOOLE W. F., 1970a, in: *Morphology of Experimental Respiratory Carcinogenesis*, Edited by NETTESHEIM P., HANNA M. G., Jr. and DEATHERAGE J. W., Jr., U. S. Atomic Energy Commission Report CONF-700501, AEC Symposium Series 21 (Springfield, Virginia: Office of Information Services), p. 383.
- LITTLE J. B., GROSSMAN B. N. and O'TOOLE W. F., 1970b, *Radiation Res.* 43, 261 (abstract).
- LITTLE J. B., GROSSMAN B. N. and O'TOOLE W. F., 1973, in: *Radionuclide Carcinogenesis*, Edited by SANDERS C. L., BUSCH R. H., BALLOU J. E. and MAHLUM D. D., U. S. Atomic Energy Commission Report CONF-720505, AEC Symposium Series 29 (Springfield, Virginia: Office of Information Services), p. 119.
- MOSKALEV Y. I., 1972, *Health Phys.* 22, 723.
- National Academy of Sciences-National Research Council, 1961, *Effects of Inhaled Radioactive Particles*, Publication 848.
- PASSONNEAU J. V., BRUES A. M., HAMILTON K. A. and KISIELESKI W. E., 1952, *Carcinogenic Effects of Diffuse and Point Source Beta Irradiation on Rat Skin: Final Summary*, U. S. Atomic Energy Commission Report ANL-4932, p. 31.
- RICHMOND C. R. and VOELZ G. L., editors, 1972, *Annual Report of the Biological and Medical Research Group (H-4) of the LASL Health Division, January through December 1971*, Los Alamos Scientific Laboratory Report LA-4923-PR.
- RICHMOND C. R. and VOELZ G. L., editors, 1973, *Annual Report of the Biological and Medical Research Group (H-4) of the LASL Health Division, January through December 1972*, Los Alamos Scientific Laboratory Report LA-5227-PR.
- RICHMOND C. R. and SULLIVAN E. M., editors, 1974, *Annual Report of the Biomedical and Environmental Research Program of the LASL Health Division, January through December 1973*, Los Alamos Scientific Laboratory Report LA-5633-PR.

- RICHMOND C. R., LANGHAM J. and STONE R. S., 1970, *Health Phys.* 18, 401.
- RICHMOND C. R., HOLLAND L. M., DRAKE G. A. and WILSON J. S., 1974, in:
Abstracts Presented at the Nineteenth Annual Meeting of the Health Physics Society, Houston, Texas (July 8-12, 1974), p. 33 (abstract P/127).
- SANDERS C. L., 1973, *Radiation Res.* 56, 540.
- ROSENTHAL M. W. and LINDENBAUM A., 1969, in: *Delayed Effects of Bone-Seeking Radionuclides*, Edited by MAYS C. W., JEE W. S. S., LLOYD R. D., STOVER B. J., DOUGHERTY J. H. and TAYLOR G. N. (Salt Lake City: University of Utah Press), pp. 371-386.
- TAMPLIN A. R. and COCHRAN T. B., 1974, *Radiation Standards for Hot Particles: A Report on the Inadequacy of Existing Radiation Protection Standards Related to Internal Exposure of Man to Insoluble Particles of Plutonium and Other Alpha-Emitting Hot Particles*, Natural Resources Defense Council Report (Washington, D. C.).
- ZALMANZON Yu. F. and CHUTKIN O. A., 1971, *Pogloshcheniya Doza V Legkikh Obuslovljennaya Radioaktivnimi Aerosolyami*, Union Scientific Research Instrument Engineering Institute (translated as *Radiation Absorbed Dose in the Lungs from Radioactive Aerosols* by The Ralph McElroy Co., Custom Division, Austin Texas, Code No. 124-3188-2).

*Table 1. Classes of nonuniformity of radiation dose**

-
1. Partial irradiation of an organ or tissue where the part irradiated is representative of the whole organ or tissue (e.g., external irradiation of skin or bone marrow)
 2. Partial irradiation where the part irradiated is not representative of the whole (e.g., bone-seeking radionuclides in bone) -- tritiated thymidine in DNA is special case
 3. *Irradiation from radioactive materials in particulate form*
-

*See ICRP (1969).

Table 2. Activity and number of $^{239}\text{PuO}_2$ and $^{238}\text{PuO}_2$ particles as a function of geometric diameter

Geometric diameter (μm)	$^{239}\text{PuO}_2$		$^{238}\text{PuO}_2$	
	nCi/particle	Particles/16 nCi	nCi/particle	Particles/16 nCi
0.01	3.0×10^{-10}	5.4×10^{10}	8.0×10^{-8}	2.0×10^8
0.1	3.0×10^{-7}	5.4×10^7	8.0×10^{-5}	2.0×10^5
1.0*	3.0×10^{-4}	5.4×10^4	8.0×10^{-2}	2.0×10^2

* Dose rate at surface of $^{238}\text{PuO}_2$ particle is $\sim 10^5$ rad/hr.

Table 3. Relationship of particle size to number of cells
at risk for a static lung burden of 16 nCi $^{239}\text{PuO}_2$ *

Particle diameter (μm)	Number of particles	Activity per particle (pCi)	Cells at risk	Fraction of lung (%)
0.1	5.4×10^7	3×10^{-4}	3.0×10^{11}	30
0.3	2.0×10^6	0.01	1.3×10^{10}	1
0.7	1.8×10^5	0.08	1.2×10^9	0.1
1.0	5.4×10^4	0.3	3.6×10^8	0.03

* Assuming static particles in a structureless human lung of uniform density $0.2 \text{ g} \cdot \text{cm}^{-3}$ with an average cell volume of $10^3 \mu\text{m}^3$. Cells at risk are taken to be those in a sphere of radius equal to the alpha range (200 μm at the assumed density).

*Table 4. Experimental exposure conditions used
by Albert et al. for skin irradiation experiments**

Radiation		Aperture	Volume (mm ³) [†]
β and E^-	Mask	25 cm ²	2500
	Grid (slots)	1 x 32 mm	32
	Sieve (pores)	2.5 mm diameter	4.5
Proton	Pores	5 mm diameter	19.6
		2 mm diameter	3.14
		0.5 mm diameter	0.2
Alpha			2.7×10^{-4}

*See ALBERT *et al.* (1967) and BURNS *et al.* (1972).

[†]Assuming average of 1 mm depth irradiated.

Table 5. Tumor production in rat skin following exposure to flat plate and point sources of $^{90}\text{Sr}/^{90}\text{Y}$ *

Source	Activity	Number of rats	Number of tumors		Relative Efficiency	
			Total	Per bead		
Flat Plate						
1000 μCi	28.6 $\mu\text{Ci}/\text{cm}^2$	71	89	~ 0	0.00049	1.59
1500 μCi	42.9 $\mu\text{Ci}/\text{cm}^2$	73				
50 Beads	30 $\mu\text{Ci}/\text{bead}$	58	27	0.009	0.00031	1.00
20 Beads	75 $\mu\text{Ci}/\text{bead}$	77	24	0.016	0.00021	0.671
10 Beads	150 $\mu\text{Ci}/\text{bead}$	74	16	0.022	0.00014	0.464

* Modified from PASSONNEAU *et al.* (1952) and information given in NAS-NRC publication 848 (1961).

Table 6. Summary of experiments by Cember and co-workers
involving intratracheal injection of beta-emitting radionuclides in rats

Reference	Radionuclide	Activity	Number of tumors/animals	% Tumor incidence	Tumor-bearing rats		
					Days to death Earliest	Average	Lung dose (rad)
CEMBER and WATSON (1958a)	Ba ³⁵ SO ₄	375 Ci/wk x 10	2/24	8	313	315	12 000-20 000
CEMBER <i>et al.</i> (1959)	¹⁴⁴ CeF ₃	5-50 µCi	13/93	14	48	> 100	2400-> 21 000
CEMBER (1963)	¹⁴⁴ CeF ₃	0.5-4 µCi	32/161	20	367	> 400	600-> 4000
CEMBER and STEMMER (1964)	¹⁴⁴ CeCl ₃	10-30 µCi	67/181	37	70	~ 300	14 000-25 000

*Table 7. Plutonium content of microspheres
for preliminary LASL "hot particle" experiment*

Isotope	Level	Specific activity (pCi/sphere)	Equivalent diameter of pure $^{238}\text{PuO}_2$ (μm)	PuO_2 weight fraction
^{239}Pu	1	0.07	0.09	4.3×10^{-4}
	2	0.22	0.13	1.4×10^{-3}
	2A	0.42	0.16	2.9×10^{-3}
	3	0.91	0.21	5.8×10^{-3}
	3A	1.6	0.26	1.1×10^{-2}
^{238}Pu	3A	2.1	0.28	4.8×10^{-5}
	4	4.3	0.36	1.0×10^{-4}
	4A	8.9	0.46	2.0×10^{-4}
	5	13.3	0.52	3.3×10^{-4}
	6	59.4	0.86	1.3×10^{-3}

Table 8. Exposure conditions for preliminary LASL "hot particle" experiment*

Level	nCi/hamster	Average dose rate [†] (rad/yr)	Local dose rate at Surface of sphere (rad/hr)	40 μ m from center (rad/hr)	Possible lung tumor incidence [¶] (tumors/group)
1	0.14	13	4.2×10^1	6.8×10^{-1}	2
2	0.44	42	1.2×10^2	2.2×10^0	10
2A	0.84	81	2.5×10^2	4.1×10^0	40
3	1.82	175	5.5×10^2	1.0×10^1	60
3A	3.2	310	1.0×10^3	1.7×10^1	40
4	8.6	875	2.5×10^3	4.2×10^1	10
5	26.6	2710	8.4×10^3	1.3×10^2	0
6	119.0	12 100	3.6×10^4	5.8×10^2	0

* 2000 spheres/hamster, 60 hamsters/group.

[†] Assuming 1 g lung irradiated.

[¶] Using NUS structure lung with density of 0.19 g/cm³ (COLEMAN and PEREZ, 1969).

FIG. 1. Graphic representation of portions of hair follicle irradiated with greater than 50% of the Bragg peak dose: (A) 0.12 mm penetration; (B) 0.35 mm penetration; and (C) 0.55 mm penetration.

FIG. 2. (a) Autoradiograph of hamster lung sacrificed 7 days after a single intratracheal injection of 0.2 μCi ^{210}Po on hematite. (b) Autoradiograph of hamster lung sacrificed 7 days after a single intratracheal injection of 0.2 μCi ^{210}Po alone in saline. The distribution of the alpha tracks is much more random than the aggregates shown in (a).

FIG. 3. Photomicrograph of microlesion caused by $^{238}\text{PuO}_2$ microsphere embedded in rat lung tissue for 21 days. A network of loosely packed, circumferentially arranged collagen fibers is present. Radiation-induced changes such as nuclear condensation and vacuolization are apparent in the portion of the respiratory bronchiole adjacent to the lesion. Grossly normal-appearing alveoli and intra-alveolar septa appear around the lesion.

FIG. 4. Photomicrograph of microlesion caused by $^{238}\text{PuO}_2$ microsphere in rat lung tissue for 180 days. The collagen bundles are now fused into dense hyaline-appearing anastomosing masses. This zone of circumferential fibrosis rapidly undergoes transition into a wide region of looser fibrosis where the general alveolar structure persists.

FIG. 5. Distribution of exposure groups with respect to number of spheres per animal (ordinate) and specific activity of spheres (abscissa). The lines are loci of constant lung burden and are labeled with nCi of plutonium per animal. Symbols indicate year of injection: (-●-) 1971; (-■-) 1972; and (-▲-) 1973.

FIG. 6. Autoradiograph of ceramic microspheres of ZrO_2 loaded with $^{238}\text{PuO}_2$.

FIG. 7. Photomicrograph of hamster lung section showing a single 10- μ m microsphere in alveolar sac near termination of the cuboidal epithelium in a respiratory bronchiole.

FIG. 8. Photomicrographs of lung sections obtained from a young control hamster (upper left), an older control animal in which hemosiderin-laden macrophages have accumulated in an alveolar space (upper right), a small focal reaction in which macrophages have surrounded a microsphere which has been extruded into the alveolar space (lower left), and a higher magnification showing more detail (lower right).

FIG. 9. Microsphere in a lung capillary with normal alveolar septal tissue in contact with the sphere.

FIG. 10. Microsphere containing 0.42 pCi ^{239}Pu that has occluded a capillary and bulges into the alveolar space. The microsphere is still covered by vascular adventitial stroma.