BROOKHAVEN NATIONAL LABORATORY

Quarterly Progress Report *July 1 - September 30, 1954*

Unclassified Section

Associated Universities, Inc.
under contract with the
United States Atomic Energy Commission

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

II

Printed at Upton, N.Y., for distribution to
individuals and organizations associated
with the national atomic energy program.

December, 1954

1050 copies

FOREWORD

This report, the nineteenth in this series of Quarterly reports, covers the unclassified research progress of the Laboratory. Abstracts only are given for research activities being reported concurrently in the scientific journals. The appropriate journal or document reference is also given in such cases. A research activity that has not yet reached the stage of journal publication will be reported on in such detail as its progress warrants.

The classified work of the Laboratory is covered in the classified quarterly reports, issued as of the 15th of February, May, August and November. It is in order to expedite the publication of all progress reports that their publication does not coincide with that of the unclassified quarterlies.

The AEC budget activity numbers for the various researches are given in parentheses at the right hand side of the page, alongside the headings to which they pertain. The symbol (†) is used to designate a supporting activity whose cost is distributed to several research activities.

Previous Quarterly Progress Reports (Unclassified)

BNL 51 (S-5)	January 1 - March 31, 1950
BNL 64 (S-6)	April 1 - June 30, 1950
BNL 82 (S-7)	July 1 - September 30, 1950
BNL 93 (S-8)	October 1 - December 31, 1950
BNL 103 (S-9)	January 1 - March 31, 1951
BNL 117 (S-10)	April 1 - June 30, 1951
BNL 132 (S-11)	July 1 - September 30, 1951
BNL 149 (S-12)	October 1 - December 31, 1951
BNL 177 (S-13)	January 1 - March 31, 1952
BNL 189 (S-14)	April 1 - June 30, 1952
BNL 209 (S-15)	July 1 - September 30, 1952
BNL 219 (S-16)	October 1 - December 31, 1952
BNL 232 (S-17)	January 1 - March 31, 1953
BNL 243 (S-18)	April 1 - June 30, 1953
BNL 259 (S-19)	July 1 - September 30, 1953
BNL 270 (S-20)	October 1 - December 31, 1953
BNL 289 (S-21)	January 1 - March 31, 1954
BNL 299 (S-22)	April 1 - June 30, 1954

IV

CONTENTS

PHYSICS DEPARTMENT

Inelastic scattering of 500-Mev negative pions in emulsion nuclei - - -	1
Vacuum polarization in a strong coulomb field - - - - -	1
K-particle production by protons 2.2 and 3.0 Bev - - - - -	1
The ratio of Γ_n^0/D for slow neutron resonances - - - - -	2
Neutron capture cross section of Ta ¹⁸² - - - - -	2
Color centers induced in Al ₂ O ₃ by reactor and gamma-ray irradiation - - - - -	3
A thin, wide-aperture, proportional counter - - - - -	3
A microsecond shutter for use in separating neutrons of various speeds - - - - -	4

COSMOTRON DEPARTMENT

Research - - - - -	5
Operation - - - - -	6

ACCELERATOR DEVELOPMENT DEPARTMENT

Electron Analogue - - - - -	8
AGS Project - - - - -	8

CHEMISTRY DEPARTMENT

Disintegration of bismuth by 2.2 Bev protons - - - - -	11
Excitation functions of radioactive products from the interaction of high energy protons with lead - - - - -	11
Radiochemical studies of nuclear reactions at Bev energies - - - - -	11
An attempt to detect the antineutrinos from a nuclear reactor by the Cl ³⁷ (ν^- , e ⁻)A ³⁷ reaction - - - - -	12
K-electron ejection in the beta decay of S ³⁵ - - - - -	12
Radioactive anthracene-C ¹⁴ and acridine-C ¹⁴ from the neutron irradiation of acridine - - - - -	12
A one-step method of synthesizing tritium-labeled compounds - - - - -	12
Chemical effects of neutron capture in some cobaltic hexammines - - -	13
The radiation chemistry of nitrite ion in aqueous solution - - - - -	13
Photoinitiated oxidation of benzoic acid by ferric ions in aqueous solution - - - - -	14
Backscattering from targets of low atomic number bombarded with 1- to 2-Mev electrons - - - - -	14
A semi-empirical study of the H ₂ Cl transition complex through the use of hydrogen isotope effects - - - - -	14
Complex ions and electron exchange - - - - -	15
The mechanism of the ring contraction of tropolones - - - - -	15

The acid catalyzed rearrangement of α -cinenic acid - - - - -	16
The effect of isotopic substitution on the mass spectra of molecules.	
II. Oxygen and carbon dioxide, experimental - - - - -	16
The effect of isotopic substitution on the mass spectra of molecules.	
III. Carbon dioxide, theoretical interpretations - - - - -	16
The isotope abundances of chlorine from various sources - - - - -	16
The mass spectrum of lithium iodide - - - - -	17
Neutron diffraction studies of ferrimagnetic spinels - - - - -	17

REACTOR DEPARTMENT

Foreword and General Notes - - - - -	18
Operations - - - - -	19
Isotope Distribution and Special Materials Group - - - - -	19
Neutron Physics - - - - -	20
Neutron Physics -- Cooperative Programs - - - - -	21
Shielding - - - - -	22
Meteorology - - - - -	22
Publications - - - - -	23

NUCLEAR ENGINEERING DEPARTMENT

Special Radioisotopes Development - - - - -	25
Radiochemical Analytical Section - - - - -	26
Hot Laboratory Operations - - - - -	28
Ultimate Waste Disposal - - - - -	29
Fission Products Utilization - - - - -	30
Heat Transfer in Spray Columns - - - - -	31

INSTRUMENTATION AND HEALTH PHYSICS DEPARTMENT

Instrumentation Division - - - - -	33
Health Physics Division - - - - -	34
Publications - - - - -	36

BIOLOGY DEPARTMENT

Preliminary Results - - - - -	37
Submitted for Publication - - - - -	37
Improved storage quality of potato tubers following exposure to gamma radiation from cobalt-60 - - - - -	37
The effects of adrenal hormones on the capillary circulation and on capillary permeability in perfused rat tissues - - - - -	37
Stimulation by chronic gamma irradiation of the spontaneous rates of heritable somatic instabilities in a clone of "Nicotiana" - - - - -	38
Mechanism of action of carbohydriase - - - - -	38
Cytogenetic analysis of a spontaneous pre-meiotic mutation in maize - - - - -	38
The sensitivity of developing "Habrobracon" to oxygen - - - - -	39

An improved sealing wax for temporary smear preparations - - - - -	39
The effect of certain gases on the radiosensitivity of "Habrobracon" during development - - - - -	39
Treatment by black-widow spiderlings of Sr^{90} - Y^{90} in their prey - - -	40
Isotopic exchange criteria for enzyme mechanisms - - - - -	40
The effect of a yeast upon the incorporation of phosphorus into the tissue of adult fruit flies - - - - -	40
Quantitative studies on the phosphorus balance of adult female fruit flies feeding upon yeast - - - - -	41
Vertebrate gonadotrophins - - - - -	41
A mechanism of gene interaction - - - - -	42
An inverse relationship between the water content of seeds and their sensitivity to x-rays - - - - -	42
Some aspects of the comparative biochemistry of iodine utilization and the evolution of thyroidal function - - - - -	42
Bacterial virus: radiation and function - - - - -	43
The relation of genome number to radiosensitivity in "Habrobracon" - -	43
General mechanisms for enzyme action - - - - -	43
Histopathological changes in "Triturus viridescens" after x-irradiation, with survival prolonged by low temperature - - - -	44
The retention of metabolic radiocarbonate - - - - -	44

MEDICAL DEPARTMENT

Division of Pathology - - - - -	45
Effects of cortisone on the thyroid function in white leghorn cockerels -	47
Division of Bacteriology and Virology - - - - -	47
Division of Biochemistry - - - - -	48
Division of Physiology - - - - -	50
Preparations in soluble monoamine oxidase - - - - -	55
The experimental application of neutron capture therapy to glioblastoma multiforme - - - - -	55

BNL-314

Summaries of papers submitted for publication by members of the Physics Department during the period July 1 - September 30, 1954 are given below. Titles of papers presented at various meetings during the review period are given at the end of each section. Detailed summaries of these papers will be included in a future progress report when the completed papers are in the process of being published.

High Energy Events, Mesons

INELASTIC SCATTERING OF 500-MEV NEGATIVE PIONS IN EMULSION NUCLEI. M. Blau and M. Caulton.** Phys. Rev. 96, 150 (1954).

Inelastic scattering of 500-Mev negative pions has been investigated in 1700 meson-induced events in emulsion nuclei. Large energy losses and strong angular relationships of the emitted mesons have been observed, similar to those in experiments with lower energy mesons. The possible causes for this behavior are discussed. In 1 to 3 percent of the events, production of charged mesons has been found. The cross section for meson production on free nucleons is estimated to be between 3.5 and 10 mb, considering the possible absorption of one or both mesons. The angular and energy distributions of mesons and recoiling nucleons in events with meson production are discussed. Six events of π^0 production on free protons (or edge nucleons) have been found. The results are compared with observations on meson scattering and meson production in cosmic-ray experiments.

**Rensselaer Polytechnic Institute, Troy, N.Y.

VACUUM POLARIZATION IN A STRONG COULOMB FIELD. E.H. Wichmann** and N.M. Kroll.† Phys. Rev. 96, 232 (1954). Letter.

The problem of the polarization charge density induced by a coulomb field is discussed; radiative corrections are omitted, but the external field is treated exactly. A relatively tractable expression is obtained for the Laplace transform of $r^2 \rho_p(r)$. The expression is used to obtain the small distance behavior of the polarization potential, which is of particular interest for mesic atoms. Numerical results are obtained for all values of the point charge source of the field for which the Dirac equation has allowable solutions. Even for the largest allowable charge, the corrections to the Uehling formula are small. The expansion of the Laplace transform in powers of the charge converges for all allowable values of the charge.

**Columbia University, New York, N.Y.

†Permanent address: Columbia University, New York, N.Y.

K-PARTICLE PRODUCTION BY PROTONS 2.2 AND 3.0 BEV. R.D. Hill,** E.O. Salant, and M. Widgoff. Phys. Rev. 94, 1794 (1954). Letter.

A TAU MESON PRODUCED BY 3.0-BEV PROTONS. R.D. Hill,** E.O. Salant, and M. Widgoff. Phys. Rev. 95, 1699 (1954). Letter.

Different experimental arrangements for the production of heavy mesons detected with nuclear emulsions were tried out, and one proved satisfactory enough for continued use. It consisted essentially of placing the emulsions close to a target struck by

*For explanation of these numbers, see Foreword.

the circulating beam (2.2 or 3 Bev) of the Cosmotron, so that secondary radiations emitted from the target at 90° (or at 45°) to the beam traversed the emulsions roughly parallel to their planes.

Ten K^+ -mesons, decaying at rest into either π^- or μ^- -mesons, one K^- -meson captured at rest and producing a star consisting of a proton and a 50-Mev pion, and one τ -meson decaying into three pions were observed. The τ -meson proved particularly valuable in establishing the masses of the other K-mesons, which agreed within experimental error with the mass of the τ -meson ($965 m_e$).

One of the K^+ -mesons gave a secondary that remained in one plate long enough (5.25 cm) for such a good measurement of its scattering that its characteristics could be determined reliably. This secondary was shown to be a pion of a momentum corresponding to a decay of the K-meson into this positive pion and a neutral pion of the same momentum. With this decay scheme, the mass of the K-meson was computed to be $(950 \pm 40) m_e$ and the Q-value of the decay was computed to be (211 ± 20) Mev. These values agree closely with the corresponding values of the θ^0 particle observed in cloud chambers, which decays into a positive and a negative pion; thus this positive K-meson is the exact charged counterpart of that neutral meson, i.e., it can be identified as θ^+ .

*University of Illinois, Urbana, Ill.

Papers Presented

Chicago APS Meeting.

Chi Meson Produced by 3-Bev Protons. R.D. Hill,* E.O. Salant and M. Widgoff.

*University of Illinois, Urbana, Ill.

Nuclear Reactions and Energy Levels

M. Goldhaber and A.W. Sunyar have written a chapter which will be included in a forthcoming book entitled "Beta and Gamma Ray Spectroscopy," edited by Professor K. Siegbahn and to be published by the North Holland Publishing Co. The chapter (No. 16, entitled "Classification of Isomers") covers a review of the present status of knowledge of nuclear isomers, with emphasis on empirical regularities and tentative theoretical interpretations.

THE RATIO OF $\overline{\Gamma_n^0}/\overline{D}$ FOR SLOW NEUTRON RESONANCES. R.S. Carter, J.A. Harvey, D.J. Hughes, and V.E. Pilcher. Phys. Rev. 96, 113 (1954).

A survey has been made of resonance parameters of the heavy elements which have been measured recently. The variation of the ratio $\overline{\Gamma_n^0}/\overline{D}$ with atomic weight has been compared with theoretical predictions. A maximum was found in the ratio $\overline{\Gamma_n^0}/\overline{D}$ at atomic weight of approximately 160, which is in fair agreement with the more recent theories of neutron scattering by nuclei. The experimental data, however, are not sufficiently accurate to permit a choice to be made between the new theories which have been proposed.

NEUTRON CAPTURE CROSS SECTION OF Ta^{182} . E. der Mateosian. Phys. Rev., in press.

The neutron capture cross section of Ta^{182} was measured by comparing the beta activity of Ta^{182} with that of Ta^{183} in a sample of Ta irradiated in the Brookhaven reactor. The beta activities were compared by growing a NaI(Tl) single crystal which

contained a trace of irradiated Ta and using this crystal as a scintillation detector for its internal source. This technique, called "internal source spectrometry," makes it possible to realize a 4π solid angle geometry, ~ 100 percent efficiency for particles and effectively zero source thickness, and is thus eminently suited to problems in which it is desirable to perform a comparison of the intensities of two or more beta activities over periods of time equal to weeks or months. A knowledge of the decay scheme is not necessary for this method, provided there are no "delayed" internal conversion electrons.

A value of $59,000 \pm 10,000$ barns was obtained for the capture cross section of Ta^{182} for pile neutrons.

Papers Presented

International Nuclear Physics Conference, University of Glasgow.

Gamma Transitions and the Shell Model. M. Goldhaber.

Chicago APS Meeting.

The Importance of Parentage of Nuclear States for Nuclear Reaction Phenomena.
D.H. Wilkinson* and A.M. Lane.**

*On leave from the Cavendish Laboratory, Cambridge, England.

**On leave from AERE, Harwell, England to MIT.

Solid State Physics

COLOR CENTERS INDUCED IN Al_2O_3 BY REACTOR AND GAMMA-RAY IRRADIATION. P.W. Levy and G.J. Dienes. Proc. Phys. Soc. (London), Bristol Conf., in press.

The absorption bands induced in $\alpha\text{-Al}_2\text{O}_3$ by gamma and reactor irradiation have been studied. The slight coloration due to γ -rays saturates. Bands specific to reactor irradiation have been found and their growth studied.

Papers Presented

Am. Crystal. Assoc. Meeting.

A Correlation of Diffraction and Transmission Experiments for X-Ray and Neutron Elastic Scattering. D.T. Keating and J.J. Antal.*

*Watertown Arsenal, Watertown, Mass.

Experimental Techniques

A THIN, WIDE-APERTURE, PROPORTIONAL COUNTER. R.M. Eisberg and H.E. Wegner. Rev. Sci. Instr., in press. Letter.

In detecting charged particles with a proportional counter, it is often convenient to use a counter which is thin in the direction of the beam of particles and wide in the directions transverse to the beam. A new design for a counter satisfying these requirements has been developed and tested, and is described in the note. The performance of the counter is compared with that of wide-aperture counters previously described in the literature. The comparison shows that this design is inherently superior to previous designs.

A MICROSECOND SHUTTER FOR USE IN SEPARATING NEUTRONS OF VARIOUS SPEEDS. F.G.P. Seidl. BNL-278 (T-46).

The factors involved in the design, construction, setting up and operation of a high speed neutron shutter are described in this report. Points covered in this report are as follows: the principle of operation of the chopper; the means of placing very small samples in the neutron beam; the design of the neutron slit system with emphasis on the slits in the rotor; the reasons for choosing a particular type of mechanical design; the rotor structure; the method of mounting and spinning the rotor; power requirements for accelerating the rotor and maintaining it at constant speed; the stresses and strains to be expected in various parts of the spinning rotor; critical speeds and dynamic stability; particular difficulties in construction; and spin tests.

Papers Presented

Chicago APS Meeting.

A New Technique for Observing Cyclotron Phase-Grouping. S.D. Bloom.

Research

The following summary of the research activities of groups using the Cosmotron includes the work of other departments of the Laboratory and of other institutions as well as work by the staff of the Cosmotron Department. The radiochemistry research performed with Cosmotron beams is included in the Chemistry Department section of this report and is not discussed here.

Measurements of the proton-proton elastic scattering angular distributions are in progress. Measurements are made by bombarding thin targets of carbon and polyethylene with the internal proton beam of the Cosmotron and detecting coincidences between the two scattered protons. A striking isotropy in the angular distribution in the center of mass system is found for incident proton energies below 400 Mev. This condition no longer holds at 1 Bev where the cross section rises strongly at small angles. At 90° in the center of mass system the differential cross section at 1 Bev is still fairly large, about 1 millibarn per steradian.

A group at the University of Wisconsin has found, in emulsions exposed to the 3-Bev internal proton beam, an excited heavy fragment which has a decay energy too large to be attributed to the decay of a bound Λ^0 particle. It is now suggested that charged hyperons, which have a Q value about 80 Mev greater than the Λ^0 , may also be found in nuclear fragments.

Brookhaven emulsion groups have continued the study of K-mesons. An event has been observed which is interpreted as the decay of a θ^+ -meson. A charged K-meson stops and decays to produce a charged π -meson, which travels sufficiently far in one emulsion that accurate measurements can be made of grain density vs range. These measurements indicate that the decay particle is definitely a π -meson and, with the assumption that a π^0 -meson is also involved in the decay, they give a Q value in excellent agreement with that of the θ^0 derived from cloud chamber measurements.

Six stopped K^- -mesons have been observed, five in a magnetically analyzed beam. Five produced stars with one black prong and one grey prong. One produced a star with but one grey prong and no black prongs. This prong distribution seems very different from that of the K^- -mesons observed from cosmic rays. A particularly interesting event is under investigation. A K^- -meson stops and produces one black prong and one grey prong of about 3 times minimum ionization which travels 8 mm with little change in grain density and then decays in flight to produce a near minimum particle. The small change in grain density suggests that the mass of the particle emitted from the star is about 2 and greater than that of a charged hyperon.

The Brookhaven group continues to operate their two diffusion cloud chambers and many of the experiments being run are in collaboration with physicists from outside institutions. In all, some ten experiments are in progress which are concerned with the scattering and interaction of various energy protons and mesons with nucleons. A pentane bubble chamber with a volume of about 500 cc has been put into operation by a University of Michigan group. Excellent pictures have been taken in the external proton beam.

A counter experiment is in progress to detect γ -rays which originate from heavy mesons or hyperons which decay at a point some distance from the target where they are produced. The intensity of the γ -rays with increasing energy of incident protons

*For explanation of these numbers, see Foreword.

on a copper target shows a threshold for these γ -rays at 1.1 - 1.2 Bev. This value is consistent with associated production by the process $p + N \rightarrow \Lambda + \theta + p$.

The theoretical group has calculated the polarization P of nucleons elastically scattered from nuclei for the nuclear potential of Woods and Saxon which has the radial dependence $\{1 + \exp[(r - r_0)/a]\}^{-1}$ where r_0 is the nuclear radius and a is a parameter which determines the length of the exponential tail. For this potential, P remains positive throughout the first diffraction minimum. The variation of P with k_1/K and with the strength of the spin-orbit coupling was investigated for the case of aluminum. For the Berkeley data for the polarization of 300-Mev protons scattered by Al, a reasonable fit has been obtained for the region inside the first diffraction peak. Theoretical studies are being made of meson photoproduction and of the integral equations of the covariant Tamm-Dancoff treatment of the meson-nucleon system.

Operation

The experimental program has proceeded mainly as it did during the previous quarter. In scheduling the practice is being followed of giving extra time to those experiments which are in the final data-taking stage; in this way two or three long-standing experiments will be brought to a conclusion fairly soon. To accomplish maintenance work experience has shown that the most effective schedule is a two-week shutdown period for this purpose separated by six to eight weeks of operation.

During the last two-week shutdown, the base of the big motor generator set was grouted and the coupling in the main shaft of the set was tightened by heating the bolts and using a sledge hammer on the end of a large socket wrench. Modifications were also made to the high voltage terminal of the injection Van de Graaff with the result that the pulsed ion source now delivers 3 ma of separated protons to the electrostatic deflection plates.

A careful series of measurements was made to determine what changes in the positions of the magnet blocks had occurred during the past two years. The data clearly show (see Figure 1) a sag in the foundation ring; there are no magnet blocks in the 10-ft intervals across the straight sections. Also there is more settlement in quadrants two and three, which is caused mostly by the weight of the radiation shield. Settlement measurements made nine months after erection (before the shield was in place) show that the ring was settling evenly. The rate of settlement can be judged from Table 1.

Figure 1. Settlement of magnet blocks four years after erection. The sagging of each quadrant near its middle is evident, as is increased sinking due to the pressure of the radiation shield behind the second and third quadrant.

Table 1				
Settling of Magnet Quadrants with Time (in thousandths of an inch)				
Time After Erection	First Quadrant	Second Quadrant	Third Quadrant	Fourth Quadrant
3/4 years	34	28	28	28
4 years	48	61	62	42

If the settlement of outer radius of the magnet is different from that of inner, the magnet blocks will tilt out of level. This can have serious consequences because the average position of the beam will no longer be midway between the top and bottom of the vacuum chamber, and serious vertical oscillations will be introduced. Measurements upon the vertical oscillations for the first few turns after injection do show that the mean position of the beam is about $1/2$ in. above the center of the vacuum chamber.

After the last shutdown a restudy of the injected beam was made to determine what effect the changes in the magnet may have had on the proton orbits. The amplitude of betatron oscillations was found to be greater by about $1/2$ in. than when measurements were made initially; on the average the beam was found to be about $1/2$ in. high. This result is consistent with a downward tilt of the gap of about .050 in. and a first harmonic in elevation of about .050°.

The Department has begun to acquire certain new facilities. Bids have been requested for the fabrication of the steel needed in the strong focusing pair and a supplier for the coils is being sought. A search is being made for a liquifier capable of producing 15 to 25 liters per hour of liquid hydrogen to replace the Chemistry Department's Collins cryostat which has proved inadequate to meet the need for liquid hydrogen in the high energy program. Both low and high pressure liquifiers are being considered.

A test has been completed which shows that it will be possible to extract some 50 percent of the internal circulating protons and have these particles available in a concentrated beam at a distance from the Cosmotron. In this test a newly constructed magnet was located on the inside of the south straight section. After passing through a special lip, the circulating protons on their next turn were displaced into the gap of this magnet; it then deflected them outward so that in their travel through the next quadrant they moved diagonally to the outside of the east straight section. At this point their intensity was measured and half of the protons in the initial beam were found within an area that will be occupied by the gap of a second magnet which will deflect them still further away from the Cosmotron. This second magnet is now being designed. Ultimately the strong focusing pair should be available to focus this beam at various distances. Since some 10^{10} protons per pulse may be expected, this external beam presents a new set of shielding problems.

Activities in the Department were split evenly between the preliminary design and development work on the 25-Bev strong focusing machine and the actual construction of the electron analogue. Detailed description of these activities is given below under their respective headings.

A representative of the Midwestern Universities Research Association, Inc. worked during the summer with the Department and at the same time, several members of the Department spent some time with this midwestern organization.

It was found necessary to reactivate another barracks building to provide the type of accommodations required by the AGS magnet group for precision magnetic measurements. With this new addition, the Department is now being housed in seven barrack-type buildings.

Electron Analogue

Construction of the electron analogue was accelerated since practically all the mechanical components for the machine were delivered during this report period.

All guiding lens elements (see Figure 1) were assembled. However, some difficulty was experienced in getting manufacturers to meet specified tolerances on all lens components, and consequently, it was deemed necessary to make detailed measurements on all complete lens assemblies before their installation on the ring of supporting girders. For this purpose several measuring jigs were designed and are being built at Brookhaven. It is now planned to analyze the dimension measurements of the various lens assemblies and then selectively mount the lenses in such a fashion as to produce minimum deviations of the electron beam from a precalculated equilibrium orbit.

The vacuum control wiring is practically completed. The freon refrigeration system, as well as the chilled water system for the diffusion pump units, was installed. All the sixteen 10-in. mercury diffusion pumps were delivered, and one of these units is now undergoing tests on one complete section of vacuum enclosure. Tests at the factory showed that a blanked-off 10-in. diffusion pump unit will go down to a vacuum of 5×10^{-7} mm of Hg within 24 hr.

The rf accelerating cavity was installed on the accelerator ring. The power amplifier which will drive the cavity was constructed, connected, and tested. Initial squeaking and power supply difficulties were eliminated. The system is now delivering 800 v with a good wave shape which should provide ample safety factors for the 300-v/turn energy which will be required for machine operation.

The construction and testing program of the injection and inflection equipment continued. Apulsing system was incorporated in the electron Van de Graaff injector. Efforts are now being made to reduce the effective size of the pulsed Van de Graaff beam.

Control wiring progressed at a rapid rate. The control console and its associated racks were set up, and many accelerator components have been wired up to this focal control point.

AGS Project

Development work on all phases of the synchrotron continued. The Stone and Webster Engineering Corp. has been retained to do the architect engineering work for the

*For explanation of these numbers, see Foreword.

Figure 1.

AGS building complex. Three engineers of this organization have been in residence at Brookhaven since August 2, 1954 and will remain here for the duration of the job, since it is felt to be imperative that the design of the buildings be performed in close cooperation with the Accelerator Development staff.

Magnet

The magnet group is continuing and refining the development of magnetic measuring and standardization equipment. Extensive design and analysis were done over the summer to determine the most appropriate search coil shapes and combinations for the dynamic measuring program. The servo-fluxmeter was completely rebuilt and now has a sensitivity of about 15 lines.

Measurements were completed on Model Magnet No. 2, Configuration 2. Some of these measurements were described in previous quarterly reports. The latest data indicate that the reversal of the pole-face contours, which will be necessary to obtain the strong focusing effects, does not cause change in the fields in the magnet gap that is to any extent detrimental to the proton orbit stability. The information gained from the dc models and some theoretical calculations have been utilized in the design of the first dynamic model. This design has been completed and the dynamic model fabrication will start in the near future. The model will have two sets of coils (144 turns and 32 turns) in order to make possible eddy current measurements as well as remanent field studies. Power systems for both study programs have been designed and are now being assembled.

Linear Accelerator

The linear accelerator group completed the construction of the ion source and the pre-accelerator for the model linac. This system is now undergoing tests to determine operating conditions for maximum beam intensity. A first model of an actual linac section containing 46 drift tubes is now under construction in the shop. At the same time a 90-mc power oscillator is being designed to supply rf power to this model. It is expected that valuable information will be obtained on linac power losses and proton dynamics.

A review of possible power sources for the linac made it evident that a single 3-Mw, 200-mc klystron supply would be desirable. However, no klystrons in this frequency range and at this power level are in existence at this time, and it was decided that a power supply utilizing 3W10,000A3 triodes should be built for the machine. Such a power system was designed and a model triode power oscillator amplifier is now under construction.

The development program of permanent magnet focusing rings has produced cylinders of Cunico and Alnico V, 5 in. long, 1 in. ID and 1-5/8 in. OD, which provide gradients of 700 to 800 gauss/cm in the center. From present experiments, it seems that a similar cylinder made of Ferroxdur ferrite will produce gradients of 1000 gauss/cm before it saturates. These gradients will be sufficiently large to produce the required focusing inside the linac drift tubes.

Radiofrequency

A model ferrite ring (20 cm ID and 35 cm OD) was received from the Philips Corporation of Eindhoven, Holland. This ring was tested and showed the same rf characteristics which had been measured on smaller samples of the same ferrite. Forty similar rings for a model accelerator cavity have been ordered and are expected to be delivered early in January 1955.

Theoretical

A preliminary analysis was made of possible eddy currents in a 0.050-in. thick stainless steel vacuum chamber. The results indicated that these eddy currents would not be negligible, but that they could probably be corrected with small correction windings.

Relaxation method studies showed good agreement with the magnetic measurements taken on Model Magnet No. 2. Consequently, this method of analysis was used to determine the pole contours for the next set of pole faces to be used on the dc model.

Studies are now in progress to develop a good method of analysis to provide a "stacking order" of strong focusing magnets or electrostatic lenses. It is expected that this analysis will determine the selective mounting which will be used in the erection of the magnet ring of the AGS or the electrostatic lens element of the electron analogue.

Computations of proton orbits under various operational conditions are continuing on the New York University Univac electronic computer installation.

The BNL Remington Rand electronic computer is being used to determine the most effective lengths of AGS magnets and straight sections for maximum orbit stability.

The following abstracts summarize work submitted for journal publication or presentation at scientific meetings by members of the Chemistry Department, including visitors, during the period July 1 - September 30, 1954. The subjects covered lie in the fields of: nuclear chemistry; hot atom chemistry; radiation chemistry; isotope chemistry and tracer studies, mass spectrometry; and neutron diffraction.

DISINTEGRATION OF BISMUTH BY 2.2-BEV PROTONS. N. Sugarman,** R.B. Duffield,† G. Friedlander, and J.M. Miller.†† Submitted to Phys. Rev. Letter.

Cross sections are reported for the formation of about 20 nuclides produced by the bombardment of bismuth with 2.2-Bev protons at the Cosmotron. The results indicate that spallation with the emission of a large number of nucleons is much more probable at this energy than at a bombarding energy of ~ 400 Mev, whereas fission into two fragments of approximately equal mass is less probable at the higher energy. The yield-vs-mass curve does not show a clear division between the spallation and fission regions. Some data on recoil ranges of radioactive products are reported.

**Institute for Nuclear Studies, University of Chicago, Chicago, Ill

†Physics Department, University of Illinois, Urbana, Ill.

††Chemistry Department, Columbia University, New York, N.Y.

EXCITATION FUNCTIONS OF RADIOACTIVE PRODUCTS FROM THE INTERACTION OF HIGH ENERGY PROTONS WITH LEAD. E. Baker, A. Caretto, J. Cumming, G. Friedlander, J. Hudis, and R. Wolfgang. Submitted for presentation at the Am. Phys. Soc. meeting, November 1954.

Excitation functions of radionuclides produced by proton bombardment of lead at energies ranging between 0.6 and 3 Bev are presented. Yields of isotopes of F, Na, Mg, and P increase rapidly from the order of 10 microbarns at 600 Mev to the order of a millibarn at about 2 Bev. Thus, at these high energies production of such light elements, which cannot result from the usual spallation or fission processes, becomes a relatively probable event. In the medium mass region, from Mn to Ba, cross sections of neutron excess isotopes, as produced by 340-Mev Bi fission (W.F. Biller, UCRL 2067), remain approximately constant or decrease by a factor of about two with increasing energy. By contrast, yields of neutron-deficient nuclides increase rapidly until at about 2 Bev they become comparable with those of the neutron excess nuclides. No clearly defined fission peak like that which appears with Bi at 340 Mev was found with Pb at Bev energies.

RADIOCHEMICAL STUDIES OF NUCLEAR REACTIONS AT BEV ENERGIES. G. Friedlander. To be presented at Natl. Acad. Sci. meeting, November 1954.

Determinations of the yields of radioactive end products have been used in the study of interactions of complex nuclei with protons of 0.4- to 3.0-Bev kinetic energy. In the bombardment of both copper and lead targets with 2.2-Bev protons, products 20 or more mass numbers below the target mass are formed with much higher cross sections than those obtained in similar bombardments with 300- to 400-Mev protons. In the case of lead targets, even such low-mass nuclides as F^{18} , Na^{24} , Mg^{28} , and P^{32} are formed with cross sections of the order of 1 millibarn. In the yield-vs-mass

*For explanation of these numbers, see Foreword

curve of the products from lead bombardments, there is no pronounced peak which could be clearly associated with fission into two almost equal fragments. Measurements of recoil energies of certain products such as barium isotopes also show that these are not formed by a fission process as they are at lower bombarding energies. The excitation functions for a number of reactions, such as the formation of the low-mass products from lead, rise steeply with proton energy above about 0.5 Bev, whereas the cross sections for relatively simple reactions in heavy elements decrease with increasing proton energies. The results may be interpreted to indicate that very large energy transfers (upward of hundreds of Mev) to the struck nuclei are quite probable with bombarding energies in the Bev region. Some characteristic reactions, particularly the formation of some α -emitting rare earth activities in heavy element targets, are useful as monitors for high energy beams.

AN ATTEMPT TO DETECT THE ANTINEUTRINOS FROM A NUCLEAR REACTOR BY THE $\text{Cl}^{37}(\bar{\nu}, e^-)\text{A}^{37}$ REACTION. R. Davis, Jr. Submitted to Phys. Rev.

Tanks containing 200 and 3900 liters of carbon tetrachloride were irradiated outside the shield of the Brookhaven reactor in an attempt to induce the reaction $\text{Cl}^{37}(\bar{\nu}, e^-)\text{A}^{37}$ with fission product antineutrinos. The experiments served to place an upper limit on the antineutrino capture cross section for the reaction of $2 \times 10^{-42} \text{ cm}^2$ per atom. Cosmic-ray-induced A^{37} was observed and the production rate was measured at an altitude of 14,100 ft and at sea level. Measurements with the 3900-liter container shielded from cosmic rays by 19 ft of earth permit placing an upper limit on the neutrino flux from the sun.

K-ELECTRON EJECTION IN THE BETA DECAY OF S^{35} . W. Robinson and J.J. Howland, Jr.* Submitted to Phys. Rev.

By means of proportional-counter pulse-height analysis, it is found that S^{35} sources emit K-photons of chlorine with a probability of $2.47 (\pm 0.6) \times 10^{-4}$ photons/beta (the indicated error being an extreme), equivalent to a probability for K-hole production of $2.3 (\pm 0.7) \times 10^{-3}$ K-holes/beta. The theory, which has about the same accuracy as the experiments, predicts 3.0×10^{-3} K-holes/beta.

*Present address: Radiation Laboratory, University of California, Berkeley, California.

RADIOACTIVE ANTHRACENE- C^{14} AND ACRIDINE- C^{14} FROM THE NEUTRON IRRADIATION OF ACRIDINE. A.P. Wolf and R.C. Anderson. Submitted to J. Am. Chem. Soc.

Anthracene- C^{14} has been produced by neutron irradiation of the nitrogen homolog, acridine. The recoiling C^{14} entities produced by the $\text{N}^{14}(\text{n}, \text{p})\text{C}^{14}$ reaction were also found in acridine molecules, where the nitrogen had not undergone neutron capture during the course of irradiation. The ratio of activity as acridine to the activity as anthracene was about 19:1. Of the total activity produced in the samples, 3.5 percent was found as acridine, and 0.2 percent was found as anthracene.

A ONE-STEP METHOD OF SYNTHESIZING TRITIUM-LABELED COMPOUNDS. F.S. Rowland,* C.N. Turton,* and R.L. Wolfgang. Submitted for presentation at the Am. Chem. Soc. meeting, September 1954.

A method of general applicability for the one-step synthesis of tritium-labeled compounds is proposed. The material to be tagged and a lithium compound are mixed together in a finely divided state, such as a solution, emulsion, slurry or fine powder, and are then exposed to neutrons. The $\text{Li}^6(\text{n}, \alpha)\text{T}$ reaction has a large slow-neutron cross section for the production of quite energetic tritons. The range of these particles

is sufficiently large that they are able to escape from the matrix of the lithium compound into that of the desired material. Some of these tritons, while possessing chemically significant amounts of energy, can then substitute into this compound for ordinary hydrogen without otherwise disrupting the molecule.

This method may have useful applications for the introduction of radioactivity into complicated molecules, especially for those materials that are difficult to synthesize chemically and that have a very low yield biosynthetically.

The technique has been used to synthesize a labeled aldohexose with an activity of 10^6 dps/mole of sugar. The compound contained in nonlabile positions is approximately 1 percent of the total number of tritons produced. Specific activities higher by several factors of ten can probably be produced under appropriate conditions.

*Princeton University, Princeton, N.J.

CHEMICAL EFFECTS OF NEUTRON CAPTURE IN SOME COBALTC HEXAMMINES. A.V. Zuber* and J.M. Miller.** Submitted for presentation at the Am. Chem. Soc. meeting, September 1954.

The chemical fate of radioactive Co^{60} formed through neutron capture by Co^{59} in cobaltic hexammine, tris-ethylenediamine, and bis-diethylenetriamine, both as crystals and in aqueous solution, has been investigated. In the irradiation of each of the three complex ions in aqueous solution, it was found that essentially all the Co^{60} formed can be carried down on a precipitate of cobalt hydroxide and is thus in a different chemical form from the parent compound. The radioactive products formed in the irradiation of solid cobaltic tris-ethylenediamine nitrate can be separated into three distinct chemical fractions after the target material has been dissolved in water:

- 1) Co^{60} carried along with a precipitate of cobalt hydroxide;
- 2) Co^{60} not carried by a cobalt hydroxide precipitate, but separable from the parent compound by fractional crystallization;
- 3) Co^{60} in a form that cannot be chemically distinguished from the parent compound.

The amount of the radioactive cobalt found in the third chemical fraction, R_3 , grows in the irradiated solid at the expense of the activity in the first fraction at a rate that can be expressed as

$$R_3 = A_1(1 - e^{-k_1 t}) + A_2(1 - e^{-k_1 t})$$

Investigation of the radioactive products formed in the neutron irradiation of solid dextro-rotatory cobaltic tris-ethylenediamine nitrate showed that more than 90 percent of the radioactive cobalt that reforms the parent compound does so with retention of the original optical configuration.

*Present address: Union Carbide and Carbon Corp., 30 East 42nd St., New York, N.Y.

**Department of Chemistry, Columbia University, New York, N.Y.

THE RADIATION CHEMISTRY OF NITRITE ION IN AQUEOUS SOLUTION. H.A. Schwarz and A.O. Allen. Submitted to J. Am. Chem. Soc.

When de-aerated solutions of KNO_2 at their natural pH are irradiated with γ - or x-rays, the main initial reaction is a formation of hydrogen and hydrogen peroxide. As peroxide builds up, a radiation-induced reaction between nitrite and peroxide sets in, which results in the formation of nitrate. This reaction is slower the higher the nitrite concentration. The kinetics of the nitrite-peroxide reaction were studied in detail by the irradiation of synthetic mixtures, and the results were satisfactorily explained

by a free radical mechanism. The main feature of the mechanism is a competition between H_2O_2 and NO_2^- for reaction with H atoms generated in the water. The yield for H-atom formation is found to be 2.4 per 100 ev. The initial yield of peroxide in the pure nitrite solutions is greater than the yield of hydrogen, and decreases with increasing nitrite concentration, at concentrations below 0.002 M. The effect of nitrite on the initial peroxide yield is qualitatively similar to the effect of bromide reported by Sworski. In nitrite solutions containing oxygen, an immediate nitrite oxidation occurs on irradiation, with a yield which increases with increasing ratio of the concentration of nitrite to that of oxygen.

PHOTOINITIATED OXIDATION OF BENZOIC ACID BY FERRIC IONS IN AQUEOUS SOLUTION. J. Saldick and A.O. Allen. Submitted to J. Am. Chem. Soc.

The photochemical action of $\text{Fe}(\text{OH})^{+2}$ on organic substrates was thought by Evans and co-workers to proceed by free OH radicals. The kinetic data of these workers on such reactions were consistent with this idea, but they failed to rule out the possibility that the organic molecule may react with an excited ion without the free radical intermediary. It is here reported that the rate of the photoinduced reduction of ferric ion by benzoic acid is independent of the concentrations of the reactants as long as the ratio of these concentrations is kept constant. This result is in agreement with the free radical hypothesis but not in agreement with the hypothesis of direct reaction with a photoactivated ion.

BACKSCATTERING FROM TARGETS OF LOW ATOMIC NUMBER BOMBARDED WITH 1- TO 2-MEV ELECTRONS. J. Saldick and A.O. Allen. Submitted to J. Chem. Phys.

Data for 1- and 2-Mev monoergic electrons on the fraction backscattered at perpendicular incidence, from targets of Mg, C, Be, and (at 1-Mev only) Li, show that the fraction increases nonlinearly with Z in the low Z range.

A SEMI-EMPIRICAL STUDY OF THE H_2Cl TRANSITION COMPLEX THROUGH THE USE OF HYDROGEN ISOTOPE EFFECTS. J. Bigeleisen and M. Wolfsberg. To be presented at Natl. Acad. Sci. meeting, November 1954.

Linear and triangular structures for the H_2Cl transition complex have been analyzed in terms of normal vibration theory and by an evaluation of the experimental data on the effect of hydrogen isotope substitution on the rate of reaction of hydrogen molecules and chlorine atoms. The frequencies of the bending vibrations are probably sufficiently small that their contribution to the relative rates can be expressed in terms of a small quantum correction of the order of

$$\left(\frac{hc}{kT}\right)^2 \left(\frac{\omega_{\text{H}}^2 - \omega_{\text{D,T}}^2}{24}\right)$$

The near classical behaviour of the bending frequencies for both linear and triangular structures of the H_2Cl complex serves to reduce the ratio of the frequency factors, $A_{\text{H}}/A_{\text{D,T}}$, in the Arrhenius equation below that expected from structural considerations alone. Quantitative agreement is found between the calculated and experimentally determined frequency factor ratios. The "symmetrical" stretching frequencies, for both linear and triangular complexes, respectively, have been evaluated from the relative rates of reactions of H_2 and HT with chlorine atoms at 0°C and the known properties of the isotopic hydrogen molecules. This empirically evaluated parameter, together with thermodynamic data and an estimate of the tunnel correction, suffices for a calculation of the relative rates of reaction of the isotopic hydrogen molecules with

chlorine atoms as a function of temperature. Good agreement is found between the calculated rates for both linear and triangular structures and the available experimental data on HD, D₂, and HT.

COMPLEX IONS AND ELECTRON EXCHANGE. R.W. Dodson. To be presented at Natl. Acad. Sci. meeting, November 1954.

Some results of a continuing investigation of the kinetics of electron exchange reactions between different oxidation states of elements in aqueous solution will be presented. These reactions, like many nonequilibrium oxidation processes, are markedly affected by anions capable of forming complexes with the reactants. In most of the known cases the electron exchange is accelerated. In some cases, however, a strong inhibition is observed. It proves convenient to relate the kinetic effects to the equilibria of complex formation.

The two-electron transfer between thalious and thallic ions has been studied from this point of view in the presence of chloride and of cyanide. The one-electron transfer between ferrous and ferric ions has been investigated in the presence of thiocyanate, bromide, and oxalate. The results are compared with earlier work on the thallium, iron, and other exchange systems, and, to the extent possible, with existing theory on the rates of electron transfer. Some rather striking relations between the heats and entropies of activation in the different systems are found.

THE MECHANISM OF THE RING CONTRACTION OF TROPOLONES. W. von E. Doering* and D.B. Denney.** Submitted for presentation at the Am. Chem. Soc. meeting, September 1954.

Two mechanisms have been proposed for the rearrangement of tropolones to benzoic acid derivatives (W.E. von Doering and L.H. Knox, J. Am. Chem. Soc. 73, 828 (1951); 74, 5683 (1952)). The first mechanism, A, envisions an attack by base on the carbonyl carbon followed by rearrangement with expulsion of X⁻ to give the product. The second mechanism, B, suggests an attack by base at carbon-2 followed by a concerted shift of atomic positions to yield the benzoic acid derivative. The use of carbon-14 provides a means of distinguishing between the two proposed mechanisms, and is the subject of this study.

Rearrangement of 2,4,7-tribromotropone-1-C¹⁴ to a mixture of 2,4- and 2,5-dibromobenzoic acids, followed by hydrogenolysis of this mixture, yielded benzoic acid, which underwent a Schmidt reaction to carbon dioxide and aniline (isolated as benzanilide). All the radioactivity was found in the carbon dioxide, whereas none was found in the benzanilide. Similarly, 2,7-dibromotropone-1-C¹⁴ was rearranged to o-bromobenzoic acid, which gave carbon dioxide containing all the radioactivity, and o-bromoaniline (isolated as the benzanilide) devoid of activity. These results definitely exclude mechanism B and are completely consistent with mechanism A, which is presumably the correct one.

*Sterling Chemistry Laboratory, Yale University, New Haven, Conn.

**Hickrill Chemical Research Foundation, Katonah, N.Y.

THE ACID CATALYZED REARRANGEMENT OF α -CINENIC ACID. J. Meinwald.*
Submitted to J. Am. Chem. Soc.

Tracer studies reveal that the isomerization of α -cinenic acid (I) to geronic acid (II) involves carboxyl transfer rather than methyl migration. Some mechanistic implications of this finding are discussed.

*Permanent address: Department of Chemistry, Cornell University, Ithaca, N.Y.

THE EFFECT OF ISOTOPIC SUBSTITUTION ON THE MASS SPECTRA OF MOLECULES. II. OXYGEN AND CARBON DIOXIDE, EXPERIMENTAL. O.A. Schaeffer and H.R. Owen. Submitted to J. Chem. Phys.

The relative changes in the probability of bond breaking following electron impact have been studied for the molecules $O^{16}O^{16}$ and $O^{16}O^{18}$, $O^{16}C^{12}O^{16}$, $O^{16}C^{13}O^{16}$ and $O^{16}C^{12}O^{18}$. The experimental results were obtained with a 60° mass spectrometer and with a source designed to equalize the collection efficiency of the fragment ions formed. The relative changes were studied systematically as a function of the various ion source voltages. In the case of oxygen, the ratio of bond breaking in $O^{16}O^{18}$ divided by that in $O^{16}O^{16}$ is 0.98. The formation of C^+ from $O^{16}C^{13}O^{16}$ is 1.022 times the formation of C^+ from $O^{16}C^{12}O^{16}$, while the formation of C^+ from $O^{16}C^{12}O^{18}$ is 0.997 times the formation of C^+ from $O^{16}C^{12}O^{16}$. The formation of CO^+ from $O^{16}C^{13}O^{16}$ is 0.974 times that from $O^{16}C^{12}O^{16}$, while the relative formation of $C^{12}O^{18+}$ from $O^{16}C^{12}O^{18}$ is 1.03 times and that of $C^{12}O^{16+}$ is 0.97 times the formation of CO^+ from $O^{16}C^{12}O^{16}$. Finally the relative formation of O^{16+} is 1.05 times and that of O^{18+} is 0.98 times the formation of O^+ from $O^{16}C^{12}O^{16}$. The values for $O^{16}C^{13}O^{16}$ are compared with previous values in the literature.

THE EFFECT OF ISOTOPIC SUBSTITUTION ON THE MASS SPECTRA OF MOLECULES. III. CARBON DIOXIDE, THEORETICAL INTERPRETATION. O.A. Schaeffer. Submitted to J. Chem. Phys.

The relative changes in pattern for O^{18} and C^{13} substituted carbon dioxide are calculated. The frequencies for the isotopic carbon dioxide molecules and the associated normal coordinates are obtained for a harmonic potential function. These are then used as the basis for the ground state vibrational wave functions. The bondbreaking probability integrals are evaluated for various values of the "critical" internuclear separation, r_c , and a study is made of the change in predicted bond breaking ratios as a function of r_c . The values of r_c are chosen which give the experimental pattern for $O^{16}C^{12}O^{16}$, and then the relative changes in pattern are calculated for the other molecules and are compared with experimental values. Especially striking is the fact that the calculation faithfully reproduces the change in pattern for the case in which a heavier molecule is more likely to fragment than a lighter one, as well as for the reverse case. It is concluded that a Franck-Condon type calculation adequately describes the electron impact process in carbon dioxide.

THE ISOTOPE ABUNDANCES OF CHLORINE FROM VARIOUS SOURCES. H.R. Owen and O.A. Schaeffer. Submitted to J. Am. Chem. Soc.

The isotope abundance of chlorine from marine, igneous, and sedimentary sources has been investigated. Rocks whose chlorine content ranged from less than .01 percent to 6.8 percent were leached with dilute nitric acid and the chloride converted to ammonium chloride, which was then reacted with sulphuric acid and methyl alcohol to form methyl chloride. The isotopic analyses were made with a 6-in. 60° spectrometer. Additional studies were made to insure against fractionation during sample preparation. Within the limits of experimental error (± 0.2 percent), no variation in the isotopic

abundances was observed. The abundances are compared with previous results and the origin of the samples. The average value of the $\text{Cl}^{35}/\text{Cl}^{37}$ ratio of over 100 determinations on 10 samples of different origin is 3.13.

THE MASS SPECTRUM OF LITHIUM IODIDE. L. Friedman. Submitted to J. Chem. Phys.

The mass spectrum of lithium iodide vapor produced from salt heated in an oven in the ion source of an analytical mass spectrometer was investigated as a function of oven temperature. Evidence was obtained for the existence in the vapor of both monomer and dimer. Partial heats of sublimation to monomer and dimer at 0°K were found to be 44.5 ± 7 and 45.0 ± 2 kcal, respectively. The heat of dissociation of $(\text{LiI})_2$ into gaseous monomers was found to be 44.0 ± 2.2 kcal. The partial heat of sublimation of monomer was used with supplementary thermochemical data to obtain values for the lattice energy and the heat of dissociation of the diatomic molecule.

Appearance potentials for the major ions in the spectrum were determined. Upper limits for the energy of dissociation of the gaseous diatomic molecule were derived from the appearance potentials of Li^+ and I^+ , which were slightly higher than the thermochemical value.

NEUTRON DIFFRACTION STUDIES OF FERRIMAGNETIC SPINELS. L.M. Corliss and J.M. Hastings. To be presented at the Naval Ordnance Laboratory Conference on Ferrimagnetism, October 1954.

A brief review of the theory of neutron scattering by magnetic substances will be presented. The available data on pure and mixed ferrimagnetic spinels will be summarized. In addition, a new low temperature magnetic transition in zinc ferrite and chromite will be discussed, and their application to the investigation of the B-B interaction will be indicated.

REACTOR DEPARTMENT

Foreword and General Notes

Detailed accounts of work being performed by the various sections comprising this Department are given below. Also included, under the heading, Neutron Physics -- Cooperative Programs, is the status of research being sponsored at Brookhaven by Columbia University and the General Electric Company's Research Laboratory. The spectrometer installation for use by the Bell Telephone Laboratories, Inc. has been completed and research is scheduled to begin shortly. Under the auspices of this Department but not detailed below, programs of research have been carried out by the California Research Corporation, Monsanto Chemical Company, Massachusetts Institute of Technology, Knolls Atomic Power Laboratory, and Westinghouse Atomic Power Division.

The US Weather Bureau continues the support of a unit, working with the Brookhaven Meteorology group. The US Geological Survey group stationed at this site has concluded compilation of a final report on research performed during the past few years. The final draft of this report has not yet been presented.

The summer brought the usual influx of visitors and summer staff members, many of whom engaged in research with the reactor facilities.

Expositions of the model of the reactor were given for: teachers taking summer courses at New York University; foreign students attending Massachusetts Institute of Technology; members of the Naval Reserve Seminar (which was held at Brookhaven); visitors from the US Naval War College; faculty members of the US Merchant Marine Academy; members of the Special Services group of the Armed Forces; personnel of Brookhaven's Medical Department; members of Congress, the US Atomic Energy Commission, the American Chemical Society, the Visiting Committee for the Engineering Departments, the Board of Trustees, The Research Corporation, the Pacific Northwest Power Group; and representatives of the respective atomic energy organizations from Australia, Brazil, England, Germany, India, Spain, Sweden, and Switzerland.

Discussions have been held on proposed cooperative research and/or the use of BNL's facilities with representatives of the Mound Laboratory, Westinghouse Atomic Power Division, Knolls Atomic Power Laboratory, Colgate-Palmolive-Peet Company, Hercules Powder Company, Glenco Corporation, Baker Platinum Company, duPont, American Smelting and Refining Company, the Oklahoma Agriculture and Mining School, and the General Electric General Engineering Laboratory. Discussions were also held with AEC personnel on general classification problems and the possibility of further declassification of Brookhaven's facilities.

Usage of the Brookhaven reactor was reviewed for a survey being made by the New England Committee on Atomic Energy. A forecast of capital equipment requirements has been prepared which covers the next fifteen months.

Preliminary plans have been discussed for participation in an international conference in connection with the "atom pool" proposal, originally outlined by President Eisenhower last December. Several specific topics covering programs in this Department which are considered classifiable and appropriate for such a conference have been outlined.

Several members of the Department submitted topics for presentation at the Symposium on Fission Physics and Classified Nuclear Physics for Reactors and Shielding, scheduled for October at Oak Ridge.

Members of the Meteorology group participated in a mathematical seminar on time series spectra at Storrs, Connecticut and at the meeting of the American Meteorological

Society in Columbus, Ohio. The Department was recently represented at the New York Meeting of the Atomic Industrial Forum, Inc.

Operations

(†)*

The reactor facilities continue to be utilized at almost capacity level by members of the various Laboratory scientific departments and outside organizations. Megawatt days accumulated from July 1 to September 30 totaled 2012. The maximum power level reached was 24.7 Mw and the average operating level was 23.92 Mw. Downtime, averaged over the period, equaled 8.8 percent.

Hurricane Carol, of August 31, caused no damage to the reactor other than an unscheduled shutdown because of power failure.

The water-cooled irradiation facility is being redesigned and fabricated. This is required because of the development of a leak in the original equipment. Temporary measures have been taken to allow continued use of this facility until the improved equipment is installed. However, the heavy demand for use of this facility has dictated the need for another one which is presently under construction.

Primary fan No. 3 was inspected, cleaned, and, in generally good condition, placed back in service.

Shipments of irradiated fuel to Oak Ridge have been continued. A cost reduction plan involving these shipments has been proposed for AEC approval.

Instrumentation improvements have included work on the galvanometer, the emergency shutdown system, the instrument power supply, and the rod position indicators.

Studies continue on the proposed reloading of the reactor. In this connection, consultations were held with personnel of the Hanford Atomic Products Division at that installation, and were followed by discussions with members of the AEC's Advisory Committee on Reactor Safeguards in Washington, D.C.

A schedule has been formulated for a repetition of last year's special maintenance experiment on the reactor, which will be carried out during the early part of October.

Isotope Distribution and Special Materials Group

(†)

AEC-sponsored discussions have continued on the formulation of a satisfactory over-all pricing policy. The long postponed issue of the brochure describing Brookhaven's services and facilities is being revised again in line with the above discussions.

A forecast of BNL's requirements for MTR irradiation services was prepared at AEC request.

Special and basic quota requirements for S/F materials for the calendar year of 1955 have been determined and submitted to the AEC. By request, this Department submitted a review of a proposed AEC "Isotope Order Form" and also a survey of Brookhaven's needs with respect to a proposed expansion of the stable isotopes production program at Oak Ridge.

In response to a request by the University of North Carolina, a review of Brookhaven's procedures and forms for scheduling and handling irradiations was prepared to serve as a guide for setting up similar operations at that institution.

Return of S/F materials, now surplus, will be scheduled in the near future.

Table 1 gives a summary of the service irradiations and reactor rentals for the first quarter of fiscal year 1954-1955.

On September 1, 1954 a new price schedule was put into effect which covers standard reactor units as suggested by ORNL. This price schedule allowed for a premium

*For explanation of this symbol, see Foreword.

Table 1		
Off-Site Irradiations and Rentals		
	No. of Shipments	\$ Billed
Target Conveyor	199	10,050.39
Pneumatic Tube	20	826.71
Air-Cooled Holes	5	1,293.00
Water-Cooled Holes	1	85.00
Thermal Column	6	212.00
Experimental Holes	10	2,459.40
Multicurie Co ⁶⁰	2	7,740.00
Animal Tunnel	1	568.80
Core Hole	1	925.00
I ¹³²	3	445.00
Special	14	522.59
Cyclotron	1	55.00
Rentals	(8)	13,414.00
	263	38,596.89
On-Site Service Irradiations for the Same Period		
Target Conveyor	154	10,658.00
Pneumatic Tubes	306	1,530.00

extensive effort is being made to obtain accurate radiation widths of resonances. The interest lies both in the variation of radiation width with atomic weight and in the variation from resonance to resonance with a particular isotope. Resonances in In¹¹⁵, Eu¹⁵¹, Ir¹⁹¹, and Ir¹⁹³ have been studied extensively. A program for obtaining highly accurate measurements of potential scattering as a function of atomic weight is in progress. These results are of particular interest at present because of their relation to contemporary theories of nuclear models.

Scattering Cross-Section Measurements

During this period scattering data runs on the 1.260-ev resonance in Rh were completed. (Before analysis can be completed a high energy total cross-section measurement is needed with a suitably thick sample. This thick sample is not presently available.)

Development work was undertaken on a new scattering experiment, in which the scattering sample will be mounted on the spectrometer arm at a glancing angle to the monochromatic neutron beam. This arrangement then presents a thick target to the incident neutron beam and a thin section to the detector at right angles to the monochromatic beam. This "thick-thin" experimental technique permits the observations of σ_s/σ_t and appears to present a simpler analysis than the previously used "bright-line" technique. Also, higher absolute counting rates are realized as well as a more favorable ratio of net to background counts.

All qualitative work has been completed and data runs are to be started the latter part of October upon completion of the equipment. (H.L. Foote, Jr. and J. Moore)

to be placed upon higher flux reactors. At the same time, Brookhaven's handling charge was reduced from \$25.00 to \$20.00.

Neutron Physics (5200)*

Instrument Development

The new automatic control system for the crystal spectrometer has been installed and is in routine operation. Data accumulated by the spectrometer are now recorded directly on punched cards. The data reduction is subsequently carried out on the Remington-Rand computer.

Total Cross-Section Measurements

The major emphasis on the total cross-section program has been shifted from survey-type measurements to detailed measurements of individual resonances which for one reason or another are of special interest. For example, an

*For explanation of these numbers, see Foreword

Analysis of Data

The computing program for analysis of resonant structures is progressing at a discouraging rate. The bottleneck lies in the lack of flexibility of the Remington-Rand computer which is used for the computations. In particular the small storage capacity (memory) and the limited number of program steps on each computing routine makes the programs awkward and limits the cases which can be analyzed. Efforts are under way to improve the existing programs in order to increase the output of results; however the primary limitation is machine capacity rather than technique in programming.

Neutron Physics -- Cooperative Programs

(5200)

Columbia University Crystal Spectrometer

A. Nonmagnetic Neutron-Electron Interaction - The liquid bismuth sample holder for the determination of the nonmagnetic neutron-electron interaction has been completed at Columbia University and will be installed on the spectrometer. In preparation for this experiment, the total cross section of solid bismuth has been measured over the energy range, 0.1 to 10.0 ev. The data obtained at energies above 0.3 ev, which is the neutron energy interval of interest for the liquid bismuth measurement, were reproducible to 0.15 percent. At neutron energies lower than 0.3 ev, the variation in the transmission of the solid bismuth sample was larger since each measurement depended upon the crystal orientation in the irradiated volume of the sample. (B.M. Rustad, E. Melkonian, and W.W. Havens, Jr.)

B. Scattering Experiment - Measurements have been made of the neutron scattering from gold with a rough scattering apparatus. The scattering cross section σ_s was obtained by detecting neutrons scattered from thin gold targets with a BF_3 counter which was well shielded from the monochromatic neutron beam. The ratio σ_s/σ_t was also determined by measuring the scattering from thick targets of Au and Ni_2B at several energies. Preliminary results are in agreement with published data and indicate that the neutron intensity of the crystal spectrometer is sufficient for precise scattering measurements. (J. Moore, W.W. Havens, Jr., B.M. Rustad, and C. Sheer)

C. Transmission Measurements at Long Wavelengths - The program, in conjunction with the Physical Chemistry Department of Columbia University, to study the effect of molecular binding on the proton cross section at long neutron wavelengths is being continued. A detailed investigation is being made of neutron filters to attenuate the higher order Bragg reflections from the mica crystal monochromator. Present results show that neutron beams with less than 4 percent order contamination can be produced between 4-7 Å with filters of Be or Be in combination with BeO. Other materials such as Bi, Mg, MgO, and graphite are being studied in order to obtain suitable filters in the 7-12 Å range. (F. Gould and T.I. Taylor)

He⁶ Beta-Ray Spectrum (Columbia University)

The shape of the beta-ray spectrum from He⁶ is being reinvestigated to determine the relative magnitude of the coupling constants, G_T , G_A , and G_P of the tensor, axial vector, and pseudoscalar invariant respectively in the beta-decay interaction. Previous measurements (Wu, Rustad, Perez-Mendez, Lidofsky, Phys. Rev. 87, 1140 (1952)) determined the endpoint of the spectrum to be 3.50 ± 0.05 Mev, but primarily because of the lack of available source strength, small deviations from a purely statistical beta-spectrum shape could not be detected. The present He⁶ generating apparatus at the Brookhaven reactor delivers approximately 8×10^7 He⁶ atoms/sec, which is

sufficient for a precise measurement with the Columbia 180⁰ beta spectrometer. The spectrometer has been carefully calibrated and is now being installed at the reactor. (A. Schwartzchild, O. Kistner, and B.M. Rustad)

General Electric Neutron Diffraction Group Investigations

Structure of Magnetic Perovskites - A wide variety of compounds of nominal composition RMO_3 which crystallize in a pseudo-perovskite (CaTiO_3) structure can be prepared in which R is a large positive ion (Ca, Sr, Ba, La) and M is a small positive ion (V, Cr, Mn, Fe, Co, Ni). The pure materials are antiferromagnetic, but solid solutions may be ferromagnetic, and the saturation magnetization and Curie temperatures are monatomic functions of the composition. To ascertain the nature of the magnetic interactions in these systems, neutron diffraction data have been obtained from powder samples of LaMnO_3 , BaMnO_3 , $\text{La}_{0.7}\text{Ba}_{0.3}\text{MnO}_3$, LaFeO_3 and LaCrO_3 .

Single Crystal Neutron Diffraction Study of NiO - Nickel oxide is antiferromagnetic and the magnetic structure, as deduced from neutron diffraction data obtained from the powder, has been reported by Shull, Strauser, and Woolan (Phys. Rev. 83, 333 (1951)). Neutron diffraction data from single crystals of NiO (supplied by G.L. Clark of the Linde Research Laboratory) have been obtained, and interpretive techniques for correcting the intensities for extinction have been developed. The single crystal data give evidence for the presence of antiferromagnetic domains in the crystals, and the investigation is being continued in order to deduce more fully the nature of the magnetic ordering. (W.L. Roth)

Shielding

(4570)

Progress of this program is reported in the classified section, BNL-309.

At the AEC's request, sections of the proposed Shielding Handbook were reviewed by this group. Several declassification requests are being prepared as a result of a survey made of the classified shielding literature. The personnel recruitment program has resulted in the addition of two scientists to the staff of this group.

Meteorology

(6500)

Oil Fog Program

Final arrangements for the computation of the power spectra of oil fog concentrations have been made and approved. The Meteorology group will utilize approximately 40 hr on the Atomic Energy Commission Univac during the current fiscal year to determine the spectra associated with approximately twenty-five 2-hour field tests. In this project, it is most desirable to obtain and process simultaneous wind and oil fog data, but the task is a formidable one even with the assistance of the best computers. For this reason, the US Weather Bureau Office at Upton and the Meteorology Group will pool their efforts and resources in a cooperative project, the former handling the wind data, and the latter the oil fog concentration data. The Scientific Services Division of the Weather Bureau has arranged for 40 hr on the Census Univac, so that between the two organizations complete analyses for each run will be made.

The second portion of the oil fog sampling program, involving the preparation and use of filter samplers, has been expanded to include 30 sampling units. These units are now completely assembled and ready for field operations. A flat-bed trailer has been altered to simplify transporting the equipment.

Hurricane Data

The Brookhaven area was subjected to unusually high winds on two occasions, August 31 and September 11. In the first storm, winds at the top of the 410-ft tower reached 125 mph, and in the latter 96 mph. Such events are fortunately extremely rare, but even more rare are detailed records of the wind fluctuations associated with them. In both of these cases, it was possible to obtain accurate records from several levels of the 410-ft tower at a maximum chart speed (3 in./min). This provides an open record from which it is possible to evaluate individual gusts of 1- to 2-sec duration.

Although work on this type of material is in a sense a diversion from the established program of the group, it was decided that intensive effort should be devoted to these particular wind data. Knowledge of short term gusts is of considerable importance in the design of structures that must withstand heavy wind loads. Although it is comparatively easy to take care of direct sustained thrusts, little is known of the frequency of strong gusts, and therefore it is difficult to design for the resonance problem. The data from both hurricanes are suitable for this type of analysis, and the results of the study will be published.

DAF Meteorology Study

A new program involving the processing of the punch cards obtained in an earlier study (BNL 288) was undertaken officially on July 1. This work is also sponsored by the Geophysics Research Division of the US Air Force. It will involve a large amount of machine processing to prepare the climatological studies, and the decision has been made to accomplish as much of this work as possible by means of the Laboratory computer facility. Outside assistance will be utilized only when the work load becomes excessive, or for operations that cannot be performed efficiently within the Laboratory.

The first portion of the work has consisted of providing the punch cards with additional index and selector groups for specific use in the new project. Most of these have been relatively simple to complete, but the development of a cloud index combining present and past cloudiness has become a rather involved computation.

Soil Temperature Measurements

Soil temperatures are being obtained routinely from four stations on the perimeter of the new accelerator site. Temperature differences between stations have not exceeded 3.0°C at identical levels below the surface, but the long period variations are already an item of special interest. The thermal diffusivity of the soil, when calculated on the basis of short term (diurnal) changes, agrees closely with values obtained by others for sandy soil. However, this quantity cannot be used to predict the annual temperature cycle, which shows a much higher diffusivity, of the order of granite.

The effect is difficult to account for at present, but it is of considerable importance in terms of foundation design, since it implies larger temperature variations at great depths than would have been anticipated on the basis of theoretical calculations.

Publications

"A summary of neutron resonances in the rare earth elements," V. Sailor, H. Landon, and H. Foote, submitted to Phys. Rev.

"Radiation width variations in the neutron resonances of Eu and In," H. Landon and V. Sailor, Bull. Am. Phys. Soc. 29, 7, B10 (1954). ✓

"Problems in the analysis of neutron resonance data," V. Sailor, H. Landon, H. Foote, and R. Wood, Bull. Am. Phys. Soc. 29, 7, B9 (1954).

"Instrumentation and control of the Brookhaven reactor," J.E. Binns, submitted to Inst. Radio Engrs.

"Computation of soil moisture at Brookhaven National Laboratory," R.M. Brown, submitted to Monthly Weather Review.

"Availability of meteorology punch card data," M.E. Smith and I.A. Singer, submitted to Am. Meteorol. Soc.

NUCLEAR ENGINEERING DEPARTMENT

This report includes some items of general interest as well as the unclassified work of the Nuclear Engineering Department on fission products utilization, polymerization, the fixation of fission products in clays for ultimate waste disposal, heat transfer in spray columns, the preparation of special radioisotopes, and the Hot Laboratory operations.

The classified work of the Department on exponential lattices, water and slightly enriched uranium, the uranium-bismuth liquid metal fuel reactor components development, various fuel processing systems, and radiation effects on reactor materials, is covered in a separate classified progress report. The last one, BNL 309, was issued as of August 15, 1954.

As a result of numerous discussions with the staff of the Reactor Division, Washington AEC, and with the AUI Visiting Committee to the Department, a special group is being organized to evaluate the work done to date on the LMFR and to report on the feasibility of constructing such a system for power production. The study group will consist of experienced engineers and scientists from several fields related to reactor development. It is hoped that one engineering organization will be able to assemble and organize the group and to be responsible for the prosecution of the work and the submission of a report to the Laboratory and the AEC by June 30, 1955. All pertinent research and development work done to date will be surveyed and an analysis prepared, not only of what has been accomplished but also of what remains to be done before an LMFR of this type can be built. The final report should contain:

1. A preliminary design for an LMFR system which will include the relative arrangements of core and blanket vessel, heat exchangers, pumps, piping, and process plants.
- 2.. Recommendations covering the development work which remains to be done and indicating the most critical items to be completed a) before an LMFR experiment is justified, and b) before mechanical engineering design of a full-scale plant is undertaken.
3. Preliminary evaluation of the feasibility and economic potentialities of LMFR.

All the papers on LMFR which were declassified for presentation by the Nuclear Engineering Department at the meeting of the AIChE at Ann Arbor in June were published as a group in a slightly condensed form in the July 1954 issue of Nucleonics. The abstracts of these papers were given in an earlier progress report (BNL-289). In addition, an unclassified paper was published, entitled "The energy-dependent Boltzmann equation applied to criticality calculations for bare graphite-moderated reactors," by J.A. Fleck (BNL 292).

Special Radioisotopes Development

(†)*

Iodine-133

Present results indicate that high yields of iodine can be obtained by oxidizing the irradiated uranium and heating to at least 900°C for at least one or two hours. Although this method is feasible and will be used to meet the present demand for this isotope, alternative methods are also being considered for the future.

*For explanation of this symbol, see Foreword.

Yttrium-90

The problem of making hard uniform clay spheres, 0.035 in. in diameter, containing stable yttrium was solved. Such spheres can now be made available for medical research and other purposes. The diameter may be varied within limits. These spheres when irradiated in the Brookhaven reactor will have about 0.5 to 1.0 mc of Y^{90} per mg of finished particle. Approximately 70 such irradiated beads were made and supplied to a medical research group at the Columbia Presbyterian Hospital in New York City.

A miniature solvent extractor was fabricated to be used in developing a simple milking system for obtaining fission product Y^{90} .

Fluorine-18

Development work was completed on the production of physiologically compatible solutions of 112-min F^{18} , and as soon as an interim price is approved, this isotope can be made generally available.

Cesium-137

Some work was started in an effort to develop an economical method for the extraction of cesium-137 from molten sodium nitrate.

Beta Coating for Balloons

The problem of applying relatively large amounts of pure beta emitter to the surface of a thin rubber balloon was undertaken. A balloon coated completely or in part could be introduced into any of the various body cavities and when inflated would put the active material in direct contact with tumors growing on the inner surface of such cavities.

Magnesium-28

The next isotope to be developed will be Mg^{28} in which several medical investigators have expressed very keen interest.

(L.G. Stang, Jr., W.D. Tucker, H.O. Banks, Jr., R.F. Doering, T.H. Mills)

Radiochemical Analytical Section

The following three problems were undertaken during the last review period.

Determination of the Insensitive Volume of the End Window Proportional Counter

The counting rate at some distance from a geiger tube is always higher than the calculated rate based purely on a geometrical extrapolation from the observed counting rate close to the tube. An investigation was undertaken to determine whether the end window proportional counter would display this same sort of behavior.

Essentially weightless samples of C^{14} and P^{32} were mounted on very thin backing and counted at ten different distances (from 0.236 cm to 5.316 cm) from the window of a proportional tube. The counting rates were corrected, first for absorption and then for geometry, by means of the factor $2R/h$, where R is the distance from a point source to a point on the circumference of the window, h is equal to $R - x$, and x is the distance from the source to the center of the window. The data indicate that, as with the geiger tube, the observed counting rate at a given distance, after all corrections are made, is

higher than the rate calculated by extrapolation of the count observed at some closer distance.

These results can be explained either on the basis of a fixed dead volume in the tube or on the assumption that a particle must traverse a certain minimum distance through the counting gas (P-10 in this case) to be detected.

In order to test these hypotheses, a series of 15 brass rings with diameters ranging from 1.270 to 2.200 cm were placed so as to define the window opening. The counting rate was determined with each ring with the source at each of seven different positions. If the increased counting rate noted above is the result of a necessary minimum trajectory of the beta particle through the counting gas, then the minimum possible path length of the beta should be constant at various distances from the tube when the aperture is made just small enough to result in 100 percent counting efficiency for that particular distance. This was found to be the case, and the average value for this minimum trajectory was calculated to be $0.43 \text{ cm} \pm 0.04 \text{ cm}$.

Polarographic Determination of Zirconium

A polarographic determination of zirconium can be carried out in aqueous solution if the zirconium is complexed by fluoride and the pH is sufficiently high to separate the zirconium wave from the hydrogen wave without precipitation of hydrous oxides. These conditions are attained at pH 7.0 and at a fluoride concentration of approximately 0.1 molar. Conventional polarographic equipment was used: Leeds and Northrup Electrochemograph Type E, capillaries of 4.3-sec drop time at the half-wave potential, with an M value of 1.31 mg sec^{-1} at a mercury pressure of 45 cm. The cells used were small H-type cells with calomel reference electrode, agar backed by a fritted disc, and a sample capacity of 2 to 3 ml. Clearly defined curves were obtained which showed a half-wave potential of -1.44 v (vs S.C.E.), not corrected for I.R. drop. Curve heights could be measured easily down to $5 \mu\text{g}$ of zirconium per 10 ml solution ($\sim 5 \times 10^{-6}$ moles/liter). A straight-line calibration plot was obtained between 5 and $300 \mu\text{g}$ of zirconium per 10 ml.

An Investigation of the Separation and Determination of Small Amounts of Zirconium in Bismuth

The solvent extraction procedure of Scudder and Ballou (Anal. Chem. 25, 1602 (1953)) for separation of zirconium involves the re-extraction of zirconium from the organic phase with dilute HF, which interferes with the subsequent spectrophotometric determination with "Quercetin" as "Thoron" reagent. Addition of perchloric acid and evaporation to remove HF yielded satisfactory results with "Thoron" reagent only in the absence of bismuth.

It has been demonstrated that anion exchange resins when prepared as the chloride complex will effectively remove bismuth. An attempt was made to separate zirconium from bismuth on a column of Amberlite IRA-400, which had been treated with HCl. However, subsequent determination of zirconium with "Thoron" yielded erratic results.

The polarographic method of Dr. Arno H.A. Heyn was applied to the determination of zirconium after separation by each of the two methods. However, in both cases the presence of a wave at a potential of approximately -1.0 v (vs S.C.E.) caused interference. (H.L. Finston, G. Samos, M. Kinsley, S. Bellanca, T. Prach, A.H.A. Heyn,* M. Paikoff,** R. Chasman†)

*Summer visitor from Boston University.

**Summer student from Purdue University.

†Summer student from Harvard University.

Hot Laboratory Operations

Radioisotope Production

Iodine-132 - Production of iodine-132 was transferred to the Operations group this quarter and is now entering the routine phase. Some minor modifications continue to be made: The iodine "generator" was redesigned to reduce holdup of spray by replacing the monel Yorkmesh spray tray with glass beads. The plastic casting around the generator assembly was increased in height so that all glass parts, including tubing, are completely encased in order to prevent breakage due to excessively rough handling during shipping. Wooden shipping crates, previously thought unnecessary for the intermediate size shipment (100 mc), were designed and fabricated. Several minor repairs to the tellurium production equipment were required, several shipping assemblies were constructed, and a shielded product-dispensing apparatus for loading shipping assemblies was built.

During the past quarter, eleven shipments were made as follows:

<u>Customer</u>	<u>Quantity*</u>
Sloan-Kettering Institute for Cancer Research, New York City	three 1000-mc shipments
Oak Ridge Institute of Nuclear Studies	four 100-mc shipments
Mayo Clinic, Rochester, Minnesota	one 10-mc shipment
University of California at Los Angeles	one 10-mc shipment
Notre Dame University	two 1-mc shipments

An exhibit describing the production of iodine-132 was sent to the annual Berlin Industrial Fair in Germany. A paper describing this production appeared in the August issue of Nucleonics by Stang, Tucker, Banks, Doering, and Mills. (W. Campbell, P. Richards, W. Tucker,** H. Banks,** R. Doering,** T. Mills**)

Yttrium-90 Micro Beads - A process based on modifications of a method by Ginell and Simon (Nucleonics 11, 3, 49 (1953)) for making small beads containing Y^{90} was set up, and 10 mg of beads in the 3- to 8- μ diameter range were supplied to a medical research group at Montefiore Hospital. The composition of these beads and the method of irradiation are the same as for the larger beads referred to above under "Yttrium-90," and they were shipped as a suspension in 12-ml of sterile saline solution. The total activity of the 10-mg shipment was approximately 2.3 mc. Leaching tests done on similar preparations indicate that the yttrium is firmly bound to the clay, and that less than 0.02 percent of the activity is being leached off. (P. Richards, J. Baranosky, P. Schneider†)

Radiography

The development of a rapid radiographic method mentioned in the last quarterly progress report (BNL 299 (S-22)) was continued and polaroid self-developing film was found to be practical when rapid results without fine detail are required. Total time for exposure and developing to a finished picture by this method is about three minutes. This technique was used to determine liquid levels in closed metal loops.

This radiographic service has been extended to include the location of internal reinforcing bars and conduits embedded in floors and walls prior to drilling or cutting

*Millicurie figures are nominal ones.

**Temporary assistance during transfer of work to Operations group.

†Summer student now at Harvard University.

holes. This new application was used when the machinery in the Hot Laboratory staff shop was moved to a new location, and it was possible to locate the equipment in such a way that metal already embedded in the floor did not interfere with bolt-down.

A source has been mounted permanently in one of the experimental "loops" of another group in this Department in order to speed up and simplify repeated radiography of a section of this loop by standardizing source-to-film distance and eliminating repeated transportation of a heavy shield.

Sources now in use include Co^{60} , Ir^{192} , and Tm^{170} in varying sizes and activities, and an 8-curie Cs^{137} source is on order.

The amount of service requested of radiography has already increased beyond that which it is possible to supply with the present manpower, and indications are that it will continue to increase. The following figures show the rate of expansion:

<u>Quarter (1954)</u>	<u>Total Number of Service Radiographs</u>
January - March	218
April - June	290
July - September	377

(P. Richards, J.C. Austin)

Beta Sources

Work has continued on the improvement and calibration of beta-emitting sources to be used in medical research. At the present time, three such thallium-204 sources have been made and are now in use:

1. A 3-mm diameter source in an 8-mm diameter holder, delivering at contact 12,000 rep/hr, for Dr. Stoner of the Medical Department.
2. A 2-mm diameter source in a 4-mm diameter holder, delivering at contact 30,000 rep/hr, for Dr. Edelmann of the Biology Department.
3. A 3-prong device used for irradiating 5-ml samples in a test tube, delivering approximately 200 rep/hr, for Dr. Drew of the Medical Department.

More recently, a process has been devised for producing extremely high beta radiation levels of the order of 300,000 rep/hr and higher. Tracer preparations indicate that such sources can be made as small as 2 to 3 mm diameter with fission product strontium-90. Such sources are much more highly concentrated than are those available commercially. (P. Richards, J. Baranosky)

Ultimate Waste Disposal

(4583)*

The removal of mixed fission products by ion exchange on montmorillonite clay in the presence of near saturation concentrations of boric acid has been demonstrated in a series of tests. The decontamination factors (about 150 with a single clay column 18 in. long) were found to be due largely to the inability of the clay to remove ruthenium. The addition of a weak-base anion exchange resin column in series with the clay gave an over-all decontamination factor of roughly 10^4 . The decontamination factors showed little or no change during the course of any run in which a volume of liquid several hundred times the volume of the clay passed through the column. /

In conjunction with the plan to remove nitric acid from wastes by the use of semi-permeable membranes prior to adsorption of fission products on clay, there is, of course, the potential advantage in removal of large first cuts of the acid by distillation. A number of tests on low pressure distillation at 60°C has again shown the feasibility

*For explanation of these numbers, see Foreword.

of this method of acid removal without any measurable carry-over of volatile ruthenium. It is now planned that removal of acid by distillation as well as by membranes will be used in pilot plant operation.

Further investigation has been made of the use of semipermeable membranes for removal of large concentrations of aluminum from waste solutions; by this method, much of the occluded fission product content can be recovered for adsorption on clay.

Some attempts have been made to remove ruthenium from waste solutions in a separate step following adsorption of the other fission products on the clay. More than 90 percent removal of ruthenium has been achieved, but the results are far from satisfactory. Further work on ruthenium removal and its ultimate disposition is planned. (L.P. Hatch, G.P. Simon, W.H. Regan)

The study of the strontium-cesium equilibria on montmorillonite has been completed. In this work the equilibrium column method has been extensively used. The system for $c_0 = 0.05$ N has been studied from 5° to 75°C . At the higher temperatures the selectivity of the clay for Cs markedly decreases.

The exchange equilibria of the alkalis on attapulgite is still under investigation. This work is nearing completion.

A fundamental study of self-diffusion in a resinous ion-exchanger is being made. This work shows that the free anion content of the cation exchanger significantly influences the diffusion constant of the cation.

The column method for the determination of exchange capacities of clays is being tested on a wide variety of clays. (Yale group: H. Thomas and associates)

Fission Products Utilization

(4582)

Polymerization Studies

N-Vinylpyrrolidone - The effect of the variables -- gamma-ray intensity, temperature, solvent composition, and monomer concentration -- on rate and molecular weight has been studied. The rate was found to be virtually independent of monomer concentration, to increase with temperature and intensity, and to decrease with addition of alcohols or chloroform to the water solvent. Molecular weight is controlled mainly by changes in solvent composition and monomer concentration: it increases with monomer concentration and decreases with addition of nonaqueous solvents. The molecular weight distribution of the polymer is significantly different from that of the commercial material. Irradiation of the polymer causes a decrease in molecular weight.

A thorough paper study of the possibility of graft polymerization has been made, and an experimental program has been initiated.

Solid state polymerization studies were continued, and a detailed study of the acrylamide system is fairly complete. An x-ray analysis of the unit cell structure indicates that the vinyl groups are in close proximity to each other so that head to tail addition should be simple. The rate was found to depend upon the first power of the intensity over a range of 3000 to 900,000 r/hr. Over the same intensity range, molecular weight was found to be relatively constant.

Work has been initiated on the use of the ionization produced by the fission fragments to promote chemical reactions. An in-pile experiment is planned to test the feasibility of converting methanol to ethylene glycol. (B. Manowitz, D.S. Ballantine, A. Glines and P. Colombo)

Radiation Application Engineering

Seven cobalt castings measuring 1/16 in. in wall thickness were received. These were encased in stainless steel and are ready to be placed in the reactor. Seventy-two

of the 200 cobalt strips ordered have been received and will be prepared for pile irradiation. The following sources were discharged from the reactor: Cobalt #23, #33 and nested source Cobalt #21.

Cobalt #23, a standard type source, was sold, together with a container, to Dow Chemical Co. Cobalt #33 was shipped to California Oil Development Co. as a replacement for the source they were using. This is to compensate for the deficit in curies in the source originally shipped to them.

Nested source Cobalt #21 is being used under water in the corrosion test setup.

Inquiries, letters of intent, and purchase orders have been received from five companies as follows:

General Electric	1 kilocurie standard source
Diamond Alkali	1 kilocurie standard source
Reynolds Tobacco	4 flat sources
Minnesota Mining	1 kilocurie standard source
Rock Island Arsenal	500 curie standard source

During the last review period two standard type cobalt sources were put into the reactor.

Use of gamma irradiation facilities during this quarter was as follows:

	<u>Source Days</u>
Fission Products Utilization Project	197
Other BNL departments	111
Outside organizations	104

Four underwater radiation chambers equipped with furnaces capable of elevating the temperature to 500°C have been built and installed in the water pit. Instrumentation which permits the control and recording of the temperature in the chambers has also been completed. This setup was designed primarily for current corrosion studies, but will also prove useful for other irradiations. (O.A. Kuhl, R. Oriez, W. Shalvoy)

Heat Transfer In Spray Columns (4550)

Heat transfer rates between hot mercury drops and water are being measured in a counter-current spray column. Data have been taken on 1-in. and 2-in. glass columns in which the drop size varied from 0.020 to 0.25 in., the water flow from 18 to 110 gal/(min)(ft²), the mercury flow rate from 400 to 6000 lb/(min)(ft²), and the volumetric holdup of mercury from 0.4 to 15 percent.

Figure 1 presents some typical results for a water rate of 37 gal/(min)(ft²). The plot shows mainly the effects of mercury rate and drop size on the volumetric over-all heat transfer coefficient, U_a , which is defined as follows:

$$U_a = \frac{q}{V (\Delta t)_{lm}} ,$$

where q = heat transfer rate, Btu/(hr)
 V = column volume, ft³

$(\Delta t)_{lm}$ = logarithmic mean temperature differences between bulk temperatures of mercury and water at both ends of the column, °F.

Figure 1 also presents a few results comparing columns of different lengths and diameters. The falling mercury drops produced considerable vertical mixing in the water phase, depending upon the relative flow rates and column diameter. Under the worst conditions, the temperature rise of the water on entering the base of the column amounted to about 75 percent of the total temperature rise across the column.

Figure 1. Volumetric heat transfer coefficients.
Water rate: 37 gal/(min)(ft²).

As shown in Figure 1, the heat transfer coefficient varies directly as the first power of the mercury flow rate. Since the mercury holdup is roughly proportional to its flow rate, it appears that the increase in the coefficient is mainly due to increase in holdup. The effect of water rate was found to be very slight -- what might be expected from the change in holdup resulting from the variation in water rate. The relationship

$$H = \frac{(V_m)}{v_m A}$$

exists where

H = fractional holdup of mercury on volume basis

A = cross-sectional area of column, ft²

v_m = average linear velocity of mercury, ft/sec

V_m = volumetric flow rate of mercury, ft³/sec

It will be noticed from the plot that the 19.4-in. column gave significantly lower coefficients than the 13.2-in. column, which may be due, in part at least,

to an end effect, although the equipment was designed to minimize turbulence at the ends of the column. The effect of drop size is not great, because as drop size is increased there are two counterbalancing effects: the surface area is reduced, but the drops fall faster, and decrease the water film resistance, which is believed to be the controlling resistance. The first of these two effects predominates.

As far as the heat transfer capacity of the column is concerned, the degree of vertical mixing of the continuous phase which occurs is a big factor. This mixing can be reduced by the use of packing. The extent of mixing is found to go through a maximum as the mercury rate is varied. Under certain conditions, mercury rates around 3 tons/(min)(ft²) appear to produce the same degree of mixing as flow rates around 1 ton/(min)(ft²). (R.D. Pierce and O.E. Dwyer)

INSTRUMENTATION AND HEALTH PHYSICS DEPARTMENT

Instrumentation Division*

(†)**

The Instrumentation Division had six summer visitors: three graduates and three undergraduates. The undergraduate visitors worked on the following: fast scalar circuit, stable wide range oscillator, and CRT pulse height analyzer.

The audiofrequency noise in tubes and transistors was studied; of special interest was the flicker noise in the low audio range. It was found that 6AK5, 6AG5, and 6CB6 tubes have similar characteristics. Some transistors are less noisy in the 20 to 100 cycle region. The most interesting discovery was that the Z729 (British Industries, Ltd.) was much quieter than other tubes and transistors in the low frequency region.† Techniques for β -ray energy measurements were developed in which scintillation counters were used with sources of energy 1.7, .96, and .167 Mev. The measured energies fall within ± 25 kev of the accepted value which shows that this scintillation method may be used for measurements of short-lived or low activity sources that are difficult to measure on a lens spectrometer.†

Current Indicator and Integrator -- EHI-641.

A new current indicator and integrator (EHI-641) with input current scales of 10^{-4} to 10^{-10} amp was built to replace EHI-306 (described in Rev. Sci. Instr. 22, 688 (1951)) for use with the Chemistry Department's Van de Graaff generator. Two feedback loops are used. The input current goes through a stabilized resistor (switched for different scales); the input end of the resistor is connected to the grid of the first stage of the first feedback amplifier, and the other end is connected to the output of the loop. The output voltage of this amplifier is proportional to the input current and is integrated in a second feedback loop of similar design. When the integrating capacitor is charged to a predetermined voltage, it is discharged by a Stevens-Arnold msec relay and a count is recorded. The two-loop arrangement avoids the necessity of separate charge time adjustments for high input currents. The uncertainty in the integrator zero setting is less than 0.01 v, which corresponds to an integration error of less than 1 cycle in 1000 sec. A full-scale (10-v) signal from the first loop results in 1 integration cycle per sec. A second integrating condenser, .01 times the normally used condenser, provides a means of zero setting the integrator for least drift. An integrating interpolation meter is provided for reading fractions of full counts. (S. Rankowitz)

Scintillation Pair Spectrometer

A pair spectrometer utilizing a gray wedge pulse-height analyzer was completed. The instrument includes a triple 0.1- μ sec coincidence circuit and two single-channel pulse-height analyzers for selecting annihilation radiation in the side crystals. The short time resolution permits the use of intense sources so that measurements can be made on gamma rays only slightly above the pair threshold in a reasonable length of time. A very presentable pair spectrum of the Co^{60} gamma rays (1.17 and 1.33 Mev) was obtained in 18 hr. (R.L. Chase)

*Formerly Electronics Division.

**For explanation of this symbol, see Foreword.

†A few copies of reports on these items are available from the Instrumentation Division.

Improved Pulse Amplifier

A new nonoverloading amplifier for scintillation counting has been designed. It features an improved output stage (based on an Oak Ridge modification of the A-1 amplifier) which is linear for more than 95 v of the total 100-v output. The low level stages are long-tailed-pair feedback loops using 12BZ7's. The gain is 1500, and the rise time is 0.1 μ sec. (R.L. Chase)

Health Physics Division

Dosimetry for Rat Irradiations

(6500)*

A dosimetric study has been made of an experimental arrangement for exposing rats to thermal neutrons in the Biology Department's thermal column at the pile. A lucite irradiation box with square cross section was divided into eight symmetrically arranged compartments serving to keep each rat in a relatively fixed position, and thus to equalize dosage.

Measurements of thermal neutron flux were made with a gold foil in one compartment and with rats of varying weights in the other seven spaces. A plot of the logarithm of flux vs total weight of rats was found to be linear, the flux depression due to 1050 grams of rats amounting to only 10 percent. The neutron flux to which a rat is exposed will be taken to be that measured as just described and corrected for the additional depression caused by the eighth rat.

A similar procedure was established for gamma exposure, in which a Bureau of Standards type film dosimeter was used. The sensitivity of this dosimeter to thermal neutrons was known from previous work, so that the thermal neutron part of the response could be subtracted to obtain the gamma dose rate in the empty eighth compartment. This rate was found to be independent of the mass of rats in the other compartments, within the precision of the measurements. Thus, no correction is necessary for the effect of the eighth rat on gamma dosage rate. Presumably, the increased gamma radiation to be expected as a result of the increased weight of rats is approximately cancelled by the flux depression. For a thermal neutron flux of 9.5×10^8 n/cm²/sec, the total gamma dose rate amounts to 220 r/hr, of which 95 r/hr originates in the other rats and 125 r/hr is characteristic of the empty facility.

A gold foil located above the irradiation box will be used for standardization of individual experiments, since there are bound to be changes in the conditions of pile operation and in the weights of the rats being irradiated. The ratio of activity of this foil to that measured in one empty space was found to be 0.606, independent of the weight of rats in the other compartments. (L. Lewis)

The measurement of gamma radiation in the empty facility by the lithium extrapolation method, reported last quarter, is nearly complete. A lead absorption curve was taken by means of cylindrical lead shells around a thimble chamber with enough shells of lithium outside the lead to absorb all thermal neutrons. The effective energy of the gammas was shown to be 3 to 4 Mev, which indicates that they come primarily from the graphite and that no appreciable soft component is present.

The initial extrapolation was made with a solid inner shield of lithium and shells with 1/16-in. spacing. Some doubt was cast on the validity of the extrapolation because of this spacing, so it was repeated with negligible spacing. Very little difference was noted in the gamma-ray level obtained. (F.B. Oleson)

*For explanation of these numbers, see Foreword.

High Level Calibration Facility

(†)

A facility for calibration of high range instruments has been set up in part of the former officers' club. A Co^{60} source giving 29.9 r/hr at a meter may be raised from a storage shield into a lead housing with an opening directed along a lightweight track with arrangements for distance measurements. The facility was badly needed for calibration of the upper ranges of various Health Physics survey instruments and will be useful for instrument development projects and for the exposure of high range calibration films for personnel monitoring. (J. Handloser)

A 1-curie Ra-Be neutron source in a 7-in. thick lead shield has been moved to the same area and is being used for fast neutron calibration of NTA films and survey meters. The neutron yield through the shield, measured with a Hanson counter, amounts to $1/14 \times 10^7$ n/sec with a gamma-ray dose rate of less than 1 mr/hr at 1 meter from the source center. The neutron spectrum is considerably degraded in energy compared with the Po-Be sources used in the past, but is still acceptable for calibration purposes in view of the expense involved in maintaining a large Po-Be source. The Po-Be equivalent of this source, as far as NTA tracks are concerned, has been determined experimentally so that the exposure standard, based on the known spectrum of Po-Be, can be retained. (F.B. Oleson)

Waste Disposal

(†)

A test has been made of the feasibility of using active concentrate from the liquid waste evaporation plant in mixing concrete for sea disposal. The resulting barrels of concrete gave relatively low external radiation levels even when full strength concentrate was used in the mix. A number of improvements in the mechanical arrangements will be necessary before this method of packaging the concentrate can be used routinely. Data on liquid wastes are presented in Table 1.

It was found that much more active objects could be packaged for sea burial if heavy concrete was used. A 14-fold improvement over ordinary concrete was achieved with a 10-in. thick liner of limonite concrete loaded with steel punchings. Up to 23 curies of Co^{60} can now be packaged in a 50-gal drum without exceeding a level of 1 r/hr at the surface. (L. Gemmell, S. Pearsall)

Miscellaneous Projects

(†)

The annual 9-week course in Health Physics activities, given for the Rochester-Brookhaven group of AEC fellows in radiological physics, was completed. Members of the group carried out a number of special projects in addition to their routine training. For instance, one student determined the size distribution of particulates in the stack effluent from the pile. It was found that practically all particles were less than 1 micron in diameter, which confirms the efficiency of the exit air filters in removing larger particles.

A continuous dust monitor has been provided for monitoring the pile stack effluent. Stack air is sampled isokinetically at the base of the stack, deposited on a moving filter paper strip, and counted with a scintillation counter. Only an approximate calibration of the monitor has been made thus far.

An intense beam of 3-Bev protons, recently brought out into the Cosmotron experimental area, has posed new problems of safety. Only preliminary surveys are available as yet, but they indicate that personnel must be rigidly excluded from the beam itself, at least within the building. A detailed study is planned as soon as the operating schedule of the machine permits.

Table 1				
Summary of Liquid Waste Disposal Data				
	July	August	September	3-Mo. Average
Input to filter beds (gal/day)	401,000	420,000	383,000	401,000
Output from filter beds and discharge to stream (gal/day)	292,000	386,000	383,000	354,000
Net loss in filter beds (%)	27	8.1	0	11.7
Stream above discharge point (gal/day)	14,000	13,000	355,000	127,000
Stream at site boundary (gal/day)	284,000	442,000	789,000	505,000
Average activity concentration at input to filter beds (c/cc)	3.7×10^{-13}	5.0×10^{-13}	3.9×10^{-13}	4.2×10^{-13}
Average activity concentration at output from filter beds (c/cc)	1.3×10^{-13}	1.4×10^{-13}	0.7×10^{-13}	1.1×10^{-13}
Reduction in activity concentration effected by filter beds (%)	65	72	82	73
Activity in sewer effluent (mc)	17	22.5	16.8	18.8
Percent of permissible	12.9	16	14	14.3

Publications

"Some remarks on data handling systems," R.L. Chase. Abstract of talk given at meeting of IRE Professional Group on Nuclear Science, October 1954.

"The effect of feedback on noise in a pulse amplifier," R.L. Chase. Submitted to Rev. Sci. Instr.

"Determination of total carbon and its radioactivity," Sinex, Plazin, Clareus, Bernstein, Van Slyke, and Chase. Submitted to J. Biol. Chem.

Preliminary Results

When Drosophila are fed on a medium containing the agent EDTA which chelates calcium and magnesium, the frequency of chromosome rearrangements is significantly increased. This is the first instance of an alteration of this type being produced by a chemical agent. (D.M. Steffensen)

It was previously noted here that Tradescantia microspores from plants grown in media deficient in calcium show a higher rate of chromosome abnormalities than those grown on a normal medium. It has now been found that when these microspores were subjected to x-irradiation, the ones from calcium-deficient plants exhibited a higher rate of increase of chromosome abnormalities than normal microspores. This definitely suggests that the x-ray breakage of chromosomes is not an "all-or-none" affair but depends upon the condition of the chromosomal material at the time of irradiation. (D.M. Steffensen)

A further example of the radiomimetic action of nitrogen mustard has been found. It had previously been reported that thyroidectomized mice given a large but non-lethal dose of radiation will develop pituitary tumors. It has now been found that a dose of nitrogen mustard will produce the same type of tumor in thyroidectomized mice. (A. Edelmänn)

The work on the characterization and identification of the toxin produced in rats by x-irradiation has continued and been reconfirmed. In addition it has been shown that this toxin will increase the liver glycogen when injected into fasted, intact rats. This response is similar to that of these animals to x-irradiation, and is the first indication that this toxin resembles the radiation syndrome in intact animals. (A. Edelmänn)

Submitted for Publication

IMPROVED STORAGE QUALITY OF POTATO TUBERS FOLLOWING EXPOSURE TO GAMMA RADIATION FROM COBALT-60. A.H. Sparrow and E. Christensen. *Nucleonics* 12, 8 (1954). Abstract.

Potato tubers of the Katahdin variety were exposed to gamma-ray dosages of approximately 1,250, 5,000, 10,000, 20,000, 80,000, and 106,250 r. The effect of these dosages on taste, sprouting, shrinkage, internal black spotting and texture was followed for 18 months. The taste panel reported a preference for the irradiated samples over the controls after eight months but no undesirable taste in any of the irradiated samples even after 18 months of storage. Sprouting was reduced at the two lower dosages and completely inhibited at doses of 20,000 r or higher. Weight loss during storage was greatly reduced in all but the lowest dosage of the irradiated tubers. After prolonged storage, the lowest dosage sample and the controls showed severe softening, and the highest doses produced some softening. The potatoes irradiated with 20,000 r were still firm and excellent in appearance after 18 months of storage. It is concluded that gamma irradiation, particularly in the dosage range 5,000 to 20,000 r, can greatly prolong the storage life of potatoes.

THE EFFECTS OF ADRENAL HORMONES ON THE CAPILLARY CIRCULATION AND ON CAPILLARY PERMEABILITY IN PERFUSED RAT TISSUES. E.M. Renkin and B.D. Zaun.** Submitted to Am. J. Physiol. Abstract.

*For explanation of these numbers, see Foreword.

Capillary permeability to fluid, protein and sucrose, and precapillary and post-capillary hydrodynamic resistance were measured in the perfused hindquarters of normal and adrenalectomized rats. No permeability differences between the two groups were observed, nor was permeability affected by adding adrenal extract to the perfusion fluid. Aqueous adrenal cortical extract was found to contain a vasoconstrictor agent readily destroyed by oxidation in alkaline solutions and therefore presumed to be epinephrine. The reported effects of such extracts on capillary permeability are attributed to their vasoconstrictor action.

**Mount Holyoke College, South Hadley, Mass.

✓
STIMULATION BY CHRONIC GAMMA IRRADIATION OF THE SPONTANEOUS RATES OF HERITABLE SOMATIC INSTABILITIES IN A CLONE OF "NICOTIANA." S.A. Sand,* H.H. Smith, and A.H. Sparrow. Submitted to Genetics. Abstract.

A *Nicotiana* clone, S5230-5, selected from cultures of interspecific hybrid origin is employed. Flowers of this clone show a high, spontaneous, somatic "mutation" rate (10^{-3}) for one heterozygous, genetic locus, and a much lower rate (10^{-5}) for another heterozygous locus. The phenotypic consequences of these instabilities are sectors of speckled and purple tissue, respectively, on the normally red petals. These sectors are counted per cm^2 petal area of estimated epidermal cell-number.

Clonal replicates were grown at Brookhaven throughout the summer (1953) at ten levels of chronic gamma irradiation (Co^{60}) varying from control to 300 r per day. Analysis of variance and graphical presentation are applied to data from samples of flowers scored during July and again during August.

The purple-speckled-sector frequency is nonlinear, and gives radiation-induced increments ten times those of the purple-sector frequency in the range below 12 r per day. Significant differences exist in speckled but not in purple sector frequency between clonal plants subjected to the same irradiation level. Mean frequencies obtained in August are about two-thirds those for July from the same plants.

In somatic tissue the speckled locus or a related physiological system is 1) hypersensitive to differences in gamma irradiation level at low dosage, and 2) also responsive to differences in other, potentially separable, external, environmental factors and/or to internal environment tentatively assumed to be correlated with metabolism and development.

*Cornell University.

MECHANISM OF ACTION OF CARBOHYDRASES. D.E. Koshland, Jr. Submitted to J. Am. Chem. Soc. Abstract.

The reactions catalyzed by the carbohydrases can be interpreted on the basis of displacement mechanisms in which a single bond is formed with the attacking reagent at the same time that the single bond in one of the substrates is broken. The attacking reagent may be the second substrate or an electron-sharing site on the enzyme surface. With a few simple mechanisms it is possible a) to explain the observed stereochemistry of carbohydrase action in terms that are consistent with physical organic chemistry, b) to correlate the cleavage point of some enzymes on the basis of their specificity patterns, and c) to explain the hydrolytic and transferase activities of many of the carbohydrases. The mechanisms also suggest the kind of groups which may be active on the enzyme surface and provide an explanation for the observed pH dependence of the enzyme activity for a number of these enzymes.

✓
CYTOGENETIC ANALYSIS OF A SPONTANEOUS PRE-MEIOtic MUTATION IN MAIZE. E.J. Dollinger* and W.R. Singleton. Submitted to J. Heredity. Abstract.

Cytogenetic tests of a spontaneous premeiotic R r mutation reveal that it is an

autonomous mutation at the R locus. Genetic tests indicate that no factor other than the R locus itself is involved in mutation and that the mutation is stable. No evidence was found for chromosomal alteration in meiosis, pollen or transmission sterility.

*National Cancer Institute Postdoctoral Fellow.

THE SENSITIVITY OF DEVELOPING "HABROBRACON" TO OXYGEN. A.M. Clark* and E.B. Herr, Jr.** Submitted to Biol. Bull. Abstract.

1. Groups of Habrobracon were exposed during postembryonic development to air, nitrogen, oxygen and carbon dioxide and studied with regard to development and oxygen uptake. Deleterious effects from oxygen and from carbon dioxide were observed after exposure at the white pupal stage, but not after exposure at the larva-in-cocoon stage. No deleterious effects from exposure to nitrogen were found.

2. Exposure of white pupae to 1 atmos of oxygen or carbon dioxide caused cessation of development at the pigmented pupal stage in a high proportion of the individuals. The incidence of emergence from cocoons as adults was higher for males than for females. Adults with wing and antennal abnormalities occurred. Exposure to 2 atmos of oxygen or carbon dioxide resulted in greater deleterious effects on development than did exposure to 1 atmos.

3. The oxygen uptake of unpigmented pupae exposed to 2 atmos of oxygen is markedly reduced. No such reduction was observed following treatment with 2 atmos of nitrogen or carbon dioxide or 1 atmos of oxygen.

*The Marine Biological Laboratory, Woods Hole, Mass , and the University of Delaware, Newark, Del

**Cornell University, Ithaca, N Y

AN IMPROVED SEALING WAX FOR TEMPORARY SMEAR PREPARATIONS. E. Christensen. Submitted to Stain Technol. Abstract.

A mixture of polyethylene glycol 4000* and gum mastic as an improvement over the classic paraffin-mastic seal for temporary slides offers the following advantages: no stratification of components; delayed solidification permitting effective application of weights for flattening cells, solubility in 3:1 ethanol-acetic acid, to eliminate scraping for removal and so reduce the number of broken cover slips and paraffin artifacts. Hardness may be varied by using other grades of polyglycol. Gum mastic is purified by treatment with acetone and hot water, after which it is mixed with molten polyglycol. The author calls the product "Glycomastic."

*"Carbowax" 4000, also available in grades 6000, 1540, 1500, 1000 from Carbide and Carbon Chemicals Co., a division of Union Carbide and Carbon Co , 30 E. 42 St., New York, 17, N Y. "Polyglycol" E9000, E6000, E4000, E2000, E1000 from Dow Chemical Co , Midland, Mich. Dow products were not tested in preparing Glycomastic, but manufacturers' specifications for the various grades indicate they are much the same as the Carbowaxes.

THE EFFECT OF CERTAIN GASES ON THE RADIOSENSITIVITY OF "HABROBRACON" DURING DEVELOPMENT. A.M. Clark* and E.B. Herr, Jr.** Submitted to Radiation Research. Abstract.

Groups of Habrobracon during postembryonic development were exposed to x-rays in the presence of oxygen, carbon dioxide, nitrogen and hydrogen. The effects on development to adulthood were observed. Pupae exposed to 100 percent oxygen (without x-radiation) were arrested in their development. Diploid female pupae were more resistant to x-rays and more sensitive to oxygen than were comparable haploid male pupae. Groups of Habrobracon irradiated during the larva-in-cocoon stage in the presence of nitrogen or carbon dioxide were about three times as radioresistant as comparable groups x-rayed in air or oxygen. The amount of protection from x-radiation

by nitrogen was the same for oxygen-sensitive and oxygen-resistant stages. There does not appear to be a clear relationship between oxygen poisoning and x-radiation damage. Groups of Habrobracon during the larva-in-cocoon stage were more radio-resistant when x-rayed in the presence of hydrogen than in the presence of nitrogen or carbon dioxide than when x-rayed in hydrogen. These experiments show that the radiosensitivity of Habrobracon can be modified by the presence of hydrogen.

*The Marine Biological Laboratory, Woods Hole, Mass., and the University of Delaware, Newark, Del.
 **Cornell University, Ithaca, N.Y.

TREATMENT BY BLACK-WIDOW SPIDERLINGS OF Sr^{90} - Y^{90} IN THEIR PREY. V.T. Bowen* and A.C. Rubinson. BNL 1965. Submitted to Science. Abstract.

Young black-widow spiders fed on Drosophila labelled with radioactive strontium-90 and yttrium-90 provided evidence that spiders have developed methods of control of their heavy metal content which act at the level of absorption. No evidence was obtained of their discrimination between the two elements, either in uptake or excretion.

*Present address: Woods Hole Oceanographic Institution, Woods Hole, Mass. Mass.

ISOTOPIC EXCHANGE CRITERIA FOR ENZYME MECHANISMS. D.E. Koshland, Jr. BNL 1972. To be given orally at the August 1955 meeting of the Faraday Society.

A wide variety of enzymatic reactions obey the stoichiometry $\text{BX} + \text{Y} \rightarrow \text{BY} + \text{X}$ and have the characteristics of organic substitution reactions. It is postulated that a few simple mechanisms can explain the fundamental acts of bond formation and bond cleavage for this diverse group and that these mechanisms involve displacements in which a single bond is broken as a single bond is formed. A combination of stereochemical, specificity and isotopic exchange evidence can be applied to distinguish between the alternative mechanisms in individual cases and to characterize the role of the enzyme in the reaction. Of these the isotopic exchange criterion is the most universally applicable. Isotopic exchange experiments of the types

can give important qualitative and quantitative information on the presence or absence of enzyme-B intermediates, the stability of enzyme-B intermediates, and the necessity or lack of necessity for the simultaneous presence of both substrates for enzymatic action. The information obtained from these exchange experiments and their limitations will be discussed. The criteria will be applied to a number of specific enzymes, such as alkaline phosphatase, acetyl cholinesterase, beta glucosidase, and glutamine synthetase.

THE EFFECT OF A YEAST UPON THE INCORPORATION OF PHOSPHORUS INTO THE TISSUE OF ADULT FRUIT FLIES. R.C. King. BNL 1981. Submitted to Evolution. Abstract.

Adult Drosophila melanogaster feeding upon a synthetic medium incorporate large amounts of exogenous phosphorus into tissue, if the medium contains live yeast (Saccharomyces cerevisiae). Insects on nonyeasted medium must ingest ten times as much food as those on yeasted medium to account for their observed exogenous phosphorus contents. It is suggested that the association of many insects with yeasts may have originated to insure efficient utilization of the phosphorus of the environment.

QUANTITATIVE STUDIES ON THE PHOSPHORUS BALANCE OF ADULT FEMALE FRUIT FLIES FEEDING UPON YEAST. R.C. King and L.P. Wilson.* BNL 1985. Submitted to Anat. Record. Abstract

A study was made of the movement of the phosphorus of yeast (*Saccharomyces cerevisiae*) ingested by adult female *Drosophila melanogaster*. It was found that 70-80 percent of the phosphorus in yeast is extracted during its passage along the gut. An average of 90 percent of the extractable phosphorus of the meal has passed through the gut to other tissues by the time the digestive tract has emptied. Incorporation of medium P into eggs is very rapid. Eggs laid by week-old females contain over thirty times as much exogenous phosphorus as endogenous phosphorus. Only a small fraction of the endogenous phosphorus of the egg can represent phosphorus originating as DNAP contributions from the nurse cells, trophocyte and primary oocyte. It is demonstrated that a *Drosophila melanogaster* female may ingest daily an amount of yeast equal to two times its body weight. During the day an amount of phosphorus equal to the amount of phosphorus in the ingested yeast is lost. Twenty percent of the phosphorus lost is nonabsorbed phosphorus of the meal. Fourteen percent of the phosphorus lost is in eggs. Sixty-six percent of the phosphorus lost is in forms other than eggs (probably in secretions of the salivary glands, digestive tract and malpighian tubes). The relation between endogenous phosphorus remaining and time follows a two-phase system where the slow phase has a half-time 12-13 times longer than the half-time of the fast phase. Male and female *Drosophila* differ markedly in the pattern of phosphorus turnover.

*Wellesley College.

VERTEBRATE GONADOTROPHINS.* E. Witschi.** BNL 1983. Submitted to the J. Endocrinol., Suppl. 1 (British). Abstract.

In the bioassay of gonadotrophic preparations, it is important to find reactions that are specific for one hormone only and which are not inhibited or augmented by other hormones that may simultaneously be present. The value of the weaver-finch reaction for the identification and quantitative assay of the luteinizing hormone (LH) is demonstrated in the results of assays of a variety of preparations. It is concluded that the male hypophysis of man and of rats never liberates LH hormone, and that its release in the female normally is restricted to that stage of the sex cycle at which the ovarian follicles reach maximal size. The LH hormone induces ovulation and corpus luteum formation.

In a second series of experiments evidence is presented in favor of the separate nature, existence, and function of an interstitial cell stimulating hormone (ICSH). While FSH alone is stimulating the follicular apparatus of the gonads, that is, the sustentacular cells of the testis and the granulosa of the ovary respectively, ICSH controls the growth and androgenic endocrine functions of the theca and interstitial cell complex.

In rats and mice, implantation at birth of a testis destroys the faculty of LH release and of cyclic function of the hypophysis of females. Attention is directed to the important role played by the hypophysis in the control of reproductive cycles of vertebrates.

The comparative study of hypophyseal contents of representatives of all vertebrate classes shows an almost even level of LH per milligram dry weight. Only in the horse and in man the stores are decidedly low. On the other hand, FSH contents are very low in cattle, birds, and the coldblooded species but rise to high levels in some mammals, particularly the horse and man.

The question of taxonomic specificity of hypophyseal hormones is discussed shortly but the few and scattered data now available do not lend themselves to generalization.

It is indicated that improved assay methods for FSH and ICSH may contribute toward the resolution of this controversial subject.

*This work was in part supported by grants from the National Institutes of Health, Public Health Service, and from the National Science Foundation.

**The State University of Iowa, Iowa City, Iowa.

A MECHANISM OF GENE INTERACTION. B.S. Strauss.* Submitted to Am. Naturalist. Abstract.

The properties of multi-enzyme systems are discussed and a mechanism of coupling such systems by means of mutual inhibition is suggested. Coupling by inhibition is a characteristic of varied types of biological phenomena. Genes control the component step reactions of multi-enzyme systems. Gene interaction is the result of the interaction of the multi-enzyme systems controlled by the genes rather than by interaction of the genes themselves.

*Syracuse University, Syracuse, N.Y.

AN INVERSE RELATIONSHIP BETWEEN THE WATER CONTENT OF SEEDS AND THEIR SENSITIVITY TO X-RAYS. R.S. Caldecott*. Submitted to Science. Abstract.

When dormant seeds of Himalaya barley were stored at different relative humidities, their total water was increased between 7 and 20 percent. When seeds with various water content were subjected to x-radiation, their sensitivity in terms of seedling growth at 7 days was inversely related to the water content.

*Present address: Agricultural Research Service, Field Crops Research Branch, U.S. Department of Agriculture.

SOME ASPECTS OF THE COMPARATIVE BIOCHEMISTRY OF IODINE UTILIZATION AND THE EVOLUTION OF THYROIDAL FUNCTION. A. Gorbman.* Submitted to Physiol. Rev. Abstract.

It seems clear that the iodotyrosines and thyroxine are easily formed in nature and are especially rich in hard cuticular iodoproteins of invertebrates. It follows that the hormone thyroxine preceded the evolution of the thyroid gland. In at least three of the phyla we have studied (annelida, mollusca, arthropoda), there are iodoproteins in or very near the mouth or pharyngeal cavity. If this at first accidentally available thyroxine became involved in the metabolic activities of a prechordate group of animals, it would be conceivably advantageous to such organisms, if evolution permitted, to develop larger and more dependable sources of iodoprotein. A series of probable changes can be imagined that would localize iodine metabolism and thyroxine formation in the thyroid area. If the first physiologically useful thyroxine came from iodoprotein in the anterior digestive tract (structures like pharyngeal teeth) then the hormone would have been freed from its protein bondage by intestinal digestion. This appears to be the case for ammocoetes larvae, in which the source of iodoprotein is no longer a cuticular structure, and in which it is localized in the endostyle. Although the lamprey's endostyle has a protease of its own, it appears more likely that the actual hydrolysis occurs in the digestive tract. At any rate, it is clear that when the endostylar duct closes at metamorphosis, the newly completed thyroid of the lamprey already contains a thyroprotein-splitting enzyme similar to that of mammals. The iodoprotein-filled cells, or the secreted protein of these cells, may pass into the pharynx through the short endostylar duct, and thence to the intestine for digestion. This dependence upon digestive enzymes for liberating thyroid hormone no longer is obligatory when the

iodoprotein-forming cells can also supply a protease to convert the hormone to a diffusible form. When this happens the final step in evolution of the thyroid can occur. The iodoprotein-forming sac, or gland, can lose its duct and become a true endocrine organ, since the hormone is able to diffuse directly into the blood. Evidence that holocrine secretion, as is seen in some ammocoetes larvae, is a primitive mode of secretion, can be adduced from the frequency with which the whole secretory cells are found in the thyroidal colloid of lower vertebrates.

*Barnard College, Columbia University, New York, N.Y.

BACTERIAL VIRUS: RADIATION AND FUNCTION. D.J. Fluke and E.C. Pollard.* Submitted to the N.Y. Acad. Sci. Abstract.

T-1 bacteriophage has been bombarded with monoenergetic particles of widely varying ion density and scored for survival of ability to kill the host E. coli cells, for survival of ability to multiply in the host cells, and for damage to the first growth cycle of survivors. The results suggest a threshold in local energy loss both for killing and multiplying ability, and the sub-threshold events which predominate for low ion density particles lead to considerable damage to the first growth cycle of survivors of such bombardment.

*Yale University, New Haven, Conn.

THE RELATION OF GENOME NUMBER TO RADIOSENSITIVITY IN "HABROBRACON." A.M. Clark.* Submitted to the Proc. Natl. Acad. Sci. Abstract.

Haploids and diploids of Habrobracon have been compared with regard to their sensitivity to damage by x-rays. No constant difference in radiosensitivity was found. The relative radiosensitivities of haploids and diploids were different for different stages of development. These studies indicate that no simple relationship between genome number and radiosensitivity exists. The bearing of these data on determining the site of injury from x-radiation is discussed.

*University of Delaware, Newark, Del.

GENERAL MECHANISMS FOR ENZYME ACTION. D.E. Koshland, Jr. BNL 2015.

A wide variety of enzymatic reactions which perform diverse biological functions and present apparently dissimilar chemical reactions, nevertheless conform to the stoichiometry $Bx + y \rightarrow By + x$. Included in this group are the phosphatases, phosphorylases, esterases, lipases, proteases, glycosidases, branching enzymes, etc. It is postulated that this large group of enzymes acts catalytically through a few basic mechanisms. These have been named the single displacement mechanism, the front-side displacement mechanism and the double displacement mechanism. The first two of these require that both substrates be present simultaneously on the enzyme surface, whereas the latter allows a primary electron-sharing attack by the enzyme on one substrate (BX) even in the absence of the second substrate.

The fundamental characteristic of all of these mechanisms is that they proceed by steps in which a single bond is broken as a single bond is formed, in direct analogy to the well-known displacement mechanisms of organic chemistry. With these assumptions it is possible to explain many observed phenomena such as the stereochemistry of the enzymatic reactions, the correlation of bond breaking with specificity, the occurrence or absence of exchange between BX and radioactive X, the simultaneous transferase and hydrolytic properties of some enzymes, and a rough explanation of the pH dependence of enzymatic activity. Specific illustrations have been obtained with

5'nucleotidase, beta glucosidase, acetylcholinesterase, the phosphorylases and glutamine synthetase.

HISTOPATHOLOGICAL CHANGES IN "TRITURUS VIRIDESCENS" AFTER X-IRRADIATION, WITH SURVIVAL PROLONGED BY LOW TEMPERATURE.

S. Jakowska* and R.F. Nigrelli.** To be given orally before the American Society of Zoologists, December, 1954.

Salamanders exposed to single-dose, whole-body x-irradiation of 1700 r, 4700 r, 6000 r, and 9000 r were kept at 5°C for 1, 2 and 3 months. A progressive degeneration occurred in the liver, kidney and spleen of the animals given the lower doses. "Albuminous" material accumulated in the lumen of the tubules and in the glomeruli; later, hyalin droplets were present in the cytoplasm of the cells of the urinary ducts, tubules and of liver. In the first month, the granulocytopoietic layer of the liver appeared reduced in thickness, but it was enlarged and showed considerable mitotic activity in the second and third month. A gradual increase in the number of neutrophils and typical basophilic granulocytes (mast leukocytes) with coarse metachromatic granules was noted in the peripheral blood. Three months after exposure to 1700 r, large numbers of giant basophils (up to 60 x 68 microns) with finer granules were also present. Animals exposed to 6000 r and 9000 r showed a pathological picture similar to that seen at the end of 3 months in animals exposed to lower doses of irradiation. Hyalin droplets were particularly prominent in the parenchyma of the liver and kidney. Leukocytosis, especially an increase in the number of basophils, and pronounced anisocytosis were observed in the peripheral blood. There was no apparent influence on pre-existing or developing pseudoerythroplastic anemia, as seen in some newts. The main pathological effects observed in the present studies appeared to involve cytoplasmic changes rather than nuclear aberrations.

*College of Mt. St. Vincent.

**New York Aquarium, N.Y. Zoological Society.

THE RETENTION OF METABOLIC RADIOCARBONATE. R. Steele. Submitted to Biochem. J.

For a somewhat idealized situation it is possible to calculate, from the retention equation for intravenously administered (C^{14}) bicarbonate carbon, the quantitative characteristics of the system of carbonate pools in the animal for which the retention data were collected. This procedure is demonstrated for the cat, with the data of Kornberg and Kornberg, Davies and Wood, and the degree of correspondence between the "ideal" cat and the real cat is noted. The analysis of the system of carbonate pools furnishes simultaneously a new equation which expresses the retention of (C^{14}) carbon dioxide arising in the soft tissue compartment of the animal, and it is shown that this equation is more suitable than the intravenous (C^{14}) bicarbonate retention equation for a calculation of the rate at which carbon dioxide is produced from an "ideal" metabolite which is burned uniformly throughout the soft tissues. If it is not known whether a particular metabolite is converted to carbon dioxide generally throughout the soft tissues or is converted to carbon dioxide mostly in a single small organ, this information can be obtained by making an additional observation. The observation required is the ratio of the (C^{14}) carbon dioxide expired to the total acid-volatile (C^{14}) carbonate retained in the body at one point in time subsequent to the initiation of (C^{14}) carbon dioxide production from the (C^{14}) metabolite in question. This ratio reveals whether the (C^{14}) carbon dioxide retention is more nearly described by the equation governing intravenous (C^{14}) bicarbonate injection or by the equation governing (C^{14}) carbon dioxide arising throughout the soft tissues.

MEDICAL DEPARTMENT

Division of Pathology

Histological Effects of Various Small Doses of I^{131} on Selected Tissues of the Rat

(6300)*

Rats given single various small doses of I^{131} have been observed throughout their life span in an effort to determine a possible carcinogenic effect. All animals have died or been sacrificed and the microscopic study of the various tissues, particularly the thyroid and pituitary, is under way.

Histological Studies of Effects of Neutron Capture Therapy

(6210)

The histological study has been completed of eight brains from patients receiving neutron capture therapy. Differences are observed in the preirradiated and postirradiated specimens which are suggestive of irradiation effects. The scalp of one patient and that of another case not included in this report show marked irradiation effects. Submission of a report for publication is expected shortly.

Long Term Effects of 400 μ c of I^{131} in Long-Evans and Sprague-Dawley Rats

(6300)

Long-Evans and Sprague-Dawley rats were given a single dose of 400 μ c of I^{131} and allowed to live out their life span to ascertain whether carcinogenic effects in the thyroid appeared.
(J.T. Godwin)

Uptake of Astatine-211 by the Rat Thyroid

(6300)

Further studies of the accumulation of At^{211} by the rat thyroid gland have shown that the potassium thiocyanate decreases the thyroidal accumulation of At^{211} and that a low iodine diet enhances the accumulation of At^{211} . Attempts to determine what oxidation states of astatine are present in the thyroid have not been successful.

Effect of Thiodiphenylamine on the Rat Thyroid

(6300)

It has been shown that impurities in the preparation of thiodiphenylamine are responsible for the lowered thyroidal accumulation of I^{131} found in rats treated with the preparation. Further experiments are in progress to determine just what these impurities might be.
(C.J. Shellabarger)

Rate of Incorporation of S^{35} -Labeled Methionine in the Chick Embryo

(6300)

An analysis was made of the rate of incorporation of sulfur-labeled methionine in several tissues of the chick embryo after administration via the yolk sac at nine days of incubation. Attention was centered on the rate of disappearance from the yolk sac and incorporation into kidney, liver, skin, and spleen. Additional data are available

*For explanation of these numbers, see Foreword

concerning incorporation into blood, brain, chorioallantoic membranes and heart. For purposes of comparison, a similar program of experiments was conducted in which labeled sodium sulfate was used. Analyses were made at 12-hr intervals between 9 and 14 days of incubation and at 17 days of incubation. Additional experiments on the effects of nonradioactive grafts of adult chicken spleen and adult duck spleen on the rate of incorporation in homologous and heterologous tissues of labeled methionine (via the yolk sac) offered further support to the view previously advanced that the rate of incorporation is not altered appreciably by the presence of a nonradioactive graft. A comparison was made of the effects of sulfur-labeled transplants of adult chicken and adult duck spleen. In agreement with earlier findings, after labeled chicken spleen grafts, selective incorporation of radioactivity in the homologous host organ was observed. There is no evidence of selective transfer following labeled duck spleen grafts. (J.D. Ebert* and A.M. Mun*)

Distribution of I^{131} and I^{131} -Labeled Thyroxine and Triiodothyronine in Rabbit Brain

(6300)

An attempt was made to corroborate reports of the selective localization of radioactivity in the hypothalamus of the rabbit after intravenous injection of I^{131} and I^{131} -labeled thyroxine. Thyroxine was administered in a dosage of .6 $\mu\text{g}/\text{kg}$. Areas other than the hypothalamus were studied for comparison. Tissue-plasma activity ratios were determined for sections cut from the pars distalis and neural lobe of the hypophysis, supraoptic and periventricular areas of the hypothalamus, cerebral and cerebellar cortex, medulla oblongata pons, striatal complex, and olfactory tract. Radioautographs were prepared from one-half of the hypothalamus and hypophysis of each animal, the contralateral half serving for activity determinations by counting procedures. Sacrifice was by vascular perfusion with saline followed by Bouin's fluid.

Within an hour after thyroxine injection, a concentration of radioactivity in the neural lobe was apparent. The tissue/plasma ratio varied from 2 to 11, depending on the time after injection. The tissue/plasma ratios of most other neural tissues were usually about .1. The adenohypophysis was from one-fourth to one-seventh as active as the neural lobe. Radioautographs were negative. These data of distribution in the neural lobe and adenohypophysis are in agreement with those previously reported.

No such localization was noted after injection of carrier-free I^{131} , even as long as seven hours after injection of 1 mc/kg. Radioautographs were negative.

The experiment was repeated with I^{131} -labeled triiodothyronine (.6 $\mu\text{g}/\text{kg}$) intravenously administered. Within an hour, a concentration in the neural lobe was apparent. At peak uptake, tissue-plasma ratios of the neural lobe rose as high as 35 and 45, and never fell below 15 at any time up to six hours after injection. Other tissues listed in the first paragraph showed ratios from .5 to 2. The ratio for adenohypophyseal tissue was from 3 to 7, depending on the time after injection. Radioautographs are yet to be examined.

Although it has not been demonstrated that the radioactivity observed was due to iodine remaining incorporated in the thyronine molecule, it is probably not amiss to speculate on the possibility that thyroxine and triiodothyronine are involved in the metabolism and endocrine function of the neurohypophysis (with signal intimacy, for example, in a neurohumoral link between the central nervous system and the adenohypophysis). (A.H. Milofsky**)

*Guest of Medical Department from University of Indiana.

**Student guest of Medical Department from Yale University School of Medicine.

Electrophoretic Separation of Serum Components

(6300)

The following preliminary investigations were initiated:

1. Electro-immunophoresis methods have been evaluated. An electro-immunophoresis apparatus like that described by Dr. Gitlin (personal communication) was constructed. Separation of serum components on filter paper placed between glass plates or suspended in a manner described by Darrum was obtained. Migration of serum components stained with brom-phenol blue through 2 percent and .3 percent phosphate-buffered agar was also observed. A power supply providing 300 v at 6 ma was used. Further investigation of antigens of embryonic origin will be conducted. (A.M. Mun*)

2. A study is being made of the incorporation of sodium chromate (Cr^{51}) into adult chicken tissues and the effects of chromium-labeled grafts in host embryonic tissues. (W.L. Hughes** and A.M. Mun*)

(6300)

EFFECTS OF CORTISONE ON THE THYROID FUNCTION IN WHITE LEGHORN COCKERELS. C.J. Shellabarger. Endocrinology 55, 100 (1954). Abstract

In the starved, nonwatered, 2- or 3-day-old White Leghorn cockerel, cortisone (E alcohol) inhibited the thyroïdal uptake of radioactive iodine in the presence or absence of TSH. E acetate was also effective in the latter respect.

The mechanism of cortisone action is discussed, and the data are interpreted as not being the result of pituitary inhibition of TSH but rather of a direct action of cortisone on the thyroid gland, although other mechanisms cannot be excluded.

Division of Bacteriology and Virology

(6300)

Incorporation of C^{14} -Labeled Amino Acids by "Trichinella Spiralis" Larvae

A series of experiments in collaboration with Dr. L.V. Hanks of the Division of Biochemistry has demonstrated the uptake of glycine-2- C^{14} and dl -alanine-2- C^{14} by *Trichinella spiralis* larvae. Newborn larvae and encysted muscle larvae were tested for their ability to incorporate C^{14} -labeled amino acids by feeding the infected host the C^{14} -labeled diets. Several groups of animals with *Trichinella* infections of two and eight weeks' duration were fed a micro-mixed Purina diet containing either glycine-2- C^{14} or dl -alanine-2- C^{14} . The mice were maintained on the diets for five to seven days and were then sacrificed for recovery of the parasites by pepsin digest. Immature larvae from the 2-week-old infection and encysted muscle larvae from the 8-week-old infection were frozen and dried under high vacuum prior to analysis for carbon-14 content. Incorporation of glycine-2- C^{14} and dl -alanine-2- C^{14} was found by C^{14} analysis of the immature larvae and the encysted muscle larvae. Further studies are under way to determine rates of amino acid C^{14} incorporation of *Trichinella* compared with those of the body tissues and blood components of the host. (R.D. Stoner)

Radiosensitivity of Tissues

Preliminary work is under way in a study of the sensitivity of intraocular transplants to beta irradiation. A thallium-204 beta source is being used to irradiate tissue implants growing in the anterior chamber of the eyes of mice. (R.D. Stoner)

*Guest of Medical Department from University of Indiana.

**Guest of Medical Department from McCollum-Pratt Institute, Johns Hopkins University.

Tetanus Antitoxin Formation by Young and Old Thymus Tissues

Work is continuing on a study of antibody formation to tetanus toxoid by intraocular transplants of young and old thymus donor tissues. Donor mice were sensitized with tetanus toxoid when 3 weeks and 6.5 months of age. Transplants of the anterior chamber of the eye were made when the sensitized donor mice were 13 weeks and 8 months of age. Both young and old thymus tissue produced significant amounts of tetanus antitoxin while growing as intraocular transplants. (R.D. Stoner)

Radiosensitivity of Bacteria

A thallium-204 beta radiation source has been developed to study the characteristics of bacterial growth during continuous irradiation. (R.M. Drew)

Division of Biochemistry

(6300)

The Reaction of Astatine with Protein

Astatine, element 85, is a halogen occurring below iodine in the periodic table. The relationship of astatine and iodine suggested the possibility of combining astatine with protein by a procedure similar to that used in the preparation of iodoalbumin. The radioactive isotope, At^{211} , has a half-life of 7.5 hr, and emits a 5.86-Mev alpha particle. This suggests that if astatine were incorporated into antibody protein, it might serve as a source of radioactivity which could be specifically directed against bacteria, viruses, or tumor cells. Since the path length of the alpha particle in tissue would be approximately 70 microns, there would be a good chance that a large portion of the 5.86 Mev of energy would be delivered in the immediate vicinity of the bacterium, virus, or tumor cell with which the antibody was combined.

The alpha particle also possesses unusual possibilities in radioautographic techniques, since its path in photographic emulsions is clearly defined and of constant length. This should make it possible to determine the precise intracellular location of any substance which can be specifically tagged with astatine. Such tagging could be carried out before injection of the substance in vivo, or the astatine could be incorporated into a compound known to interact specifically with some cellular substance (such as an antibody to a cellular protein), and the resulting compound added during the histological preparation of a tissue section.

An improved procedure was developed for the removal of At^{211} from the bismuth targets, which were prepared by W. Harris and the members of the 60-in. cyclotron crew. The targets were dissolved in nitric acid, and the astatine was reduced to zero valence with ferrous ion and extracted into ether. The astatine was then oxidized to a valence of plus one and back extracted into water with hypochlorous acid.

Tyrosine was astatinated, by means of astatine and carrier N-iodosuccinimide. Preliminary experiments with proteins indicate that although proteins may be astatinated, a considerable portion of the astatine is combined in a form more labile than in astato-tyrosine. Astato-gamma-globulin was prepared and injected intravenously into a rabbit. The astato-gamma-globulin disappeared from the plasma at approximately twice the rate of a subsequent injection of a similarly iodinated preparation of the same gamma-globulin. (W.L. Hughes* and D. Gitlin**)

*Guest of the Medical Department from McCollum-Pratt Institute, Johns Hopkins University.

**Guest of the Medical Department from the Children's Hospital, Boston, Mass.

The Absorption and Metabolism of Glycerides

Further studies on the metabolism of C^{14} -glycerol-labeled triolein have revealed that about 10 percent of the ingested fat was completely split in the lumen of the intestine, as shown by the isolation of radioactive glycerol whose specific activity was about 36 percent of that of the ingested triolein. Also isolated from the contents of the small intestine were small amounts of radioactive phosphatides with a specific activity that was about 80 percent of that of the triolein. The activity of the glycerides isolated from the lumen was equal to that of the triolein. These data suggest that the phosphatides were formed directly from the glycerides in the lumen without free glycerol being an intermediate. (L.I. Gidez)

Studies on Liver Lipides

Investigations have begun on the composition of the liver glycerides of normal rats and rats fed a fat-free diet, and the incorporation of C^{14} -glycerol into these glycerides. Methods have been developed for the separation of the glycerides from as little as 600 mg of the acetone-soluble fat of the liver into as many as seven fractions by low temperature crystallization from anhydrous acetone. The following determinations were made on these fractions: the saponification equivalent, the amount of unsaponifiable material, the iodine value, the amount of poly-unsaturated fatty acids, and the total carbon content. The specific activity of each of the fractions was also determined.

Animals suffering from a severe fat deficiency had enlarged fatty livers which contained a relatively low percentage of phosphatides. The iodine values obtained for the fatty acids of the total acetone soluble fat from fat-deficient animals was 60.5, compared with the value of 80.0 for the control group. The seven fractions of glycerides from the deficient rats had iodine values ranging from 80.5, for the glycerides soluble in acetone at -55° , to 33.1 for the glycerides soluble in acetone at 25° . Preliminary determinations on the carbon-14 content of these fractions suggest that they are turning over at different rates. (L.I. Gidez and R.A. Meigs*)

Studies on Nephrotic Syndrome

Seven children with nephrotic syndrome are currently being studied in the Medical Department. Although the metabolism of calcium, magnesium, and lipides is known to be abnormal in the nephrotic syndrome, the dynamic processes involved remain largely unexplored. As a basis for a better understanding of these processes, a study has been undertaken which will include the following:

1. The frog heart technique of McClean and Hastings (J. Biol. Chem. 107, 337 (1934)) is being used to determine active calcium as it occurs in the serum and other extracellular fluids. The inactive calcium bound to protein, citrate, and possibly lipides of lipoproteins is being determined from the difference between the frog heart value and the total calcium.
2. The proteins and lipoproteins of the plasma are being separated by electrophoresis on filter paper and, after staining, are being determined by photodensitometry.
3. Total protein, total serum lipides, cholesterol, neutral fat, phospholipides, and unsaturated fatty acids, as well as serum sodium, potassium, chloride, and phosphate are being measured. The integration of the data obtained from samples of blood from such a series of patients, all under the same regimen, will be of primary value in understanding the nephrotic syndrome. (P.E. Carson and L.I. Gidez)

*Student guest of the Medical Department from Harvard University, Boston, Mass.

The Fractionation of Histones

Histones are basic proteins which occur in the nuclei of the somatic cells of all organisms, in intimate association with the genetic material. Nothing is known about their function in the cell. Published experiments show clearly that all histone preparations which have been studied are mixtures. To understand the nature and function of these proteins, it will be necessary to isolate the individual components of such mixtures by modern methods of separation. The histone of duck erythrocytes is relatively easy to prepare. A systematic search is being made for conditions which will resolve duck erythrocyte histone into pure components, either by counter-current distribution or by ion exchange chromatography. (E.A. Popenoe)

Preparation of N-(2-Amino-Benzoyl)-Glycine and N-(2-Amino-Benzoyl)-Alanine

C¹⁴-labeled anthranilic acid has been fed to rabbits and rats. When the urine from these animals was chromatographed on filter paper, and radioautographs were made of the chromatograms, several spots were obtained which could not be identified. It was felt that two of these spots might be N-(2-amino-benzoyl)-glycine and N-(2-amino-benzoyl)-alanine. The N-(2-amino-benzoyl)-glycine has never been described in the literature. These two compounds have now been synthesized and a comparison of the natural and synthetic products is being made. (L.V. Hanks)

Division of Physiology

Tissue Physiology

(6300)

Scrutiny of the methods available for assay of acetylcholinesterase (ACHE) in tissues revealed that none was specific or sensitive enough for the study of irradiation effects on ACHE-containing tissues. A rapid, specific, and flexible ultramicromethod was therefore developed which is directly applicable to the quantitation of acetate elaborated by the enzyme even in the presence of very labile acetate compounds. In conjunction with the procedure for the determination of monoamine oxidase activity (Cotzias and Dole, J. Biol. Chem. 190, 665 (1951)), it now appears possible to study the kinetics of both enzymes' inactivation by irradiation. Experiments with rat brain homogenates revealed that both the esterase and the oxidase activities declined in linear proportion to the irradiation dose, the plots with the two enzymes being superimposable. This was true with various dose rates of γ -rays from a Co⁶⁰ source. The two enzymes did not display their individuality in the irradiation field, but behaved as if the properties of their "milieu" determined their fate. Soluble monoamine oxidase (Cotzias, Serlin, and Greenough, Science 120, 144 (1954)) is considerably more susceptible to irradiation than its mitochondrial counterpart, and clearly shows a nonlinear, first order reaction of inactivation. Preliminary data with the mitochondrial enzyme show linear proportionality between dose and inactivation.

In view of the fact that a long-chain alcohol has been used to solubilize the enzyme, the effects of normal alcohols on this system were studied with and without irradiation. It was found that the alcohol per se is not responsible for the phenomena under study. Another series of phenomena revealed themselves, however. It appears that n-alcohols of a chain length of 5-8 carbon atoms are monoamine oxidase inhibitors, whereas shorter or longer alcohols have questionable effects, if any. This type of inhibition is clearly temperature dependent, being more marked at lower temperatures. (G.C. Cotzias and I. Serlin)

Studies in Humans with I^{131} Serum Albumin

(6300)

Results of extension of the metabolic studies in humans with I^{131} serum albumin in which plasma specific activity and urinary excretion have been followed up to 60 days following injection make it appear reasonable to mathematically treat deiodination of the protein and excretion of the tag and the internal loss of tag from the readily exchangeable albumin pool to a slowly turned-over extravascular compartment as exponential but non-feed-back processes. When the total retention curve is appropriately corrected and plasma specific activity expressed as plasma fraction of radioactivity retained in the exchangeable albumin pool per gram of albumin, the plasma specific activity curve reaches an equilibrium value in three to four days and can be broken down into an equilibrium value and two exponential terms, and subjected to classical analysis of a three-compartment closed system.

A heavily collimated counter shield of high resolution has been designed and constructed for determination of transfer rates of locally injected I^{131} serum albumin and other substances tagged with gamma-emitting isotopes.

The study of long-term therapy of thyroid carcinoma with repeated doses of I^{131} is being continued. An analysis of hematologic response to initial therapeutic radioiodine compared with the response to subsequent isotope administration is under way. (C.G. Lewallen)

Physiology of Manganese

(6210)

Investigation of the in vivo behavior of radiomanganese (Mn^{56}) has continued, with the aim of studying the basic physiology of this element and developing possible therapeutic applications.

The excretion of injected manganese in large amounts by the biliary system has been confirmed, and the evidence suggests that this excreted manganese may be bound in the form of a coordination complex to some factor present in bile. In view of the unusual nature of this route of excretion, further studies are being pursued.

With the possibility in mind that the uptake of Mn^{56} by the liver might be enhanced, and its usefulness as a therapeutic agent in malignant disease of the liver in humans thereby improved, investigations of the effects of complexing agents on the absorption and distribution of injected Mn^{56} have been initiated. Results of preliminary experiments suggest that citrate and maleate complexes enhance the uptake of Mn^{56} by the liver, and that pyrophosphate and versenate markedly inhibit it.

Further trials of Mn^{56} as a therapeutic agent in liver carcinoma in humans will be made in the near future. (L.S. Maynard)

Irradiation Damage to Skin

(6210)

Among the brain tumor patients who underwent capture therapy, some developed delayed skin ulcerations at the site of the neutron entry. The ulcerations were confined to the skin flap which had been created during craniotomy two weeks earlier. The site and behavior of these ulcerations suggested that a sensitization to irradiation of only the manipulated skin had been induced during the operative procedure. In view of the clinical importance of this phenomenon, studies of animals have been initiated which are directed toward its elucidation and control. (W.H. Perkins)

Potassium Transport in the Heart and Brain

(6300)

A technique for in vivo evaluation of potassium transport between plasma and extracellular fluid (ECF) and between extracellular and intracellular fluid (ICF) in

individual organs has been evolved. This technique utilized a continuous simultaneous measurement for 30 minutes of radiopotassium (K^{42}) activity in the vascular influx and outflux of the organ under study. In the myocardium of the anesthetized dog, the plasma-ECF exchange rate was found to average 1.9 mEq/kg of left ventricle (wet weight)/min and the ECF-ICF exchange rate was found to average 4.2 mEq/kg/min. The plasma-ECF rate is of the proper magnitude expected if the process were blood flow limited. This concept was given further corroboration by the reduction of plasma-ECF exchange to one-half normal in dogs having hemorrhagic shock when coronary plasma flow was reduced to one-half normal. Studies of brain K^{42} transport are not sufficiently advanced for proper quantitation, but it is definite that K^{42} exchange between blood and cerebro-spinal fluid is much slower than that between blood and ECF, presumably because of arachnoid membrane interference. It is also apparent that exchange between CSF and brain is slower than between ECF and ICF, presumably because of pia membrane interference. The results finally suggest that most of the cerebro-spinal fluid is elucidated from the choroid plexus and from the capillaries deep in the brain which are surrounded by Virchow-Robin spaces. (H.L. Conn, Jr.)

Chloroform (Cl^{38}) and Xe^{133} Blood-Brain Kinetics in the Dog

Venous and arterial levels of anesthetic agents were taken after administration by tapping the vessels to provide continuous flow sampling for a scintillation detector followed by a logarithmic counting rate meter. The initial phase to saturation appears to be blood flow limited, which indicates that diffusion does not control the process. In the case of chloroform, the desaturation curves were atypical and further study was required to determine the nature of processes which are altering the release pattern, either chemical or physiological. Correlation of tissue concentrations was corroborated by taking samples from the brain at termination. The uptake process will be analyzed as a two-compartment system. (R.M. Featherstone,* C.B. Pittinger,* and H.L. Conn)

Boron Toxicity in Mice

(6210)

A number of factors which may or may not modify boron toxicity in mice have been investigated or are under investigation. The factors under scrutiny which may alter borax toxicity are: intravenous administration rate, hypodermic needle size under intravenous administration, temperature of the administered boron preparation, selection of a chemical class for a vehicle, variation in the vehicle preparation within a given chemical class, molar relationships of borax to the vehicle being evaluated, pretreatment with some selected vehicle preparation, population age, variation among populations, and animal environmental temperature. In connection with the variation in the vehicle within a chemical class, toxicological data are being obtained on those materials which are independent of their incorporation with borax. (O.D. Easterday)

Lithium Toxicity in Mice

(6210)

The study of problems associated with inorganic lithium and boron derivatives with respect to their toxicity and chemical characteristics was continued. (O.D. Easterday and L.E. Farr)

Borax Kinetics

(6210)

Boron blood levels and the associated kinetics involved in the removal of intravenously administered boron from the circulation system are being studied in the dog and

*State University of Iowa.

mouse. In the dog studies, blood samples are withdrawn at varying intervals of time after the administration of borax and are analyzed for boron content. In the mouse studies, at varying time intervals after the administration of borax the animals are sacrificed, bled, and the boron blood levels are determined. The influence of the vehicle under these conditions is also being evaluated. (O.D. Easterday and L.E. Farr)

Electrocardiography of Boron (6210)

Preliminary studies are being conducted in order to evaluate the effect of boron on the mouse heart. These studies have not progressed sufficiently so that a comment may be included in this report. (O.D. Easterday)

Physical Properties of Boron (6210)

Several physical chemical studies have been completed or are at present in progress. These include an investigation of the stability of borax vehicle preparation and its solubility characteristics.

Because of the critical nature of the borax solubility in these studies, a system has been devised whereby the temperature of the injectable solutions may be controlled quite accurately. This system merely incorporates the use of a water jacket on a 0.25-ml tuberculin syringe or vial and the circulation of water of a constant temperature through the system. Heating mantles have also been employed in controlling the temperature of the boron solutions. (O.D. Easterday)

Modifications of Equipment Used in the Medical Facility of the Reactor (6210)

Attempts are being made to improve the physical conditions under which animals are irradiated in the medical facility of the Brookhaven reactor. Problems related to the air supply, body heat, exposure uniformity, urine absorption, and radiation filters are under further study. (O.D. Easterday and L.E. Farr)

Ion Transport Across Cell Surfaces (6300)

When human red cells were suspended in media containing 136 mEq/liter K and 2 mEq/liter Na at 25°C, the following changes in sodium transport took place (as compared to controls in simulated plasma media containing 133 mEq/liter Na and 5 mEq/liter K): 1) Inward rate constant, k^i_{Na} , increased fourfold; inward flux decreased 17-fold. 2) Outward rate constant, k^o_{Na} , as well as Na outflux decreased progressively, while a net cell sodium loss took place for at least 40 hr, at which time cell sodium had decreased from an initial 8 mEq/liter cells to 0.5 mEq/liter cells. Media were tagged with Na^{24} for influx measurements and cells were pretagged, washed, and re-suspended in nonradioactive media for outflux measurements.

In the outflux determination, a semilogarithmic plot of cell Na^{24} activity against time resulted in a curve which could be expressed as the sum of two exponential terms. The two resulting k^o_{Na} 's were $k_1 = 0.21 \text{ hr}^{-1}$, and $k_2 = 0.045 \text{ hr}^{-1}$. Other possible interpretations include a time course reduction in k^o_{Na} (in vitro) or a dependency of k^o_{Na} on cell Na concentration. Metabolic activity, as determined by lactate production, is little, if any, diminished in "high K" media. Further experiments, designed to distinguish between the above alternatives, are in progress. It appears quite possible, however, that "nonexchangeable" red cell sodium which has been reported by several investigators is, in fact, a more slowly exchanging component.

Since cell sodium analysis by flame photometry is subject to considerable error at low cell sodium levels, another method of analysis has been used, with the assistance of the Biophysics Division and the Chemistry Department. This entails neutron

bombardment of unknowns and standards in the medical thermal column, and subsequent scintillation counting of high energy (2.8 Mev) Na^{24} gamma emissions, with beta exclusion by aluminum absorption, and gamma (below 2 Mev) exclusion by means of a pulse height discriminator. Virtually complete exclusion of all activated elements present in red cells other than sodium was achieved. Gamma counting by pulse height discrimination also appears to offer substantial advantages over other methods presently employed to distinguish between Na^{24} and K^{42} in biological tracer studies in which simultaneous use of these two isotopes is desirable. (E.T. Dunham)

Reactor Radiation in Tissue

(6210)

A handling device has been built for a set of tissue-equivalent ionization chambers to provide for complete scanning of the radiation pattern throughout a liquid phantom. The phantom is also specifically compounded to be completely tissue-equivalent for all components of the mixed radiation emerging from the reactor. This experiment measures the total radiation dose in tissue at all points and provides separate measurements of the contribution to the total dose by the proton resulting from neutron capture in the nitrogen constituent of tissue as well as the additional effect in tissue containing boron at the levels used in the neutron capture therapy experiment. Previous work has given limited data of this nature in studies in a horizontal collimated beam of neutrons of relatively low flux. The present apparatus is built to explore the situation under actual conditions used in the therapy experiment. (E. Stickley in collaboration with H.H. Rossi,* W.A. Higinbotham,** and J.B.H. Kuper**)

Solubility of Xenon

(6300)

The investigation of the anesthetic properties of xenon gas required further measurements of the specific solubility of the gas in various chemical and physiological systems. This work was completed and is being evaluated with special attention to the possible influence of radioactive contaminants suspected in the earlier experiments. (E. Stickley and L. Levy†)

Irradiation of "E. coli" by the B^{10} (n, α) Li Reactions

(6300)

Cultures of E. coli were given a calculated dose of 134,000 rads of heavy particle radiation by exposure to slow neutrons at the medical treatment facility in a medium containing 100 μg of B^{10} per ml. On subsequent recultivation, these specimens had a survival of 0.1 percent which may be compared with nonirradiated controls with or without boron present. A culture irradiated without boron showed over 50 percent survival. This work was carried out as a step in the development of a biological measure of mixed radiation effects in tissue. Tissue-equivalent solid and liquid media are being developed to provide means for dispersing and sampling suitable microorganisms throughout a volume of tissue under conditions of the neutron capture therapy experiment. (B. Seife,†† R.M. Drew, and E. Stickley)

Neutron Capture Therapy of Experimental Tumors Localized in Organs Other than Brain

(6210)

In order further to investigate the applicability of neutron capture therapy to

*Radiological Research Laboratory, Columbia University.

**Instrumentation and Health Physics Department.

†Georgetown University.

††Student guest from University of Rochester College of Medicine and Dentistry.

organs other than brain, a laboratory model is desirable. The methylcholanthrene-induced glial tumor, used by this Division in brain tumor studies, is again being used since considerable data are available regarding its transplantability and susceptibility to gamma radiation. The lung was chosen as the organ site for implantation, in part because its physiological characteristics are markedly different from brain. A procedure had to be developed which promised reasonable success in transplantation in this organ, since none was found in the literature. Four methods of implantation have been investigated:

1. Inhalation - drops of tumor suspension placed on nostrils of anesthetized mice.
2. Intratracheal - hypodermic needle placed inside trachea.
3. Intrabronchial - tubing inserted through trachea and into one primary bronchus.
4. Intrathoracic - tumor injected through chest wall into lungs.
5. Intravenous.

In terms of ease of operation, yield, and reproducibility, the data indicate thus far that the inhalation and intratracheal methods are superior to the others. (R.H. Tamis*)

(6300)

PREPARATION OF SOLUBLE MONOAMINE OXIDASE. G.C. Cotzias, I. Serlin and J.J. Greenough. *Science* 120, 144 (1954). Abstract.

The difficulties encountered in attempts to purify monoamine oxidase have led to the assumption that this system is comprised of many enzymes acting in sequence or in unison. The fact that this enzyme is tightly bound to the mitochondria (Cotzias and Dole, *Proc. Soc. Exptl. Biol. Med.* 78, 157 (1951)) led to a search for a procedure which would allow its dissociation from these particles. Several detergents were found to be capable of bringing on this effect. Of these, the most active was Isooctylphenoxyethoxyethanol. Evidence of actual solubilization of this enzyme is presented. The fact that several simple intracellular enzymes were also solubilized while the succinoxidase complex was destroyed by this procedure suggested that monoamine oxidase is probably a single enzyme rather than a complex.

(6210)

THE EXPERIMENTAL APPLICATION OF NEUTRON CAPTURE THERAPY TO GLIOBLASTOMA MULTIFORME. L.E. Farr. Vith International Cancer Congress, Sao Paulo, Brazil, July 26, 1954. Abstract of talk.

The physics and physiology of neutron capture therapy were briefly reviewed in relation to the use of boron-10 as the capture element. Preliminary interim results were given regarding results observed in four patients with four months postradiation. These patients were the first to receive treatment with thermal neutrons at a flux of 2×10^9 neutrons/cm²/sec at the skin surface. This flux is approximately 20 times that used for the first 10 patients treated. The results over a 4-month interval were encouraging. The development of skin complications and means to prevent or ameliorate them were briefly discussed.

*Student guest from New York University College of Medicine.