AUG 14 1963

ON THE UNIQUENESS OF THE PARTIAL WAVE AMPLITUDES

A. P. Balachandran

MASTER

THE UNIVERSITY OF CHICAGO

THE ENRICO FERMI INSTITUTE FOR NUCLEAR STUDIES

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

ON THE UNIQUENESS OF THE PARTIAL WAVE AMPLITUDES

A. P. Balachandran

The Enrico Fermi Institute for Nuclear Studies
and Department of Physics
The University of Chicago, Chicago, Illinois

Facsimile Price \$ 1.10
Microfilm Price \$.88

Available from the
Office of Technical Services
Department of Commerce
Washington 25, D. C.

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

Contract No. AT(11-1)-264

ON THE UNIQUENESS OF THE PARTIAL WAVE AMPLITUDES*

A. P. Balachandran[†]

The Enrico Fermi Institute for Nuclear Studies
and Department of Physics
The University of Chicago, Chicago, Illinois

Important insight into the structure of strong interactions of elementary particles has been achieved recently by the study of the ambiguities associated with the solutions of the Mandelstam representation. In this field, which was initiated by Froissart,¹ it is customary to assume^{1,2} that the double-spectral functions associated with the process are given for some suitable values of the energy variables. Here, we will consider a related problem where it will instead be assumed that all but a finite number N of partial waves associated with a two-body scattering amplitude are given and the ambiguities associated with the remaining N partial waves will be examined. It will be proved that the remaining N partial waves are in fact uniquely determined except possibly for an additive s -wave constant if either a) the scattering amplitude has crossing symmetry, or b) in one of the crossed channels, there exists an

*Work supported by the U. S. Atomic Energy Commission.

[†]On leave of absence from the Institute of Mathematical Sciences, Madras, India.

energy region where the scattering is purely elastic.³⁾ In case a), a knowledge of the precise nature of the analyticity of the scattering amplitude is not required. It need only be assumed that the analytic continuation of the same amplitude determines the two processes we are interested in.

For simplicity, we will consider the scattering of two spinless particles of equal mass m . Let s denote the square of the center-of-mass energy in one of the three Mandelstam channels, and let θ be the corresponding scattering angle. The invariant momentum transfer t is defined through the usual relation

$$t = -2k^2 (1 - \cos \theta) \quad (1a),$$

$$\text{i.e. } \cos \theta = 1 + \frac{2t}{s - 4m^2} \quad (1b)$$

where k denotes the centre-of-mass momentum in this channel.

The partial wave amplitude $F_\ell(s)$ is defined as

$$F_\ell(s) = \left[\frac{s}{s - 4m^2} \right]^{\frac{1}{2}} \frac{[\eta_\ell(s) e^{2i\delta_\ell(s)} - 1]}{2i} \quad (2)$$

in a familiar notation.

To proceed further, let us assume that all the F_ℓ 's for ℓ greater than some integer N are given. The proof goes through with trivial alterations when all but any finite number of the F_ℓ 's are given. The ambiguity we wish to resolve is associated with the possibility that for $\ell \leq N$, there may exist

3.

at least two amplitudes, say $F_{\ell}^{(1)}(s)$ and $F_{\ell}^{(2)}(s)$, which may describe the scattering for these ℓ 's. If this were possible, we may construct two scattering amplitudes $F^{(1)}(s, t)$ and $F^{(2)}(s, t)$ when $s \gg 4m^2$ and $+1 \leq \cos\theta \leq -1$ through the definitions

$$F^{(1)}(s, t) = \sum_{\ell=0}^N (2\ell+1) F_{\ell}^{(1)}(s) P_{\ell}\left(1 + \frac{2t}{s-4m^2}\right) + \sum_{\ell=N+1}^{\infty} (2\ell+1) F_{\ell}(s) P_{\ell}\left(1 + \frac{2t}{s-4m^2}\right),$$

$$F^{(2)}(s, t) = \sum_{\ell=0}^N (2\ell+1) F_{\ell}^{(2)}(s) P_{\ell}\left(1 + \frac{2t}{s-4m^2}\right) + \sum_{\ell=N+1}^{\infty} (2\ell+1) F_{\ell}(s) P_{\ell}\left(1 + \frac{2t}{s-4m^2}\right)$$

(3)

Eq. (3) shows that $F^{(1)}(s, t) - F^{(2)}(s, t)$ is a polynomial in t of degree N :

$$F^{(1)}(s, t) - F^{(2)}(s, t) = \sum_{n=0}^N b_n(s) t^n \quad (4)$$

where the b_n 's are linear combinations of $F_{\ell}^{(1)}(s) - F_{\ell}^{(2)}(s)$ with coefficients which remain bounded as s tends to infinity.

Let us now examine case a). Crossing symmetry implies that $F^{(1)}(s, t) = F^{(1)}(t, s)$, $F^{(2)}(s, t) = F^{(2)}(t, s)$ where we have assumed without loss of generality that the t -channel is the one that is crossing symmetric to the s -channel. (4) now gives

$$\sum_{n=0}^N b_n(s) t^n = \sum_{n=0}^N b_n(t) s^n \quad (5)$$

or $b_n(s)$ is a polynomial of degree N in s :

$$b_n(s) = \sum_{r=0}^N a_{n,r} s^r \quad (6)$$

But the partial waves are bounded at infinity and hence the b_n 's must be constants for all n . (5) then shows that only $b_0(s)$ can be a non-zero constant. Thus $F^{(1)}(s,t)$ and $F^{(2)}(s,t)$ can differ at most by a constant which is what we set out to prove. If we assume at this point that the η_0 's go to zero as s tends to infinity, which seems a physically plausible assumption (dominance of inelastic processes at high energies), we see that this constant also must be zero.

To discuss case b), we note that (4) implies that $F^{(1)}(s,t) - F^{(2)}(s,t)$ is regular for every finite t and fixed s . The absorptive part of $F^{(1)}(s,t) - F^{(2)}(s,t)$ in the t -channel is therefore zero. If $\varphi_l^{(1)}(t)$ and $\varphi_l^{(2)}(t)$ denote the partial wave amplitudes in the t -channel, it follows that

$$\text{Im } \varphi_l^{(1)}(t) = \text{Im } \varphi_l^{(2)}(t) \quad (7)$$

for all l and $t \gg 4m^2$. By hypothesis, there is an interval $4m^2 \leq t < t_1$ in the t -channel where elastic unitarity is applicable. In this interval, as a consequence (7), one finds

$$\text{Re } \varphi_l^{(1)}(t) = \pm \text{Re } \varphi_l^{(2)}(t). \quad (8)$$

Let M denote the number of subtractions in the Mandelstam representation. The case $\text{Re } \varphi_l^{(1)}(t) = -\text{Re } \varphi_l^{(2)}(t)$ may then be

eliminated for $l > M$ by appealing to an elegant argument due to Martin.² Thus

$$\varphi_l^{(1)}(t) = \varphi_l^{(2)}(t) \quad (9)$$

for $l > M$ and $4m^2 \leq t < t_1$ and by analytic continuation, for all t . $F^{(1)}(s, t) - F^{(2)}(s, t)$ now reduces to a polynomial of degree M in s and the rest of the argument proceeds as before.

Let t_2 be the threshold for the second inelastic process in the t -channel and let the first inelastic channel be also a two-body process. A simple application of partial wave unitarity then reveals that the partial waves in this new channel are also uniquely determined up to a phase for $t_1 < t < t_2$. However, it seems difficult to carry this argument further since the phase may become complex for $t > t_2$ or $t < t_1$.

We may comment in conclusion on the relevance of the preceding remarks to the hypothesis that all particles are Regge poles.⁴ The partial wave amplitudes satisfy standard dispersion relations which however have a multiplicity of solutions.⁵ It has been seen that for a class of processes, if these amplitudes are given for $l > N$, there is one preferred solution for $l \leq N$. This is equivalent to the statement that if we are given the interpolating partial wave amplitude $F(s, l)$ of complex angular momentum theory,⁶ which is a priori defined only for $\text{Re } l$ greater than some integer M , the physical amplitudes for $l \leq M$ are in fact uniquely determined. The crucial point then is whether or not the analytic continuation of $F(s, l)$ for $\text{Re } l \leq M$ coincides

with the preferred solutions of the partial wave dispersion relations.⁷ It is important to observe in this connection that if we define an amplitude

$$F'(s,t) = \sum_{l=0}^{\infty} (2l+1) F(s,l) P_l(\cos\theta) \quad (10),$$

there is apparently no reason to insist that $F'(s,t)$ satisfy either crossing symmetry or unitarity in one of the crossed channels.⁸ Otherwise, it would be a simple matter to prove that all particles are Regge poles.

The author wishes to thank Professor R. H. Dalitz and Professor R. Oehme for discussions. He is particularly indebted to Dr. F. von Hippel for his generous comments and criticisms.

References

1. M. Froissart, Phys. Rev. 123, 1053 (1961).
2. A. Martin, Phys. Rev. Letters 9, 410 (1962).
3. The result in case b) has been independently obtained by A. Martin and by R. Oehme. The author is indebted to Professor Oehme for informing him of his and Martin's results.
4. G. F. Chew and S. C. Frautschi, Phys. Rev. Letters 1, 394 (1961); 8, 41 (1962).
5. L. Castillejo, R. H. Dalitz and F. J. Dyson, Phys. Rev. 101, 453 (1956).
6. M. Froissart, La Jolla Conference on the Theory of Weak and Strong Interactions, 1961 (unpublished); K. Bardakci, Phys. Rev. 127, 1832 (1962).
7. Since all the poles of $F(s, \ell)$ are Regge poles. See R. Oehme, Phys. Rev. Letters 9, 358 (1962); Phys. Rev. 130, 424 (1963).
8. This point was repeatedly emphasized to the author by Professor R. Oehme. Similar remarks have also been made by A. Martin (private communication to Professor R. Oehme).