MASTER

Progress Report of Contract With
U.S. Atomic Energy Commission
Division of Biology and Medicine

Contract No.: AT-(11-1)-1783   October 14, 1971

Institution: Case-Western Reserve University

Department: Biochemistry

Title: The Role of Inorganic Pyrophosphate in the Metabolism
of Phosphoenolpyruvate

Principal Investigator: Harland G. Wood

INTRODUCTION

Inorganic pyrophosphate (PP_i) is a metabolic product of many reactions of biological importance. In the past, it has generally been considered to be removed by hydrolysis, thereby promoting synthetic reactions by pulling the reaction to completion. However, through the work of Baltscheffsky [Science 153, 1120 (1966)]; Keister [Bookhaven Sympos. Biol. 19, 246 (1966)]; Nordlie [J. Biol. Chem. 245, 6078 (1970)]; Reeves [Biochem. Biophys. Acta 220, 346 (1970)] and others, concepts are being developed whereby PP_i is believed to play an important role in energy conservation and utilization in many organisms. An excellent example of one aspect of these concepts is the scheme proposed by Reeves [Biochim. Biophys. Acta 220, 346 (1970)] to explain the conversion of P-enolpyruvate to pyruvate in Entamoeba histolytica, an organism lacking pyruvate kinase. This scheme is outlined in Figure 1, and involves the two new and interesting reactions, P-enolpyruvate carboxy-transphosphorylase and pyruvate, phosphate dikinase. Both of these enzymes have been found in the propionic acid bacteria, as well as the amoeba. The carboxy-transphosphorylase from Propionibacterium shermanii catalyzes the reactions:

Reaction (1), referred to as the oxaloacetate reaction, is freely reversible and is stimulated by thiols. In the reverse reaction, the energy of PP_i is being utilized to synthesize the biologically important compound, P-enolpyruvate. Reaction (2), referred to as the pyruvate reaction, is irreversible and is inhibited by thiols and by CO₂. It is interesting to note that the enzyme from the amoeba does not catalyze the pyruvate reaction (Reeves, R. E., Personal Communication).

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

Figure 1

Metabolism of P-enolpyruvate in *Entamoeba histolytica* as Proposed by Reeves

Pyruvate, phosphate dikinase (PPDK) catalyzes the reaction:

The enzyme is referred to as a dikinase, since the reaction represents the phosphorylation of both P_i and pyruvate, when viewed in the reverse reaction.

In last year's Proposal we planned to study the following aspects of these two enzymes: (I) The Mechanism of the Carboxytransphosphorylase Reaction, which included (A) Subunits of Carboxytransphosphorylase, (B) Detection of Enzyme Bound Intermediate Compounds and Examination with ^{18}O of the Mechanism of the Carboxytransphosphorylase Reaction, (C) Kinetic Analysis of the Carboxytransphosphorylase Reaction and (II) Catalysis and Mechanism of the Pyruvate Phosphate Dikinase Reaction. Item (C) still has not been attempted, due to difficulties with other aspects of the problem.

SUMMARY

Catalysis by carboxytransphosphorylase appears to be a two-step process and several mechanisms have been considered which fit this observation and are outlined in Figure 2a, b, c (Pages 6-8). These mechanisms involve: (a) pyrophosphoenolpyruvate or (b) a cyclic mixed anhydride of P-enolpyruvate as an intermediate or (c) a displacement reaction at the P-enolpyruvate carbonyl. All of these are hydrolytic steps and predict the incorporation of solvent water into: (a) pyrophosphate, (b) the α -carboxyl of the keto acid product or (c) the carbonyl of the keto acid product respectively. We have investigated the incorporation of isotopic water into the products of the carboxytransphosphorylase reaction and have found that none of the products are labeled. By use of ^{18}O -labeled P-enolpyruvate, we have found that the reaction proceeds with cleavage of the oxygen-phosphorus bond of P-enolpyruvate. These studies lead to the conclusion that the reaction mechanism does not involve a hydrolytic step, as predicted by the above mechanisms. We believe that the mechanism proceeds via classical attack of inorganic phosphate on the P-enolpyruvate phosphoryl leading to the formation of pyrophosphate and an enzyme enolpyruvate complex. In a second step, the enolate species is carboxylated by nucleophilic attack on CO_2 to yield oxaloacetate (Reaction 1) or is released to the medium, where it undergoes protonation to keto-pyruvate (Reaction 2).

The subunit structure of carboxytransphosphorylase has been extensively studied and a manuscript has been prepared for publication on this subject. The enzyme was found to exist as three species, a tetramer, a dimer, and a monomer. It appears that all three species are active in catalysis of the oxaloacetate reaction, whereas only the monomeric and tetrameric species are active in the pyruvate reaction. As yet, it has not been possible to determine if there is a difference in the specific activities of the three species, thus the relationship of this dissociation phenomenon to a possible control mechanism for the enzyme is not known at this time.

The studies with pyruvate, phosphate dikinase have been designed to verify which of two proposed mechanisms of reactions for the pyruvate, phosphate dikinase is operative in the case of Propionibacterium shermanii and Bacteriodes symbiosus enzymes. The Andrew-Hatch mechanism (reactions (8), (9) p. 18) can be summarized

as involving two partial reactions in the first of which (reaction (8) p. 18) there is an attack of ATP and P_i on an active group in the enzyme with the formation of monophosphorylated enzyme forms. The Evans-Wood mechanism, on the other hand, postulates three partial reactions (reaction (4), (5), and (6) p. 17) in which each substrate adds to the enzyme sequentially and is converted to the diphosphorylated (or pyrophosphorylated) enzyme (reaction (4) p. 17) and then the monophosphorylated form (reaction (5) p. 17).

Three lines of evidence supporting the Evans-Wood mechanism have been obtained: a. Steady state kinetics; b. Inhibition patterns; c. Exchange reactions and exchange kinetics.

a. The steady state kinetics studies showed the reaction involves a Tri (Uni Uni) ping pong mechanism. This supports strongly the Evans-Wood mechanism: ATP adds to the enzyme, a modified enzyme form is formed (postulated to be the diphosphorylated form) and AMP leaves it (reaction (4)) then P_i adds to the enzyme when a second form of the enzyme is formed (reaction (5)) and PP_i leaves the enzyme etc. Should the Andrews and Hatch mechanism be true in these microbial enzymes, a Bi Bi Uni Uni ping pong mechanism should have been found. The steady state kinetics does not, however, tell us anything about the nature of the enzyme forms.

b. The inhibition profile of the PPDK by products of the reactions could be explained only by assuming a non-classical ping pong mechanism in which there are three sites for the substrates/product pairs AMP/ATP, pyruvate/P-enolpyruvate and P_i/PP_i . The observed inhibition profile could be explained well on the assumption of competition of various substrates and products of three different enzyme forms (presumably unphosphorylated, monophosphorylated and diphosphorylated enzyme forms).

c. The exchange reaction was investigated and is the best proof available of the validity of Evans-Wood mechanism. The three isolated partial reactions could be investigated kinetically and were found to conform to the ping pong type mechanism. The expected inhibition profile discussed above could be verified by conducting the exchange reactions at equilibrium concentrations of other substrates -- indeed a full conformation to the predicted pattern was obtained thus demonstrating once again the Evans-Wood mechanism.

DETAILED REPORT

I. The Mechanism of Reaction of Carboxytransphosphorylase--R. Singleton, Jr. and H. G. Wood. [In part in collaboration with Dr. Irwin Rose (Cancer Institute, Philadelphia, Penn.), Dr. James Willard (University of Vermont, School of Medicine), and Dr. S. J. Benkovic (Penn. State University)].

Potential Mechanisms. As has been discussed in previous reports and in publications from this laboratory, the carboxytransphosphorylase reaction appears to be a two step process, i.e. the carboxylation and phosphorylation steps are separate. Any mechanisms proposed to explain the reaction must ultimately account for the following observations: (1) The enzyme catalyzes an exchange reaction between oxaloacetate and $^{14}\text{CO}_2$, which requires the presence of PP_i and Mg^{+2} and is not inhibited by EDTA, (2) under conditions of the CO_2 /oxaloacetate exchange in the presence of EDTA, no exchange occurs between $^{32}\text{P}_i$ and PP_i or $^{32}\text{PP}_i$ and P-enolpyruvate, (3) no exchange occurs between pyruvate and oxaloacetate or P-enolpyruvate under any condition, and (4) the pyruvate reaction (reaction 2) is irreversible.

The mechanisms outlined in Figure 2 (a, b, and c) are all hydrolytic mechanisms and fit the above observations. Mechanism I (Figure 2A) was originally proposed by Wood, *et al.* [Biochem 8, 3145 (1969)] and involves the formation of pyrophosphoenolpyruvate as an intermediate. The mechanism predicts the incorporation of ^{18}O from water into pyrophosphate, via incorporation of solvent water in the hydrolytic step.

Mechanism II (Figure 2b) involves the formation of a cyclic anhydride intermediate involving the P-enolpyruvate carboxyl and its phosphoryl moiety. This species is then phosphorylated by nucleophilic attack of phosphate, generating carboxypyrophosphoenolpyruvate, which can either be carboxylated or protonated to the corresponding carboxy pyrophosphoryl- α -keto acid. This compound spontaneously hydrolyzes to pyrophosphate and the α -keto acid. Cyclic anhydride intermediates of this type have been suggested by Benkovic [JACS 93, 2522 (1971)] and others [JACS 85, 3705 (1963)] for the hydrolysis of various P-enolpyruvate ester derivatives. This mechanism predicts the incorporation of ^{18}O from water into the carboxyl group of the keto acid product. Benkovic [personal communication] has recently found that in some cases, cyclic anhydride intermediates of this type will undergo ring opening leaving the phosphoryl moiety on the carbonyl oxygen, rather than on the carboxyl. In the case of the carboxytransphosphorylase reaction, this would lead to the generation of a pyrophosphoenolpyruvate intermediate of the type discussed in Mechanism I.

Recently, we have been considering a third hydrolytic mechanism, which is outlined in Mechanism III (Figure 2c). This mechanism involves an initial concerted attack of hydroxyl ion on the carbonyl carbon, with displacement of the phosphoryl to inorganic phosphate, leaving enol-pyruvate. The mechanism predicts the incorporation of ^{18}O from water into the carbonyl oxygen of the keto acid. Furthermore,

Figure 2a

Pyrophosphoenolpyruvate as an Intermediate of the Carboxytransphosphorylase Reaction
(Proposed by Wood, et al. (Mechanism I).

Figure 2b

Cyclic Anhydride as an Intermediate of the
Carboxytransphosphorylase Reaction,
(Mechanism II)

Figure 2c

Double Displacement as a Mechanism for the
Carboxytransphosphorylase Reaction.

(Mechanism III)

5

unlike the previous mechanisms, this mechanism involves carbon-oxygen bond cleavage, rather than phosphorus-oxygen cleavage.

Reaction in H_2^{18}O . The possible involvement of these mechanisms was tested by running the carboxytransphosphorylase reaction in ^{18}O -enriched water and examining the products for incorporated ^{18}O . The results are summarized in Table I. Experiment 1 was run in the oxaloacetate reaction, and malate dehydrogenase was included in the reaction mixture to pull the reaction by removal of oxaloacetate. Experiments 2, 3, and 4 were both run in the pyruvate reaction and yeast inorganic pyrophosphatase was included in the reaction mixture to pull the reaction and to remove PP_i which inhibits the reaction. In experiment 3, lactate dehydrogenase was added to the reaction mixture to convert pyruvate to lactate, in an attempt to prevent any exchange of the carbonyl oxygen. In experiment 4, H_2O_2 was added to the reaction mixture to convert pyruvate to acetate. In all experiments, the product was isolated by exchange chromatography on QAE-Sephadex using a linear gradient of either triethylammonium acetate or bicarbonate. The ^{18}O content of PP_i and lactate was determined by pyrolysis with guanidine HCl . The ^{18}O content of the carboxyl carbon of pyruvate was determined on CO_2 formed by specific decarboxylation of the silver salt of pyruvate. The ^{18}O content of acetate was determined directly in the mass spectrometer. The results of these experiments are summarized in Table I.

There was no incorporation of ^{18}O into either pyrophosphate or the pyruvate carboxyl, as shown in experiments 1 and 2. In experiment 3 and 4 there was some incorporation of ^{18}O into the carbonyl oxygen; however, in light of the difficulties in preventing exchange of the carbonyl oxygen, we do not feel this incorporation is mechanistically significant. This conclusion is further justified in light of the experiments with $[\text{}^{18}\text{O}]$ P-enolpyruvate discussed below.

On the basis of these experiments it appears that a hydrolytic step is not involved in the carboxytransphosphorylase reaction, and that the mechanisms described above are incorrect as written. The exception to this conclusion would be if $^{18}\text{OH}^-$ was sequestered at the enzyme site and did not exchange with the solvent water as discussed below.

Effect of Hydrogen Peroxide. In the course of developing methods for determination of the ^{18}O content of the pyruvate carbonyl group, an interesting phenomenon was noted which may be related to the mechanism of the reaction. Initially, attempts were made to run the reaction in the presence of stoichiometric amounts of hydrogen peroxide. This was required, since the pyruvate carbonyl group hydrates very rapidly and therefore will exchange with solvent water. This could lead to an erroneous finding of ^{18}O incorporation. By converting the pyruvate to acetate with peroxide, this exchange can be reduced and the acetate may be used directly in the mass spectrometer for ^{18}O measurement. Such techniques have been used by Sprinson and his workers in their studies on the isocitrate dehydrogenase [J. Biol. Chem. 241, 1067 (1966)] and pyruvate kinase [BBRC 40, 1464, (1970)] reactions.

As seen in Figure 3, the utilization of P-enolpyruvate was the same in the presence or absence of peroxide. However, the acetate yield never exceeded 50% of the theoretical. When the pyruvate formed in the control reaction (absence of peroxide) was treated with the same amount of peroxide as used when the pyruvate was

Table 1

 ^{18}O Content of Products of Carboxytransphosphorylase in H_2^{18}O

Experiment No.	Product Examined	^{18}O Atom % Excess	
		Expected	Found
1 ^a	PP_i	0.173	0.00
2 ^b	Pyruvate Carboxyl	2.00	0.00
3 ^c	Pyruvate Carbonyl	2.00	0.32
4 ^d	Pyruvate Carbonyl	2.00	0.25

- a) Reaction mixture contained μmoles : P-enolpyruvate, 20; P_i , 100; MgCl_2 , 120; KHCO_3 , 300; CoCl_2 , 1; mercaptoethanol, 10; NADH, 20 and malate dehydrogenase, 200 units and carboxytransphosphorylase, 30 units. H_2^{18}O = 1.39 atom % excess. Volume = 10 ml. Incubation at 25° under CO_2 for 0.5 hours. PP_i yield = 16.5 μmoles .
- b) Reaction mixture contained in μmoles : P-enolpyruvate, 40; P_i , 200; MgCl_2 , 80; and PP_i ase, 4 units; carboxytransphosphorylase, 1.5 units. H_2^{18}O = 4 atom % excess. Volume = 20 ml. Incubation at 25° under N_2 for 5.0 hours. Pyruvate yield = 36 μmoles .
- c) Reaction mixture contained in μmoles : P-enolpyruvate, 100; P_i , 100; MgCl_2 , 40; NADH, 100 and PP_i ase, 4 units; lactate dehydrogenase, 36 units; carboxytransphosphorylase, 5 units. Volume = 10 ml. Incubation at 25° under N_2 for 9.0 hours. Lactate yield = 95 μmoles .
- d) Reaction mixture contained in μmoles : As in c, except in place of lactate dehydrogenase and NADH, 100 μmoles H_2O_2 . Acetate yield = 38 μmoles . Incubated at 25° under N_2 for 2 hours 45 minutes.

Figure 3

Effect of H_2O_2 on the carboxytransphosphorylase Reactions: The mixture contained in μ moles: P-enolpyruvate, 27; P_i , 20; $MgCl_2$, 8; H_2O_2 , 20 and 4 units, PP_i ase and 0.5 unit carboxytransphosphorylase. Incubation at 20° under N_2 . Volume = 2 ml

oxidized as it was produced, acetate was formed in 90% yield. This experiment suggests that the initial product of the enzyme reaction is some product other than keto-pyruvate, which is not oxidized to acetate by peroxide. We postulate that this initial product of the enzyme reaction is enol-pyruvate, which is in part oxidized by peroxide to some product other than acetate before it is tautomerized to keto-pyruvate. We are currently in the process of trying to identify this product.

Sequestered Water. In all of the above mechanisms, it will be noted that a hydroxyl ion is displaced from the substrate at some time prior to the hydrolytic step. If this hydroxyl species could somehow be sequestered from solvent water, and then be involved as the attacking species in the hydrolytic step, incorporation of ^{18}O from solvent water would not be found. Thus, the conclusion reached regarding the above experiments must be qualified to this extent.

Last year we proposed to approach a study of this problem, with respect to Mechanism I, as follows. Close examination of Mechanism I (Figure 2a) demonstrates that the hydroxyl group is transferred from one phosphate of pyrophosphate to the other, if one assumes sequestering of water. Thus, if one were able to distinguish between the source of the phosphates in pyrophosphate, it would be feasible to study the possibility of sequestering of water. The experiment to test this hypothesis is outlined in Figure 4, using ^{18}O phosphothioate. If the reaction occurred as in Mechanism I with sequestering of OH^- then one atom of ^{18}O would be introduced in the phosphate and two would remain in the phosphothioate.

Figure 4

Outline of Experiment Testing Role of Sequestered Water in Mechanism I.

It is obvious that this experiment involves assumptions which would need to be confirmed before the experiment could be attempted, not the least of which is that phosphothioate would serve as a substrate for the enzyme. This was tested by assaying the enzyme in the oxaloacetate reaction in the presence and absence of phosphate or phosphothioate (note: since the enzyme has a very high affinity for phosphate and since phosphate was always available in trace amounts, it was impossible to eliminate completely some reaction in the absence of added phosphate). The results of this test are summarized in Table II.

Table II
Effect of Phosphothioate on the Carboxytransphosphorylase Reaction^a

Phosphothioate ^b (μmoles/ml)	Phosphate (μmoles/ml)	% Activity	% Inactivation
3.0	---	17.0	---
15.0	---	9.5	---
30.0	---	6.2	---
0	9.0	100	0
3.0	9.0	---	34.0
15.0	9.0	---	73.0
30.0	9.0	---	85.0

^aEnzyme assayed in presence or absence of phosphothioate under standard conditions of the oxaloacetate reaction, except that phosphate was omitted from the reaction mixture where indicated. Trace amounts of phosphate arose from solutions.

^bPhosphothioate prepared by S. J. Benkovic. Purity demonstrated by paper chromatography. Structure confirmed by IR analysis.

It is apparent from these results that phosphothioate does not serve as a substrate for the enzyme; indeed it may very well be a good competitive inhibitor for the reaction. For this reason, the experiment as outlined in Figure 4 was not feasible.

Reaction with [¹⁸O] P-enolpyruvate. We have utilized P-enolpyruvate (supplied by Dr. D. Sprinson of Columbia University) labeled as shown below, to determine the position of bond cleavage in the carboxytransphosphorylase reaction. The experiment was run essentially as described above for experiment 4, except that pyrophosphatase

Figure 5.

Use of ¹⁸O-enriched P-enolpyruvate to Show Possible Cleavage Points

was omitted from the reaction mixture. Enrichment of ^{18}O in the PP_i and lactate was determined as discussed above. There was no excess in the PP_i . The ^{18}O content of the acetate was 0.78 atom % excess. According to Dr. Sprinson, the bridge oxygen in the ^{18}O -P-enolpyruvate should have been enriched to around 11 atom %. The value we found in the acetate carboxyl is much below this value and suggests that perhaps the pyruvate was not immediately converted to acetate and the carbonyl oxygen had an opportunity to exchange out. The complete absence of ^{18}O in the PP_i suggests that the carboxytransphosphorylase reaction proceeds with oxygen-phosphorus bond cleavage and thereby eliminates mechanism 3. No conclusions can be made regarding mechanism 1 and 2 on the basis of this experiment, since they both predict oxygen-phosphorus bond cleavage.

Proposed Mechanism for the Carboxytransphosphorylase Reaction. On the basis of the data currently available and on new evidence concerning the stereochemistry of the reaction obtained by Drs. I. A. Rose, Cancer Institute, Philadelphia, Pennsylvania and J. M. Willard, University of Vermont, Burlington, Vermont, we feel that the mechanism of the carboxytransphosphorylase reaction proceeds as outlined in Figure 6. This mechanism involves a classical nucleophilic attack of phosphate on the P-enolpyruvate phosphoryl generating PP_i and an enolate species. The enolate is then carboxylated by nucleophilic attack of $^{14}\text{CO}_2$ to yield oxaloacetate. The PP_i remains enzyme bound during this process, thus satisfying the conditions of the oxaloacetate/ CO_2 exchange reaction. Under conditions of the pyruvate reaction, the enolate species is released from the enzyme surface and is protonated in solution. In this respect the enzyme resembles the 6-P-gluconic dehydrase studied by Meloche and Wood [J. Biol. Chem. 239, 3505 (1964)]. The latter aspect of the mechanism is required for several reasons. First, as previously noted, the pyruvate reaction is completely irreversible and no exchange between radioactive pyruvate and oxaloacetate or P-enolpyruvate was found. With Drs. Rose and Willard, it has been shown that the enzyme does not catalyze a loss of tritium from 3- ^3H -pyruvate, thus the enzyme does not catalyze a keto/enol tautomerization. The reason the pyruvate reaction is irreversible and pyruvate does not exchange into P-enolpyruvate or oxaloacetate is most likely due to the very low concentration of enol-pyruvate in solution (around 99% as the keto form). In the reverse of the pyruvate reaction, if the true substrate is enol-pyruvate, rather than keto-pyruvate under normal conditions the enol concentration would be far below the K_m ; therefore reversibility would not be observed. Second, with Drs. Rose and Willard, we have studied the stereochemistry of the pyruvate reaction. It has been found that unlike the carboxylation reaction, which is entirely stereospecific [I. A. Rose, et al. J. Biol. Chem. 244, 6130 (1969)] the protonation step is completely random. This observation is consistent with the hypothesis that the protonation occurs free in solution, rather than being enzyme directed.

It should be noted that all of the mechanisms previously discussed can lead to an enolate species. However, we feel that the proposed mechanism is more correct for several reasons. First, it does not involve a hydrolytic step and therefore predicts no incorporation of ^{18}O from water into products. This is consistent with the experimental findings reported here. Second, all of the previously discussed mechanisms require the energetically unfavorable expulsion of hydroxyl ion from the P-enolpyruvate phosphoryl upon nucleophilic attack of phosphate (Mechanism 1) or carboxylate (Mechanism 2). The proposed mechanism, on the other hand, involves the displacement of enolate from the phosphoryl, a highly favorable leaving group.

Figure 6

Enolate as Intermediate of the Carboxy-transphosphorylase Reaction (Mechanism IV)

One puzzling factor is that the enzyme does not catalyze the decarboxylation of oxaloacetate to pyruvate and CO_2 under conditions in which $^{14}\text{CO}_2$ is rapidly exchanged into oxaloacetate. The reaction mixture contained oxaloacetate, $^{14}\text{CO}_2$, PP_i , Mg^{+2} , mercaptoethanol and EDTA. The EDTA prevents phosphoryl transfer and thus catalysis of the overall reaction. If the enzyme-enolpyruvate is an intermediate of the exchange it apparently remains bound to the enzyme and does not dissociate to free enolpyruvate. It was partly for this reason that pyrophosphoenolpyruvate was originally proposed as an intermediate of the reaction.

Dr. Willard has recently examined the rate of the decarboxylation reaction under optimum conditions for formation of pyruvate from P-enolpyruvate, i.e. CO_2 is omitted as well as mercaptoethanol and, EDTA. The mixture contained enzyme, PP_i , oxaloacetate and Mg^{+2} . Under these conditions there was decarboxylation to pyruvate but at about 1/10 the rate of formation of pyruvate from P-enolpyruvate in the forward reaction. One explanation of the difference in rates may reside in the equilibria shown schematically in Fig. 7.

Figure 7

Equilibria Involved in Proposed Mechanism of Carboxytransphosphorylase Reaction

Thus, if the concentration of (PP_i -enzyme-enol-pyruvate) were small when approached from the oxaloacetate direction as compared with approach from P-enolpyruvate and P_i , then the release of pyruvate might be slower in the reverse reaction than in the forward reaction.

The failure of the enzyme to catalyze oxaloacetate decarboxylation with EDTA-treated enzyme in the presence of CO_2 and mercaptoethanol may result because EDTA treatment somehow interferes with the ability of the enolate to be released from the enzyme. A complete clarification of this problem will be attempted in the kinetic study of the enzyme, which we hope will allow us to determine the concentration of enolate enzyme.

II. SUBUNITS OF CARBOXYTRANSPHOSPHORYLASE--M. E. Haberland and H. G. Wood

The work on the subunit nature of carboxytransphosphorylase has now been completed and is being submitted for publication. A copy of the manuscript is appended, therefore our comments in this report will be confined to a brief summary of the work.

The native enzyme was found to have a molecular weight of 4×10^5 g/mole and a sedimentation coefficient of 15.2S. In the presence of substrates (P-enolpyruvate, P_i , and CO_2) a 10S species was formed, with CO_2 being required for the conversion. In addition on prolonged dialysis against dilute phosphate there was dissociation to a 7.1 species ($\sim 1 \times 10^5$ g/mole). In the presence of guanidine-HCl a species having a molecular weight of 0.9×10^5 g/mole was formed. On the basis of these molecular weight determinations and on the basis of electron micrographs and other data it was concluded that the subunit composition of the enzyme is as follows: the native 15S species exists as a tetramer, the 10S species is a dimer, and the 7S species is a single polypeptide chain monomer.

By means of techniques which allow the determination of the sedimentation coefficient of the catalytically active species, it was found that all three forms of the enzyme are active in the oxaloacetate reaction; only the monomeric and tetrameric species were shown to be active in the pyruvate reaction. It was not possible to determine the relative specific activities of the three species. A catalytically discrete subunit species of carboxytransphosphorylase may play an important role in the regulation of enzymatic activity, as will be discussed in this year's Proposal.

III. CATALYSIS AND MECHANISM OF PYRUVATE PHOSPHATE DIKINASE REACTION--Y. Milner and H. G. Wood

Mechanisms Proposed for the Dikinase Reaction--The mechanism of the reaction catalyzed by pyruvate, phosphate, dikinase (PPDK) was investigated by Evans and Wood [Proc. Nat. Acad. Sci. 61, 1448 (1968).] and by Andrews and Hatch [Biochem. J. 114, 117 (1969).] using labeled substrates and identifying partial reactions catalyzed by the enzyme. Evans and Wood on the basis of exchange studies, proposed the following mechanism, composed of three partial reactions:

The mechanism includes a pyrophosphoryl-enzyme intermediate (or diphosphoryl enzyme) in addition to the monophosphorylated form. The monophosphorylated form was demonstrated by Evans and Wood (1968) by gel filtration.

Andrews and Hatch proposed an alternative mechanism for plant PPK. They found that $\text{ATP} \rightleftharpoons \text{AMP}$ exchange occurred only in the presence of inorganic phosphate and proposed, therefore, only two partial reactions:

This mechanism involves an attack of ATP and P_i on an active group of the enzyme with the formation of enzyme-P and pyrophosphate without formation of a pyrophosphoryl enzyme complex.

It is interesting to compare here the P-enolpyruvate synthase from *Escherichia coli* described by Cooper and Kornberg [Biochem. Biophys. Acta 104, 620 (1965); 141, 211 (1967); Biochem. J. 105, 49C, (1967).] They have proposed that the enzyme catalyzes a very similar reaction sequence:

Water (hydrolytic step) substitutes for P_i (phosphorolytic step) in this reaction. With this enzyme, P_i was found to be required for the $\text{AMP} \rightleftharpoons \text{ATP}$ exchange (reaction 10) which can be explained on the assumption that the hydrolytic step (reaction 11) is very favorable and rapid hence a high level of P_i is needed to reverse reaction (11) and to maintain enzyme-PP at a substantial level for $\text{AMP} \rightleftharpoons \text{ATP}$ exchange. However, this enzyme was reinvestigated by Berman and Cohn [J. Biol. Chem. 245, 5309, 5319, (1970).] using ATP analogs (such as β - γ methylene ATP and β - γ amine ATP in which the oxygen bridge between β and γ phosphates is replaced by CH_2 - and NH_2 -groups respectively). They compared ratios of $\text{H}_2\text{O}^{18} \rightleftharpoons \text{P}_i$ exchange and $^{32}\text{P}_i \rightleftharpoons \text{ATP}$ exchange. They concluded that a free enzyme-PP complex is not formed and propose a mechanism in which the following reaction occurs.

In this mechanism the pyrophosphoryl group occurs attached to the enzyme but only as an enzyme complex with AMP also attached (never as the free E-PP).

The observation of Andrews and Hatch might be explained by a similar mechanism with a phosphorolytic cleavage rather than hydrolysis as illustrated in reaction (15).

The free pyrophospho-enzyme would not occur in this case and AMP exchange into ATP would only occur in the presence of P_i . Thus the difference between the pyruvate, phosphate dikinase of plants and that of bacteria might be that the AMP remains bound to the enzyme prior to the phosphorolytic cleavage and the free enzyme-pyrophosphate intermediate is not a component as it is with the enzyme from propionibacteria.

Another enzyme which has been considered to involve a pyrophosphoryl-enzyme is phosphorylribosyl pyrophosphate synthase [Harris and Koshland, *Science* **172**, 1253 (1971)]. Switzer found [*J. Biol. Chem.* **245**, 483 (1970)] an ordered sequential BiBi mechanism for the reaction and concluded that the pyrophosphoryl-enzyme form, if it does exist at all, is not significantly kinetically.

The findings regarding PPDK from plants, and phosphorylribosyl pyrophosphate synthase from *Salmonella typhimorium* made it necessary to reexamine the validity of the Evans-Wood mechanism and our report here will deal, primarily, with evidence by steady state kinetics, inhibition profiles, exchange reactions, and exchange kinetics.

Purification of Enzymes. Last year we reported purification of the enzyme from *Propionibacterium shermanii* and proposed to attempt purification from *Bacteriodes symbiosus* in which the enzyme is present in much higher levels than in the propionibacteria [Reeves, et al, *J. Biol. Chem.* **243**, 5486 (1968)]: We also undertook the purification of enzyme from propionibacteria because of the routine availability of these cells in our laboratory. Starting from about 300 grams of glycerol grown cells (wet packed mass) the previous procedure [Evans and Wood, *Biochem.* **10**, 721 (1971)] was modified somewhat for purification and produced enzyme (see summary in Table III) with a higher specific activity than observed previously. The preparation with specific activity of 0.5 (step 8, Table III) contained adenylate kinase and ATPase activities in minute but significant amounts. However, these contaminants were removed by further purification to specific activity of 0.8 (step 10). The elimination of these activities is of utmost importance to study the enzymatic mechanism using exchange reactions. For example, the presence of ATPase could furnish inorganic phosphate in the $\text{ATP} \rightleftharpoons \text{AMP}$ exchange experiment making it impossible to distinguish between the Evans-Wood and Andrews-Hatch mechanisms (reaction 4 vs reaction 8). The preparation of specific activity 0.8, showed one major band upon isoelectric focussing on acrylamide gels.

Table III

Summary of Purification of PPK from P. Shermanii
(300 g of wet packed cells were the starting material)

Stage of Purification	Volume (ml)	Total units ($\frac{\mu\text{moles}}{\text{min}}$)	Total protein (mg)	Specific activity (unit/mg)	Recovery (%)
3. 0.05M P_i pH = 6.8 eluant of Cell. P (see Evans-Wood Biochem. 1971) precipitated at 90% AmSO_4 and dissolved	200	492	49,200	0.01	100
4. 35-65% AmSO_4 Cut dissolved in buffer pH = 6.6	120	250.9	11,400	0.022	51
5. DEAE-Sephadex A-50* chromatograph (pooled fractions)	760	29.52	118.80	0.25	6
6. Sephadex G-200 pooled fractions	35	9.84	22.30	0.44	2
7. DEAE cellulose chromatograph (pooled fractions)	180	3.98	11.42	0.35	0.8
8. Sephadex G-200 II	11	2.25	4.50	0.50	0.46
9. DEAE cellulose (micro-granular) chromatograph	135	1.2	1.60	0.75	0.24
10. 64% saturation with AmSO_4 (dissolved precipitate)	0.6	0.7	0.78	0.85	0.14

Notes: From 7 onward the enzyme was assayed for adenylate kinase by determining production of ^{14}C -ADP from ^{14}C -ATP and ^{12}C -AMP and ATPase by production of ^{14}C -ADP from ^{14}C -ATP. Additional steps of purification were taken whenever those activities were found in significant amounts (above 1% of the PPK activity). Protein was determined by the micro-Biuret Method.

* - chromatography performed at room temperature, other stages at 4°C.

Bacteriodes symbiosus enzyme was purified starting from 1035 grams of wet packed cells grown by Prof. Richard Reeves, Department of Biochemistry, Louisiana State University, School of Medicine. The purification procedure is summarized in Table IV. All purification steps were carried out at 4°C (except step 6 which was done at room temperature) using 0.01M imidazole buffer pH 7.0 and 1 mM EDTA. Yields and specific activities are much higher than for the enzyme from propionibacteria. Stage 6 enzyme was essentially devoided of adenylate kinase and ATPase activities. Stage 7 enzyme gave one faint band upon isoelectric focussing electrophoresis and two major bands and three minor bands on gel-electrophoresis in sodium lauryl sulphate.

Properties of the enzymatic preparations. Table III demonstrates there is a very low yield of the enzyme from propionibacteria which is due to its inherent instability. We have attempted to determine the origin of this instability. The pH-activity and stability profiles were determined, and as a result the purification procedures were performed at pH 6.0-7.0 to minimize pH inactivation. Dicarboxylic acids and especially malate, were found to protect the enzyme against heat inactivation after a wide variety of compounds were tested. Hence Tris-malate buffer was used throughout most of the purification procedures. The enzyme was found to be stabilized against heat inactivation by AMP (Fig. 8, c and d), and both AMP and ATP stabilize the enzyme against proteolysis by trypsin (Fig. 8a). None of the substrates prevented inhibition by p-hydroxy mercuribenzoate except for a slight protection afforded by P-enolpyruvate (Fig. 8b). Although ATP protects against proteolysis it appears to destabilize the enzyme somewhat towards heat. These observations suggest that AMP and ATP interact specifically with the enzyme without the presence of other substrates and perhaps change the enzyme conformation. However, none of these compounds stabilized the enzyme during the purification procedures. Stability during storage was improved several fold compared to that reported by Evans and Wood, 1971. The enzyme solution in 0.01 M Tris-malate pH = 6.6 and 1 mM EDTA could be stored at 4°C without apparent loss of activity for at least 2 months. Furthermore the solution could be frozen and thawed several times without loss of activity.

The stability of the B. symbiosus during storage was inferior to the propionibacteria enzyme. The pure enzyme was stabilized by keeping it as a suspension in 65% saturated $(\text{NH}_4)_2\text{SO}_4$. Dialysis caused considerable inactivation.

Molecular weights of both enzymes seems to be around 150,000 g/mole as judged by gel-filtration profiles and sedimentation behavior in the analytical ultracentrifuge.

Steady State Kinetics of the PPDK from Propionibacteria. The mechanism proposed by Evans and Wood (see page 17) demands a ping pong mechanism as defined by Cleland ["The Enzymes" Vol. 2, 1, (1970): J. Biol. Chem. 241, 673 (1966)]. In such a mechanism, (see scheme below) one substrate (A) interacts with the enzyme to produce a modified enzyme (F) (with part of (A) remaining bound to it) and release of the first product (P). The second substrate (B) combines with the modified enzyme producing a third enzyme species (G) and release of the second product (Q).

Table IV

Summary of the Purification of PPDK from B. symbiosus
(Preparation from 1032 g of wet cells)

Stage of Purification	Volume (ml)	Total protein (mg)	Total units $\frac{\mu\text{moles}}{\text{min}}$	Specific activity $\frac{\text{units}}{\text{mg}}$	Recovery (%)
1. Lysosyme digest	18,000	110,000	19,100	0.17	100
2. 38%-65% AmSO ₄ sat. cuts (3times) repeatedly	5,000	20,000	13,100	0.65	68
3. DEAE cellulose chromatog (Fibrouse type)	8,200	2,950	8,010	2.70	42
4. 35%-65% AmSO ₄ sat. cut (precipitate dissolved)	12	2,400	6,000	2.50	31
5. Sephadex G-200 chromatog (pooled fractions)	45	1,000	5,020	5.00	26
6. DEAE cellulose (microgranular) chromatog. (pooled fractions)	350	310	2,800	9.00	14.7
7. Isoionic chromatograph on microgranular DEAE cell	300	117	1,400	12.00	7.3
8. 65% AmSO ₄ saturation precipitate suspended ⁴ in 65% AmSO ₄	20	41	500	12.20	2.6

Notes: Stage 2 presented the greatest difficulty owing to inability to precipitate the lysed cells and nucleic acids. Stage 7 was not really necessary since at stage 6 the enzyme was free essentially from adenylate kinase (ATPase was very low at the begining). Protein was estimated by 280 nm absorbtion assuming 1 O. D. = 1 mg protein.

Figure 8.

Stability Towards Heat, Proteolysis, and p-Hydroxymercuribenzoate in the Presence of Various Substrates

Proteolytic stability was determined as described below (c and d) but contained, in addition, vari

amounts of trypsin and chymotrypsin (equal amounts) as indicated in Figure 8a. Mixtures were incubated at 24°C for 15 minutes and the residual activity* determined immediately. Control: same ingredients as above omitting proteolytic enzymes.

b. pHMB stability was determined with the same ingredients as below with inclusion of pHMB and incubation for 15 minutes at 24°C with the concentrations of pHMB as indicated in Figure 8b. Residual activity* was measured as below. Control: tube containing same ingredients as above incubated at 24°C without pHMB.

c and d. Heat stability was determined in tubes contained in final volume of 0.2 ml: Tris-malate buffer pH 6.6, 20 μ moles; $MgCl_2$, 5 μ moles; enzyme (*P. shermanii* PPDK, specific activity=0.35) 0.02 enzymes units** and, when indicated; ATP, 1 μ mole or AMP, 1.2 μ moles; PEP, 1.0 μ moles; 3',5'-cyclic AMP, 0.85 μ moles. The mixtures were incubated at 49°C and at time intervals indicated the tube was withdrawn and placed in ice-bucket. Residual activity was measured in standard assay*. Activity was compared to controls containing the same ingredients but incubated in ice-bucket. Controls were compared to untreated enzyme.

*Standard assay. Enzymatic samples in 0.2 ml final volume. The following solutions were added to the samples: 0.1 ml of 0.03 M PP_i , 0.05 ml of 0.06M AMP and 0.3 ml starting mixture which contained: Tris-malate buffer pH = 6.6, 20 μ moles; $MgCl_2$, 6 μ moles; $(NH_4)_2SO_4$, 2 μ moles; P-enolpyruvate, 3 μ moles. The reaction was at 24°C and was terminated by addition of 0.3 ml of 5 mM dinitrophenylhydrazine in 2N HCl. Mixtures were incubated at 24°C for 15 minutes, then 0.8 ml of 5 N NaOH and 4.0 ml of water were added and the color read at 420 nm in a Klett-Summerson photocolormeter against suitable blanks and standards.

** The *B. symbiosus* enzyme was specific activity of 0.6 and 0.025 enzyme units were added.

The third substrate (C) combines with (G) generating the final product (R) and the original enzyme form (E). Using the shorthand notation of Cleland [Biochem. Biophys. Acta 67, 104 (1963)], we can describe the PDK reaction sequence in terms of a Tri (Uni Uni) ping pong mechanism.

The A→P sequence represents reaction (4) which can be measured as the $\text{ATP} \rightleftharpoons \text{AMP}$ exchange reaction which should be independent of other substrates. The sequence, B→Q, is reaction (5) which is dependent on the phosphorylated enzyme form (F) [hence dependent on addition of A (ATP) or R (PEP) to phosphorylate the enzyme] and can be measured as the $\text{P}_i \rightleftharpoons \text{PP}_i$ exchange reaction. The C→R sequence is reaction (6) represented by $\text{PEP} \rightleftharpoons \text{Pyr}$ exchange which should be independent of addition of other products and substrates. The general kinetic profile of such a mechanism with $1/v$ plots against $1/[A]$, $1/[B]$ or $1/[C]$ at various fixed second or third substrates concentrations should give parallel straight lines when the concentration of [A] is varied vs different concentrations of [B] or [C] or [B] vs [A] and [C] etc. This occurs because the release of products (P, Q and R) interrupts reversibility (since initial velocity is measured and thus the products are at a very low level) between [A] and [B] or [C]. Thus when velocity is measured by appearance of R the rate is dependent on the level of F (dependent on [A]) and on the level of G (dependent on [B]). Each [B] has its level of G hence no slope effect will be seen in reciprocal plots, but a series of parallel lines (for each [B]) intersecting according to the [B] level (when [A] vs [B] are plotted). The same holds true for other combination of substrates.

When the Andrews Hatch mechanism is considered with the same terminology and logic one obtains:

This is a BiBi UniUni ping pong mechanism in which: (1) no $A \rightleftharpoons P$ exchange ($AMP \rightleftharpoons ATP$) is possible without B and Q (P_i and PP_i) present, (2) a reciprocal plot of $1/v$ vs $1/A$ at different fixed levels of B will give intersecting lines, (3) $1/v$ vs $1/[B]$ at different fixed C levels will give parallel lines since B and C combination is interrupted by P, Q release (Q will not be released unless A and B are present in proper alignment).

The kinetic patterns obtained with the PPDK from propionibacteria agree completely with the Tri (Uni Uni) ping pong mechanism concept. Typical kinetic patterns are shown in Figure 9 for the forward reaction and in Figure 10 for the reverse reaction. The following K_m values were obtained from intercept replots: $K_{PP_i} = 0.071$ mM, $K_{AMP} = 0.035$ and 0.1 mM, $K_P = 1$ mM and $K_{Pyr} = 0.38$ and 0.25 mM in good agreement with the values published by Evans and Wood (1971). There are irregularities in the pattern notably in the presence of high concentration of PP_i where a tendency towards convergence of the lines was apparent. This probably arises from contamination of PP_i with a minute amount of P_i (the PP_i was purified on a QAE-Sephadex column but nevertheless hydrolysis does occur). When unpurified PP_i (8-10% contamination of P_i) was used the pattern was intersecting (except for the very low PP_i concentration) as seen in Figure 11. This phenomenon is in support of the general Tri (Uni Uni) ping pong mechanism since the presence of substrates tends to establish the reversibility for all other sequences and will induce a slope effect (Q is in a key position in the mechanism) in the ping pong mechanism.

Product Inhibition Profile. The steady state kinetics presented show that there are three partial reactions in the PPDK reaction which can be described as a Tri (Uni Uni) ping pong mechanism. There are, however, at least two types of ping pong mechanisms: 1) The classical type in which sites for binding of all substrates and products on the enzyme are all the same or partially overlapping as in the case of dinucleotide kinase (Graces, E., Cleland, W. W., Biochem. 8, 663, (1969)). 2) The non-classical type ping pong mechanism described in this laboratory [Northrup, D., Wood, H. G., J. Biol. Chem. 244, 5808 (1969)] in which there are separate independent sites for each substrate as in the case of methylmalonyl-CoA transcarboxylase [Northrop, D., J. Biol. Chem. 244, 5808 (1969)] or pyruvate carboxylase (Barden, R., et al in press). Based on the latter two enzymes the rules for non-classical ping pong mechanisms were established for product inhibition: 1) competitive patterns are observed only when inhibitor and substrate belong to the same partial reaction and 2) non-competitive and uncompetitive patterns are expected when inhibitor and substrates belong to different partial reactions. The inhibition profile for PPDK from Propionibacterium shermanii are summarized in Table V. The results obtained agree well with the non-classical type mechanism (except AMP vs P_i) and support the Evans-Wood proposal of a ping pong mechanism. The results also indicate that the enzyme contains three different sites for substrates/products pairs: AMP/ATP, P_i/PP_i and pyruvate/P-enolpyruvate. This also explains why substrate inhibition was not observed. The origin and type of inhibition is clearly understood if we consider the enzyme forms that are involved. Reaction with P-enolpyruvate will be inhibited by ATP because both compete for the "E" form of the enzyme at the phosphorylation site rather than at the PEP binding site and are therefore non-competitive. ATP will be inhibited competitively by AMP, because they are mutually excluded from the same binding site. In the same way P_i will inhibit the reaction with AMP (both

Figure 9.

Steady State Kinetics of the Forward Reaction (*P. shermanii*)

Reaction mixtures were made to the proper concentrations indicated in the figure and brought to 0.3 ml final volume with water. The reaction was started by adding 0.2 ml of solution containing: imidazol buffer pH = 7.0, 20 μmoles ; pyrophosphatase (Worthington), 1.6 units; ^{14}C -pyruvate, 0.0125 μmoles 1.2×10^7 dpm/ μmole and 0.2 μmoles ^{12}C -pyruvate; 1.2 μl enzyme (0.0026 enzyme units). Reaction was performed at 24°C for 60 minutes and terminated by adding 0.03 ml 2 N HCl. Samples of 5 μl from each tube were spotted on No. 3 Whatman paper, dried, carrier ^{12}C -P-enolpyruvate and ^{12}C -Pyr were spotted on each sample and the papers were developed for 6 hours using water:methanol mixture (20:80 v/v) as developer. Pyruvate and P-enolpyruvate spots were visualized by uv light, marked and cut to scintillation vials containing 10 ml of scintillation liquid (3.6 gr POP+0.4 gr POPOP in 1 liter toluene). Samples were counted for 4 minutes in the Unilux counter (efficiency of 66% when upper channel A-30, Lower-G-20). Amounts of P-enolpyruvate formed were calculated according to the percent distribution of counts between pyruvate and P-enolpyruvate.

Figure 10.

Steady State Kinetics of the Back Reaction

Reaction mixtures in final volume of 1.0 ml contained the variable substrates as indicated in the figures and 3 μ moles AMP. 1.0 ml of starting mixture was added to start the reaction. Starting mixture contained per ml; Tris malate pH=6.6, 20 μ moles; $(\text{NH}_4)_2\text{SO}_4$, 1.25 μ moles and 0.25 μ l enzyme (0.0002 enzyme units). Mg^{++} was present always as 2/1 ratio to PP_i . Reaction performed at 24°C for 60-70 minutes and terminated by adding 0.1 ml dinitrophenyl hydrazine reagent (5 mM in 2 N HCl) and 0.4 ml HCl 2N. The mixtures were allowed to stand 2 hours at 24°C and then read at 390 nm against suitable blanks and standards.

In this procedure 1 μ mole pyruvate = 8.0 O.D. at 390 nm.

Figure 11.

Steady State Kinetics of the Back Reaction

See Figure 10 for experimental details.

Table V

Summary of Inhibition Profiles Observed for
Pyruvate, Phosphate, Dikinase from *P. shermanii*

Inhibitor	Substrates	Type of inhibition in classical ping-pong	Type of inhibition in non-classical ping-pong	Type of inhibition found	Figure*
PEP	Pyruvate	NC	C	--	--
PEP	Pi	NC	NC	--	--
PEP	ATP	C	NC	NC	12, 16
PPi	Pi	NC	C	C	14
PPi	Pyruvate	C	NC	NC	--
PPi	ATP	NC	NC	NC	14
AMP	ATP	NC	C	C	13
AMP	Pyruvate	NC	NC	NC	--
AMP	Pi	C	NC	UC	--

*Not all inhibition experiments are presented in this report owing to lack of space.

C-competitive inhibition, NC-non-competitive inhibition, UC-uncompetitive inhibition

Figure 12.

Inhibition-Pattern in the PPDK Reaction (Referred to in Table V).

Reaction mixtures in final volumes of 0.3 ml, contained the indicated concentrations of various substrates. In addition, all mixtures contained; 1.2 mM AMP and 0.0003 enzyme units (propionibacteria). The reaction was started by adding starting mixture, 0.2 ml, containing, Tris-malate pH=6, 5-20 μmoles ; PP_i , 1.5 μmoles ; MgCl_2 , 3 μmoles . Reaction performed for 40 minutes at 24°C and terminated by adding 2.0 ml of 0.25 mM dinitrophenyl hydrazine in 1 N HCl.

Figure 13

Inhibition of the Back Reaction by ATP (Referred to in Table V)

In addition to the indicated substrate levels, PEP, 1 μmoles and starting mixture were as in Fig. 12. 0.0002 (propionibacteria) reactions were terminated after 60 minutes.

16
 react with E-PP from different sites) and pyruvate will be inhibited by PP_i (reaction with E-P). P_i will be inhibited competitively by PP_i and pyruvate by PEP because they interact with the same binding sites on the enzyme (for each pair). It is of interest to note that the enzyme is competitively inhibited with respect to PP_i (Figure 14) in the back reaction (and probably in the forward reaction also) by α - β -methylene-ATP (in which the bridge oxygen between the terminal phosphates is replaced by $-CH_2-$) but this inhibition is non-competitive with respect to AMP. Methylene pyrophosphate (P-C-P) is a competitive inhibitor with respect to PP_i (Figure 14). This is in contrast (next page) to the inhibition exerted by the β - γ -methylene ATP which is probably exerting its inhibition through the adenyl moiety, rather than through the P-C-P moiety which is responsible for the inhibition patterns discussed above. From this work the various K_i 's were calculated: $K_m(PP_i) = 0.12$ mM, $K_i(P-C-P) = 0.74$ mM, and $K_m(ATP) = 0.027$ mM, $K_i(\alpha\beta$ methylene ATP) = 0.02 mM, and $K_i(\beta\gamma$ methylene ATP) = 0.027 mM. These data show that some binding constants are determined probably by the intact adenyl moiety in the inhibitors.

In conclusion, the kinetic and inhibitor profiles shown here are in accord with the proposed Evans-Wood mechanism and represents Tri (Uni Uni) ping pong mechanism of a non-classical type in which three different substrate sites probably exist on the enzymes surface involving the binding of pyruvate/PEP, PP_i/P_i , ATP/AMP pairs.

Exchange Reactions Catalyzed by PPDK from *P. shermanii* and *B. symbiosus*.

We have verified with pure enzymes that the exchange reactions described by Evans and Wood (1969) occur with both enzymes. The study performed also included a detailed kinetic analysis of the exchange reactions.

a. The AMP $\rightleftharpoons$ ATP Exchange Reaction. This exchange is of crucial importance to distinguish between the Andrews-Hatch and Evans-Wood mechanism. As shown in Tables VI and VII AMP $\rightleftharpoons$ ATP exchange occurs at the same rate whether or not exogeneous phosphate is added. If phosphate were required, one would expect an increase in the rate to occur with exogenously added phosphate. Trapping of trace amounts of phosphate which might arise from contamination of AMP or ATP using a starch/phosphorylase a system, did not abolish the exchange. Furthermore, in the Andrews-Hatch mechanism, pyrophosphate is the product of the first partial reaction (reactions 8, 15) and therefore is necessary for the exchange just as P_i is essential. However, the presence of inorganic pyrophosphatase in the exchange mixture did not abolish the exchange. The divalent metal (Mg^{+2}) requirement was found to be satisfied by Co^{+2} or Ni^{+2} , although they were less effective. P-enolpyruvate inhibits the exchange non-competitively as expected. In general agreement with the ping pong type mechanism discussed above, a plot of AMP concentrations vs ATP gives parallel lines on reciprocal plots (Figure 15) up to the point of mutual competitive inhibition (arrow, Figure 15).

The AMP $\rightleftharpoons$ ATP exchange was also conducted in the presence of all substrates at equilibrium concentrations and pairs of reactants were varied at a constant ratio ($P_i/PP_i = 33.5$, Pyr/PEP = 24.2 and ATP/AMP = 1.41). It is observed (Figure 16) that there was increased exchange rates with increase in the concentration of the ATP/AMP pair. This is to be expected since the enzyme is held as E and E-PP, the latter is required for the exchange. There was strong inhibition by increasing concentrations of the Pyr/PEP pair. This inhibition is predicted since the pyruvate and PEP would convert the enzyme to the E and E-P and do not generate E-PP required for the ATP $\rightleftharpoons$ AMP exchange. Further experiments are needed to determine if higher concentrations of P_i/PP_i (or P_i) is indeed inhibiting the AMP $\rightleftharpoons$ ATP exchange since curve C does not show it clearly. Actually one should expect inhibition by varying P_i alone.

Figure 14.

Inhibition Pattern of P_i , PP_i and ATP Analogues

Mixtures in 0.3 ml final volume contained the indicated substrates and inhibitors. Mg^{++} was present as 2:1 ratio of PP_i to ATP. The mixture also contained 1.1 μ moles AMP and 1.0 μ moles PEP and .0001 enzyme units (propionibacteria). The reaction was started and terminated as described in Figure 12. The pyruvate formed was determined in all mixtures by incubation for 90 minutes at 24°C with the dinitrophenylhydrazine and reading at 390 nm. 1 μ moles pyruvate = 10.5 O.D. at 390 nm.

Table VI

AMP—ATP exchange catalyzed by PPDK from propionibacteria

System	Time (Min.)	Counts per 10 min. in ATP	Counts per 10 min. in ADP	Counts per 10 min. in AMP	*μmoles AMP incorporated to ATP
1. AMP + ATP + Mg	20	720	655	20,099	0.700
"	60	1,524	682	17,841	3.100
2. AMP + ATP + Mg	20	667	637	17,017	0.900
+ phosphorylase + starch	60	1,293	891	17,521	2.820
3. Complete + pyrophosphatase 1.6 units	20	813	614	22,625	0.480
	60	1,473	793	36,292	1.380
4. Complete + P _i 1.0 mM	20	791	---	19,739	0.840
	60	1,155	---	21,388	1.860
5. AMP + ATP + Ni ⁺² 0.5 mM	20	716	---	19,874	0.600
	60	1,381	---	21,081	2.400
6. AMP + ATP + Co ⁺² 0.5 mM	60	1,229	---	26,596	1.260
	60				
7. Complete + 2.5 mM PEP	20	840	---	26,076	0.360
	30	868	---	26,310	0.420
8. AMP + β-γ CH ₂ ATP + Mg	20	1,151	---	11,450	0.552
	60	1,668	---	17,740	1.440
9. Complete No Enzyme	20	256	604	25,788	0
	60	242	633	26,077	0

* All values reported are corrected by zero time subtraction.

Experimental Procedure:

System for the exchange contained in the volume of 0.015 ml, ATP, 0.005 μmoles, ¹⁴C-AMP, 0.06 μmoles (2.5 μmoles - 26,000 cpm gCl₂, 0.01 μmoles; or where indicated: NaP_i, pH 6.8, 0.02 μmoles; inorganic pyrophosphatase (W. ngton).

Table VI (cont.)

1.6 units; pyrophosphorylase a, 0.15 units + 2 mg soluble starch; PEP, 0.05 μ moles; $\text{Co}(\text{NO}_3)_2$ or $\text{Ni}(\text{NO}_3)_2$, 0.01 μ moles each when indicated; β - γ methylene ATP, 0.02 μ moles². The reaction was started by addition of 0.005 ml solution containing: Tris malate buffer pH = 6.6, 3 μ moles; $(\text{NH}_4)_2\text{SO}_4$, 0.125 μ moles, and .0008 enzyme units of propionibacteria or 0.0011 units of enzyme of *B. symbiosus* (Table IV). Reaction was run at 24°C and at various time intervals, samples of 2-5 μ l were spotted on Whatman No. 3 chromatography paper and dried immediately with hot air. After application of carriers AMP and ATP on the same spots to be separated, the papers were chromatographed for 20 hours in descending technique using ethanol (95%): ammonium acetate (1.0M, pH 7.5) (7:3 v/v) as developer. The papers were dried, UV spots identified and the AMP and ATP (occasionally the ADP spots too) were cut out and counted. Scintillation fluid was 3.6 gm POP + 0.4 gm POPOP in 1 liter toluene. Counting was done in the Unilux counter for 10 minutes. The efficiency was 64-66%. Since the amount of solution spotted was not measured accurately the total counts were used to normalize all counts to the same basis.

Table VII

AMP $\rightleftharpoons$ ATP exchange catalyzed by PPDK from B. symbiosus

System	Time min.	Counts per 2 min. in ATP	Counts per 2 min. in ADP	Counts per 2 min. in AMP	μ moles AMP incorporated into ATP
1. AMP + ATP + Mg + 4	60	711	317	30,964	4.50
	90	2,126	562	56,385	8.10
2. Complete + Phosphory- lase a + starch	60	868	309	24,196	7.40
	90	2,382	417	50,701	10.50
3. Complete + Pyrophos- phatase	60	668	307	23,534	5.40
	90	2,208	512	45,182	10.80
4. Complete + P _i	15	636	---	40,643	1.80
	90	1,650	---	35,864	10.20
5. AMP + ADP + Mg	15	415	685	59,780	0.60*
	90	407	542	59,859	0*
6. AMP + ATP + Co ⁺²	90	661	---	60,884	0.21
7. AMP + ATP + Ni ⁺²	90	304	---	21,701	5.20
8. AMP + ATP + Mg ⁺²	60	238		32,702	0
No Enzyme	90	254		26,809	0

See Table VI for experimental procedure. All values have been corrected by subtracting counts of AMP with no enzyme system. System contained ¹⁴C-AMP 15 μ c/ μ mole.

* - μ moles AMP incorporated to ADP.

Figure 15.

AMP, ATP Exchange at Equilibrium (P. Shermanii).

Experimental details are as described in the legend of Table VI (0.0003 enzyme units present).

Figure 16

AMP $\rightleftharpoons$ ATP Exchange Rates at Equilibrium

Reactions, mixtures and determination of the exchange rates were similar to those reported in Table V. Reactions were conducted in the presence of all substrates and products at equilibrium concentrations (taken from Reeves *et al.* J. Biol. Chem. 243, 5486 (1968). Tracer [^{14}C] AMP was added after 15 min preincubation to the equilibrium mixture including enzyme. Substrates and products pairs, at constant ratio, were varied: Line A - Pyruvate/PEP = 24.2 at increasing concentrations (indicated as $1/(\text{Pyr})$ on abscissa and $1/v$ as A ordinate). Line B - ATP/AMP = 1.41 at increasing concentrations as $1/\text{AMP}$ (B) abscissa and rates as B on ordinate. Line C - $\text{P}_i/\text{PP}_i = 33.5$ (indicated as $1/\text{P}_i$ on C abscissa and rates as C on ordinate. The non-varied pairs were kept constant at concentrations given in the points second from the left of the Figure.

Maximal velocities of $\text{AMP} \rightleftharpoons \text{ATP}$ exchange were recorded as: 0.105 $\mu\text{moles}/\text{min}/\text{Enz. unit}$ in equilibrium exchange, and 0.095 $\mu\text{moles}/\text{min}/\text{Enz. unit}$ in the regular Lineweaver Burk plot (Figure 15). The $\text{AMP} \rightleftharpoons \text{ATP}$ exchange observed is therefore additional evidence supporting the proposed Evans-Wood mechanism.

b. The Pyruvate-P-enolpyruvate Exchange. The enzyme from propionibacteria catalyzes a rapid $\text{Pyr} \rightleftharpoons \text{PEP}$ exchange without a requirement for other substrates (Table VIII) and, as expected from the inhibition profile presented, PP_i inhibits this exchange driving the E-P formed to E-PP thus reducing the available enzyme (non-competitive inhibition). Likewise ATP competes with P-enolpyruvate for the E form of the enzyme. Of special interest, is the observation that Mg^{+2} is needed for the exchange and Co^{+2} and Ni^{+2} could substitute in a lower efficiency. The most interesting finding is that β - γ -methylene-ATP inhibits the PEP-Pyr exchange. This suggests that the ATP analogue interacts with the enzyme to form E-P- CH_2 -P (analogous to ATP) which is unhydrolysable (P-C-P is inhibitor and not substrate!). Indeed ATP inhibition is much less than the β - γ analog. The reason, probably, is that the E-PP can be converted to E-P by traces of phosphate present while E-P-C-P cannot. The V_m of Pyr-PEP exchange is 5-10 $\mu\text{moles}/\text{min}/\text{Enz. unit}$ which is higher than the overall rate of the reaction catalyzed by the enzyme and varies according to the P-enolpyruvate (Figure 17 experimental details see Table VIII). The rate of exchange is linear for a certain time but drops later (Figure 17). It is linear with enzyme concentration (Figure 17). We have examined the catalysis of $\text{Pyr} \rightleftharpoons \text{PEP}$ exchange using the purified enzyme from *B. symbiosus* (Table IX). This enzyme also catalyzes the exchange in the absence of ATP, P_i , PP_i , AMP provided that Mg^{+2} is present, thereby verifying the general form of reaction (4) in the Evans-Wood mechanism. In this case the inhibition by PP_i and ATP is not apparent, however, β - γ -methylene-ATP does inhibit. The reason why ATP does not inhibit is not yet apparent. The V_m for exchange in this case is also high -- 2.5 $\mu\text{moles}/\text{min}/\text{Enz. unit}$, however, this might be a minimum value since all the exchange systems were in 0.15 M $(\text{NH}_4)_2\text{SO}_4$ final concentration since the enzyme was suspended in 65% sat. $(\text{NH}_4)_2\text{SO}_4$. Under optimum conditions it might approach that of the enzyme of the propionibacteria.

c. The Phosphate-Pyrophosphate Exchange Reaction. This partial reaction, in agreement with the Evans-Wood mechanism, was catalyzed by both purified enzymes (Tables X, XI). The exchange was facilitated by incubation with ATP or P-enolpyruvate (ATP is seemingly more efficient) as demanded by the sequence of partial reactions (4, 5, 6). The exchange is inhibited by AMP (dephosphorylation of the enzyme) and pyruvate as predicted. Ni^{+2} and Co^{+2} could substitute with lower efficiency for Mg^{+2} . The kinetics of this reaction were further investigated using the calculation of Yagil and Huberman [Biochem. 8, 352 (1969)], for equilibrium velocity which was found to be 0.078 $\mu\text{moles}/\text{min}/\text{Enz. unit}$ for the enzyme from propionibacteria. It was some 20 fold less with the *Bacteriodes* enzyme (again probably because of the high $(\text{NH}_4)_2\text{SO}_4$ concentration). The ping pong mechanism is manifested by the dependence of the rate of exchange of P_i on the PP_i concentration [(Figure 18) parallel lines until competitive inhibition between the pair is observed]. A non-competitive inhibition by AMP and bromopyruvate of $\text{P}_i \rightleftharpoons \text{PP}_i$ exchange was also found and is, again, in agreement to the general mechanism proposed; as is the inhibition by pyruvate (non-competitive) and methylene pyrophosphate (competitive with respect PP_i). Exchange rates measured at equilibrium concentrations of all substrates and products

Table VIII

Pyruvate $\rightleftharpoons$ PEP exchange with the enzyme from propionibacteria

System	Time min.	CPM in PEP	CPM in pyruvate	μ moles pyruvate exchanged to PEP
1. Pyruvate + PEP + Mg^{+2}		1,001	6,512	0
	5	1,841	6,936	10.8
	10	3,046	6,298	24.5
	20	4,495	5,000	41.1
	30	6,004	4,622	51.9
2. Complete + 1.5 mM PP _i	20	4,708	12,134	23.4
	30	7,049	13,631	30.7
3. Complete + 2.5 mM ATP	20	3,542	5,164	33.8
	30	4,167	4,549	41.7
4. Py + PEP + Co^{+2} (0.5 mM)	20	1,366	3,909	11.6
	30	3,475	7,712	25.3
5. Py + PEP + Ni^{+2} (0.5 mM)	20	2,370	7,673	15.1
	30	3,387	7,653	23.7
6. Py + PEP No Metal	20	1,132	9,453	2.3
	30	1,844	9,191	9.3
7. Py + PEP + 2.5 mM β γ methylene ATP	20	1,965	9,763	11.4
	30	3,099	11,048	16.6

Experimental details: Reaction mixtures in total volume of 0.02 ml contained Tris malate buffer pH 6.6, 3 μ moles; $(NH_4)_2SO_4$, 0.01 μ moles; pyruvate, 0.1 μ moles ^{12}C + 0.0025 μ moles ^{14}C -pyruvate, 5×10^5 cpm; P-enolpyruvate .05 μ moles; $MgCl_2$ (except where noted), 0.02 μ moles; enzyme (SA=0.6), 0.00065 units. At various time intervals 2-4 μ l samples were withdrawn and dried by hot air on Whatman No. 3 paper. Chromatography was for 6 hours in methanol:water (80:20) after applying drops of 1.0 M pyr and PEP on the same spots as the samples to be separated. PEP and Pyr spots were visualized by UV light, marked and cut out. The papers were counted in scintillation liquid (3.6 gm POP + 0.4 gm POPOP in 1 liter toluene) in the Unilux counter with an efficiency of 64-66%. Since the samples were taken at variable volumes there sometimes were different counts in different spots. Pyruvate incorporated into PEP was calculated from % distribution of counts in the spots.

Figure 17.

The Pyruvate $\rightleftharpoons$ P-enolpyruvate Exchange

Kinetics of the phenolpyruvate $\rightleftharpoons$ pyruvate exchange reaction (*P. shermanii*).
Experimental details similar to that in table VIII.

Table IX

The Pyruvate $\rightleftharpoons$ PEP exchange catalyzed by B. symbiosus PPDK

System	Time min.	CPM in PEP	CPM in pyruvate	μ moles pyruvate incorporated into PEP
1. Pyr + PEP + Mg ⁺²	0	1,198	6,918	0
	20	1,629	6,607	4.70
	60	2,345	6,236	14.00
2. Pyr + PEP + Mg ⁺² +1.5 mM PP _i	20	1,304	6,153	7.00
	60	1,892	5,788	16.60
3. Pyr + PEP + Mg ⁺² +2.5 mM ATP	20	2,072	10,301	9.40
4. Pyr + PEP + Co ⁺² (0.5 mM)	20	1,385	9,257	4.00
	60	1,772	10,142	7.20
5. Pyr + PEP + Ni ⁺² (0.5 mM)	60	990	7,099	4.75
6. Pyr + PEP + Mg ⁺² +2.5 mM β -methy- lene ATP	20	1,200	6,591	2.20
	60	1,629	7,083	7.10
7. Pyr + PEP + Mg ⁺² No Enzyme	20	1,091	9,561	0
	60	1,072	11,921	0

Experimental details as in Table VIII except 0.0002 enzyme units were used.

Table X

 $P_i \rightleftharpoons PP_i$ exchange catalyzed by PPDK from propionibacteria

System	Time (min)	CPM in P_i	CPM in PP_i	mμmoles P_i incorporated in PP_i
1. $P_i + PP_i + Mg^{+2}$ + <u>ATP 2.5 mM</u>	10 90	508,677 255,586	4,242 19,443	0.60 6.80
2. $P_i + PP_i + Mg^{+2}$ + <u>PEP 2.0 mM</u>	10 60	298,606 440,345	3,031 10,755	0.80 2.10
3. $P_i + PP_i + Mg^{+2}$	10 60	409,343 473,850	904 1,839	0.02 0.20
4. $P_i + PP_i + Mg^{+2}$ + <u>2.5 mM ATP + 4 mM</u> pyruvate	10 90	259,248 340,043	443 1,477	0.00 0.20
5. $P_i + PP_i + Mg^{+2}$ + ATP + <u>2.4 mM AMP</u>	10 90	357,225 544,590	1,120 1,374	0.10 0.10
6. $P_i + PP_i + Mg^{+2}$ + ATP <u>No enzyme</u>	10 90	421,227 347,781	717 1,072	0.00 0.10
7. $P_i + PP_i + ATP$ + Co^{+2} <u>2.5 mM</u>	60	551,801	4,713	0.60
8. $P_i + PP_i + ATP$ + Ni^{+2} <u>2.5 mM</u>	60	421,811	2,081	0.20

Exchange mixture contained in 0.015 ml final volume: $NaPP_i$, pH 6.6, 90 mμmoles; NaP_i , pH 6.7, 100 mμmoles; $MgCl_2$, 200 mμmoles; ATP (when included), 50 mμmoles; PEP (if used), 50 mμmoles; Pyr (if used), 100 mμmoles; AMP (if used), 60 mμmoles; $Ni(NO_3)_2$ or $Co(NO_3)_2$, 100 μmoles; $^{32}P_i$ 5.5×10^6 cpm (0.002 μmoles). To start the reaction starting mixture was added which contained: Tris-malate pH 6.6, 4 μmoles; $(NH_4)_2 SO_4$ 0.2 μmoles enzyme from propionibacteria, 0.5 μl, 0.00065 enzyme units or from *Bacteroides symbiosus* 5 μl, 0.001 enzyme units suspension in 65% saturation $(NH_4)_2 SO_4$. At zero time and other times 1-2 μl samples were transferred to Whatman No. 3 paper and dried immediately with hot air.

Developing and counting: The paper was developed, after applying carrier P_i and PP_i on the same spots, in Isopropanol: TCA 20%: NH_4OH 27% (75:25:25 (v/v)) for 10 hours. The P_i and PP_i spots were visualized by using the Hanes-Isherwood spray (Nature, 764 1107 (1969)).

P_i and PP_i spots were cut out and counted on paper in the same way as in Table VI. Efficiency of reading is approximately 90%-95%.

Table XI

$P_i \rightleftharpoons PP_i$ exchange catalyzed by PPDK from B. symbiosus

System	Time (min)	CPM in P_i	CPM in PP_i	μmoles P_i incorporated into PP_i
1. complete + ATP (2.0 mM)	10 60	429,261 322,209	700 1,894	0.140 0.587
2. complete + PEP (2.0 mM)	10 90	257,283 214,578	523 979	0.207 0.587
3. complete + ATP + Pyr (4.0 mM)	20 60	395,045 406,890	782 1,214	0.192 0.292
4. complete + ATP + AMP (2.4 mM)	10 60	409,445 298,983	432 610	0.105 0.204
5. complete <u>No ATP or PEP</u>	10 90	387,685 308,562	447 525	0.100 0.100
6. complete + ATP No enzyme	10 90	551,172 471,557	504 377	<0.05 <0.05

Experimental details as in Table X.

Figure 18.

Steady State Kinetics of $P_i \rightleftharpoons PP_i$ Exchange

$P_i \rightleftharpoons PP_i$ exchange at equilibrium. Experimental details are similar to Table XII.

(Figure 19) are again in general agreement with the overall mechanism discussed; the rate of ^{32}P incorporation to PP_i increases with P_i/PP_i or P_i/AMP concentration. The rate of exchange is decreased when the pyruvate is increased and more when both pyruvate and AMP are increased. The V_m for exchange was determined at equilibrium to be $0.155 \mu\text{moles/min/Enz. unit}$ and $0.115 \mu\text{moles/min/Enz. unit}$ from steady state exchange kinetics.

Publications

1. Evans, Herbert J., Wood, Harland G.
1971 Purification and Properties of Pyruvate
Phosphate Dikinase from Propionic Acid Bacteria
Biochemistry, 10, 721-729.
2. Schulman, Marvin and Wood, H. G.
1971 Determination and Degradation of Microquantities
of Acetate¹
Anal. Biochem., 39, 505-520.
3. Ghambeer, Rajinder K., Wood, H. G., Schulman, M. and Ljungdahl, L.
1971 Total Synthesis of Acetate from CO_2
III. Inhibition by Alkylhalides of the Synthesis from CO_2 ,
Methyltetrahydrofolate, and Methyl- B_{12} by Clostridium
thermoaceticum
Arch. of Biochem. and Biophysics, 143, 417-484.
4. Cooper, T. G. and Wood, H. G.
1971 The Carboxylation of Phosphoenolpyruvate and Pyruvate
II. The Active Species of " CO_2 " Utilized by
Phosphoenolpyruvate Carboxylase and Pyruvate Carboxylase
J. Biol. Chem., 246, 5488-5490.

Figure 19.

$P_i \rightleftharpoons PP_i$ Exchange Rates at Equilibrium

Figure 19 $\text{Pi} \rightleftharpoons \text{PPi}$ Exchange Rates at Equilibrium

Reaction mixtures and determination were done similar to the experiments in Table X. The exchange was conducted at equilibrium concentrations of all products and substrates (see Figure 16). Tracer ^{32}P was added after 15 min. preincubation at 24°C to the mixtures including enzyme. The varied pairs of substrate/product ratios and the non varied pairs are summarized for each line of the figure in the table below:

Line	Varied pair	First constant pair		Second constant pair	
	Ratio	Ratio	mM	Ratio	mM
A	$\frac{\text{Pyr}}{\text{AMP}} = 1.2$	$\frac{\text{ATP}}{\text{PEP}} = 0.25$	$\text{ATP} = 0.25$	$\frac{\text{Pi}}{\text{PPi}} = 33.5$	$\text{Pi} = 6.0$
B	$\frac{\text{Pi}}{\text{PPi}} = 33.5$	$\frac{\text{Pyr}}{\text{PEP}} = 24.2$	$\text{Pyr} = 2.25$	$\frac{\text{ATP}}{\text{AMP}} = 1.41$	$\text{ATP} = 0.25$
	$\frac{\text{Pi}}{\text{AMP}} = 32$	$\frac{\text{ATP}}{\text{PPi}} = 1.34$	$\text{ATP} = 0.25$	$\frac{\text{Py}}{\text{PEP}} = 24.2$	$\text{Pyr} = 2.25$
C	$\frac{\text{ATP}}{\text{PEP}} = 0.25$	$\frac{\text{Py}}{\text{PPi}} = 12$	$\text{Pyr} = 2.25$	$\frac{\text{Pi}}{\text{AMP}} = 32$	$\text{Pi} = 6$
D	$\frac{\text{ATP}}{\text{PPi}} = 1.34$	$\frac{\text{Pi}}{\text{AMP}} = 32$	$\text{Pi} = 6$	$\frac{\text{Pyr}}{\text{PEP}} = 24.2$	$\text{Pyr} = 2.25$
E	$\frac{\text{Pyr}}{\text{PEP}} = 24.2$	$\frac{\text{Pi}}{\text{PPi}} = 33.5$	$\text{Pi} = 6$	$\frac{\text{ATP}}{\text{AMP}} = 1.41$	$\text{ATP} = 0.25$

Abscissa shows $1/\text{S}$ of one of the substrates in the varied pair as indicated by letters. Rates of exchange are given on the ordinate, again indicated by letters.