MASTER

C60-2089-1

FINAL REPORT

AEC CONTRACT AT(11-1)-2089

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

This contract was made to enable us to conduct a spark-chamber experiment on $K_{\mu 3}$ decay at the Princeton-Pennsylvania accelerator, and to analyze the resulting film by automatic scanning techniques at Washington University.

Participating in this experiment were the following faculty members:

John Dobbs
John Scandrett
Malcolm Whatley
Richard Zacher

Two graduate students were involved, who are writing their Ph.D. dissertations on this experiment:

Jerry Nurenberg
Ernst Zinner

The following persons also assisted in the initial phases of the experiment:

Fred Geisler
Charles Lady
Peter Phillips
Don Ragan
Ron Rider

The equipment was built in less than a year at Washington University, and taken to Princeton in the spring of 1969. The experiment was run in the summer of that year, yielding 176,000 pictures of stopping K^+ undergoing various modes of decay. $K_{\mu 3}$ events often resulted in pictures containing a

muon track and two converting gamma rays. 15,000 of the pictures are intentional background pictures, obtained by mistiming the logic. 30,000 are $K_{\mu 2}$ pictures, which are helpful in determining the efficiency of the chambers. The rest are $K_{\mu 2}$, K_{e3} , $K_{\mu 3}$ or accidental background. About 30% of the pictures with a $K_{\mu 3}$ signature were taken with a 1" Platinum target in the external proton beam, the rest with a 3" target. The background was severe in the 3" target film, and it has proved impracticable to scan it. About 60% of the 1" target film has been scanned; the rest probably will not, because of financial and manpower limitations.

The film has been scanned and measured automatically using a computer-controlled C.R.T. scanning device designed and built by us. For this device a complete system of utility programs has been written. These automatic scanning and measuring programs yield about 95% of the events that would be found by hand scanners and take an average of 5.6 seconds of CPU time per event. Measuring precision is quite satisfactory for this experiment.

To be acceptable for complete scanning and measuring an event must have just one charged track in the chamber which satisfies the charged-track logic, and just one shower in each of the other two chambers. Missing sparks are allowed in at most 2 gaps for the charged-particle track. A shower is defined as at least 2 sparks, the first two of which have to be close to one another in 1 to 3 consecutive gaps, the rest inside a defined cone after the first spark (conversion point).

Frames violating these scanning criteria are abandoned, and the type of violation recorded. The scanning and measuring program records the position of the beginning,

part of the solution be zero. Most $K_{\mu 3}$ events give two real solutions; we assign these solutions weights by computing a probability for each solution using gamma-ray shower information.

At present we compute this probability simply from the number of sparks in the shower. Figure 1 shows the average number of sparks in a shower as a function of gamma-ray energy. This graph averages over varying conversion points of the gamma ray and therefore over a varying possible length of the shower until it leaves the end of the chamber as well as over different angles of incidence. A more refined method takes these variables for an individual shower into account by computing first a number of expected sparks, n_{exp} , from the given energy, conversion point and angle of incidence, and then finding the probability of getting the observed number of sparks given this value of n_{exp} .

Figure 2 shows a typical probability distribution for (n_{obs}/n_{exp}) . As indicated by the error bars in Fig. 1, such distributions correspond to an energy resolution of from 32% for 200 MeV gamma rays to 70% for 50 MeV.

Figure 3 is a histogram of the probability of one of the two solutions for a sample of $K_{\mu 3}$ events. This probability is normalized so that $p(\text{solution 1}) + p(\text{solution 2}) = 1$. This method will allow us to find the right solution in about 75-80% of the cases. In many additional cases the π^0 energy difference between the two solutions will be small, so that a wrong choice will introduce only a small error.

We have studied our accidental background by photographing 15,000 events with the logic set up so that one of the particles of the event is known to have arrived 33 nsec before the rest. The pictures taken were of all possible combinations of mistimings. Extensive studies using just the counters were made to measure the relative rates of these combinations of accidental events. The background photographs have been studied both by hand scanning and automatic measuring followed by kinematic fitting.

A first analysis of the film has now been made, and variations of it are being considered. The analysis is capable of yielding two parameters, ξ and Λ_+ . If we are conservative and fit for only one, we hold Λ_+ fixed at the conventional value of +.03, and find a value of ξ around -1. This is in agreement with some other experiments. However, if we fit for both parameters at once we get $\Lambda_+ \approx -0.15$ and $\xi \approx -3$. We are now trying to determine whether any systematic errors remain in the data, and whether the statistical errors have been correctly estimated. A slight problem also remains with the Monte Carlo calculation. It predicts that of the candidates for $K_{\mu 3}$ production, 23% should actually be $K_{e 3}$. Hand scanning indicates that a better number is around 7%. We are searching for the cause of this discrepancy.

Two Ph.D. theses will be presented in June 1972. One, by Ernst Zinner, will be concerned with the determination of the parameters of $K_{\mu 3}$ decay. The other, by Jerry Nurenberg, will concentrate on the problem of using electron showers to determine gamma ray energies. Both students are being supported on a science development grant from NSF during the first six months of 1972.

We are grateful to the AEC for their support of this experiment, which we believe will contribute to the information available on $K_{\mu 3}$ decays. We regret that it has not been possible to continue this work, either through an AEC contract or by means of a grant from NSF. In these difficult circumstances, we have no immediate plans to do more work in this area.

Fig. 1 Average number of sparks in shower,
for present chambers.

Fig. 2 Probability p for getting various numbers of sparks in a shower when the most probable number is seven.

Fig. 3 Histogram of number of events against the probability that solution number 1 is the correct choice.