COO-1780-4

RECEIVED BY DTIC DEC 9 1969

1. U. S. Atomic Energy Commission

Division of Research

Washington, D. C.

MASTER

PROGRESS REPORT

CONTRACT NO. AT-(11-1)-1780

DOCUMENT NUMBER COO-1780-4

entitled

"STABLE ISOTOPE TRACER STUDIES"

1 March 1969 to 28 February 1970

Submitted by

Gilbert Gordon

Department of Chemistry

University of Iowa

Iowa City, Iowa 52240

21 November 1969

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

leg

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

I. Personnel

During the period from the inception of the research contract on 1 March 1969 through the period 28 February 1970, the following persons will have contributed to the program:

Research Assistants:

Robert Buchacek, 12 months, half-time

Eugene Grimley, 12 months, half-time

Robert J. Seyse, 12 months, quarter-time

Robert Silverman, 6 months, quarter-time

Research Associates:

Dr. Donald Hoppenjans, 12 months, full-time

Research Supervisor:

Gilbert Gordon, 12 months, 1/5-time

II. Research Accomplished and in Progress

A variety of projects have been continued during the past year and several manuscripts are in various stages of preparation to be submitted for publication early in 1970. Several projects are close to completion and titles of three of the manuscripts in preparation are given below:

- (1) "The Chromium(II) Assisted Aquation of Halotetraamminechromium(III) Complexes in Aqueous Solution" by Donald Hoppenjans and Gilbert Gordon
- (2) "The Trans Effect in Some Iodoaquotetraamminechromium(III) Complexes" by Donald Hoppenjans and Gilbert Gordon
- (3) "The Kinetics of the Reactions between Oxovanadium(IV) and Chlorate and Chlorite Ions and Chlorine Dioxide" by William Melvin and Gilbert Gordon

The first two manuscripts are a result of our attempts to unravel the com-

plexities of the effect of "bridging ligands" on the chromium(II)-uranyl ion reaction. Oxygen-18 tracer experiments¹ indicate very efficient transfer of oxygen from uranyl ion to the chromium(III) product. The stoichiometry of the reaction

is relatively unaffected by the addition of anions such as fluoride, chloride, bromide, and azide and the principle product of the reaction is $\text{Cr(H}_2\text{O)}_6^{+3}$, and only small amounts of $\text{Cr(H}_2\text{O)}_5\text{X}^{+2}$ are formed. On the other hand, when thiocyanate ion is used, a marked change in mechanism, stoichiometry and rate of reaction is observed. The most interesting observation is that the ratio of $\text{Cr(H}_2\text{O)}_5\text{SCN}^{+2}/\text{Cr(H}_2\text{O)}_6^{+3}$ is greater than 1000.

The stoichiometry of the reaction is followed by means of ion exchange studies and subsequent spectral observation. We have also studied the rate of formation and decomposition of the intermediate, $[\text{Cr}^{\text{III}}-\text{O}-\text{U}^{\text{V}}-\text{O}]$, in the presence and absence of thiocyanate ion. Considerable data have been collected which show a definite retardation of the reaction in the presence of thiocyanate ion. It is surprising that there is no apparent kinetic dependence on the concentration of thiocyanate ion. All of these observations are consistent with an interpretation which suggests the primary role of thiocyanate ion is that of a "non-bridging" rather than a "bridging" ligand. In either case, the electron transfer process is markedly affected by the presence or absence of thiocyanate ion.

In an attempt to improve our understanding of the role of non-bridging ligands in electron transfer processes and to allow for a quantitative interpretation of the chromium(II)-uranyl ion reaction in the presence of thiocyanate ion, we have undertaken a series of aquation and chromium(II) assisted

aquation reactions of various tetraamminechromium(III) complexes.

The object of this investigation is to study the labilizing effect of ligands bound in cis and trans positions. It is well known that iodide ion exhibits a trans effect in square planar complexes of platinum(II). Do similar effects exist for chromium(III) complexes? Moore, Basolo and Pearson², employing H_2O^{18} , have shown that hexaquo-chromium(III) produced from the aquation of iodo-pentaquo-chromium(III) ion contains two labeled water molecules. They have suggested that the water molecule in the position trans to the iodide ion is labilized which allows for the penetration of the first coordination sphere of chromium(III) by a labeled water molecule prior to the loss of iodide ion.

The recent synthesis in this laboratory of cis and trans iodo-aqua-tetraamminechromium(III) ion suggested a series of experiments in which attention could be focused on the two relatively labile sites on the chromium(III) complex. These two sites are occupied by water molecules in either the cis or trans position. In the case of the aquation of halo-pentaquo-chromium(III) ions, the results were interpreted in terms of a long-lived penta-coordinated intermediate and a strong labilizing effect exerted by iodide ion. Experimentally, the five aquo ligands are indistinguishable which results in some ambiguity. We have chosen the tetraamminechromium(III) series in that this ambiguity is removed (the four ammonia ligands are relatively inert) and it should be possible to distinguish both cis and trans effects.

We have prepared both the cis and trans $Cr(NH_3)_4(OH_2)I^{+2}$ and trans $Cr(NH_3)_4ICl^+$ ions. The stoichiometry and kinetics of the aquation reactions have been investigated in a perchlorate media over a wide range of hydrogen ion concentrations. We always observe retention of configuration as is shown

below:

First-order rate constants and activation parameters for the loss of coordinated iodide are: $k = 1.72 \times 10^{-4} \text{sec}^{-1}$, $\Delta H^\ddagger = 21.8 \text{ kcal/mole}$, $\Delta S^\ddagger = -1.82 \text{ eu}$ at $30.0^\circ \pm 0.1^\circ\text{C}$ for $\text{trans Cr(NH}_3)_4(\text{OH}_2)\text{I}^{+2}$; $k = 1.13 \times 10^{-3} \text{sec}^{-1}$, $\Delta H^\ddagger = 20.4 \text{ kcalmole}^{-1}$, $\Delta S^\ddagger = -6.06 \text{ eu}$ at $25.0 \pm 1^\circ\text{C}$ for $\text{cis Cr(NH}_3)_4(\text{OH}_2)\text{I}^{+2}$; $k = 0.365 \text{ sec}^{-1}$, $\Delta H^\ddagger = 19.2 \text{ kcalmole}^{-1}$, $\Delta S^\ddagger = +5.0 \text{ eu}$ at $10.0 \pm 0.1^\circ\text{C}$ for $\text{trans Cr(NH}_3)_4(\text{OH})\text{I}^+$; $k = 1.27 \text{ sec}^{-1}$, $\Delta H^\ddagger = 18.2 \text{ kcalmole}^{-1}$, $\Delta S^\ddagger = +3.8 \text{ eu}$ at $10.0 \pm 0.1^\circ\text{C}$ for $\text{cis Cr(NH}_3)_4(\text{OH})\text{I}^+$; $k = 3.28 \times 10^{-3} \text{sec}^{-1}$, $\Delta H^\ddagger = 18.6 \text{ kcalmole}^{-1}$, $\Delta S^\ddagger = -3.2 \text{ eu}$ at $25.0 \pm 0.1^\circ\text{C}$ for $\text{trans Cr(NH}_3)_4\text{ICl}^+$, all at ionic strength 1.0M. The aquation of $\text{trans Cr(NH}_3)_4\text{ICl}^+$ is not dependent on hydrogen ion in the concentration range 0.1 to 1.0M.

The chromium(II) catalyzed aquation of the cis and trans iodoaquo ions was observed to produce iodopentaquo chromium(III) as a primary product. Second-order rate constants and activation parameters for the chromium(II) reductions are: $k = 14.6 \text{ M}^{-1} \text{sec}^{-1}$, $\Delta H^\ddagger = 8.4 \text{ kcalmole}^{-1}$, $\Delta S^\ddagger = -25 \text{ calmole}^{-1} \text{deg}^{-1}$ at $25.0 \pm 0.1^\circ\text{C}$ for $\text{cis Cr(NH}_3)_4(\text{OH}_2)\text{I}^{+2}$; and $k = 165 \text{ M}^{-1} \text{sec}^{-1}$, $\Delta H^\ddagger = 7.5 \text{ kcalmole}^{-1}$, $\Delta S^\ddagger = -25 \text{ calmole}^{-1} \text{deg}^{-1}$ at 25.0°C for $\text{trans Cr(NH}_3)_4(\text{OH}_2)\text{I}^{+2}$, both at unit ionic strength and 1 M HClO_4 .

These results are most consistent with a mechanism in which a pentacoordinated chromium(III) intermediate is formed prior to the formation of the final chromium(III) product. Additional experiments in the presence of competing anions are presently in progress and are described in the new proposal.

As part of our study of the reduction of chlorine-containing oxidizing agents by metal ions which may undergo two-electron oxidations, we felt it would be of interest to extend this study to metal ions which are limited to

a single electron oxidation. Since the chlorine-containing halogenates are known to undergo both one- and two-electron transfer reactions with $V(II)$,³ $Fe(II)$,⁴ $Cr(II)$,^{5,6} $U(IV)$, and $Fe(phen)_3^{+2}$, it may be possible to restrict these halates to exclusively a single electron transfer process with a metal ion such as $V(IV)$. In the complete reduction of chlorate ion or chlorine dioxide, do a series of stepwise reductions occur or do inter-halate reactions provide alternate pathways for product formation and possible catalytic activity? Furthermore, what role do bridging and non-bridging ligands have on rates and mechanisms of these electron-transfer reactions?

The reactions we have studied are examples of a general type of reaction where an oxyhalogen anion or molecule oxidizes a metal ion or metal complex and is itself reduced to its lowest oxidation state. Thus, the overall process involves rupture of all the oxygen bonds of the oxyhalogen anion or molecule. The rate determining step could be electron transfer from the central metal ion to an orbital on the halogen. On the other hand, it is possible that atom transfer takes place by means of an oxygen atom in a two-equivalent change or by means of an $[O^-]$ ion in a one-equivalent change.

The reactions between oxovanadium(IV) and chlorate ion and chlorine dioxide are in their final stages of completion. Both of these systems are most amenable owing to the pronounced differences in rates with which they react with VO^{+2} . For example, the kinetics of the reactions with chlorate ion can be followed by conventional techniques. In these studies, we have demonstrated the existence of ClO_2 , both as an intermediate and first reduction product. Stopped-flow techniques, on the other hand, have been employed to study the subsequent reactions of this intermediate species. Rapid-mixing techniques are employed to gather direct information on the reduction of ClO_2 by VO^{+2} .

The reaction

follows an apparent rate law of the general form

$$\frac{-d[\text{V(IV)}]}{dt} = \frac{k_1 + k_2[\text{H}^+]}{1 + K[\text{Cl}^-]} [\text{V(IV)}][\text{ClO}_3^-]$$

The chloride ion dependence lends support to the fact that VO^{+2} is partially complexed in the presence of chloride ions. The kinetics show that as the concentration of chloride ion increases, the rate of reaction decreases to a significant degree. This indicates that the more reactive species in solution is VO^{+2} rather than VOCl^+ .

A detailed mechanism involving the nature of elementary steps and the geometry of the activated complex has not yet been established, but the results strongly suggest that the rate of reduction of the chlorine-oxidants is governed by the structural properties of the metal ion complex.

Our studies have confirmed that ClO_2 is the first of several intermediate species arising from the reductive restriction imposed by oxovanadium(IV) on chlorate. Direct kinetic and stoichiometric measurements on the reduction of ClO_2 show ~~that apparently~~ a successive single electron transfer does occur. Experiments were performed to determine the stoichiometric coefficient, α , in the following reaction:

It was found that the reaction has a variable stoichiometry depending on the absence or presence of chloride ion, i.e., $\alpha = 1.5$ or 1.0 , respectively.

This indicates that reductions occur to form a Cl(III) species and an unstable Cl(II) species. Interhalate reactions occur during this period of reduction to produce an oxyanion of increased oxidation state and the initial product decomposes at its characteristic slow rate. Preliminary studies have also

indicated that the reduction of chlorite proceeds via the same intermediate as does ClO_2 and is consistent with this mechanism.

Preliminary kinetic runs on the chlorine(I)-tris(1,10-phenanthroline)-iron(II) system indicated that it might be possible to study the system in acid solution, where the predominant oxidizing species would be hypochlorous acid. It was observed in acid sulfate solution that the rate increases as the reaction proceeds. We attributed this autocatalytic behavior to the formation of Cl^- , as a product of the reduction of HOCl , followed by the reaction of Cl^- with HOCl to give Cl_2 according to the following equilibrium:

Chlorine is estimated⁷ to react about 1000 times faster than HOCl with $\text{Fe}(\text{phen})_3^{++}$. Consequently, even trace amounts of chloride, produced in the first 5% of the reaction, were found to contribute to an increase in the overall reaction rate.

The most promising results have been obtained from a study of the system under slightly alkaline conditions. $\text{Fe}(\text{phen})_3^{++}$ is stable in the pH region 2-9, which means that neutral or slightly basic conditions may be utilized without encountering the problem of base hydrolysis of the complex and in this region, interference from chlorine is essentially nonexistent. The problem of two oxidizing species does exist--namely OCl^- and HOCl . It seems likely that the dependence upon each of these oxidants can be determined by varying the pH.

The reaction has been followed by observing the change in absorption at 5100 Å. Although deviations from pseudo first order behavior appear after about 40% reaction, the pseudo first order rate constants calculated from the initial 40% of the reaction are constant and vary only with hypochlorite

ion and hydrogen ion concentration. A graph of k_{obs} as a function of the hypochlorite ion concentration is linear with an intercept (k_0) of $7.3 \pm 0.2 \times 10^{-5} \text{sec}^{-1}$ and a slope ($k_1 + k_2$) of $3.09 \pm 0.14 \times 10^{-3} \text{M}^{-1} \text{sec}^{-1}$. The dominant oxidizing species is hypochlorite ion and additional experiments in which the hydrogen ion concentration was varied suggest that the rate law is comprised of three terms as given below:

$$\text{Rate} = \{k_0 + k_1[\text{OCl}^-] + k_2[\text{HOCl}]\} \{[\text{Fe}(\text{phen})_3^{++}]\}$$

The existence of three terms in the rate law is consistent with two different mechanisms; one dissociative and one associative. This is the first time we have been able to observe both mechanisms for the reduction of chlorine-containing oxidizing agents. We are hopeful that continued investigation of the details of this reaction will give us considerable insight into the distinguishing characteristics of these two mechanisms.

Robert Buchacek and Gene Grimley are working in the general area of the effect of scavengers on the removal of intermediates in reactions which involve the reduction of various chlorine-containing oxidizing agents. Phenol appears to be a very effective scavenger for the removal of hypochlorous acid and the proposed Cl_2O_2 intermediate in the uranium(IV)-chlorous acid reaction.⁸ In the presence of phenol, the uranium(IV) reaction can be written as follows

Initial oxygen-18 tracer experiments in the presence of phenol indicate that more than one chlorite oxygen is transferred to the uranyl ion product. In view of the large difference in oxygen exchange lability between uranium(V) and uranium(VI) and the stoichiometry shown in the above equations, we suggest that the reaction is indeed a two electron exchange reaction between uranium(IV) and chlorite ion proceeding through direct coordination of the

chlorite oxygen to the uranium(IV) reactant.

This is particularly interesting in that it suggests that two electron processes in the case of the chlorine-containing oxidizing agents proceed via oxygen bridging whereas one electron transfer processes occur via coordination through chlorine. Several additional reactions in the presence of other scavengers and the effect of these scavengers on the chlorine-containing intermediates must be carried out before the microscopic details of these reactions are clearly understood. However, it appears that we may have a distinct tool for distinguishing between one and two electron transfer processes which involve the reduction of chlorine-containing oxidizing agents. Work will be continued along these general directions.

The final project which deserves some comment in this progress report is our double-tagged tracer study of the decomposition of uranyl peroxide dihydrate which decomposes thermally as is shown below:

The method we are using to study this decomposition is to enrich half of the peroxide oxygen atoms with oxygen-18 and to mix this with an equivalent amount of hydrogen peroxide of normal oxygen-18 content. The resulting uranyl peroxide is decomposed thermally between 150 and 400°. By studying the mass 32, 34, and 36 peaks with our Nuclide analysis double beam isotope ratio mass spectrometer it is possible to determine the amount of mixing of adjacent oxygen atoms during the decomposition process.

Our results indicate that less than 10% mixing of adjacent peroxide molecules occurs during the decomposition. This would indicate that the following unimolecular mechanism predominates during the solid state decomposition:

This mechanism is supported by an additional tracer study which indicates that peroxide oxygen also becomes part of the UO_3 product without exchange with the product water.⁹

A possible explanation for the small amount of mixing of adjacent peroxide molecules during the thermal decomposition could be oxygen exchange with the product UO_3 or initial uranium peroxide dihydrate reactant prior to decomposition. In an attempt to explore these possibilities we have installed an automatic toeppler pump on our vacuum system in direct link with the mass spectrometer and are hopeful of being able to rapidly separate the products from the reactants during the thermal decomposition. The final experiments should be finished in the next several months and will form one part of the Ph.D. thesis for Mr. Robert Seyse. A preprint of the manuscript describing these experiments will be submitted separately in the near future.

References

1. G. Gordon, Inorg. Chem., 2, 1279 (1963).
2. P. Moore, F. Basolo, and R. Pearson, Inorg. Chem., 5, 223 (1966).
3. G. Gordon and P. H. Tewari, J. Phys. Chem., 70, 200 (1966).
4. B. Z. Shakhashiri and G. Gordon, J. Am. Chem. Soc., 91, 1103 (1969).
5. R. C. Thompson and G. Gordon, Inorg. Chem., 5, 557 (1966).
6. R. C. Thompson and G. Gordon, Inorg. Chem., 5, 562 (1966).
7. B. Z. Shakhashiri and G. Gordon, Inorg. Chem., 7, 2454 (1968).
8. G. Gordon and F. Feldman, Inorg. Chem., 3, 1728 (1964).
9. G. Gordon and H. Taube, Inorg. Chem., 1, 69 (1962).

III. Incident Reports

In compliance with the AEC regulation that reports be submitted on accidents, fires, and unusual operational incidents which occur in conjunction with work being performed under AEC contract, the following are reported:

1. Serious accidents	- none
2. Fires, explosions, etc.	- none
3. Other property damage	- none
4. Personal injuries	- none