UNIVERSITY OF CALIFORNIA - BERKELEY

UCRL-764

~~SECRET~~

**DO NOT
PHOTOSTAT**

CAUTION

~~This document contains information affecting the
National Defense of the United States.
Its transmission or the disclosure of its contents in
any manner to an unauthorized person is prohibited
and may result in severe criminal penalties under
applicable Federal Laws~~

RADIATION LABORATORY

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

Cover Sheet
Do not remove

INDEX NO.

This is copy 73 of 96 Series A

~~SECRET~~

Each person who receives this document must sign the cover sheet in the space below

[illegible]

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. It is the property of the U.S. Atomic Energy Commission, and its use is limited to the Commission, nor any person acting on its behalf. It is not to be distributed, copied, or represented, express or implied, with reference to the use of any information, apparatus, method, or process disclosed in this report, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report. As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission to the extent that such employee or contractor prepares, handles or distributes, or provides access to, any information pursuant to his employment or contract with the Commission.

UCRL-764
Chemistry-General Distribution

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

CLASSIFICATION CANCELLED
DATE 12-8-55
For The Atomic Energy Commission

H. S. Crandall
Chief, Declassification Branch *let*

THE DISTRIBUTION OF TRACE URANIUM (VI) IN
NITRIC ACID--TTA--ORGANIC SOLVENT MIXTURES

D. L. Heisig and H. W. Crandall

June 30, 1950

Photostat Price \$ 4.80
Microfilm Price \$ 2.70

Available from the
Office of Technical Services
Department of Commerce
Washington 25, D. C.

Berkeley, California

866 001

THE DISTRIBUTION OF TRACE URANIUM (VI) IN
NITRIC ACID--TTA--ORGANIC SOLVENT MIXTURES

D. L. Heisig and H. W. Crandall
Radiation Laboratory
University of California, Berkeley, California

June 30, 1950

INTRODUCTION

The distribution of uranium (VI) in benzene-water-perchloric acid-TTA¹ mixtures has been studied by E. L. King² and W. C. Orr.³ Orr reported the value of the equilibrium constant for the reaction

at $\mu = 1.9$ of lithium perchlorate and perchloric acid as

$$K = \frac{(\text{UO}_2\text{K}_2)(\gamma_{\text{UO}_2\text{K}_2})}{(\text{UO}_2^{++})} \frac{(\text{H}^+)^2}{(\text{HK})^2} \frac{1}{\gamma_{\text{HK}}^2} = .0055$$

Similar values are obtained from King's data. The distribution coefficients of trace uranium (VI) in the above system when TTA is 0.2M and nitric acid replaces perchloric acid have been measured by E. K. Hyde and J. Tolmach.⁴

In this report the distribution coefficients of trace uranium (VI) in mixtures of dilute nitric acid and solutions of TTA in hexone, cyclohexanone, and pentaether are presented.

EXPERIMENTAL

In all the experiments U^{233} was used, generally at about 5×10^{-6} to 10^{-5} M. The experiments with hexone and many of those with cyclohexanone were run at room temperature on a 4 cc. scale in small flasks equipped with motor driven stirrers. In the pentaether experiments 2 cc. volumetric flasks were filled and placed on a revolving wheel in a 25° thermostat. In all the experiments the volume ratio was one.

Several experiments in hexone and cyclohexanone, using the motor driven stirrers, were followed as a function of time to determine the rate of equilibration. As one would expect from the fact that two to three hours are required for TTA initially in dry benzene to reach equilibrium between benzene and water,⁵ the equilibration of uranium (VI), TTA, nitric acid, and hexone is slow, two to three hours, if the aqueous and organic phases are not equilibrated before the addition of the uranium (VI) (Table I). Equilibrium of the uranium (VI) in the preequilibrated phases was reached in less than fifteen minutes when a mechanical stirrer was used, although the rate varied with the speed of stirring. When samples were shaken on the revolving wheel, the phases were not equilibrated before the addition of uranium (VI). By making plates at the end of three and four hours it was shown that equilibrium had been reached by the end of three hours.

Plates were made from 50 to 100 portions of each phase and were counted with $\pm 1.5\%$ accuracy.

Commercial grade hexone was used without any purification. Commercial cyclohexanone and pentaether were redistilled. The cyclohexanone used boiled at 153.1°C . while the pentaether boiled at 190°C at 9 mm. to 193°C . at 11 mm. The TTA was synthesized by M. W. Davis, Jr. and H. R. Lehman. The molarity of the solutions of TTA calculated from the weight of TTA used and from direct titration differed by less than 2%.

DISCUSSION

When nitric acid, TTA, uranium (VI) and an organic solvent are mixed a number of reactions may take place.

In the aqueous phase:

In the two phase system:

E. L. King's data² and those of W. C. Orr³ with perchloric acid--TTA--benzene mixtures indicate that little complexing of uranyl ion with TTA occurs in the aqueous phase, Reactions (c) and (d), when the hydrogen ion and TTA concentrations in the aqueous phase are in the range of those used in the experiments in this report. King found that a plot of $\log E' = \log \frac{(\text{U org.}) \gamma_{\text{UO}_2\text{K}_2}}{(\text{U aq.})}$ vs. $\log (\text{TTA}) (\gamma_{\text{TTA}})$ at constant acid gave a straight line with a positive slope of two which is the theoretical slope for Reaction (3) involving the uncomplexed uranyl ion. Orr estimated that the value of the equilibrium constant for Reaction (c) is about 0.15. With this value the fraction of UO_2^{++} complexed as UO_2K^+ in the aqueous phase was calculated and found to be less than one percent under the most favorable complexing conditions used, e.g., Experiment C29 with 0.0745M nitric acid and 0.001M TTA in the aqueous phase. The concentration of the TTA in the water phase was calculated from the distribution coefficient of TTA in a mixture of pentaether and 0.157M nitric acid, which was found to be approximately 300 by H. W. Crandall and D. L. Heisig.⁶ From these considerations it was concluded that Reactions (c) and (d) are of little importance in this system.

At 8.0M total salt concentration the constants for Reactions (a) and (b) have been calculated from distribution measurements between benzene and an aqueous

phase and reported as⁷

$$K_a = \frac{(UO_2NO_3^+)}{(UO_2^{++})(NO_3^-)} = .116 \text{ and}$$

$$K_b = \frac{UO_2(NO_3)_2}{(UO_2^{++})(NO_3^-)^2} = .031$$

More recently R. H. Betts and Rita K. Michels⁸ have determined the value of K_a from absorption data. Their value is 0.21 at $\mu = 5.4$, and these authors state that the presence of a second complex is doubtful at this salt concentration.

All of the uranium in the aqueous phase can be represented by the equation

$$(5) U_{aq.} = (UO_2^{++}) + (UO_2NO_3^+) + (UO_2(NO_3)_2) = (UO_2^{++}) [1 + K_a (NO_3^-) + K_b (NO_3^-)^2]$$

where activity effects are neglected. This equation shows that constants or distribution coefficients calculated on the basis that $U_{aq.} = (UO_2^{++})$ will be in error from 1 to 2% at 0.1M (NO_3^-) to 15 to 20% at 1M (NO_3^-) depending on which constants are used. For a given nitrate ion concentration the fraction of uranyl ion complexed is constant, and in this report corrections for obtaining the fraction of UO_2^{++} uncomplexed by nitrate ion are not used. Nitrate complexing will account for part of the variation of the constants K_1 , K_2 , and K_3 with nitric acid concentrations.

Corresponding to Equations (1), (2), (3), and (4) are the following extraction coefficients and equilibrium constants where subscript_o designates the concentration in moles per liter in the organic phase.

$$(6) K_1 = \frac{(UO_2(NO_3)_2)_o}{(UO_2^{++})(NO_3^-)^2} = K_1^o \frac{\gamma_{UO_2^{++}} \gamma_{NO_3^-}^2}{\gamma_{UO_2(NO_3)_2}} = \frac{E_1}{(NO_3^-)^2}$$

$$(7) K_2 = \frac{(UO_2NO_3K)_o(H^+)}{(UO_2^{++})(NO_3^-)(HK)_o} = K_2^o \frac{\gamma_{UO_2^{++}} \gamma_{NO_3^-} \gamma_{HK}}{\gamma_{UO_2NO_3K} \gamma_{H^+}}$$

$$(8) K_3 = \frac{(UO_2K_2)_o (H^+)^2}{(UO_2^{++})(HK)_o^2} = K_3^o \frac{\gamma_{UO_2^{++}} \gamma_{HK}^2}{\gamma_{UO_2K_2} \gamma_{H^+}^2}$$

$$(9) K_4 = \frac{(HNO_3)_o}{(H^+)(NO_3^-)} = \frac{E_{HNO_3}}{(NO_3^-)}$$

$$(10) E_M = \frac{c/m/100\lambda \text{ in org. phase}}{c/m/100\lambda \text{ in aq. phase}}$$

In general

$$(11) E_M = \frac{(UO_2K_2)_o + (UO_2KNO_3)_o + (UO_2(NO_3)_2)_o}{(UO_2^{++})} = \frac{(UO_2K_2)_o + (UO_2KNO_3)_o}{(UO_2^{++})} + E_1$$

Combination of Equations (11), (7), and (8) gives

$$(12) E_M = \frac{(HK)_o^2}{(H^+)^2} \frac{\gamma_{HK}^2}{\gamma_{UO_2K_2}} \frac{\gamma_{UO_2^{++}}}{\gamma_{H^+}^2} K_3^o + \frac{(HK)_o(NO_3^-)}{(H^+)} \frac{\gamma_{UO_2^{++}} \gamma_{HK} \gamma_{NO_3^-}}{\gamma_{H^+} \gamma_{UO_2NO_3K}} K_2^o + E_1$$

$$\text{or } (13) E_M - E_1 = \frac{(HK)_o^2}{(H^+)^2} K_3 + \frac{(HK)_o(NO_3^-)}{(H^+)} K_2$$

The nitric acid distribution, Reaction (4), between water and the organic liquid was determined by titration or had been determined previously.^{9,10} These values together with the distribution coefficients, E_1 , of uranyl nitrate between dilute nitric acid and each of the organic solvents in the absence of TTA are recorded in Columns 8 and 9 of Table II. Volume changes were known in the penta-ether system only and in this system no corrections were necessary except at nitric acid concentrations greater than one molar. Since both the nitric acid and uranium (VI) distribution coefficients were determined by analyzing equal volumes of the organic and aqueous phases only the TTA concentration had to be corrected for the volume changes. If sodium nitrate was used in the experiment, E_1 , differed from the value obtained at the same equilibrium nitrate concentration in the aqueous phase with nitric acid alone. This was especially noticeable at high ionic strengths and is probably due to activity effects and to differences in the

concentration of the nitric acid in the organic phase. In hexone, E_1 was neglected since it is small compared to the corresponding E_M .

That both Reactions (2) and (3) contribute to the distribution coefficient of uranium (VI) in these solvents can be seen from the data for both pentaether and for cyclohexanone. (Tables III to VII, and Figures 1 to 7.) In Figures 1, 2, 3, 4, and 6 the experimental data are represented by points while the lines drawn are theoretical curves calculated from the constants described in a later section of this report. At high TTA concentrations and low nitric acid concentrations the first term in the right hand side of Equation (13) should be large compared to the second, providing K_2 and K_3 are of comparable magnitude. Under these conditions and at constant nitrate and hydrogen ion concentrations, neglecting activity effects, Equation (13) becomes

$$(14) \log (E_M - E_1) = 2 \log HK + \text{Constant}.$$

The experimental points along Curves A and B, Figures 1 and 2, taken from data in Tables III and V satisfy the above conditions of low nitrate and high TTA, and lie on curves whose slope is plus two. Only an approximation is expected as activity coefficients are omitted.

Similarly, at high nitric acid concentration and low TTA concentration the second term in Equation (13) predominates and at constant hydrogen and nitrate ion concentration, neglecting activity effects, Equation (13) becomes

$$(15) \log (E_M - E_1) = \log HK + \text{Constant}.$$

The experimental points along Curve D, Figure 2, and Curve E, Figure 1, satisfy the above conditions, and the slopes of the curves drawn through the experimental points approach plus one. Figures 1 and 2 show that as the nitric acid concentration is increased the TTA dependence on $(E_M - E_1)$ at constant ionic strength decreases from two to one in agreement with Equation (13).

Figures 3 and 4 are plots of $\log (H^+)$ vs. $\log (E_M - E_1)$ at constant TTA concentration using the data in Tables III and V. From Equation (13) two extremes are predictable for the hydrogen ion dependence on $(E_M - E_1)$ where the TTA concentration is constant and activity effects are neglected. At low hydrogen ion concentration and high TTA concentration the first term on the right hand side of Equation (13) predominates and

$$(16) \log (E_M - E_1) = -2 \log (H^+) + \text{Constant}.$$

The slopes of curves connecting the experimental points at low nitric acid concentration in Figures 3 and 4 approach a limit of minus two. At high nitric acid concentration and low TTA concentration Equation (13) reduces to

$$(17) \log (E_M - E_1) = 0 \log (H^+) + \text{Constant}$$

provided that nitric acid is the only source of nitrate and hydrogen ions. The slopes of the curves connecting the experimental points at high nitric acid concentration in Figures 3 and 4 approaches zero.

The dependence of $(E_M - E_1)$ on the hydrogen ion concentration when the ionic strength and the nitrate ion concentration are maintained constant is shown in Figure 6 from the data in Tables IV and VI.

Slopes of -1.1 in cyclohexanone and -2.0 in pentaether are obtained in regions where the slopes predicted from Equation (13) using the constants obtained at the same ionic strength but at constant nitric acid concentration are -1.0 and -2.0, respectively. Figures 3, 4, and 6 show that as the nitric acid concentration is increased, the hydrogen ion dependence on $(E_M - E_1)$ at a given TTA concentration increases from minus two to zero if nitric acid is the only source of nitrate and hydrogen ions and to minus one if sodium nitrate is present. Thus, they support the mechanism summarized by Equation (13).

The nitrate dependence at high nitrate ion concentration and low TTA concentration is shown by Figure 7 from the data in Tables IV and VI. In Figure 7 the

lines shown connect the experimental points and are not calculated. The data used in Tables IV and VI are less accurate than most of the other data. In the case of the experiments in pentaether ($E_M - E_1$) represents the difference between two distribution coefficients of comparable size, e.g., $E_1 = 0.485$ and $E_M = 0.730$. Furthermore, E_1 and E_M are calculated from counting plates which contain sodium nitrate in varying amounts. The values of E_1 in cyclohexanone which are recorded in Table VI are extrapolated from a graph of E_1 measured when nitric acid was the only nitrate source against the concentration of nitric acid in the aqueous phase after equilibrium had been reached. It was assumed that E_1 for a given nitrate concentration was independent of the cation. This is not exactly true, but gives an indication of ($E_M - E_1$) for the experiments in Table VI. The counting for these experiments was also done in the presence of sodium nitrate which reduces the counting accuracy. The ionic strength varied from 0.6 to 0.9 in the nitrate dependence experiments in pentaether, and from 0.4 to 0.6 and 0.6 to 0.9 in cyclohexanone. With this change in ionic strength the activity coefficients of nitric acid¹¹ as well as those of uranyl nitrate¹² vary less than ten percent, but it is not known how the activities of the uranium complexes in the organic phases vary. However, since the over-all change in the ionic strength is rather small for each series, and since the nitrate dependence is observed in regions in which it was independently predicted from the variation of ($E_M - E_1$) with TTA and hydrogen ion concentrations, it is believed that the nitrate dependence curves are real.

The maximum experimental error in these determinations is probably not more than 20% and reproducibility seems to be nearer 10%. If the two extreme points in Curve C, Figure 7 are taken and if it is assumed that the upper point is actually 20% less and the lower curve is 20% greater than that recorded a line of slope 0.3 is obtained, an indication that the inaccuracies of the measurements cannot account for the nitrate dependence curves.

The hexone data in Table VIII are insufficient to demonstrate that both Equations (2) and (3) apply to the hexone system. However, in Figure 5 a plot of $\log E_M$ vs. $\log TTA$ at two hydrogen ion concentrations shows a decrease of slope from 1.8 to 1.6 as the equilibrium nitric acid concentration increases from 0.23 to 0.43 paralleling the trend in Figures 1 and 2 for pentaether and cyclohexanone. This supports the supposition that the extraction mechanism of uranium (VI) by TTA is the same in hexone as it is in pentaether and cyclohexanone and can be represented by Equations (1), (2), and (3).

ACTIVITY EFFECTS

Insertion of activity coefficients in any of the equations mentioned will not alter the qualitative picture just presented of the mechanism of extraction of uranium (VI) into cyclohexanone, hexone, and pentaether. When the ionic strength and the nitric acid concentration are constant, Figures 1 and 2, the only activities changing are those of TTA, UO_2K_2 , UO_2KNO_3 , and $UO_2(NO_3)_2$ in the organic phase. The activity coefficients of TTA in pentaether which has been equilibrated with 0.5M nitric acid have been measured.⁶ In going from 0.045M TTA to 0.378M TTA the activity coefficient of TTA increases from 1.01 to 1.19. If $\log (E_M - E_1)$ is plotted against $\log (TTA) \gamma_{TTA}$ using the data in Curves A and B, Figure 1, a slope of 1.9 as compared to 2.0 without the activity coefficients is obtained.

It is not known whether the activity coefficient of the uranium complexes in the organic phase increases or decreases with TTA. In benzene the form is UO_2K_2 and its activity coefficient is known to decrease with TTA presumably due to the formation of $UO_2K_2 \cdot HK$.^{2,3} However, TTA is known to interact strongly with pentaether ($\frac{TTA_{org.}}{TTA_{aq.}} = 300$ in pentaether as compared to 40 in benzene),⁶ and UO_2K_2 as evidenced from its one hundred fold increase in extractability in pentaether as compared to benzene, also appears to interact strongly with

pentaether. Hence, it is probable that the activity coefficient of UO_2K_2 will decrease much less rapidly with the TTA concentration in pentaether than in benzene since the complex $\text{UO}_2\text{K}_2 \cdot \text{HK}$ is less likely to form. In fact, like the activity coefficient of TTA in pentaether, that of UO_2K_2 may increase with increasing TTA concentrations, and it seems reasonable to assume that the behavior of UO_2KNO_3 is similar to that of UO_2K_2 under these conditions. In this event the changes in the activity coefficients of UO_2K_2 and UO_2KNO_3 tend to cancel the increase of γ_{TTA} and the activity effects at constant aqueous phase composition would be small. Further discussion of the interaction of TTA and pentaether will be found in another report from this laboratory.⁶

On the other hand when the TTA concentration is held constant and the nitric acid concentration is varied, the activities of the TTA, UO_2K_2 , and UO_2KNO_3 in the organic phase are not constant since it is known that the composition of the organic phase as well as the activity of TTA in pentaether changes with the concentration of nitric acid in the aqueous phase. Large variations of the constants K_2 and K_3 are observed as the ionic strength increases from 0.1 to 1.0.

However, the experiments in Figure 6 from the data in Tables IV and VI were carried out at constant ionic strength as well as constant TTA concentration. Under these conditions the activity coefficients in both phases should be fairly constant, especially since the total nitrate ion concentration is constant and the hydrogen ion concentration is varied over a limited range only. Since the activity effects at constant ionic strength are small, any changes in the plot of $\log (E_M - E_1)$ vs. $\log (H^+)$ in Figure 6 that would result from the use of activity corrections must also be small, and so could not alter the qualitative agreement already noted between the data in Figure 6 and the extraction mechanism for uranium (VI) summarized by Equation (13). Furthermore, since the experiments in Figures 3 and 4 where the ionic strength is varied agree qualitatively with the results obtained at

constant ionic strength, it is believed that the use of activity coefficients where the ionic strength is varied will not invalidate the trends previously observed in Figures 3 and 4.

CALCULATION OF CONSTANTS

Although the data presented here indicate that the distribution of uranium (VI) between dilute nitric acid and cyclohexanone, hexone, or pentaether is represented by Equations (1), (2), and (3), the absolute values of the constants K_1^0 , K_2^0 , and K_3^0 cannot be determined since the only activity coefficients known in these systems are those of TTA in pentaether. For pentaether and cyclohexanone, values for K_2 at high nitrate and low TTA concentrations and for K_3 at high TTA and low nitric acid concentrations can be calculated directly from the data, and these constants are given in Columns 7 and 8 of Tables III and V, and Column 11 of Tables IV and VI. It was found in all cases where data were available that at a given ionic strength the same value of a given constant was obtained when nitric acid was present alone in the aqueous phase as when nitric acid and sodium nitrate were both present.

Furthermore, it can be shown from Equation (13) that under the conditions where each of these constants is calculated the term in Equation (13) involving the other constant at its maximum value contributes less than 10 percent of the measured value of $(E_M - E_1)$ and can justifiably be neglected in the calculation. The starred values of K_2 and K_3 are calculated under conditions where the term in Equation (13) involving the other constant at the value recorded in Column 6 of Tables III and V, and Column 10 of Tables IV and VI contributes less than 10 percent of the measured value of $(E_M - E_1)$. Examples of these calculations are shown in Table VIII.

The constant values of K_3 at $\mu = 0.0745$ in pentaether and those of K_2 at $\mu = 0.385$ in cyclohexanone indicate that for TTA concentrations from 0.04 to 0.4M

$\frac{\gamma_{\text{UO}_2\text{K}_2}}{\gamma_{\text{HK}}^2}$ and $\frac{\gamma_{\text{UO}_2\text{KNO}_3}}{\gamma_{\text{HK}}}$ are constant within the experimental error if the ionic strength is constant. By assuming as a first approximation that K_2 and K_3 were independent of the TTA concentration rough values of K_2 and K_3 were assigned at various nitric acid concentrations where they could not be calculated directly from the data. These constants, which are recorded in Column 6 of Tables III and V, and Column 10 of Tables IV and VI are of the proper order of magnitude only and in some cases can be varied up to 20% and fit the data as well.

Figure 8 is a plot of K_2 and K_3 as well as the measured values of E_1 vs. the nitric acid concentration in the aqueous phase at equilibrium from the data in Columns 3 and 6 of Tables III and V and Columns 3 and 10 of Table VI. From these constants values of E_M can be calculated with an accuracy of 15% when the nitric acid and TTA concentrations are 0 to 0.9M and 0 to 0.6M, respectively, in pentaether and 0 to 0.8M and 0 to 0.5M in cyclohexanone. The calculated values of $(E_M - E_1)$ are shown in Column 6 of Tables III and V and in Column 10 of Tables IV and VI and are represented by the curves drawn in Figures 1, 2, 3, 4, and 6.

SUMMARY

phenyl trifluoroacetone

1. The distribution of uranium (VI) in systems of nitric acid, *(TTA)* and hexone, cyclohexanone, or pentaether has been shown to depend upon three *specific reactions*

2. The values of the equilibrium constants in pentaether and cyclohexanone when activity effects are neglected have been estimated at various nitric acid concentrations.

3. These constants can be used to calculate the distribution coefficient of uranium (VI) with an accuracy of 15 ⁷⁰ percent in pentaether-nitric acid mixtures containing up to 1.42M total nitric acid and 0.6M TTA and in cyclohexanone-nitric acid mixtures containing up to 1.0M total nitric acid and 0.5M TTA. (auth)

Table I
Equilibration of Uranium (VI) in
Hexone-TTA-Nitric Acid Systems

Experiment No. A5

Conditions: 0.444M HNO_3 , 0.375M TTA in hexone. No equilibration of phases, prior to the addition of U^{233} .

Time Min.	Counts/Min. Per 100 $\angle$ * Aq. Phase	Time Min.	Counts/Min. Per 100 $\angle$ Aq. Phase
0	1300	15	540
.5	469	60	729
1.0	328	126	834
2.0	354	210	850

Experiment No. A10

Conditions: 0.444M HNO_3 , 0.375M TTA in hexone. Phases equilibrated several hours before addition of U^{233} .

Time Min.	Counts/Min. Per 100 $\angle$ Aq. Phase
0	1280
.5	868
1.5	819
3.0	790
20.0	797

* The rise in the U^{233} concentration in the aqueous phase following its first dip occurs as TTA diffuses into the aqueous phase.

Table II

Distribution of Uranium Nitrate and Nitric Acid Between Equal Volumes
of Water and Various Organic Solvents in the Absence of TTA

Solvent	Ex.No.	Initial Aqueous Phase			Equilibrium		$E_{aq}^{org.}$	
		$UO_2(NO_3)_2$ M	HNO_3 M	$NaNO_3$ M	HNO_3 M (B) Org. Aq.		E_1 $UO_2(NO_3)_2$	E HNO_3
Hexone (A)			.252		.008	.244		.033
			.504		.029	.475		.061
	B48	4×10^{-6}	.350		.011	.339	.0055	.032
Cyclo- hexanone			.157		.013	.144		.090
			.157		.012	.145		.083
			.467		.087	.380		.229
			.467		.088	.379		.232
			.467		.093	.374		.249
			1.269		.422	.847(AA)		.498
	B47	4×10^{-6}	.101		.007	.094	.017	.074
	B17	4×10^{-6}	.234		.031	.203	.058	.152
	B16	4×10^{-6}	.479		.094	.385	.182	.244
	B46	4×10^{-6}	.985		.246	.739	.549	.333
	B27	4×10^{-6}	.985		.246	.739	.522	.333
	B39	4×10^{-6}	0	.948			.73	
	B29	4×10^{-6}	0	.948			.68	
Penta- ether	C30	10^{-5}	.0788		.0043	.0745	.0085	.058
	C12	10^{-5}	.158		.015	.142	.028	.106
	C2	4×10^{-6}	.473		.088	.384	.144	.229
	C4	4×10^{-6}	.473		.088	.384	.136	.229
	C3	4×10^{-6}	.788		.199	.588	.337	.338
	C1	4×10^{-6}	.788		.199	.588	.336	.338
	C41	10^{-5}	1.42		.49	.92	.879	.533
	C55	10^{-5}	.0315	.110	.0015	.030	.033	.050
	C56	10^{-5}	.788	.157	.199	.588	.485	.338
	C57	10^{-5}	.788	.259	.199	.588	.566	
	C67	10^{-5}	.788	.259	.199	.588	.572	
	C58	10^{-5}	.788	.315	.199	.588	.638	
	C68	10^{-5}	.788	.315	.199	.588	.626	
	C59	10^{-5}	.788	.047	.199	.588	.372	
	C63	10^{-5}	.788	.102	.199	.588	.422	
	C65	10^{-5}	.788	.102	.199	.588	.427	
	C69	10^{-5}	.788	.205	.199	.588	.537	

(A) See CN-3525, Fig. 18.

(AA) Determined by Peter C. Mayer

(B) Calculated from the data of Stover and Crandall, UCRL-649. It was assumed that the nitric acid distribution did not change with the addition of $NaNO_3$.

Table III

Distribution of Uranium (VI) in Pentaether-TTA-Nitric Acid Mixtures

Ex.No.	Initial HNO_3M	Equil. HNO_3M	TTA M	$(E_M - E_1)$	$(E_M - E_1)\text{calc.}$	K_2	K_3
					$K_2 = .15$ $K_3 = 1.15$		
C24	.0788	.0745	.0453	.411	.43		1.11*
C25			.0755	1.10	1.20		1.07
C26			.151	5.00	4.75		1.22
C27			.227	10.2	10.7		1.11
C28			.302	20.6	18.9		1.26
C29			.378	31.5	29.7		1.22
					$K_2 = .2$ $K_3 = 1.1$		
C13	.158	.142	.0755	.306	.32		1.08*
C14			.151	1.26	1.25		1.11*
C15			.227	2.63	2.80		1.03
C16			.302	4.92	4.93		1.09
C17			.378	7.71	7.75		1.09
					$K_2 = .5$ $K_3 = 1.0$		
C6	.473	.384	.0755	.085	.077		
C5			.151	.208	.23		
C7			.227	.397	.45		
C8			.302	.737	.77		
C9			.378	1.20	1.16		
C10			.453	1.65	1.62		
C11			.529	2.05	2.16		
					$K_2 = .8$ $K_3 = .9$		
C36	.788	.588	.0453	.037	.041	.82	
C37			.0755	.068	.075		
C38			.128	.153	.15		
C18			.151	.179	.18		
C39			.227	.331	.32		
C19			.302	.473	.48		
C40			.378	.665	.67		
C20			.453	.885	.90		
C21			.604	1.37	1.43		
C22			.755	1.84	2.09		
C23			.906	2.45	2.85		
					$K_2 = 1.2$ $K_3 = .8$		
(A) C42	1.42	.92	.0718	.084	.091	1.17	
C43			.144	.192	.193	1.20*	
C44			.216	.305	.30		
C45			.288	.416	.43		
C46			.360	.608	.56		

* Explained in Calculation of Constants, p.13.

(A) In Experiments C42-C46 the TTA concentration has been corrected for the volume change of 1 org./ 1 aq. to 1.05 org./ .95 aq.

860 C17

Table IV
Distribution of Uranium (VI) in Pentaether-TTA-Nitric Acid Mixtures
Nitrate Ion Dependence

Ex.No.	Initial HNO_3M	Equil. HNO_3M	NaNO_3M	TTAM	Total NO_3M	E_M	E_1	$(E_M - E_1)$	$(E_M - E_1)$ calc.	K_3
C18	.788	.588	0	.151	.588	.515	.336	.179		
C47			.047		.635	.575	.372	.203		
C48			.102		.690	.665	.425	.240		
C64			.102		.690	.665	.425	.240		
C49			.157		.745	.730	.485	.245		
C50			.205		.793	.822	.537	.285		
C60			.259		.847	.885	.569	.316		
C66			.259		.847	.866	.569	.297		
C52			.315		.903	.948	.632	.316		
C61			.315		.903	.940	.632	.308		

Hydrogen Ion Dependence

									$K_2 = .2$	
									$K_3 = 1.1$	
C31	.0315	.030	.110	.0755	.140	6.56	.03	6.53	7.0	1.03*
C32	.0473	.045	.095		.140	2.95		2.92	3.15	1.04*
C33	.0788	.075	.063		.138	1.10		1.07	1.15	1.06*
C34	.110	.103	.039		.142	.59		.56	.61	1.04*
C35	.134	.124	.016		.140	.37		.34	.43	.92
C13	.158	.142	0		.142	.33		.30	.33	1.06

* Explained in Calculation of Constants, p. 13.

Table V

Distribution of Uranium (VI) in Cyclohexanone-TTA-Nitric Acid Mixtures

Ex. No.	Initial HNO_3M	Equil. HNO_3M	TTA M	$(E_M - E_1)$	$(E_M - E_1)$ calc.	K_2	K_3
$K_2 = 1.5$							
$K_3 = 1.09$							
B55	.0504	.0479	.0383	.709	.76		1.11*
B56			.0765	2.79	2.9		1.09*
B57			.115	6.19	6.4		1.08
B58			.153	12.2	11.4		1.19
B59			.191	16.5	16.3		1.04
B60			.230	23.8	23.4		1.03
$K_2 = 2.0$							
$K_3 = 1.1$							
B49	.1008	.0935	.0383	.247	.26		
B50			.0765	.898	.89		
B51			.115	1.96	1.89		
B52			.153	3.39	3.18		
B53			.191	4.36	4.98		
B54			.230	7.09	7.04		
$K_2 = 3.2$							
$K_3 = .7$							
B28	.234	.203	.0383	.157	.15		
B6			.0765	.420	.34		
B5			.115	.624	.60		
B4			.153	.945	.89		
B3			.191	1.24	1.23		
B2			.230	1.45	1.64		
B18			.268	1.99	2.08		
B7			.268	2.12	2.08		
B19			.306	2.75	2.58		
$K_2 = 5.0$							
$K_3 = .10$							
B14	.479	.385	.0382	.184	.19	4.82	
B13			.0765	.395	.39	5.16	
B12			.153	.792	.78	5.18*	
B8			.230	1.18	1.19	5.14*	
B9			.306	1.51	1.59	4.93*	
B10			.383	1.92	2.02	5.02*	
B11			.459	2.21	2.44	4.81*	
B15			.536	2.81	2.87	5.24*	

*Explained in Calculation of Constants, p. 13.

Table VI

Distribution of Uranium (VI) in Cyclohexanone-TTA-Nitric Acid Mixtures

Hydrogen Ion Dependence

Ex.No.	Initial HNO_3M	Equil. HNO_3M	NaNO_3 M	TTA M	Equil. NO_3^-M	E_M	E_1	$(E_M - E_1)$	$(E_M - E_1)$ calc.	K_2
$K_2 = 5.2$										
B43	.473	.380	.362	.153	.742	2.13	.54	1.59	1.55	5.32*
B42	.630	.491	.248	.153	.739	1.74	.54	1.20	1.20	5.14
B41	.788	.602	.142	.153	.744	1.49	.54	.95	.98	5.04
B25	.985	.739	0	.153	.739	1.27	.54	.73	.80	4.78
B38	.985	.739	0	.153	.739	1.42	.54	.88	.80	5.74

Nitrate Ion Dependence

B43	.473	.380	.362	.153	.742	2.13	.54	1.59		5.32*
B45	.473	.380	.263	.153	.616	1.65	.41	1.24		4.98*
B12	.473	.380	.0	.153	.380	.97	.18	.79		5.16*
B41	.788	.602	.142	.153	.744	1.49	.54	.95		5.04
B33	.788	.602	.291	.153	.893	2.13	.64	1.49		6.56
B24	.788	.602	.197	.153	.799	1.62	.57	1.05		5.19
B40	.788	.602	.00	.153	.602	1.17	.41	.76		4.97

*Explained in Calculation of Constants, P. 13.

Table VII

Distribution of Uranium (VI) in Hexone-TTA-Nitric Acid Mixtures

Ex.No.	Initial HNO_3M	Equil. HNO_3M	TTA M	E_M
A19	.479	.454	.563	1.08
A5	.444	.423	.375	.554
A10	.444	.423	.375	.604
A9	.444	.423	.563	1.100
A8	.444	.423	.751	1.790
A12	.350	.339	.375	.834
A17	.350	.339	.563	1.46
A18	.350	.339	.563	1.52
A20	.350	.339	.563	1.86
A16	.234	.229	.150	.336
A15	.234	.229	.225	.756
A11	.234	.229	.375	1.71
A13	.234	.229	.563	3.83
A14	.234	.229	.563	3.74

Table VIII

Comparison of the Contributions of the Terms

$$K_3 \left(\frac{HK}{H^+}\right)^2 \text{ and } \frac{(HK)(NO_3^-)}{(H^+)} K_2 \text{ to } (E_M - E_1)$$

Measured in Pentaether

$$(E_M - E_1) = \left(\frac{HK}{H^+}\right)^2 K_3 + \frac{(HK)(NO_3^-)}{(H^+)} K_2$$

$K_3 \text{ Max.} = 1.15$
 $K_2 \text{ Max.} = 1.2$

Ex.No.	Equil. HNO_3 $\underline{M}$	TTA $\underline{M}$	$\frac{(HK)(NO_3^-)}{(H^+)} K_2$	$\left(\frac{HK}{H^+}\right)^2 K_3$	$(E_M - E_1)$ Measured	Max. Contri- bution of $\left(\frac{HK}{H^+}\right)(NO_3^-) K_2$ to $(E_M - E_1)$ measured %
C24	.0745	.0453	.054	.426	.411	13
C25		.0755	.091	1.18	1.10	8
C26		.151	.181	4.71	5.00	4
C27		.227	.272	10.7	10.2	3
C28		.302	.362	18.9	20.6	2
C29		.378	.454	29.7	31.5	1.5
						Max. Contri- bution of $\left(\frac{HK}{H^+}\right)^2 K_3$ to $(E_M - E_1)$ measured %
C42	.92	.072	.086	.0071	.084	9
C43		.144	.173	.028	.192	15

REFERENCES

1. TTA is thenoyltrifluoroacetone,

The symbol HK represents the TTA molecule while K^- represents that part of the molecule which remains after the removal of a proton.

2. E. L. King, Manhattan Project Metallurgical Laboratory Report CC-3618 (Aug., 1946).
3. W. C. Orr, UCRL-196.
4. E. K. Hyde and J. Tolmach, ANL-4248.
5. E. L. King and W. H. Reas, AECD-2573.
6. H. W. Crandall and D. L. Heisig, UCRL-778.
7. KAPL-94, p. 22.
8. R. H. Betts and Rita K. Michels, JCS, Supplementary Issue No. 2 (1949), p. 286.
9. Manhattan Project Metallurgical Laboratory Report CN-3525, Figure 18.
10. C. N. Stover, Jr. and H. W. Crandall, UCRL-649.
11. E. Abel, O. Redlick, B. Lengyel, Z. Phys. Chem. 132, 189
12. R. A. Robinson, J. M. Wilson, and H. S. Ayling, J. Am. Chem. Soc. 64, 1469 (1942).

(FROM THE DATA IN TABLE III)
VARIATION OF THE URANIUM (VI) DISTRIBUTION COEFFICIENT IN
PENTAETHER WITH THE TTA CONCENTRATION.

FIG. 1

MU 462

161001

(FROM THE DATA IN TABLE V)
VARIATION OF THE URANIUM(VI) DISTRIBUTION COEFFICIENT IN
CYCLOHEXANONE WITH THE TTA CONCENTRATION.

MU 463

FIG. 2

(6109)

M/L HNO_3 IN THE AQUEOUS PHASE AT EQUILIBRIUM
VARIATION OF THE URANIUM(VI) DISTRIBUTION COEFFICIENT
IN PENTAETHER WITH THE NITRIC ACID CONCENTRATION.
(FROM THE DATA IN TABLE III)

FIG. 3

MU 464

16110

(FROM THE DATA IN TABLE V)
VARIATION OF THE URANIUM (VI) DISTRIBUTION COEFFICIENT IN
CYCLOHEXANONE WITH THE NITRIC ACID CONCENTRATION

FIG. 4

MU 465

16411

(FROM THE DATA IN TABLE VII)
VARIATION OF THE URANIUM (VI) DISTRIBUTION COEFFICIENT
IN HEXONE WITH THE TTA CONCENTRATION.
MU 466

FIG. 5

161121

86C 027

(FROM THE DATA IN TABLES IV AND VI)
VARIATION OF THE URANIUM (VI) DISTRIBUTION COEFFICIENT
WITH THE HYDROGEN ION CONCENTRATION.

MU 467

FIG. 6

161131

(FROM THE DATA IN TABLES IV AND VI)
VARIATION OF THE URANIUM (VI) DISTRIBUTION COEFFICIENT
WITH NITRATE ION CONCENTRATION.

MU 468

FIG. 7

860 C29

SUMMARY OF CONSTANTS
AT VARIOUS NITRIC ACID CONCENTRATIONS

FIG. 8 MU 469

866 030