LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

LA-AC-6824

NOT FOR PUBLIC RELEASE - OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

EXCLUDED FROM BE MADE OFFICIAL

JAN 15 1968

A Hydrodynamic Model of He II Near T_{λ}^*

83550

Philip J. Bendt

Los Alamos Scientific Laboratory, University of California

Los Alamos, New Mexico

MASTER

The temperature dependence of the torque on a Rayleigh disk in rotating helium (Fig. 1) has remained unexplained since the observations were published by J. R. Pellam⁽¹⁾ in 1960. Pellam suspended a 3 mm disk at 45° to the tangential velocity field in a vessel of liquid He⁴, which was rotated at 2 rpm. P. P. Craig⁽²⁾ has analyzed the hydrodynamics of Rayleigh disk experiments in helium, and concludes that the torque in steady state experiments is probably due to Helmholtz flow. In Helmholtz flow, torque results from Bernoulli pressure at one stagnation point only, on the front surface of the disk; eddies which peel off the edges of the disk flow into the region behind it.

We will describe a model of He II, which may explain Pellam's observations. In this model, He II consists of two interpenetrating components (A and B), both of which rotate with the vessel.⁽³⁾ A moves past the disk with Helmholtz flow, and applies the observed torque, while B does not apply a torque. It is reasonable to assume the flow of component B is constrained, so that it does not respond freely to hydrodynamic pressure. If component B yields only slightly

* Work performed under the auspices of the U. S. Atomic Energy Commission.

RELEASED FOR ANNOUNCEMENT

IN NUCLEAR SCIENCE ABSTRACTS

LASL-AEC-OFFICIAL

LASL-AEC-OFFICIAL

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

under hydrodynamic pressure, volume elements of B will remain very nearly in place (in rotating coordinates) until they make contact with the surface of the disk. The velocity of B will then be uniform across the surface of the disk, and there will be no stagnation point or Bernoulli pressure to apply torque.⁽⁴⁾

To explain the temperature dependence of component B, we propose that part of the entropy of He II is in the form of vortex rings. This modification of the Landau model was suggested by R. P. Feynman,⁽⁵⁾ who wrote "If rotons are the smallest ring vortices, and those of lowest energy Δ , then there are states of high energy corresponding to larger rings.... For a long line there are an enormous number of shapes and orientations available.... It may be likened to a chain of a finite number of links. Adding one link requires an energy ϵ , say of order Δ , but increases the number of orientations by some factor, asymptotically, say s . In equilibrium then, the number of chains of $n + 1$ links is a factor $s \exp(-\epsilon/kT)$ times the number with n links."

We have tried to make Feynman's description semi-quantitative by writing a very approximate spectrum for the vortex rings. Lacking a better value, we use $\Delta(t)$ in the Boltzmann probability.⁽⁶⁾ The variable n is then approximately the length of a vortex ring in units of the circumference of a roton. We also use what appears to be the asymptotic value of s for small rings as well as large rings, and write

$$I(n) = \alpha(t) \exp(n\Delta_{\lambda}/kT_{\lambda}) \exp(-n\Delta/kT). \quad (1)$$

$\alpha(t)$ is a normalization factor, to be evaluated by Eqs. (2) or (3) below. The first exponential is $s^n = (13.2)^n$. The value of s was determined from the criterion that vortex rings of all lengths would have the same probability at T_λ . The helium atoms are then almost entirely in the largest rings. The innumerable orientations of these very long loops become the uncorrelated disorder which exists above T_λ . The relative normal fluid density is equated to

$$\rho_n/\rho = \sum_{n=1}^{\infty} n I(n) \quad (2)$$

if n has only integer values, or to

$$\rho_n/\rho = I(1) + \int_m^{\infty} n I(n) dn \quad (3)$$

where $m \geq 1$, if n is a continuous variable. $I(1)$ gives the relative density of rotons, $I(2)$ of vortex rings having twice the circumference, etc. The vortex rings are much smaller than the Rayleigh disk, except immediately below the lambda transition. Since vortex rings are unimportant below 1.6°K , and the Landau model breaks down above this temperature, ⁽⁷⁾ neither neutron scattering data nor specific heat measurements preclude consideration of this model. The hydrodynamic measurements (viscosity, drag coefficient, and mutual friction) are unable to distinguish between this model and Landau's.

We show by dashed lines in Fig. 1 the relative superfluid density ρ_s/ρ , as well as the fractions of normal fluid contributed by the first few terms in Eq. (2). ⁽⁸⁾ It is apparent that curve 1,

which is the sum of the superfluid and roton densities, is in reasonable agreement with Pellam's experimental points. We therefore identify the superfluid and rotons with component A, and the remaining density, which consists of larger vortex rings, with component B.

The most striking feature of Pellam's data is the sudden drop to zero torque just below the lambda transition. In this model, both the superfluid and roton densities are zero just below T_λ , the entropy being entirely in the form of vortex rings. If we assume that the flow of vortex rings is constrained, so they do not apply a torque, then the sudden drop of Pellam's data to zero torque is explained.

The superfluid part of rotating helium He II is expected to contain an array of vortex lines, roughly parallel with the vertical axis of rotation.⁽⁵⁾ These lines provide quantization of the superfluid circulation. The lines are attached to the bottom of the vessel, and extend to the top surface of the liquid, where they are attached to the surface-tension film. Since they have an innate tension, the flow of these lines is constrained. However, when the local velocity field of the vortex lines is compared with the relative velocity of the liquid helium and Rayleigh disk (0.37 cm/s), we find that only about 2×10^{-4} of the superfluid is bound to the vortex lines.⁽⁹⁾ Therefore the bulk of the superfluid, as well as the rotons, moves past the Rayleigh disk with Helmholtz flow.

We now discuss a possible mechanism for constraining the flow of vortex rings in rotating helium. The situation can only be solved by quantum mechanics, so the classical description below will only indicate what may happen. Normal fluid viscosity prevents the vortex rings from moving with the local velocity field of superfluid vortex lines. Therefore a Magnus force⁽¹⁰⁾ tends to orient rings which are close to a vortex line, into a plane containing the line. If the vortex rings are large enough, the gradient of the local velocity field results in the attractive Magnus force, on the side of the ring which is nearest to a line, being larger than the repelling Magnus force, on the far side of the ring. We propose that the vortex rings are constrained by becoming attached to superfluid vortex lines. Since rotons are only a few times larger than the interatomic separation, the Magnus force on them will be very small, perhaps zero. They have sufficient kinetic energy to avoid being trapped in the field of the vortex lines.⁽¹¹⁾

This completes the description of our model. We hope these ideas will stimulate further study of the very interesting phenomena discovered by Pellam. An important experiment will be to move the Rayleigh disk through nonrotating helium. To avoid superfluid turbulence, this might be done in a "superfluid wind tunnel,"⁽¹²⁾ with the normal fluid at rest. The disk would then be moved linearly with the same velocity as the superfluid. Our model predicts vortex loops will then contribute to the torque on the disk.

It is a pleasure to acknowledge helpful discussions with Professor J. R. Pellam, Dr. T. R. Roberts, and Dr. T. A. Oliphant.

Figure Caption

Fig. 1. The experimental points were observed by J. R. Pellam, and are proportional to the torque on a Rayleigh disk suspended in a rotating vessel of liquid He^4 . The difference between curve 1 and ρ_s/ρ is the fraction of the density consisting of rotons, according to Eq. (2). The difference between curve 2 and ρ_s/ρ is the fraction due to the first two terms in Eq. (2), etc.

References

- (1) J. R. Pellam, Phys. Rev. Letters 5, 189 (1960); Ninth Inter. Conf. on Low Temp. Physics, Columbus, Ohio, Aug. 31-Sept. 4, 1964, paper CB-8.
- (2) P. P. Craig, Phys. Rev. Letters 13, 708 (1964).
- (3) The components A and B are not the same as the usual division of He II into normal fluid and superfluid.
- (4) A familiar example will clarify what we mean: consider the Rayleigh disk to be like a plane, tilted snowplow moving slowly through new-fallen snow (the analog of component B). The snow does not flow like a liquid, but remains essentially in place until contacted by the plow. The velocity of the snow is uniform across the surface of the plow; there is nothing resembling the stagnation point of liquid flow.
- (5) R. P. Feynman, Progr. in Low Temp. Physics (C. J. Gorter, ed., North-Holland Publishing Co., Amsterdam, 1955) Vol. I, p. 17.
- (6) $\Delta(t)$ is the roton energy at the minimum of the excitation curve, and Δ_λ is the value at T_λ ($\Delta_\lambda/k = 5.6^\circ\text{K}$). See D. G. Henshaw and A. D. B. Woods, Phys. Rev. 121, 1266 (1961).
- (7) K. R. Atkins, Liquid Helium (Cambridge University Press, Cambridge, 1959) pp. 63, 69.
- (8) If Eq. (3) is used, with $m = 1$, the calculated relative-density curves lie slightly below those shown in Fig. 1.

- (9) To be bound to a vortex line, we assume the local velocity of a volume element, due to the vortex line, must be larger than the relative velocity of the liquid helium and Rayleigh disk.
- (10) The Magnus force $\underline{F} = \rho \underline{v} \times \underline{k}$, where ρ is the helium density, $\underline{v}$ is the local superfluid velocity due to the vortex line, and $\underline{k}$ is the circulation around the core of the vortex loop.
- (11) The probability that the small vortex rings ($n = 2, 3$, etc.) become attached to vortex lines may be less than unity, and may be temperature dependent. The spectrum given by Eqs. (1) and (2) is too approximate to draw conclusions.
- (12) P. P. Craig and J. R. Pellam, Phys. Rev. 108, 1109 (1957).

6

Fig. 1