33
-21

2120

LA-4892

MASTER

Analyzing Power in $\text{Si}(\vec{p}, p)$ below 15 MeV

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

Printed in the United States of America. Available from
National Technical Information Service
U. S. Department of Commerce
5285 Port Royal Road
Springfield, Virginia 22151
Price: Printed Copy \$3.00; Microfiche \$0.95

LA-4892

UC-34

ISSUED: March 1972

Analyzing Power in $\text{Si}(\vec{p}, p)$ below 15 MeV

by

R. A. Hardekopf

D. D. Armstrong

P. W. Keaton, Jr.

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

ANALYZING POWER IN $\text{Si}(\vec{p}, p)$ BELOW 15 MeV

by

R. A. Hardekopf, D. D. Armstrong, and P. W. Keaton, Jr.

ABSTRACT

Analyzing powers, A_y , were measured for protons elastically scattered from a natural silicon target in the energy range 5 to 15 MeV. Angular distributions of A_y were obtained at 5.16, 7.50, and 12.50 MeV using a thin ($\sim 0.24 \text{ mg/cm}^2$) target. Excitation functions of A_y were measured with a 6 mg/cm^2 target at several angles. The results indicate that silicon has only limited usefulness as a polarimeter for protons in the energy range of 5 to 15 MeV.

I. INTRODUCTION

Our search for an efficient polarization analyzer for protons has led us to investigate the analyzing power of silicon in the energy range 5 to 15 MeV. There is a significant advantage to the use of silicon as an analyzer. When a silicon detector is used as the second scatterer, pulses are produced in both the analyzing detector and the side detector. A coincidence between these pulses can be performed to reduce background, and the pulses can be summed to give the total energy of the proton incident on the analyzer. This allows the use of a thick analyzer without appreciable loss in resolution.

The use of a silicon detector as an analyzer was first proposed by Miller¹ in 1965. Although the analyzing power of silicon was known at only a few energies, several groups have utilized extrapolation of existing data to design, build, and test silicon polarimeters. Martin and Levesque² chose an analyzing angle of $148^\circ \pm 16.5^\circ$, based on 10.5 and 14.5 MeV polarization data of Rosen et al.³ and optical model calculations. Bangert et al.⁴ reported measurements of $\text{Si}(p, p)$ polarization at 130° and found useful values for a polarimeter from 8.3 to 10.9 MeV at this angle. Frois et al.⁵ have tested

a polarimeter using forward scattering angles for energies greater than 20 MeV. In view of the ease with which the analyzing power in elastic scattering can be determined with a beam from a polarized ion source, it seemed reasonable for us to obtain more data on silicon before evaluating the usefulness of a silicon polarimeter.

II. PROCEDURE AND RESULTS

The data reported here were obtained by bombarding a natural silicon target with 5 to 15 MeV polarized protons from the Los Alamos Scientific Laboratory's (LASL) polarized ion source⁶ and tandem accelerator. The beam polarization, obtained by the quench-ratio method,⁷ was about 0.92.

To supplement the data of Rosen et al.³ at 10.5 and 14.5 MeV, we measured angular distributions of the analyzing power, A_y (Basel Convention with y-axis along $\vec{k}_{in} \times \vec{k}_{out}$), at 5.16, 7.50, and 12.50 MeV using a $1\text{-}\mu\text{m}$ thick ($\sim 0.24 \text{ mg/cm}^2$) silicon crystal. These distributions are shown in Fig. 1 along with the earlier data. Relative yields indicated in general that the differential cross section was a minimum around 90 to 120° (as low as 2 mb/sr) and increased at back angles, often by an order of magnitude over the minimum. Therefore, only the peaks

in the analyzing power near 55° and near 155° seemed to be useful candidates for an analyzer, and we investigated these angles in more detail. Using a 25- μm thick target ($\sim 6 \text{ mg/cm}^2$), we obtained values for A_y at 140° , 150° , and 160° in 500-keV steps from 5.0 to 15.0 MeV. These data are shown in Fig. 2. We also measured analyzing powers near 55° and 155° as a function of energy, such that the energy spread in the target (from 330 keV at 5 MeV to 150 keV at 15 MeV) was greater than or equal to the energy steps. The resulting analyzing powers and relative yields are shown in Fig. 3. Numerical values for all data are listed in Tables I-III. Although statistical errors for A_y were generally less than ± 0.01 , we assign errors of ± 0.02 to all the data because of the survey nature of the experiment.

III. CONCLUSION

From the data in Figs. 1 and 2, one might conclude that the analyzing power varies slowly with energy except near 7.5 MeV. Figure 3, however, shows that considerable fluctuations in the analyzing power and cross section exist from 5 to 15 MeV, and it is clearly inappropriate to describe the reaction by an optical model in this energy range. Cross-section measurements for $^{28}\text{Si}(p,p)$ and $^{28}\text{Si}(p,p')$ from 12 to 15 MeV by Shotter et al.⁸ also indicated intermediate structure in this region of compound nucleus excitation.

Although there are energy regions of a few MeV where the average analyzing power of silicon is fairly large, it is clear that the fluctuations severely limit the usefulness of a silicon polarimeter below 15 MeV. Such a polarimeter would be practical only over carefully selected energy and angular ranges and would be very sensitive to small shifts in energy. For the cases where these limitations are acceptable, the present data should be valuable as a guide in polarimeter design and implementation.

REFERENCES

1. D. W. Miller, "Experiments on Polarization in Nuclear Stripping Reactions," in Proc. 2nd Intern. Symp. on Polarization Phenomena of Nucleons, Karlsruhe, September 6-10, 1965, P. Huber and H. Schopper, Eds. (Birkhäuser Verlag Basel, 1966), pp. 410-432.
2. J. P. Martin and R. J. A. Lévesque, "A Double Scattering Proton Polarimeter with High Resolution and Good Efficiency," Nucl. Instr. and Meth. **84**, 211-217 (1970).
3. J. Rosen, J. G. Beery, A. S. Goldhaber, and E. H. Auerbach, "Elastic Scattering of 10.5- and 14.5-MeV Polarized Protons from Nuclei and the Optical Model Potential at Intermediate Energies," Ann. Phys. **34**, 96-152 (1965).
4. R. Bangert, B. Gonsior, P. Rösner, M. Roth, B. Steinmetz, and A. Strömich, "Polarization Experiments on Si and Si-Polarimeters," in Polarization Phenomena in Nuclear Reactions, H. H. Barschall and W. Haeblerli, Eds., Proc. 3rd Intern. Symp., Madison, Wisconsin, 1970 (University of Wisconsin Press, 1971) pp. 646-648.
5. B. Frois, J. Bircha'll, R. Lamontagne, R. Roy, and R. J. Slobodrian, "A High Efficiency Proton Polarimeter with Silicon Detector Analyser," Nucl. Instr. and Meth. **96**, 431-436 (1971).
6. J. L. McKibben, G. P. Lawrence, and G. G. Ohlsen, "Performance of the LASS Polarized Source," in Polarization Phenomena in Nuclear Reactions, H. H. Barschall and W. Haeblerli, Eds., Proc. 3rd Intern. Symp., Madison, Wisconsin, 1970 (University of Wisconsin Press, 1971) pp. 828-831.
7. G. G. Ohlsen, J. L. McKibben, G. P. Lawrence, P. W. Keaton, Jr., and D. D. Armstrong, "Precise Proton-Polarization Standards Determined with a Lamb-Shift Ion Source Incorporating a Nuclear Spin Filter," Phys. Rev. Lett. **27**, 599-602 (1971).
8. A. C. Shotter, P. S. Fisher, and D. K. Scott, "Intermediate Structure and the Reaction $^{28}\text{Si}(p,p')^{28}\text{Si}^*$," Nucl. Phys. **A159**, 577-597 (1970).

Fig. 1. Angular distributions of analyzing power, A_y , for $\text{Si}(p,p)$. The dots are from the present work and the squares are from Ref. 3. The broken lines are hand-drawn interpolations of the data.

Fig. 2. Analyzing power for $\text{Si}(p,p)$ near 150° . The numbers on the graph are incident proton energies in MeV. The lines are hand-drawn interpolations of the data.

Fig. 3. Analyzing power (dots) and relative yield (squares, arbitrary units) for $\text{Si}(\vec{p},p)$ as a function of proton energy. The lines are drawn between data points for clarity.

TABLE I
ANGULAR DISTRIBUTIONS OF $\text{Si}(\text{p},\text{p})$ ANALYZING POWER

θ_{lab} (deg)	$\theta_{\text{c.m.}}$ (deg)	$E_{\text{p}} = 5.16 \text{ MeV}$	$E_{\text{p}} = 7.50 \text{ MeV}$	$E_{\text{p}} = 12.50 \text{ MeV}$
		A_{y}	A_{y}	A_{y}
30.0	31.0	-0.04	0.05	-0.1
40.0	41.3	-0.07	0.10	-0.01
50.0	51.6	-0.05	0.22	0.55
60.0	61.8	0.02	0.30	0.23
70.0	71.9	0.06	0.31	-0.14
80.0	82.0	-0.09	0.32	-0.36
90.0	92.1	-0.20	0.34	-0.56
100.00	102.0	-0.14	0.43	-0.50
110.0	111.9	0.02	0.50	-0.05
115.0	116.9	0.11		
120.0	121.8	0.21	0.52	-0.26
130.0	131.6	0.35	0.22	-0.16
140.0	141.3	0.39	-0.17	0.29
150.0	151.0	0.37	-0.40	0.65
160.0	160.7	0.28	-0.36	0.62

TABLE II
 $\text{Si}(\text{p},\text{p})$ ANALYZING POWERS AT 140° , 150° , AND 160°

E_{p} (MeV)	$\theta_{\text{lab}} \rightarrow 140^\circ$	150°	160°
5.0	0.24	0.26	0.21
5.5	0.48	0.39	0.26
6.0	0.00	-0.03	-0.03
6.5	0.34	0.26	0.19
7.0	0.68	0.56	0.40
7.5	0.31	0.08	-0.03
8.0	-0.05	0.00	0.01
8.5	0.58	0.33	0.17
9.0	0.65	0.67	0.58
9.5	0.68	0.58	0.42
10.0	0.57	0.65	0.56
10.5	0.38	0.44	0.43
11.0	0.42	0.52	0.51
11.5	0.12	0.40	0.56
12.0	0.14	0.37	0.53
12.5	0.49	0.71	0.65
13.0	0.35	0.58	0.78
13.5	0.26	0.45	0.55
14.0	-0.07	0.58	0.66
14.5	0.36	0.64	0.51
15.0	-0.29	0.45	0.92

TABLE III
Si($\vec{p},p$) EXCITATION FUNCTION FOR A_y

$\theta_{lab} = 55^\circ$			$\theta_{lab} = 155^\circ$			$\theta_{lab} = 160^\circ$		
E_p (MeV)	A_y	Yield (relative units)	E_p (MeV)	A_y	Yield (relative units)	E_p (MeV)	A_y	Yield (relative units)
9.70	0.05	47.1	5.00	0.24	61.3	10.80	0.31	34.6
9.90	0.21	42.4	5.25	0.36	81.3	10.95	0.57	27.7
10.10	-0.07	27.9	5.50	0.32	136.1	11.10	0.31	11.3
10.25	0.13	27.5	5.75	-0.17	120.6	11.25	0.55	28.1
10.40	-0.06	33.3	6.00	-0.03	67.5	11.40	0.64	16.2
10.55	-0.40	43.5	6.25	0.00	49.2	11.55	0.49	13.0
10.65	0.22	42.5	6.50	0.22	52.3	11.70	0.53	28.8
11.00	0.41	34.9	6.75	0.23	40.8	11.85	0.63	32.4
11.15	0.09	30.8	7.00	0.48	66.0	12.00	0.60	18.0
11.30	0.24	19.0	7.25	0.40	94.9	12.15	0.00	22.0
11.45	0.37	31.8	7.50	0.01	59.9	12.25	-0.07	25.9
11.60	0.28	25.8	7.65	-0.50	37.7	12.30	0.05	27.5
11.75	0.40	25.9	7.80	-0.54	49.1	12.45	0.48	20.6
11.90	0.58	29.4	7.95	-0.24	37.0	12.65	0.68	25.9
12.05	0.53	30.2	8.10	0.34	28.8	12.80	0.77	25.2
12.20	0.64	27.8	8.25	0.28	39.0	12.95	0.84	15.1
12.35	0.51	24.5	8.40	0.10	77.4	13.10	0.75	11.6
12.50	0.52	24.4	8.55	0.42	56.6	13.25	0.22	18.5
12.65	0.33	25.7	8.70	0.72	43.9	13.40	0.59	18.5
12.80	0.37	14.7	8.85	0.61	15.4	13.55	0.49	27.3
12.95	0.60	18.4	9.00	0.64	11.3	13.70	0.59	36.7
13.10	0.62	22.9	9.15	0.57	13.1	13.85	0.38	32.3
13.25	0.40	18.2	9.30	0.26	19.5	14.00	0.68	29.5
13.40	0.34	14.3	9.45	0.44	35.2	14.15	0.66	27.9
13.55	0.49	12.9	9.60	0.49	22.8	14.30	0.55	18.4
13.70	0.51	19.3	9.75	0.29	49.8	14.45	0.47	15.6
13.85	0.46	27.6	9.90	0.50	36.7	14.60	0.87	11.0
14.00	0.42	22.0	10.05	0.64	18.2	14.75	0.74	16.4
14.15	0.30	27.0	10.20	0.63	23.0	14.90	0.85	17.6
14.30	0.44	28.1	10.35	0.49	35.0	15.05	0.93	10.8
14.45	0.37	25.4	10.50	0.46	36.6	15.20	0.52	6.0
14.60	0.32	17.4	10.65	0.13	51.3			
14.75	0.27	19.2						
14.90	0.06	19.0						
15.05	0.17	15.1						
15.20	0.18	11.7						