

UCLA - 12 - 8 37

MASTER

UNIVERSITY OF CALIFORNIA, LOS ANGELES

RECEIVED BY DTIE OCT 18 1971



LABORATORY OF NUCLEAR MEDICINE AND RADIATION BIOLOGY

CONTRACT NO. AT (04-1) GEN-12

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

COMPARATIVE MYOCARDIAL UPTAKE AND CLEARANCE
CHARACTERISTICS OF POTASSIUM AND CESIUM

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

Norman D. Poe, M.D.

Department of Radiological Sciences,
Center for Health Sciences
and
Laboratory of Nuclear Medicine and Radiation Biology
University of California, Los Angeles, California

Running Title: Myocardial Uptake of Potassium and Cesium

Address Correspondence to:

Norman D. Poe, M.D.
Division of Nuclear Medicine
UCLA Medical Center
Los Angeles, California 90024

This study was supported by Contract AT(04-1) GEN 12 between the U.S. Atomic Energy Commission and the University of California

Myocardial scanning is potentially an important clinical tool for localization and quantitation of damaged cardiac tissue. Early attempts at scanning with potassium analogs, while encouraging, did not prove satisfactory because of the unsuitable gamma energies of available radionuclides (1,2). Recently, two new radionuclides have become accessible: potassium-43 and cesium-129 (3,4). Both have relatively short half-lives and dominant gamma emissions in the iodine-131 range. As these two nuclides can be produced by the newly installed UCLA Biomedical Cyclotron, laboratory investigations were undertaken to compare the uptake and clearance characteristics of potassium and cesium after intravenous and coronary arterial injection. Observations were made with the prospect of utilizing one or both of these agents for quantitative measurements of regional myocardial blood flow and function.

MATERIALS AND METHODS

Experiments were performed using potassium-42 and cesium-131. When both radionuclides were to be compared, they were mixed and administered as a single solution. In vivo clearance data was obtained in medium sized mongrel dogs under pentobarbital anesthesia, 30 mg/Kg. Recordings were made with the animals in the right lateral decubitus position with well shielded probes placed over the head and precordium. Each probe was connected to a dual spectrometer-recorder system. For the intravenous experiments, quantitative injections were made through indwelling jugular vein catheters. Blood samples were drawn from an indwelling catheter in the opposite jugular. Continuous urine samples were collected from an indwelling bladder catheter. For the intracoronary injections, the left chest was opened and a 27 gauge needle with fine polyethylene tubing

attached was inserted into the anterior descending coronary artery. The tubing was preloaded with tracer in a 0.1 cc volume and kept outside the detector field. At the appropriate time, a bolus injection was made by flushing the tubing with saline. All observations in dogs were made with 50-100 μ Ci of the two radionuclides.

In vitro clearance studies were made in rats (approximately 300 gm) which were anesthetized with intraperitoneal pentobarbital. A femoral vein was exposed and then 15 μ Ci doses of radionuclides were quantitatively injected. Hearts were removed at selected intervals and counted in a well counter.

RESULTS

INTRAVENOUS

Myocardium: A definite time differential is found between maximum uptake of potassium and cesium (Fig. 1). Potassium reaches a plateau in the heart (with slight chest wall contribution) within 5-20 minutes, then decreases during the next 90 minutes. A more gradual negative phase follows. The $T_{1/2}$ based on this latter curve in 3 animals ranges between 5.6 and 7.7 hours (average 6.5 hours). In contrast, cesium initially decreases reaching a nadir at 5 minutes, then peaks between 120 and 180 minutes before a very gradual clearance phase develops. Although the 3 animals were followed for 5 hours, a clear-cut $T_{1/2}$ could not be identified. Further measurements were obtained at 24 hours. Based on this data, the $T_{1/2}$ ranges from 16.5 to 43.0 hours (average 27.3 hours). One animal followed for a week still retained 10% of the peak level.

Head: Concentration increases over the head for the first 10 minutes with both tracers. Then a steady level of uptake is maintained for the

following 180 minutes. The head represents a combination of both skeletal muscle and blood pools.

Blood: A marked difference in blood clearance is seen between potassium and cesium (Fig. 2). At 2 minutes 42% of the cesium remains in the blood but only 22% of the potassium remains. At 1 hour these values drop to 3.8% and 1.6% respectively. The disappearance curves are multiphasic, probably containing 3 basic components representing ion exchange in the intravascular, interstitial and intracellular compartments.

Urine: Cesium clearance only was measured. In 3 animals 3.5% to 5% of the injected dose was excreted in 5 hours.

In Vitro: The data in 12 rats is summarized in Table 1. These values confirm the myocardial uptake pattern described in dogs above. In the rat the heart represents only 0.35% of the body weight in contrast to 0.9% in the dog.

INTRA-ARTERIAL

The dichotomy between potassium and cesium is further exemplified in 3 animals after bolus injection of the two radionuclides into the anterior descending coronary artery (Fig. 3). Potassium is highly extracted with almost three times the efficiency of cesium (average 71% vs. 22%) on the first circulation through the coronary capillary bed. Likewise, potassium is cleared more rapidly from the heart. Based on the slope of the curve from 10 to 60 minutes, the T 1/2 for potassium is 78 minutes and for cesium 390 minutes.

DISCUSSION

The above observations show distinct differences in the myocardial concentration and blood clearance patterns between potassium-42 and

cesium-131. These findings are in accord with reports by previous investigators (2,3,5,6). Potassium is rapidly cleared from the blood and concentrated by the myocardium. Its high extraction on a single circulation through the vascular bed makes it a suitable indicator for coronary blood flow measurements. Conversely, the blood clearance and myocardial uptake of cesium are slower and the intracellular retention much longer even though the maximum myocardial concentration of the two analogs is approximately the same. The differences in rate probably can be attributed to the inability of the cell membrane transport mechanism to handle the cesium ion with the efficiency of the potassium ion. Although the early reports of Carr and others (2,7) leave little doubt that radiocesium is perfectly adequate for qualitatively localizing sites of established myocardial infarcts, the behavior of this ion during acute and evolving infarction is not established. It has been demonstrated that the egress of potassium from ischemic myocardium is dependent both on the magnitude and duration of the ischemia (8,9). Presumably cesium will behave similarly but until the effects of ischemia on the kinetics of this ion have been established, quantitative measurements with cesium during the developing stages of infarction are not justified.

The less efficient extraction of cesium following intracoronary arterial injection is a definite disadvantage in quantitative studies when the radionuclide is administered by this route. At least 75% of the injected dose will be free to recirculate throughout the body and return to the myocardium in proportion to coronary blood flow. Although the myocardium receives only about 5% on the cardiac output, the amount of cesium finally reaching the heart by recirculation could exceed 15% of the amount extracted on the initial injection. Comparative scans

made after intracoronary injection of cesium-131 and technetium-99m labeled particles which have no recirculation do demonstrate substantial and readily detectable concentrations of cesium throughout the myocardium (10). This is not a problem with potassium because the percent available for recirculation is inconsequential.

The whole principle of the use of radiopotassium and related substances for regional flow measurements in damaged myocardium is open to question. Cellular extraction of these ions is not only a function of flow but also of cell membrane integrity. The capability of cells to concentrate potassium can remain nearly unimpaired with flow reductions up to 50% of normal (9). After ischemia has been produced, maintenance or recovery of function is related to the extent and duration of the hypoxia. During the acute and evolving stages of a myocardial infarction, significant inequalities between flow and cell damage very likely coexist. Under these circumstances erroneously low perfusion deficits could conceivably be obtained in the presence of normal perfusion.

After the intravenous administration of potassium, scanning can commence immediately as maximum counts are reached by 5 minutes. For cesium a delay of 1 to $1\frac{1}{2}$ hours is required to attain maximum count rates. The rapid clearance of potassium after intracoronary injection necessitates almost immediate scanning, preferably performed rapidly. Cesium clearance by this route is slow, making scanning possible any time within the first several hours after injection. For myocardial dosimetry calculations with intravenous potassium, the calculated 6.5 hour T $1/2$ should be sufficiently accurate. However, the cesium data was less precise and it is suggested that the 32 hour physical T $1/2$ be utilized.

SUMMARY AND CONCLUSIONS

Potassium-43 and its analog, cesium-129, have recently been suggested as suitable radiopharmaceuticals for myocardial scanning. Studies with potassium-42 and cesium-131 were undertaken in dogs to determine the myocardial turnover characteristics of these two ions. After simultaneous intravenous injection, potassium clears rapidly from the blood, reaches a plateau in the myocardium within 5-20 minutes, and then clears with a $T_{1/2}$ of 6.5 hours. Initial cesium clearance is slower. A peak is reached gradually in the myocardium within 1-3 hours and the clearance $T_{1/2}$ is 27.3 hours. Following intracoronary injection, potassium is extracted with 71% efficiency on a single circulation, but cesium is extracted with only 22% efficiency. It is concluded that cesium should be utilized with caution as a substitute for potassium in quantitative myocardial blood flow and function measurements, but that comparable results can be obtained with either radionuclide in localizing ischemic myocardium.

REFERENCES


1. Carr EA, Beierwaltes WH, Wegst AV, et al: Myocardial scanning with rubidium-86. J Nucl Med 3:76-82, 1962.
2. Carr EA, Gleason G, Shaw J, et al: The direct diagnosis of myocardial infarction by photoscanning after administration of cesium-131. Am Heart J 68:627-636, 1964.
3. Hurley PJ, Cooper M, Reba RC, et al: ^{43}KCl : a new radiopharmaceutical for imaging the heart. J Nucl Med 12:516-519, 1971.
4. Yano Y, Van Dyke D, Budinger TF, et al: Myocardial uptake studies with ^{129}Cs and the scintillation camera. J Nucl Med 11:663-668, 1970.
5. Love WD, Ishihara Y, Lyon LD, et al: Differences in the relationships between coronary blood flow and myocardial clearance of isotopes of potassium, rubidium, and cesium. Am Heart J 76:353-355, 1968.
6. Langer GA, Brady AJ: Potassium in dog ventricular muscle: kinetic studies of distribution and effects of varying frequency of contraction and potassium concentration of perfusate. Circ Res 18:164-177, 1966.
7. McGeehan JT, Rodriguez-Antunez A, Lewis RC: Cesium-131 photoscans. JAMA 204:585-589, 1968.
8. Jennings RB, Sommers HM, Kaltenbach JP, et al: Electrolyte alterations in acute myocardial ischemic injury. Circ Res 14:260-269, 1964.
9. Case RB, Nasser MG, Crampton RS, et al: Biochemical aspects of early myocardial ischemia. Am J Cardiol 24:766-775, 1969.
10. Poe ND: Regional myocardial blood flow measurements using retained radioactive substances: comparison of radiocesium and labeled particles. In preparation.


LEGENDS

Figure 1 After intravenous injection different precordial uptake curves are observed for potassium and cesium. Potassium reaches a peak concentration between 5 and 20 minutes. Cesium concentration is lowest at 5 minutes, then gradually rises to a peak between 120 and 180 minutes. Values are means for 3 animals \pm SD and represent the total activity in the myocardium, the heart blood pool, and the overlying muscle.

Figure 2 Comparative blood clearance curves demonstrate more rapid and efficient removal of potassium from the vascular space and earlier intracellular equilibration. The slower cesium clearance and consequent higher blood levels correlate with the delayed precordial cesium peak seen in Figure 1. Values expressed as means \pm SD.

Figure 3 Myocardial extraction of potassium after a bolus injection directly into a coronary artery is approximately three times that of cesium. This difference indicates that a much larger percentage of cesium is free to recirculate to other regions of the myocardium. (Mean \pm SD)








TABLE 1 Percent Myocardial Uptake in Rats Following Intravenous Injection: Mean (Range) n=3

	<u>5 min</u>	<u>15 min</u>	<u>30 min</u>	<u>60 min</u>
¹³¹ Cs	1.45 (1.34-1.51)	2.13 (1.82-2.31)	1.99 (1.79-2.25)	2.02 (1.90-2.25)
⁴² K	2.54 (2.49-2.59)	2.39 (2.30-2.49)	1.86 (1.76-1.97)	1.04 (0.95-1.10)

ACKNOWLEDGMENTS

The author wishes to acknowledge the technical assistance of Mr. Carl Selin and Mr. Emery Terao.

