AUG 1 1963

3256ys.

MASTER

ORNL-3471

UC-4 - Chemistry

TID-4500 (20th ed., Rev.)

THE ACTIVITY COEFFICIENTS OF HYDROCHLORIC
ACID AND SODIUM CHLORIDE IN
HYDROCHLORIC ACID-SODIUM
CHLORIDE MIXTURES

M. H. Lietzke
R. W. Stoughton

OAK RIDGE NATIONAL LABORATORY

operated by

UNION CARBIDE CORPORATION

for the

U.S. ATOMIC ENERGY COMMISSION

Facsimile Price \$ 1.10

Microfilm Price \$.80

Available from the
Office of Technical Services
Department of Commerce
Washington 25, D. C.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

Contract No. W-7405-eng-26

CHEMISTRY DIVISION

THE ACTIVITY COEFFICIENTS OF HYDROCHLORIC ACID
AND SODIUM CHLORIDE IN HYDROCHLORIC
ACID-SODIUM CHLORIDE MIXTURES

M. H. Lietzke and R. W. Stoughton

Date Issued

JUL 29 1963

OAK RIDGE NATIONAL LABORATORY
Oak Ridge, Tennessee
operated by
UNION CARBIDE CORPORATION
for the
U. S. ATOMIC ENERGY COMMISSION

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

The Activity Coefficients of Hydrochloric Acid
and Sodium Chloride in Hydrochloric
Acid-Sodium Chloride Mixtures

M. H. Lietzke and R. W. Stoughton

ABSTRACT

The activity coefficients of HCl and NaCl in HCl-NaCl mixtures have been computed from literature data. The calculations are based on the observation that at constant ionic strength and temperature the logarithm of the activity coefficient of HCl in HCl-NaCl mixtures varies linearly with NaCl concentration.

The Activity Coefficients of Hydrochloric Acid
and Sodium Chloride in Hydrochloric
Acid-Sodium Chloride Mixtures *

M. H. Lietzke and R. W. Stoughton

In a recent study¹ of the cell

it was shown that at constant temperature and ionic strength the logarithm of the activity coefficient of HBr in HBr-KBr mixtures varies linearly with the molality of KBr. It was also shown that with the parameters describing this variation it is possible to calculate the activity coefficient of the KBr in the mixtures. Previously it had been shown² that at constant ionic strength and temperature the logarithm of the activity coefficient of HCl in HCl-NaCl mixtures varies linearly with NaCl molality. This paper deals with the calculation of the parameters describing the activity coefficients of both HCl and NaCl in HCl-NaCl mixtures by using literature data on the activity coefficient of HCl in pure solutions³ and in NaCl solutions⁴ and of NaCl^{2,2a} in pure solutions. For these calculations the values of the activity coefficient of HCl($\underline{m} = 0.01$) in NaCl solutions⁴ for 0 to 50° were extrapolated to 60°.

In carrying out the computation it was assumed that the ionic strength dependence of the logarithms of the activity coefficients of HCl, of NaCl, and of HCl($\underline{m} = .01$) in NaCl solutions could be described at each temperature

*This work was supported in part by the Office of Saline Water, U. S. Department of the Interior.

by equation (1)

$$\ln \gamma_{\pm} = \frac{-\mathcal{S} \sqrt{I}}{1 + 1.5 \sqrt{I}} + BI + CI^2 + DI^3 \quad (1)$$

In this equation $\mathcal{S}$ is the appropriate Debye-Hückel limiting slope which contains as a factor the square root of the density of water; I is the ionic strength (on a molality basis); and B , C , and D are adjustable parameters. The values of the parameters were determined in each instance by the method of least squares and are given in Table I. For convenience, the value of the parameter in the denominator of the Debye-Hückel expression was taken in all cases to be 1.5.

Table I

Values of the Parameters of Equation (1) Giving the Ionic Strength Dependence of the Activity Coefficients of HCl and NaCl

Temp.	NaCl*			
	$B \times 10^2$	$C \times 10^2$	$D \times 10^3$	σ_{fit}^2 **
0	-3.73852	3.56644	-3.03349	1.5×10^{-5}
10	-0.701016	2.95583	-2.43814	8.9×10^{-6}
20	1.02915	2.81899	-2.50491	7.2×10^{-6}
25	2.41390	2.29369	-1.96781	1.7×10^{-6}
30	2.65655	2.56721	-2.45514	8.1×10^{-6}
40	4.17905	2.11154	-1.97467	9.1×10^{-6}
50	5.53079	1.68468	-1.53191	8.5×10^{-6}
60	6.28909	1.95709	-2.28398	5.3×10^{-6}

* The equations hold to 4 m.

** Variance of fit.

Table I (Contd.)

HCl*

<u>Temp.</u>	<u>B x 10</u>	<u>C x 10²</u>	<u>D x 10³</u>	<u>σ^2_{fit}</u>
0	2.53049	2.33598	-2.02789	1.7×10^{-5}
10	2.47702	2.13754	-1.83070	6.2×10^{-6}
20	2.41086	1.92351	-1.56655	2.9×10^{-6}
25	2.37579	1.82248	-1.45024	2.8×10^{-6}
30	2.22354	3.38269	-6.73504	2.2×10^{-6}
40	2.18233	2.50886	-4.14886	3.2×10^{-6}
50	2.14575	1.82417	-1.81932	1.7×10^{-6}
60	2.13183	0.589944	2.96181	2.1×10^{-6}

* At 25° and below the equations hold to 4 m, above 25° to 2 m.

HCl (m=.01) in NaCl*

<u>Temp.</u>	<u>B x 10</u>	<u>C x 10²</u>	<u>D x 10³</u>	<u>σ^2_{fit}</u>
0	1.39450	4.28343	-6.23885	1.9×10^{-5}
10	1.48713	3.59159	-4.98155	8.5×10^{-6}
20	1.50904	3.33731	-4.62235	7.0×10^{-6}
25	1.56031	2.91288	-3.89802	4.4×10^{-6}
30	1.54654	2.95854	-4.11460	6.5×10^{-6}
40	1.57406	2.57192	-3.64777	5.7×10^{-6}
50	1.60227	2.07142	-2.89576	1.1×10^{-5}
60	1.63973	1.61486	-2.50417	1.7×10^{-5}

* The equations hold to 3 m.

At constant ionic strength and temperature it may be assumed that

$$\ln \gamma_{\text{HCl}} = \ln \gamma_{\text{HCl}}^{\circ} - \alpha m_{\text{NaCl}}, \quad (2)$$

where α is a function only of total ionic strength and temperature. In equation (2) γ_{HCl} is the activity coefficient of HCl in the HCl-NaCl mixture and $\gamma_{\text{HCl}}^{\circ}$ is the activity coefficient of HCl in an HCl solution of the same ionic strength. The parameters in Table I giving the ionic strength dependence of the logarithm of the activity coefficients of HCl and HCl($m=.01$) in NaCl may be used to calculate values of the parameter α in equation (2). These values are given in Table II.

Table II

Values of the Parameter α in Equation (2) for HCl-NaCl Mixtures

$t(^{\circ}\text{C})$	0	10	20	25	30	40	50	60
.02	.22642	.19740	.17980	.16266	.13557	.12163	.10860	.09801
.03	.16953	.14783	.13464	.12183	.10174	.09121	.08141	.07336
.06	.13493	.11775	.10721	.09708	.08153	.07294	.06504	.05834
.11	.12266	.10717	.09753	.08842	.07495	.06683	.05950	.05296
.21	.11518	.10088	.09172	.08333	.07190	.06371	.05657	.04966
.51	.10686	.09424	.08544	.07816	.07057	.06158	.05443	.04629
1.01	.09921	.08839	.07981	.07376	.07000	.06028	.05348	.04478
2.01	.09193	.08290	.07448	.06987	.06597	.05782	.05400	.05054
3.01	.09344	.08405	.07556	.07114	.05691	.05457	.05685	.06721

In order that the values of α shown in Table II could be used to compute the activity coefficient of NaCl in the HCl-NaCl mixtures they were fitted by the method of least squares to equation (3) for ionic strength values above 0.11.

$$\alpha = a_0 + a_1 I + a_2 I^2 + a_3 t + a_4 I t \quad (3)$$

Because the activity coefficient values appeared to be more precise at 25° and below two separate fits of the α -values were made: (1) using the α -values from 0-25°, and (2) using the α -values from 25-60°. The parameters of equation (3) for the two fits are given in Table III.

Table III

Parameters of Equation (3)*

$$\alpha = a_0 + a_1 I + a_2 I^2 + a_3 t + a_4 I t$$

Temp. range	$a_0 \times 10$	$a_1 \times 10^2$	$a_2 \times 10^3$	$a_3 \times 10^3$	$a_4 \times 10^4$
0 - 25°	1.20661	-2.53234	5.32570	-1.22881	1.40681
30 - 60°	1.08192	-2.11277	4.45109	-0.962928	1.88598

* Equation holds over the ionic strength range .11 to 3.01 m at 0-25° and .11 to 2.01 above 25°.

Equation (3) may be used in equation (2) to calculate the values of γ_{HCl} in HCl-NaCl mixtures between 0 and 60° for a total ionic strength as low as 0.11. It may also be used in calculating the activity coefficient of NaCl in the HCl-NaCl mixtures, since according to the rules of partial differentiation it follows from the equation for the total free energy of the system in terms of the partial molal quantities that⁵

$$\left(\frac{\partial \ln a_3}{\partial m_2} \right)_{m_3} = \left(\frac{\partial \ln a_2}{\partial m_3} \right)_{m_2} \quad (4)$$

This equation may be expressed in terms of the variable I :

$$\left(\frac{\partial \ln \gamma_3}{\partial m_2} \right)_{m_3} = \left(\frac{\partial \ln \gamma_3}{\partial I} \right)_{m_3} = \left(\frac{\partial \ln \gamma_2}{\partial m_3} \right)_{m_2} = \left(\frac{\partial \ln \gamma_2}{\partial I} \right)_{m_2} \quad (5)$$

Upon differentiating⁶ equation (2) with respect to the total ionic strength I at constant m_2

$$\left(\frac{\partial \ln \gamma_2}{\partial I} \right)_{m_2} = \frac{d \ln \gamma_2^0}{d I} - \alpha - m_3 \frac{d\alpha}{dI} \quad (6)$$

Equation (6) may be inserted into equation (5) and the latter integrated with respect to I between the limits m_3 and $m_2 + m_3$; whence

$$\begin{aligned} \ln \gamma_3 (m_2 + m_3) - \ln \gamma_3^0 (m_3) &= \ln \gamma_2^0 (m_2 + m_3) - \ln \gamma_2^0 (m_3) \\ &- \int_{I=m_3}^{I=m_2 + m_3} \alpha dI - \int_{I=m_3}^{I=m_2 + m_3} m_3 \frac{d\alpha}{dI} dI \end{aligned} \quad (7)$$

The second term on the left and the first two terms on the right of equation (7) may be evaluated by the use of equation (1) and the parameters given in Table I for the pure components HCl and NaCl. The substitution of equation (3), which gives the ionic strength and temperature dependence of α , into equation (7) permits the evaluation of the remaining integrals and hence the calculation of the activity coefficient of NaCl in HCl-NaCl mixtures. Values of the activity coefficients of HCl and NaCl in pure solution and in a 50-50% mixture at 0, 10, 25, 40, and 60° are given in Table IV.

As expected, (1) at any temperature and ionic strength the activity coefficients are in the order (greatest to least) γ_{HCl}^0 , $\gamma_{\text{HCl}}^{(m_2 = m_3)}$, $\gamma_{\text{NaCl}}^{(m_2 = m_3)}$, γ_{NaCl}^0 ; (2) at any temperature the changes of activity coefficients with ionic strength are in the same order.

Table IV

The Activity Coefficients of HCl and NaCl in Pure
Solution and in a 50-50% Mixture

$t = 0^{\circ}$

	$m_{\text{HCl}} = I$	$m_{\text{HCl}} = m_{\text{NaCl}} = \frac{1}{2} I$	$m_{\text{NaCl}} = I$
I	γ_{HCl}	γ_{HCl}	γ_{NaCl}
.2	.780	.771	.749
.5	.777	.756	.704
1.0	.840	.799	.694
1.5	.943	.878	.714
2.0	1.078	.984	.748
2.5	1.247	1.114	.793
3.0	1.455	1.266	.843

$t = 10^{\circ}$

.2	.775	.767	.748	.737
.5	.769	.750	.706	.679
1.0	.826	.790	.700	.648
1.5	.921	.865	.722	.642
2.0	1.046	.964	.759	.650
2.5	1.202	1.085	.805	.668
3.0	1.391	1.225	.856	.692

$t = 25^{\circ}$

.2	.768	.762	.746	.736
.5	.758	.743	.705	.682
1.0	.809	.780	.703	.656
1.5	.895	.850	.728	.656
2.0	1.008	.942	.767	.668
2.5	1.148	1.054	.815	.688
3.0	1.318	1.182	.866	.715

$t = 40^{\circ}$

.2	.758	.753	.740	.732
.5	.743	.731	.699	.679
1.0	.785	.762	.698	.657
1.5	.861	.824	.722	.660
2.0	.960	.904	.757	.675

$t = 60^{\circ}$

.2	.745	.741	.730	.723
.5	.723	.715	.689	.673
1.0	.754	.737	.686	.654
1.5	.818	.790	.710	.661
2.0	.907	.864	.749	.680

References

- (1) M. H. Lietzke and R. W. Stoughton, J. Phys. Chem., in press.
- (2) H. S. Harned and B. B. Owen, "The Physical Chemistry of Electrolytic Solutions", Chap. 14 and Appendix A, Third Edition, Reinhold Publ. Corp., New York, N. Y., 1958.
- (2a) R. A. Robinson, Trans. Farad. Soc. 35, 1222 (1939).
- (3) R. S. Greeley, William T. Smith, Jr., M. H. Lietzke, and R. W. Stoughton, J. Phys. Chem. 64, 1445 (1960); H. S. Harned and R. W. Ehlers, J. Am. Chem. Soc. 55, 2179 (1933).
- (4) H. S. Harned and G. E. Mannweiler, Ibid 57, 1873 (1935).
- (5) N. Bjerrum, Z. physik. Chem. 104, 406 (1923).
- (6) H. A. C. McKay, Trans. Farad. Soc. 51, 903 (1955).

ORNL-3471
UC-4 - Chemistry
TID-4500 (20th ed., Rev.)

INTERNAL DISTRIBUTION

1. Biology Library	51. C. E. Larson
2-3. Central Research Library	52-76. M. H. Lietzke
4. Reactor Division Library	77. S. Lindenbaum
5-6. ORNL - Y-12 Technical Library	78. H. F. McDuffie
Document Reference Section	79. R. M. Rush
7-41. Laboratory Records Department	80. C. S. Shoup, Jr.
42. Laboratory Records, ORNL R.C.	81. M. J. Skinner
43. P. B. Bien	82-106. R. W. Stoughton
44. R. E. Biggers	107. J. A. Swartout
45. C. F. Coleman	108. E. H. Taylor
46. Wallace Davis, Jr.	109. A. M. Weinberg
47. F. F. Dyer	110. T. H. Davies (consultant)
48. A. A. Grau	111. D. F. Hornig (consultant)
49. Donald Jacobs	112. I. Perlman (consultant)
50. K. A. Kraus	113. J. D. Roberts (consultant)

EXTERNAL DISTRIBUTION

114. Research and Development Division, AEC, ORO
115-623. Given distribution as shown in TID-4500 (20th ed., Rev.) under Chemistry category