October 5, 1973

Dr - 380

17-
10-23-73

23,718

THE ENTHALPY OF SOLUTION OF
URANIUM IN NITRIC ACID

James D. Navratil

Franklin L. Oetting

DOW CHEMICAL U.S.A.
ROCKY FLATS DIVISION
P. O. BOX 888
GOLDEN, COLORADO 80401

U. S. ATOMIC ENERGY COMMISSION
CONTRACT AT(29-1)-1106

MASTER

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

Printed in the United States of America
Available from the
National Technical Information Service
U. S. Department of Commerce
Springfield, Virginia 22151
Price: Printed Copy \$4.00 Microfiche \$0.95

Printed
October 5, 1973

RFP-2097
UC-4 CHEMISTRY
TID-4500-R60

**THE ENTHALPY OF SOLUTION OF
URANIUM IN NITRIC ACID**

James D. Navratil

Franklin L. Oetting

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

**DOW CHEMICAL U.S.A.
ROCKY FLATS DIVISION
P. O. BOX 888
GOLDEN, COLORADO 80401**

SUBJECT DESCRIPTORS

Nitric Acid
Thermodynamics
Uranium

Prepared under Contract AT(29-1)-1106
for the
Albuquerque Operations Office
U. S. Atomic Energy Commission

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

CONTENTS

Abstract	1
Introduction	1
Summary and Conclusions	1
Experimental	1
Reagents	1
Calorimeter	1
Procedure	2
Results and Discussion	2
References	4

THE ENTHALPY OF SOLUTION OF URANIUM IN NITRIC ACID

James D. Navratil and Franklin L. Oetting

Abstract. The enthalpies of solution of uranium in 15.9 and 12.7M HNO₃ were measured calorimetrically at 25.0°C as -250.8 ± 3.3 and -242.0 ± 0.9 Kcal·mol⁻¹, respectively. Calorimetric measurements were also determined for the heat of solution of uranium in nitric acid at 38.0°C, in nitric acid containing uranium, and for different uranium-to-nitric acid ratios.

INTRODUCTION

Nitric acid (12.7M) at ~60°C is presently used in production operations to clean the surface of uranium metal. In the pickling process the generated nitrogen oxides are scrubbed, and control of the exothermic reaction is necessary to ensure that the NO_x generated does not exceed the capacity of the scrubbing system. Thus, the reaction enthalpies were measured under a variety of conditions to facilitate proper design of cooling coils for reaction control.

SUMMARY AND CONCLUSIONS

The enthalpy of solution of uranium (0.24-0.36 g) at 25.0°C in 15.9 and 12.7M HNO₃ (100 ml) was determined with an LKB reaction calorimeter as -250.8 ± 3.3 and -242.0 ± 0.9 Kcal·mol⁻¹, respectively. The latter value was unchanged using 12.7M HNO₃ containing 4.5 g/l uranium, and went to -249.2 ± 1.9 and -251.4 ± 0.7 Kcal·mol⁻¹ when the temperature was increased to 38.0°C and the uranium increased from 0.24-0.36 to 0.45 g per 100 ml of nitric acid, respectively. The uranium reaction times were too erratic for correlation with the experimental conditions. Analysis of the nitrogen oxides generated in the uranium-nitric acid reaction are reported as well as the recommended operating conditions for the uranium-nitric acid pickling operation.

EXPERIMENTAL

Reagents

The uranium powder was obtained from Alfa Products, Division of Ventron Corporation. The impurities were determined by spark source mass spectroscopy and emission spectroscopy prior to the heat measurements. The former analysis was also repeated after the measurements were taken. Since there was no change in analysis, all three determinations were averaged and are (ppm): Al, 20; C, 210; Ca, 200; Cl, 16; F, 14; Fe, 200; Mo, 15; N, 3200; Nb, 30; Ni, 125; O, 295; Si, 70; and Th, 74. The other 25 elements detected were <84 ppm total. X-ray diffraction confirmed that the uranium was all in the alpha phase and detected traces of UO₂ and UN. The percent uranium (93.078) was taken by difference of the total impurities (69217 ppm) where C, Ca, Cl, F, N, O, and Si were assumed to be chemically combined with the uranium.

Allied Chemical, B&A reagent ACS (70.0-71.0%) nitric acid (15.9M) was used in the experiments and to prepare 12.7M HNO₃ (with distilled water).

Calorimeter

A modified LKB reaction calorimeter of the isoperibol type was used for the measurements. Details of the description and operation of the calorimeter may be found elsewhere.¹ The precision and accuracy of the calorimeter was determined by one standard fast reaction and a known slow reaction. The enthalpy of the calorimetric standard reaction, the neutralization of TRIS (tris-hydroxymethylaminomethane) SRM 724 in excess 0.1M hydrochloric acid ($\Delta t \approx 1$ minute), was determined under conditions recommended by the National Bureau of Standards. Our value of -7108 ± 4 cal·mol⁻¹ is in excellent agreement with reported values.¹

A slower known reaction ($\Delta t \cong 24$ minutes) was also measured since it had a comparable reaction time to the uranium-nitric acid reaction time. The enthalpy of dissolution of beryllium (0.12 g/l) in 1.0-1.2M sulfuric acid was determined at 25.0°C as $-91.814 \pm 0.085 \text{ Kcal} \cdot \text{mol}^{-1}$. This value is only in fair agreement with the reported value of $-92.83 \pm 0.11 \text{ Kcal} \cdot \text{mol}^{-1}$ (Reference 2); however, the calculated heats of formation for beryllium sulfate and its hydrates resulting from our enthalpy of dissolution results were found to be in better agreement with other studies.³ This indicated that the calorimeter was performing properly under these conditions.

Procedure

The uranium was handled and loaded into one milliliter glass ampoules under a nitrogen atmosphere. A typical run consisted of breaking the ampoule into 99.8 ± 0.1 ml of nitric acid after establishment of the pre-reaction time-temperature rating period. The temperature rise was then

followed until the steady-state after period was established. Calibration runs followed the experimental runs and consisted of supplying to the heater a known amount of power to match the temperature rise of the reaction. The corrected temperature rise was then determined via a computer program¹ by the Regnault-Pfaundler calculation method.

RESULTS AND DISCUSSION

All results are expressed in defined calories (1 calorie = 4.184 abs joules) and the formula weight of $238.03 \text{ g} \cdot \text{mol}^{-1}$ for uranium was used. Uncertainties stated are twice the standard deviation of the mean, $2\{\sum d^2/n(n-1)\}^{1/2}$.

Table 1 contains the experimental enthalpy data of uranium and nitric acid. Experiment numbers 1-5 and 6-14 were performed under various conditions using 15.9 and 12.7M HNO_3 , respectively. Experiment number 11 was not included because of instrument problems during the calibration run.

Table 1. Enthalpy of Solution Data for Uranium in 99.8 ml of Nitric Acid.

Experiment No. and Designation of Conditions ^a	Weight of Uranium ^b (mg)	e^c (cal·K ⁻¹)	ΔT^d (K ⁻¹)	Δt^e (min.)	$-\Delta H$ (Kcal·mol ⁻¹)	$\Delta t/g$ (min·g ⁻¹)
5-A	226.2	89.58	2.699	51.54	254.4	228
1-A	234.6	89.77	2.774	81.92	252.6	349
4-A	257.0	89.30	3.018	68.34	249.6	266
2-A	317.7	90.02	3.659	85.46	246.8	269
					250.8 ± 3.3	278 ± 51
3-AT	233.2	94.00	2.825	70.64	271.1	303
13-B	234.3	94.54	2.525	84.45	242.5	360
10-B	249.8	94.15	2.693	69.10	241.6	277
					242.0 ± 0.9	318 ± 84
6-B	419.5	96.18	4.599	76.60	251.0	182
7-B	442.3	96.83	4.831	72.55	251.7	164
					251.4 ± 0.7	173 ± 18
8-BT	273.5	98.33	2.922	86.03	250.2	314
9-BT	319.1	98.30	3.386	54.24	248.3	170
					249.2 ± 1.9	242 ± 145
14-BU	224.0	94.37	2.402	90.21	240.8	403
12-BU	300.7	94.50	3.237	59.33	242.1	197
					241.4 ± 1.3	300 ± 207

^a 15.9M HNO_3 (A), 12.7M HNO_3 (B), $38.75 \pm 0.03^\circ\text{C}$ (T), 12.7M HNO_3 + 4.5 g/l uranium (U).

^b Corrected for impurities.

^c e is the electrical energy equivalent.

^d ΔT is the corrected reaction temperature change.

^e Δt is the corrected reaction time.

he enthalpy values in Table 1 were not corrected for the heat absorbed when water was vaporized by the escaping nitrogen oxides because the non-corrected value is more representative of the heat generated during the metal pickling operation. The heat of ampoule breaking and heat of vaporization were found insignificant for the magnitude of the enthalpy values reported here. The corrections for the contribution of the impurity heats of solution were not made because enthalpy data on many of the uranium compounds were not known. However, if an average heat of solution value of $-100 \text{ Kcal}\cdot\text{mol}^{-1}$ is assumed, the estimated correction would be $-6.4 \text{ Kcal}\cdot\text{mol}^{-1}$. Therefore, the results in Table 1 are maximum values, whereas the results minus $-6.4 \text{ Kcal}\cdot\text{mol}^{-1}$ would be the extreme minimum if $-100 \text{ Kcal}\cdot\text{mol}^{-1}$ is assumed for impurities.

The average heat of solution of uranium at 25.0°C in $15.9M \text{ HNO}_3$ (experiments numbers 1, 2, 4, 5) is $-250.8 \pm 3.3 \text{ Kcal}\cdot\text{mol}^{-1}$. Uranium concentration dependence on the heat of solution of uranium metal is apparently insignificant in the 0.23 to 0.32 g U per 100 ml HNO_3 range. The enthalpy of solution of uranium in $15.9M \text{ HNO}_3$ increases to about $-271 \text{ Kcal}\cdot\text{mol}^{-1}$ when the reaction temperature is increased to 39°C (experiment number 3). However, since only one run was made at this temperature, the value should be taken as approximate only.

The enthalpy of solution of uranium in $12.7M \text{ HNO}_3$, representative of the pickling solution, was measured under various conditions (experiment numbers 6-14). The value obtained at 25°C and using $\sim 0.24 \text{ g U}$ per 100 ml HNO_3 , $-242.0 \pm 0.9 \text{ Kcal}\cdot\text{mol}^{-1}$, is less exothermic than the value obtained in $15.9M \text{ HNO}_3$ ($-250.8 \pm 3.3 \text{ Kcal}\cdot\text{mol}^{-1}$). In experiments 6 and 7, an increased amount of uranium ($\sim 0.43 \text{ g U}$ per 100 ml HNO_3) was used which increased the heat of solution to $-251.4 \pm 0.7 \text{ Kcal}\cdot\text{mol}^{-1}$. Increasing the temperature to 39°C (experiments 8 and 9) also increased the enthalpy ($-249.2 \pm 1.9 \text{ Kcal}\cdot\text{mol}^{-1}$). When uranium was dissolved in $12.7M \text{ HNO}_3$ containing 5 g/l U (experiments 14 and 15), the enthalpy was $-241.4 \pm 1.3 \text{ Kcal}\cdot\text{mol}^{-1}$ —little change from uranium-free nitric acid.

The last column in Table 1 gives the corrected reaction time per gram of uranium reacted ($\Delta t/\text{g}$). The data have such large error limits that no correlation of the reaction times with reaction conditions can be made, except that faster reaction times were observed when larger amounts of uranium were dissolved (experiments 6 and 7).

The above enthalpy data in $12.7M \text{ HNO}_3$ are a valuable aid in the design of cooling coils or heat exchangers for control of the exothermic uranium-nitric acid reaction. The heats of solution were found to vary in a normal manner; increased enthalpy with increased temperature and ratio of uranium to nitric acid; the latter condition also increased the time of reaction, as expected. The presence of uranium in the nitric acid was found to have a small effect on the enthalpy value. Thus, the major controlling factor of the reaction, for a specified uranium to nitric acid ratio, is temperature. In the present pickling operation, the reaction is not temperature controlled. With heat exchangers to regulate the pickling temperature, the rate of NO_x generated could be controlled and the capacity of the present NO_x scrubbing system could be maintained.

The rate of dissolution of uranium in nitric acid was reported as decreasing with increasing acidity, unlike most metals.⁴ Therefore, the use of $15.9M$ instead of $12.7M \text{ HNO}_3$ in the pickling operation might be tried to slow the reaction; although the experiments described above did not indicate any advantages with increased nitric acid concentration.

Information on the composition of nitrogen oxides generated in the uranium-nitric acid reaction was also desired for optimization of the scrubbing system. The stoichiometry of the uranium-nitric acid reaction was reported as being dependent on the acid concentration.⁴ (Reactions 1 and 2 represent 8 and $13M \text{ HNO}_3$, respectively):

Since these stoichiometries were cited without references,⁴ an attempt was made to check the latter stoichiometry by analyzing the nitrogen oxides by infrared techniques.⁵ Preliminary analysis of the gases generated in the uranium-15.9M HNO₃ reaction showed that only NO₂ and N₂O were generated at a mole ratio of 8:1 (five runs gave a relative standard deviation of 18%). No NO was observed during the analysis. However, nitric oxide was found to be converted to NO₂ when NO was purged through the nitric acid.⁵ Therefore, it was concluded that NO is probably also generated in the reaction and the exact composition of the evolved nitrogen oxides could not be determined. Knowledge of the general composition of the reaction gases may be deduced from the above information which may be useful for the gas scrubbing operation.

REFERENCES

1. J. D. Navratil and F. L. Oetting, "The Exothermic Enthalpy of TRIS with 0.1M Hydrochloric Acid - Standard Reaction for Solution Calorimetry," USAEC RFP-1976, Rocky Flats Division, Dow Chemical U.S.A., 1973.
2. A. R. Taylor, Jr., B. B. Letson, and D. F. Smith, U.S. Department of the Interior, Bureau of Mines, RI 6724 (1966).
3. J. D. Navratil and F. L. Oetting, *J. Inorg. Nucl. Chem.*, in press.
4. E. H. P. Cordfunke, "The Chemistry of Uranium," Elsevier, New York, 1969, p 33.
5. R. E. Chichorz and T. L. McFeters, "Laboratory Services Semiannual Progress Report," USAEC RFP-2053, May 31, 1973