SEP 8 1965

100. 773-4

FEB 5 1965

SCF CALCULATIONS OF α - AND β -NAPHTHOL⁺Leslie S. Forster and Kichisuke Nishimoto[‡]

MASTER

12597

Department of Chemistry

University of Arizona

Tucson, Arizona

and

Mellon Institute, Pittsburgh, Pennsylvania

DE-FASE - OFFICIAL

This paper was submitted for publication in the open literature at least 6 months prior to the issuance date of this Microcard. Since the U.S.A.E.C. has no evidence that it has been published, the paper is being distributed in Microcard form as a preprint.

⁺Supported by the U. S. Atomic Energy Commission[‡]Permanent address, Department of Chemistry, Faculty of Science, Osaka City University, Osaka, Japan.

ABSTRACT

The Pariser-Parr-Pople method (SCF formalism) is applied to the computation of transition energies and intensities and charge distributions in the ground and excited states of α and β -naphthol. The calculated energy levels are in good agreement with experiment and fairly insensitive to the parameters employed. The transition intensities and charge distributions are sensitive to these parameters but qualitative trends may be correctly predicted.

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

RELEASED FOR ANNOUNCEMENT
IN NUCLEAR SCIENCE ABSTRACTS

FEB 8 1965

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

INTRODUCTION

In spite of serious theoretical shortcomings, the Hückel method has for many years provided a useful conceptual framework for the interpretation of diverse phenomena exhibited by π -electron systems.¹

-
- (1) A. Streitwieser, Molecular Orbital Theory for Organic Chemists, Wiley, New York (1961).
-

This widespread usage of an admittedly defective method is an illustration of the oft encountered principle that an imperfect theory is preferable to no theory at all. The availability of digital computers has made routine the calculation of orbital energies, electron densities, bond orders, etc., for molecules containing 30 (or more) π -electrons. The conceptual and computational simplicities of the Hückel method have been strong factors in its continued use and a comprehensive monograph¹ was published some 30 years after this approximation was introduced.

In the "simple" Hückel theory, electron repulsion is not introduced explicitly, but in 1953 this defect was removed without sacrifice of the intuitive appeal that characterized the Hückel scheme^{2,3}. In the

-
- (2) R. Pariser and R. G. Parr, J. Chem. Phys. 21, 466, 767 (1953).

- (3) J. A. Pople, Trans. Faraday Soc. 49, 1375 (1953).
-

intervening decade hundreds of papers concerned with the so-called Pariser-

Parr-Pople method (or modifications thereof) have been published⁴ and a

-
- (4) An extensive bibliography is contained in R. G. Parr, Quantum Theory of Molecular Electronic Structure, Benjamin, New York (1963).
-

rather detailed description of the method has recently been included in a text book⁵. It is now evident that the computational and conceptual tools

-
- (5) K. B. Wiberg, Physical Organic Chemistry, pp. 108-122, Wiley, New York (1964).
-

for such calculations are (or will soon be) available to a large number of workers.

The success of any semi-empirical method depends upon the evaluation of suitable parameters, transferable from molecule to molecule. It is precisely in this respect that the Hückel method has failed when applied to π -electron systems containing heteroatoms. In this and succeeding papers we adopt a frankly pragmatic outlook in an attempt to determine suitable parameters. When a sufficient number of studies of this type have been made it will be possible to judge the utility of the method and to delimit the domain in which it is applicable. It is particularly important to determine the effect of parameter variation on the energy levels and the π -electron distributions in the ground and excited states.

The naphthols are well suited for the evaluation of the effect of the -OH group on the properties of aromatic hydrocarbons as the transition energies and electronic distributions depend upon the position of the substituent.

METHOD

The self-consistent-field (SCF) variant³ of the Pariser-Parr-Pople method is well adapted for machine calculations, and all of the results described herein have been obtained by this technique. Similar calculations by the ASMO-CI method, the other commonly employed procedure, have been made for naphthalene and the naphthols.⁶ One purpose of this work is to compare the results of the two methods.

(6) K. Nishimoto, J. Phys. Chem. 67, 1443 (1963).

The notation and approximations employed have been previously described.⁷ One important difference must be noted, however.

(7) L. S. Forster, J. Am. Chem. Soc. 86, 3001 (1964).

The expression for the core integral is

$$\alpha_p = -H_p - \sum_{q \neq p} N_q X_{pq}$$

Where $N_q = 2$ for the oxygen atom.⁸

(8) R. L. Miller, P. G. Lykos and H. N. Schmeising, J. Am. Chem. Soc. 84, 4623 (1962).

Ten iterations are sufficient to obtain convergence to ± 0.003 in the charge densities. The filled SCF molecular orbitals obtained are labeled $\phi_1 - \phi_k$ (ϕ_1 is the highest filled orbital) while the unoccupied orbitals are designated $\phi_1 - \phi_n$ (ϕ_1 is the lowest unoccupied orbital), where $2k$ is the number of π -electrons and $k+n$ is the number of atoms in

4

the π -framework. The configurational functions for the one electron excitations, $\phi_i \rightarrow \phi_j$ are then used as a basis for a configuration interaction calculation. This CI calculation is made with a program that evaluates the resultant state energies and wave functions, E_a and $\Psi_a = \sum_m b_{am} \Theta_m$, as well as the intensities of the transitions $\Psi_a \leftarrow \Psi_0$ and the charge distributions in the excited states. A decision must be made regarding the number of configurations to be included in this calculation. Since we are primarily interested in the lower excited states, only configurations within 3 ev of the lowest excited configuration were included (this corresponds to an energy approximately 7 ev above the ground state). In the cases of α - and β -naphthol this encompasses 10 (or 11) of the 30 singly excited configurations. Increasing this interval to 3.4 ev (15 configurations) was without appreciable effect on any of the transitions with energies <6 ev.

Geometry

The geometry of naphthalene⁹ was assumed to be unchanged upon

-
- (9) S. C. Abrahams, J. M. Robertson and J. G. White, Acta Cryst. 2, 238 (1949).
-

substitution and the C-O distance was set at $1.39 \text{ \AA}$ ¹⁰. In α -naphthol,

-
- (10) H. C. Watson and A. Hargreaves, Acta Cryst. 11, 556 (1958).
-

O₁₁ (fig. 1) was placed in line with C₁ and C₄ while in β -naphthol, $\angle O_{11}-C_2-C_3 = 120^\circ$.

Parameter Choice

Assignments of the semi-empirical parameters, W_p , β_{pq} and α_p vary considerably. This is partly due to the relative insensitivity of the

of the experimentally most accessible quantities, the transition energies, to the parameters utilized. If only transition energies are needed for spectral assignments, then this situation is favorable. If, however, properties involving wave functions (intensities, dipole moments, etc.) are to be calculated, it is quite likely that the choice of parameters will be more critical. This is one of the questions which we will examine below.

Valence state ionization energies, W_p . These quantities are set equal to 11.42 ev for all carbon atoms in naphthalene. For oxygen atoms donating two π -electrons, two very different values of W_O have been used, 34.95¹¹ and 23.5¹² ev. The smaller value was used in the calculation of

(11) J. E. Bloor and F. Paradejordi, Theor. Chem. Acta 1, 83 (1962).

(12) M. Orloff and D. Pitts, J. Chem. Phys. 38, 2334 (1963).

furan but in another study of the same molecule, 34.16 ev was employed¹³.

(13) D. Sappinfield and M. Kresvov, Tetrahedron 19 (Suppl. 2), 157 (1963).

We treat this quantity as a free parameter and determine the "best" value.

One center repulsion integrals. For carbon atoms, γ_{cc} is assigned as 10.84 ev (within the range, 10.53-11.13 ev, generally used). Two values for the γ_{oo} integral have been suggested, 21.532¹⁴ and 16.86 ev¹⁵

(14) K. Nishimoto and R. Fujishiro, Bull. Chem. Soc. Japan 31, 1036 (1958).

(15). L. Paolini, Nuovo Cimento 4, 410 (1956).

and both of these were used.

Two center repulsion integrals. The use of $\bar{J}_{pq}$ calculated from theoretical formulae¹⁶ with adjusted exponents^{7,15} leads to an incorrect

(16) C. C. J. Roothaan, J. Chem. Phys. 19, 1445 (1951).

ordering of the two lowest excited states of naphthalene. Reducing the nearest neighbor repulsion integrals by 0.8 ev ($\bar{J}_{cc} = 7.00$ ev), arbitrarily, corrects this problem. It has been shown that the differences in $\bar{J}_{pq}$ are the important quantities rather than the values of individual integrals¹⁷.

(17) P. G. Lykos, J. Chem. Phys. 35, 1249 (1961).

The 0.8 ev reduction leads to differences in closer agreement with those suggested by Lykos¹⁷.

While several other prescriptions for the evaluation of $\bar{J}_{pq}$ yield results not very different from those obtained with the Roothaan formulae^{2,18}, a rather different set of $\bar{J}_{pq}$ are obtained by the method of

(18) J. R. Hoyland and L. Goodman, J. Chem. Phys. 36, 12 (1962).

Nishimoto and Mataga¹⁹. One aim of the present investigation is to compare

(19) K. Nishimoto and N. Mataga, Z. Physik Chem. (Frankfurt) 12, 335 (1957).

the results obtained from these two sets of $\bar{J}_{pq}$ which are designated theoretical and NM.

Resonance integrals, β_{pq} . For carbon-carbon bonds ($r = 1.39 \text{ \AA}$), $\beta_{cc} = -2.236 \text{ ev}$ has been used to fit the naphthalene spectrum⁶. This quantity, somewhat smaller than the parameter commonly used (-2.40 ev), was obtained with the NM repulsion integrals. The naphthalene spectrum can also be fitted with the same β_{cc} and the theoretical integrals. The β_{co} value is rather uncertain. For carbonyl C-O bonds ($r = 1.23 \text{ \AA}$), $\beta_{co} = -2.7 \text{ ev}$ has been used²⁰

(20) R. D. Brown and M. L. Heffernan, Trans. Faraday Soc. 54, 757 (1958).

and a much smaller value, -1.66 ev , has been used for molecules with hydroxy substituents²¹. We have employed an intermediate value, -2.115 ev ¹⁴ in most

(21) K. Nishimoto and R. Fujishiro, Bull. Chem. Soc. Japan, in press.

of the calculations but have varied this parameter over a wide range to evaluate the sensitivity of the results to this quantity.

RESULTS AND DISCUSSION

Naphthalene

The naphthol results are to be compared with the naphthalene calculations. Two naphthalene calculations, differing only in the γ_{pq} integrals, have been made. These results are compared with experiment in Table I. Neither calculation is demonstrably superior to the other.

α - and β -Naphthol

The parameters used in the calculations are listed in Table II. In assessing the reliability of the SCF method and delimiting the domain of

its applicability, we may use several criteria, in order of increasing stringency. At the lowest level, reasonable agreement between the observed and calculated transition energies is acceptable if the calculated transition energies are not too sensitive to parameter choice. This latter requirement minimizes the ad hoc character of the calculations and promotes the hope that the parameters may be transferable from molecule to molecule.

Singlet Transition Energies. In the weak sense outlined above the calculated singlet state energies (Figs. 2 and 3) are encouraging and these may be used for a preliminary screening of the parameters. The following points may be emphasized:

(1) From calculations 1-4 and 5-8 the effect of varying W_0 (NM and theoretical $\bar{U}_{pq}$ are used, respectively, in the two sets of calculations) may be evaluated. The results are not too sensitive to W_0 but it is clear that reducing this quantity below 30 ev leads to poor agreement with experiment²². The "best" value is 33 ev, very close to that obtained from

(22) H. Baba and S. Suzuki, Bull. Chem. Soc. Japan 34, 82 (1961).

Paolini's formula¹⁵, 32.89; and 33.182 ev¹⁴. This parameter was set at 33 ev for the remainder of the calculations.

(2) The variations of β_{CO} (-1.70 to -2.40 ev) and $\bar{U}_{-00}$ (16.860-21.532 ev) have little effect on the singlet transition energies (c.f. calcs. 6, 9, 11, 12, 14).

The results of similar calculations on aniline are equally satisfactory when viewed in this rather non-restrictive manner²³.

(23) J. E. Bloor, P. N. Daykin and P. Boltwood, Can. J. Chem. 42, 121 (1964).

A more demanding criterion for the success of the method is quantitative agreement between the calculated and observed changes in the naphthalene transition energies produced by hydroxyl substitution. The spectra were obtained in isooctane solution. It is found (vide infra) that the π -electron dipole moments in the 1L_a and 1L_b states are larger than the ground state moments and hence the net dipole moments in the ground and excited states differ in magnitude. This factor would tend to shift the naphthol transitions to higher or lower energies depending on the sign of the difference between the ground and excited state net moments.²⁴

(24) N. S. Bayliss and E. G. McRae, J. Phys. Chem. 58, 1002 (1954).

The observed shifts should be corrected for this effect before comparison with the calculated shifts. This uncertainty precludes a strict test of the calculations by this shift criterion, but it is nonetheless of interest to make such a comparison for the 1L_a and 1L_b bands (Table III). It should be emphasized that the calculated naphthol energies are compared to the calculated naphthalene energies obtained with the same set of $\tilde{U}_{pq}$ for the aromatic moiety. On this basis the "best" parameters are $W_0 = 31.33$ ev, $\beta_{CO} = -2.115$ ev and $\tilde{U}_{OO} = 21.532$ ev. Either set of $\tilde{U}_{pq}$ may be used.

Triplet States. The triplet state energies (Figs. 4 and 5) are more sensitive to W_0 than the corresponding singlet state energies. With all parameter combinations, the 3L_a energy is somewhat smaller for α -naphthol than for β -naphthol. The calculations indicate a reduction in the 3L_a energy by hydroxyl substitution.

Ionization Potentials and Electron Affinities. The ionization potentials and electron affinities are $-\epsilon_1$ and $-\xi_1$, respectively. These

quantities are shown in Fig. 6. The ionization potential of naphthalene is 8.26 eV²⁵ and although the experimental electron affinity is unknown, it is

(25) M. E. Wacks and V. H. Dibeler, J. Chem. Phys. 31, 1557 (1959).

very small (or negative)²⁶. The quantities are not known for the naphthols,

(26) R. S. Becker and W. E. Wentworth, J. Am. Chem. Soc. 85, 2210 (1963).

but the ionization potential of phenol is about 0.75 eV less than that of benzene²⁷, a situation roughly in accord with the calculated differences

(27) K. Watanabe, J. Chem. Phys. 26, 542 (1957).

in the naphthalene and naphthol values. We may estimate the naphthol ionization potentials by adding the appropriate differences in the naphthol and naphthalene SCF orbital energies to the observed ionization potential of naphthalene. This leads to the following values: α -naphthol, 7.85 (calc. 2) and 7.67 eV (calc. 6); β -naphthol, 7.99 (calc. 2) and 7.85 eV (calc. 6).

Ground State π -Electron Distribution. The next higher level for testing the adequacy of the SCF formalism involves the calculation of charge distributions. The ground state π -electron distribution is reflected in the ground state dipole moment (μ_{π}). These are quite sensitive to the parameter choice (Fig. 7). Since the magnitude of μ_{π} is unknown, these results cannot be used to choose the best parameters. It is, however, significant that all of the calculations predict that μ_{π} of β -naphthol is somewhat larger (5-10%) than the α -naphthol value. The Hückel method leads to the opposite result.

$\mu_{\pi} =$
 $\mu_{\pi} \text{ sub } \pi$

Similarly, although the charge densities (Table IV) are dependent on the parameters, the order of the charge densities is $q_2 > q_4 > q_3$ for α -naphthol and $q_1 > q_3 > q_4$ for β -naphthol. Hence, any correlation (or lack thereof) of π -electron densities and substitution patterns in electrophilic reactions is not dependent on the parameters used.

Transition Intensities. It has long been apparent that molecular orbital theories are not very useful for the calculation of the intensities of π -electron transitions.²⁸ It is not surprising, therefore, that the

(28) C. M. Moser, J. Chem. Soc. 1954, 3455.

intensities are not in accord with experiment (Table V).

The 1L_a and 1L_b transitions, although weak, will always be observable because they will not be hidden by more intense bands. This is not true for weak transitions of higher energy. The distinction between observable and hidden bands in Figs. 2 and 3 is somewhat arbitrary as can be seen in Table V. However, the number of "observable" bands predicted by the theory is not unreasonable in light of the experimental spectra.²²

The relative ordering of the intensities is quite dependent on the parameters employed. Calculations 6 and 9 both give a reasonable pattern for the relative intensity distribution.

Comparison with Other Studies. One feature that distinguishes the present work from previous treatments of α - and β -naphthol^{6,21} is the increased configuration interaction. In addition to mixing between $\Theta_{1-1'}$, $\Theta_{2-2'}$, $\Theta_{2-1'}$, and $\Theta_{1-2'}$, the configurations represented by $\Theta_{4-1'}$, $\Theta_{3-2'}$, $\Theta_{1-3'}$, $\Theta_{3-1'}$

$\Theta_{1-4'}$, $\Theta_{2-3'}$ and sometimes $\Theta_{4-2'}$ were included. The effect of including these additional configurations is not too large on the transition energies, but the transition intensities and especially the polarization of the 1L_b band are quite sensitive to the amount of CI. The conclusion that the α -naphthol Ψ_1 and Ψ_2 resemble closely the corresponding naphthalene wave functions while the β -naphthol wave functions are very different⁶ is still valid, but a small difference in Ψ_1 can result in a markedly different polarization of 1L_b (Table VI). The effect is especially pronounced in this weak transition (forbidden in naphthalene) and it is uncertain that calculations of this type will be useful for such polarization predictions.

In general, the results of the SCF calculations are similar to those obtained by the ASMO-CI variant of the Pariser-Parr-Pople method. A comparison of several of the SCF calculations with an ASMO-CI calculation is made in Table VI. The availability of a high digital computer comparable in speed and storage to the IBM 7072 or 7090 would favor the use of the SCF method while the ASMO-CI method would be more feasible if such a computer were not available.

Tabular presentation of the results and a description of the computer programs are available from the authors.

Acknowledgements. We are grateful to the University of Arizona Numerical Analysis Laboratory for providing machine time and to Professor Robert L. Baker for programming assistance.

Table I

Cited State	Excitation Energy (ev)			Oscillator Strength (f)		
	Calc. 1 ^a	Calc. 2 ^b	Observed ^c	Calc. 1	Calc. 2	Observed ^c
1L_b	4.121	4.008	3.99	0	0	0.022
1L_a	4.269	4.287	4.51	0.191	0.270	0.11
1B_b	5.860	5.540	5.62	2.002	1.944	1.70
1B_a	6.274	6.178	-----	0.981	0.852	-----
3L_a	2.773	2.343	2.64			

(a) Theoretical $\bar{\epsilon}_{pq}$ (nearest neighbor $\bar{\epsilon}_{cc} = 7.00$ ev)

(b) Nishimoto and Mataga $\bar{\epsilon}_{pq}$ ¹⁹ (equal bond lengths, $r_{cc} = 1.39$ Å)

(c) Ref. 22.

Table II

KEY TO CALCULATIONS

Calc. No.	W_0 (ev)	$\bar{U}_{00}$ (ev)	$\bar{U}_{pq}^a$	E_{CO} (ev)
1	34.95	21.532	I	-2.115
2	33.00	21.532	I	2.115
3	31.00	21.532	I	-2.115
4	23.50	21.532	I	-2.115
5	34.95	21.532	II	-2.115
6	33.00	21.532	II	-2.115
7	31.00	21.532	II	-2.115
8	29.00	21.532	II	-2.115
9	33.00	21.532	II	-2.400
10	33.00	16.860	II	-2.115
11	33.00	21.532	II'	-2.115
12	33.00	21.532	II'	-1.700
13	33.00	21.532	I'	-2.115
14	33.00	16.860	II'	-2.115

a. $\bar{U}_{pq}$ are designed as follows:

I Wishimoto and Mataga integrals¹⁹ ($r_{CO} = 1.39 \text{ \AA}$)

I' Same as I for $r_{CO} = 1.46 \text{ \AA}$

II Theoretical integrals; nearest neighbor $\bar{U}_{CC} = 7.00 \text{ ev}$

II' Same as II but nearest neighbor $\bar{U}_{CO}$ reduced by 0.8 ev

All calculations with I and I' $\bar{U}_{pq}$ are to be compared with naphthalene 2 and all others with naphthalene 1.

Table III

SPECTRAL SHIFTS (RELATIVE TO NAPHTHALENE) (ev)

Calc. No.	<u>α-Naphthol</u>		<u>β-Naphthol</u>	
	1L_b	1L_a	1L_b	1L_a
Observed ^a	-0.13	-0.20	-0.21	+0.03
1	-0.141	-0.132	-0.078	-0.031
2	-0.078	-0.237	-0.154	-0.055
3	-0.168	-0.423	-0.332	-0.096
5	-0.051	-0.103	-0.091	-0.017
6	-0.106	-0.180	-0.186	-0.040
7	-0.233	-0.303	-0.384	-0.078
8	-0.439	-0.457	-0.679	-0.107
9	-0.125	-0.201	-0.216	-0.025
10	-0.023	-0.053	-0.042	-0.011
11	-0.100	-0.180	-0.176	-0.040
12	-0.073	-0.149	-0.133	-0.041
13	-0.081	-0.237	-0.158	-0.055
14	-0.022	-0.054	-0.041	-0.012

a. Ref. 22

Table IV

<u>α-Naphthol</u>					<u>β-Naphthol</u>			
Calc. No.	$\mu_{\Sigma}(D)$	q_2	q_3	q_4	$\mu_{\Sigma}(D)$	q_1	q_3	q_4
1	0.94	1.048	0.994	1.024	0.96	1.048	1.029	0.995
2	1.36	1.063	0.992	1.033	1.40	1.063	1.037	0.993
3	2.05	1.086	0.990	1.047	2.09	1.087	1.050	0.991
4	7.25	1.219	0.979	1.143	7.63	1.240	1.119	0.982
5	1.03	1.064	0.986	1.031	1.09	1.065	1.040	0.986
6	1.51	1.085	0.980	1.042	1.60	1.087	1.052	0.982
7	2.30	1.117	0.972	1.060	2.52	1.120	1.070	0.975
8	3.43	1.158	0.962	1.084	3.61	1.165	1.093	0.966
9	1.82	1.102	0.976	1.051	1.87	1.105	1.063	0.978
10	0.67	1.046	0.990	1.022	0.72	1.046	1.029	0.990
11	1.44	1.083	0.981	1.041	1.49	1.085	1.051	0.982
12	1.02	1.059	0.986	1.029	1.05	1.060	1.036	0.987
13	1.38	1.064	0.992	1.034	1.44	1.065	1.038	0.993
14	0.64	1.045	0.990	1.021	0.66	1.046	1.029	0.991

Table V
TRANSITION INTENSITIES (f)

	$\Psi_0 \rightarrow \Psi_1$	Ψ_2	Ψ_3	Ψ_4	Ψ_5	Ψ_6	Ψ_7
Calc. No.							
<u>α-Naphthol</u>							
Observed ^a	0.016(1L_b)	0.102(1L_a)	0.328 and	0.892			
1	0.0113	0.294	0.750	0.0170	1.111	0.0054	0.875
2	0.0226	0.304	0.774	0.0196	0.230	0.806	0.895
3	0.0486	0.311	0.658	0.0729	0.0333	0.962	0.938
5	0.0153	0.225	0.0062	0.250	0.0597	1.630	1.000
6	0.0391	0.237	0.0133	0.423	0.00981	1.421	1.045
7	0.0681	0.241	0.0365	0.469	0.00089	1.231	1.099
8	0.0867	0.284	0.170	0.287	0.0209	1.081	0.719
9	0.0498	0.237	0.0160	0.418	0.00799	1.406	1.064
10	0.00651	0.209	0.00241	0.120	1.843	0.0373	0.979
11	0.0411	0.231	0.0110	0.441	0.0102	1.415	1.036
12	0.0254	0.218	0.00816	0.515	0.0115	1.369	1.010
13	0.0235	0.305	0.764	0.0196	0.214	0.829	0.896
14	0.0638	0.182	0.00112	0.117	1.848	0.0220	0.977
<u>β-Naphthol</u>							
Observed ^a	0.0211(1L_b)	0.0811(1L_a)	1.06				
1	0.0311	0.220	1.150	0.381	0.432	0.0322	0.806
2	0.0562	0.186	0.970	0.558	0.408	0.191	0.592
3	0.0950	0.149	0.533	1.023	0.284	0.562	0.115
5	0.0464	0.134	0.0670	0.0234	1.859	0.125	0.919

Table V Continued

$\Psi_0 \rightarrow \Psi_1$	Ψ_2	Ψ_3	Ψ_4	Ψ_5	Ψ_6	Ψ_7	
Calc. No.							
6	0.0718	0.105	0.155	0.0512	1.725	0.205	0.790
70	0.104	0.0808	0.192	0.355	1.351	0.217	0.693
8	0.123	0.0828	0.0965	0.775	0.772	0.490	0.744
9	0.0857	0.0985	0.154	0.0587	1.687	0.273	0.781
10	0.0254	0.158	0.0190	0.0131	1.939	0.0683	0.961
11	0.0709	0.105	0.158	0.0480	1.724	0.187	0.799
12	0.0503	0.123	0.168	0.0425	1.707	0.122	0.834
13	0.0577	0.184	0.888	0.634	0.414	0.205	0.579
14	0.0251	0.158	0.0185	0.00714	1.939	0.0652	0.966

a. Ref. 22

Table VI

COMPARISON OF SCF AND ASMO-CI RESULTS

 α -Naphthol β -Naphthol

A. Transition Energies (ev)

Excited State	SCF		ASMO-CI ⁶		SCF		ASMO-CI ⁶	
	Calc. 6	Calc. 2		Obs. ²²	Calc. 6	Calc. 2		Obs. ²²
¹ Ψ_1	4.015	3.930	3.928	3.86	3.935	3.854	3.880	3.78
¹ Ψ_2	4.089	4.050	4.010	4.31	4.229	4.232	4.233	4.54
¹ Ψ_3	5.221	5.374	-----	-----	5.246	5.295	-----	-----
¹ Ψ_4	5.383	5.257	5.490	5.40	5.484	5.377	-----	-----
¹ Ψ_5	5.696	5.606	-----	-----	5.707	5.535	5.414	5.53
¹ Ψ_6	5.903	5.637	-----	5.80	5.963	5.873	5.896	-----
¹ Ψ_7	6.337	6.198	6.169	-----	6.115	5.972	-----	-----
³ Ψ_1	2.512	2.213	2.197	-----	2.688	2.300	2.273	-----
³ Ψ_2	3.480	3.116	-----	-----	3.439	3.088	-----	-----

Table VI (Cont.)

 α -Naphthol β -Naphthol

A. Transition Energies (ev) (Cont.)

SCF

SCF

Excited State	Calc. 6	Calc. 2	ASMO-CI ⁶	Obs. ²²	Calc. 6	Calc. 2	ASMO-CI ⁶	Obs.
$^3\Psi_3$	3.617	3.397	3.388	----	3.663	3.393	3.388	----
$^3\Psi_4$	4.140	3.983	3.955	----	3.956	3.761	3.792	----
$^3\Psi_5$	4.453	4.061	4.080	----	4.443	4.145	4.200	----
$^3\Psi_6$	5.500	5.137	-----	----	4.810	4.563	-----	----

B. Transition Intensities (f) and Polarizations

$^1\Psi_1$	0.0391(-50.0°) ^a	0.0226(2.7°)	0.013(x)	0.016	0.0718(-60.1°)	0.0562(-62.7°)	0.066(-64°)	0.021
$^1\Psi_2$	0.237(78.8°)	0.304(84.7°)	0.332(y)	0.102	0.105(75.0°)	0.186(-74.5°)	0.195(-76°)	0.081
$^1\Psi_3$	0.0133(76.4°)	0.0196(74.1°)	-----	----	0.155(1.3°)	0.970(4.8°)	-----	-----

Table VI (Cont.)

α -Naphthol				β -Naphthol				
B.	Transition Intensities (f) and Polarizations (Cont.)							
	SCF				SCF			
Excited State	Calc. 6	Calc. 2	ASMO-CI ⁶	Obs. ²²	Calc. 6	Calc. 2	ASMO-CI ⁶	Obs.
¹ Ψ_4	0.423(1.8°)	0.774(-0.3°)	1.949(x)	0.328	0.0512(85.2°)	0.558(20.5°)	-----	-----
¹ Ψ_5	0.0098(-68.8°)	0.230(-13.8°)	-----	-----	1.725(11.3°)	0.408(12.8°)	2.037(10°)	1.06
¹ Ψ_6	1.421(-2.1°)	0.806(1.0°)	-----	0.892	0.205(44.3°)	0.191(-47.3°)	0.763(-14°)	-----
¹ Ψ_7	1.045(89.3°)	0.895(89.7°)	0.802(y)	-----	0.790(-67.8°)	0.592(-81.0°)	-----	-----
C.	Calculated μ_x (D)							
Ψ_0	1.51(87.5°) ^b	1.36(86.5°)	1.34(91°)		1.60(16.5°)	1.40(17.8°)	1.19(13°)	
¹ Ψ_1	3.22(46.7°)	2.70(46.3°)	2.71(21°)		4.14(25.5°)	3.51(22.4°)	3.75(21°)	
¹ Ψ_2	3.87(39.2°)	3.49(36.6°)	3.70(18°)		4.13(26.0°)	3.41(30.2°)	3.31(33°)	

a. Angle between transition moment vector and +x axis.

b. Direction of dipole moment relative to +x axis.

LEGENDS

- Fig. 1 Core structure of naphthols.
- Fig. 2 α -naphthol singlet energy levels. _____ and - - - - - represent transitions likely to be observed and unobserved, respectively. (See text and Table V). Experimental naphthalene and α -naphthol transitions are represented by $\bigcirc$ and $\odot$ respectively.
- a. No configuration interaction, only lowest configurational energy indicated.
- b. 1' and 2' refer to naphthalene 1 and 2.
- Fig. 3 β -naphthol singlet energy levels. Same conventions as in Fig. 2.
- Fig. 4 α -naphthol triplet energy levels.
- Fig. 5 β -naphthol triplet energy levels.
- Fig. 6 Ionization potentials (solid lines) and electron affinities (broken lines) for α -naphthol. The corresponding ionization potentials for β -naphthol are 0.1 - 0.2 ev larger and the electron affinities are 0.03 - 0.1 ev larger.
- Fig. 7 Molecular diagrams. π -electron densities and bond orders.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig 6

Calc. 6

Calc. 2

Fig. 7

ADDENDUM TO NAPHTHOL SCF CALC.

SINGLET TRANSITION ENERGIES (ev)

Calc. No.	E ₁	E ₂	E ₃	E ₄	E ₅	E ₆	E ₇
α -Naphthol Observed ^a	3.86(¹ L _b)	4.31(¹ L _a)	5.40 and 5.80(¹ B _b)				
1	3.967	4.155	5.350	5.451	5.595	5.746	6.188
2	3.930	4.050	5.257	5.374	5.606	5.637	6.198
3	3.840	3.864	5.093	5.238	5.413	5.694	6.213
4	<3.109	No Configuration Interaction Calculation					
5	4.070	4.166	5.304	5.438	5.837	5.867	6.315
6	4.015	4.089	5.221	5.383	5.696	5.903	6.337
7	3.888	3.966	5.119	5.246	5.485	5.965	6.360
8	3.682	3.812	5.018	5.114	5.307	5.996	6.330
9	3.996	4.068	5.226	5.344	5.679	5.914	6.352
10	4.098	4.216	5.364	5.474	5.858	5.932	6.297
11	4.021	4.089	5.220	5.403	5.711	5.910	6.335
12	4.048	4.120	5.211	5.493	5.746	5.908	6.314
13	3.927	4.050	5.253	5.374	5.604	5.635	6.198
14	4.099	4.215	5.361	5.476	5.858	5.937	6.296
β -Naphthol Observed ^a	3.78(¹ L _b)	4.54(¹ L _a)	5.53(¹ B _b)				
1	3.930	4.256	5.410	5.443	5.558	5.898	6.056
2	3.854	4.232	5.295	5.377	5.535	5.873	5.972
3	3.676	4.191	5.102	5.223	5.444	5.811	5.933
4	<2.749	No Configuration Interaction Calculation					
5	4.030	4.252	5.361	5.502	5.771	6.004	6.169
6	3.935	4.229	5.246	5.484	5.707	5.963	6.115

Calc. No.	E_1	E_2	E_3	E_4	E_5	E_6	E_7
7	3.737	4.192	5.083	5.397	5.617	5.861	6.098
8	3.442	4.162	4.953	5.287	5.528	5.744	6.145
9	3.905	4.244	5.273	5.493	5.698	5.952	6.129
10	4.079	4.258	5.399	5.507	5.813	6.019	6.219
11	3.945	4.229	5.251	5.491	5.709	5.970	6.122
12	3.988	4.228	5.273	5.506	5.726	5.989	6.117
13	3.850	4.232	5.289	5.374	5.537	5.877	5.970
14	4.080	4.257	5.400	5.507	5.813	6.019	6.221

(a) Ref. 22

TRIPLET STATE ENERGIES (ev)

Calc. No.	E ₁	E ₂	E ₃	E ₄	E ₅	E ₆
<u>α-Naphthol</u>						
1	2.282	3.110	3.456	3.993	4.056	5.279
2	2.213	3.116	3.397	3.983	4.061	5.137
3	2.020	3.076	3.269	3.956	4.072	4.855
5	2.655	3.499	3.695	4.137	4.443	5.613
6	2.512	3.480	3.617	4.140	4.453	5.500
7	2.148	3.399	3.503	4.141	4.484	5.289
8	1.515	3.257	3.293	4.134	4.575	4.978
9	2.479	3.461	3.568	4.157	4.461	5.446
10	2.724	3.513	3.743	4.131	4.437	5.688
11	2.567	3.535	3.627	4.142	4.451	5.512
12	2.621	3.604	3.716	4.121	4.445	5.663
13	2.208	3.105	3.392	3.982	4.062	5.131
14	2.732	3.523	3.744	4.132	4.437	5.696
<u>β-Naphthol</u>						
1	2.327	3.120	3.467	3.847	4.135	5.385
2	2.300	3.088	3.393	3.761	4.145	4.563
3	2.164	2.929	3.272	3.431	3.855	4.264
5	2.740	3.537	3.716	4.030	4.453	5.758
6	2.688	3.439	3.663	3.956	4.443	4.810
7	2.369	3.148	3.564	3.653	4.156	4.583
8	1.647	2.906	3.236	3.621	4.150	4.629
9	2.671	3.416	3.655	3.913	4.462	4.774
10	2.760	3.534	3.753	4.073	4.439	5.766

Continued

Calc. No.	E_1	E_2	E_3	E_4	E_5	E_6
11	2.703	3.462	3.672	3.982	4.473	4.837
12	2.725	3.492	3.693	4.035	4.476	4.940
13	2.999	3.085	3.388	3.752	4.141	4.500
14	2.761	3.535	3.757	4.076	4.443	5.768

Change Densities and Dipole Moments

<u>α</u> -Naphthol					<u>β</u> -Naphthol			
Calc. No.	$\mu_{\pi}(D)$	q_2	q_3	q_4	$\mu_{\pi}(D)$	q_1	q_3	q_4
1	0.94	1.048	0.994	1.024	0.96	1.048	1.029	0.995
2	1.36	1.063	0.992	1.033	1.40	1.063	1.037	0.993
3	2.05	1.086	0.990	1.047	2.09	1.087	1.050	0.991
4	7.25	1.219	0.979	1.143	7.63	1.240	1.119	0.982
5	1.03	1.064	0.986	1.031	1.09	1.065	1.040	0.986
6	1.51	1.085	0.980	1.042	1.60	1.087	1.052	0.982
7	2.30	1.117	0.972	1.060	2.52	1.120	1.070	0.975
8	3.43	1.158	0.962	1.084	3.61	1.165	1.093	0.966
9	1.82	1.102	0.976	1.051	1.87	1.105	1.063	0.978
10	0.67	1.046	0.990	1.022	0.72	1.046	1.029	0.990
11	1.44	1.083	0.981	1.041	1.49	1.085	1.051	0.982
12	1.02	1.059	0.986	1.029	1.05	1.060	1.036	0.987
13	1.38	1.064	0.992	1.034	1.44	1.065	1.038	0.993
14	0.64	1.045	0.990	1.021	0.66	1.046	1.029	0.991

Orbital Energies (ev)

Calc. No.	Naphthalene			
	ϵ_1	$\epsilon_{1'}$		
1	-10.18649	-1.81368		
2	-9.49766	-2.50249		
	α -Naphthol		β -Naphthol	
	ϵ_1	$\epsilon_{1'}$	ϵ_1	$\epsilon_{1'}$
1	-9.23282	-2.35715	-9.33113	-2.39738
4	-7.23412	-1.70495	-7.15098	-1.81317
5	-9.82067	-1.62617	-9.95035	-1.66899
6	-9.60183	-1.55575	-9.77631	-1.60498
7	-9.22442	-1.44695	-9.40293	-1.50103
8	-8.72959	-1.29689	-8.84918	-1.35057
2	-9.08575	-2.30743	-9.22338	-2.35677
3	-8.81324	-2.23054	-8.97495	-2.29159
9	-9.51613	-1.50302	-9.69997	-1.56469
10	-9.96858	-1.68343	-10.05139	-1.71851
14	-9.97605	-1.68600	-10.05670	-1.72169
11	-9.63024	-1.56455	-9.80126	-1.61692
12	-9.75868	-1.63730	-9.90827	-1.67772
13	-9.08590	-2.30641	-9.22609	-2.35767