FEB 2 1968

O.R. AEC Conference on Nucl. Med
Cleveland 9/27/67

ACRH--1000-34
CONF-670958--1

MASTER

BONE SCANNING

Alexander Gottschalk, M.D.

Associate Professor of Radiology, Chief of the Section of
Nuclear Medicine, Department of Radiology, University
of Chicago Hospitals and Clinics, and
Director, Argonne Cancer Research Hospital
(operated by the University of Chicago for the
United States Atomic Energy Commission)

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

leg

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

BONE SCANNING

Alexander Gottschalk, M.D.

Associate Professor of Radiology, Chief of the Section of

Nuclear Medicine, Department of Radiology, University

of Chicago Hospitals and Clinics, and

Director, Argonne Cancer Research Hospital

(operated by the University of Chicago for the

United States Atomic Energy Commission)

Chicago, Illinois

Abstract

This article is a brief critical review of bone scanning. A basic discussion of bone physiology as it relates to bone scanning is presented, and the radionuclides most frequently used for bone scanning are characterized. Since no ideal scanning agent is available, the value and limitations of each agent are discussed. The rationale for bone scanning is examined and evaluated, and future developments of the technique are suggested.

Introduction

This presentation reviews the salient features of bone scanning as they have developed to date. A complete treatise on the subject is not attempted, but this effort is rather aimed at calling attention to some of the essential features and limitations of the technique as it is currently practiced.

The British surgeon, John Belchier, was probably the first to use a bone tracer of any kind, when in 1736 he noted that the bones of animals fed extracts of madder root became red¹. In 1935, Chiewitz and de Hevesy found that there was a rapid uptake of phosphorus-32 in bone². In 1942 Treadwell and colleagues at the Donner Laboratory demonstrated that strontium-90 concentrated in neoplastic osseous tissue³. In the 1950's, Bauer and colleagues began extensive studies utilizing external counting with strontium-85 and calcium-47⁴. In 1961, the first strontium-85 bone scan was made by Fleming *et al.*⁵ at Bethesda, and in 1962, the first fluorine-18 bone scan was performed by Blau and colleagues⁶.

Physiology

The physiology of bone is an extremely complicated subject, and only the fundamentals pertaining to bone scanning will be considered here. The

interested reader is referred to the monograph by McLean and Urist⁷ for a starting point in this complicated field. The substance of bone is composed of organic matrix, crystalline bone salt, and associated water. Our concern with scanning agents leads us to consider primarily the mineral phase of bone since as yet no matrix tracers have been investigated for scanning. Bone mineral has crystals that are ultramicroscopic in size, chemically nonhomogeneous, and arranged in a basic structure that has been shown by X-ray diffraction patterns to be an apatite crystal, generally agreed to be hydroxyapatite. The crystals are arranged so that a large surface area is presented to the body fluids, with the result that extremely rapid exchanges are possible in areas of metabolic activity. In particular, ionic substitution or exchange as well as dissolution and recrystallization may occur readily.

For bone scanning, our attention is particularly directed to ion transfer and ion exchange. In particular, the process of heteroionic exchange, i.e., the exchange of one ion for a different ion, is important in the scanning mechanism. It is postulated that an ion may have a choice of one of four positions and still be in relation to the hydroxyapatite in bone: (1) It may be in the interior of the bone crystal, therefore relatively stable; (2) it may be on the surface of the bone crystal; (3) it may be adsorbed to the surface of the crystal; or, (4) it may be in solution in the hydration shell in immediate proximity to the crystalline surface⁷. It is generally believed that it takes at least several days, and possibly weeks, for an ion to become placed reasonably securely in the interior of the crystalline surface itself. The surface and exchange phenomenon is an extremely dynamic one, and a basic premise of bone scanning is that this phenomenon occurs most avidly where bone metabolism is active.

Bone Scanning Agents

The properties of the four most commonly used bone scanning agents are summarized in Table 1. It is important to recognize from this Table that an ideal bone scanning agent does NOT exist. If the basic composition of hydroxyapatite consists of hydroxyl groups, calcium, and phosphorus, then any isotopes of these elements would seem to be logical choices. Unfortunately, no ideal isotopes of these elements are available with physical properties well suited to scanning, and calcium-47 is the only "natural" nuclide that has even been investigated as a scanning agent. Calcium-47 has a gamma energy of 1.3 MeV, which is extremely high, and requires special scanning equipment. Isotopes of strontium have been shown to behave somewhat similarly to calcium, and strontium has been used effectively as a bone scanning agent. The physical characteristics of strontium-85 are such that in the doses employed (and these are limited by the rather high dose delivered to bone, stemming primarily from the long half-life of strontium-85) the scan times are extremely long due to count rate limitations. This deficiency is overcome somewhat by the use of strontium-87m, but here the half-life is so short that a scan must be done within hours following injection of the nuclide. Work at the Sloan-Kettering Institute⁸ has shown that the tumor to normal bone ratio in a small series of patients was such that tumor scans done with strontium at 5 hours appeared normal in 10% of the cases, ultimately becoming positive at 5 days. One might theorize that the ion exchange on the bone mineral surface of these patients is not always rapid for strontium, but ultimately a diagnostically usable portion of the nuclide will be incorporated into the interior of the bone crystal. The Sloan-Kettering study would seem to provide important empiric evidence that strontium scanning will yield the highest results when the scan is made days (not hours) after administration of the nuclide. As a result, serious misgivings about the use of strontium-87m as an effective bone scanning agent seem warranted.

Fluorine-18 may ultimately turn out to be the best compromise. The disadvantage of fluorine-18 at the moment is that it requires reactor or cyclotron

production and chemical processing, both of which require skills beyond those found in many nuclear facilities. The short half-life prohibits shipping of the nuclide for great distances, but as production methods become commercialized, the fact that this material can be produced in very large curie quantities gives hope that shipping from a central commercial supplier to a rather wide peripheral region may be accomplished. The Sloan-Kettering group has also shown that for EARLY bone scanning, fluorine-18 is as good as, or better than, calcium when the ratios of abnormal to normal bone uptake are compared⁹. Again, the quantification has been done in only a relatively small group of patients, but these results are encouraging. It should be pointed out, however, that the highest abnormal to normal bone ratios were always obtained with calcium in approximately five days following administration of the isotope. The advantages in using fluorine-18 are that it can be administered in millicurie rather than microcurie quantities, and the radiation dose to the patient is modest.

In summary, then, if scanning is to be started early after the administration of a nuclide, fluorine-18 would seem to be the agent of choice. The best abnormal to normal bone ratios can be obtained with calcium-47 if satisfactory instrumentation is available. If strontium is to be used, strontium-85 scanned at approximately 5 days following administration seems to give better results than strontium-87m scanned in the first few hours.

A Critical Appraisal of the Rationale for Bone Scanning

When considering bone scanning, it is always important to realize that the lesions demonstrated are entirely nonspecific. A partial list of the etiology of positive bone scans is given in Table 2. To date, much of the bone scanning done has been a part of the work-up of the known or suspected cancer patient. In these patients, the complaint of bone pain is likely to be the most common underlying stimulus for bone scanning or roentgen examination.

There seems to be no question that the bone scan will demonstrate lesions that are invisible on roentgenograms (see Figure 1). For example, in some series of selected cancer patients with pain or other indication of bone metastases, approximately 60% of those with negative roentgenograms had positive bone scans, whereas only about 20% or less of those with definite roentgen lesions had negative bone scans^{10,11}. Radioautography of different bone lesions shows that although the lesion itself may not be osteoblastic in character, it stimulates bone metabolism around its periphery--presumably the body's effort at healing--and this metabolic bone activity is responsible for a nuclide uptake which causes the bone scan to be positive^{12,13}. In short, both osteoblastic and osteolytic metastases can be demonstrated by scanning.

It is not infrequently stated that the correlation of the size of a lesion as shown on bone scan and its actual dimensions is quite good¹⁴. Although this is undoubtedly true for some cases, it must be recognized that bone scanning is done most often with a coarse focused collimator which may display a lesion with a magnification approximately ten times its actual size. As a result, judging the true size of any lesion from its bone scan is a hazardous undertaking. The extent of the pathologic process can often be more clearly defined by scanning, since regions thought to be normal on roentgenogram will be shown to contain lesions on scan^{11,15}.

The scan has been used to aid the radiation therapist to plan adequate treatment portals. While this is certainly of benefit, it must be recognized that innumerable patients with widespread bony metastases undergo therapy, and the experienced radiation therapist will "treat them where they hurt," recognizing that the metastatic disease is certainly more extensive than is likely to be demonstrated by any diagnostic technique, whether roentgenography or bone scanning. And, finally, the cancer patient may well be helped by using the bone scan to help evaluate therapeutic response.

The overall experience with bone scanning to date suggests that one of the most useful patient groups to examine is the preoperative cancer suspect. The bone scan is known to be useful for detecting metastatic lesions prior to their roentgenographic appearance. (It has long been recognized that 30-50% bony destruction must be present before a lesion can be seen on roentgenogram¹⁶.) It is reasonable to argue that the addition of bone scanning to the work-up of the preoperative patient would be useful as an additional diagnostic tool to evaluate metastatic disease prior to operation. Unfortunately, no such series exists. Sklaroff and Charkes however, have recently reported a series of stage II breast carcinoma patients who were scanned shortly following operation¹⁷. In the 64 women described, 10 (16%) had lesions demonstrated on bone scanning that were not detected on X rays. Of these 10, four died with autopsy proven bone lesions. Three of these four never had positive X rays. Although it is to be stressed again that bone scanning is not specific, certainly an area positive on the bone scan, not positive on a roentgenogram, in a preoperative patient, can be biopsied. With this technique it should be possible to further diagnose metastatic disease prior to surgery, and avoid useless operation.

Bone scanning has also been used to evaluate a variety of non-neoplastic disorders. One of the most imaginative of these studies has been the investigation of traumatic compression fractures of the spine in an effort to determine whether these are recent or old. A series of such cases has been studied at Miami¹⁸. Since both recent and old vertebral fractures can cause virtually indistinguishable vertebral "wedging" on roentgenograms, differentiation is often difficult. A recent lesion, however, should have a positive bone scan, and an old lesion should not. In addition, this group found that the bone scan of compression fractures taken from 6 to 24 months to return to normal.

Finally, the investigation of arthritis continues. Bauer and colleagues recently demonstrated the appearance of strontium in the osteoarthritic knee, and a suggestion that the nuclide distribution is related to the stress forces in the joint has been advanced¹⁹.

General Considerations of Technique and Interpretation

The first point to be stressed is that scanning agents accumulate unevenly throughout the skeleton in the normal patient, being high in areas of normally active metabolic bone (such as the joint regions) and low in other areas. It is urged therefore that the nuclear physician familiarize himself with the normal scan pattern before attempting to evaluate potentially abnormal cases, since techniques of film development for photoscanning will vary from institution to institution, and scan parameters such as line-spacing, speed, time constant, and background erase may also differ.

With strontium-85 (the isotope most commonly used at this time), accumulation of the nuclide in the large bowel in the first few days following administration is to be expected. As a result, scanning either should be undertaken at a later period, e.g., at 5 to 7 days, or a cleansing enema should be administered in order to avoid overlap of colonic activity. Wherever possible, comparison of both sides is always useful. For example, if a left knee is the region of suspicion, a scan of the normal right side should also be undertaken. Finally, with strontium-85 slow scan speeds are necessary since count rates less than twice background are common.

Conclusions

In my judgment, the emphasis in bone scanning to date has been somewhat misplaced. In particular, the game "find the metastatic lesion the diagnostic radiologist has missed" is probably overplayed, since the good therapeutic radiologist is accustomed to giving palliative treatment for complaints which

in his judgment represent metastatic disease, whether or not these can be substantiated by either roentgenography or bone scanning. Bone scanning should be more important in the work-up of the preoperative patient where occult metastatic disease can be discovered and useless surgery avoided.

Too little has been done to utilize bone scanning to investigate bone physiology in a variety of normal and pathologic situations. Van Dyke, for example, has amassed evidence that fluorine-18 (and this is likely to be true of all bone scanning agents) is dependent upon bone blood flow in a large part for its selective localization in regions of bone pathology²⁰. This work should be followed up in a variety of clinical situations, for example, the aseptic necroses.

The extent and scope of bone scanning, like so much of nuclear medicine, has grown appreciably in the past few years. The fact that there are still "lions to slay" makes this a challenging area. Let us hope that bone scanning receives the creative attention it merits so that we may stalk more lions in the future.

Acknowledgment

The author is indebted to Dr. W. S. Tirman and his colleagues in the Department of Radiology at the Memorial Hospital, South Bend, Indiana, for the case material shown in Figure 1.

References

1. J. Belchier, An account of the bones of animals being changed to a red colour by aliment only, Phil. Trans., 39: 287 (1732).
2. O. Chiewitz, and G. de Hevesy, Radioactive indicators in the study of phosphorus metabolism in rats, Nature, 136: 754 (1935).
3. A. deG. Treadwell, B. V. A. Low-Beer, H. L. Friedell, and J. H. Lawrence, Metabolic studies on neoplasm of bone with the aid of radioactive strontium, Am. J. Med. Sci., 204: 521 (1942).
4. G. C. H. Bauer, and B. Wendeborg, External counting of Ca^{47} and Sr^{85} in studies of localised skeletal lesions in man, J. Bone Joint Surg., 41B: 558 (1959).
5. W. H. Fleming, J. D. McIlraith, and E. R. King, Photoscanning of bone lesions utilizing strontium-85, Radiology, 77: 635 (1961).
6. M. Blau, W. Nagler, and M. A. Bender, Fluorine-18: New isotope for bone scanning, J. Nucl. Med., 3: 332 (1962).
7. F. C. McLean, and M. R. Urist, Bone: An introduction to the physiology of skeletal tissue, 2nd ed., University of Chicago Press, Chicago, 1961.
8. D. A. Weber, Sloan-Kettering, personal communication, 1966.
9. E. Greenberg, Sloan-Kettering, personal communication, 1966.
10. W. J. Simpson, and R. P. Orange, Total body scanning with ^{85}Sr in the diagnosis of metastatic bone disease, Can. Med. Assoc. J., 93: 1237 (1965).
11. D. F. Williams, and Wm. H. Blahd, The diagnostic and prognostic value of strontium-85 photoscanning in carcinoma of the prostate, J. Urol., 97: 1070 (1967).

12. E. Greenberg (discussant), in Radioisotopes and Bone, F. C. McLean, P. Lacroix, and Ann M. Budy (Eds.), p. 103, F. A. Davis Co., Philadelphia, 1962.
13. K. R. Corey, P. Kenny, E. Greenberg, A. Pazianos, O. H. Pearson, and J. S. Laughlin, The use of Ca^{47} in diagnostic studies of patients with bone lesions, Am. J. Roentgenol. Radium Therapy Nucl. Med., 85: 955 (1961).
14. D. M. Sklaroff, and N. D. Charkes, The detection of bone metastases by photoscanning with radioactive strontium, in Scintillation Scanning in Clinical Medicine, J. L. Quinn, III (Ed.), p. 69, W. B. Saunders Co., Philadelphia, 1964.
15. N. D. Charkes, D. M. Sklaroff, and I. Young, A critical analysis of strontium bone scanning for detection of metastatic cancer, Am. J. Roentgenol. Radium Therapy Nucl. Med., 96: 647 (1966).
16. J. Borak, Relationship between clinical and roentgenological findings in bone metastases, Surg. Gynecol. Obstet., 75: 599 (1942).
17. D. M. Sklaroff, and N. D. Charkes, The early detection of bone metastases with Stage II carcinoma of the breast at time of radical surgery (abst.), J. Nucl. Med., 8: 272 (1967).
18. M. F. Hall, R. C. Flipse, and A. J. Gilson, Evaluation of compression of vertebral bodies employing use of strontium-85 (abst.), J. Nucl. Med., 8: 272 (1967).
19. D. G. Anderson, G. C. H. Bauer, and E. M. Smith, Pattern of distribution of SR-85 in osteoarthritis of the knee (abst.), J. Nucl. Med., 8: 273 (1967).
20. D. Van Dyke, H. O. Anger, Y. Yano, and C. Bozzini, Bone blood flow shown with F^{18} and the positron camera, Am. J. Physiol., 209: 65 (1965).

Figure 1.

Left--Roentgenogram of the pelvis of a woman after radical mastectomy for breast cancer, but with no proven metastases. This was initially interpreted as demonstrating some osteoporosis, but no metastases. Even in retrospect, after viewing the bone scan, three professors of Radiology at the University of Chicago still judged that the roentgenogram showed no lesion.

Center--Bone scan with strontium-85. An obvious lesion is seen in the right ischial ramus.

Right--Bone scan with pelvic outline for orientation--note residual activity in the sigmoid colon and cecum as well as normal uptake in the L₅ vertebra and sacroiliac joints.

The lesion in the right ischial ramus was subsequently proven when films 5 months later (following oophorectomy) showed definite reactive sclerosis about the metastasis.

TABLE 1

PROPERTIES OF BONE SCANNING AGENTS

Nuclide	^{47}Ca	^{85}Sr	$^{87\text{m}}\text{Sr}$	^{18}F
Half-life	4.7 days	64 days	2.8 hours	1.87 hours
Gamma Energy (keV)	1310	513	388	510 (B+)
Chemical Form	Chloride	Nitrate	Citrate	Fluoride
Dose (μCi)	20-40	100	250	1000-2000
Usual Time Starting Scan	2-7 days	2-7 days	45 minutes	1 hour
Patient Dose (in millirads)				
a) Bone	600	1500-4500	25	30-60
b) Body	60	700	5	225-450
Comment	Need special scanner	Slow scan times	Generator produced (from ^{87}Y)	Reactor produced <u>or</u> cyclotron produced

TABLE 2

SOME OF THE LESIONS PRODUCING POSITIVE BONE SCANS

Bone Metastases

Primary Bone Neoplasms

Fractures

Pyogenic Osteomyelitis

Spondylitis Tuberculosa

Fibrous Dysplasia

Degenerative Arthritis

Rheumatoid Arthritis (and Ankylosing Spondylitis)

Hyperostosis Frontalis Interna

Paget's Disease

Chronic Sinusitis with Osteitis

Surgical Osteotomy

Growing Epiphyses

Myositis Ossificans

Myelomatosis

Kutschalla Fig 2 Top

Bone scanning