SEP 10 1962

MASTER

Submitted to
Journal of Mathematical Physics

July 23, 1962

BNL 6294

Available from the
Office of Technical Services
Department of Commerce
Washington 25, D. C.

Facsimile Price \$ 1.10
Microfilm Price \$ 1.80

Mathematical Deductions From
Some Rules Concerning High Energy Total Cross Sections*

C. N. Yang

Brookhaven National Laboratory, Upton, New York

1. In 1956 Pomeranchuk¹ and Okun and Pomeranchuk² discussed a very interesting rule, which we shall call the Pomeranchuk-Okun rule. It states that at very high energies, in the forward direction, the exchange scattering amplitudes (such as charge exchange, strangeness exchange, etc.) are infinitesimal compared with the elastic amplitude. Two years later Pomeranchuk³ discussed another very interesting rule, which we shall call the Pomeranchuk rule. It states that the total cross section in an $A+B$ collision is the same as that of $\bar{A}+B$ at very high energies. Both rules seem to have increasing experimental support in recent years.

In this note we want to discuss the mathematical consequences of each of the two rules, when one takes into consideration the isotopic spin symmetry of the strong interactions. The results are valid only in so far as the effects of the electromagnetic and weak interactions

* Work performed under the auspices of the U.S. Atomic Energy Commission

/ Permanent address: The Institute for Advanced Study, Princeton, New Jersey

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights, or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

This paper was submitted for publication in the open literature at least 6 months prior to the issuance date of this Microcard. Since the U.S.A.E.C. has no evidence that it has been published, the paper is being distributed in Microcard form as a preprint.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

are negligible. Higher symmetries are touched upon briefly in Sec.5.

2. We shall use slightly generalized versions of the two rules: Each of the two colliding particles is supposed to be a coherent mixture of the various components of an isotopic spin multiplet, (e.g. 70 percent proton plus 30 percent neutron). The same reasons for believing the original rules support this generalized version of the rules.

3. Theorem 1: The Pomeranchuk-Okun rule implies that at ∞ energy the total cross section in a collision $A+B$ is independent of the total isotopic spin. I.e.

$$\sigma_t(A+B) = \sigma_t(A'+B')$$

if A and A' , B and B' belong to the same isotopic multiplet. Proof:

Let $a = 1, 2, \dots$, $b = 1, 2, \dots$ denote the m -quantum numbers of the particles A and B respectively. Let

$$\langle a', b' | \alpha | a, b \rangle \quad (1)$$

denote the forward scattering amplitude (times $4\pi\lambda$) from $A+B$ to $A'+B'$.

Then the Pomeranchuk-Okun rule states that

$$\langle a', b' | \alpha | a, b \rangle = \text{diagonal}.$$

The rule states in fact that (1) still holds if one changes the repre-

sentations of the particles A by a unitary transformation U and that of B by a unitary transformation V:

$$\langle a', b' | UV \alpha V^{-1} U^{-1} | ab \rangle = \text{diagonal} . \quad (2)$$

It follows easily from (2) that α is a multiple of the unit matrix.

Taking its imaginary part one obtains the theorem.

4. In this section we shall discuss the consequences of the Pomeranchuk rule irrespective of the validity of the Pomeranchuk-Okun rule.

Theorem 2: The Pomeranchuk rule is equivalent to the rule that at ∞ energy, $\sigma_J = \sum_{J'} M'_{JJ'} \bar{\sigma}_{J'}$, where $M'_{JJ'} = (-1)^{2A+2B} (2J'+1) \begin{Bmatrix} ABJ \\ ABJ' \end{Bmatrix}$, A and B stand for the total isotopic spins of particles A and B (as well as for the particles themselves), the $\begin{Bmatrix} \end{Bmatrix}$ for the 6J symbol, and σ_J , $\bar{\sigma}_J$ respectively for the total cross sections with total isotopic spin J in the collisions A+B and A+B̄. Proof: (a) Take the forward scattering amplitude matrix α of (1) and let

$$\beta = -\frac{1}{2}i(\alpha - \alpha^\dagger) . \quad (3)$$

β is thus the imaginary part of the forward scattering amplitude. Its diagonal elements are the total cross sections. We define the corresponding matrices $\bar{\alpha}$ and $\bar{\beta}$ for A+B̄ collisions. The Pomeranchuk rule states that

$$\text{diagonal elements of } (\beta - \bar{\beta}) = 0 . \quad (4)$$

Under a unitary transformation U on the states of A and a unitary transformation V on the states of B, the matrices β and $\bar{\beta}$ transform by:

$$\beta \rightarrow UV\beta V^{-1}U^{-1}, \quad \bar{\beta} \rightarrow U V^* \bar{\beta} V^{-1}U^{-1}.$$

Thus the generalized Pomeranchuk rule states that

$$\text{diagonal element of } (UV\beta V^{-1}U^{-1} - UV^* \bar{\beta} V^{-1}U^{-1}) = 0. \quad (5)$$

Since U is an arbitrary unitary matrix operating on the states of A, (5) is equivalent to

$$\langle ab | V\beta V^{-1} - V^* \bar{\beta} V^{-1} | a'b \rangle = 0. \quad (6)$$

Since V is unitary, (6) is equivalent to

$$\langle ab | V\beta V^{-1} | a'b \rangle = \langle a'b | V^* \bar{\beta} V^{-1} | ab \rangle \quad (7)$$

Since U is an arbitrary unitary matrix operating on the states of b (7) is equivalent to

$$\langle ab | \beta | a'b' \rangle = \langle a'b | \bar{\beta} | ab' \rangle.$$

Or to

$$\langle ab | \beta | a'b' \rangle = \langle ab' | \bar{\beta} | a'b \rangle. \quad (8)$$

(b) ₄ We can express β in terms of σ_j and Clebsch Gordon coefficients

$$\langle ab|\beta|a'b'\rangle = \sum_{Jm} \langle ab|Jm\rangle \sigma_J \langle Jm|a'b'\rangle. \quad (9)$$

We can also express $\bar{\beta}$ in terms of $\bar{\sigma}_J$, Clebsch Gordon coefficients and the symbols $\begin{smallmatrix} J \\ mn \end{smallmatrix}$:

$$\langle ab'|\bar{\beta}|a'b\rangle = \sum_{\substack{b''b''', \\ J'm'}} \begin{pmatrix} B \\ b'b'' \end{pmatrix} \begin{pmatrix} B \\ bb''' \end{pmatrix} \langle ab''|J'm'\rangle \bar{\sigma}_J \langle J'm'|a'b'''\rangle. \quad (10)$$

The occurrence of the symbols $\begin{pmatrix} J \\ mn \end{pmatrix}$ is due to the definition of $\bar{\beta}$ in which the rows and columns are labelled to correspond to the rows and columns of β .

(c) Substituting (9) and (10) into (8) one obtains after some straightforward algebra the relationship between σ_J and $\bar{\sigma}_J$, given by theorem 2. The use of $\begin{smallmatrix} 6J \\ 5 \end{smallmatrix}$ symbols for crossing relations was made by Wick.

We now discuss in detail the consequences of theorem 2 for the following three cases:

(A) A and $\bar{A}$ do not belong to the same isotopic spin multiplet, and B and $\bar{B}$ do not belong to the same isotopic spin multiplet. In this case we have four sets of σ_J 's: for $A+B$, $A+\bar{B}$, $\bar{A}+B$ and $\bar{A}+\bar{B}$. By repeated application of theorem 2 one obtains

$$\sigma_J(A+B) = \sigma_J(\bar{A}+\bar{B}), \quad \sigma_J(A+\bar{B}) = \sigma_J(\bar{A}+B).$$

We have made use of the fact that the square matrix M' multiplied by itself is unity, as can be easily verified from the properties of the $\begin{smallmatrix} 6J \\ 5 \end{smallmatrix}$ symbols. Furthermore, $\sigma_J(A+B)$ and $\sigma_J(A+\bar{B})$ are related by theorem 2. There are thus altogether $2I+1$ linearly independent total cross sections

where I is the smaller of the total isotopic spins of A and B .

(B) A and $\bar{A}$ do not belong to the same isotopic spin multiplet, B and $\bar{B}$ do. In this case, by definition,

$$\sigma_J(A+B) = \sigma_J(A+\bar{B}) .$$

Thus theorem 2 shows that $\sigma_J(\overline{=}\sigma_J)$ satisfies a set of linear homogeneous equations. These equations have been completely and explicitly solved. In particular the number of linearly independent total cross sections is

$$\begin{aligned} B+1 & \text{ for } A \geq B, \\ A+\frac{1}{2} & \text{ for } A < B, \quad 2A = \text{odd}, \\ A+1 & \text{ for } A < B, \quad 2A = \text{even}. \end{aligned}$$

It is obvious by repeated application of theorem 2 that

$$\sigma_J(A+B) = \sigma_J(\bar{A}+B).$$

(c) A and $\bar{A}$ belong to the same isotopic spin multiplet. B and $\bar{B}$ belong to the same isotopic spin multiplet. In this case, the same conclusions as in case (B) obtain.

5. If one takes both the Pomeranchuk and the Pomeranchuk-Okun rule, one obtains, of course, trivially that all σ_J are the same for all J and for $A+B$, $\bar{A}+B$, etc.

6. The above considerations can clearly be made also for any compact group of symmetries of the strong interactions. To obtain a general solution, however, one must first have a proper definition of the $6J$ symbols, and one must know certain symmetry and summation properties of the $6J$ symbols. This is an unsolved problem.

It is a pleasure to thank the members of the Physics Department of the Brookhaven National Laboratory for the hospitality extended to me during my visit.

REFERENCES

1. I.Ia. Pomeranchuk, J. Exptl. Theoret. Phys. USSR 30, 423 (1956).
English translation: Soviet Phys. JETP 3, 306 (1956).
2. L.B. Okun and I.Ia. Pomeranchuk, J. Exptl. Theoret. Phys. USSR 30, 424 (1956). English translation: Soviet Phys. JETP 3, 307 (1956).
3. I.Ia. Pomeranchuk, J. Exptl. Theoret. Phys. USSR 34, 725 (1958).
English Translation; Soviet Phys. JETP 7, 499 (1958).
4. See e.g., A.R. Edmonds, Angular Momentum in Quantum Mechanics, Princeton University Press (1957).
5. G.C. Wick, Brookhaven Lectures, 1960.
6. M.E. Rose and C.N. Yang, J. Math. Phys. 3, 106 (1962).
7. Cf. W.R. Sharp, Princeton University Thesis (1960).