COLUMBIA
UNIVERSITY

CORNELL
UNIVERSITY

HARVARD
UNIVERSITY

JOHNS HOPKINS
UNIVERSITY

MASSACHUSETTS
INSTITUTE OF
TECHNOLOGY

PRINCETON
UNIVERSITY

UNIVERSITY
OF
PENNSYLVANIA

UNIVERSITY
OF
ROCHESTER

YALE
UNIVERSITY

ASSOCIATED
UNIVERSITIES,
INC.

Annual Report

July 1, 1962

BROOKHAVEN NATIONAL LABORATORY

Upton, Long Island, New York

Brookhaven National Laboratory is operated under a contract between the United States Atomic Energy Commission and Associated Universities, Inc. This, the thirteenth in a series of unclassified Annual Reports, gives an account of the progress of the Laboratory during the period July 1, 1961 - June 30, 1962, and its plans for the future. It is submitted under the terms of Contract No. AT-30-2-GEN-16 between Associated Universities, Inc., and the Atomic Energy Commission.

Previous reports in this series are:

BNL 74 (AS-4) July 1, 1950
BNL 131 (AS-5) July 1, 1951
BNL 196 (AS-6) July 1, 1952
BNL 246 (AS-7) July 1, 1953
BNL 303 (AS-8) July 1, 1954
BNL 364 (AS-9) July 1, 1955
BNL 426 (AS-10) July 1, 1956
BNL 462 (AS-11) July 1, 1957
BNL 523 (AS-12) July 1, 1958
BNL 560 (AS-13) July 1, 1959
BNL 632 (AS-14) July 1, 1960
BNL 682 (AS-15) July 1, 1961

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

PRINTED IN USA
PRICE \$3.00

Available from the
Office of Technical Services
Department of Commerce
Washington 25, D.C.

November 1962

2200 copies

CONTENTS

INTRODUCTION.....	ix
-------------------	----

Physical Sciences and Engineering

PHYSICS.....	3
HIGH ENERGY ACCELERATORS.....	27
INSTRUMENTATION.....	37
CHEMISTRY.....	43
NUCLEAR ENGINEERING.....	59
APPLIED MATHEMATICS.....	82

Life Sciences

BIOLOGY.....	89
MEDICAL RESEARCH.....	105

Supporting Activities

TECHNICAL OPERATIONS AND SERVICES.....	123
ADMINISTRATION AND OPERATIONS.....	138
APPENDIX A. PUBLICATIONS.....	147
APPENDIX B. OFFICERS AND SCIENTIFIC STAFF.....	171

Brookhaven National Laboratory is operated by Associated Universities, Inc., under contract with the United States Atomic Energy Commission. Nine northeastern universities sponsor Associated Universities, Inc. Two representatives from each university, one a principal administrative or corporate officer and the other a scientist, form the Board of Trustees of AUI. In addition, the Board itself selects three Trustees-at-Large. Their purpose is to insure adequate representation of any scientific discipline in which the corporation may undertake work and to provide general flexibility in methods of operation. The composition of the Board as of June 1962 is given below. The officers of AUI are shown on the organization chart for the Laboratory (Figure 1). AUI has appointed five visiting committees to Brookhaven National Laboratory to provide a continuing independent evaluation of the research program. The committees cover the fields of biology, chemistry, medicine, nuclear engineering, and physics, and report annually to the Board of Trustees.

BOARD OF TRUSTEES

OF

ASSOCIATED UNIVERSITIES, INC.

AS OF JUNE 30, 1962

RALPH S. HALFORD, <i>Chairman</i>	Columbia University	I.I. RABI
ROBERT L. SPROULL	Cornell University	SANFORD S. ATWOOD
WILLIAM S. SWEET	Harvard University	L. GARD WIGGINS
P. STEWART MACAULAY	Johns Hopkins University	WILLIAM D. McELROY
CARL F. FLOE	Massachusetts Institute of Technology	PETER T. DEMOS
VAL L. FITCH	Princeton University	JOSEPH C. ELGIN
CARL C. CHAMBERS	University of Pennsylvania	JONATHAN E. RHOADS
ELMER H. STOTZ	University of Rochester	LARoy B. THOMPSON
C. HAMILTON SANFORD	Yale University	WILLIAM W. WATSON

Trustees-at-Large

ALFRED E. WHITFORD, Lick Observatory EDWARD M. PURCELL, Harvard University
HAROLD F. WEAVER, University of California at Berkeley

Annual Trustees

ROBERT F. BACHER LEO GOLDBERG EDWARD REYNOLDS

Honorary Trustee

FRANK D. FACKENTHAL

VISITING COMMITTEES

Biology

SEVERO OCHOA, *Chairman*
New York University

DAVID M. BONNER
University of California

ARMIN C. BRAUN
The Rockefeller Institute

MARCUS M. RHOADES
Indiana University

CARROLL M. WILLIAMS
Harvard University

RAY F. DAWSON
Columbia University

FRANCIS J. RYAN
Columbia University

ARTHUR K. SOLOMON
Harvard Medical School

Chemistry

WALTER J. KAUFMANN, *Chairman*
Princeton University

ROBERT E. CONNICK
University of California

PHILIP H. ABELSON
Carnegie Institution of Washington

WILLIAM O. BAKER
Bell Telephone Laboratories

ROBERT W. TAFT, JR.
Pennsylvania State University

CLYDE A. HUTCHISON, JR.
University of Chicago

Medicine

ROBERT A. MOORE, *Chairman*
State University of New York

JOSEPH E. SMADEL
National Institutes of Health

A. BAIRD HASTINGS
Scripps Clinic and Research Foundation

LOUIS H. HEMPELMANN
University of Rochester

W.T.S. THORP
University of Minnesota

MAXWELL M. WINTROBE
Salt Lake County General Hospital

BRONSON S. RAY
New York Hospital

WILLIAM B. WOOD, JR.
Johns Hopkins University

Nuclear Engineering

W.E. SHOUPP, *Chairman*
Westinghouse Electric Corporation

ERIC R. JETTE*
Union Carbide Corporation

CHARLES D. CORYELL
Massachusetts Institute of Technology

IRVING KAPLAN
Massachusetts Institute of Technology

HAROLD ETHERINGTON
Allis-Chalmers Manufacturing Company

STEPHEN LAWROSKI
Argonne National Laboratory

SIDNEY SIEGEL
Atomics International

ALBERT R. KAUFMANN
Nuclear Metals, Inc.

*Retired from Union Carbide Corporation in May 1962.

Physics

CLIFFORD G. SHULL, *Chairman*
Massachusetts Institute of Technology

ROBERT R. WILSON
Cornell University

R.E. MARSHAK
University of Rochester

POLYKARP KUSCH
Columbia University

RICHARD L. GARWIN
Columbia University

JOSEPH L. FOWLER
Oak Ridge National Laboratory

ASSOCIATED UNIVERSITIES, INC.

Figure 1.

ORGANIZATION CHART
JUNE 30, 1962

THE NUMBER OF EMPLOYEES IN EACH ORGANIZATIONAL UNIT ON JUNE 30, 1962, IS SHOWN; TOTAL FOR THE LABORATORY, 2593.

Dr. Leland J. Haworth delivering the dedicatory address at dedication ceremonies for the Alternating Gradient Synchrotron on September 13, 1961. Seated on the platform are (left to right) Dr. G.K. Green, Chairman, Accelerator Department; Mr. E.L. Van Horn, Manager, AEC Brookhaven Office; Dr. I.I. Rabi, President, Associated Universities, Inc.; and Dr. M. Goldhaber, Director, Brookhaven National Laboratory.

Introduction

The program and activities of Brookhaven National Laboratory during the fiscal year 1962 are described in this annual report. The progress and trends of the research program are presented together with a description of the operational, service, and administrative activities of the Laboratory. The scientific and technical details of the many research and development activities are covered more fully in scientific and technical periodicals and special reports of the Laboratory. A list of all publications published or submitted for publication during the year may be found in Appendix A.

RESEARCH PROGRAM

The Brookhaven research program, which covers a wide range of subjects in the physical and

biological sciences and in engineering, has as its central motif the development and exploitation of nuclear science and technology. It can be broadly described under five main headings:

1. FUNDAMENTAL STUDIES OF ATOMIC NUCLEI, THE PARTICLES THAT CONSTITUTE THEM, AND THE FORCES INVOLVED IN THEIR STRUCTURE. These studies, which are basic to all nuclear science and technology, involve the use of all the major machines of the Laboratory. They range from measurement of the properties of undisturbed nuclei to study of the violent disruptions resulting from nuclear fission or from bombardment with high energy particles from the Laboratory's 30-Bev proton accelerator, the new Alternating Gradient Synchrotron (AGS).

2. STUDIES OF THE PHYSICAL, CHEMICAL, AND BIOLOGICAL EFFECTS OF NUCLEAR RADIATION. The

Figure 2.

Figure 3. Infrared photograph of the gamma forest, looking toward the metal pipe housing the 9500-curie source of Ce^{137} . Close to the source the trees have been killed by chronic exposure to gamma irradiation for ≈ 6 months. In the foreground, oaks further from the source show 50% normal leafage. This experiment is designed to run for many years and to yield information not only on the sensitivity of plants to damage from ionizing radiation, but also on longer-term genetic effects, the various physiological characteristics of organisms growing in natural arrays, and such biological interactions as host-parasite relationships.

effects of radiation are utilized to determine the characteristics of physical and chemical structures and to elucidate the more complex properties of living systems and the changes they undergo in their life and reproductive cycles. Of direct practical interest are the possible deleterious effects of radiation upon biological systems and upon various materials, for example, those involved in reactor construction. Of equal interest are the

potentially beneficial effects of radiation in the treatment of disease, in the induction of genetic changes in plants or animals, and in the improvement of manufacturing processes.

3. THE USE OF NUCLEAR TOOLS, SUCH AS NEUTRONS, CHARGED PARTICLES, GAMMA RAYS, AND ISOTOPIC TRACERS, IN ALL BRANCHES OF SCIENTIFIC RESEARCH. In this broad and diversified field, which overlaps the one just mentioned, nuclear

particles and radiations are used as tools in studying physical, chemical, and biological systems in their undisturbed states. Neutrons and other particles are used as probes; by observing their penetration and scattering, minute details of physical and chemical structure can be studied. Wide use is made of isotopic tracers in many fields to yield information on such matters as the mechanisms and rates of chemical reactions and biological processes.

4. **RESEARCH AND DEVELOPMENT, NOT NECESSARILY ITSELF OF A NUCLEAR NATURE, DIRECTED TOWARD SOLVING THE PROBLEMS OF ATOMIC ENERGY DEVELOPMENT.** The more basic aspects of this category include the chemistry of elements and compounds of special interest, isotope effects, the metallurgy of materials used in nuclear reactors and other devices of importance in the atomic energy program, the neutron scattering and absorbing properties of substances used in reactors and other nuclear devices, and similar subjects. Among its applications are the development of reactor components such as fuels, structures, and shields, the development of components and processes for the chemical processing and useful recovery of reactor products, and theoretical and experimental studies in the field of reactor physics.

5. **THE DEVELOPMENT OF SPECIFIC DEVICES FOR USE AS RESEARCH TOOLS OR IN PRACTICAL APPLICATIONS OF ATOMIC ENERGY.** This category includes both the design and development of special research tools, such as accelerators, reactors, and other technical equipment, and the development of devices for practical applications of atomic energy, such as the preparation of special isotopes, the development and packaging of high intensity radiation sources, and basic development work on power reactor systems.

The involvement of the various scientific disciplines in these areas of study and development results in a coherence and mutuality of interest that enhance the entire program of the Laboratory in both tangible and intangible ways. Specific research projects are described in those sections of this report devoted to the particular discipline or organizational unit in which the work is being carried out. It will be noted that research interests overlap from one department to another, and similar problems are being attacked from different points of view.

Figure 4. X-ray photograph of a normal rat, showing most of the important skeletal landmarks used in connection with Biology Department studies of the effects of radiation on the nervous system.

RESEARCH FACILITIES

Research at Brookhaven is centered on, although not confined to, the use of several large machines and other special facilities, some of which are described briefly below.

Alternating Gradient Synchrotron

The AGS, the world's largest accelerator, operating at a nominal energy of 30 Bev, has continued to demonstrate its capabilities as an efficient, effective

tive tool for performing high energy physics research. At the end of the fiscal year, the backlog of approved experiments amounted to more than a year's running time. The continuing efforts devoted to equipment development and modification have resulted in increased beam intensity and machine reliability, with a consequent increase in the availability of the facility for experiments. Peak beam intensities of 4.0×10^{11} protons/pulse have been achieved, and intensities of 2 to 3×10^{11} protons/pulse are routine.

On January 15, 1962, operation of the AGS was increased from 2 to 3 shifts per day, 5 days a week. Of the more than 3400 hr available for machine operation during the year, over half were used for research in high energy physics. During the second half of the year, as the AGS settled down to steady operation, $>60\%$ of all available time could be utilized for research. Accelerator studies, machine start-up and shutdown, experimental setup and takedown, and machine downtime accounted for the rest.

The East Target Building and a small addition to the Service Building were completed in August 1961. The new Target Building, an L-shaped addition to the original building, provides additional experimental area needed for the efficient arrangement of the complex beam transport systems for both primary and parasitic experiments scheduled in the high energy physics program. In April 1962 the building to house the 80-in. bubble chamber was completed, and on June 11 the synchrotron was shut down for ≈ 4 months so that the main magnet tunnel wall could be breached to allow particle beams to be guided to the 80-in. Bubble Chamber Building. This work is scheduled for completion in October 1962.

Additional information on the AGS will be found in the section on High Energy Accelerators.

Cosmotron

The demand for experimental time on the Cosmotron, a 3-Bev proton synchrotron, continues to be high. The present backlog of approved experiments amounts to ≈ 1 year of running time. Efforts are continuing to increase the intensity of the beam and to make better provision for simultaneous experiments.

Except for a 4-month scheduled shutdown for machine improvements and additional experimental facilities, the Cosmotron operated con-

tinuously 3 shifts per day, 7 days a week, with ≈ 2 shifts a week scheduled for routine maintenance. The scheduled 4-month shutdown was used primarily for the installation of new straight sections in the machine; the improved magnetic field configuration has approximately doubled the beam intensity. Maximum beam intensity is now 8×10^{11} protons/pulse, the highest intensity reported by any proton synchrotron. Of the ≈ 8000 hr available for machine operation during the year, almost 65% were used for high energy physics research. Additional information on the Cosmotron is included in the section on High Energy Accelerators.

60-in. Cyclotron

The 60-in. cyclotron accelerates protons to 10 Mev, deuterons to 20 Mev, and helium ions to 40 Mev. Deflected beams up to $50 \mu\text{a}$ are available as well as internal and externally focused beams. A small external beam with momentum analysis to 1 part in 1000 is routinely used. The cyclotron has been used by more than 50 people, 5 Brookhaven departments, and 10 outside institutions. Work for outside institutions and work by guests and collaborators from outside institutions accounts for over a third of the machine time. The principal activities have involved nuclear reaction studies, mainly with the analyzed beam, production of radioactive materials for a wide variety of purposes, and the direct use of the beam for various studies of biological systems. There has been some activity directed toward future modifications and modernization.

18-in. Cyclotron

The 18-in. cyclotron accelerates protons to an energy of 3 Mev. The external beam is focused on a Zr-T target 30 feet from the machine, and monoenergetic neutrons are produced through the $T(p,n)\text{He}^3$ reaction. The beam-bunching inherent in the acceleration process in the cyclotron results in neutron pulses from the target of 1 to 2×10^{-9} -sec duration. Neutron scattering experiments are carried out which utilize the time-of-flight techniques made possible by the pulsed character of the neutron beam. A proton beam current of 130 to $150 \mu\text{a}$ is used routinely, but current as high as $200 \mu\text{a}$ has been used occasionally. The peak current in the pulse of protons can therefore be as high as 10 ma.

Electrostatic Generator

This accelerator has been operated for research on a routine basis with protons, deuterons, alpha particles, and He^3 ions, beam currents up to $75 \mu\text{a}$, and energies up to 3 Mev. During the past year the machine was operated for research during 429 8-hr shifts. The highest number of shifts in any one month was 70, and the highest in one week was 21. Approximately 10% of the time was utilized for fast neutron irradiations for the Medical and Nuclear Engineering Departments. The remainder of the time was used mainly for physics research consisting of a variety of experimental measurements on the properties of energy levels in light nuclei. Part of this work made use of the external

beam-pulsing system with which it is possible to achieve a time resolution of $2\tau = 5 \times 10^{-10}$ sec.

Graphite Research Reactor

The in-pile and beam facilities of the Brookhaven Graphite Research Reactor (BGRR) are used in a variety of programs by Brookhaven personnel and by visiting scientists and engineers from other institutions. The maximum thermal neutron flux available is $\approx 2 \times 10^{13}$ neutrons/cm²-sec. The reactor loading was increased from 575 to 607 channels to meet reactivity requirements. Fuel elements removed from the reactor averaged 41.7% burn-up of the original U^{235} .

A new graphite annealing procedure to repair graphite radiation damage was approved in May

Figure 5. Experimental area on top of the Graphite Research Reactor. In the center foreground a technician is inserting a container of seeds into the Biology thermal column, as part of the program to study radiation-induced mutations in plants. To the right of the thermal column is the drum-shaped housing for a slow-neutron chopper; at the end of the 15-ft arm is a detector. The neutron scattering angle is varied by means of the upright frame.

1962. It calls for more frequent annealings at lower temperatures and is based on a new model of the mechanism of radiation damage developed and checked experimentally at BNL.

Approximately 27 reactor holes are assigned to BNL research departments. Other reactor holes are assigned for the use of the eight outside organizations that have availed themselves of reactor facilities (other than routine irradiation services) under various cooperative arrangements.

Medical Research Reactor

The Medical Research Reactor (MRR) was constructed for the sole purpose of exploring the possible applications of nuclear reactors to the study of man and his diseases. Each salient feature of the reactor was designed in relation to its use for therapy and diagnosis, or in the advancement of basic medical science. Operation on an intermittent basis is demanded by the nature of the research program. Operating power levels up to 3 Mw have been approved for continuous operation, and levels up to 5 Mw are permitted for intermittent periods not to exceed 10 minutes.

The MRR was operated 256 times during the year for a total of 297 Mw-hr.

Hot Laboratory

The Hot Laboratory, which is adjacent to the BGRR, contains extensive facilities for the analysis, processing, and development of highly radioactive materials. It includes three hot cells in which chemical operations can be performed remotely while under observation by periscope, and a larger hot cell of the cave type for the physical examination of materials, especially metals of high activity.

Critical Assembly Laboratory

The Critical Assembly Laboratory provides specialized facilities for research in reactor physics and for reactor development studies. Prior to completion of the Critical Facility Addition at the end of the year, the laboratory consisted of two critical assembly areas, each with an operating console and assembly room; a small, water-cooled, graphite-reflected neutron source reactor for exponential studies contained in one of the assembly rooms; and a pulsed Van de Graaff neutron source for diffusion experiments. The Critical Facility Addi-

tion provides two more critical assembly cells with improved arrangements and greater flexibility for the handling of critical experiments and the necessary storage facilities, and fuel storage, office, and computational space.

MAJOR RESEARCH FACILITIES UNDER DESIGN AND CONSTRUCTION

High Flux Beam Research Reactor

Ground was broken for the Brookhaven High Flux Beam Research Reactor (HFBR) on September 22, 1961. This reactor is designed to meet the need for very high neutron fluxes, especially in the epithermal range. The HFBR is cooled and moderated by heavy water and contains a heavy water reflector; it is fueled with enriched uranium. The core consists of 28 ETR-type flat-plate fuel elements housed in a shroud in the lower spherical portion of the reactor vessel. Cool D₂O, pumped into the upper cylindrical part of the vessel, flows downward inside the shroud through the fuel elements and is discharged into the bottom of the vessel; it then flows upward outside the core and leaves from an outlet pipe at the upper part of the vessel to be cooled by heat exchange with water from cooling towers.

Outside the core vessel is a water-cooled thermal shield. This secondary vessel provides emergency containment to keep the core covered with D₂O in the event of a leak in the primary vessel. The 16 control rods are divided, with 8 main rods operating above the core and 8 auxiliary rods operating below the core. There are 6 standard horizontal beam tubes; 5 are tangent and 1 is radial with respect to the core. The radial tube will be used for fast chopper time-of-flight experiments in which neutrons of resonance energy are required. Other beam tubes are expected to be used for diffraction work. There are also 2 dual beam tubes, designed to accommodate 2 diffraction setups, and 1 cold neutron beam tube. In addition, 3 thermal and 4 fast vertical irradiation thimbles are provided.

The HFBR will be housed in a 3-story, circular, domed, gas-tight building. The bottom floor will house the operating machinery for the reactor; the second or ground floor will be reserved for beam experiments and laboratories; and the top floor will accommodate the control room, irradiation experiments, and fuel handling facilities.

This reactor, which is expected to deliver a peak thermal flux of $\approx 7 \times 10^{14}$ neutrons/cm²-sec and epithermal fluxes of ≈ 1 to 2×10^{15} neutrons/cm²-sec, will be used primarily as a source of high-intensity external thermal and epithermal neutron beams for studies in the physical sciences.

The architect-engineer work is being done by the Lummus Company under contract to the Atomic Energy Commission. The reactor is being constructed on Rutherford Hill, southeast of the BGRR's cooling tower. At the end of the year the project was approximately half finished, with completion expected late in 1963. A preliminary summary hazards report was submitted to the AEC and subsequently to the Advisory Committee on Reactor Safeguards (ACRS). After a hearing the ACRS concluded in July 1961 that the construction of a reactor of this type at Brookhaven would raise no special problems.

80-in. Hydrogen Bubble Chamber

Construction of the BNL 80-in. liquid hydrogen bubble chamber has reached the final stages. The building to house the chamber, compressors, controls, and power supplies has been completed to-

gether with outdoor auxiliary equipment such as the safety sphere, cooling tower, gas storage tanks, and power station. In June 1962 work was begun on an 86-ft-wide opening in the AGS tunnel, with a beam channel to conduct particle beams to the 80-in. bubble chamber. Fabrication and assembly of the bubble chamber itself was $\approx 80\%$ complete at the end of the fiscal year. The major components of the chamber have been delivered and are being assembled and installed. These include the magnet undercarriage (used for translating, rotating, and elevating the 450-ton magnet assembly), the steel yoke, copper coils, and vacuum chamber, which are all being assembled into two units that can be separated to permit installation and servicing of the hydrogen chamber. The 1500-liter stainless steel chamber body with safety chamber will be mounted by the 36-in.-diameter expansion cylinder and carried by one of the two magnet sections. The first two ultra-lightweight Inconel honeycomb pistons, which decouple the chamber liquid from the hydrogen gas expansion system, have been received. This system requires excellent timing controls, and exhaustive component tests have been performed to assure proper operation.

Figure 6. Construction of the containment vessel for the High Flux Beam Research Reactor, June 1962.

Figure 7. Construction of the 80-in. Bubble Chamber Building. In the foreground is the gas storage area; to the left is the cooling tower, and to the right is the safety sphere for the emergency containment of hydrogen. To the rear of the Bubble Chamber Building can be seen the AGS magnet tunnel, with the Target and Service Buildings at the upper left.

An $81 \times 30 \times 6\frac{1}{2}$ -in. tempered glass plate for the chamber window has been received and has been polished to flatness within an accuracy of a few light wavelengths. The illuminating system specially designed to avoid undesirable reflections is being assembled. This system makes use of "coat hangers," a type of shaped reflector developed at Lawrence Radiation Laboratory. The refrigeration components (for hydrogen, nitrogen, and freon), vacuum pump, and gas purification systems are being installed, and many of the hydraulic and electric controls have been prefabricated. The vacuum chamber and the four 70-mm pulsed cameras are nearing completion. The 80-in. bubble chamber is expected to be ready for operation early in 1963.

High Intensity Radiation Development Laboratory

The High Intensity Radiation Development Laboratory (HIRDL) has been completed except for special equipment which is in the process of being installed. This laboratory will be used to obtain engineering data on a variety of radiation sources in the million-curie range and to develop more efficient techniques for handling large-scale radiation sources. This information is essential for the design of future irradiation facilities needed for a wide variety of applications of radiation energy.

The total level of radiation expected to be employed in this laboratory is ≈ 2 million curies. Most of the sources will be of Co^{60} and Cs^{137} ; spent

Figure 8. Inside the 80-in. Bubble Chamber Building, showing the refrigeration platform and part of the refrigeration equipment. At the floor level is the hydraulic control panel for elevation and rotation of the entire bubble chamber structure.

Figure 9. The newly completed High Intensity Radiation Development Laboratory (HIRDL).

Figure 10. Interior views of the High Intensity Radiation Development Laboratory. Top: In the foreground is the preparation cell, and beyond it are the canal and the irradiation cell; openings for windows and manipulators can be seen in the north faces of both cells. Bottom: South faces of the preparation (left) and irradiation cells, showing door and window openings and manipulator plugs; the recess between the cells accommodates a turntable.

Figure 11. Rear view of the interior of one of the Metallurgy Hot Cells. The Model 8 master-slave manipulators are being operated from the far side of the heavy lead glass shielding window.

reactor fuel elements will also be used as sources of radiation.

The main design features of the HIRDL are two unique cells, one an irradiation cell for the experimental work with radiation sources, and the other a work preparation cell in which various types of sources will be prepared for experimental use.

GENERAL CONSTRUCTION PROGRESS

As previously mentioned, the East Target Building and a small addition to the Service Building at the AGS were completed in August 1961, the building to house the 80-in. bubble chamber was finished in April 1962, and in June 1962 work was begun on the opening in the AGS tunnel to allow beams to be guided to the 80-in. bubble chamber.

The facilities of the Nuclear Engineering Department were expanded by completion of the Critical Facility Addition, mentioned earlier, the

Metallurgy Building Addition, and the Volatility Studies Laboratory. The Metallurgy Building Addition, finished in June 1962, provides 10 general laboratories and office facilities with a total floor area of 13,800 sq. ft. The Volatility Studies Laboratory, begun in August 1961 and completed in June 1962, will be used for the development of new chemical processes for the recovery of uranium and plutonium from irradiated reactor fuel based on the volatility of the hexafluorides of these two elements. The laboratory provides almost 2600 sq. ft. of experimental area. Facilities include a small chemistry laboratory and a mechanical equipment room.

The Physics Building was completed in March 1962. It provides $\approx 104,000$ sq. ft. of laboratory and office space for activities of the Physics Department formerly housed in 10 temporary buildings. The building includes a 2-story laboratory section containing 27 general laboratories together

Figure 12. Aerial view of the new Physics Building. At the right is the seminar room.

Figure 13. Front of the new Physics Building.

with special laboratories for work in nuclear moments and solid state physics, for scanning and data analysis work connected with the emulsion and bubble chamber programs in high energy physics, and for other specialized supporting work. It is already clear that the space provided by this building will be inadequate to meet the total needs of the rapidly growing physics program, especially in the field of high energy physics.

The 1 and 2-room efficiency apartments to house visiting scientists were finished in March 1962 in time for use during the summer. These units comprise a total of 60 rooms that can be used in various combinations to accommodate 38 small family groups and supplement the quarters converted from Army medical units in 1947 which are inadequate to meet the present needs for visitors' housing.

The master building plan for Brookhaven National Laboratory is being implemented as rapidly as is practical, with first priority being given to the provision of permanent laboratories of a type and size to meet the current and future needs of the Laboratory. Buildings and facilities under design by architect-engineers at the end of the year were permanent laboratories for the Chemistry and Instrumentation and Health Physics Departments, and a small addition to the Cosmotron complex to provide laboratories, offices, and shop space. The request for 1963 now before Congress includes a Physics and Mathematics addition to the Physics Building and a Controlled Environment Laboratory for the Biology Department. Detailed planning for many other laboratories and facilities needed to meet the future needs of the Laboratory is currently under way. Brookhaven will request a continuation of this building program for future years in order that the former Army buildings still in use, which were modified after the Laboratory took over the Camp Upton site, will ultimately be replaced by suitable permanent constructions.

PERSONNEL

The total number of employees at Brookhaven National Laboratory on June 30, 1962, excluding temporary appointees, Research Collaborators, and guests, was 2593. This represents a net increase of 205, or slightly less than 8.6% for fiscal 1962. During this period, the regular salaried

scientific staff (Ph.D. or equivalent) increased from 334 to 336, Research Associates decreased from 78 to 76, and salaried visitors decreased from 31 to 27 for a net decrease in the total salaried scientific staff from 443 to 439. The nonsalaried scientific staff, however, increased from 406 in 1961 to 479 in 1962, as of May 31 of these years. These visitors are not at Brookhaven on a full-time basis. About one-fourth to one-third of them are on site at any given time. About 500 (as compared to 450 the previous year) visiting scientists and students (salaried and nonsalaried) worked at the Laboratory during the year; this number does not include Research Associates. These individuals were affiliated with 185 different institutions. The total number of professional employees (B.S. or equivalent) increased during this period from 194 to 229.

Summer visitors continue to play a substantial role in the Laboratory's program. Arrangements have been made for 270 visiting scientists and students to work at Brookhaven during the summer of 1962. Of this number, 132 are staff members from colleges, universities, industrial organizations, and other institutions, and 138 are students.

ADMINISTRATION

The organization of the Laboratory as of July 1, 1962, is given in Figure 1. On July 20, 1961, Dr. M. Goldhaber was appointed Director of Brookhaven National Laboratory. Dr. G.F. Tape, the Deputy Director of BNL, had been serving as Acting Director since Dr. L. J. Haworth's appointment to the U.S. Atomic Energy Commission in April 1961. On January 18, 1962, the Board of Trustees appointed Dr. Tape a Vice President of AUI, but he did not assume the duties of his new position full time until May 1962. On March 7, 1962, the appointment, effective May 1, of Dr. C. Williams as Deputy Director of BNL was announced. Dr. Williams had been Chairman of the Nuclear Engineering Department since its formation in 1952.

Dr. C.E. Falk was appointed Associate Director on April 1, 1962, with main responsibility in the area of programs, plans, and budgets. Effective May 1, 1962, Dr. G.N. Glasoe was appointed an Assistant Director with principal responsibilities in the area of technical services. Dr. Glasoe had been Associate Chairman of the Physics Department.

Dr. W. J. Catacosinos was appointed Business Manager of the Laboratory, effective May 21, 1962. Dr. Catacosinos replaced Mr. M.M. Small who had been appointed by AUI as Project Manager for the 140-ft radio telescope to be erected at the National Radio Astronomy Observatory at Green Bank, West Virginia.

On June 25, 1962, Dr. W.E. Winsche assumed the Chairmanship of the Nuclear Engineering Department, succeeding Dr. Williams. Dr. Winsche, who has spent the last eleven years with E.I. du Pont de Nemours and Company, mainly at the Savannah River Laboratory, was Head of the Chemistry and Chemical Engineering Division of the Nuclear Engineering Department when he left BNL in 1951.

Effective July 2, 1962, Dr. V.P. Bond was appointed Chairman of the Medical Department, replacing Dr. L.E. Farr who resigned to assume

the newly created Chair of Nuclear Medicine at the M.D. Anderson Hospital and Tumor Institute, University of Texas, Houston, Texas. Dr. Bond was previously an Assistant Chairman of the Medical Department.

FINANCE

As in the past, the Laboratory's AEC-supported research was financed by five Divisions of the AEC, namely, Reactor Development, Research (Physical Sciences), Biology and Medicine (Life Sciences), Isotope Development, and Training and Education. The operating costs associated with the work for each program are shown in Figure 14; the major categories of operating expenditures are also shown. Table 1 shows in detail the operating expenditures of the Laboratory on a broad operational basis.

OPERATING COST DISTRIBUTION BY PROGRAMS							
FISCAL YEAR	PHYSICAL SCIENCES	REACTOR DEVELOPMENT	LIFE SCIENCES	ISOTOPE DEVELOPMENT	TRAINING & EDUCATION	OTHER (NET)	TOTAL COST
1962	18,930,903	3,892,953	5,216,925	547,238	145,587	929,043	29,662,649
1961	16,560,779	3,198,809	4,845,371	381,462	97,211	584,134	25,667,766
1960	11,775,522	3,421,837	4,584,337	315,310	166,300	671,205	20,934,511

* MAJOR CATEGORIES OF OPERATING EXPENDITURES					
FISCAL YEAR	SALARIES AND WAGES	MATERIALS & SUPPLIES SUBCONTRACTS, TRAVEL & SPECIAL PROCUREMENTS	POWER	MISCELLANEOUS (NET)	OPERATING TOTAL
1962	20,881,773	6,831,784	1,047,376	901,716	29,662,649
1961	18,630,185	5,521,786	975,191	540,604	25,667,766
1960	15,125,492	4,295,397	868,432	645,190	20,934,511

Figure 14.

Table 1

Organizational Expenditures - Fiscal 1960, 1961, 1962

(Includes Operating, Services to Fixed Assets, and Work for Others. Direct Costs of AGS and Other Fixed Assets and Additions to Inventory Are Not Included; See Tables 3 & 4)

		Salaries, Wages, Insurance			Material & Supplies	Sub- contracts & Special Procure- ments		Miscel- laneous (Net)	Total Organi- zational Costs	% of Total	Man-Years		
		Staff	Consultants & Temporary Employees	Travel			Power				Scientific & Professional (Incl. Guests)	Others	Total
Physics & Chemistry Research	1962	6,703,649	229,532	163,881	2,399,307	250,803	402,092	(1,000)	10,148,264	34.2	375.0	438.0	813.0
	1961	5,831,438	191,771	159,766	2,196,171	110,902	342,521	—	8,832,569	34.4	339.5	372.0	711.5
	1960	3,867,825	202,370	129,020	1,088,434	130,681	130,386	—	5,548,716	26.5	275.0	241.5	516.5
Biology, Medicine & Biophysics Research	1962	2,493,351	86,324	81,635	603,116	770	—	(5,303)	3,259,893	11.0	140.5	194.5	335.0
	1961	2,301,162	76,928	90,853	554,921	3,642	—	(6,125)	3,021,381	11.8	135.0	183.0	318.0
	1960	2,002,118	77,678	79,408	580,596	2,562	—	(5,158)	2,737,204	13.1	129.0	164.0	293.0
Nuclear Engineering Research	1962	2,208,169	38,870	85,757	592,567	113,116	—	—	3,038,479	10.2	113.0	129.0	242.0
	1961	1,865,931	44,166	74,706	443,491	144,350	—	—	2,572,644	10.0	104.5	111.5	216.0
	1960	1,663,191	44,176	46,563	436,454	148,173	(179)	—	2,338,378	11.2	94.0	104.0	198.0
Isotope Development	1962	193,727	6,897	10,165	67,888	47,119	—	—	325,796	1.1	13.5	10.5	24.0
	1961	136,682	6,075	7,795	60,949	(2,198)	—	—	209,303	0.8	11.0	5.5	16.5
	1960	96,401	3,047	7,947	66,538	13,396	—	—	187,329	0.9	7.5	3.5	11.0
Training & Education	1962	47,838	3,419	26,789	30,185	2,093	—	(6,800)	103,524	0.4	3.5	0.5	4.0
	1961	29,635	3,643	21,167	15,291	5,664	—	(2,000)	73,400	0.3	3.5	2.5	6.0
	1960	27,180	411	32,736	44,486	4,277	—	—	109,090	0.5	2.5	3.5	6.0
Radiation Protection	1962	475,329	2,258	5,677	85,619	—	—	—	568,883	1.9	3.0	47.0	50.0
	1961	435,473	4,083	3,617	75,080	—	—	(1,115)	517,138	2.0	11.5	46.5	58.0
	1960	365,589	2,886	2,838	88,498	—	—	(1,985)	457,826	2.2	10.0	40.5	50.5
Supporting Scientific & Technical Services	1962	3,607,195	22,184	26,812	1,116,269	19,470	281,880	(129,824)	4,943,986	16.7	67.0	365.5	432.5
	1961	3,172,192	14,664	22,724	622,241	4,665	321,217	(167,786)	3,989,917	15.5	53.0	326.5	379.5
	1960	2,683,242	15,364	25,108	460,831	—	390,407	(191,770)	3,383,182	16.1	48.0	296.0	344.0
Security & Plant Protection	1962	580,958	552	876	9,841	—	—	2,331	594,558	2.0	—	75.0	75.0
	1961	596,392	833	502	11,620	—	—	1,946	611,293	2.4	—	77.0	77.0
	1960	563,172	754	795	13,371	—	—	2,894	580,986	2.8	—	77.5	77.5
Miscellaneous (including Lighting, T & T, Heating Fuels, Special Maintenance, etc.)	1962	—	—	908	183,800	(8,827)	363,404	838,735	1,378,020	4.6	—	—	—
	1961	—	—	(6,640)	172,465	—	311,453	697,911	1,175,189	4.6	—	—	—
	1960	—	—	2,416	175,618	—	347,818	728,851	1,254,703	6.0	—	—	—
General and Administrative	1962	4,053,265	51,371	151,104	572,470	15,200	—	(27,123)	4,816,287	16.2	—	563.5	563.5
	1961	3,805,697	35,614	126,328	505,433	(15,234)	—	(191,927)	4,265,911	16.6	—	539.5	539.5
	1960	3,424,033	19,715	92,011	559,725	(9,211)	—	(65,642)	4,020,631	19.2	—	507.5	507.5
Laboratory Total	1962	20,363,481	441,407	553,604	5,661,062	439,744	1,047,376	671,016	29,177,690	98.3	715.5	1,823.5	2,539.0
	1961	18,174,602	377,777	500,818	4,657,662	251,791	975,191	330,904	25,268,745	98.4	658.0	1,664.0	2,322.0
	1960	14,692,751	366,401	418,842	3,514,551	289,878	868,432	467,190	20,618,045	98.5	566.0	1,438.0	2,004.0
AUI Administration	1962	—	—	—	—	—	—	230,700	230,700	0.8	—	—	—
	1961	—	—	—	—	—	—	209,700	209,700	0.8	—	—	—
	1960	—	—	—	—	—	—	178,000	178,000	0.8	—	—	—
Total AUI and BNL	1962	20,363,481	441,407	553,604	5,661,062	439,744	1,047,376	901,716	29,408,390	99.1	715.5	1,823.5	2,539.0
	1961	18,174,602	377,777	500,818	4,657,662	251,791	975,191	540,604	25,478,445	99.3	658.0	1,664.0	2,322.0
	1960	14,692,751	366,401	418,842	3,514,551	289,878	868,432	645,190	20,796,045	99.3	566.0	1,438.0	2,004.0
Work for Others, Direct Costs Only	1962	75,079	1,806	4,015	111,789	61,570	—	—	254,259	0.9	7.5	7.5	15.0
	1961	75,896	1,910	3,388	45,537	62,590	—	—	189,321	0.7	9.0	5.5	14.5
	1960	64,051	2,289	4,841	60,127	7,158	—	—	138,466	0.7	7.5	5.0	12.5
Grand Total	1962	20,438,560	443,213	557,619	5,772,851	501,314	1,047,376	901,716	29,662,649*	100.0	723.0	1,831.0	2,554.0
	1961	18,250,498	379,687	504,206	4,703,199	314,381	975,191	540,604	25,667,766**	100.0	667.0	1,669.5	2,336.5
	1960	14,756,802	368,690	423,683	3,574,678	297,036	868,432	645,190	20,934,511†	100.0	573.5	1,443.0	2,016.5

* \$712,038 of this total was distributed to Fixed Assets and as services to Work for Others and Inventory.

** \$433,401 of this total was distributed to Fixed Assets and as services to Work for Others and Inventory.

† \$553,081 of this total was distributed to Fixed Assets and as services to Work for Others and Inventory.

Table 2
Capital Equipment Expenditures
(Including Charges From Organizational Units; See Table 1)

	FY 1962		FY 1961		FY 1960	
	\$	%	\$	%	\$	%
Scientific & hospital	6,604,023	93.6	5,220,768	92.0	2,760,196	85.2
Automotive & heavy mobile	154,852	2.2	198,411	3.5	201,234	6.2
Office machines & furniture	87,252	1.2	94,738	1.7	75,971	2.3
Shop equipment	159,313	2.3	136,201	2.4	142,141	4.4
Miscellaneous	46,661	0.7	24,462	0.4	62,842	1.9
Expenditures, Total	7,052,101	100.0	5,674,580	100.0	3,242,384	100.0
Proceeds from sales	(12,665)		(22,036)		(39,958)	
Expenditures, Net	7,039,436		5,652,544		3,202,426	

Table 3
Costs Incurred for Fixed Assets
(Including Charges From Organizational Units; See Table 1)

	FY 1962			FY 1961			FY 1960		
	Man-years			Man-years			Man-years		
	Costs, \$	Sci.	Others	Costs, \$	Sci.	Others	Costs, \$	Sci.	Others
<u>Alternating Gradient Synchrotron</u>									
Direct									
Salaries, wages, insurance	—	—	—	192,317	4.5	20.0	1,248,309	36.0	119.0
Materials, construction, etc.	1,226,016			2,486,877			2,249,780		
Subtotal direct	1,226,016			2,679,194			3,498,089		
Charges from organizational units	22,231			65,834			253,025		
Total	1,248,247			2,745,028			3,751,114		
<u>Other, Including High Flux Beam Research Reactor and 80-in. Bubble Chamber</u>									
Direct									
Salaries, wages, insurance	776,466	32.5	49.5	454,041	24.0	28.0	277,001	15.0	11.0
Materials, construction, etc.	10,079,560			7,174,253			4,446,120		
Subtotal direct	10,856,026			7,628,294			4,723,121		
Charges from organizational units	476,206			230,004			160,256		
Total	11,332,232			7,858,298			4,883,377		

The increase in total expenditures for fiscal 1962 is associated with a larger staff and with the acquisition of materials, supplies, and equipment, especially for high energy research. Of the total increase in expenditures for operations and equipment, $\approx \$3,000,000$, or $\approx 65\%$, was associated with the high energy physics program. This particular increase coincided with the first full year

of AGS research operation. Table 2 shows expenditures for capital equipment. Here again most of the increase is for scientific equipment essential to the program in high energy physics. A summary of expenditures for fixed assets (plant and equipment) is presented in Table 3. Table 4 sets forth inventories for which the Laboratory is responsible.

Table 4

Inventory at Close of Fiscal Year

Type of inventory	Fiscal year		
	1962	1961	1960
General stores*	\$ 704,424	\$ 524,004	\$ 448,487
Precious metals and radium	96,683	84,648	87,668
Stable isotopes	38,793	37,644	23,413
Heavy water	683,039	672,330	679,535
Total	\$1,522,939	\$1,318,626	\$1,239,103

*The number of months investment was 3.5 in 1962, 3.0 in 1961, and 3.2 in 1960.

BROOKHAVEN NATIONAL LABORATORY
COMPARATIVE BALANCE SHEET

	June 30, 1962	June 30, 1961
<u>Assets</u>		
Cash	\$ 435,663	\$ 260,753
Accounts receivable	187,917	88,071
Advances and prepaid expenses	113,667	89,930
Deposits	76,008	71,388
Property, plant, and equipment (less reserves of \$39,176,059 at June 30, 1962, and \$33,089,120 at June 30, 1961)	83,524,884	76,589,179
Construction in progress	17,851,202	9,414,288
Research materials and supplies	6,142,313	6,129,188
Total assets	<u>\$108,331,654</u>	<u>\$92,642,797</u>
<u>Liabilities</u>		
Accounts payable	4,301,914	2,922,184
Accrued payroll	177,688	214,270
Atomic Energy Commission	103,852,052	89,506,343
Total liabilities	<u>\$108,331,654</u>	<u>\$92,642,797</u>

NOTE: Although the Laboratory has custody and use of the assets, title remains vested in the United States Government.

**CONFERENCES, PUBLIC INFORMATION,
AND EDUCATION**

The third annual series of George B. Pegram Lectures was delivered at the Laboratory on October 2, 3, 5, and 6 by Dr. Charles A. Coulson of Oxford University. These four public lectures, with the title "The Scientist and Society," dealt with the general subject of the scientist's responsibility in politics and human relations. Because of a change in schedule, the fourth in the series of Pegram Lectures was also held during the fiscal year, on June 19, 22, 27, and 29. Professor Derek

J. de Solla Price, Chairman of the Department of History of Science and Medicine at Yale University, was the lecturer on this occasion. His subject, under the general title "Little Science, Big Science," was the changing position of the scientist in a situation in which research had frequently evolved from a solitary occupation into big, cooperative projects. Books based on both the above lecture series will be published by Columbia University Press.

Eleven formal conferences were conducted at the Laboratory during the year. Among these were the Seventh AEC Air Cleaning Conference,

October 10-12; the Sixth Conclave in the Series on Nuclear Energy in Medicine, October 23-24; an International Conference on Neutron Thermalization, April 30-May 2; Brookhaven Biology Symposium Number 15, on Enzyme Models and Enzyme Structure, June 4-6; and a Molecular Beams Conference, June 11-13.

In addition to these meetings, the Laboratory served as host organization for the 1961 International Conference on High Energy Accelerators, September 6-12, sponsored jointly by the International Union of Pure and Applied Physics, the US AEC, and AUI. The first part of the conference, from September 6 through 9, was held in New York City, in the theater of the Barbizon-Plaza Hotel. On the evening of the 10th, the 200-odd participants were transported to Brookhaven for the final two days of the conference. On September 13, dedication ceremonies for the AGS were held in the new section of the AGS experimental building. Appropriately, the dedicatory address was delivered by Dr. Leland J. Haworth, Commissioner, US AEC, who was BNL's Director from 1948 until April 1961. In the latter capacity, he was largely responsible for the design and construction of the AGS.

At the Laboratory's invitation, the annual winter conference of the Science Teachers' Association of New York State (STANYS) was held at Brookhaven, January 25-27. This was the first time that STANYS, in the 66 years of its existence, had held its annual conference on Long Island. The attendance of over 700 primary and secondary science teachers from all parts of the State was the highest on record. Brookhaven scientists participated in many of the program sessions, and tours of the major research facilities were arranged for the visitors.

The Ninth Annual Naval Reserve Nuclear Science Seminar, on The Effects of Radiation on Human Populations, and Civil Defense and Management of Nuclear Casualties, was attended by 40 reserve officers, September 18-20.

The twelfth annual Visitors' Day for the general public, held on October 21, was attended by 7135 individuals. The eighth High School Visitors' Day was held on October 28 and was attended by 4273 juniors and seniors from 151 high schools. The fourth College Visitors' Day for junior and senior college classes, held on October 27, was attended by 1445 students representing 74 colleges and uni-

versities. On February 8, the Laboratory held its eighth Science Youth Day in observance of Thomas Alva Edison's birthday. Two hundred sixty-five selected science students and editors of school papers from 35 Suffolk County high schools responded to the invitation.

During the year, 787 persons representing 26 professional and technical groups paid one-day visits to the Laboratory to view its facilities and to learn about its research program. Additional visitors included 330 high school and college teachers receiving special training in the nuclear sciences at summer institutes sponsored jointly by the National Science Foundation and the AEC. Four groups of graduate students, comprising 42 individuals, visited BNL to discuss with scientists research projects in their fields of interest. More than 200 professional, governmental, and industrial representatives from foreign countries made short visits during the year. In addition, two Japanese study teams, the Reactor Study Group and the Special Committee for Science and Technology of the Japanese House of Representatives, were at the Laboratory on October 17 and November 21, respectively.

A recently completed Exhibit Hall is proving extremely useful as an adjunct to the special tours arranged for technical and professional visitors to the Laboratory. Among the exhibits on display here are a number of illuminated panels showing various aspects of Brookhaven's research program, a small operating spark chamber, and a reactor model. This hall is also available for meetings and seminars and can serve as a reception center for large groups of visitors.

The Public Information Office arranged for two formal press conferences during the year. One was held in New York City during the International Conference on High Energy Accelerators. The other, on June 20, was held at Brookhaven to enable science reporters to obtain detailed advance information on the joint Columbia-Brookhaven discovery of the existence in nature of two different types of neutrino, which was to be announced in the July 1, 1962, issue of *Physical Review Letters*. Both Brookhaven and Columbia scientists were available at the conference to describe the experiment and the significance of the discovery. In spite of the long lead time, all the news media honored faithfully the July 1 release date, and the story was covered accurately and adequately. In the field

of education, the activities of the Public Information Office continued to grow, mainly because of the increasing emphasis on the nuclear sciences in school science curricula. This is producing a heavy demand for literature and other advice and as-

sistance on science projects and term papers. The Laboratory continued to provide speakers for neighboring civic, educational, and semiprofessional audiences; 29 staff members responded to such requests during the past year.

PHYSICAL
SCIENCES
AND
ENGINEERING

Physics

The research program in physics consists of experimental and theoretical studies concerned with the structure and fundamental properties of matter. Various features of the complex structure and properties of matter can be most effectively studied by observing the interactions of charged particles, neutral particles, and radiation with matter. Some of these studies deal with the interactions involving individual atoms, atomic nuclei, or nucleons, and others with conglomerates of atoms and molecules in bulk matter. The Alternating Gradient Synchrotron (AGS), Cosmotron, 60-in. cyclotron, Van de Graaff accelerator, 18-in. cyclotron, and the Graphite Research Reactor at Brookhaven provide a wide range of energies and diversity of particles and radiation with which to carry out experimental investigations. The researches reported here will be described under the categories of particle physics, nuclear structure, neutron physics, atomic and molecular physics, and solid state physics. The theoretical scientists conduct their investigations in close association with the experimental scientists and provide stimuli for new approaches to the problem.

The work of Brookhaven scientists is enhanced by the presence of a number of visiting and guest scientists on leave from other institutions in this country and abroad. These visitors and guests are attracted to Brookhaven not only by the availability of its facilities but also by the opportunity to collaborate with Brookhaven scientists. The cross-fertilization of ideas and experience resulting from the presence of these visiting scientists is very important to an active research program.

PARTICLE PHYSICS

Research at the Cosmotron

Because Brookhaven National Laboratory was established to provide facilities for visitors from other institutions as well as for physicists connected with the Laboratory, the attack on the problems of the structure of matter is necessarily diverse. Some general trends are, however, clear. The problem of the nuclear force continues to receive

much attention via a study of pion-nucleon and pion-pion interactions. The latter are difficult to study because of the instability of the pion. The investigation of the resonant states between pions and between pions and protons is an indirect approach that seems to hold promise.

These resonant states appear as peaks in the spectra of the total production cross sections. Resonances, whether or not they are considered as "particles," have the characteristic that a state with well-defined quantum numbers (such as spin, isospin, and parity) must be the major contributor to the peak. Experiments that permit these quantum numbers to be directly measured or deduced are thus vital in obtaining information about the particles. The energy value at which the resonance peak appears and its width establish the mass and lifetime of the "particle." At present, most of the parameters are not well known, and a concerted attempt is being made to provide this information.

The experiments next described are the work of Brookhaven groups using the BNL 20-in. hydrogen bubble chamber.

The reactions $\pi^- + p \rightarrow n + \pi^+ + \pi^-$ and $\pi^- + p \rightarrow p + \pi^0 + \pi^-$ were investigated with 1.25-Bev incident pion energy. A ρ^0 resonance was seen in the first reaction, and a ρ^- resonance was seen in the second. Evidence for isospin state $I=1$ and spin state $J=1$ was shown, and the pion-pion cross sections were evaluated as a function of energy by means of the Chew-Low theory. Subsequent study of the data has shown that the ρ^0 resonance can be understood as interference between the ρ -meson and a two-pion state of the ω -meson.

The ρ^+ -meson was observed in the reaction

$$p + p \rightarrow d + \pi^+ + \pi^0,$$

using 2-Bev protons, which is further and definitive evidence for the isospin state $I=1$. It follows from symmetry arguments that the spin state is $J=1$. Another peak in the pion mass distributions appeared at ≈ 565 Mev, but, because of background and low statistics, it cannot be considered as established. If this finding is supported by further ob-

servations, it is consistent with the ζ particle, whose existence was suggested by workers at Saclay.

Another resonance, the three-pion η , was measured in the reaction

Dalitz plots were made and were shown to be consistent with the quantum numbers (isospin = 0, odd spin parity, and even G parity), but more data are required to make a definitive assignment.

In proton-proton collisions at a kinetic energy of 2.85 Bev in the laboratory system, an analysis of the four-charged-prong events has shown clear evidence for production of the resonant-state ω -mesons that appear in the reaction $p + p \rightarrow p + p + \omega$, where the ω decays very quickly into $\pi^+ + \pi^- + \pi^0$. By plotting the effective mass of the last three particles, the ω -resonance is found near 785 Mev with a half-width of ≈ 15 Mev, in agreement with observations previously obtained in antiproton and meson collisions with protons. Other resonances, such as ρ , η , ξ , and α , were not observed in this experiment. The cross section for ω production in proton-proton collisions at 2.85 Bev is found to be $48 \pm 13 \mu\text{b}$.

Another approach to the π - π interaction is made through the analysis of pion-nucleon scattering and pion production in pion-nucleon and nucleon-nucleon collisions. The isobar and one-pion exchange (OPE) models continue to provide phenomenological mechanisms for the analysis. According to the isobar model, the proton is raised to a metastable "isobaric" state by the collision and emits a pion when it decays to the ground state. The OPE theory, on the other hand, treats the process as an interaction of the incident particle with a single pion in the peripheral pion "cloud" surrounding the nucleon. Again the BNL 20-in. bubble chamber has been the tool used by Brookhaven experimenters in exploring these models.

In the reaction $p + p \rightarrow p + n + \pi^+$ observed with incident protons at 2-Bev kinetic energy, the neutron momentum spectrum shows a strong peak due to the $\frac{3}{2}, \frac{3}{2}$ isobar. The observed variation of cross section with energy is given fairly well, but not perfectly, by the Selleri equation (derived from OPE). It is further shown, about as well as can be expected from the statistics, that the scat-

tering of the virtual pion is like that of a real pion.

These p - p interactions were also studied with regard to momentum spectra, angular distributions, and cross sections; results for elastic scattering, Λ production, and single-pion production were obtained. The data were analyzed in relation to the extended isobar model and to OPE. Single-pion production was dominated by the $\frac{3}{2}, \frac{3}{2}$ isobaric resonance. The center-of-mass nucleonic angular distributions were peaked forward and backward, a distribution to be expected in peripheral collisions. In the reaction producing $pn\pi^+$, the neutron kinetic energy spectrum computed from the OPE theory agreed with the observed spectrum in events of low four-momentum transfer, but was too large in events of high four-momentum transfer.

Similar measurements of multiple-pion production were reported and showed fair agreement with the extended isobar model. The $\frac{3}{2}, \frac{3}{2}$ resonant state was shown clearly, but effects of some other state, possibly $I = \frac{1}{2}$, were also evident. In double-pion production, nucleon angular distributions showed the forward-backward peaking expected of peripheral collisions, but were nearly isotropic in triple-pion production.

In $p + p$ collisions at 1.5 Bev single π -meson production was investigated. The expected $p + \pi^+$, $T = \frac{3}{2}$ resonance was observed, as well as a strongly forward-backward peaked distribution of the inelastically scattered nucleons. (This peaking increases further at higher energies, as expected if the mesons are made in peripheral collisions.) The data were analyzed kinematically on the basis of the OPE model, and no disagreement was found. Production of deuterons was relatively high, with a cross section of almost 0.7 mb from all available reaction channels.

Angular distributions of the particles in the events contained in the ρ peak from 1.25-Bev $\pi^- + p$ interactions reported above were analyzed for consistency with the demands of the OPE model, according to the criteria developed by Treiman and Yang. It appears that OPE does not account entirely for ρ production, but the strength of the $\frac{3}{2}, \frac{3}{2}$ isobar at this incident energy introduces ambiguities into the analysis. Measurements at other incident energies are required to provide an answer to the interesting question of whether or not some other mechanism than isobars or OPE is influencing the results.

A Brookhaven counter group in collaboration with a University of Rochester group have observed pion production from proton-proton collisions at 2.9-Bev incident energy. They have obtained detailed momentum spectra of positive and negative pions at several angles. The shape of the spectra provides a critical test for the isobar and OPE models. There is good evidence for the dominance of the well-known $T = J = \frac{3}{2}$ isobaric level in single-pion production. The angular distribution is sharply peaked in the forward direction, and close agreement with the isobar model predictions can be obtained. Further, some evidence for the participation in multiple-meson production of the higher energy isobaric states of the $T = \frac{1}{2}$, π - p system is also found. The calculated total cross sections are in agreement with results from Brookhaven bubble chamber work.

With the aid of a specially designed threshold Čerenkov counter, another Brookhaven counter group working with the neutral beam at the Cosmotron have demonstrated that a peak exists in the energy spectrum of neutrons having the same energy as the primary protons (3 Bev). This peak arises from charge exchange and is clearly separated from the inelastically scattered neutrons. From theoretical predictions it is expected that the cross section will have a $1/E^2$ energy dependence, where E is the center-of-mass energy of the two particles. Attempts in the past to identify decisively these nucleon charge exchange events in the Bev energy region have not been successful. The angular distribution of the elastic events is sharply peaked in the forward direction. A preliminary estimate of the charge exchange cross section at 3 Bev is ≈ 1 mb. This in principle is the most straightforward measurement of the one-meson charge exchange at these energies and therefore has evoked much interest, especially because of the present theoretical work on peripheral interactions. Further work is in progress.

The Cambridge Bubble Chamber Group have partially completed their analysis of neutral pion production by 1.14-Bev/c π -meson collisions with protons. In order to observe the π^0 , they have used a methyl iodide-loaded propane chamber which has a high efficiency for conversion of the gamma rays from π^0 decay into electron pairs. The differential cross section found for the charge exchange reaction, $\pi^- + p \rightarrow \pi^0 + n$, satisfied the triangular inequalities required by charge independence and

showed behavior quite similar to that of the angular distribution found in elastic $\pi^- + p$ scattering. The integrated cross section of this reaction was found to be 6.4 ± 0.6 mb. In the case of interactions with no visible secondaries it was concluded that the branching ratio of one- π^0 , two- π^0 , and three- π^0 production was 1:0.3:0.05.

A team of collaborators from Brookhaven, Yale University, and Indiana University used the 14-in. BNL hydrogen bubble chamber to observe interactions of π^- -mesons with protons at pion kinetic energies of 771 and 761 Mev above and below the Λ - K^0 threshold. At both energies the 3,3 π - N isobar is prominent. The ζ -meson is not apparent at either energy in the π^- - π^0 or in the π^+ - π^- effective mass distributions.

The study of the production and decay of the strange particles – the kaons and hyperons – provides another important approach to the understanding of the structure of matter. This type of study permits a determination of their intrinsic properties: charge, rest mass, spin, lifetime, decay modes, isospin, strangeness, magnetic moment, and parity. The existence of pion-hyperon resonances similar to the pion-nucleon resonances was predicted on the basis of global symmetry, and such resonances have been observed. Analysis of such resonances is useful in determining the properties of the strange particles.

Confirming theoretical predictions, the associated production of hyperons and heavy mesons in pion-proton collisions was first observed in the BNL diffusion cloud chamber in 1953. This important subject has received much attention since then. Recently a group from BNL have studied the process in considerable detail at lower incident π -meson energies than heretofore. The 20-in. BNL hydrogen bubble chamber was used for the experiment. The $\Lambda + K^0$ production cross section, when plotted at the four energies 775, 791, 829, and 871 Mev (which were chosen immediately above the threshold for the reaction $\pi^- + p \rightarrow \Lambda + K^0$, but below the threshold for $\pi^+ + p \rightarrow \Sigma^0 + K^0$), does not show the abrupt rise expected if production occurred in an s -state only. The observed very gradual rise from zero at threshold indicates that higher states are also involved. Angular differential cross sections were then fitted by partial wave analysis. Only s and p -waves were necessary for good fits except at the energy of 829 Mev, where d -waves (angular momenta with $\ell=2$) also had to

be used. This behavior may indicate a $\Lambda + K^0$ resonant state at ≈ 1650 Mev. Analysis is being continued on this important point.

A University of Michigan counter group have verified the existence of the Υ_1^{0*} resonance at 1.385 Bev in an investigation of the reaction $\pi^- + p \rightarrow \Upsilon_1^{0*} + K^0$, $\Upsilon_1^{0*} \rightarrow \Lambda + \pi^0$. The decays of the Λ and K^0 were observed in a large, thin-plate spark chamber triggered by a counter array. The half-width appeared to be not much more than the experimental resolution of 17 Mev.

Analysis of the first successful measurement of the magnetic moment of the Λ -hyperon has been completed by a group from BNL, ANL, MIT, and NYU. Polarized Λ -hyperons were produced by π^+ -mesons incident on a beryllium target in the reaction $\pi^+ + n \rightarrow \Lambda + K^+$. A pulsed high-field magnet was used to precess the Λ 's, which were observed to decay in a parallel-plate spark chamber triggered by a counter logic system to select the desired events. Because of the nonconservation of parity, the Λ decay has an asymmetry in the decay products with respect to the direction of the spin polarization. This asymmetry indicates the direction of polarization to which the magnetic moment is either parallel or antiparallel. The magnetic field and flight path were such that a magnetic moment of 1 Bohr nucleon magneton would precess 16.5° in a plane perpendicular to the Λ trajectory. Measurement of the angle of rotation of the decay pattern with the field on, off, and reversed gave the magnitude of the moment. The sense of rotation yielded the sign. The direction of the magnetic moment was found to be antiparallel to the direction of polarization. The experiment yielded a value $\mu_\Lambda = -1.5 \pm 0.5$ Bohr nucleon magnetons. A more precise measurement is being scheduled. While no truly reliable theoretical prediction of the magnetic moment exists with which to compare the experimental value, the unexpectedly large and negative value of the Λ magnetic moment is of considerable interest in a theory of strong interactions and should provide a valuable check point for the testing of new ideas.

The Λ -decay parameters themselves have been the subject of a searching experiment by a Princeton University group. A new technique made the experiment simple and straightforward but yielded precise results. π^- -Mesons incident on protons produced Λ 's in a spark chamber array. The polarization of the Λ -decay proton was observed

by the scattering in large carbon plates. They found that the decay asymmetry parameter $\alpha = -0.62 \pm 0.07$, which gives the proton a positive helicity in contradiction to the direct application of the universal Fermi interaction to the Λ decay. They also found that the decay proceeds predominantly through the s -wave channel and that the results are in agreement with time reversal invariance. This experiment should lead to an increased understanding of nonleptonic decays of hyperons.

A beta-decay angular correlation experiment involving the decay of the K -meson has been performed by another Princeton University group. It is now well known that the beta-decay process in strangeness-conserving interactions is of basically vector nature. It was the purpose of the experiment to determine whether or not the strangeness-nonconserving decays were of the same type. By using counter techniques the angular relations between the π^0 and the electron in the decay process $K^+ \rightarrow e^+ + \pi^0 + \nu$ were investigated. The results indicate that the interaction is vector. The decay is unique in beta decay in that extraordinarily high four-momentum transfers are involved. No measurable dependence of the interaction on the four-momentum transfer was observed.

A group from the University of Pennsylvania used a special technique to obtain information on the decay of the positive kaon. They employed a chamber composed of layers of scintillation filaments viewed by an image intensifier and photomultipliers to provide oscilloscope displays of the time sequence and spatial relations of events occurring in the chamber. Sections of the chamber were separated by gold plates to provide sufficient stopping power for the products resulting from the kaon decay. The measurement of the range spectrum of the decay muon in the $K_{\mu 3}$ decay resulted in a determination of the ratio of the form factors $\xi = f^-/f^+$. Previously ξ had been found from branching ratio measurements to be in either interval $-9.7 < \xi < 7.7$ or $1.0 < \xi < 3.0$. This measurement gave $\xi = -9$ with 95% certainty. The experimental spectrum is consistent with the assumption that the form factors are not strongly energy dependent.

A hitherto unobserved rare decay mode of the Λ particle was discovered in a photograph taken in the BNL 20-in. hydrogen bubble chamber by a group from Columbia University and Brookhaven.

The decay is $\Lambda \rightarrow p + \mu^- + \bar{\nu}$. The chamber had been exposed to a beam of 900-Mev π^- -mesons.

A method of gaining information about cosmic rays has recently been developed in the study of meteorites. A group from the Carnegie Institute of Technology have used the 3-Bev Cosmotron proton beam to irradiate heavy targets to aid in the analysis of nuclear effects of cosmic rays in meteorites.

Eight percent of the Cosmotron's experimental time has been devoted to radiochemistry experiments, and a small fraction has been used for animal studies by the Medical Department. Groups from more than 17 universities, in addition to BNL groups, participated in the program.

Research at the Alternating Gradient Synchrotron

One of the most important experiments in recent years in high energy physics was performed at the AGS this year. The Columbia-Brookhaven group established a high intensity, high energy neutrino beam to study neutrino interactions.

Pauli first postulated the existence of the neutrino in 1930 in an effort to salvage the fundamental conservation laws of energy and momentum which were apparently violated by nuclear beta decay. Fermi's beta-decay theory using Pauli's neutrino was a beautiful mathematical solution to the problem. It satisfactorily accounted for all the observed facts by endowing the neutrino with zero rest mass, zero charge, and intrinsic angular momentum $= \hbar/2$. These characteristics and the fact that the neutrinos participate only in weak interactions have made them extremely elusive from an experimental point of view. It was only in 1955 that workers from Los Alamos Scientific Laboratory were able to directly observe the effects of neutrino interactions. One of the most interesting characteristics of weak interactions is the fact that whenever a muon is involved in such an interaction, so is a neutrino or an antineutrino. The same holds true for the electron. Thus, if either an electron or a negative muon is absorbed by a nucleus, a neutrino is emitted. In this sense both electrons and muons seem to be explicitly coupled to neutrinos in the weak interactions. In recent years, however, a major difficulty has arisen. Calculation has shown that if both the muon and electron are connected with the same type of neutrino, a muon should spontaneously decay some fraction of the time into an electron and a gamma ray. The branching ratio should be

$$\frac{\mu \rightarrow e + \gamma}{\mu \rightarrow e + \nu + \bar{\nu}} \approx \frac{1}{10^4}.$$

However, the gamma-decay mode has never been seen, although $>10^8$ decays have been experimentally observed. Feinberg, Lee, and Yang have pointed out that the most natural explanation for this difficulty is that the neutrinos coupled to muons are not the same as the neutrinos coupled to electrons. If this were true, then the muon would never decay into an electron and a gamma ray.

A clear test of the identity of the neutrinos is provided by observing the capture by nuclei of neutrinos arising from the decay of pions. If these neutrinos are the same as the neutrinos involved in beta decay, they should produce as many electrons as muons. If they are different, they can in principle produce only muons. The reactions are

$$\nu + n \rightarrow \mu^- + p \text{ or } e^- + p,$$

$$\bar{\nu} + p \rightarrow \mu^+ + n \text{ or } e^+ + n.$$

This experiment has only become feasible with the advent of the 33-Bev AGS and 28-Bev CERN accelerators, which can provide beams of high energy neutrinos of sufficient intensity (the interaction cross section of neutrinos increases rapidly with neutrino energy). The experiment at the AGS was performed by using a ten-ton spark chamber shielded from all particles but neutrinos by ≈ 5000 tons of iron. The running time amounted to ≈ 800 hr, with the AGS delivering $\approx 4 \times 10^{17}$ protons in the primary beam. The integrated flux through the spark chamber of $\approx 10^{14}$ neutrinos resulted in only 50 interactions producing energetic events. Of these, 24 showed only a single energetic muon being produced, while the remainder showed muons being produced along with other particles. In no case was a single energetic electron produced. It is therefore concluded that the neutrinos arising from pion decay are different from those involved in beta decay.

This elegant experiment has demonstrated the value of the AGS in providing beams of high energy neutrinos as a probe to investigate the nature of weak interactions. It is undoubtedly the forerunner of other work with high energy neutrinos and opens up an entirely new and exciting field of investigation.

The other major effort at the AGS was the electromagnetic separated beam program. High intensity beams of antiprotons, kaons, and antikaons

Figure 1. A view inside the heavily shielded room containing the ten-ton spark chamber. Each of the chamber's 10 units consists of 9 aluminum plates $44 \times 44 \times 1$ in. thick separated by $\frac{3}{8}$ -in. Lucite spacers and is driven by a specially designed high-pressure spark gap. The spark chamber is triggered by scintillation counters. Anticoincidence counters are provided on top, in back, and in front of the chamber to reduce the effect of muons from the AGS and cosmic rays that penetrate the heavy shielding.

Figure 2. A neutrino event as observed in the spark chamber. The long lower track is interpreted to be a muon. It is accompanied by what appears to be an electron pair from a gamma ray, pointing back to the origin.

Figure 3. A photograph of the event that resulted in the discovery of the predicted Ξ^- . The event $\bar{p} + p \rightarrow \Xi^- + \Xi^+$ was observed in the 20-in. liquid hydrogen bubble chamber. The accompanying sketch identifies the tracks according to the most likely mass interpretation.

of high purity became available and were enthusiastically utilized by experimenters. These beams stimulated renewed effort in the search for resonances of all kinds, besides providing a new attack on the measurement of the properties and production mechanisms of strange particles. The 20-in. hydrogen bubble chamber took more than 700,000 photographs during this program. Analysis of most of these experiments has only started, but some of the early results are given below.

The first observation of a $\Xi^- + \Xi^+$ cascade hyperon pair was made at the AGS, and simultaneously a similar observation was made at the CERN laboratory in Europe. This confirmed the existence of the Ξ^- , one of the last predicted antiparticles. Two such events have now been observed in the $\bar{p} + p$ exposure, the cross section probably

amounting to at most a few microbarns. This was a joint experiment with Yale University in which hydrogen was exposed to a beam of 3.25-Bev/ c antiprotons.

A larger number of $\Lambda + \bar{\Lambda}$, $\Sigma^+ + \bar{\Sigma}^+$, and $\Sigma^- + \bar{\Sigma}^-$ pairs has also been observed, as well as examples of $\Lambda + \bar{\Sigma}^0$, $\bar{\Lambda} + \Sigma^0$, Υ (hyperon) + $\bar{\Upsilon}$ (antihyperon) + (π or 2π), and $K + n + \bar{\Upsilon}$ or $K + \bar{n} + \Upsilon$ production. The cross section for the reaction $\bar{p} + p \rightarrow \bar{\Lambda} + \Lambda$ was found to be $\approx 94 \mu\text{b}$ with an uncertainty of at least $20 \mu\text{b}$. Most striking is the very strongly forward-peaked angular distribution of the $\bar{\Lambda}$ (and correspondingly backward-peaked distribution of the Λ) in the center-of-mass system. This distribution is clearly inconsistent with a production model invoking the exchange of a single pseudo-scalar K -meson. Production of $\Sigma^+ + \bar{\Sigma}^+$ pairs shows a sim-

ilar trend, with the $\bar{\Sigma}^+$ traveling forward in the center-of-mass system. Since the mechanisms for production of $\Sigma^+ + \bar{\Sigma}^+$ and $\Sigma^- + \bar{\Sigma}^-$ may be quite different ($\Sigma^+ + \bar{\Sigma}^+$ can be produced by single K exchange, $\Sigma^- + \bar{\Sigma}^-$ cannot), further study of these reactions resulting in charged hyperons is of particular importance.

In the same exposure events are being analyzed for elastic and inelastic interactions (including annihilations) resulting in pions, hyperons, and K -mesons. The total $\bar{p} + p$ cross section at the mentioned momentum is $\sigma_{\text{tot}} = 74.2 \pm 1.5$ mb, and the elastic cross section is $\sigma_{\text{el}} = 22.9 \pm 1.1$ mb. Annihilation resulting in two charged prongs amounted to $\leq 13 \pm 1$ mb, mostly consisting of pion production; without annihilation, pion production amounted to $\geq 8 \pm 1$ mb. The angular distribution of the elastically scattered antiprotons is strongly peaked forward, although a few antiprotons were scattered backward in the center-of-mass system. By use of the simple optical model and of the optical theorem for the forward amplitude, an interaction radius of 1.3 fermi, an opacity of 0.96 ± 0.05 , and a total inelastic cross section of $\sigma_{\text{inel}} = 46.8$ mb, in agreement with the observed value of $\sigma_{\text{inel}} = 51.3 \pm 2.6$ mb, were calculated.

Analysis of events with four charged prongs led to cross sections for annihilations resulting in four, five, or more than five pions, of 0.85 ± 0.16 , 4.92 ± 0.38 , and 10.86 ± 0.60 mb, respectively. All other four-prong events (nonannihilation and strange particle production) amounted to 4.61 ± 0.37 mb. The forward-to-backward charged pion ratios for four, five, and seven-pion production were 1.44 ± 0.25 , 1.20 ± 0.10 , and 1.11 ± 0.06 , respectively, with the negative charges tending to travel forward and the positive charges backward in the center-of-mass system. It is thus indicated that even in such very strong interactions as $\bar{p} + p$ annihilations some characteristics of the initial conditions concerning charge are preserved. Finally, a search for resonant states between pions revealed small effects amounting to at most 2.5 ± 0.5 mb for ρ^0 , 2.5 ± 0.4 mb for $\rho^\pm$, and 0.4 ± 0.1 mb for ω production. Lower limits of a few hundred microbarns for production of these resonant states were also established. No evidence for new resonant states has been found so far.

In a joint experiment with Syracuse University, hydrogen was exposed to K^- -mesons at energies of 2.24 and 2.5 BeV/c. Seventy-four charged cascade

hyperons, Ξ^- , were seen to decay into $\Lambda + \pi^-$. Eleven more were inferred from analysis. These were produced in the reactions $K^- + p \rightarrow \Xi^- + K^+$, $\Xi^- + K^+ + \pi^0$, and $\Xi^- + K^0 + \pi^+$. With this sample the Ξ^- mass (1321.0 ± 0.5 MeV) and lifetime [$(1.16 \pm_{0.17}^{0.26}) \times 10^{-10}$ sec] were redetermined more accurately than in previous observations. The data are consistent with a spin of $\frac{1}{2}$ for the Ξ^- , although a spin of $\frac{3}{2}$ is not ruled out. An asymmetry parameter $\alpha_\Xi = 1.0 \pm_{0.33}^{0.0}$ is found, which indicates strong parity nonconservation in the Ξ^- decay. This indicates at the same time that the Λ from the Ξ^- decay is completely longitudinally polarized. In addition it was found that (1) charge conjugation is violated in Ξ^- decay, and (2) time reversal invariance is indicated. These results are similar to results for the other hyperons. The ratio of $\alpha_\Lambda/\alpha_\Xi = -0.6 \pm 0.3$ disagrees with a prediction of some weak interaction theories making use of "global symmetry."

In the same exposure a leptonic decay of either a Σ^- or a Ξ^- into a $\Lambda + e^- + \bar{\nu}$ was observed. Interpretation as leptonic decay of a Ξ^- is favored by odds amounting to roughly 70 : 1. But, whether a Ξ^- or a Σ^- is indeed involved, the event is of equal interest because of the unusual combination of decay products.

In the $K^- + p$ exposure at 2.24 BeV/c a number of resonant states between π -mesons, K -mesons, and strange particles were observed. Two new resonant states were indicated. One is a $K + \bar{K}$ resonance occurring in the reactions $K^- + p \rightarrow \Lambda + K^0 + \bar{K}^0$ and $\Lambda + K^+ + K^-$. The resonant state decays into $K^0 + \bar{K}^0$ or $K^+ + K^-$ with an effective mass of ≈ 1020 MeV and a full width of this resonance of 40 MeV. These values are still somewhat tentative. The other new resonance was found in the reactions $K^- + p \rightarrow \Xi^- + \pi^0 + K^+$ and $\Xi^- + \pi^+ + K^0$, where $\Xi^* \rightarrow \Xi^- + \pi^0$ at 1535 MeV with full width of 50 MeV.

Confirming previous results, the Y^* decaying fast into $\Sigma^+ + \pi^-$ (effective mass, 1520 MeV), and into $\Lambda + (\pi^+ \text{ or } \pi^- \text{ or } \pi^0)$ at 1390 MeV was seen. Also observed was the $\omega \rightarrow \pi^+ + \pi^- + \pi^0$ (at 785 MeV) and the $K^* \rightarrow K^0 + \pi^-$, with a mass of 900 MeV, which is ≈ 15 MeV higher than that found previously.

$K^- + p$ interaction cross sections at 2.24 BeV/c were determined for many of the open reaction channels. Of particular interest is the relatively large cross section for $K^- + p \rightarrow (\Lambda \text{ or } \Sigma) + K + \bar{K}$,

which has a lower limit of $\approx 180 \mu\text{b}$, and the large cross section for $K^- + p \rightarrow \Xi^- + K^+$ or $\Xi^- + K^+ + \pi^0$ or $\Xi^- + K^0 + \pi^+$, with the combined value of $218 \pm 20 \mu\text{b}$. The latter reaction proceeds in $\approx 30\%$ of the cases through the resonant state $\Xi^* \rightarrow \Xi^- + (\pi^0 \text{ or } \pi^+)$. In addition it was found that the Ξ^- from $K^- + p \rightarrow \Xi^- + K^+$, the two-body production mode, are polarized, while those from the three-body modes are not. The angular distribution of the Ξ^- in the center-of-mass system is strongly peaked forward, reminiscent of the distribution of Σ^- -hyperons produced by π^- on protons.

A Brookhaven group studied interactions of 4.7-Bev/c π^- -mesons in hydrogen. Cross sections for associated production of strange particles, hyperons, and K-mesons have been studied. The different possible channels ($\gamma^0 + K^0$, $\gamma^0 + K^\pm$, and $K^0 + K^\pm$) are represented approximately equally, with lower limits for the partial cross sections from 0.1 to 0.3 mb. Several cases of negative cascade hyperon (Ξ^-) production were also seen.

In the same set of photographs the interactions resulting in four charged prongs were searched for two and three-body combinations exhibiting resonances. In the reaction $\pi^- + p \rightarrow p + \pi^+ + 2\pi^-$, the well-known (at lower energies) $T, J = \frac{3}{2}$ isobar involving the proton and the π^+ -meson was found at the effective mass value of $1220 \pm 20 \text{ Mev}$, with a total width of $90 \pm 20 \text{ Mev}$. In addition the ρ^0 resonance between π^+ and π^- was found at $740 \pm 30 \text{ Mev}$ with a width of $170 \pm 30 \text{ Mev}$, which confirmed similar results obtained at much lower energies. A small excess of events having an effective mass at 400 Mev for two π 's was observed, which suggests a new resonance. Besides these resonances, in the reaction $\pi^- + p \rightarrow p + \pi^+ + 2\pi^- + \pi^0$ the $\rho^\pm$ resonance ($\pi^\pm + \pi^0$) is seen as well as the characteristically sharp-peaked ω involving 3π -mesons ($\pi^+ + \pi^- + \pi^0$). Finally, in the reaction $\pi^- + p \rightarrow n + 2\pi^+ + 2\pi^-$, no ρ resonance is observed, but instead evidence is found for a resonance between π^+ and π^- near 400 Mev. If confirmed, the latter resonance is new and must be added to the already established ρ , ω , and η resonant states.

Many other groups used the 20-in. bubble chamber to obtain exposures which are still being analyzed. A large fraction of these exposures were obtained parasitically while the beam was being used for the neutrino experiment. This chamber, which makes possible high precision measure-

ments because of its special design and controls, has produced ≈ 2 million pictures since it was first put into operation.

Simultaneously with the work of the 20-in. bubble chamber, the BNL 14-in. hydrogen bubble chamber was set up in a neutral beam, and a Brookhaven group obtained 200,000 photographs of K_2^0 particle interaction with protons while operating parasitically with the neutrino experiment. A preliminary analysis indicates that this beam was very rich in kaons. It was possible to make a direct observation of the decay $K_2^0 \rightarrow \pi^+ + \pi^- + \pi^0$. This decay had hitherto been inferred from observation of the charge pions in kinematic analysis.

A Brookhaven counter group measured the total interaction cross sections for π^+ and π^- -mesons and protons from 8 to 20 Bev/c. These measurements are of interest in connection with the Pomeranchuk high energy limiting theorem, which states that at high enough energies particle and antiparticle total cross sections in matter become constant and equal to each other. Similar work was done last year on the p - p and $\bar{p}$ - p cross sections and on the K and $\bar{K}$ - p cross sections. Comparison of the cross sections of the π^+ - π^- particle-antiparticle pair is especially interesting since, in addition to the general Pomeranchuk theorem mentioned above, another specific argument can be applied to the behavior of the difference $\sigma_t(\pi^- + p) - \sigma_t(\pi^+ + p)$ at high enough energies but cannot be applied to the behavior of the difference $\sigma_t(\bar{p} + p) - \sigma_t(p + p)$ or the difference $\sigma_t(K^- + p) - \sigma_t(K^+ + p)$. This argument, proposed by Okun and Pomeranchuk, states that at high enough energies $\sigma_t(\pi^- + p) = \sigma_t(\pi^+ + p)$. The argument does not depend on the constancy of cross sections at high energies, which is a necessary requirement of the Pomeranchuk theorem. Furthermore, it does not depend on the validity of the forward dispersion relations.

The results show that both $\sigma_{\text{total}}(\pi^+ + p)$ and $\sigma_{\text{total}}(\pi^- + p)$ decrease monotonically between 4 and 20 Bev/c. The difference in the two cross sections does not appear to change appreciably throughout this energy region, with a $\Delta\sigma \approx 2 \text{ mb}$. The best empirical fit to the $\sigma_{\text{total}}(\pi^- + p)$ is found to be $\sigma_t = a + b/p$, where a and b are constants and p is the momentum of the pion. Thus it is seen that neither of the pion cross sections seems to satisfy the Pomeranchuk theorem at this energy.

Considerable interest has been shown recently in cross section experiments from the point of view

of the S -matrix theory as developed by Chew, Frautschi, Udgaonkar, and others. This theory relates the behavior of the high energy total and elastic cross sections to the residues of the so-called Regge poles in the S -matrix which correspond to particles and resonances. Based on this theory, it has been shown that the sum of the $\pi+p$ and $\pi-p$ total cross sections has a $1/E$ dependence, i.e., $\sigma_{\text{total}}(\pi^+p) + \sigma_{\text{total}}(\pi^-p) = a + b/E$, which is in agreement with the experimental results.

Further analysis has been done on the survey of secondary particle beams completed last year. The data show interesting features of the process of production in the new energy range made available by the AGS accelerator. For example, momentum transfers from 30-Bev collisions are small. The production mechanism is highly asymmetric in the center-of-mass system, and the production of antiprotons and K^- -mesons is much more copious at 30 Bev than at 6 Bev. In addition, a relatively large abundance of deuterons is produced at 90° .

The surprisingly low cross section for high energy collisions with large four-momentum transfers is being further investigated. This type of collision should reveal the presence of a central core in the nucleon. The proton structure in proton-proton elastic scattering is being studied by a Cornell University-Brookhaven group. Their attention has been focused upon elastic scatterings in which very large momentum transfers are involved. Such collisions are rare and correspond to nonperipheral, small impact parameter interactions. The elastic cross section has been determined at a momentum transfer of 2.5 Bev/ c . An upper limit of $5 \times 10^{-35} \text{ cm}^2/\text{sterad}$ has been established at 3.3 Bev/ c . These data, together with recent measurements at CERN, show that the p - p cross section is varying rapidly with the momentum transfer. In contrast, the variation as a function of the primary energy is considerably less in extent. This evidence is suggestive of a hard core proton structure.

As an alternative framework, the p - p strong interaction may be analyzed in terms of the trajectories of Regge poles. It is then found that the dominant trajectory ceases to decrease and approaches $\alpha = -0.5$ at large momentum transfers.

The new 80-in. liquid hydrogen bubble chamber facility is being made ready to go into operation by the end of 1962. Concurrently, higher

energy separated beams are being designed. Well-separated antiproton beams up to 8.5 Bev/ c and $K^\pm$ -meson beams between 3.0 and 5.5 Bev/ c (here the lower limit is set by the relatively short lifetime of the $K^\pm$) of useful intensity will be available. Experiments with such beams in hydrogen and deuterium will, for instance, open up many new channels for production of antihyperons, including anticascade-hyperons, for further study. Also, in addition to already discovered resonances between pions and pions, and pions and strange particles, resonances between strange particles alone (KK , YK , $\bar{Y}K$, etc.) can be studied further. The large size of the 80-in. chamber will aid greatly in these studies with respect to statistics and accuracy obtainable.

Data processing for the BNL Bubble Chamber Group has been accomplished by means of nine scanning machines and five semiautomatic coordinate-measuring machines. The IBM 7090 computer now installed at BNL has greatly facilitated this work. Programs are being written continually to make more and more effective use of this machine. The main delays in the flow of data are now encountered in the measuring phase. Many additional measuring machines would be required to improve this situation sufficiently. A new fully automatic device, the Hough-Powell Flying-Spot Digitizer (FSD), which is nearing completion, is expected to help eliminate the bottleneck. A 20- μ light spot scans the film for bubble images, which are then located by comparison with the number of lines crossed on a grating by a coherent light spot. The other coordinate is found from the position of the precision measuring stage that advances the film while it is being scanned. Thus a typical photograph can be measured in 6 sec. A prototype FSD was extensively tested in collaboration with the CERN and Berkeley laboratories during the first half of the fiscal year. The Brookhaven 35-mm FSD for analysis of 20-in. chamber film is in the final stages of assembly and testing. The electronic circuits for direct transmission of coordinates from the machine to the IBM 7090 computer have been finished and in the main tested. The mechanical spot generator and high-pressure mercury arc for illumination are operating properly. A new high-speed film transport has been completed and partially tested. Some tasks remaining are the installation of the Moire-fringe system for measurement of displacements of the

precision stage, the testing of the system for reading picture numbers from a binary code on the film, and trials of the system for computer control of stage and film.

During the first year or two, measurements by the FSD will be guided by rough measurements taken at scanning tables equipped to record coordinates of desired points (hereafter called rough-digitized scan tables). The first such table has been completed and is undergoing trials. An advanced form of rough digitizer is under development by the Instrumentation Division and if successful will be used to equip subsequent scan tables. Five additional rough-digitized scan tables are to be provided for the 80-in. chamber film.

A 70-mm FSD for analysis of 80-in. chamber film is in the late stages of design and is planned for completion early in 1963. This machine embodies several distinct advantages in optical and mechanical design discovered during tests of the prototype.

Computer programs for gating and filtering the high flux of precision coordinates from the FSD are largely complete. They are nonetheless expected to require appreciable debugging and revision in the light of experience with real measurement runs. Very considerable bookkeeping and physics problems are involved in simply understanding, organizing, and making deductions from a great many analyzed events. The Berkeley computing system FOG-CLOUDY-FAIR is particularly designed to handle large numbers of events, and this system is therefore being adapted to handle results from the Brookhaven bubble chambers.

Much work is being done, mostly by members of the Applied Mathematics Division, in pure "pattern recognition," i.e., organization of the FSD coordinate information by the IBM 7090 without human guidance. Results to date are encouraging and may lead to the transfer of some scanning functions to the computer somewhat earlier than expected.

The Theoretical Physics Group at Brookhaven devotes much of its attention to particle physics. A considerable fraction of the work has been concerned with the interactions and decay properties of particles produced in high energy reactions. The aim of this work has been to understand experimental results on K -mesons, hyperons, ω and η -mesons, etc., obtained at Brookhaven and other laboratories.

A general method of analysis into angular momentum states has been developed, applicable to relativistic three-body reactions.

Dispersion theory techniques have been applied in the following cases:

1. It was shown that a consistent picture of the π spectrum in τ decay emerges if the effects of the known p -wave π - π interactions are included in an otherwise standard treatment.

2. The θ^+/θ^0 branching ratio was calculated. For a large class of $\pi^+\pi^0$ s -wave scattering phase shifts it is easy to get agreement with the observed branching ratio. In particular, if the ζ -meson exists and has spin parity 0^- , the predicted theoretical result agrees with the experimental ratio to within $\approx 30\%$.

3. It was shown that a simple explanation of the cross sections for large momentum transfer in the reactions $p+p \rightarrow \pi+p+p$ and $\pi+p \rightarrow \pi+\pi+p$ emerges if a pionlike resonance exists in, say, the 3π spectrum. As a candidate for this object (Π), the bump in the experimental data of Pickup, Robinson, and Salant was suggested. The decay modes and probable branching ratios for its decay were listed. Its strength of coupling to the nucleon was shown to be approximately the same as that of the pion.

4. Radiative decay of neutral vector bosons was considered.

Other work of a related nature was concerned with electromagnetic-mixing effects between the ω and ρ -meson, which may have a bearing on recent experimental results.

Resonant-state calculations have been made on the so-called Lee model. The possibility of a resonant state V^* in N - θ scattering in the Lee model has been examined. (The existence of both V and V^* yields an eigenvalue equation for the V^* mass which can be solved numerically. This should lead to further extensions of the model.)

Preliminary estimates were made of corrections to the energy levels of mu mesic atoms.

Mass corrections to the ratio of the hyperfine splittings of the metastable ($2s$) state and the ground state of hydrogen atoms were computed by using a modified Dirac equation and the Foldy-Wouthuysen transformation. The procedure was shown to be equivalent to the covariant calculation.

NUCLEAR STRUCTURE

The researches in this field may be divided into two broad classifications: those dealing with studies

of the characteristics of unstable (radioactive) nuclei, and those concerned with the characteristics of the instantaneous products of a nuclear reaction. Studies of the ways in which an unstable nucleus is produced and reverts to a stable nucleus, through the radioactive decay process, yield information on the characteristics of the energy levels of the nuclei involved. Characteristics such as the spin, parity, and lifetime are sought and analyzed to obtain the systematics of the decay process. Researches dealing with observations of the instantaneous products of a nuclear reaction also yield information about the characteristics of energy levels of nuclei. Some energy levels can be observed both in radioactive decay and in nuclear reaction studies, while others can be observed only in nuclear reactions, because of their very short lifetimes. The ways in which nuclear energy levels may be excited and the systematics of the level structure, in isotopes consisting of different numbers of protons and neutrons, are the basis for the formulation of theories of the fundamental structure of atomic nuclei.

Considerable effort has been expended on the improvement of the delayed coincidence technique for the direct measurement of very short lifetimes of excited nuclear states, and upon the understanding of the factors that contribute to the finite time resolution of fast coincidence systems. It has proved possible to observe in detail the exponential decay of nuclear states with half-lives of tenths of millimicroseconds with accuracies of a few percent.

The fast coincidence technique has been applied in the direct determination of the reduced $E2$ gamma-ray transition probability of the transition between the $4+$ and $2+$ rotational states in a few strongly deformed nuclei. The strong coupling model predicts a value of $10/7$ for the ratio of reduced transition probabilities

$$\frac{B(E2) 4+ \rightarrow 2+}{B(E2) 2+ \rightarrow 0+}.$$

Experimentally measured lifetimes and values of this ratio for Dy^{162} , Er^{166} , Er^{168} , and Hf^{180} are listed in Table 1.

The decay of long-lived ($T_{1/2} > 5$ yr) Ag^{108m} has been investigated. Conversion electrons of a highly converted isomeric transition (31 ± 1 kev) are followed by an 80-kev transition to the Ag^{108} ground state. This transition is identified as electric dipole

from a measurement of its K -shell conversion coefficient. The intensity of the isomeric branch of Ag^{108m} is 8.5%. The remaining 91.5% of the decays from the isomeric state take place by electron capture to a state at 1776 kev in Pd^{108} . The energy release in this electron capture branch has been determined to be 45 ± 5 kev from the measured fraction (0.65) of electron captures that take place from the K -shell. Directional correlation measurements involving the three cascade gamma rays of 727, 617, and 432 kev that follow the electron capture are consistent with the spin sequence $6 \rightarrow 4 \rightarrow 2 \rightarrow 0$ for the cascade from the 1776-kev state to the ground state of Pd^{108} . It has been known for some time that the first and second $2+$ states in Pd^{108} occur at 432 and 940 kev respectively, while a $0+$ state occurs at ≈ 1030 kev. Recent Coulomb excitation results at other laboratories make it probable that the spin 4 state observed in this decay is de-excited by the 617-kev gamma ray, which places it at 1040 kev. It thus appears that the excitation spectrum of Pd^{108} exhibits a closely spaced triplet near 1 Mev with spins $0+$, $2+$, and $4+$. This may be interpreted as the two-phonon triplet predicted by the near-vibrational model of collective nuclear motion.

Considerations involving the electron capture rate, the probable $M4$ nature of the 31-kev isomeric transition in Ag^{108m} , the $E1$ nature of the 80-kev transition, and the decay characteristics of 2.3-min Ag^{108} suggest a $6+$ spin and parity assignment for Ag^{108m} and a $6+$ spin and parity assignment for the 1776-kev state in Pd^{108} .

A six-gap Orange beta-ray spectrometer of the Copenhagen design, whose components were purchased in Denmark by the United States Army, is being assembled and instrumented. Since this type of spectrometer combines very high transmission with moderate resolution, the instrument will be especially useful for electron-gamma-ray coincidence studies and will complement other spec-

Table 1

Isotope	$T_{1/2}(2+)$ ($\times 10^{-9}$ sec)	$T_{1/2}(4+)$ ($\times 10^{-10}$ sec)	$\frac{B(E2) 4+ \rightarrow 2+}{B(E2) 2+ \rightarrow 0+}$
Dy^{162}	2.25 ± 0.11	1.32 ± 0.08	1.42 ± 0.11
Er^{166}	1.83 ± 0.2	1.2 ± 0.1	1.42 ± 0.17
Er^{168}	1.90 ± 0.1	1.21 ± 0.08	1.46 ± 0.12
Hf^{180}	1.53 ± 0.06	0.75 ± 0.10	1.44 ± 0.20

trometers now in use at this Laboratory. This instrument will be used cooperatively with Frankford Arsenal personnel at Brookhaven.

A study has been made of the Mössbauer absorption spectrum of Fe^{57} impurity atoms in beryllium. For this purpose, a source of Co^{57} diffused into Cu provided a single-line emission spectrum with a width approximately equal to the natural line width of the 14.4-keV gamma ray. The absorption of this line by Fe^{57} bound as an impurity atom in polycrystalline beryllium showed, after annealing, a pattern suggestive of a quadrupole doublet. The application of a transverse magnetic field of 8600 gauss broadened the absorption lines asymmetrically, a result consistent with the quadrupole interpretation. The broadening was largest for the higher energy absorption line. When the quadrupole interaction energy is large compared to the magnetic interaction energy, the process of averaging over all angles of the magnetic field direction with respect to the direction of the electric field gradient at the Fe^{57} nucleus in polycrystalline beryllium causes the $m = \pm \frac{1}{2}$ state to broaden more; therefore it is concluded that it is the state of highest energy. Consequently, the quadrupole interaction term, e^2qQ , is negative for Fe^{57m} in a beryllium lattice.

These conclusions have been reinforced by studies of the absorption pattern resulting when a beryllium single crystal containing Fe^{57} impurity atoms is oriented with the c -axis either perpendicular or parallel to the direction of the incident gamma-ray. The expected intensity ratio for absorption into the $\pm \frac{3}{2}$ state as compared to that into the $\pm \frac{1}{2}$ state is $\frac{1}{3}$ for the perpendicular case and $\frac{2}{3}$ for the parallel case. Experimentally observed ratios are close to these values, which supports the interpretation that the $m = \pm \frac{1}{2}$ states lie higher.

An external magnetic field of 8600 gauss has been applied parallel to the c -axis of the beryllium single crystal. The magnetic splitting of the $m = \pm \frac{3}{2}$ state then is considerably larger than that for the $m = \pm \frac{1}{2}$ state. This observation proves conclusively that the $m = \pm \frac{3}{2}$ state is the lower energy state in the quadrupole pattern, and that e^2qQ is indeed negative.

An estimate may be made of the quadrupole moment Q of the 14.4-keV state of Fe^{57m} by combining the measured quadrupole interaction, $e^2qQ = -(5.57 \pm 0.19) \times 10^{-8}$ ev, with a theoretical

calculation of the pure beryllium ion contribution to the electric field gradient and with a reasonable estimate of the Sternheimer antishielding factor ($\gamma_{\text{Fe}} = \gamma_{\infty(\text{Fe})} = -15$). This results in the value $Q_{\text{Fe}^{57m}} \cong +2 \times 10^{-25} \text{ cm}^2$.

The study of level schemes in the osmium-iridium region has been continued. Progress was made with the construction and analysis of the level scheme of Os^{186} , which was studied in cooperation with the Chemistry Department. A ground-state band 0, 2, 4, 6, 8 (even parity), a " $K=2$ " band 2, 3, 4, 5, 6, 7 (even parity), and a number of additional levels have been established. These are shown in Figure 4. The ratios of the transition probabilities between two levels of the two bands were analyzed in terms of the Davydov asymmetric top model and of the rotation-vibration interaction method. It can be shown that the latter is mathematically almost identical with the former, and that the best fit to the data is achieved with an asymmetry parameter ($\gamma = 12^\circ$) considerably smaller than that derived from the analysis of the level energies (16.5°). This is in contrast to the case of Os^{190} in which approximately the same γ ($\approx 21^\circ$) fits the energies and transition probabilities.

Figure 4. Levels of Os^{186} populated in the decay of Ir^{186} . On the left are shown the levels of the ground-state ($K=0$) band; in the center, levels of the gamma ($K=2$) band; and on the right, additional levels whose character is not completely clear at this time.

Figure 5. Proposed decay scheme of Sb^{122m} . Figure 6. Plot of the isomeric cross section ratios ($\sigma_{\text{low spin}}/\sigma_{\text{high spin}}$) vs A of the target nuclei.

The most energetic positron branch was found to have a maximum energy $E = 1.94 \pm 0.02$ Mev and to feed the $6+$ state of the ground-state band. Hence a value of 3.83 Mev is obtained for the total disintegration energy of 15.5-hr Ir^{186} .

Several theoretical groups are working on an extension of the asymmetric top model to odd- A nuclei, but experimental information on odd- A osmium nuclei is so far scarce. In an experimental study of the energy levels of Os^{189} its parent, Re^{189} , has been produced for the first time. This new isotope has a half-life of 23 hr and decays by emission of beta particles with a maximum energy of 980 keV.

A better understanding has been achieved of isomeric transitions in Sb^{122} . The decay scheme,

as now established, is shown in Figure 5. In the past year the apparent two-step decay of the 4.2-min isomer was studied and reported. Because a half-life of 4.2 min for the 75-keV transition from the 136-keV state is several orders of magnitude greater than known lifetimes of similar $E2$ transitions, it was proposed that this lifetime might actually belong to an additional low energy transition. Since then, at another laboratory, a half-life of 530 ± 40 μsec has been measured for the 136-keV state produced by the $\text{Sb}^{123}(\gamma, n)\text{Sb}^{122m}$ reaction. Another group of workers has recently observed a 19.5-keV electron in the decay of Sb^{122m} . These results have been confirmed here. The electron line has been identified as the L -conversion line of a 25-keV transition and has been shown by means of delayed coincidence methods to precede the 530- μsec transition.

Isomeric cross section ratios for neutron capture reactions have been measured for the even-even target nuclei Te^{120} , Te^{130} , Ce^{128} , Pt^{196} , and Hg^{196} . In addition, the cross sections for the reactions $\text{Pt}^{194}(n, \gamma)\text{Pt}^{195m}$ and $\text{Pt}^{192}(n, \gamma)\text{Pt}^{193m}$ have been measured. The even-even target nuclei were chosen to avoid any ambiguity in spin of the compound nucleus, i.e., in each of these cases the compound nucleus has a spin and parity of $\frac{1}{2}+$. It is found that isomeric cross section ratios ($\sigma_{\text{low spin}}/\sigma_{\text{high spin}}$) tend to increase at the magic neutron and proton numbers. Away from the magic numbers, these ratios obey an empirical rule which states, "The isomeric cross section ratios for various even-even isotopes in the same element remain approximately the same provided the spins of the compound nuclei and of the isomeric states are the same for all the isotopes." The trends of these and other results are illustrated in Figure 6.

The investigation of the gamma rays emitted following thermal neutron capture in various elements has continued under the auspices of Brookhaven National Laboratory and the Aeronautical Systems Division of the United States Air Force. In this experimental setup a small sample of the material being studied is placed in a thermal neutron beam outside the reactor, and coincidences are measured between high energy capture gamma rays detected by a three-crystal pair spectrometer and lower energy gamma rays detected by a single scintillator. Recent improvements in the experimental arrangement, particularly lower background counting rates and improvement of

the resolution of the pair spectrometer to 2.5% in the energy range from 3 to 10 Mev, have increased the quality of the results and the scope of possible experiments.

Studies of the capture gamma rays of $U^{238} + n$ have led to the construction of an energy level scheme for U^{239} . Two components of the strong 4.06-Mev transition (previously found to consist of at least three individual transitions) populate levels whose observed properties suggest that they may be the $\frac{1}{2}-$ and $\frac{3}{2}-$ states of a band resulting from the coupling of the $\frac{1}{2}+$ (631) single-particle level with a $K=0$ octopole excitation of the type found in neighboring even-even nuclei. The possibility of the existence of such states in odd- A nuclei had previously been suggested.

Coincidence studies of the capture gamma rays of scandium ($Sc^{45} + n$), coupled with precision measurements of the capture gamma rays and (d, p) experiments, both carried out at MIT, have led to a level scheme for Sc^{46} .

Coincidence studies of $Se^{76} + n$ and $Se^{77} + n$ carried out with samples enriched in Se^{76} and Se^{77} have permitted the assignment of individual capture gamma rays to Se^{77} and Se^{78} . (Previously existing information was obtained only with the use of the natural element, so that the isotope of origin of many capture gamma rays was unknown.) Level schemes have been constructed for Se^{77} and Se^{78} .

It is well known that, because of the uncertainty principle, a spontaneous nuclear decay may proceed in a two-step process via a virtual state. Since such a process is expected to have a very low probability, it can only be detected if competing transitions are highly retarded.

Ni^{59} was selected as a promising case for experimental study (Figure 7). The direct electron capture from the Ni^{59} ground state to the Co^{59} ground state is second forbidden, the lifetime of Ni^{59} being 8×10^4 years in spite of the ≈ 1 -Mev decay energy available. On the other hand, the transition from the Ni^{59} ground state to the first excited state of Co^{59} is allowed.

A careful study of the gamma spectrum of Ni^{59} was made; the results are consistent with the assumption that the virtual e^- capture indeed takes place and that its intensity is of the order of 1% of the intensity of the inner bremsstrahlung associated with the direct decay. A more definite conclusion from this experiment must await more

Figure 7. Decay scheme of Ni^{59} . The dotted line connecting the ground state of Ni^{59} with the 1.100-Mev state of Co^{59} indicates the virtual orbital electron capture process.

precise knowledge of the shape of the second forbidden inner bremsstrahlung.

The experiment to determine the beta-neutrino angular correlation coefficient in the decay of He^6 , conducted under the Columbia University-BNL cooperative program, is nearing completion. Preliminary results indicate an angular correlation coefficient of $\lambda = -0.29 \pm 0.06$. This is compatible with an axial vector interaction in the beta decay of He^6 . A final value of the correlation coefficient to an accuracy of ± 0.02 will soon be obtained.

Detailed studies have been carried out with the Van de Graaff accelerator on the properties of energy levels in the nuclei Be^{10} , B^{10} , C^{14} , and N^{14} . Part of this work made use of the intermediate-image pair spectrometer to measure the radiative transitions from Be^9 and C^{13} targets bombarded with deuterons. It was shown that the ground-state transitions from the 6.18-Mev level in Be^{10} and from the 6.58-Mev level in C^{14} are not Doppler-shifted. This requires lifetimes for these states greater than the slowing-down time of the recoil product nuclei and suggests electric monopole radiation. Conclusive proof of the $E0$ nature of the C^{14} transition and the consequent assignment of a spin and parity of $0+$ to the 6.58-Mev state were obtained by a comparison of the internal and external pair conversion spectra where it was observed that gamma-ray emission does not occur.

The pair spectrometer was also used to determine the parity of the 2.14-Mev first excited state of B^{11} by a determination of the internal pair conversion coefficient of the transition. The state was excited by the $B^{11}(p, p')B^{11*}$ reaction. It was shown that the transition is predominantly magnetic dipole and that the parity of the 2.14-Mev state must therefore be odd.

In another phase of these studies the lifetime of the Be^{10} 3.37-Mev first excited state was measured

by means of two newly developed Doppler-shift techniques. One method utilized a fixed NaI detector to measure the energies of gamma rays emitted first from an upstream target and then from a downstream target located in a long target tube. By measuring the forward-to-backward energy shift of the 3.37-Mev radiation from targets of beryllium and of BeCu bombarded with deuterons, the lifetime of the level was obtained in terms of the characteristic slowing-down times of the Be^{10} nucleus in beryllium and in copper. In the second method two NaI counters were used, and the 3.37-Mev radiation was detected in coincidence with 2.6-Mev gamma rays from the 5.96-Mev state of Be^{10} . By holding one counter in a fixed position and varying the other in angle with respect to the beam, a difference in the shifts of the 2.6 and 3.37-Mev radiations was observed. This led to a value for the 3.37-Mev-level lifetime in terms of the slowing-down time. The results of both methods were in agreement. The lifetime of the 3.37-Mev state of Be^{10} is 1.4×10^{-13} sec.

Lifetime measurements on first excited states of B^{10} and C^{15} have been carried out by means of the 3-Mev Van de Graaff beam-pulsing facility. The technique consists of electrostatic deflection of the beam by an external radio-frequency oscillator, simultaneous terminal pulsing to reduce background effects, and the use of the so-called slanted target method to produce short time resolution. A resolving time of $2\tau = 5 \times 10^{-10}$ sec has been achieved for prompt radiations. The mean lives measured are 1.04×10^{-9} sec for B^{10} and 3.73×10^{-9} sec for C^{15} .

Experiments on the $\text{B}^{10}(\text{He}^3, p)\text{C}^{12}$ reaction, undertaken to detect the 3.2-Mev gamma-ray decay of the 7.66-Mev level in C^{12} , were successfully completed. This level is thought to be involved in the helium-burning process in red giant stars. As a further check on this de-excitation mode, experiments have been done on the B^{12} beta decay which has a partial branch to the 7.66-Mev state. After gamma-gamma coincidence measurements proved unsuccessful it was decided to attempt to detect beta-gamma-gamma triple coincidences. Although positive indications for the presence of the 3.2-Mev gamma ray have been obtained, the results are not yet conclusive and the work is being continued.

For some time it has been known empirically that anomalies may occur in the internal conversion of nuclear electric dipole radiation in the L -

shell of an atom. A formalism has been developed that explains the general pattern of these observed anomalies. The computations necessary to make this formalism quantitative have been carried out with the help of the Frankford Arsenal Computing Center. The explicit extraction of the anomalous matrix elements and a comparison with model predictions are in progress.

A similar program for setting up the quantitative formalism required for analysis of anomalous magnetic dipole internal conversion in the L -shell is in progress. The calculations have been carried through for the interesting case of Tl^{203} and are being compared with preliminary experiments done elsewhere. The electron wave functions used in this work have also been applied to other nuclear-electron interaction problems.

Studies of the moment of inertia of a large system whose states are described by the "superconducting" model, and work on nuclear models, paralleling experiments done in this Laboratory, have also been carried out.

The electronic dipole polarizabilities α_d of the alkali atoms have been calculated with use of appropriate wave functions for the valence electrons involved. The theoretical results for α_d are in reasonably good agreement with the observed values from two recent experiments. In connection with the possible existence of a nuclear electric hexadecapole (16-pole) moment, various anti-shielding effects for ions in crystals have been investigated. A new orthogonality property of the radial wave functions for hydrogenlike atoms has been proved.

NEUTRON PHYSICS

Studies of the interactions between neutrons and atomic nuclei continue to yield information important to the better understanding of the properties of matter. In a continuing program of research in neutron physics at the Brookhaven Graphite Research Reactor, investigations have been made of the scattering of very slow or "cold" neutrons by liquids and solids.

Cold neutron scattering measurements of the motions of the ammonium ion in the ammonium halide molecular crystals NH_4Cl , NH_4Br , and NH_4I have been completed.

The torsional oscillations and optical vibration frequencies of the NH_4^+ ion in the different halogen

lattices both below and above the order-disorder transition have been determined. The neutron data indicate that the ammonium ion is performing torsional oscillations both below and above the transition point. The torsional line shifts to a slightly higher frequency below the transition and is sharper than the line observed above the transition.

showed that they could not be fitted to any model having free rotation. In order to substantiate this conclusion, neutron measurements were made as a function of sample temperature for three different temperatures in the face-centered phase. The observed spectra broadened as the temperature was increased, but the peak of the observed spectra

Figure 8. The present arrangement of the cold neutron facility.

In the ammonium halides a high temperature phase transition is known to exist, in which the structure changes from body- to face-centered cubic. Previous infrared and neutron diffraction measurements had led to the conclusion that in this high temperature phase the ammonium ion was undergoing free rotation. Neutron measurements made in this high temperature phase of NH_4I resulted in data markedly different from those observed in the body-centered phase. A careful analysis of the data, based on the Sachs-Teller mass tensor theory for rotational motions,

remained essentially at the same energy. For a freely rotating ion, the peak would be expected to shift with temperature because of the Boltzmann population factor. It is therefore concluded that in the high temperature phase NH_4I is not freely rotating, but most likely is undergoing a very large amplitude, anharmonic torsional oscillation.

The cold neutron facility has been modified to permit the angle of neutron scattering to be changed and thereby give a wide range of momentum transfers. The present arrangement is shown in Figure 8.

At the joint BNL-Chalk River fast chopper project, transmission measurements using the 88-meter flight path were carried out to determine average total neutron cross sections in the 10 to 100-keV energy region. The results of these measurements are being compared with the neutron cross sections predicted by an optical model with a diffuse edge potential. Measurements have been carried out on Ni, Mo, Rh, Ag, Cd, In, and Y. The interest in this region of mass number lies in the observation of the expected splitting of the p -wave giant resonance by a spin-orbit coupling term in the nuclear potential. At energies of 10 to 100 keV only the $\ell=0$ and $\ell=1$ partial waves contribute to the cross section.

Two interesting results have come from this work: First, it is found that in order to get a reasonable optical model fit to the data, the diffuseness parameter of the potential must be ≈ 0.3 f. The data cannot be fitted with a diffuseness parameter of 0.5 f, the value used by Feshbach, Porter, and Weisskopf in fitting the lower energy s -wave data. A re-examination of previously published s -wave strength function data, by Dr. Eric Vogt of Chalk River, has led to the conclusion that the diffuseness parameter is 0.34 f. Second, it is found that the p -wave strength function, when plotted as a function of atomic weight, shows the effect of spin-orbit splitting. The best fit to the data indicates that the strength of the spin-orbit potential is more nearly 16 MeV than the 8 MeV obtained from polarization experiments performed in the MeV energy region.

Studies of the angular distribution of elastically scattered neutrons from oxygen, utilizing the neutrons produced by the $T(p,n)He^3$ reaction from the 3-MeV protons accelerated in the 18-in. cyclotron, have continued. Results have been obtained at incident neutron energies of 2.250, 2.030, 1.750, and 1.510 MeV. The mean energy of the protons incident on the tritium target was monitored by the time-of-flight method, and the spread in energy of these neutrons was determined to be ± 33 , 57, 63, and 63 keV respectively. The scattered neutron intensities were measured at 15° intervals over the range of 20° to 165° in the laboratory system. The appreciable polarization of the $T(p,n)He^3$ neutrons in the measurements at 2.030, 1.750, and 1.510 MeV made it possible to determine the polarization analyzing power of oxygen at these energies. In a separate experiment the

polarization of the neutrons from the $T(p,n)He^3$ reaction is being determined by measurements of the scattering from helium. The data obtained for the scattering from oxygen together with further measurements at lower neutron energy will also be used to determine the $T(p,n)He^3$ polarization. This will provide a comparison of oxygen and helium as polarization analyzers and hence assist in evaluating the accuracy of phase shifts derived from oxygen scattering data.

In a cooperative program of Columbia University, Bell Telephone Laboratories, and the Chemistry Department and Instrumentation Division of Brookhaven National Laboratory, measurements of the correlated energy spectrum of fission fragments from thermal neutron fission of Pu^{239} , U^{235} , and U^{233} have been made. A thin target of the fissile isotope is placed between two p-n junction semiconductor detectors. Coincident amplified pulses from the two detectors are fed into the two axes of a 64×64 two-dimensional pulse height analyzer. In all three isotopes fine structure has been observed in the correlated kinetic energy distributions which is reflected in the fine structure of the calculated fission mass distributions. In addition, structure has been observed in the total mass distribution because the mass yield curves for different total kinetic energy tend to fall into three groups. In the case of U^{235} for total kinetic energies < 162 MeV, the most probable light fragment mass number is 92; for energies between 164 and 171 MeV, ≈ 96 ; and for energies > 173 MeV, ≈ 101 .

Single kinetic energy and total kinetic energy distributions of fission fragments associated with the emission of long-range alpha particles in the thermal neutron fission of U^{235} have been investigated with the use of solid state detectors. The results obtained show that the most probable energies for the light and heavy fragment peaks associated with the emission of long-range alpha particles are 89 ± 1 MeV and 61 ± 1 MeV, respectively. Compared with binary fission the light fragment peak loses more energy than the heavy fragment peak. The mean total energy in "ternary" fission is equal to the mean total energy in binary fission.

In a similar experiment solid state detectors in coincidence and anticoincidence arrangements have been used to determine the binary to ternary fission ratio in U^{235} to a higher precision than has

hitherto been attained. The ratio of binary to ternary fission in U^{235} for pile neutrons with a cadmium ratio of 30 is 490 ± 20 .

In cooperation with the Chemistry Department of Columbia University, studies have been continued on the slow-neutron scattering cross sections of various chemical compounds. During the past year, cross section measurements have been completed for four ammonium halide salts and NH_3 gas over a neutron energy range from 0.0007 to 2 ev. These measurements have also been extended to the low temperature crystal structure phases of NH_4I and NH_4Cl . A definite correlation is observed between the slopes of the scattering cross section vs neutron wavelength curves and the rotational freedom of the ammonium ion in these compounds.

The presence of rotational states at low temperature for exchange of energy with neutrons, together with the comparatively low absorption cross sections, makes ammonium salts such as NH_4PF_6 (N^{15} -enriched) and methyl-substituted compounds such as $CH_3C \equiv CCH_3$ of interest as moderators for cold neutron sources.

Slow-neutron cross section measurements have also been carried out on NH_4PF_6 , $(NH_4)_2S_2O_8$, $(NH_4)_2CrO_4$, NH_4SO_3F , and $CH_3C \equiv CCH_3$.

Experiments utilizing polarized neutrons and polarized nuclei are proceeding on a routine basis. Monochromatic, polarized neutrons obtained by Bragg reflection from a magnetized cobalt-iron crystal are passed through nuclear samples contained in a demagnetization cryostat on the spectrometer arm. Transmission measurements with the neutrons polarized parallel and antiparallel to the nuclei give the spin dependence of the neutron cross section and information on the hyperfine splitting and magnetic structure of the target.

Work on samarium has been completed. Targets used were samarium ethyl sulfate, samarium double nitrate, enriched Sm^{149} double nitrate, and samarium metal. The spins of the four resonances at 0.0976, 0.87, 4.93, and 8.9 ev in Sm^{149} are all $J = I + 1/2$. Analysis of the double nitrate data led to theoretical work on nuclear polarization in powdered samples of anisotropic salts. Failure to observe any polarization in samarium metal at $0.1^\circ K$ indicates that the metal is antiferromagnetic.

An alloy sample containing 8 atomic percent indium in holmium was prepared and studied. The nuclear polarization, f_N , in both components

Table 2

Target nucleus	E_0 , ev	J -value
Dy^{161}	2.72	3
	3.69	2
	4.33	2
	7.73	3
Dy^{163}	1.71	2
Er^{167}	0.460	4
	0.584	3
	6.10	3
	9.6	3

was measured as a function of temperature over the range 4° to $0.07^\circ K$. The holmium results were fitted to an equation of the form $f_N = CB_I(A/2kT)$, with B_I the Brillouin function and C a constant. The derived value of $A/k = +0.58^\circ K$ corresponds to a field of $+8.4 \times 10^6$ oe at the holmium nucleus, in excellent agreement with results of specific heat measurements and theory. Owing to the strong magnetic anisotropy and incomplete magnetic saturation of the holmium, only a fraction of the true polarization in each domain is observed. These effects are accounted for by the constant C . The experimental value $C = 0.48$ is in reasonable agreement with that calculated on the basis of a simple magnetic model for holmium. The indium results show a field of -130 koe at the indium nucleus in this alloy. This "ferromagnetic host" method appears to be very promising as a nuclear polarization technique.

Transmission experiments have also been performed on samples of erbium and dysprosium metal and on erbium ethyl sulfate. Spins were assigned for the resonances at energy E_0 (see Table 2). Preliminary results for the effective fields at the nuclei are: Er , $H_{eff} = +7.8 \times 10^6$ oe; Dy , $H_{eff} = 8.2 \times 10^6$ oe (unsigned, but presumably positive). The spin dependence of the cross section in the region of the 0.460 to 0.584-ev resonance doublet in Er^{167} was investigated in detail. Measurements were also made of the depolarization of the neutron beam by the metal samples. It was found that there was serious depolarization at energies up to 0.5 ev, the highest energy at which the neutron polarization could be measured in a double-crystal arrangement.

A number of nuclei were studied by brute force polarization. Although the nuclear polarizations are small it was possible to get measurable effects

by using thick samples and by working at thermal energies where the counting rates are high. It was found that transmission changes due to nuclear polarization as small as 0.2% could be reliably measured in this way. The results are all preliminary. The protons in a sample of PdH were polarized as a first step toward a polarized deuteron experiment. The success of this experiment has led to the preparation of samples of ZrH_2 and ZrD_2 which will be studied in the near future. The thermal cross section of Li^6 appears to be $90 \pm 10\%$ from the $J=I-\frac{1}{2}$ state, while that of B^{10} is $90 \pm 10\%$ from the $J=I+\frac{1}{2}$ state. In Be^9 the two states make equal contributions, as expected from the fact that the incoherent scattering cross section is zero and thus $b_+ = b_-$. This work was greatly facilitated by the development of a new sample holder that can be moved perpendicular to the neutron beam while at low temperatures. Several samples can be studied at once and transmission measurements can be made. Further, from the transmission of an indium sample included in the assembly, the true temperature of the sample can be derived.

ATOMIC AND MOLECULAR PHYSICS

An examination of the hyperfine structure of the 3.7-hr isomer of Lu^{176} is in progress. The stable

Figure 9. "Isothermals" representing thermal spikes in copper, following a 100-ev knock-on event. Time $= 3.27 \times 10^{-13}$ sec = two Debye periods. Figures give absolute temperatures computed by averaging the kinetic energies of the atoms over two Debye periods.

nucleus Lu^{175} has previously been studied with high precision by others. Both lutetium isotopes have very large electric quadrupole interactions. At operating temperatures both the $D_{3/2}$ and $D_{5/2}$ states of the lutetium atom have appreciable populations. It should be possible, therefore, to compare the interactions of each nucleus with each electronic state with the assurance that each electronic state will be essentially the same for each nucleus. This work may provide new information about the effects of polarization of the electron core by the nuclear electric quadrupole moment. In preliminary measurements the interaction constants have been determined to an accuracy of ≈ 1 part in 10^4 . Further work is in progress involving precision of the order of a few parts in 10^7 . A group from the Lawrence Radiation Laboratory, who have done low precision work on Lu^{176m} , have cooperated in this project.

SOLID STATE PHYSICS

Theory

Most of the theoretical work during the year has been on the production and properties of defects in crystals. Several calculations have reached the stage of giving important results.

Machine calculations of radiation damage events in a model simulating metallic copper have been described in earlier BNL annual reports. This technique has been exploited further during the last year by applying it to thermal spike models of radiation damage. Immediately after the most violent stages of a displacement event individual atoms appear to have agitations sufficiently chaotic to constitute thermal energy. By averaging the kinetic energy of an atom over a time somewhat longer than one Debye period, an effective temperature T can be attributed to that atom. A typical result is illustrated in Figure 9 where isothermal contours are shown for the plane of major action after about two Debye periods. Temperatures often varied erratically from one lattice site to the next, and thus it is an idealization to draw continuous isothermals. Nevertheless the general trend of the results is most easily conveyed in this manner. Very drastic heating, far above the boiling point, is seen in Figure 9. The temperature distribution at this time is far from uniform, and there is strong correlation between very hot spots and collision chains. As the event is

followed, a certain over-all resemblance is found to cooling according to classical laws of heat conduction for a continuum, but striking differences also exist. This is to be expected, since the lattice certainly is not a continuum. An effective thermal diffusivity can be derived in a variety of ways; it is not clear which, if any, is meaningful for macroscopic comparisons. Roughly, a thermal diffusivity of $\approx 0.01 \text{ cm}^2/\text{sec}$ seems operative. This is about an order of magnitude higher than values that have commonly been used in thermal spike calculations, and the major part of the cooling occurs, in these calculations, at least an order of magnitude faster than has usually been supposed. The very ragged distributions of effective temperature, spreading extremely rapidly and according to complex laws, that are revealed in these studies do not inspire confidence in the significance of conventional discussions of thermal spikes.

The above-mentioned computing machine program has also been used to study changes in the vibrational spectrum when an impurity is introduced into the copper lattice. Preliminary results have been obtained for the frequency of a defect mode of vibration for a simple defect, a substitutional impurity differing only in mass from the copper atoms. The frequency of this defect mode has been determined as a function of the mass ratio of the impurity atom to the copper atom. The minimum mass ratio necessary to produce a defect mode of vibration as determined by the machine program agrees with the results derived from analytical expressions. The average displacement from equilibrium of the impurity atom for a given initial velocity has also been obtained as a function of the mass ratio. These latter results are of theoretical interest, and useful for interpretation of recent Mössbauer experiments and cold neutron inelastic scattering experiments.

Point defects tend to anneal out of any crystal when their concentration exceeds that of equilibrium. Single point defects can also cluster or become attached to impurity atoms. An extensive theoretical study of annealing processes is in progress at Brookhaven based on the isolation and combination of simple kinetic steps. Where analytic solutions could not be found, computer solutions have been used to obtain approximations and determine their regions of validity. In quenching experiments only vacancies and their simple clusters are introduced. The simplest idealized model of

the subsequent annealing process involved the simultaneous migration of single and divacancies to sinks and the formation and decomposition of the divacancies. The over-all annealing process is kinetically complex, but under certain circumstances simple quadratic decay and a quadratic plus linear decay are predicted theoretically, with the rates controlled by the migration energy of the single vacancy. In a subsequent calculation, clusters of up to six vacancies were taken into account. The rates of vacancy loss and cluster formation are still controlled by the single vacancy. The final concentration of the various clusters is independent of the annealing temperature but is very sensitive to the initial vacancy concentration and to the sink concentration. For single-vacancy annealing with vacancy-impurity complex formation, with no other interactions permitted, the decay curve is exponential after an initial transient, and the decay constant is related, but not equal, to the rate constant for vacancy migration, since it involves the binding energy of the vacancy to the impurity atom. Preliminary computer solutions of an annealing scheme that allows the formation of clusters of up to six vacancies and the same sized clusters of vacancies attached to impurity atoms show that the rate of this complicated process is controlled by the energy of migration of the single vacancy.

In irradiated metals the basic types of defect are the interstitial and the vacancy. These can recombine, disappear at sinks, cluster, and interact with impurities. Three of these processes have been studied theoretically, the assumption in each case being that the interstitial is the more mobile defect. Random recombination of vacancies and interstitials is bimolecular and therefore obeys quadratic decay. If interstitials are permitted also to disappear at sinks, the simple quadratic behavior is changed to a much more complex decay. However, there is an analytic relation between the concentration of interstitials and vacancies that is valid at all times and independent of the annealing temperature, but a function of the sink concentration and initial defect concentration. Vacancy interstitial annihilation with di-interstitial formation was investigated. No closed solutions and no simple analytic approximations have been found for this kinetic scheme. However, the computer solutions can be used to establish the range of temperature and binding energy in which di-

interstitial formation must be taken into account. A complete analytic solution has been obtained for the kinetics of vacancy-interstitial annihilation with impurity-interstitial trapping. After an initial transient the decay is a simple quadratic that is governed by the decomposition of the interstitial-impurity complex, and its effective activation energy is, therefore, the sum of the interstitial migration energy and its binding energy to the impurity.

Radiation Effects

Radiation damage and other disturbances of perfect periodicity in solids are under investigation with diverse techniques. This is one of the major experimental research activities in solid state physics.

The study of solid state reactions controlled by diffusion and nucleation is continuing. Recent experiments have considerably clarified the role of radiation in influencing the kinetics of the disappearance of carbon from solution in iron. Measurements of the rate of decay of the internal friction peak in quenched iron-carbon after a 12-day low temperature irradiation show that the carbon disappears from solid solution over a thousand times as fast as in unirradiated iron. Complementary measurements of electrical resistivity show a decrease in the same temperature range (40° to 50°C). Analysis of the data from both types of measurements shows that the process of carbon disappearance obeys second-order kinetics with an activation energy of 19.8 kcal/mole, the activation energy for the diffusion of carbon in iron. Calculations indicate that not enough extra precipitation nuclei could have been created by these irradiation doses to cause such an extremely rapid disappearance of the carbon from solution. Further, the thermal precipitation of carbon is governed by nearly first-order rather than second-order kinetics. These two difficulties can be resolved if it is assumed that the carbon is trapped by the vacancies or the interstitials produced by the irradiation. This model is confirmed by irradiations of different total exposure, which all yield second-order kinetics for different initial concentrations of components. If the carbon is trapped by point defects, extra energy must be added to allow the carbon atoms to boil away from the traps and migrate to precipitation sites. Evidence for this is obtained by resistivity measurements during isochronal annealing of irradiated and unirradiated specimens

in that the resistivity decay associated with thermal precipitation of carbon occurs at a higher temperature in the irradiated specimens. Electron-microscope observation of irradiated samples that have been annealed isochronally up to 100°C reveals no precipitate particles, although this time and temperature are sufficient to complete the decay of both the internal friction peak and the electrical resistivity of specimens irradiated at low temperature. This indicates that the carbon is being trapped and that low temperature irradiation retards carbon precipitation.

A study of gamma and x-ray-irradiated Rochelle salt has been completed. The prevailing view has been that the marked changes in ferroelectric properties with irradiation originate in macrostructural effects, that is, in effects involving domain mobility, but unrelated to the crystal structure. Neutron diffraction and optical studies have shown that important structural changes do occur, although domain mobility must also be considered. The behavior of crystals irradiated outside the ferroelectric temperature range differs from that obtained with crystals irradiated in the ferroelectric phase. The results suggest that two basic processes are involved when the ferroelectric phase is irradiated, one related intrinsically to the crystal chemical peculiarities of Rochelle salt, and the other to domain pinning. The competition between these processes leads to locked monoclinic domains, stable even at temperatures outside the normally ferroelectric range, but a decreased spontaneous strain. On the other hand, when crystals are irradiated outside the ferroelectric range, there are no domains to be locked and no domain walls to trap damage products, and the crystal structure effects dominate. After sufficient dosage, such a crystal will remain orthorhombic and nonferroelectric at all temperatures.

A study of the formation and annealing of the color centers in natural quartz is nearing completion. For the first time the absorption spectra, which consist of a superposition of bands of unknown shape, have been successfully resolved into the individual component bands. The bands are non-Gaussian and nonsymmetric. This resolution is essential for a detailed interpretation of the annealing steps. One of these steps, that near 240°C in gamma-irradiated quartz, is particularly interesting, since it correlates with a "peak" in the ultrasonic attenuation obtained when irradiated

quartz is heated (experiment done in cooperation with Brown University). Since it is unlikely that any atomic motion is involved in the production or disappearance (on heating) of the coloration produced by gamma rays alone, these effects most likely involve the motion of electrons and holes. Thus the large ultrasonic attenuation peak observed when the irradiated quartz is heated must be electronic in origin.

Research on irradiated insulating crystals with electron spin resonance techniques (in cooperation with Picatinny Arsenal and the University of Connecticut) is beginning to yield important information on two systems, Al_2O_3 and NaN_3 . Electron spin resonances in Al_2O_3 have been induced by ultraviolet, gamma-ray, reactor, and high energy electron irradiations. At least five different paramagnetic centers have been detected. Ultraviolet and gamma-ray irradiations produce the same resonance absorption line, which saturates with dose. Reactor and 1.8-Mev electron irradiations produce different centers whose intensities do not seem to saturate with dose. It appears that the electron irradiation causes the displacement of only one atom, whereas the reactor irradiation causes more complicated displacements. Gamma-ray irradiation of potassium azide crystals at 77°K creates several paramagnetic defects in addition to the N_4^- molecule-ion previously reported. One of these defects has been identified by its fine and hyperfine structure as a trapped nitrogen atom. Another of the radiation-induced defects has been classified as an N_2^- molecule ion. The trapped nitrogen atoms and the N_2^- molecule ions decrease rapidly upon warming, but the N_4^- is observed to grow in intensity. At room temperature the N_4^- decays with a half-life of ≈ 16 hr.

Photoconductivity measurements represent another important experimental technique for studying insulating crystals. The first extensive measurements with the recently completed photoconductivity apparatus are being done on the radiation-induced changes in the photoconductivity of TiO_2 . Although this material has been studied before, several new aspects have emerged. New photoconductive peaks have been observed at 6200 and 5500 Å which can be populated with ultraviolet light corresponding to the band gap. The photo current (I) at each peak can be expressed as a function of light intensity (L) by the relation $I = \text{constant} \times L^n$. With exposure to

gamma-ray or reactor irradiations the exponent n is altered. The dark current is also changed (by factors of 10 to 1000) by irradiation as well as by oxidation and reduction.

The basic study of radiation effects in easily decomposable materials is continuing. The investigation of the effect of gamma-ray and reactor irradiations on the subsequent thermal decomposition of sodium bromate has been completed. The decomposition is accelerated in a way that suggests that irradiation has enhanced the melting that occurs prior to decomposition. Lead azide has also been studied; the radiation effects on the kinetics of decomposition are extremely marked in this case. The over-all rate increased by a factor of 10, the activation energy decreased from 35 to 25 kcal/mole, and the pre-exponential factor was reduced by about a factor of 10.

Structure of Solids

Another major research activity is the use of neutron and x-ray diffraction in structural and extrastructural studies of crystals. As in the past, these have largely been confined to magnetism and ferroelectricity. The work in these areas has been greatly strengthened by experiments involving the Mössbauer effect.

The emphasis in the magnetic electron density studies has shifted somewhat from metallic materials to inorganic compounds, partly as a result of the lack of suitable single-crystal specimens, but also because of the possibility of more direct interpretations of the spin-density maps. Of particular interest is the change in the spatial distributions of the unpaired electrons in Fe^{3+} , Mn^{2+} , and Ni^{2+} from the free atom state to one in which they are interacting with their surroundings in a solid. For example, the extensive series of measurements carried out on NiO over the past two years has not only demonstrated that the magnetic electrons are significantly contracted toward the nucleus as compared to the calculation for the free atom, but in addition shows that the over-all moment derived from the neutron scattering is $\approx 20\%$ lower than expected on the basis of the g value for the Ni^{2+} atoms. Such differences could be explained by strong covalent coupling between the Ni^{2+} atoms and their surrounding oxygen atoms. These covalent effects were further investigated in single crystals of MnF_2 . The results show the presence of an unpaired electron on the fluorine atoms and

also show the spatial distribution of these ligand magnetic electrons. This electron charge constitutes $<2\%$ of the total unpaired electronic charge associated with each of the Mn^{2+} atoms. The magnetic electron projections now being prepared will permit comparison with previous nuclear magnetic resonance work on MnF_2 and also with current theoretical ideas on the nature of the interactions between the magnetic and nonmagnetic atoms. It is possible to study such spin-density distributions (magnetic moment, ≈ 0.05 Bohr magneton) only because of the increased sensitivity of the polarized beam technique. These experiments also show that (1) the distribution of antiferromagnetic domains is not uniform throughout the bulk of the crystal, (2) it can be related to the local perfection in the crystal, and (3) in a given region of the crystal the domain distribution remains constant right up to the Néel point.

Recently, much interest has been shown in alloys of the iron group elements with platinum, palladium, and rhodium. Polarized neutron techniques have been used to establish the division of the total moment between the constituent elements and to determine the radial extension of the induced unpaired electrons on the atoms of the nontransition elements. In Pd_3Fe , Pt_3Mn , and RhFe a ferromagnetic arrangement is found, whereas in Pt_3Cr there is a ferrimagnetic grouping of the Pt and Cr atoms. The largest induced moment occurs in RhFe with 1.1 Bohr magnetons on the rhodium and 3.1 Bohr magnetons on the iron atoms. This alloy is of special interest since it is the first case among the iron alloys in which the hyperfine field at the iron nucleus, as determined by Mössbauer experiments, is not proportional to the intrinsic moment on the iron atoms. This effect may be related to the fact that the unpaired electrons on the rhodium atoms are significantly more spread out than the $3d$ electrons in the iron group series.

The manner in which a spiral spin configuration changes as the stoichiometry is altered has been investigated by analyzing the neutron patterns of $\text{MnFe}_{2-t}\text{Cr}_t\text{O}_4$. The superlattice peaks in the ferromagnetic spiral structure previously found in MnCr_2O_4 begin to appear at around $t=1.5$. For this composition evidence for short-range ordering is found. Below $t=1.5$ there is no evidence for the spirals. Above $t=1.8$ sharp peaks denoting the presence of long-range spirals are present. From

these measurements it may be concluded that the presence of a significant percentage of iron atoms serves to break up the spirals leading to inhomogeneous magnetic structures at the iron-rich end.

The magnetic structures of several alloys of the AB_2 type have been studied. The alloys have Laves-type structures and heretofore little had been known about their magnetic behavior. TiFe_2 and ZrFe_2 have been found to be ferromagnetic, while ZrCo_2 has an antiparallel spin arrangement. The most interesting of these alloys is TiFe_2 . No moment is found on the titanium atoms, while the iron atoms, located on two nonequivalent sites, have room-temperature moments of 0.2 ± 0.05 and 0.1 ± 0.05 Bohr magneton. The Mössbauer pattern of TiFe_2 taken together with these neutron results demonstrates nicely the linear relation between the magnetic field at the iron nucleus and the intrinsic iron moment.

Earlier studies of the magnetic structures of anhydrous sulfates of the divalent iron-group metals have been extended to include MnSO_4 , CuSO_4 , and the high temperature modification of CoSO_4 . MnSO_4 appears to have a canted spin structure which is weakly ferromagnetic. The CuSO_4 structure can be accurately determined only with single crystals, but an approximate structure that is reasonable and consistent with magnetic data has been derived from powder data. The case of CoSO_4 is especially interesting. This compound can be prepared in two modifications, both becoming antiferromagnetic at about the same temperature. The stable low temperature form, as reported earlier, has a structure with two spin axes that at first seemed to offer a very reasonable explanation for the Russian discovery of a field-induced ferromagnetic transition. It is now clear that the Russians were working on the high temperature modification, which turns out to have a structure with four spin axes.

A cold neutron inelastic scattering experiment on nickel-palladium alloys has demonstrated the existence of a simple defect mode of vibration arising from nickel impurities in a palladium lattice. The frequency of the defect mode of vibration is ≈ 1.1 times the maximum frequency expected for pure palladium lattice. The effect of concentration-broadening on the defect mode was not observable in the experiment because of poor instrumental resolution.

High Energy Accelerators

The responsibility of the Accelerator Department for the two high energy accelerators at Brookhaven, the 3-Bev Cosmotron and the 30-Bev Alternating Gradient Synchrotron, is threefold. At present, by far the greatest emphasis is on the operation, maintenance, improvement, and exploitation of the two accelerators. The second area of responsibility involves the development of the so-called community property equipment for the experimental program (beam separators, magnets, etc.). The accelerator development program, which encompasses the theoretical and practical aspects of advanced machines and techniques in

the high energy accelerator field, constitutes the third area.

The following sections summarize the current status of activities in the Department.

ALTERNATING GRADIENT SYNCHROTRON

Operations

The Alternating Gradient Synchrotron (AGS) has smoothly made the transition from a project under construction to a facility for efficiently conducting research in the high energy field. This is evidenced by the increased effectiveness of the

Figure 1. The AGS complex, showing the new 80-in. Bubble Chamber Building and the North Experimental Area under construction.

AGS as a source of particles and the change in emphasis in the experimental program from survey to quantitative experiments. For example, a separated beam capable of delivering antiprotons, *K*-mesons or pions to the 20-in. hydrogen bubble chamber is now available, and neutrino interactions are being observed in a 10-ton spark chamber.

During the first part of the fiscal year, the machine was operated on the basis of two 8-hr shifts per day, 5 days per week. On January 15, 1962, the work day was extended to three 8-hr shifts. A typical weekly schedule is as follows: 1 shift for routine machine maintenance, 2 shifts for AGS studies, and 12 shifts for the experimental program. The maintenance period, generally the first 8 hr of the week, is utilized for work that must be completed while unrestricted access to the main magnet enclosure is possible. AGS studies are concerned with obtaining data on machine performance and characteristics, the evaluation of newly developed auxiliary equipment, and the training of operators in new machine techniques.

The AGS has achieved peak circulating intensities of $\approx 4.0 \times 10^{11}$ protons/pulse. However, the synchrotron is routinely operated at intensities of 2 to 3×10^{11} protons/pulse. During the year, 3465 hr of operating time were available and were utilized as follows: 55%, high energy physics experiments; 10%, AGS studies; 7.4%, start-up and shutdown procedures; 26%, linac and AGS downtime; 1.3%, experimental downtime; and 0.3%,

experimental area downtime. The linear accelerator (linac) was responsible for much of the synchrotron downtime. However, design changes and system modifications that are being installed as rapidly as practicable should reduce the machine outages. The synchrotron magnet power supply has been pulsed $> 8,126,000$ times with no major system difficulty. The AGS rf system has proved to be extremely reliable after elimination of the usual early component failures. In this area, the current effort is toward long-term reliability. Evapor-Ion pumps (titanium evaporating) continue to be the workhorses of the machine's high vacuum systems. A good operation-maintenance schedule has been achieved. The development of controls for machine operation, monitoring, and safety is a continuing effort throughout the machine complex. The next major phase involves the new East and North Experimental Areas as well as an expanded access control system for the synchrotron. The iron content of the primary cooling water system shows no sign of decreasing; consequently, chemical treatment of this water will be continued. Auxiliary water systems are being modified to be compatible with this water quality. Surveying data have been assembled throughout the year whenever large loading changes were made in and around the machine complex. These data showed that the changes in survey elevations of the machine are well within the expected range. Design and construction of auxiliary experimental

Figure 2. Current arrangement of equipment on the floor of the Target Building for separated beam and neutrino experiments as well as several parasitic experiments.

apparatus continues to account for a substantial portion of the Department's effort. At present, this work includes magnets, power supplies, beam separators, and equipment for circulating beam extraction.

On June 11, 1962, the synchrotron was shut down for an estimated 3-month period, chiefly to permit the main magnet enclosure to be breached for a beam channel to service the North Experimental Area, including the 80-in. bubble chamber facility. This opportunity to service, renew, or improve all components of the synchrotron is being exploited to the fullest.

On July 1, 1961, personnel associated with the AGS numbered 181. By July 1, 1962, the total had reached 214. Further growth is planned in anticipation of machine operation on the basis of 21 shifts per week.

Experimental Programs

During July 1961 the total cross section experiments mentioned in last year's annual report ($\bar{p}$ - p , p - p , π^+ - p , and K^+ - p) were completed. In the months following (except for ≈ 3 months of setup time), two main experimental programs have been in operation: (1) the study of neutrino interactions at the G-10 target location; and (2) the photographing of interactions, in the 20-in. hydrogen bubble chamber, involving separated antiprotons, K -mesons, and pions and the separated-beam transport system originating at the F-10 target location.

The neutrino experiment required the erection of ≈ 4500 tons of steel, 2000 tons of heavy concrete, and 60 tons of lead shielding. Somewhat more than 800 hr of primary running time was provided for this experiment. However, during this time considerable parasitic experimentation was accomplished. The separated-beam system from the F-10 target and the 20-in. hydrogen bubble chamber were used in this way to obtain $\approx 160,000$ π^- pictures at various momenta for groups from the Universities of Pennsylvania and Wisconsin (a joint project), the University of Illinois, and Brookhaven; and to obtain $\approx 200,000$ π^+ pictures for groups from Yale University, Columbia University, the University of California (San Diego), and two Brookhaven groups.

Similarly, secondary beams from the G-10 target were used for several other experiments during the neutrino experiment. About 150,000 pictures were

obtained in the 14-in. hydrogen bubble chamber for the study of neutral beam interactions at $+21^\circ$. At -15° and $+22^\circ$, a study was made by a joint Cornell University-Brookhaven group of p - p elastic scattering (at $\approx 90^\circ$ in the center-of-mass system). This experiment also used ≈ 100 hr of prime experimental time. At -15° , the K - p total and differential cross sections are being measured at momenta around 5 Bev/ c . At -30° , a group from MIT has been studying K^- - p scattering to obtain both the elastic and inelastic distributions at momenta near 1.2 Bev/ c .

The separated beam system can provide antiprotons at momenta of up to ≈ 3.5 to 3.7 Bev/ c , and $\approx 100,000$ pictures were taken in the 20-in. bubble chamber of $\bar{p}$ - p interactions. The beam transport system is ≈ 270 ft long and consists of one pair of 12-in.-diam, 30-in.-long quadrupoles; four pairs of 8-in.-diam, 24-in.-long quadrupoles; two deflection magnets (aperture, 6 in. high $\times$ 18 in. wide $\times$ 72 in. long); and two electromagnetic velocity spectrometers. To avoid multiple scattering, an almost continuous vacuum pipe extends from the synchrotron to the final mass-resolving slit. Precise magnetic measurements of the quadrupoles and bending magnets were made prior to installation, and from these the currents required for optimum tuning were calculated. During the actual tuning procedure it was found that the best focus conditions were obtained with these calculated currents. A pair of 4-in.-diam quadrupoles and a bending magnet shape and bend the beam for entrance into the bubble chamber. The antiprotons transmitted are ≈ 12 per 10^{10} protons of circulating beam in the AGS. The measured separation factor is ≈ 500 .

This beam also provides separated K^+ -mesons, up to ≈ 2.3 to 2.5 Bev/ c . About 100,000 K^- pictures were taken (at maximum momentum) by a Syracuse University-Brookhaven group, and $\approx 50,000$ K^- pictures were taken at ≈ 1.9 Bev/ c by a Yale University-Brookhaven group. A parasitic emulsion exposure in the K^- beam was made by a group from the University of Maryland. About 20,000 K^+ pictures were obtained at 1.9 Bev/ c by a group from the University of California (Berkeley), and 42,000 K^+ pictures were taken at 2.3 Bev/ c by a group from the Johns Hopkins University.

During most of the operating period, ≈ 2 hr per week have been used to irradiate targets for radiochemistry studies by the Chemistry Department.

For future experiments, orbit calculations have been made for use in designing extracted proton beams for both sharp (one-turn) and slower extraction. Work is in progress on the necessary magnets and mechanical and electrical devices. The design for the external-beam transport system and the transport system for separated beams of higher momenta (≈ 8 Bev/ c for antiprotons) have required extensive collaboration with the Bubble Chamber Group in the Physics Department. These systems will be used in conjunction with the 80-in. bubble chamber in the North Experimental Area.

Linear Accelerator

A "Duo-Plasmatron" ion source, installed in the preinjector in September, increased the preinjector output to between 60 and 80 ma. The linac output increased, but not in proportion, probably because of poorer emittance properties and current-loading effects in the low energy end of the accelerating tube and focusing system of the preinjector. One evidence of this was the extremely high x-ray radiation of several hundred mr/hr measured in the linac control room during operation of this source. This required the installation of lead and lead-glass shielding. However, filament life was short (< 1 week), and the output pulse of protons was often heavily modulated with noise and frequencies up to 10 mc/sec. For these reasons the PIG source was reinstalled, and the Duo-Plasmatron was again available for continued development.

In September, during a shutdown period, the linac was opened and the position of the drift tubes measured. Displacements of several ten-thousandths of an inch had occurred in many of them. This may have been due to relief of stresses in their supporting stems. The displacements did not seem to have a large adverse effect upon the linac operation. Nevertheless, ≈ 20 of the units were repositioned until all lay upon a smooth curve which departed ≈ 0.007 in. from a straight line. During this time the controls for the drift-tube vacuum system were improved to give added protection against possible damage caused by loss of vacuum.

The linac was designed with a large cylindrical alumina window, through which radio frequency is fed from a waveguide into an evacuated coaxial transmission line and linac coupling loop. This window had failed on several occasions because of electrical punctures of the alumina and the ac-

cumulation of sputtered copper upon its inside surface. The window was replaced by a thick Teflon disk in the coaxial line, located almost in the plane of the tank wall. The new window has been operating satisfactorily for > 8 months with no evidence of deterioration.

Much difficulty has been experienced with electrical breakdown at the ends of the RG 18 coaxial dielectric filled cables used for rf interconnections of the large FTH power amplifiers. Changing the insulating material in the connectors of these cables from polyethylene to petroleum jelly has eliminated small cracks. A long run of RG 18 coaxial cable has been replaced with rigid air coaxial line. Both changes have reduced the machine downtime. Failure of the grid coupling loops in the FTH amplifiers still constitutes a major difficulty. These loops are fabricated from RG 18 cable and the end assembly is cast in polyethylene. A system of air-insulated loops fed by rigid air insulated coaxial cable has been developed, and the components will be installed as soon as possible.

A new low-level rf and frequency multiplying system has proved to be quite reliable under actual operating conditions.

In July 1961 a buncher was installed between the preinjector and the linac, and a new fast energy control was installed on the preinjector. The buncher increased the capture efficiency of the linac by a factor of ≈ 2 . The buncher gap was initially equipped with grids of 5-mil tungsten wire which were eventually destroyed by the proton beam. These have been replaced by tantalum strips ≈ 0.010 in. thick.

The preinjector, using a PIG ion source, delivers ≈ 25 to 30 ma to the linac. The linac output is ≈ 8 to 10 ma.

Development and Engineering Programs

Power. The main magnet power supply operated without any major difficulty from July to December 1961 on the basis of 2 shifts per day, 5 days per week, with $\approx 895,000$ pulses in 612 running hours, and from January 1962 on a 3-shift, 5-day schedule with $\approx 4,000,000$ pulses in 1905 running hours.

The design and construction of two new 13.8-kv feeders for the 80-in. bubble chamber and the new experimental areas have been completed. An additional feeder capacity of 20 Mv-a is now available to the complex. Six unit substations of 2.5

Mv-a each and the 4.8-Mv-a bubble chamber substation were connected to these feeders. The high-voltage incoming side of each substation consists of a load-break oil-immersed disconnect switch which can be connected in three positions such that the power (1) is supplied to the substation only, (2) bypasses the substation and is supplied to the other feeder loads, or (3) is connected to the substation and the other feeder loads at the same time. This type of installation makes possible the operation of each feeder in a radial fashion with the open end at the most desirable point with respect to load demands; it also provides a method of feeding power from an alternate feeder breaker or permits operation as a loop feeder.

Five of these 2.5-Mv-a unit substations provide an additional capacity of 7.5 Mv-a for the experimental equipment which has utilized $\approx 100\%$ of the power in the Target Building (a calculated load of 6079 kv-a being supplied from 6000 kv-a of available power). Five experimental power feeders installed in the East Experimental Building provide 5 Mv-a of power for this area. The design of experimental power distribution for the conjunction tunnel of the North Experimental Area was completed, and the construction contract has been awarded. The completion of this installation early next fiscal year will provide 7 Mv-a of fixed power and 2.5 Mv-a that can be diverted from the East Experimental Building, or a total of 9.5 Mv-a of experimental power in the conjunction tunnel of the North Experimental Area. Forced-air cooling fans ordered for these transformers will increase the available power to 11.5 Mv-a.

The programmed control circuit and field winding power supplies associated with the motor generator sets for the quadrupole and sextupole correcting magnets have been completed, and operational tests are under way.

Mechanical - General. The second group of experimental magnets for beam transport systems is on hand. Deflecting magnets include seven 36-in.-long and nine 72-in.-long units, each with 6×18 -in. apertures. Quadrupoles include four 24-in.-long and eight 48-in.-long units of 8-in. diam, and four 30-in.-long and eight 60-in.-long units of 12-in. diam. The water circuits in all the magnets in this group have been modified to permit higher fields and lower temperature gradients. The water circuits in $\approx 1/3$ of the first group ordered have also been modified.

Several new magnetic-measurement devices have been built to make measurement of the experimental magnets more accurate and easier. Outstanding among these are three harmonic search coils used to measure precisely the fields of the quadrupoles. With the equipment now available, the various magnets can quickly be inter-compared for uniformity. In addition, absolute field strength measurements can be made for each type of magnet. To construct more precise beams, three-dimensional field measurements are needed, and the equipment to obtain these data is now available.

Work has begun on the design of components for an external beam at the AGS. A model of the first septum magnet, made to the full cross section dimensions, has been built and successfully tested. At a distance of 1 mm off the septum a field of < 2 gauss was measured with a 2000-gauss level in the main field of the magnet. Various components of the hydraulic ram system for positioning the magnet at the appropriate time in the acceleration cycle are being assembled for testing.

Flip-target mechanisms of a new type, suitable for heavy and light targets, have been developed and installed in the AGS. Their many advantages include low power consumption, quick blade ejection, ease of speed control, and high reliability. One of the mechanisms has operated for $> 1,200,000$ cycles without mechanical failure or noticeable signs of wear. All targets are now installed in air locks to allow quick replacement of blades with minimum operational downtime.

Figure 3. Target blade shown in UP position (beam path).

During the year a rapid beam deflector was built and installed in the AGS. It is capable of displacing the proton beam ≈ 1 in. off orbit horizontally and has been used primarily in conjunction with targets to produce a rapid spill for high energy physics experiments. A second deflector has been fabricated and is ready for installation.

Vacuum. During fiscal 1962 the vacuum system of the AGS has been in operation on a 24-hr/day basis except for planned shutdowns involving the whole complex. The average pressure in the linac tank is 1×10^{-6} Torr, and that in the ring is 5×10^{-7} Torr. The Evapor-Ion pumps evaporate titanium for ≈ 15 min of every 6 hr; the rest of the time they are ion-pumping only. However, in the target areas where high gas loads exist from newly prepared targets and from target chamber locks, the titanium feed must be operated for 15 min every 2 or 3 hr to keep the pressure at 1×10^{-6} Torr or below.

Over 80 Evapor-Ion pumps are in use on the machine and beam separators. Experience now indicates that ≈ 2 or 3 pumps per week must be removed for regular maintenance. This maintenance program is as good as or better than would be expected with an equal number of diffusion pump systems together with their auxiliary equipment.

During the year a new method of depositing titanium in the pumps was devised. The development and testing of these units is being continued.

An entirely new injection line vacuum system has been designed and ordered for installation as soon as possible. It has been calculated that pressure in the injection pipe will be 1×10^{-6} Torr instead of the current 2×10^{-5} Torr.

Controls. A personnel safety and security system has been designed for the machine complex, and installation has been started. The activation of the East Experimental Area has required the extension of control, communication, information, and safety systems to this area. Circuit designs and equipment arrangements have been completed. The experimental areas will be supervised from a control room in the new facility. The operator will have available all information relative to machine status, details of experimental apparatus, and power usage.

Operating experience with the high-level rf system in the magnet enclosure indicated the need for modification of the turn-on sequence and certain main control room displays of system status.

Accordingly, control circuit changes were made and satisfactory operation has been achieved.

A control system for the linac drift-tube vacuum system was designed and installed. Because of the technique used in assembling the drift tubes, a significant pressure differential between the interior of these units and the linac tank vacuum would destroy some of the vacuum joints. This system provides the necessary safety features together with operational flexibility.

The timing system in the main control room has been revised to insure orderly operational procedures. Sequencing, testing, patching, and trouble-shooting of the timing system elements (pre-determined counters, multiplexing units, etc.) have been facilitated by means of a uniquely devised cable interconnecting rack.

Beam Separators. In order to produce separated beams for bubble chambers and other experiments, seven parallel-plate velocity spectrometers or beam separators, each ≈ 5 m or 200 in. long, are being constructed. Magnetic field coils have been installed on the seven tanks, and the magnetic field uniformity has been measured. In addition, several of the tanks have been conditioned electrically under high voltage, and some difficulties in holding the design voltages have been encountered. Development work is continuing in this area.

Neither of the 600-kv power supply systems has yet passed acceptance tests; they are being improved by the manufacturer to meet the original BNL specifications.

The first of the new separators is expected to be installed at the Cosmotron for use in several experiments scheduled to start during the summer. As additional units become available during the AGS shutdown, another will be installed in the shielding wall inside the ring for low-momentum K -meson studies. A beam that will ultimately require six of the new separators has been planned for separating antiprotons for use in the new 80-in. bubble chamber. Some elements of this array will be installed in the conjunction section to be built this summer and the rest in the new North Experimental Area.

The two original "Cosmotron-type" beam separators (glass negative electrodes) have been in constant use at the AGS. A number of changes in equipment are planned that should further improve their reliability.

Figure 4. Beam separator of the AGS type, showing electrodes extended from tank.

Electronics. The principal concern of the Electronics Group continues to be the improvement and increased reliability of the AGS electronic systems. Because of excessive radiation damage to power transistors in the rf accelerating stations, studies are being made in cooperation with the Health Physics Division to determine dose rates and resultant transistor degradation. With the set-up for the neutrino experiment, very high radiation levels are observed in the vicinity of the first rf station downstream from the Target Building. Consequently, this area was selected as the test site, and a shield of high density concrete was erected nearby. A number of the transistors were exposed to the radiation inside and outside the shield. Preliminary results indicate that the transistor lifetime to be expected inside the shield, under the most adverse conditions, is ≈ 1 year; the current life outside the shield is 2 weeks.

Since bubble chambers are sensitive for an extremely short period of time, rapid beam deflectors are used to move the beam onto a target at a precise moment. The first such deflector has been in operation with the 20-in. and 14-in. bubble chambers for ≈ 1 year. The electronic components of a second deflector have been constructed and tested, and it is scheduled for installation during the present shutdown.

To facilitate vertical positioning of the beam, a "tickler coil" has been modified for vertical steer-

ing. This coil originally was installed to excite betatron oscillations for machine study purposes. It has now been powered to provide vertical motions of ≈ 0.2 in. above and below centerline to compensate for possible orbit perturbations.

In order to bring out an external proton beam, extremely high peak power pulses must be introduced into an ejection system. The present design utilizes a hydrogen thyratron and a pulse-forming network to develop 50 Mw of pulsed power for several microseconds. This power is fed into a ferrite line surrounding the beam and is expected to provide enough motion to move the beam into an ejection magnet. A full-scale experimental setup of the electronic system is being tested. Several types of terminating resistors capable of handling these peak powers have been designed and are being constructed for evaluation.

Survey. A complete level circuit of the machine was run in the fall of 1961. When the machine was not operating, supplemental circuits were run in the vicinity of load changes. The leveling shows that the machine components are stable within ± 0.010 in. when loads are not changed (Monuments A to F).

When the 2600 tons of shielding for the neutrino experiment was installed, the pile caps in the vicinity settled ≈ 0.132 in. They settled an additional 0.030 in. when another 500 tons was installed in the same area. To compensate for this

motion the magnets from *F*-18 to *G*-20 were re-leveled to the standard position. The releveling also corrected the maximum radial displacement of +0.013 in. measured after the total neutrino shield loading. The excavations for the Target Building extensions and the 80-in. bubble chamber had caused a rebound or rise of a maximum of 0.053 in. in the area of superperiods *H* to *K* by February 1962, but the magnets were not re-adjusted for this. Extensive surveying and releveling will be done before, during, and after connection of the magnet enclosure to the North Experimental Area to obtain data on the effect of changing loads on the pile cap foundations and to restore the machine to standard orbit on completion of construction.

The techniques for setting experimental apparatus required some development, and it was necessary to obtain plans and specifications including tolerances and setting marks from the experimenters in advance of the work. For efficient operation, separated beams with slits of 0.50-in. height and collimators of 0.25-in. aperture require optical traverses, screw adjustable supports, and definite reference marks as well as accurately known target positions. Apparatus and shielding already placed often interfered with sight lines. The 100-in. grid on the Target Building floor has been very useful in overcoming these handicaps. The requirements for precise setting are now well understood by all concerned. This is important, for much of the setting must be done when the machine is shut down and time is limited.

The 100-in. grid marked by stainless steel nails, with brass bolts at 300-in. intervals, was extended to cover the floor of the East Target Building. The elevation of nails at 50-ft intervals was also determined to facilitate setting apparatus in this building.

A small first-order triangulation and traverse was run from Monument *H* on the ring through the Target Building parking lot and over the hill to locate horizontal position marks for the 80-in. Bubble Chamber Building. Levels were also run from bench marks near the linac to marks in this building.

Mechanical Services. The iron content of the AGS primary water system continues to be excessive, in the range of 2 ppm. The techniques for chemical treatment of the wells have been revised and augmented to insure the close control over the

hexametaphosphate feed rate required to achieve optimum retardation of corrosion and lessen the purge action due to excessive phosphate. Mechanical cleaning equipment also has been procured and will be used to maintain maximum flow rates through the main water lines from the wells to the complex.

The aluminum heat exchanger in the main magnet cooling system developed large corrosion pits and its failure seemed imminent. The unit has been removed from service and because of the short time available before start-up will be completely reworked with copper-lined aluminum tubing.

Long-range plans include the replacement of the entire heat exchanger with a type 304 stainless steel unit, adaptation of the existing experimental cooling water system to cooling tower operation with make-up water supplied through iron removal units, and provision for a closed system for cooling the ring vacuum pumps with a separate heat exchanger using well water as the primary coolant.

Advanced Accelerator Development

The Accelerator Development Group has been formed to undertake the development of experimental equipment for use with the AGS and to carry out studies concerned with the improvement of the AGS and the design of future accelerators.

During the summer of 1961 this group, in collaboration with the Physics Department and with considerable outside assistance, prepared two reports, *Design Study for a 300 - 1000 Bev Accelerator*, and *Experimental Program Requirements for a 300 - 1000 Bev Accelerator*. These reports were compiled at the request of the United States Atomic Energy Commission in connection with a possible cooperative accelerator project between the United States and the Soviet Union. Although the project is now dormant, the work done during these studies is providing a valuable basis for American studies directed toward a large national accelerator project.

During the summer of 1962 the group will sponsor two conferences at Brookhaven. The first, to be held in August, will be concerned with the design of linear accelerators for the billion-volt range. These accelerators are of interest both for meson and neutrino experimentation and as injectors for super-high-energy synchrotrons. Yale

University is joining in the organization of this conference, which will be attended by experts from 12 linear accelerator centers in the United States and Western Europe. The second conference, to be held early in September, will present a comparison of techniques and plans for the AGS and the CERN Proton Synchrotron. It will be attended by a group from CERN and by observers from major accelerator installations in the United States.

In addition to these activities the group has undertaken four projects. The first is the development of a radio-frequency particle separator. This device gives promise of particle separation at momenta considerably higher than those attainable by the electrostatic separators now in use. Theoretical and model studies have been started as a check on a cavity design originally proposed at CERN.

The objective of the second study is the improvement of conventional electrostatic separators through the development of methods for holding higher voltages across the separator plates. It involves both an investigation of the mechanism of electrostatic breakdown and the invention of techniques to avoid this breakdown.

The third study is theoretical and is concerned with the redesign of the low energy end of the linac injector at the AGS. Preliminary results indicate that a continuous bunching section will give much more effective prebunching of the linac beam than can be achieved with the present single-gap buncher. Successful development of this method

Figure 6. Cosmotron utilization during fiscal 1962.

should increase the linac beam intensity by a factor of 2.

The fourth project initiated by the Accelerator Development Group is a study of the effects on the synchrotron orbit of random displacements of the individual synchrotron magnets. The tolerances now set on the magnet alignment in the AGS were derived from a theoretical analysis by Courant and Snyder. The present study will provide an experimental check on this theory and will serve as a guide in future realignment of the AGS magnet and in the determination of aperture dimensions for new accelerators of the AGS type.

Figure 5. Use of the Cosmotron for experiments in high energy physics during fiscal 1962.

COSMOTRON

The trend continues toward more complicated experiments designed to investigate in detail phenomena that occur only rarely. Considerable work has been done in the investigation of the various pion resonances recently discovered. Other investigations include a measurement of the polarization of the proton arising from Λ decay, measurements of neutral pion production arising from collisions of negative pions with protons, and measurements of the decay parameters of the $K_{\mu 3}$ -meson. Further work was done on the isobaric states of the proton and on the proton-neutron charge-exchange cross section at small angles. During the year >60% of the Cosmotron's experimental time was used by outside research groups. An analysis of potential running time is given in Figures 5 and 6.

The scheduled 4-month shutdown was used primarily for installation of new straight sections on the machine, and the resultant improvement in magnetic field configuration has led to an increase by a factor of >2 in beam intensity and considerably more reliable and stable operation. A number of beam monitoring instruments added during the shutdown allow more rapid and precise internal and external beam positioning. Considerable work was also done on the Van de Graaff injector, which contributed substantially to the increased beam intensity and stability. Sixteen new experimental dc power supplies were installed, and all experimental power supplies were equipped with control circuits capable of regulating current or field to 0.02%. Because of the increased scope of the experimental program, a new experimental control room was constructed and put in operation to serve as the control point and nerve center for all activities on the experimental floor.

The program to produce monokinetic external proton beams lasting for several hundred milliseconds is progressing satisfactorily, and beams of

long duration should be in operation within a few months. Several experimental magnets were procured, and a number of magnet positioning tables were constructed. Accurate measurements of external beam ion-optics have been made over a wide range of energies as part of a continuing program to improve the efficiency and usefulness of the external proton beams. Design of a new office-laboratory wing, including improved shop facilities, has been completed, and construction will begin shortly. A pool of equipment for use in high energy electronic experiments has been established. Equipment available to experimental groups includes coincidence circuits, discriminators, power supplies, scalars, etc.

The hydrogen liquefier produced 126,783 liters of liquid hydrogen for experimental use in fiscal 1962; in addition, 35,000 liters were purchased from an outside supplier during periods of peak demand. The target group designed and constructed a number of liquid hydrogen targets, and the installation of an electrolysis plant to produce deuterium gas has begun.

Instrumentation

The staff of the Instrumentation Division changed very little during the past fiscal year, having increased from 80 to 83. As of June 30th it comprised 20 engineers and physicists, 1 mathematician, 49 electronic and mechanical technicians, 5 glassblowers, 3 draftsmen, and 5 persons engaged in secretarial and administrative activities. A number of the more experienced technicians have assumed supervisory responsibilities and participate to a substantial extent in engineering design and development. No changes were made in the physical plant during the year.

The Division operates a glass shop that fabricates glassware for the whole Laboratory. A wide variety of services are performed in a vacuum shop; for example, 300 items were coated with thin films by vacuum deposition. The calibration and repair of a great variety of commercial instruments requires the full-time services of nine technicians. Two displays were built for the Exhibit Hall. One is a spark chamber that shows cosmic-ray tracks. The other is an animated panel to illustrate the technique of neutron activation analysis used in the Medical Department. A number of more conventional design and construction jobs were carried out. Many of the major projects are summarized below.

INSTRUMENTATION FOR HIGH ENERGY PHYSICS

Increased activity at the Alternating Gradient Synchrotron called for much new instrumentation. Research on spark chambers was continued, considerable effort was expended on instrumentation for handling bubble chamber photographs, and fast circuit boards were produced in volume for counter and spark chamber experiments.

The study of pulsed spark chambers for the spatial determination of particle tracks was continued, with emphasis on improving spark chamber characteristics to meet the requirements of experiments in high energy nuclear physics.

With the aid of a counter-controlled spark chamber, a method for labeling tracks in bubble chambers in terms of the masses of the particles

producing them has been devised and experimentally tested. This method is of advantage primarily in the region above 10 Bev/ c , for which mass separators are not yet available. A particle-selecting counter system which includes a differential Čerenkov counter triggers a multisection spark chamber upon traversal by a particle of a given mass and indicates its position and direction in the beam. Separate pairs of spark chamber sections may be sequentially activated in times as short as the spark chamber resolving time, which makes possible the labeling of several particles per beam burst. By correlating the positions in the spark chamber with those in the bubble chamber, the tracks of the selected particles are indicated, which permits study of the interaction of these particles in the bubble chamber or their exclusion as undesirable contaminants of the beam.

Of importance in the utilization of spark chambers for rare particles in intense beams is reduction of the spark chamber resolving time, which is usually of the order of $\frac{1}{2}$ μ sec. One of the limiting factors in this reduction is the system delay between the passage of an ionizing particle through the particle-selecting counters and the production of a spark in the chamber. The delay times in photomultipliers, cables, coincidence mixers, high voltage pulse generators, spark gaps, spark chamber voltage rise, and spark formation were investigated, and a system was developed which results in a total delay, including that in the photomultiplier, of only about 33 nsec. This allows the spark chamber resolving time to be reduced by an order of magnitude. The shorter delay reduces the random displacement of sparks about the particle trajectory caused by diffusion of track electrons during the waiting time and decreases the transverse displacement of tracks in spark chambers operating in a magnetic field.

Several members of the Division have worked with the Bubble Chamber Group on apparatus for semiautomatic handling of the film on which tracks are recorded. This work included the design of a system for identifying successive frames on the film and for reading the information unam-

biguously at manual or automatic scanning stations, development of a film transport mechanism for use with a flying-spot film scanner, and construction of the electronic circuits associated with an optical scanner which generates digitized coordinate information and transmits it directly to an IBM 7090 computer.

The bubble chamber photographs are first surveyed by human scanners who select tracks of interest and record coordinates at a few locations on each track. The manually recorded data are used to guide the computer and flying-spot scanner in automatically measuring the selected tracks and calculating their properties. Following a suggestion of Professor H. Teager of MIT, the Instrumentation Division has developed a novel digitizing table for the manual-visual scanning operation. Photographs are projected onto a surface beneath which there is an orthogonal array of etched wires supplied with a suitably programmed pattern of current pulses. A light-weight magnetic pickup probe, manually positioned on the projection surface, produces a set of induced signals which can be interpreted to indicate the Cartesian position coordinates of the probe.

Circuits were also developed for use with existing manual film-scanning apparatus. Two different systems provide a measure of the bubble density along the track, which is related to particle velocity, and a servo system was designed to automatically align a slit with the center of a track after an operator has set the slit in the vicinity of the track. In this way a precise reading is obtained more easily and faster than is possible by hand.

Most of the counter experiments at the high energy accelerators record on scalars. Each scalar has seven decades. A system has been constructed to read 16 such scalars one after the other in the period between accelerator pulses. A printed record is produced with a parallel printer and the digits are punched sequentially on paper tape.

SEMICONDUCTOR RADIATION DETECTORS

Work has continued on the development of silicon p-n junction radiation detectors. In collaboration with the Bell Telephone Laboratories, radiation detector packages have been designed to investigate the proton and electron energy spectra in the Van Allen radiation belts. These packages are to fly in Bell Laboratories' Telstar communi-

cation satellite and also in the similar RCA vehicle. Both packages have completed all tests, and launchings are scheduled for the summer of 1962.

Investigations have been made of the physics of the lithium drift process for the production of thick junctions capable of stopping long-range particles. A method has been developed for drifting at constant power that makes use of a specially designed power supply and liquid fluorocarbon heat sink. This system enables drifting to proceed at significantly increased rates. Junctions ≈ 2.5 mm in thickness and having areas of 1 cm^2 have been produced, although this thickness is nowhere near the limit obtainable. When cooled to dry-ice acetone temperatures these detectors exhibit line widths of ≈ 10 kev. A special contacting procedure has enabled the insensitive layer on these devices to be reduced to $\approx 50 \mu\text{g}/\text{cm}^2$.

Techniques have been developed for the production of thin silicon wafers suitable for $\Delta E/\Delta X$ particle identification systems. The resulting detectors have thicknesses of $\approx 25 \mu$ and areas up to $\approx 20 \text{ mm}^2$. While the thickness uniformity leaves something to be desired, nevertheless these devices have proved adequate for particle identification in some low energy nuclear reaction studies.

Work is proceeding on improved fabrication techniques, operation of devices at low temperatures, and associated electronic problems.

AMPLITUDE AND TIME-OF-FLIGHT SPECTROSCOPY

Semiconductor radiation detectors have very high intrinsic energy resolution. With low noise amplifiers, charged particles differing in energy by as little as 1 part in 1000 may be resolved. This is much superior to the resolution obtainable with scintillation detectors. To exploit such good resolution, pulse-height analyzers with more channels were needed. Furthermore, even slight drifts in amplifier gain or analyzer sensitivity could not be tolerated. For this application a 4096-channel analyzer and automatic gain stabilization system have been developed. The analyzer converts the pulse height to a time measurement, as do most existing multichannel analyzers, but at a ten times faster rate. Fast scalars and buffers are used. Stabilization is achieved by periodically applying a pair of precisely regulated reference pulses to the preamplifier input. The pulses are measured by

the analyzer, and the digital values of the results are compared with preset intended values. Errors in the measurement of the smaller pulse are integrated digitally and are used to correct the analyzer zero bias, while errors in the measurement of the larger pulse are used to correct the energy calibration.

A similar technique has been employed in a time-of-flight analyzer. Neutrons are produced in short bursts at the BNL 18-in. cyclotron. Their energy is determined by the length of time it takes them to travel from the source to a detector. The time scale is $<0.1 \mu\text{sec}$. This short time is converted to a longer one by measuring the phase of the beat frequency between an oscillator pulsed by the detector and the cyclotron oscillator. The phase of the beat note is measured digitally and tallied in the memory of a commercial pulse-height analyzer. If either oscillator drifts, readings will be in error. It is worth while therefore to make the detector-pulsed oscillator follow the cyclotron oscillation with a fixed frequency offset. It is possible to measure one full cycle of the beat note following analysis of each neutron. Again the measurement is digital, and a scaler precesses if there is any error. The control scaler automatically adjusts the pulsed oscillator to the correct frequency.

In 1955 deWaard published a circuit for stabilizing a scintillation spectrometer system. A reference source is placed near the crystal, and a single-channel analyzer is set to receive pulses from the photopeak. The channel bias is modulated by a small signal so that it samples the counting rate just above and just below the center of the expected pulse amplitude distribution. The high voltage of the photomultiplier (which controls the gain) is adjusted to keep these two rates equal. This system has been redesigned by using transistors and an electrometer tube. In addition to the modulated single-channel analyzer, the chassis contains a bidirectional count integrator and a built-in, servo-controlled high voltage supply.

NEUTRON SPECTROSCOPY

A complex control system has been constructed to control neutron crystal spectrometers and to record the data. A program is prepared on punched paper tape. The electronic system reads instructions, one at a time, and after each one

has been executed goes back to the tape for the next instruction. In the simplest mode of operation one program consists of setting the sample to a particular angle, setting the detector at another specified angle, counting neutrons for a preset time, printing out the scaler reading, advancing the detector by a given increment, counting and recording, etc., until the detector reaches another angle specified as the end of that run. Then the sample is repositioned and another run is started. The system can adjust a second crystal (if there is one), operate with reference to a monitor detector rather than for a fixed time, transfer data from the input tape to the output, and perform a variety of other operations. With this system spectrometers can operate continuously and provide both a printed record and punched tape for computer analysis.

AUTOMATIC CONTROLS

A largely mechanical system was built to perform automatically a predetermined series of functions for accumulating and recording data in the study at low temperature of fluorescence and phosphorescence. A high-resolution beta-ray spectrometer has been equipped for automatic spectrum scanning. A more stable emission regulator was built to regulate the ion source in a mass spectrometer. An automatic recording system was installed to record temperatures at various locations among the trees irradiated by a high-intensity gamma source.

BRAIN TUMOR SCANNER

An instrument was built that makes use of positron-emitting isotopes for location of brain tumors based on the method developed by Sweet, Brownell, and Aranow at Massachusetts General Hospital, Boston, Mass. This system inherently provides more information about the distribution of radioactivity in the head in less time than existing scanners which use one or two detectors. A stationary circular array of 32 scintillation detectors scans a horizontal layer of the head from many directions simultaneously. The data, consisting of the number of counts in all possible pairs of detectors, are coded and stored in the memory of a two-dimensional pulse-height analyzer. The system includes a unique method of displaying and

interpreting the data which enables rapid approximate analysis of complex source distribution patterns.

A study program is being conducted to determine whether a mathematical solution of the problem of converting the coincidence data to a map of the radioactivity distribution with reasonable errors is feasible.

METEOROLOGY

Air-Borne Particulate Studies

Five very successful operations were conducted during fiscal 1962 in which Cu^{64} spheres and uranine dye were utilized simultaneously as tracers in particulate deposition studies. The techniques and procedures required for this work have been under development for a considerable period, and only within the last year have acceptable field data become available. The total number of runs may seem small, but it should be recognized that the 13-hr half-life of the radioactive copper imposes a minimum delay of ≈ 2 weeks between tests, and that suitable meteorological conditions occur on a somewhat indefinite schedule. Also, reduction of the basic sampling data, which include uranine and copper concentrations at 300 points, deposition measurements of two different types at ≈ 150 points, and associated meteorological records, requires 30 days of effort by several members of the Meteorology Group.

Most important is the fact that all the techniques have operated satisfactorily and require no

important changes. It has been found, for example, that the Cu^{64} particles retain their individual identity without serious agglomeration, and that the mobile carts detect true deposition on the ground and vegetation. The success of this technique suggests that the copper spheres may have application as tracers in a number of other projects, since they have relatively few of the disadvantages usually associated with radioactive isotopes. Air-borne concentrations well below maximum allowable values are readily detectable, and the purity of the isotope insures substantially no residual long-lived activity.

The data from these runs are now being compared with the theoretical treatment of Chamberlain and with the recent work of Davidson relating to polydisperse particles. It is already possible to state that the sticky paper used routinely by the Atomic Energy Commission for particle sampling throughout the United States appears to be very satisfactory for the larger (5 to 10- μ) particles. Agreement between the direct measurements of field deposition and the collection on sticky paper squares is generally within a factor of two for this size range.

A second phase of the particulate studies involves investigation of the dispersion of pollen grains in cooperation with the New York State Museum and Science Service. In fiscal 1962 natural particulates other than ragweed pollen were included in this investigation for the first time. Limited tests with 100- μ corn particles have been conducted in the sampling field west of the meteorological facility, and during the summer of 1962 the investigation will be extended to the 32- μ timothy pollen grains.

Results from the past year's work have shown, as anticipated, quite rapid deposition of the ragweed pollen, and have substantiated earlier indications that the aeroallergen problem is primarily concerned with very large source regions rather than local vegetation. Figure 1 shows the rapid reduction in pollen concentration with distance from the special source area as related to the natural background of the pollen derived from other sources and observed upwind. For example, for an initial concentration at the edge of the field reaching some 500 times the background pollen count, a reduction sufficient to bring the concentration to levels equivalent to background occurs at about 400 feet. This implies that, to experience

Figure 1. Decrease of ragweed pollen concentration directly downwind of the source. The graph shows the average distance required for pollen concentration to decrease from some multiple of background (X_0/BK) at the source to specified percentages of background.

reactions from local ragweed, it would almost be necessary for a person to live within a dense growth of it.

A new sampling field in the eastern part of the site has been equipped for measurements of airborne pollen concentration at three heights above the ground, and current work includes the development of mathematical approximations of complex ragweed area sources.

Instrumentation

Three significant instrumental developments have been pursued by the Meteorology Group within the past year, two of them associated with meteorological problems and a third with particulate sampling. Of potential general value is the development of a new anemometer cup and ring design which has an unusual combination of weight, sensitivity, and durability. This design, completed in the spring of 1961, has undergone

Figure 2. Prototype of rotoslide particulate sampler. Glass microscope slides are rotated by a drive motor at 1700 rpm. The cylindrical plastic shield is automatically raised to prevent spurious collection when the device is inoperative. The sampler can be used with an appropriate time cycle to obtain representative 24-hr samples without overloading.

continuous testing and modification throughout the past year, with various combinations of cup and ring materials. Recent testing has shown that the device has a starting speed of 0.3 mph and can withstand maximum winds as high as 150 mph. In fact, one of the units reached 175 mph before failing.

Another development for wind measurement is a digital wind vane. Wind instrumentation should, wherever possible, provide a digital output directly suitable for automatic recording and processing, and a bench model of a wind vane has been completed using a system in which the output in electrical pulses per unit time is directly proportional to the wind direction. This instrument has not been tested in the field, since considerable work is still necessary on the components, although the principle has been shown to be completely feasible.

For a number of years, members of the Group have been experimenting with a variety of sampling devices suitable for the larger air-borne particulates (10 to 100- μ). This range of particle size presents some very real sampling difficulties in the natural atmosphere. While the particles are easily collected by filter media in the sense that they do not pass through filters readily, it is exceedingly difficult to design a filter holder that does not become inefficient in strong turbulent flow. On the other hand, impaction-type samplers are subject to wind effects in the opposite sense, since these devices become almost completely inefficient at very low wind speeds.

A happy solution to this problem was developed by Perkins and Leighton of Stanford University. They designed a rotating rectangular metal bar assembly known as the "rotorod," in which the rapid movement of the bar surface through the atmosphere impacts the large particles efficiently and essentially independently of wind speed. This sampler in its original form has two important defects in terms of sampling aeroallergens. Counting particles on the rods under the microscope is difficult, and a continuously rotating rod sampler is completely impractical for long-period utilization because an excessive number of particles reduces further collection substantially. Brookhaven's "rotoslide" sampler utilizes the original principle of the Stanford development, but includes modifications that eliminate these difficulties. Figure 2 shows one of the prototypes, in which the two slides mounted on edge in metal holders are clearly visible. The plastic cover and cylinder pre-

vent collection of material when the sampler is not in operation. The cylinder is automatically positioned at this time so that the slide edges are completely protected from the natural wind flow and cease to be impaction collectors when inoperative. This device has been sufficiently successful to be submitted to the National Academy of Allergy as a possible substitute for the Durham slide collector as a standard pollen sampler.

Forest Studies

Publications within the last two years have reflected the Group's interest in the shorter-term fluctuations of air-borne concentrations from a continuous point source, and in the 1961 annual report mention was made of the initiation of a program to extend these studies to dense vegetation. Within the past fiscal year, an 84-ft tower has been installed immediately south of the main meteorological facility in an area of very uniform forest vegetation. Installation was carefully managed so that there was almost no disruption of the vegetation, and the tower is actually within the forest rather than in a cleared area surrounded by forest. Completion of the tower instrumentation was deferred until very recently, since there was little point in beginning the investigation while the

vegetation was changing rather rapidly. Consequently, the installation of 6 matched anemometers and the completion of 6 new photometric densitometers for the measurement of oil-fog smoke has just been finished. With these instruments, detailed studies of the wind profile within the forest canopy and directly above it will be conducted, and records of oil-fog concentration fluctuations from a ground-level point source will be made, both in the open field to the south of the forest and within it. From these measurements, the effect of vegetation on short-term concentrations can be delineated.

Data Processing

Automation of the Meteorology Group records has been continued throughout the year with the intent not only of making the Group's efforts more productive, but of providing data that can be readily transferred to other organizations for their use. At present, climatological data consisting of wind direction and speed at several heights, soil and air temperatures from 10 ft below ground to 410 ft above ground, and other variables such as wind gustiness, precipitation amount, and net radiation are all recorded on magnetic tape for the period from February 1960 to the present.

Chemistry

The following sections describe research in the Chemistry Department on various problems involving radiations, isotopes, nuclear properties and reactions, and related chemical and physical topics. The work reported has been done by members of the continuing staff, postdoctoral Research Associates, and visitors from other institutions.

MASS SPECTROSCOPIC STUDIES ON ION-MOLECULE REACTIONS

The Eyring-Hirschfelder-Taylor theoretical calculation of the rate constant for the reaction of H_2^+ with H_2 giving H_3^+ , modified for the case $D_2^+ + D_2 \rightarrow D_3^+$, agrees remarkably well with the subsequent experimental measurements on the latter reaction by Stevenson and Schissler. In contrast, in the case of the reaction between H_2^+ and He to give HeH^+ , Hertzberg found a rate constant that was only about 15% of the theoretical value. The H_2^+ -He reaction has been reinvestigated and Hertzberg's results have been qualitatively confirmed. The rate constants did not show a scatter found by Hertzberg, but varied systematically and reproducibly with ion velocity. It is to be noted that the H_2^+ -He reaction, unlike most gas-phase ion-molecule reactions, is endothermic by 0.95 eV.

The experimental results support the hypothesis of Polanyi that exothermic reactions produce vibrationally excited products and endothermic reactions proceed via vibrationally excited reactants. This accounts for the facts that HeH^+ is not observed as a product of the highly exothermic reaction of He^+ with H_2 and that only 20% of the H_2^+ can react with He giving HeH^+ . It can be shown that $\approx 20\%$ of the total H_2^+ formed by electron impact on H_2 has sufficient vibrational energy for the reaction.

The variation of the HeH^+ rate constant with ion velocity above a critical value is similar to that observed for the H_3^+ reaction. If a repulsive interaction between He and H_2^+ similar to that between H_2 and H_2^+ is assumed, then the results in this range of higher ion velocities are quantitatively

explained. For a narrow region of low ion velocity the reaction rate increases with increasing ion translational energy.

The HeH^+ reaction may be used as a probe to estimate the relative concentrations of H_2^+ above and below the fifth vibrational state. This has been done for H_2^+ produced by electron impact on methane. The experimental rate constant for the reaction between He and H_2^+ from methane is about four times that with H_2^+ from H_2 .

RATE CONSTANTS FOR DECOMPOSITIONS OF EXCITED MOLECULE IONS

For a number of years the dissociation patterns observed in a mass spectrometer have been explained in terms of competitive unimolecular reaction paths of excited molecule ions. For the formulation of the rate constants the statistical theory of mass spectra of Eyring and co-workers has usually been employed. This theory assumes that the excitation energy is equilibrated over all degrees of freedom. Its application requires a knowledge of the number of states accessible to both the excited molecule ion and the corresponding transition state. Eyring and co-workers proposed the use of certain semiclassical formulas for this purpose. Many difficulties have been encountered in the application of this theory, a serious one being the prediction of too low a rate constant near the threshold excitation energy. It has also been noted by several researchers that the semiclassical formulas are inadequate.

This problem has been further studied. Rate constants in the threshold region for unimolecular processes have been recalculated with procedures involving either the counting of the actual number of vibrational states accessible or the use of corresponding formulas that have been proved accurate. One of the main faults of the semiclassical theory is found to be that it ignores the ground vibrational state of the transition state. The new formulation removes many of the difficulties of the older theoretical treatment, and makes some predictions which can be tested by experiment.

KINETIC STUDIES IN SOME INORGANIC SYSTEMS

The System Mn^{2+} - Mn^{3+}

In electron transfer reactions between ions of the type $\text{A}(\text{H}_2\text{O})_6^{2+}$ and $\text{A}(\text{H}_2\text{O})_6^{3+}$ the rate for the case $\text{A} = \text{Cr}$ is many orders of magnitude less than when $\text{A} = \text{Fe}$ or Co . This behavior has been ascribed to the fact that with Cr the above reaction involves hydrated ions of markedly different radii, in contrast to the cases of Fe and Co . On the basis of this interpretation the rate for the case $\text{A} = \text{Mn}$ should also be small, since the aquo complexes of di- and trivalent Mn differ considerably in size. However, early studies of the Mn system indicate an exchange rate as great as those observed with Fe and Co . This prompted reinvestigation of the Mn system. Various precipitation and solvent extraction procedures were used to separate the Mn^{2+} and Mn^{3+} . In all instances complete exchange in 6 F perchloric acid was observed within the shortest time required to separate the reactants, which left open the question of separation-induced exchange.

An attempt was made to estimate the exchange rate by an indirect method. This can be accomplished, in principle, by studying redox reactions of the type

in which Z^{2+} is a suitable reducing agent. It follows from certain simple assumptions that if k_{12} is the rate constant for the Z^{2+} - Mn^{3+} reaction, and k_1 is the rate constant for the Mn^{2+} - Mn^{3+} exchange, then

$$k_{12} \cong (k_1 k_2 K_{12})^{1/2} \quad (1)$$

where k_2 is the rate constant for the Z^{2+} - Z^{3+} exchange, and K_{12} (which must not be too large) is the equilibrium constant for the redox reaction. Preliminary experiments with $\text{Z} = \text{Fe}$ suggest that the Mn^{2+} - Mn^{3+} exchange is indeed relatively slow, as expected from the considerations above. However, a direct measurement has not yet been possible.

Equation (1) is found to be satisfied by a variety of reactions involving Fe^{2+} , Fe^{3+} , Ce^{3+} , $\text{Ce}(\text{IV})$, $\text{Fe}(\text{phen})_3^{2+}$, and $\text{Fe}(\text{phen})_3^{3+}$ ions. It has been confirmed, for example, in a determination of the rate constant for the oxidation of $\text{Fe}(\text{phen})_3^{2+}$ by $\text{Ce}(\text{IV})$. A value of $3 \times 10^5 \text{ F}^{-1} \text{ sec}^{-1}$ was calculated for the rate constant for the $\text{Fe}(\text{phen})_3^{3+}$ - $\text{Ce}(\text{IV})$

Figure 1. Analysis of the products of $\text{Cr}(\text{II})$ - $\text{Tl}(\text{III})$ reaction in the presence of chloride ions.

reaction, which is in satisfactory agreement with the value $1.42 \times 10^5 \text{ F}^{-1} \text{ sec}^{-1}$ observed in a direct measurement made on 0.5 F sulfuric acid solutions at 25°C .

The System H_2O_2 - $\text{Ce}(\text{IV})$

The kinetics of the H_2O_2 - $\text{Ce}(\text{IV})$ reaction have been investigated in 0.8 N sulfuric acid at 25°C . The results are consistent with the reaction scheme

with $k_1 = 1.0 \times 10^6 \text{ F}^{-1} \text{ sec}^{-1}$, and $k_2/k_{-1} = 13 \pm 2$. (A measurement of the HO_2 lifetime is reported under the heading "Chemical Effects of Radiation" below.)

The System $\text{Cr}(\text{II})$ - $\text{Tl}(\text{III})$

The products of the $\text{Cr}(\text{II})$ - $\text{Tl}(\text{III})$ reaction in perchloric acid containing various concentrations of chloride ion have been separated by ion-exchange chromatography. In Figure 1 the yields of the various $\text{Cr}(\text{III})$ species are plotted as a function of the chloride concentration. In the absence of added chloride, $\approx 90\%$ of the $\text{Cr}(\text{III})$ is found in the form of a dimer, and the remainder as the hexaquo-ion. The yield of CrCl_2^{2+} increases with

increasing chloride ion concentration, which suggests that the reaction between Cr^{2+} and TiCl_2^{2+} proceeds via a chloride-bridged activated complex. No CrCl_2^{2+} is produced in the reaction.

Studies on Fephen^{2+}

The kinetics of the formation and dissociation of Fephen^{2+} has been studied in perchloric acid solutions. A mechanism consistent with the observed acid dependence of the rate involves the participation of intermediates in which the Fe(II) is attached to only one nitrogen ion. The energy of activation for the Fephen^{2+} dissociation is considerably lower, and the entropy of activation considerably more negative, than in the corresponding case of Fe(phen)_3^{2+} dissociation. This can be interpreted in terms of crystal field effects. As against Fephen^{2+} and Fe^{2+} , which are paramagnetic, the Fe(phen)_3^{2+} ion is diamagnetic, which indicates that the latter complex is stabilized to a considerable extent by crystal field effects. The corresponding relatively high activation energy for its dissociation can be ascribed to the loss of a large amount of this crystal-field stabilization energy in going from the diamagnetic Fe(phen)_3^{2+} to the paramagnetic Fe(phen)_2^{2+} ion. The high positive value of the entropy of activation for the Fe(phen)_3^{2+} dissociation can be attributed to the increase in the mobility of the ligands in forming the activated complex. Similar spin changes do not accompany the Fephen^{2+} dissociation.

REACTION MECHANISMS IN ORGANIC SYSTEMS

Disproportionation of Ethylbenzene

Experiments have continued⁹ on the mechanism of the disproportionation of ethylbenzene-1- C^{14} in contact with $\text{Al}_2\text{Br}_6\text{-HBr}$ catalyst. The distribution of activity in the ring of ethylbenzene- C^{14} recovered from the reaction mixture has been re-determined with much improved accuracy by means of a novel degradation scheme involving the reduction of ethylbenzene to 1-ethylcyclohexene, the oxidation of the latter to the keto acid, and the reduction of the keto acid to caprylic acid. This procedure permits a direct determination of the C^{14} in the *ortho*, *meta*, and *para* positions. The new results on C^{14} distribution in the ring, for two different times of contact with catalyst, are shown in the following diagrams. It is seen that the reaction path leads predominantly to *para* labeling.

The activity is introduced into the *para* position by an exchange reaction of the *para*-diethylbenzene-1- C^{14} with ethylbenzene-1- C^{14} , which is more rapid than that of ethylbenzene- C^{14} with itself. This pathway is consistent with the following related observations: (1) the ratio of *meta*-diethylbenzene to *para*-diethylbenzene changes from 1.8 at 3% reaction to 10 at 40% reaction; (2) ethylbenzene- C^{14} and *meta*-diethylbenzene- C^{14} are rapidly formed when inactive ethylbenzene is treated with *para*-diethylbenzene- C^{14} in the presence of $\text{Al}_2\text{Br}_6\text{-HBr}$ at a contact time of 3 min.

In the case of the analogous problem of the HF-BF_3 catalyzed reaction between ethylbenzene and xylenes, it has been proposed in the literature that the alkylating agent is *meta*-diethylbenzene. In view of the preceding results, this suggestion seems unlikely.

Photolysis of $\text{C}_3\text{O}_2\text{-C}_2\text{H}_4$ Mixtures

Studies on carbon recoil reactions (reported below under the heading "Chemical Effects of Nuclear Transformation") are being complemented by reaction mechanism studies on small carbon-containing fragments, since the latter may help to elucidate the former. The reaction selected for study was the photolysis of carbon suboxide-2- C^{14} in the presence of ethylene, since, as reported in the literature, the major reaction products are allene and CO , which are also found when C^{11} is produced by nuclear bombardment of ethylene.

Allene- C^{14} produced in the photolysis of $\text{C}_3\text{O}_2\text{-CH}_4$ mixtures was degraded and found to have the following distribution of activity:

The same distribution is found for 8% and 47% conversion of C_3O_2 , which indicates that a secondary photolytic rearrangement of allene-2- C^{14} is not

responsible for the incorporation of activity in the terminal carbons.

This information does not settle the question of whether ketocarbene [C_2O] or atomic carbon is the reactive intermediate in the formation of allene. If the activity distribution is a reflection of the relative rates of double bond insertion and single bond insertion, it shows a degree of discrimination incompatible with the expected voracious reactivity of atomic carbon.

Work with this new and interesting reactive intermediate is continuing.

CHEMICAL EFFECTS OF RADIATION

The Nature of the "OH" Radical Produced in the Radiation Decomposition of H_2O

Water under irradiation decomposes to reducing and oxidizing free radicals which in the past have usually been called H and OH. In last year's annual report experiments were described which showed that the reducing radical is not in fact a hydrogen atom, but a negative ion, presumably a solvated electron. The method used was to determine the effect of ionic strength on the rates of reaction of the reducing radical with positive ions, negative ions, and neutral molecules.

This method has now been used to determine the charge on the oxidizing radical, generally called OH. If "OH" is really electrically neutral, the ratio of its reaction rate with an ionic species to that with a neutral species will be unaffected by ionic strength. Convenient neutral and ionic species for this purpose are ethyl alcohol and Br^- . "OH" is known to react with ethyl alcohol to yield a radical which in turn reacts with dissolved oxygen to form hydrogen peroxide. "OH" is also known to react readily with Br^- to give bromine atoms which in turn react with hydrogen peroxide to decompose it. Thus, a measurement of the amount of H_2O_2 produced by irradiating a solution containing bromide, alcohol, and oxygen permits computation of the relative amounts of reaction which the "OH" undergoes with the bromide ion and alcohol. The experiments showed that the yield of peroxide as a function of bromide and alcohol concentrations was independent of the ionic strength when the latter was changed by addition of potassium or lithium perchlorate to the solution. This is good evidence that the oxidizing radical is a neutral body, and so should probably

be represented by the traditional formula OH. The positive ion that must be formed at the same time to maintain charge balance is presumably the hydrogen ion H^+ . Thus the radiolysis of water is best thought of as leading primarily to the formation of the three bodies OH, H^+ , and the solvated electron.

$HClO_4$ Solutions

The radiation chemistry of perchloric acid solutions has been studied up to high concentrations of perchloric acid. It is inferred from the following observations that the initial products of the action of radiation on perchlorate ion are chlorate ion and *molecular* oxygen. (1) The chlorate yield is proportional to the concentration of perchlorate; hence chlorate production is due to the direct action of radiation on the perchlorate ion rather than to secondary reactions between the perchlorate ion and radicals formed from the water. (2) The yield of hydrogen produced from the water decreases linearly with increasing concentration of perchloric acid, which probably indicates that the perchlorate ion does not lose an electron to give the radical $ClO_4^\cdot$, but rather dissociates directly to ClO_3^- and oxygen. (3) That the oxygen is produced initially in the molecular, rather than the atomic, form is indicated (but not conclusively) by the results of experiments with various substances added to the perchloric acid solutions. Perhaps the oxygen is initially produced as an electronically excited oxygen atom which immediately reacts with a ClO_4^- to form O_2 and another ClO_3^- . The perchlorate ion is apparently inert toward the free radicals formed in the radiolysis of water.

Co^{+++} Complexes

In studies on the radiation chemistry of $Co(NH_3)_6^{+++}$ and $Co(NH_3)_5H_2O^{+++}$ it was found, contrary to expectation, that the yields of divalent cobalt were very small and that the corresponding oxidized product was not peroxide or oxygen, but nitrogen. The results indicate that the reducing (solvated electron) radicals from the water do not react with these ammonia complexes of cobalt. Rather, the ammonia molecules in the complex are attacked by the oxidizing (hydroxyl) radicals. Some of the ammonia is oxidized to nitrogen, the complex breaks up, and the uncomplexed trivalent cobalt is then readily reduced to divalent cobalt

by reducing radicals. The inertness of the cobalt in these complexes toward the reducing radicals contrasts strongly with the ease with which complexes of trivalent iron are reduced by these radicals. A parallel may be drawn with the rates of the electron exchange reaction of these complexes. It is a familiar fact that electron exchange occurs relatively rapidly between divalent iron and complexes of trivalent iron but very slowly between divalent cobalt and the ammonia complexes of trivalent cobalt.

Ag Salts

Solutions of silver ion, under irradiation, are expected to react very readily with solvated electrons to give a precipitate of metallic silver. With solutions of silver nitrate or silver perchlorate, however, the yield of metal is found to be very small. This has been traced to the reoxidation of the metal to the ion by the hydroxyl radical. By adding reagents such as nitrite ion or alcohol, which react readily with OH, the yield of metallic silver was greatly increased.

The HO₂ Radical

The radical HO₂ is known to be formed in the irradiation of solutions containing oxygen and it can also be formed as an intermediate in the ordinary oxidation of H₂O₂ by Ce(IV) (see above, under the heading "Kinetic Studies in Some Inorganic Systems").

A determination of the lifetime of HO₂ produced in the latter reaction has now been made by means of electron spin resonance in order to check the measurement described in the 1960 annual report, which was made on electron-irradiated water by use of chemical detection. Rapidly mixed solutions of ceric sulfate and hydrogen peroxide, with peroxide in large excess, were made to flow rapidly through the microwave cavity of the Varian EPR machine. A spin resonance absorption was found which agreed with that expected for the HO₂ radical. The rate of decay of the radical was measured by determining the variation in strength of this signal as the time of flow from point of mixing to the cavity was varied. The radicals were found to disappear in the expected second-order process, with a rate constant of 5×10^6 liters/mole-sec at 25°C and an activation energy of 5.9 ± 0.4 kcal/mole.

The rate was much higher than that expected on the basis of the flow experiments with electron-irradiated water mentioned above. A repetition of these latter measurements has shown that the active species formed decays by a first-order rather than a second-order process. It is therefore probably not a free radical but some chemically reactive intermediate formed by the combination of free radicals. The rate of decomposition of this new species is found to be proportional to the hydrogen ion concentration in the solution, which indicates that its disappearance is an acid-catalyzed reaction. The new species is conjectured to be either H₂O₄, produced by the combination of two HO₂ radicals, which would then decompose by an acid-catalyzed reaction to H₂O₂ + O₂; or H₂O₃, formed by combination of HO₂ and OH, which would decompose to water and oxygen. Study of these reactions is continuing.

EPR STUDIES ON FREE RADICALS IN GAMMA-IRRADIATED (CH₃)₄NX

The influence of the halide ion on the radicals formed in the isomorphous crystalline tetramethyl ammonium halides under gamma irradiation has been studied by means of EPR. The powdered iodide, bromide, and chloride, irradiated and observed at 77° K, all show the presence of methyl radicals which anneal out at 150° K. This is the only radical present in the iodide. The bromide and the chloride show the presence of another radical, tentatively identified as (CH₃)₃N⁺CH₂, which, after the methyl radical is annealed out, gives the pattern of a poorly resolved triplet (1:2:1), the resolution of which did not improve when single crystals were used. It is to be expected that such a radical is more likely to be produced by bromine and chlorine than by iodine. This radical anneals out at room temperature.

In addition to these two radicals, the irradiated chloride shows a third paramagnetic species, with a very complicated pattern, which has been identified as N⁺(CH₃)₃. It has been studied at room temperature (where the two other radicals anneal out) on irradiated single crystals of N(CH₃)₄Cl (tetragonal, *P*₄/*nmm*). The EPR patterns obtained when the crystal is rotated with its fourfold axis perpendicular to the magnetic field show that in N⁺(CH₃)₃ the nitrogen coupling constant has cylindrical symmetry, with principal values of the

coupling tensor 13.2, 20.6, and 20.6 gauss; and all 9 protons are equivalent, all having the coupling constant 26.6 gauss. The intensities of all 30 peaks observed at room temperature are consistent with the 9 protons' being equivalent. Since the isotropic nitrogen coupling is fairly small, the $N^+(CH_3)_3$ radical is probably close to planar. At low temperatures its EPR spectrum is much more complicated, and the nitrogen coupling tensor is no longer cylindrically symmetrical about the four-fold axis.

FREE RADICALS PRODUCED ON ULTRAVIOLET IRRADIATION

The photochemical rearrangement of *o*-nitrobenzaldehyde to nitrosobenzoic acid is known to occur both in the crystal and in solution. A further study of this rearrangement has been made by carrying out the photolysis ($\lambda > 3800 \text{ \AA}$) within the cavity of an EPR spectrometer. Paramagnetic species were found during the photolysis of 0.1 to 1.0 *M* solutions of *o*-nitrobenzaldehyde in deoxygenated ethyl alcohol and isopropyl alcohol. The well-resolved fine structure can be readily interpreted in terms of nitrogen and proton hyperfine splitting. The observed hyperfine coupling constants are compared with those from the electrochemically produced negative ion of *o*-nitrobenzaldehyde in acetonitrile in Table 1. The photo-radical is characterized by a large coupling constant for one proton, while the nitrogen coupling constants in the two radicals are comparable. This indicates that in the photo-radical one proton is much closer to the nitrogen than in the negative ion, and possible structures for the photo-radical are

The EPR spectrum of the negative ion in acetonitrile with added alcohol shows a definite solvent effect. The change in the spectrum can be attributed to an increasing nitrogen coupling constant, other coupling constants remaining essentially unchanged.

A comparison of the steady-state free radical concentration with the amount of nitrobenzoic

Table 1
Hyperfine Coupling Constants (in Gauss)

<i>o</i> -Nitrobenzaldehyde	a_N	a_1	a_2	a_3	a_4	a_5
Photo-produced radical	8.1	11.2*	2.9**	1.0**		
Negative ion	7.2	3.4*	2.4*	1.0*	0.6*	0.3*

*Interaction with one proton.

**Interaction with two equivalent protons.

acid formed during the photolysis indicates that the photo-radical observed is not a main intermediate in the rearrangement, but, since similar photosensitive nitro compounds also give paramagnetic species on photolysis, it appears to be closely connected with the rearrangement.

Paramagnetic species were also found when the powder was photolyzed at low temperatures. Further work is in progress on single crystals.

CHEMICAL EFFECTS OF NUCLEAR TRANSFORMATION

A number of nuclear reactions, $C^{12}(p,pn)C^{11}$, $C^{12}(n,2n)C^{11}$, and $N^{14}(p,\alpha)C^{11}$, are being used to investigate the chemistry of C^{11} recoils, a program begun at Brookhaven some years ago.

The production of C^{11} recoils by $N^{14}(p,\alpha)C^{11}$ is particularly well suited to these studies since the cross section is high (as measured here, ≈ 0.65 barns for 8.5-Mev protons). Experiments can be done over a range of 10^5 in dose, so that the effects of radiation damage to the system can be separated from the C^{11} recoil reactions of interest.

C^{11} Recoils in NH_3

The proton irradiations were carried out on purified, dry, gaseous ammonia. In Figure 2 the yields of the two major reaction products, $C^{11}H_4$ and $C^{11}H_3NH_2$, are plotted as a function of the radiation dose. (Below 10^{-2} ev/molecule a third species, as yet unidentified but readily converted by chemical means to methylamine- C^{11} , appears in 20% yield.) At doses above 0.1 ev/molecule $C^{11}H_4$ is almost the sole reaction product. This is an effect of radiation damage, as shown by auxiliary experiments in which mixtures of $C^{11}H_4$ and $C^{11}H_3NH_2$ were converted to $C^{11}H_4$ by exposure to large radiation doses. Charge and/or excitation

transfer mechanisms involving NH_3 are believed to account for the reductive process. The ammonia system is being studied in aqueous, liquid, and solid phases in the presence of scavengers and further moderators.

C^{11} Recoils in Hydrocarbons

The reactions of C^{11} recoils in CH_4 have been studied with C^{11} from the $\text{C}^{12}(p,pn)$ and $\text{C}^{12}(n,2n)$ nuclear reactions. It was found that the anomalous appearance of CO in this system was not due to wall effects, as has been suggested in the literature, but to traces of O_2 in the tank methane. Reduction of the O_2 impurity to <1 part in 10^5 eliminated the formation of C^{11}O . The same distribution of reaction products is obtained with C^{11} from both nuclear reactions. In addition to the major products acetylene and ethylene, saturated products such as methane, ethane, and propane were observed at doses of 5.3×10^{-4} ev/molecule. The proportion of saturated products increased with dose, which indicated radiolytic reduction, as in the case of ammonia. The additives O_2 , C_2H_2 , and CH_3NH_2 , which have lower ionization potentials than methane, protect the recoil $\text{HC}\equiv\text{CH}$ and $\text{H}_2\text{C}=\text{CH}_2$ against reduction. Acetylene and methylamine are more effective than oxygen.

The scavenger effect of oxygen is shown in Figure 3. The curves are similar to those found for scavengers in numerous other systems involving nuclear recoil. It is clear that O_2 at very low concentration effectively suppresses the thermal products. As the O_2 concentration increases, the O_2 either competes with the methane for the species leading to acetylene and ethylene, or it may oxidize a highly excited acetylene or ethylene molecule.

Figure 2. Yields of C^{11}H_4 and $\text{C}^{11}\text{H}_3\text{NH}_2$ in proton irradiation of gaseous NH_3 , as a function of dose.

Figure 3. Effect of O_2 on the yields of C_2H_2 and C_2H_4 in the proton bombardment of CH_4 . Dose = 5.2×10^{-4} ev/molecule.

Corresponding studies on propane have shown that for most reaction products the yields in liquid targets differ from those in gas. In the liquid, the yields of acetylene and ethylene are lower and the ratio of the synthesis products butane and isobutane is statistical, 3/1 (based on available hydrogen positions in propane), a drastic change from the nonstatistical butane/isobutane = 0.76/1 observed in gas. This behavior exactly parallels that of the well-characterized hot methylene produced by photolysis, and therefore suggests that methylene is the species reacting with propane to give butanes in the C^{11} system.

Hot Atom Reactions in $\text{Co}(\text{NH}_3)_6\text{Br}_3$

Studies of recoil reactions in the ammino complexes of Co(III) have continued. New results have been obtained on the yields and annealing behavior of different radioactive and chemical species formed in crystals of $\text{Co}(\text{NH}_3)_6\text{Br}_3$ irradiated in the reactor at dry ice temperatures. The chemical entities are separated by paper electrophoresis after dissolution of the irradiated solid in 5% acetic acid at 0°C . The yields of four different radioactive species of the radiosynthetic ion $\text{Co}(\text{NH}_3)_5\text{Br}^{++}$ were found to be Co^{60} , 0.048; Br^{80} , 0.116; Br^{80m} , 0.117; and Br^{82} , 0.072. When the crystals are annealed isothermally at temperatures in the range 0° to 55°C after irradiation, the yields decrease. Activation energies for the annealing reactions are estimated to be 0.95 ev for Br^{80m} and Br^{82} and roughly 0.8 ev for the Co^{60} .

A possible mechanism for the annealing out of $\text{Co}(\text{NH}_3)_5\text{Br}^{++}$ is reduction to the Co(II) complex ion in the lattice by capture of an electron. A corresponding explanation has been given for the

observation that annealing converts Co(II) into the Co(III) complex $\text{Co}(\text{NH}_3)_6^{+++}$. Here the Co(II) is thought to be present in the lattice as $\text{Co}(\text{NH}_3)_6^{++}$, which on thermal annealing ionizes an electron to the conduction band.

The effects of the isomeric transition of Br^{80m} to Br^{80} were also studied in $\text{Co}(\text{NH}_3)_6\text{Br}_3$ crystals into which Br^{80m} had been introduced chemically. As in the case of the bromine recoils from neutron capture, Br^{80} -labeled $\text{Co}(\text{NH}_3)_5\text{Br}^{++}$ is found as a product, the yield being 5.6%. Its formation may be due to a Varley mechanism: in the isomeric decay of Br^{80m} , a negative bromine at a lattice position of minimum energy is converted into a bromine of high positive charge, which thus finds itself at an energy maximum, and so acquires enough kinetic energy to displace an NH_3 from a nearby $\text{Co}(\text{NH}_3)_6^{+++}$.

Annealing at Low Temperatures

A series of studies was made to investigate the importance of annealing reactions at low temperatures during neutron irradiation and subsequent chemical processing. Surprisingly large effects have been found in some instances. In the case of potassium permanganate crystals, for example, in which for years the accepted value of the retention (i.e., yield of Mn^{56} in the form of MnO_4^- after neutron irradiation) has been 20%, the retention is found to be 5.1% when crystals are irradiated at -195°C and dissolved (for chemical processing) in acetone at -95°C . Intermediate values are found at somewhat higher temperatures. Similar effects of smaller magnitude have been found for other crystals. It appears that thermal annealing at low temperatures can be more serious than has been thought and must be allowed for in quantitative interpretations of the yields of hot atom reactions in solids.

ISOTOPE EFFECTS ON VAPOR PRESSURE

In the theory of vapor pressures of isotopically substituted molecules, the classical partition functions are corrected by a power series of the form

$$1 + (hc/kT)^2 \sum \omega_i^2 + (hc/kT)^4 \sum \omega_i^4 + \text{higher terms},$$

where the sums are over all intermolecular and intramolecular vibrations. The relative vapor pressures of $\text{N}^{15}\text{N}^{14}\text{O}^{16}$ and $\text{N}^{14}\text{N}^{15}\text{O}^{16}$ previously measured here show that the term in ω^2 is important, and that a partition function that depends only on molecular masses is not valid.

Since then, the importance of the term in ω^4 has been investigated by measurements of the relative vapor pressures of the six isotopically substituted ethylene molecules: C_2H_4 , $\text{C}_2\text{H}_3\text{D}$, *cis*- $\text{C}_2\text{H}_2\text{D}_2$, *trans*- $\text{C}_2\text{H}_2\text{D}_2$, *gem*- $\text{C}_2\text{H}_2\text{D}_2$, and $\text{C}^{12}\text{C}^{13}\text{H}_4$. The vapor pressures of the first five of these species were determined directly by precise manometric measurements over the range 120° to 180°K . The vapor pressure of the sixth, $\text{C}^{12}\text{C}^{13}\text{H}_4$, was measured relative to that of $\text{C}^{12}_2\text{H}_4$ in distillation experiments, with $\text{C}_2\text{H}_3\text{D}$ used as an internal standard. The relative vapor pressures are in the order *trans*- $\text{C}_2\text{H}_2\text{D}_2 > \text{gem}$ - $\text{C}_2\text{H}_2\text{D}_2 > \text{cis}$ - $\text{C}_2\text{H}_2\text{D}_2 > \text{C}_2\text{H}_3\text{D} > \text{C}^{12}_2\text{H}_4 > \text{C}^{12}\text{C}^{13}\text{H}_4$.

The effect of the term in ω^4 makes itself manifest in the vapor pressure difference between geometric isomers of $\text{C}_2\text{H}_2\text{D}_2$ which cannot be ascribed to the terms in ω^2 , since the latter are identical for the different isomers. Another effect of the term in ω^4 is the observation that the isotope effect for $\text{C}_2\text{H}_2\text{D}_2$ is not exactly twice that of $\text{C}_2\text{H}_3\text{D}$. The difference between $\text{C}_2\text{H}_3\text{D}$ and $\text{C}^{12}\text{C}^{13}\text{H}_4$, which is due to terms in ω^2 , fits in with previously reported findings for N_2O .

IBM 7090 computer calculations have been made of the intramolecular and intermolecular frequencies, and their changes upon condensation to the liquid or solid phase, for the six molecules listed above and for C_2HD_3 , C_2D_4 , $\text{C}^{12}\text{C}^{14}\text{H}_4$, and $\text{C}_2\text{H}_3\text{T}$. The calculations involve the diagonalization of the full 18×18 matrix of the vibrational secular equation set up in Cartesian displacement coordinates. The computed differences between molecular vibrations in the gas and liquid phases gave vapor pressure effects in excellent agreement with the experimental ones. Further calculations are in progress to determine the precise effect of the assumed force field on these differences.

More general calculations have been made to evaluate some of the possible origins of the change in intramolecular vibrations of a molecule that enters a condensed phase. In this work it has been found extremely useful to apply a perturbation method based on the fact that the zero-point vibrational energy is the energy of the ground state in the quantum mechanical vibrational problem. The Hamiltonian is written in terms of Wilson's *F* and *G* matrices. The unperturbed system is taken to be a system of independent oscillators, and the perturbation terms (which are readily evaluated) are found to converge very rapidly.

One of the effects investigated in this way is that of van der Waals' interactions on a molecule in the condensed phase. These interactions have been found to lead to a decrease in intramolecular force constants, which explains why in some cases the heavier isotopic species has the higher vapor pressure. Good agreement is found between isotope effects calculated from this model and those found experimentally in the cases of $C^{12}Cl_4/C^{13}Cl_4$ and $C^{12}Cl^{35}_4/C^{12}Cl^{37}Cl^{35}_3$.

Another source of the changes in internal frequencies of a molecule in the condensed phase is that degrees of freedom corresponding to free rotations and translations of the molecule in the gas phase become hindered, and so couple with the intramolecular vibrations. This coupling is isotope dependent, since the nature of the hindered degrees of freedom depends on isotopic substitution, whereas the potential energy surface does not. The magnitude of the isotope effect due to these interactions cannot be neglected in treating the vapor pressures of isotopic molecules.

SECONDARY ISOTOPE EFFECTS

Thermal Decomposition of $Hg(CH_3)_2$

In general, in reactions of the type $R_1R_2C(H \text{ or } D)X \rightarrow R_1R_2C(H \text{ or } D) + X$ at room temperature, α -deuterium-labeled compounds have been found to react ≈ 10 to 15% more slowly than normal compounds. This has been attributed to the change in one of the $C-H$ bending frequencies in going from the tetrahedral to the trigonal configuration during reaction. The isotope effect results because the vibrational zero-point energy difference between the labeled and normal compounds in the reactant (tetrahedral) is different from that in the activated complex (trigonal).

To gain insight into the nature of the transition state, the isotope effect in the thermal decomposition of dimethylmercury in the presence of cyclopentane was studied. The important reactions are

Reactions (1) and (1') are rate controlling in the presence of an excess of cyclopentane. Starting with a known mixture of deuterated and natural dimethylmercury and cyclopentane, the isotope

effect is obtained by measuring the deuterium ratio in the methane produced.

Contrary to expectation, the deuterium compound reacts $\approx 7\%$ faster than the natural compound. On the other hand, a normal C^{13} isotope effect is found. The only model for the activated complex that appears consistent with these observations is one in which the effect on the bending frequency described above is not present. Instead, there must be an increase in the $C-H$ stretching frequencies when the activated complex is formed. This is in agreement with other properties of trigonal $C-H$ bonds as compared with tetrahedral $C-H$ configurations, but has not previously been invoked in explaining isotope effects of this type.

The Diels-Alder Reaction

The use of the magnitude of the secondary α -deuterium isotope effect to help elucidate mechanisms of organic reactions is being pursued. The theoretical basis for alpha effects has been briefly outlined above. If a molecule has two breakable bonds, each with deuterium in the position alpha to it, then if both bonds break simultaneously the reaction rate is expected to be ≈ 20 to 30% lower than in the corresponding case with hydrogen. On the other hand, if the two bonds break stepwise, a normal 10 to 15% would be expected.

These considerations have been applied in a study of the Diels-Alder reaction, represented in its simplest form as

since some believe that the two new bonds are formed simultaneously, and others that they are formed in stepwise fashion. The rates of the reaction

were measured, with the following compounds:

The observed isotope effects are $k_I/k_{II}=1.16$, and $k_I/k_{III}=1.08$. The existence of an isotope effect k_I/k_{III} , although smaller than expected, indicates that bond *b* is breaking in the rate-controlling step. Since little progress seems to occur along the reaction coordinate in going from adduct to transition state, as evidenced by the low value of k_I/k_{III} , it appears that the ratio $k_I/k_{II}=1.16$ is really indicative of both bonds *a* and *b* breaking in the same step. Further experiments to put this interpretation on firmer ground are in progress. The mechanism, then, for the formation of adduct would be a concerted one-step process as demanded by the principle of microscopic reversibility.

VALENCE STATES IN CRYSTALLINE TiN

The valence states of Ti and N in TiN (rock salt structure) have been investigated by measuring the relative intensities of the different diffraction peaks from TiN powders and comparing them with the theoretical values expected on the basis of various assumed Ti and N valence states. Such a study is of interest because nitride structures in general are not understood theoretically, and also because TiN is a superconductor.

The theoretical intensity calculations were made for the assumed valence states Ti^0N^0 and $Ti^{3+}N^{3-}$. For Ti^0 and Ti^{3+} , use was made of Watson and Freeman's scattering factors and Dauben and Templeton's dispersion terms. For N^0 a Hartree-Fock scattering factor was used, and for N^{3-} a factor obtained by a crude extrapolation.

The relative intensities of the first ten reflections were measured with $CuK\alpha$ radiation.

A reflection can serve to distinguish between the two valence models only if its $\sin\theta/\lambda$ value is $< \approx 0.25$ Å, and if in it the Ti and N scattering factors are out of phase. Only the (111) and (200) reflections meet the first of these conditions (the other eight observed reflections served to confirm Dauben and Templeton's dispersion term), and, of the two, only the (111) reflection meets the second. For the two models, the calculated intensities of

the (111) reflection differ by 25%, that calculated for $Ti^{3+}N^{3-}$ agreeing with the observed value.

A NEUTRON DIFFRACTION STUDY OF *o*-NITROBENZALDEHYDE AND THE C—H—O INTERACTION

The structure of the stable modification of *o*-nitrobenzaldehyde, previously determined by x-ray diffraction, has been redetermined by means of neutron diffraction. Nonhydrogen-atom parameters agree with values from the x-ray determination. The parameters of the hydrogen atoms and a new set of anisotropic temperature factors have been obtained.

The observed positions of the hydrogen atoms indicate that the intramolecular interaction between the aldehydic hydrogen atom and the adjacent oxygen of the nitro group is weak, and that no hydrogen bond exists between these two atoms.

It has been generally assumed that temperature factors obtained from refinement of both x-ray and neutron diffraction data are appreciably affected by systematic errors in the data and the model, and are therefore less accurate than their standard deviations indicate. In the present investigation comparison of the x-ray and neutron temperature factors reveals the effect of these systematic errors, and it is shown that, in this case at least, they are of the same order of magnitude as the standard deviations derived from the least-squares refinement, and also that there is no significant difference between the means of the x-ray and neutron temperature parameters.

MAGNETIC STRUCTURE OF MANGANESE CHROMITE

Numerous attempts have been made to explain the anomalous properties of chromites and of mixed spinels containing chromium on the basis of the theoretical models of Néel and of Yafet and Kittel. Recent calculations by T.A. Kaplan and co-workers, using a classical Heisenberg model, have shown that over an appreciable range of the BB to the AB interaction the ground state is a ferrimagnetic spiral, i.e., a conical spin configuration in which the transverse components progress in spiral fashion along a fixed direction in the crystal. A neutron diffraction study of $MnCr_2O_4$ was undertaken to test the suitability of the various proposed models. From room temperature

data it is readily established that MnCr_2O_4 is a normal spinel with $<1\%$ of the Mn^{2+} ions present on B sites. The Curie temperature, as determined by diffraction, is $\approx 43^\circ\text{K}$. At 18°K , additional sharp peaks appear at positions that cannot be indexed either on the original unit cell or on any reasonably enlarged cell. No change in either the positions or intensities of the fundamental lines is observed in going through this transition. Application of a magnetic field along the scattering vector decreases the magnetic contributions to the fundamentals, but increases the intensities of the extra reflections. Analysis of the data shows that while little difficulty is encountered in explaining the observed magnetization data, the Néel and Yafet-Kittel models fail to account for major qualitative features of the diffraction results. On the other hand, the ferrimagnetic spiral model is in good qualitative accord with all the observations. The theoretical parameters give reasonably good quantitative agreement with the positions and intensities of the extra lines as well as the intensities of the fundamentals, while retaining the expected values for the individual ionic moments and the macroscopic saturation moment. Although some discrepancies still remain, the spiral model appears to be a good first approximation to the ground-state spin configuration.

LOW ENERGY NUCLEAR REACTIONS

Nuclear reaction studies currently in progress include excitation function measurements, investigations of the recoil properties of residual nuclei, and measurements on the charged particles emitted in nuclear reactions. These experimental projects are complemented by theoretical studies of nuclear reactions in the 10 to 50-Mev energy range.

Alpha Particles on Cd^{106}

The reactions of Cd^{106} with 15 to 40-Mev alpha particles from the Brookhaven 60-in. cyclotron are currently being investigated. Excitation functions have been obtained for a number of reactions, and peak cross sections of 5.2, 510, and 860 mb were measured for the (α, γ) , (α, n) , and $(\alpha, 2n)$ reactions, respectively. A number of reactions involving the emission of charged particles are also under investigation. The results are being compared with Monte Carlo evaporation calculations based on the statistical theory.

Protons on Ga^{69} and Ga^{71}

The reactions of alpha particles with nuclides in the iron-germanium regions have been fairly thoroughly investigated in previous years. It was of interest to obtain detailed results for proton-induced reactions in this same mass region for comparison purposes. A joint study with a group from the Institute for Nuclear Study of Tokyo University was undertaken for this purpose. The reactions of Ga^{69} and Ga^{71} with 15 to 56-Mev protons were investigated, and excitation functions for 25 reactions were obtained. The results were compared with Monte Carlo evaporation calculations by using pairing energy and level density parameter values that gave a good fit to data for alpha-induced reactions in the same mass region. In most cases good agreement with the experimental peak cross sections was obtained, which confirmed the occurrence of compound nucleus formation. Evidence for direct interaction mechanisms was also obtained, since many of the excitation functions exhibited prominent tails at energies past the peak. The excitation functions will be compared with a combined cascade-evaporation calculation in order to assess the relative contribution of direct interaction and compound nuclear processes.

Recoils of Residual Nuclei

The measurement of the recoil properties of the residual nuclei following a nuclear reaction can also yield information about the reaction mechanism. Recoil ranges projected along the beam direction have been measured for a number of reactions of indium with 5 to 10-Mev protons and 20 to 40-Mev alpha particles. The results have been compared with values expected for compound nucleus formation, and agreement is obtained for the (α, pn) , $(p, p\gamma)$, and (p, n) reactions. In the case of the $(\alpha, \alpha\gamma)$ reaction, on the other hand, the measured ranges are a factor of 3 to 6 smaller than the calculated values, which indicates a direct interaction process. A similar study has been carried out for the $\text{Al}^{27}(\alpha, \text{Be}^7)$ reaction. The results for Be^7 at 40 Mev are consistent with approximately equal contributions from direct interaction and evaporation processes, while the results for Na^{24} from alpha particles on Al^{27} indicate compound nucleus formation. It is concluded from cross section measurements that at 40 Mev $\approx 60\%$ of the yield of Na^{24} may be attributed to the $(\alpha, \alpha\text{He}^3)$ reaction.

The interpretation of recoil studies requires a knowledge of the range-energy relations for the recoiling atoms. Such were measured for Ne^{22} ions in aluminum and for gallium atoms in copper by means of the stacked foil technique, with analysis by means of a high-sensitivity mass spectrometer. For Ne ions of energies between 1.0 and 2.8 Mev, the mean ranges projected along the beam direction varied between 0.35 and 0.67 mg/cm² of aluminum, and the half-widths of the range curves were 0.20 mg/cm². The ranges of the gallium atoms, determined from measurements using monoenergetic atoms produced in radiative capture reactions such as (p,γ) , (d,γ) , and (α,γ) , varied between 7 and 174 $\mu\text{g}/\text{cm}^2$ of copper for gallium atoms of energies between 75 and 950 kev.

Charged Particles From Nuclear Reactions

Studies of charged particles emitted in nuclear reactions are being carried out in collaboration with members of the Physics Department. Current experiments at the 60-in. cyclotron involve observation of the correlations in energy and angle of two incident charged particles by means of solid state detectors in conjunction with a two-dimensional multichannel pulse-height analyzer. The most complete data obtained are for the reaction $\text{O}^{16}(\alpha,2\alpha)\text{C}^{12}$ (ground state). Certain features of the data are consistent with a model assuming the direct knock-out of the second alpha particle, while others are in agreement with the assumption of a relatively long-lived state in O^{16} between the emission of the two alpha particles. Preliminary data for the reaction $d(p,2p)n$ indicate a resonance corresponding to the singlet state of the deuteron; for $d(p,p')d$ evidence has been seen for electromagnetic transitions representing nuclear brehmsstrahlung. The principle of charge independence of nuclear forces predicts the existence of a dineutron equivalent to the singlet state of the deuteron. Studies of the reaction $d(d,2p)2n$ are now being carried out in order to observe this particle.

Theoretical Calculations

A Monte Carlo evaporation calculation described in a previous report has been programmed for the IBM 7090 computer to provide calculated excitation functions for comparison with experimental results. The calculation has been modified to permit calculation of the momentum imparted to the residual nucleus for comparison with the re-

sults of recoil studies. Also in progress is a more detailed calculation of excitation functions which considers the effect on excitation functions of the competition between gamma-ray and particle emission in the final de-excitation step. This competition depends in turn mainly on the energies E_i of the set of product levels that is composed of the lowest level at every angular momentum. A method for making approximate calculations has been developed by using plausible assumed distributions of the E_i 's. An analysis of experimental excitation functions for the reaction pair $\text{Ag}^{109}(\alpha,n)$ and $\text{Ag}^{109}(\alpha,2n)$ indicates that when gamma-ray emission is included larger values of the level density parameter a are required to fit the data than when gamma-ray emission is not included. This calculation is being programmed for the IBM 7090 computer.

Miscellaneous Nuclear Data

In the course of other experiments new information has been obtained on the decay of several nuclides. The half-lives of Kr^{76} and Y^{83} have been redetermined and found to be 15.0 ± 0.5 hr and 7.5 min, respectively. These values are in disagreement with previously reported values of 10 and 3.7 hr. The ratios of electron capture to positron emission in the decays of Cs^{125} and Cs^{127} have been measured as 1.03 ± 0.07 and 27.2 ± 1.7 , respectively.

HIGH ENERGY NUCLEAR REACTIONS

Investigations of the interactions of high energy protons with complex nuclei have been continued at the two high energy accelerators available at Brookhaven. A substantial part of this effort during the year has involved investigation of the change in cross section of various proton-induced reactions as the proton energy is increased from 3 to 30 Bev. Such relative cross sections obtained at a single laboratory are expected to be more reliable than comparisons making use of data obtained in different laboratories. Studies of carbon, aluminum, and copper targets have now been completed, and investigations of silver, lead, and uranium targets are in progress. All the data so far obtained show that no drastic changes occur in cross sections on going from 3 to 30-Bev protons. At 30 Bev it has been possible to investigate products of larger half-life or lower cross section than at 3 Bev be-

cause of the large integrated fluxes available at the AGS.

Aluminum Targets

The cross sections for producing products ranging from Mg^{27} to Be^7 have been measured for 28-Bev protons on aluminum. Excitation functions for a number of these reactions have been extended down to 0.4 Bev. These measurements were relative to the cross section of the $\text{C}^{12}(p,pn)\text{C}^{11}$ reaction for which an absolute measurement at 28 Bev was described in last year's report. Particular emphasis was placed on obtaining the cross section for the $\text{Al}^{27}(p,3pn)\text{Na}^{24}$ reaction which has been used as a monitor reaction for other studies. The cross section of this reaction decreases from 9.1 mb at 3 Bev to a value of 8.6 mb at 28 Bev. The energy dependence between 3 and 30 Bev of the other cross sections from aluminum is not very different. There is some indication that the production of F^{18} and Na^{22} is less probable (relative to Na^{24}) and that of Be^7 somewhat more probable at 28 Bev than at 3 Bev.

Copper Targets

The formation cross sections for the spallation products from copper irradiated with 30-Bev protons differ very little from those observed at 3 Bev in the cases of products within ≈ 30 mass numbers from the target. Within this mass region the ratio of 30 to 3-Bev formation cross sections is unity for neutron-excess products, and varies from ≈ 0.7 to unity for neutron-deficient species. Only the lightest products, Be^7 , Na^{22} , and Na^{24} , seem to be made with higher probability at the higher energy, the cross section ratio varying from ≈ 1.2 for Na^{24} to 1.5 for Na^{22} . A recent CERN report on the interactions of 24-Bev protons with copper indicated that, although the yield pattern for the spallation products observed was essentially the same as at lower energies, the total spallation cross section at 24 Bev was $\approx 40\%$ lower than the inelastic interaction cross section of copper. The present data are somewhat at variance with the CERN results. They show a small shift in the spallation yield pattern between 3 and 30 Bev, but no deficiency in total cross section at the higher energy.

Silver, Lead, and Uranium Targets

A comparison of yields of selected products from silver, lead, and uranium targets at 3 and 30 Bev

indicates that the trends observed with the light target nuclei are also true of the heavier targets. Preliminary results show that many cross sections change little in the 3 to 30-Bev energy region. Some appreciable changes have been observed (e.g., in the yield of Ar^{41} from uranium targets) but these occur mostly between 3 and 10 Bev. Changes between 10 and 30 Bev are generally smaller.

Theoretical Implications

The changes in yield patterns (particularly in the case of heavy element targets) with increasing proton energies between 0.3 and 3 Bev have been previously postulated to be primarily due to the production and reabsorption of π -mesons in intranuclear collisions. The number of such mesons produced per collision is known to increase significantly as the energy is increased from 3 to 30 Bev. Hence any theoretical description of nuclear reactions at energies $> \approx 3$ Bev must account for the fact that these additional mesons do not drastically change the deposition energy spectrum or observed yield patterns of the various product nuclides.

Product Nuclides of the Same Mass as the Target Nuclide

Considerable interest has been shown recently in high energy nuclear reactions in which the product nucleus has the same mass number as the target. Examples are the $(p,p\pi^+)$, $(p,p\pi^-)$ or (p,n) , and $(p,p\pi^0)$ reactions. Cross sections for such reactions have recently been calculated from elementary nucleon-nucleon and pion-nucleon interactions by Ericson, Selleri, and Van de Walle. Cross sections for $(p,p\pi^+)$ reactions on Al^{27} and Cu^{65} have been measured for 28-Bev protons at the AGS, and both $(p,p\pi^+)$ and $(p,p\pi^0)$ cross sections of indium have been obtained at 2.9 and 30 Bev. The results of these measurements and of other available measurements are in qualitative agreement with the theoretical estimates but are high by factors of ≈ 1 to 3. (A factor of 2 in the theoretical underestimate has been ascribed by Ericson to simplifications of the theoretical treatment.) Further studies on these reactions, including those of recoil properties, are in progress.

Miscellaneous Studies at the Cosmotron

Research has continued in a number of directions suggested by previously reported experiments. Studies have been made of the isobaric yield pat-

terms in the mass region 127 to 134 for the products produced by irradiation of uranium with 1, 2, and 3-Bev protons. A number of (p, pn) cross sections on targets having 36 to 46 neutrons have been measured to determine whether the $1f_{7/2}$ neutron shell becomes unavailable for the (p, pn) reaction at some point in this mass region, and thereby leads to a sudden decrease in the reaction cross section. Studies of the recoil properties of Na^{24} produced in the irradiation of bismuth with 2.9-Bev protons have been undertaken. This reaction is typical of the "fragmentation" process first observed when proton energies were increased above 400 Mev, and it is hoped that a further understanding of the mechanism will be gained from this work. Studies of the spallation of silver at 2.9 Bev and of the recoil properties of some selected product nuclei are in progress, and the results will be compared with data obtained with nuclear emulsions.

Numerous photographs of a liquid xenon bubble chamber exposed to $\approx 1\text{-Bev}/c$ π -mesons are available from studies of elementary particle production. These are being scanned by members of the Chemistry Department in a study of the interactions of π -mesons with xenon.

Theoretical Calculations

Work on the new Monte Carlo calculations of intranuclear cascade processes mentioned in last year's report has proceeded in collaboration with members of the Applied Mathematics Division. Programming for nonmesonic processes is essentially complete, and detailed comparisons of results with the earlier calculations by Metropolis et al. show excellent agreement when the same input parameters are used. The introduction of reflection and refraction of nucleons at the nuclear boundary, and substitution of a trapezoidal for a rectangular density distribution (with corresponding change in the shape of the potential well) both have rather pronounced effects; these refinements tend to decrease some of the discrepancies between the earlier calculations and experimental data. Progress has also been made in the formulation of the cascade processes involving mesons. The observed resonance character of elementary pion-nucleon interactions has the interesting consequence that the creation of pions in nucleon-nucleon collisions can be considered a two-step process, the first step being the production of an

excited nucleon isobar, the second step the decay of this isobar into a nucleon and a pion. The calculated lifetimes of these isobars are such that they can travel appreciable distances within the nucleus before decay, and therefore a consistent scheme for treating these isobars and their interactions had to be developed for inclusion in the cascade calculations.

NUCLEAR SPECTROSCOPY

The Os^{186} Level Scheme

Analysis has been completed of the data previously obtained in measurements made on radiations from Ir^{186} decay by members of the Physics and Chemistry Departments. Approximately 60 of the 100 observed transitions have been fitted into a 24-level scheme for Os^{186} . Five of these levels may be classified as members of a ground-state rotational band and six others as members of a rotational band based on a gamma vibration. In the ground-state band the energies are not well fitted by the strong-coupling model, even with inclusion of vibration-rotation interaction; in the gamma band, although the levels show even-odd energy effects predicted by the model, the agreement between observed and theoretical energy values is not satisfactory. Values computed from the asymmetric rotor model for $\gamma = 16.5^\circ$ fit the level energies as well as the strong-coupling values do, but with fewer parameters.

Results of comparisons of intra- and interband transition probabilities appear to show, in terms of the strong-coupling model, that the interband mixing required to fit these transition probabilities is different from that which gives the best fit for energy values. Similarly, in terms of the asymmetric rotor model, different values of the asymmetry parameter, γ , are required for transition probabilities and for energies.

Studies on other isotopes of osmium are in progress, and some new data on Os^{185} , Os^{187} , and Os^{189} have been obtained.

Improvements on the Beta-Ray Spectrometer

The resolving power and transmission of the 50-cm double-focusing beta-ray spectrometer have been considerably improved by appropriate magnetic shimming. It has been possible to achieve a line width (full width at half-maximum intensity) of $< 0.05\%$; this is only about 10% greater than

the theoretical minimum for the source and detector widths employed. A gratifying increase in transmission has accompanied the improvement in resolution. Further, operation of the spectrometer and recording of the data have been made fully automatic.

ATOMIC ELECTRON EJECTION IN Po^{210} DECAY

In continuation of last year's report under the same heading, it is now reported that the probability of a lead M photon being emitted in the decay of Po^{210} is $0.91(\pm 15\%) \times 10^{-3}$, as measured by means of a proportional counter and pulse-height analyzer. In the course of this measurement the relative intensities of the component lines in the lead M spectrum from Po^{210} decay were observed to differ significantly from those in a lead M spectrum excited by conventional means. In particular the spectrum from Po^{210} contains a 3.7-keV component, constituting $\approx 4\%$ of the total spectral intensity, whose energy is consistent with the allowed, but hitherto unreported, transition $M_{\text{I}}O_{\text{II,III}}$. These intensity differences are attributed to extensive ejection of electrons from the outermost shells in the Po^{210} decay act.

Migdal's theory of electron ejection in alpha decay, which neglects the effect of nuclear recoil and calculates electron transitions only to continuum states, underestimates the observed M intensity by a factor of ≈ 150 , and, as previously reported, it underestimates the L intensity by a factor of ≈ 25 , and gives a roughly correct estimate of the K intensity. When the theory is corrected by introducing the effect of nuclear recoil and transitions to bound states, the underestimate becomes worse, the ratio of measured value to theoretical value for M , L , and K electrons being respectively 410, 270, and 15. Levinger has shown that for L electrons, but not for K electrons, one factor of 10 in the discrepancy may be attributed to the premature amputation of a series used by Migdal. Presumably this accounts for a factor of 10 in the M discrepancy also. Thus, for all three shells a discrepancy of one order of magnitude remains. As a result of a critical reworking of the theory it is now made very plausible that the remaining order-of-magnitude discrepancy is due to an unjustifiable use of hydrogenic wave functions in the theory. Comparison of a considerable body of experimental data in both alpha and beta decay with theo-

retical estimates made by use of hydrogenic wave functions reveals that the theoretical estimates are low by an order of magnitude in all cases except those in which the calculations are made by appeal to the completeness property. In the latter cases, the use of hydrogenic wave functions is justifiable.

SPATIAL CONSTANCY OF COSMIC RAYS

The intensity of cosmic rays in the neighborhood of the earth can be compared to the average intensity over the orbit of a meteorite by determining the relative activities of cosmic-ray-induced Ar^{37} (35 day) and Ar^{39} (325 yr) in the meteorite material. This method was applied to the Hamlet meteorite (see annual report for 1960) and has been repeated on 350 grams of Harleton meteorite obtained shortly after its fall on May 30, 1961.

The Harleton meteorite was a typical chondrite consisting of a stone phase interspersed with a metallic phase (87% Fe, 13% Ni, 0.4% Co) in the form of small grains 0.5 mm in diameter. The iron phase was separated magnetically from a ground sample of the meteorite to measure the $\text{Ar}^{37}/\text{Ar}^{39}$ ratio on the separated metallic phase. In this way a more certain test for variations in cosmic-ray intensity can be made because the $\text{Ar}^{37}/\text{Ar}^{39}$ ratio produced in iron by proton bombardment at the Cosmotron is accurately known and this ratio is independent of the proton energy. Measurements of the $\text{Ar}^{37}/\text{Ar}^{39}$ ratio were made separately on the whole meteorite sample, the stone phase, and the iron phase. The values are given in Table 2 together with the $\text{Ar}^{37}/\text{Ar}^{39}$ ratio for iron bombarded with 3-Bev protons at the Cosmotron. Since these ratios are the same within the limits of error it is concluded that the cosmic-ray intensity at the

Table 2
Argon Activities in the Harleton Meteorite

Sample	Ar^{39} , dis/min/kg	$\text{Ar}^{37}/\text{Ar}^{39}$	$\text{Ar}^{37}/\text{Ar}^{39}$ from 3-Bev proton bombardment
Iron phase	22.0 ± 1.1	0.84 ± 0.11	0.82 ± 0.02 measurement incomplete
Stone phase	7.0 ± 0.4	1.58 ± 0.25	
Whole meteorite	9.1 ± 0.4	1.02 ± 0.13	measurement incomplete

earth's orbit is within 15% of the time-averaged value over several astronomical units of distance from the sun.

ARCHAEOLOGICAL CHEMISTRY

Glasses Containing Antimony and Manganese

In the continuing investigation of the composition of ancient glasses special attention has been given to colorless, clear glasses containing antimony and manganese in sufficiently great concentrations to indicate that compounds of these elements were added deliberately, presumably to counteract the discoloration due to iron. The alternate and joint use of these two decolorants form an interesting and significant pattern that correlates with the geographic and chronological origins of the glasses. The observation that this use of antimony was distinctly different from the utilization of lead and tin in ancient glass tends to refute the frequently expressed opinion that ancient artisans did not differentiate between antimony and these elements.

Glasses Containing Magnesium and Potassium

Earlier data had shown that a particular type of glass, characterized by a high concentration of magnesium and potassium, was widely produced in the second millenium B.C. and again later in Islamic countries. It had appeared that the manufacture of this type of glass ceased between roughly

the seventh century B.C. and the seventh century A.D. Recent analyses indicate that in some regions of Mesopotamia glass of this composition continued to be made throughout this intervening period.

NICOTINE BIOSYNTHESIS

The biosynthesis of nicotine by tobacco roots has been found to proceed in the pyridine ring portion by incorporation of only one carbon atom from various two, three, and four-carbon compounds related either directly or indirectly to the Krebs citric acid cycle. With aliphatic acids such as acetic, succinic, and fumaric, only the carbons alpha to the carboxyl group are appreciably utilized, and with amino acids such as alanine, β -alanine, and aspartic, only the carbon beta to the amino group. Glycerol, however, appears to be incorporated partly via oxidation to acetate and partly as a two-carbon fragment. It is possible that the pyridine ring of nicotinic acid in this system may arise by reaction of citric acid or some close relative thereof (derived via the Krebs cycle from the simpler compounds mentioned above) with a two-carbon fragment from glycerol (such as glycol aldehyde) and ammonium ion from the nitrogen pool of the root system. Degradation studies are being continued to determine the specific location in the pyridine ring of activity derived from each of the important (specifically labeled) precursors used. This will clarify the mode of formation of nicotinic acid in such a plant system.

Nuclear Engineering

On May 1, 1962, Dr. Clarke Williams, who had been Chairman of the Nuclear Engineering Department since its formation in 1952, became Deputy Director of Brookhaven National Laboratory. Dr. Warren E. Winsche, who was one of the original members of the Laboratory staff but had been at Savannah River Laboratory since 1951, became Chairman of the Nuclear Engineering Department on June 25, 1962.

On January 1, 1962, the Radiation Division was formed within the Nuclear Engineering Department. Its three sections have responsibility for studies of the mechanisms of chemical reactions initiated by radiation, studies of selected radiation-induced chemical processes, and the development and characterization of large-scale radiation sources.

The Atomic Energy Commission has designated the facilities at Brookhaven as a Radiation Development Center that will provide a focal point for radiation research and development in the United States. The High Intensity Radiation Development Laboratory (HIRDL) Building has been completed, and special equipment is being installed.

The present program of the Department falls into three main categories. The first, and by far the largest in terms of man-years of effort, is basic research, which includes fundamental studies in physics, chemistry, and metallurgy. Also included are studies on liquid metals, reactor materials at high temperatures, liquid metal heat transfer, and diffusion of fission products in solids.

In the second category is the long-range work of developing components that may bring about significant advances in reactor technology and radiation application. Some of the more important phases of this work are the development of reactor fuels, high temperature volatility and fluidized bed processing schemes, radiation engineering methods, and waste processing devices.

The engineering and evaluation studies in the third category are focused on practical application of the work in the preceding categories. Studies were made during the year on settled particulate

bed fueled systems for breeder, chemonuclear, and high-temperature, high-performance reactors. Through this program assistance is also provided to the AEC in reactor evaluation. Dr. Chad Rase-man has returned to Brookhaven after two years in Washington working for the AEC in this connection. During this period he carried out assignments in 15 European and Latin American countries.

During the year, Department personnel have supervised the construction and operation of gamma irradiators for exhibits sponsored by the US AEC Division of Special Projects in Beirut, Lebanon; Lima, Peru; Mexico City, Mexico; and Athens, Greece. In Lima, the research reactor program associated with the exhibit was also headed by BNL personnel.

Additions to the Critical Assembly Facility and the Metallurgy Building were occupied in the latter part of the fiscal year.

After the Advisory Committee on Reactor Safeguards had reviewed the preliminary hazards summary report for the High Flux Beam Research Reactor (HFBR), field construction work began in the fall of 1961 with implementation of the rough clearing and grading contract. Early in 1962 the architectural and civil contractor for the structural concrete work and the building dome contractor for the steel containment shell began work. As of June 30, 1962, the steel containment shell is $\approx 60\%$ complete, and the structural concrete work is well under way. Practically all items of major equipment are on order, and deliveries are beginning to be made. All field work subcontracts have been placed, the major ones, in addition to those mentioned above, being those for the electrical, mechanical, air-conditioning and heating and ventilation work, and for construction of the cooling tower.

The Brookhaven technical staff has continued close liaison with the AEC in approval matters. Only odds and ends of detailed design remain; emphasis has shifted to operating manuals, spare parts, final hazard reports, operator training, and inspection of work in the field and in vendors' shops.

REACTOR PHYSICS

Theoretical Reactor Physics

An international conference on neutron thermalization was organized and held at Brookhaven from April 30 to May 2, 1962. Approximately 60 papers were presented, including several from the Reactor Physics Division and many from overseas Laboratories. A feature of the conference was the use of the reporter system under which the various papers submitted on a particular subject are covered by one speaker. The impressive progress in both theoretical and experimental work in this field has justified the establishment of strong reactor physics research centers abroad and in the United States.

The importance of theoretical studies of reactor stability and control has been emphasized by the inherent instability found in a number of recent reactors and by the occurrence of several nuclear incidents involving critical assemblies. During the year the study of the influence of xenon and temperature feedback on reactor stability has continued. The importance of the type of control chosen to stabilize inherently unstable reactors was reported in an invited session at the June meeting of the American Nuclear Society in Boston where a review of nonlinear aspects of reactor dynamics was also given.

Review papers on the status of the theory of enriched uranium-water lattices were presented at the IAEA Panel on Light Water Reactor Lattices held at Vienna from May 28 to June 1, 1962. The theoretical results, based strictly on the best available cross sections and rigorous calculations, yield fairly close predictions of the criticality of these systems. One paper was devoted to recent results on the thermal neutron spectrum in uranium-water lattices. The theoretical calculations now yield results of sufficient accuracy to make approximate measurements of the intrinsic properties of these systems no longer useful.

Calculations have been carried out on a number of advanced reactor concepts. Physics studies of settled bed reactors have continued with the emphasis shifting to investigations of the stability of large-scale fast-breeder reactors. Limited studies were completed on fast breeders of the Pu-Bi solution type to determine their feasibility and economic attractiveness. Studies on the "cold moderator" concept of high-conversion-ratio, plutonium-

fueled thermal reactors have continued. With large clusters of variably enriched, plutonium-fueled rods in a cold, internally pressurized, D_2O -moderated core, these reactors have optimum initial conversion ratios of ≈ 1.1 and specific powers of up to 5 Mw (thermal) per kilogram of plutonium. The rapid decrease in the value of η for Pu^{239} puts a premium on low neutron temperatures which are obtainable by slow circulation of the nearly insulated moderator. It is not claimed that true breeding can be achieved in a practical system; however, economical, near-breeding, plutonium-fueled, thermal reactors are considered to have a long-term potential. Small quantities of make-up fuel would be obtained, if necessary, from fast-breeder reactors. These studies also indicate a long-range potential for heavy water as a moderator in power reactors.

Experimental Reactor Physics

Uranium Metal- H_2O Lattices. The program of measurements on the neutron physics of slightly enriched uranium metal rods in light water has ended, but study has continued to resolve some discrepancies. The first area of discrepancy was that of the relative number of U^{235} fissions in the energy regions above and below the cadmium cutoff. By using an improved technique developed at Bettis Atomic Power Laboratory which involves activation of dilute U^{235} foils, a complete set of measurements has been carried out. The results are now consistent and agree with measurements at Bettis. This discrepancy is now considered to be removed.

Studies were performed in connection with the technique of resonance capture measurements. These were inspired by difficulties at other laboratories in measuring believable values of ρ_{28} (the ratio of resonance to thermal U^{238} captures) in uranium oxide-fueled lattices. As a first step toward resolving this question, it was thought necessary to place the measurements in metal lattices on a sound footing. A number of techniques in common use have been refined to a point of high accuracy, and the results now agree. The measurement of ρ_{28} in metal fuel lattices is now believed to be reliable and precise. The values for the many metal lattices studied will now be re-measured, and the techniques will be extended to oxide-fueled systems.

After correction for all known sources of error, there still remains a discrepancy of ≈ 3 to 6% be-

tween measured and calculated capture density disadvantage factors in metal rod lattice cells.

The measured parameters of ≈ 100 metal rod lattices were assembled and presented at the IAEA panel discussion mentioned above.

$\text{UO}_2\text{-H}_2\text{O}$ Lattices. A systematic study has been made of one lattice of 3% enriched UO_2 rods in light water. Macroscopic flux distributions were measured with pure water moderator and with three concentrations of boric acid in the water. The purpose of this study was to identify the source of spurious neutrons which had affected previous measurements and to eliminate this effect. The source was identified as photoneutrons generated in the natural D_2O of the water by gamma rays from the thermal column. Elimination of the problem has permitted a refined set of experiments on the lattice concerned, to provide good values of the buckling, the migration area, the anisotropy of migration area, and the effect of relaxation length on reflector savings. The relation between macroscopic flux distributions and the critical size has been found to be quite complex in these oxide-fueled systems. The anisotropy is found to be considerably greater than in previously studied metal-fueled lattices.

With the resolution of the problems underlying these exponential experiments and clarification of the way they must be analyzed, the experiments on the remaining assemblies should now proceed quickly.

Slab Lattice Experiments. Most of the required measurements in this program have been completed.

$\text{U}^{233}\text{O}_2\text{-ThO}_2$ Experiments. Discussions and correspondence with personnel at Oak Ridge National Laboratory have resolved all known uncertainties in specifications for the $\text{U}^{233}\text{O}_2\text{-ThO}_2$ fuel, and fabrication at Oak Ridge will begin soon. First delivery of the fuel is scheduled for December 1962.

Minimum Reflection Critical Experiments. These experiments are performed on a platform ≈ 20 ft above ground level in an unshielded silo building with thin steel walls (Figure 1). The objective is twofold: to obtain good experimental data on the critical mass of homogeneous, unreflected graphite-uranium reactor cores over a wide range of atom ratios, and to test experimentally the so-called second fundamental theorem of reactor physics.

After final approval of safety considerations by the Brookhaven Reactor and Critical Experiments

Safety Committee and the AEC, critical loading of the first assembly was achieved on April 25, 1962.

Measurements have been made of the safety rod worth and of the change in reactivity with thickness of the air gap between partial assemblies on two carts. The reactivity meter described in the 1960 annual report was used in some cases, and measurements of decay time were made with a pulsed neutron source. The latter method is more reliable in measuring the change of reactivity with gap thickness. The measurements confirm reasonably well a theoretical prediction made by Chernick in connection with the design of the Brookhaven Graphite Research Reactor (BGRR).

Measurements of flux distributions are now under way. These tentatively indicate a small amount of neutron reflection from the carts for which correction must be made in analyzing the data and drawing conclusions about flux separability.

Pulsed Neutron Measurements. The Van de Graaff generator was given a complete overhaul,

Figure 1. Minimum Reflection Critical Experiment.

and improvements were made in the pulsing system. Greater sharpness of pulse shaping and a considerable increase in signal to background neutron intensities (now $\approx 10^7$) have been achieved. With the smaller background, measurements over a much greater range of decay are possible, and some problems obscuring the understanding of neutron thermalization in graphite have been solved. The time required for effective neutron energy equilibration in graphite was found to be much longer than expected, ≈ 2 msec, and not very dependent on stack size. Measurements at all laboratories prior to this equilibration had contained data that led to incorrect values of the diffusion coefficient, absorption cross section, diffusion length, and diffusion cooling coefficient. The measurements made after the longer waiting times are consistent and have now been confirmed by measurements at other laboratories.

In an extension of the pulsed neutron techniques, diffusion length measurements were made in light water poisoned by different amounts of boric acid. The data obtained could be analyzed with use of the same assumptions as those used in pulsed neutron experiments. They led to results that agreed with the values of diffusion lengths, absorption cross section, diffusion constant, and diffusion cooling constant found in the best pulsed experiments, but the new measurements have improved the accuracy.

The Van de Graaff accelerator has been dismantled for transfer to the new addition to the Critical Assembly Facility.

High Flux Beam Research Reactor. The critical assembly mock-up of the HFBR has been reinstalled in a spherical vessel of the shape and size to be used in the reactor. This vessel has permanently installed beam tubes.

The experimentally based estimate of the core size needed to furnish the requisite excess reactivity was found to be correct. Measurements of the thermal neutron flux from beam tubes confirmed earlier measurements made with movable tubes in the cylindrical vessel. The ability of safety rods to hold down the peak excess reactivity encountered during fuel changes was also confirmed.

Measurements of the gamma heat rates in the vessel wall were made with a mocked-up thermal shield in place.

Cold Neutron Project. A slow-neutron chopper has been assembled and is being checked out on

the 4×4-ft graphite thermal column of the Neutron Source Reactor.

A small cryostat has been designed and is ready for fabrication.

A series of runs was made using a 1.65-m flight path. Intensity was adequate even at the Bragg cutoff for bismuth. The variation of count rate with rotor angle was determined and found to be triangular.

The best of all data obtained so far show that the spectrum emerging from the face of the thermal column of the Neutron Source Reactor corresponds to a 300°K Maxwellian distribution perturbed by the variations in the graphite cross section.

Reactor Safety. Emphasis by the AEC on more formal reactor safety procedures has increased greatly since the SL-1 incident. A conservative estimate indicates that at least two scientist man-years have been devoted by the Experimental Reactor Physics Group to Brookhaven reactor safety matters in fiscal 1962.

Sigma Center

Neutron Cross Section Compilation Group. A second and completely new edition of BNL-400, *Neutron Cross Sections – Angular Distributions*, is being readied for publication in 1962.

A digital computing machine program has been prepared to duplicate on magnetic tape most of the neutron cross section information in the Sigma Center's files. Such duplication should make access to the files more rapid and flexible than heretofore. Automatic plotting of selected sets of the data should also be possible. Activation measurements of thermal (n, γ) cross sections of Al^{27} , V^{51} , Ag^{107} , and Ag^{109} were made to resolve discrepancies in the previously published values of these cross sections.

The Group cooperated closely with the Nuclear Cross Sections Advisory Group of the AEC by furnishing secretarial service and preparing its reports for publication.

Reactor Cross Section Evaluation Group. Newsletters No. 5 and No. 6 were issued (BNL 694 and BNL 732) covering recently received neutron cross section data of particular interest to reactor physicists.

A least-squares analysis was made of the many interconnected measurements of the thermal neutron cross sections and other parameters of the

three fissile nuclides, U^{233} , U^{235} , and Pu^{239} . The results constitute a refinement of the World Consistent Set of 2200-m/sec parameters published in the last (1960) Supplement to BNL 325 - *Neutron Cross Sections*. They appear in BNL 722.

The digital machine program, ABACUS, for computing cross sections according to the optical model of the nucleus has been improved and extended. Transmission coefficients for all nuclei have been calculated and tabulated in a manner convenient for calculation of nuclear cross sections. The tabulation will be made generally available in report form. Inelastic neutron scattering cross sections for U^{238} have been computed by the Hauser-Feshbach method for various possible choices of energy level schemes, and the quite-good fit to experiment has been scrutinized for evidence that a spherically symmetric potential well may not represent the actual U^{238} nucleus. The optical model has also been applied with some success to the scattering of slow electrons by argon atoms. This application to a nonnuclear problem demonstrates the model's range of applicability.

Some theoretical understanding of the observed tendency for nuclear energy levels to shun or repel each other has been obtained by showing that a similar behavior characterizes the eigenvalues of certain randomly chosen quantum systems.

Reactor Evaluation and Advanced Design

Settled Bed Reactor Concept. A major effort during the past year in the settled bed reactor program has centered on the directly cooled fast reactor systems. This reactor concept involves a settled bed of spherical fuel particles directly cooled by a liquid metal flowing in either an axial or a radial direction through the bed. The reference fuel is UPuC. Fuel development, economics, and reactor physics comprised the main areas of evaluation.

Initial fuel development work is being performed with UC. The fuel development program has included procurement of UC spheres, capsule design and testing in an out-of-pile high-temperature liquid metal environment, and fabrication and testing of an in-pile capsule.

Organic Cooled Reactor Studies Brookhaven was designated as the information-coordinating AEC laboratory in the organic cooled reactor program. In addition, a broad experimental program

has been started to seek an understanding of (1) the mechanisms of fouling film formation on heated metal surfaces, (2) corrosion of metals, and (3) the thermal and radiation stability of various organic coolants.

Direct Conversion. A study has been started on the direct electrical conversion of energy from a fissionable ionized gas, UF_6 , at 3000 to 15,000°K. Ultimately 20 to 30% direct conversion may be achieved, with utilization of 35% of the remaining waste heat via a conventional steam cycle. Electrodeless electromagnetic coupling is used to extract the energy, either by a pulsed subsonic axial expansion of the heated gas into a cylindrical magnetic piston, or by a quasi-steady subsonic flow of gas between two reactors through varying magnetic fields. The alternating gas eddy currents generate a voltage in an external pickup coil. The shock technique proposed by S. Colgate is not considered because of the high losses anticipated from turbulence. The coupled nonlinear energy, momentum, Maxwell's, matter, and state equations have been derived. An analytic solution for a linear perturbation showed low efficiencies of several percent; for high efficiencies operation must be nonlinear. A numerical computer solution is needed and will be sought. A multigroup neutron diffusion analysis has shown critical concentrations of $\approx 2 \times 10^8$ molecules/cc, which correspond to plasma pressures of ≈ 1 atmosphere. Containment seems feasible with fuel stabilized by excess fluorine.

On the subject of a thermally regenerated fuel cell, analytical calculations verified by experimental measurements showed that the aqueous concentration cell was power-limited at 200 w/ft³. The analysis also provided a basis for evaluating compounds as to their potential merit. Accordingly, it was decided to examine systems that indicated capability for higher electric power density. The program is also directed toward a system without moving parts.

Chemonuclear Studies. Design and evaluation work on two new chemonuclear systems was undertaken. Both reactors are being designed for the production of CO in a CO₂ process stream by utilizing fission fragment recoil energy. Particular emphasis is being placed on an externally moderated system capable of producing 700 Mw (thermal) with a chemical output of ≈ 1200 tons of CO per day. The second concept features a two-

region core, the inner region designed for heat generation alone, and the outer, annular region for chemical production.

Fuel Cycle Analysis. Programming of two computer codes was begun. PSYCHO is a generalized fuel-cycle cost code which closely follows the method presented in the *AEC Cost Evaluation Handbook*. THORCYC is a one-dimensional multigroup-diffusion-theory fuel-cycle analysis code for the analysis of Th-U²³³ fast-breeder systems.

During the period various evaluation projects were undertaken for the AEC.

METALLURGY

Reactor and General Metallurgy

The reactor metallurgy program is concerned with the problems of containment of liquid metal coolants used in the power conversion systems of compact reactor systems, the production of controlled composition alloys for research, and general consultation on materials throughout the Laboratory.

Materials for Containing Boiling Liquid Metals.

Boiling liquid metals, especially Hg and Na, are currently of interest as working fluids in high-performance nuclear reactor-turbine systems. Three all-liquid and six boiling Hg-containing loops have been run to determine the effectiveness of Ti and Zr as inhibitors of Hg corrosion in systems fabricated of ferrous alloys. In addition, a loop made of Haynes Alloy #25 has been run using Ti-inhibited Hg, and two loops built of refractory metal are now in operation.

The loops in which only liquid was circulated were fabricated from tubing having carbon steel on the inner surface and 1.25 wt % Cr-0.5 wt % Mo steel on the outer surface for strength and air oxidation resistance. Each loop was run for ≈ 8000 hr, with a temperature difference of 204°C (528° to 316°C). The fluid in one loop was uninhibited, while the other two loops contained either Ti or Zr dissolved in the Hg. The Hg from the loops was filtered to remove corrosion products at the end of the test. On the basis of the weighed corrosion products, the Ti inhibitor reduces corrosion by a factor of 3, while the Zr inhibitor reduces it by a factor of 8. Metallographic examination shows the mode of attack in the Ti-inhibited and the uninhibited systems to be of a grain boundary type,

and that in the Zr-inhibited system to be of a transgranular nature.

The seven boiling Hg loops were operated at 538° to 593°C boiler and 607° to 704°C superheater temperatures. Materials tested were Inconel-clad carbon steel, 2.25 wt % Cr-1.0 wt % Mo steel, and Haynes Alloy #25. The Inconel-clad carbon steel and Haynes #25 loops were run with a boiler temperature of 593°C. Effectiveness of the Zr as an inhibitor was essentially the same in the boiling loops as in the all-liquid system loop. The Ti did not appear to have an inhibiting effect in the boiling system. Of the boiling loops, the Haynes Alloy #25 loop produced the greatest amount of transferred material and exhibited a leaching type of corrosion rather than the grain boundary or transgranular attack observed in the ferrous materials. No corrosion was detected in the boiler section of this loop, which indicates that the Ti perhaps has some inhibiting effect on the Haynes #25 alloy.

Boiling in these small thermal convection loops is rather unstable, which makes it difficult fully to understand the action of the inhibitors. In an attempt to study the movement of inhibitors in a boiling system, a loop has been built which will have a radioactive tracer dissolved in the Hg. By scanning with a counting device, it should be possible to follow the movement of the soluble constituent in the boiling loop system.

The extent of inhibition taking place in the condensing portions of a boiling system is unknown, but it is possible that adequate protection may be achieved by forming a protective layer prior to operating the loop at design conditions. Capsule experiments have been run to determine the time and temperature required to form an inhibiting film. Electron transmission studies on films stripped from the capsule specimens have shown the films to be ZrN when Zr was used as the inhibitor.

A series of reflux-type capsule tests is being run at 1204°C to determine the corrosion resistance of Cb-1 wt % Zr alloy to Na containing various oxygen additions and getters such as Ba and Mg. The resistance of Mo, Ta, and Cb to purified Na is also being determined in reflux capsule tests. A small thermal-convection boiling loop built of Cb-1 wt % Zr and containing purified Na is now operating at 1093°C boiler and 1204°C superheater temperatures.

Two high-vacuum chambers 4 ft in diameter and 8 ft high have been put into operation. They will contain a forced-circulation boiling Na loop and a similar Hg loop. Both loops will be built of Cb-1 wt % Zr alloy and thus will require protection from the atmosphere. A vacuum of 10^{-8} mm Hg can be attained in the chambers by using a Vac-Ion pump.

General Metallurgical Studies. A series of iron alloys made from high purity materials has been prepared in the molecular-still vacuum induction furnace. The 1-lb melts contain up to 0.30 wt % carbon and up to 1 wt % Ni or Cr. The alloys are then cold-swaged to 0.230-in. diameter for use in the radiation effects program.

Ta-Fe, U-Pt, Cu-B¹⁰, and Cu-B¹¹ alloys have been prepared by inert gas arc melting for various groups at the Laboratory. The Cu-B¹⁰ alloy (9 atom % B¹⁰), prepared for the Cross Section Group, was of sufficient uniformity to allow the determination of the predominant spin state of the B¹⁰.

HFBR Metallurgy. Aluminum alloy 6061 has been chosen as the construction material for the HFBR containment vessel. Since experience in the welding of this material in thick sections is limited, a series of tests was run to determine the suitability of alloys C809 and 4043 as weld filler materials. Test welds were made by using the manual consumable inert arc process. Mechanical testing showed the joints to have adequate strength even in the as-welded condition, and x-ray examination revealed only small amounts of porosity. Although metallographic examination revealed considerable microporosity, no cracks, tears, laminations, or inclusions were found.

Three prototype fuel elements for the HFBR have been irradiated in the Engineering Test Reactor (ETR). They contained 30, 35, and 40 wt % U²³⁵ in Al. They were mechanically assembled by a "peened-lug" method. The first two of these elements received 40 to 50% burn-up. Hot cell examination of the first element showed severe plate distortion with several channels reduced in some areas to $< \frac{1}{4}$ their original spacings. The third element received 16% burn-up. All channels in the 35 and 40 wt % elements were locally restricted to $< 75\%$ but $> 50\%$ of their original value. New tests are under way of elements fabricated by the roll-swaged method of assembly. These all contain 35 wt % U²³⁵ in Al in the fuel-bearing core. Four will be irradiated in the ETR and two in the Ma-

terials Testing Reactor (MTR) to 20 to 25% burn-up.

Irradiation of Dy₂O₃-stainless steel compacts in the MTR has been initiated. One compact containing 30 wt % Dy₂O₃ has been irradiated for three MTR cycles without noticeable damage.

Alloy Theory and Nature of Solids

Structure of Liquid and Solid Alloys. Further evaluation of the heats of solution of solute metals in liquid Bi suggests that the electronic configuration is more important than the size factor in determining the thermodynamics of dissolution. A generalized theory for metallic structures combining the molecular, orbital, and valence bond approaches was formulated. Calculations for simple molecules are in agreement with experiment.

Liquid Metal Corrosion Fundamentals. 1. *Emf effects.* An attempt is being made to determine the effects of a dc current on the diffusion of various metals in molten Bi. Experiments were performed in which irradiated Fe, Cr, and Ag diffused through capillaries containing molten Bi at 500°C while currents of 0 to 1000 amp/cm² were passed through the Bi. Fe and Ag show accelerated diffusion against the electron flow and tend to concentrate about the cathode, whereas Cr shows accelerated diffusion in the same direction as the electron flow and tends to concentrate about the anode. Preliminary calculations suggest diffusion coefficients of the order of 10^{-5} cm²/sec with a twofold increase in mobility of Ag against an electron flow of 1000 amp/cm² during the electrolysis of Bi-Ag alloys.

Measurements of the W-liquid Bi thermocouple have shown that the absolute emf of liquid Bi is not affected by Fe additions. Measurements of the emf between Fe and W electrodes in Bi have shown the potential of the iron electrode to be sluggish in establishing its steady-state value. This may be due to an effect of dissolved iron on the electrode potential (concentration cell). Currents > 25 ma can be drawn from a cell with a 5 wt % Cr-0.5 wt % Mo steel anode and a carbon steel cathode immersed in liquid Bi at 500°C. The current flow continues steadily for at least 3 hr without polarization. The polarity of this cell (compared with the electrodiffusion measurements) suggests selective attack of the 5 wt % Cr-0.5 wt % Mo steel electrode, in agreement with loop observations.

2. Surface reaction studies. An attempt was made to measure the kinetics of formation of a Zr-bearing film on a 2.25 wt % Cr-1 wt % Mo steel at 700°C during the first 4 hr of reaction by using an improved thin liquid film technique. It was found that the precision would have to be further improved for meaningful results. Analysis by error theory suggested that a potential coefficient of variation of 10% should be attainable through more precise control of the radioactive Zr tracer added. With this coefficient it should be possible to distinguish a parabolic from a cubic film growth curve.

3. Dissolution kinetics. One method of estimating the activation energy of interaction between a liquid metal and a dissolving metal is to measure the temperature dependence of the rate constant of the dissolution process. It was concluded from a literature survey that a pumped tube system would provide the best estimate of the thickness of the boundary layer, and that a rotating concentric cylinder system was next best.

Thin Film Studies. Fission events in thin, vapor-deposited metallic films produce distinct tracks, which are observable by electron microscopy. Alpha and tritium particles generated by the $\text{Li}^6(n, \alpha)\text{H}^3$ reaction have also produced tracks in 100-A-thick Li^6F films. The appearance of the tracks in both cases is strongly influenced by the thickness of the matrix material. Track characteristics in metal films are also influenced by the atomic number of the film. It has been tentatively concluded that the thermal spike mechanism is mainly responsible for the damage observed in metal films. In LiF films, both the alpha and the H^3 particles are capable of raising the temperature in excess of the melting point, since their total energies are 2.04 and 2.84 Mev respectively. This corresponds to an energy loss rate of 60 to 120 ev/A in the case of the alpha particles and 15 to 30 ev/A for the H^3 particle. The mode of energy transfer in this case is thought to be mainly by electron excitation.

Radiation Effects

Effects of Neutron Irradiation on the Mechanical Properties of Body-Centered-Cubic Metals. The effects of material selection and neutron exposure on the properties of body-centered-cubic (bcc) metals and of pure iron in particular are being studied by measurement of the mechanical prop-

erties at ambient to liquid He temperatures. The changes observed in the yielding, plastic flow, and fracture properties after fast neutron exposures from 10^{17} to 10^{19} nvt have been studied in pure iron and iron alloys as a function of the grain size, integrated exposure, testing temperature, and postirradiation annealing temperatures. Changes in the microstructure, electrical resistivity, and hardness were also analyzed.

The results obtained during the past year show that the yield stress becomes independent of the grain size after an exposure of $\approx 10^{18}$ nvt and that the plastic behavior after irradiation is similar to that of prestrained iron. Quantitatively, the yield stress of iron after fast ($E > 1.45$ Mev) neutron exposures of 2×10^{18} nvt and greater can be described by the relation

$$\sigma_{ys} = \sigma_0 \ln 1/KT,$$

where K is a function of the integrated exposure and σ_0 is a material constant. Also, the linear stress-strain curves obtained after the above exposures follow the relation

$$\sigma_e = \sigma_{ys} + \bar{K}\epsilon,$$

where $\bar{K}$ is a function of the temperature only and ϵ is the natural strain.

Exposures of 10^{18} nvt also affect the brittle to ductile transition temperature markedly, raising it by as much as 60°C. The ductility as measured by the reduction in area is lowered from 10% to 2% below the transition temperature.

A proposed explanation is that at low irradiation exposures the increase in yield strength is the result of the interaction of moving dislocations with point defects. Above a threshold of 10^{18} nvt clustering or condensation of defects occurs, and these more complex defects are capable of forming a cellular substructure which tangles with moving dislocations. This substructure has a more profound effect on the yield strength and is capable of decreasing the ductility and raising the ductile to brittle transition temperature. A 2-hr postirradiation anneal at 350°C was required to remove the radiation-induced embrittlement.

Electron-Microscopic Study of the Effects of Neutron Irradiation on the Properties of Iron and Other BCC Metals. A program has also been started to study the effects of fast neutron irradiation on the properties of iron and other bcc metals using electron transmission microscopy. Although it is un-

likely that structural defects produced by neutron irradiation can be observed directly, it is anticipated that the effects of such defects upon the behavior of dislocations during plastic deformation can be observed. Correlation of such observations with mechanical and other properties should shed light on the nature of the defects and the mechanism by which they affect the properties of bcc metals.

Irradiation Effects in Graphite. The objectives of the graphite program at Brookhaven are to understand radiation effects in graphite and to determine irradiation and annealing conditions that will minimize detrimental effects on the physical, chemical, mechanical, thermal, and electrical properties of graphite.

1. Brookhaven Graphite Research Reactor studies. In the monitoring program on the BGRR, measurements of dimensional changes, stored energy, and lattice parameters were obtained in graphite that have been extremely useful in developing a model to explain many of the observed effects of neutron bombardment on continuously irradiated and periodically irradiated and annealed graphites.

The rate of growth of graphite during neutron irradiation and its dimensional recovery after a reactor annealing operation can now be predicted. Heating (annealing) of the BGRR graphite to 150°C with greater frequency and for shorter soak times (length of time at annealing temperature) will be started soon. The revised procedure should result in increased dimensional recoveries and removal of damage already present. Frequent annealing also prevents the stored energy released at 200°C from exceeding 20 to 30 cal/g even after total fast ($E > 0.6$ Mev) neutron exposures of 10^{19} nvt.

2. Fundamental studies. The activation energy for the migration of single interstitials in neutron-irradiated graphite and the absolute rate of formation of displaced atoms have been experimentally determined from dimensional and c-axis expansions of graphite irradiated at temperatures between 77° and 490°K. The value of the activation energy E lies between 0.068 and 0.074 ev and agrees well with estimates and theoretical calculations of other investigators. The absolute rate of formation of displaced atoms (when no annealing occurs during the irradiation process) was found to be $5 \times 10^{-3}\%$ displaced atoms per 7×10^{16} nvt and is 2.5 times the resultant rate of displaced atoms at 30°C.

Periodically irradiated and annealed graphites show property changes markedly different from those produced in the first irradiation and anneal. Evidence has been obtained indicating the existence of consecutive chain reactions between interstitial complexes that lead to steady-state concentrations of the intermediate species. These concentrations result in dimensional recoveries and stored energy releases that do not increase with re-exposure as they do in continuously irradiated graphites.

Interstitial complexes formed by thermal annealing interact through chains via mobile single interstitials. Kinetic equations for chains open at both ends show that large complexes can decompose to form smaller ones. Experimental evidence supporting the calculations shows that periodic removal of the smaller complexes by low temperature annealing results in the net removal of large amounts of damage associated with the larger interstitials.

3. Kinetics of graphite-bromine reactions. A kinetic study of the reactions between graphite and bromine has been undertaken. Bromine and a number of other substances are able to penetrate between the layers of the graphite structure and form compounds of variable composition. Study of the graphite-bromine compounds has proved to be of importance in elucidating the electrical properties of graphite and in increasing the understanding of irradiation damage in graphite.

The data so far obtained indicate that the exchange of normal and radioactive bromine in graphite-bromine compounds is a diffusion-controlled process. The data also indicate that the rate of bromination of graphite is determined by the rate at which the bromine-graphite compound is able to penetrate the graphite lattice. Further efforts are being focused on obtaining quantitative information on the rate constants and activation energies involved in the rate processes.

4. Electrical properties of irradiated graphite. A program has been undertaken to study the changes occurring in the electrical properties of graphite when it is irradiated and periodically annealed. Apparatus has been constructed for the measurement of the electrical resistivity, Hall coefficient, and magnetoresistance of graphite. Preliminary data indicate that periodic annealing can be very effective in limiting the amount of electrical damage incurred in irradiated graphite.

5. Oxidation and heat transfer measurements in graphite channels. The effects of convection currents on the temperature distribution in preheated 10-ft-long graphite channels were investigated. Above 700°C, relatively rapid temperature increases occur in graphite channels due to thermal convection of air. At 650°C or lower, the heating rates are not large and the heat produced does not result in a safety problem if the graphite is cooled at a rate faster than 10°C/hr either by conduction, convection, or forced convection of air.

Xenon Diffusion in Carbides. Experiments to determine the diffusion of Xe^{133} in uranium carbide have been performed on four uranium carbide powder preparations and one uranium dicarbide powder.

The xenon release kinetics, to a large extent, have been observed to follow a diffusion mechanism, with an activation energy for diffusion of $\approx 85,000$ kcal/mole in the temperature range 1000° to 1400°C. In this range the diffusion coefficient can be represented by the equation $D = 2.1 \times 10^{-5} e^{-85,000/RT}$.

At low temperatures (up to $\approx 850^\circ\text{C}$) a considerable fraction of the gas released is released in short bursts of <1-hr duration. A "pressure burst" usually accompanies an "activity burst" when the freshly irradiated powders are heated for the first time at low temperatures. In addition, the continuing emanation of activity at these temperatures is much more rapid than that predicted by extrapolation of the high temperature results, and does not fit nearly so well the simple diffusion kinetics observed for most of the high temperature work.

CHEMISTRY AND CHEMICAL TECHNOLOGY

Reactor Chemistry

Thermodynamics of High Temperature Systems.

Much useful information on the thermodynamic properties of materials at high temperatures can be obtained from measurements on suitably designed galvanic cells. A novel type of cell was developed for systems in which a simple cell will not work because of the solubility of metal in the salt. The new cell design is illustrated by the following example:

Two electrolytes are used, a pure salt and a mixture, separated by a membrane that is imperme-

able to liquid but is an ionic conductor, the current-carrier being the cation common to the two electrolytes. The net cell reaction is the transfer of NaCl from the left-hand electrolyte compartment to that on the right, and the emf is directly proportional to the partial molar free energy of NaCl in the particular mixture used, on the right.

This cell has been tested with a number of different salt compositions at 820°C. The results obtained were reproducible and self-consistent, and in agreement with earlier data within the latter's limits of error, which are probably greater than those of the present method. The new type of cell should be applicable to a large number of systems, and further work of this nature is planned.

For determining free energies of formation, it is often convenient to use cells in which all the components, both electrodes and electrolytes, are solids. Each component has the form of a flat-surfaced tablet, and a cell consists simply of a series of such tablets pressed together. Most salts can be easily made into tablets by compressing the powdered material at a few thousand psi. The pressed products usually have sufficient conductivity at 400° to 500°C to permit the observation of emf on cells of which they are components. A typical cell measured in the past year's program was the following:

Here "Ce, CeCl₃" represents an intimate mixture of the powdered materials. The NaCl and AgCl form a solid solution in which the activity of AgCl is known from previous published work. The cell reaction involves the formation of CeCl₃, and, since all the other factors contributing to the observed emf are known, the free energy of formation of CeCl₃ can be obtained. This type of cell has also yielded precise information on the free energies of MgCl₂, UCl₃, and ThCl₄.

Spectroscopy of Fused Salts. One of the most powerful methods for investigating the structure of fused salts and following reactions in salt media is that of absorption spectroscopy. A large number of fused salt systems have been examined by other workers in the visible and infrared regions of the spectrum. In the ultraviolet, however, the extremely high absorption of most salts has hitherto prevented any detailed resolution. To open this region to exploration, a cell with a light path of $\approx 0.1 \mu$ has been developed.

The spectra of several pure salts and mixtures have been obtained with such cells, both in the liquid state and over a wide temperature range in the solid. The most interesting results were those observed on the solids. In certain cases as the temperature changed the maximum of an absorption band shifted toward longer wavelengths. Such shifts were observed with both PbCl_2 and KI for the first band encountered on going from the transparent visible region of the spectrum into the ultraviolet. Figure 2 shows several curves for PbCl_2 . This first band is often attributed to the formation of an exciton, or the transfer of an electron from an anion to a cation. It is not always possible to establish the position of its maximum precisely, but the location was clear in the spectra of PbCl_2 and KI up to the melting points (498° and 685°C , respectively). A remarkable correlation was found between the temperature-induced shifts and the enthalpy increases of the compounds over the same temperature interval. In the case of KI the energy decrement corresponding to the frequency change was exactly equal to the enthalpy increment as calculated from the heat capacity. For PbCl_2 , it equaled $\frac{1}{2}$ of the enthalpy increment. The meaning of the correlation is still obscure, but

Figure 2. Ultraviolet spectrum of PbCl_2 at various temperatures.

Table 1
 G Values for Electron Irradiation of Fluorocarbons

Compound	Irradiation temperature	G_{gas}	G_{polymer}
Hexafluorobenzene	Room	0.001	2.15
Perfluoro- <i>n</i> -heptane	Room	0.85	1.21
Perfluorodecalin	Room	0.12	1.40
Perfluorobiphenyl	100°C	≈ 0.0001	1.39

the factor of $\frac{1}{2}$ may be connected with the fact that the heat capacity involves both chloride ions in PbCl_2 , whereas the charge transfer occurs with only one chloride ion at a time.

Fluorocarbon Program. The outstanding thermal and chemical stabilities of fluorocarbons make this class of compound interesting to both the physical chemist and the chemical engineer. A program is under way to determine whether these compounds also are resistant to attack by ionizing radiation.

A number of fluorocarbons have been irradiated with 1.5-Mev Van de Graaff electrons after careful purification by preparative-scale gas-liquid chromatography. G values given in Table 1 indicate that at or near room temperature the aromatic compounds are less stable to irradiation than the corresponding hydrocarbons, whereas the aliphatic and alicyclic compounds are more stable than their hydrocarbon analogues.

Purification of starting materials by preparative-scale gas-liquid chromatography has been a difficult and laborious process, particularly with respect to compounds with high boiling points. To avoid contamination by material bleeding off the columns, work must be done at temperatures much below the boiling points of the compounds being purified. This necessitates, for an adequate degree of separation, use of relatively small samples, e.g., 0.3 g compared with 1.5 g or more for liquids with low boiling points, such as C_6F_6 .

Another factor that greatly lengthens the purification procedure is the need to use two columns for some compounds, e.g., perfluoronaphthalene. As shown by the chromatograms in Figure 3, silicone oil #702 is the only stationary phase that allows both main impurities to be observed. However, resolution is incomplete with this column, and two different stationary phases are required, one for removal of each impurity.

Figure 3. Chromatographic behavior of commercially available perfluoronaphthalene with different stationary phases. The impurity peaks are gradually shifted relative to perfluoronaphthalene from the situation in which peak A is well resolved and B entirely hidden to that in which A is entirely hidden and B resolved.

Radiochemical Analysis Development

Research and development in the area of radiochemical analysis include the following items:

The study of the enhanced extraction of thorium by thenoyltrifluoroacetone (HT) in the presence of tri-*n*-octylamine (R_3N) has been completed and the behavior explained by five equations, the significant one being

A similar study begun on the extraction of uranium by these reagents has shown that R_3N enormously enhances the extraction of U by HT. The postulated reaction

has been tentatively verified, and in the absence of HT the reaction

has been firmly established.

Controlled-potential electrochemical studies in the molten LiCl-KCl eutectic include anodization to determine nickel which had been predeposited from the eutectic onto a platinum gauze electrode, study of oxidation-reduction mechanisms such as the dissolution of zirconium, use of coulometry for determining dilute cadmium and zinc, measurement of the standard potentials of Rh^{+++}/Rh and Ir^{+++}/Ir couples, and measurement of the activity coefficient of dilute cadmium in molten bismuth.

Electrochemical studies in molten nitrate eutectics have shown that in the LiCl-KCl system lead can be studied quantitatively at the dropping mercury electrode and that previous difficulties with controlled-potential coulometry in this medium were due to oxidation of insufficiently noble metal amalgams by the nitrate melt. When potentiometric studies in the $NaNO_3$ - KNO_3 system indicated that Cu^{++} is too acidic to form stable solutions in this melt, experiments to measure indirectly the Cu^{++}/Cu potential were begun; however, difficulties attributable to the irreproducible nature of the surface of the copper sheet electrodes used were encountered.

In electrochemical studies on molten bromides and iodides, at least three adequate solvents were found for a number of metal ion-metal systems: 1-methyl pyridinium bromide, the LiBr-KBr eutectic, and a 46 M % NaI-KI mixture. The oxidation-reduction potentials for many elements in these systems were measured.

The measured abundances of gammas from Te^{129m} and Te^{129} have been found to disagree with those reported in the literature, and the decay scheme of these two isotopes is being investigated.

Zirconia was found to have excellent absorption characteristics for tellurium and molybdenum from HNO_3 solutions, and tellurium in $\leq 2 N$ HNO_3 was found to be much more strongly adsorbed than on chromatographic alumina.

Small amounts of hafnium can now be separated rapidly from large amounts of tantalum via a methyl isobutyl ketone extraction of a tantalum fluoride complex.

A procedure involving a helium sweep followed by direct counting of Xe^{133} adsorbed on charcoal was found to be satisfactory for studying the diffusion of xenon from irradiated uranium carbide.

Several procedures for determining trace quantities of elements have been developed. Chemical concentration of manganese from blood plasma, followed by arcing small samples at high currents, now permits detection of 3×10^{-10} g manganese via emission spectrography. A few ppm aluminum in steel can be determined by complexing with an organic dye followed by incremental polarography of the complex. X-ray fluorescence coupled with spectrophotometry has been used to determine traces of phosphorus. Gamma-gamma coincidence counting has been used on samples of irradiated diamond dust to accomplish what would have been more difficult and time-consuming without this technique, to wit, establishing the simultaneous presence of traces of Co^{60} and Fe^{59} and establishing the presence of traces of Sc^{46} in the midst of larger amounts of Zr^{95} which would otherwise have masked the Sc^{46} . Neutron activation has been used successfully to determine traces of iodine in dog food.

Radiation Chemistry Research

Radiation Effects in Organic Solids. The determination of the detailed crystal structure of acrylamide is essentially completed, and the structure of propionamide, butyramide, valeramide, and other homologues is being determined.

The data on postirradiation polymerization of dilute solid solutions ($<10\%$) of acrylamide in propionamide indicate that the polymer assumes a random configuration at a very low degree of polymerization and nucleates as a separate phase at very low total conversion. Although diffusion of acrylamide through the intact crystal can hardly be expected, diffusion along defects or the crystal-polymer interface may explain these results.

The radiolytic evolution of hydrogen from pure propionamide and pure acrylamide at room temperature indicates the greater radiation sensitivity of the former.

The second-order rate constant for radical decay in irradiated propionamide has been determined and found to be independent of total dose.

Polymerization Mechanisms. A technique for detecting and measuring the ions present in a con-

ventional ionic polymerization (stannic chloride-initiated styrene) has been developed. It consists of measuring at various frequencies the dielectric loss of the polymerizing mixture as a function of time, at varying concentrations of initiator and water. The data show that a steady-state ion concentration is attained early in the reaction. The loss in this state, measured as a function of frequency, is found to be due to conducting ions, which can be shown to be definitely associated with a monomer-initiator complex. The technique is now being applied to a system in which the detailed kinetics are known, so that quantitative data may be obtained.

The radiation polymerization of ultra-pure alpha-methylstyrene has proceeded, but at a very reduced rate. An attempt is being made to establish the base-line kinetic data in the pure water-free monomer in order to study the effects of various additives.

Radiation Chemistry of Aromatics. Liquid toluene has been irradiated, generally at room temperature, at total doses ranging from 5×10^5 to 5×10^8 rad. A search for dimeric products not previously identified is in progress. When this is complete, G values for the major products will be determined as a function of irradiation temperature. Equipment is now being calibrated to permit determination of the G values of the gaseous products. A flame ionization detector attachment for the vapor phase chromatograph was acquired and is being made operable. This detector will permit investigations of radiolysis products from relatively low dose irradiations.

The G values for hydrogen formation in the Co^{60} gamma radiolysis of *ortho*, *para*, and *meta*-xylenes and ethylbenzene decrease in the order given from 0.24 to 0.16 and are unaffected by the addition of iodine as a radical scavenger. The yield ($G=0.026$) of methane from ethylbenzene is approximately twice that from the xylene isomers, and for each of the compounds the value of G_{CH_4} is reduced by 80% on the addition of iodine. In the case of *p*-xylene, oxygen was found to have the same effect on the gas yields as iodine.

Radiolysis of *p*-xylene at various temperatures showed that G_{H_2} and G_{CH_4} increased slowly from -196° to 0°C , but above the melting point (13°C) the yields were considerably higher, G_{H_2} being independent of temperature up to 110°C , while the production of methane increased with

temperature. Methane yields at high temperatures decreased as a function of radiation dose.

Variation of dose rate from 8.5×10^{16} to 1.3×10^{17} $\text{ev g}^{-1} \text{sec}^{-1}$ caused no change in the G values for hydrogen and methane production from p -xylene irradiated at 20°C .

Isotope Development

The following is a partial list of accomplishments in the development of ways of making available potentially useful isotopes.

Aluminum-28 Generator. A very satisfactory generator for 2.30-min Al^{28} has been developed. It yields a 99.999% pure product in 30 sec and can be "milked" dozens of times per hour for several days.

Argon-38. This stable isotope, of interest for calibrating mass spectrometers used in geological dating experiments, is produced by reactor irradiation of Cl^{37} . The use of pure NaCl as a target produces very high purity ($>99.4\%$) Ar^{38} entirely free of radioactive Ar^{39} .

Characterization of Tin and Antimony Isotopes. The half-life of hitherto undiscovered Sn^{129} , a product of U^{235} fission, was found to be 3.5 min. New gammas have been found in the decay of Sn^{128} and Sb^{128m} , and the beta energies of these isotopes have been revised. The half-life of Sn^{130} has been found to be 3.5 min instead of the 2.6 min reported in the literature.

Cross Section Measurements. Thick-target reaction cross sections have been measured for neutrons and tritons of incident energy 14 and 2.7 Mev, respectively: $\text{Ti}^{48}(n,p)\text{Sc}^{48}$, 55 ± 1 mb; $\text{V}^{51}(n,\alpha)\text{Sc}^{48}$, 18 ± 3 mb; $\text{Ca}^{48}(n,2n)\text{Ca}^{47}$, 920 ± 50 mb; $\text{Ti}^{47}(n,p)\text{Sc}^{47}$, 99 mb; $\text{Ti}^{50}(n,\alpha)\text{Ca}^{47}$, 10 ± 2 mb; $\text{Te}^{130}(t,p)\text{Te}^{132}$, $<7\mu\text{b}$; and $\text{Mg}^{26}(t,p)\text{Mg}^{28}$, 8 mb. Although the $\text{V}^{51}(n,\alpha)\text{Sc}^{48}$ cross section disagrees with literature values (which disagree with each other), it agrees very well with a calculated value based on the assumption of compound nucleus formation.

$\text{Mg}^{26}(t,p)\text{Mg}^{28}$ Excitation Function. Thin foils of magnesium have been made, preparatory to determining the 0 to 2.7-Mev region of the excitation function of the $\text{Mg}^{26}(t,p)\text{Mg}^{28}$ reaction.

Tagging Protein With Iodine-124. Whereas protein can be readily tagged with I^{131} obtained from Oak Ridge, tagging with I^{124} produced at Brookhaven had been erratic and yields generally low. An intensive study of this problem revealed

that the iodination yield is adversely affected by use of permanganate during the preparation of I^{124} and by the presence of free chlorine, iodide ion, or traces of antimony (from the target) in the I^{124} product; moreover, the yield varies directly with the concentration of ICl (the iodinating agent). The procedure for production of I^{124} has therefore been modified, and consistently good protein iodination yields can now be obtained.

Decay Scheme of Scandium-48. The new 170-keV gamma of Sc^{48} was shown to be in coincidence with each of the 1.0-Mev gammas, with the 1.3-Mev gamma, and with a new softer (≈ 480 -keV) beta, but not with the harder (≈ 650 -keV) beta.

Half-Lives of Aluminum-28 and Fluorine-18. A new technique using the IBM 7090 computer which permits simultaneous determination of counter resolving time, sample half-life, and sample initial activity was used to measure the following half-lives: Al^{28} , 2.28 ± 0.01 min; and F^{18} , 109.9 ± 0.2 min.

Gamma-Back-Scatter Coincidences. The search for gammas in coincidence with back-scatter radiation has proved useful in deciding, when unraveling a complex decay scheme, whether or not a gamma is present that is not in coincidence with the others. It has also been found to be a useful technique for the determination of positron-emitting impurities in a beta-gamma-emitting matrix.

Radiation and Chemonuclear Processing Research

The radiation and chemonuclear processing program is concerned with investigations of the application of nuclear energy to the synthesis and production of industrial chemicals. The effort is divided into two main areas: (a) radioisotope and machine radiation studies, and (b) fission fragment chemonuclear studies.

Radiation Processing Program. Co^{60} gamma radiation has been used to investigate the following chemical systems:

1. **Anhydrous and aqueous ammonia for the production of hydrazine.** In the homogeneous anhydrous system the $G_{\text{N}_2\text{H}_4}$ value reaches a maximum of 0.13. In the aqueous system this value decreases below 0.13 with decreasing ammonia concentration and increasing doses and dose rates. Hydroxylamine is formed in the aqueous system. The $G_{\text{NH}_2\text{OH}}$ value reaches a maximum of 0.61 at 28% NH_3 concentration. It is believed that hyponitrite is also present.

When anhydrous ammonia is adsorbed on a solid substrate of pure sodium aluminum silicate, the $G_{N_2H_4}$ value is found to increase by a factor of 10, and the rate is enhanced to ≈ 3 times that obtained in the homogeneous system.

2. The polymerization of ethylene. The G values and rate of polymerization are being investigated at an ethylene gas pressure of up to 680 atm, in the temperature range of 20° to 100°C, and at dose rates from 20,000 to 500,000 rad/hr. Oxygen concentrations >60 ppm have been found to inhibit the rate of polymerization. The inhibition period decreases with decreasing oxygen content. The rate increases with the first power of the ethylene pressure and with the 0.3 to 0.5 power of the radiation intensity. At 21,000 rad/hr a maximum differential G value of 47,000 was obtained at a rate of 14.2 g/l/hr. The polyethylene formed is a fine white powder. Its properties depend on the conditions of formation, but fall generally into the category of intermediate density conventional polymer.

3. Copolymerization of ethylene with other monomers. Survey experiments have been undertaken to determine monomers that will mutually polymerize with ethylene at pressures up to 680 atm. Preliminary experiments indicate that methyl methacrylate, acrylonitrile, vinyl acetate, and acetone are of interest in this respect.

4. The reaction of uranium and propane. The effect of Co^{60} gamma radiation on this system at temperatures up to 350°C indicated a minor formation of uranium monocarbide.

5. Metal deposition studies. The radiation decomposition of dicumene chrome as a method of chromium deposition at low temperatures is being investigated. Initial experiments indicate a 25° to 100°C lowering of the thermal threshold.

A design study of a high-pressure gamma loop facility has been undertaken. The facility would be of use in investigating problems in radiation physics, chemistry, and engineering associated with promising radiation processes.

Chemonuclear Processing Program. This program is concerned with studies in fission fragment chemistry and chemonuclear reactor development.

An in-pile chemonuclear loop is being designed for installation in the BGRR to study (1) fission fragment and fast neutron chemistry in a circulating system, (2) the energy deposition efficiency of chemonuclear fuel elements, and (3) the contamination and decontamination of chemical products.

The in-pile section is being designed to incorporate movable and replaceable fuel elements. The out-of-pile section will have a gas bearing compressor for circulating gases through the in-pile section at pressures up to 68 atm and temperatures up to 538°C. Analytical monitoring of the chemical constituents in the loop will be continuous. The systems to be initially investigated are (a) nitrous oxide for dosimetry, (b) nitrogen and oxygen for nitrogen fixation, (c) carbon dioxide for carbon monoxide formation, (d) methane for acetylene production, and (e) ammonia for hydrazine formation.

In the development of thin-sectioned chemonuclear fuel, studies are being made of uranium-containing alloy foil $\frac{1}{2}$ mil or less in thickness, glass fibers in oriented arrays, oxide coatings, and various honeycomb structures. An out-of-pile test loop has been constructed for testing the thermal, mechanical, and hydrodynamic stability of these fuel elements.

Capsule experiments have been undertaken to determine the energy deposition of prospective elements in a static system. Theoretical calculations of deposition efficiency based on range-energy data are being checked by chemical dosimetry with various thicknesses of fuel.

A design study has been made of a chemonuclear reactor for the cracking of carbon dioxide to carbon monoxide. The two concepts analyzed include a two-region dual-cycle fixed-fuel reactor core and a cavity reactor core.

High Temperature Technology

High Temperature Physical Properties Program.

This program is concerned with the measurement at high temperatures of the physical properties of refractory nuclear fuel materials. It was concluded that the highest yield of information with the personnel and facilities available could be achieved by x-ray diffraction, and the initial phase of the program has been concerned almost entirely with this technique. An apparatus has been developed with which diffraction measurements can be made in vacuum and in reducing or inert atmospheres at temperatures up to 3000°C. A rhenium filament is used as a heater and specimen stage. Temperature measurements were made in the intermediate range by thermocouple and in the high range by pyrometry under essentially black body conditions. Thermal expansion coefficients for

platinum to 1550°C and for ThO₂ to 2500°C were obtained. This method should soon make lattice-parameter and thermal-expansion measurements at very high temperatures a routine procedure.

Release of Fission Products From Fuel Elements.

The objective of this program is to provide basic knowledge of the chemical and physical behavior of fission products released from fuel elements in the event of a nuclear accident.

One area of investigation was measurement of the temperatures at which iodine, cerium, and barium-lanthanum condense after release into a helium stream from the postirradiation melting of a U-Mo fuel sample. The helium stream and the released fission products passed through a decreasing temperature gradient (1200° to 30°C at 100°C/in.) and then into a liquid nitrogen charcoal trap. The temperature at which each fission product was deposited was determined by a gamma-ray spectrum analysis of the meltdown tube wall at the end of the experiment. Iodine was deposited at temperatures >250°C, which indicated that it was deposited not as elemental iodine, but in some combined form. Cerium and barium-lanthanum were found deposited in the regions 1100° to 1200°C; however, in one experiment in which the uranium was badly oxidized, cerium was found widely distributed throughout the meltdown tube from 1200° to 300°C.

The reactions between individual fission products and molten uranium were also studied. Tellurium and uranium were found to react at temperatures as low as 900°C. Several U-Te compounds have been reported in the literature (i.e., UTe, UTe₂, U₃Te₄); however, very few physical or chemical data are given, and it is impossible to predict which of these compounds are thermodynamically stable at high temperatures and would be likely to form in the temperature range of interest in the Safety Release Program (1200° to 1900° K). X-ray diffraction and fluorescence techniques are being used to study the reaction products.

Preliminary results indicate that a single phase with the approximate composition of U₃Te was formed in a reaction between uranium and excess tellurium at 950°C. This result, if verified in succeeding experiments, is of significance for the Safety Release Program, since it indicates that the stoichiometry of the reacting system does not determine the compound formed. Therefore, a simi-

lar result may be expected in an irradiated fuel when a large U/Te ratio exists. This phase was successfully indexed on the basis of a tetragonal cell, $a=11.30$ Å, $c=11.05$ Å.

Experiments are under way to determine the diffusion of fission products in the cladding materials of reactor fuel elements by measuring the uptake of the fission products by the cladding material as a function of time. The diffusion of tellurium in aluminum is being studied by using Te¹²⁹ as a tracer. The results of one series of experiments at 600°C indicate a diffusion coefficient of 5×10^{-13} cm²/sec; however, the aluminum sheet contained a surface film of Al₂O₃, and diffusion through this film may be the rate-controlling step.

Sorption and Diffusion of Fission Products in High Temperature Materials. The sorption of iodine in graphite has been investigated by using I¹³¹ tracer techniques. Two different types of graphite were studied over the temperature range 25° to 800°C.

When graphite was exposed to saturated iodine at 25°C, more than an equivalent nitrogen monolayer of I₂ was sorbed in 19 days. This iodine could not be removed by heating to 950°C in a vacuum and had not reached equilibrium.

In addition to this irreversible sorption, iodine was reversibly adsorbed in the temperature range 200° to 800°C. At 800°C, the adsorption appeared to be independent of iodine pressure in the range 2 to 30 mm Hg but was small, amounting to 1.6% of an equivalent nitrogen monolayer (0.37 m²/g). The adsorptions of the two graphite samples of the same type differed by an order of magnitude at 300°C.

The adsorption of iodine on graphite at 100°C gave a surface area equal to that obtained with nitrogen at liquid nitrogen temperature; at 12 mm of I₂ pressure, equilibrium was achieved in ≈ 2 hr.

Fuel Reprocessing

Reprocessing Reactor Fuels by Volatilization Through the Use of Inert Fluidized Beds.

1. **Stainless steel decladding with Cl₂.** A major problem in decladding stainless-steel-clad fuel by volatilization reaction with Cl₂ is the corrosion of the reactor vessel at operating temperatures ($\approx 600^\circ\text{C}$). Therefore the use of ceramic-lined vessels is being considered. From the standpoint of corrosion resistance alone high density alundum appears acceptable. To fabricate such a vessel a shrink-fitting technique is being used which in-

volves spray-coating the outer surface of the alundum tube with nickel to a thickness of $\approx \frac{1}{16}$ in., machining to give a cylindrical surface and the appropriate diameter, and rapidly inserting the coated tube into a nickel pipe preheated to $\approx 850^\circ\text{C}$ in an inert atmosphere.

Several tubes of $\approx 1\frac{1}{2}$ -in. i.d. and a clad length of ≈ 15 in. were fabricated by using the above procedure. Flanges were welded directly at the ends of the nickel pipe, and such a flanged vessel was then used to carry out the Cl_2 -stainless steel reaction in a fluidized bed of inert Al_2O_3 . The temperature of the lined section of the vessel was $\approx 700^\circ\text{C}$, while the base metal portion extending beyond the alundum liner was held at 500°C . With a peak temperature in the reaction zone of 800°C , cracks developed in both ends of the alundum tube liner. The formation of these cracks may well be due to the greater coefficient of thermal expansion of the nickel as compared with that of the ceramic. It is planned to fabricate similar reaction vessels with shrink-fitted pipes of metal having coefficients of expansion closely matched to that of alundum.

2. Zircaloy fuels. In a continuation of the studies on the recovery of uranium from Zircaloy fuels by volatilization of the zirconium by hydrochlorination and fluorination of the uranium, a series of experiments was carried out with reaction temperatures in the range 350° to 400°C . The results showed that when the initial uranium concentration in the bed after hydrochlorination was at the level that would obtain with full-scale reprocessing of spent reactor fuels, $>99\%$ of the uranium was removed from the bed by fluorination.

3. Uranium carbide fuels. The application of the inert fluidized bed process to the recovery of uranium from spent uranium carbide and graphite matrix fuels by direct volatilization is considered promising. This would be a two-step operation involving the reaction with oxygen to produce CO_2 and U_3O_8 , followed by fluorination to give volatile UF_6 . Some preliminary work on the oxidation of uranium carbide to produce U_3O_8 in an inert fluidized bed has been reported in BNL progress reports, but no work was done on the recovery of uranium as the volatile fluoride. During the past year a program was initiated to study the reprocessing of uranium carbide fuels, and several experiments were carried out to obtain quantitative data on uranium recovery. The results of three

typical experiments show the recovery of uranium to be 99.6, 98.7, and 99.1%.

Nitrofluor Volatility Process. The alternate solvents, $\text{NOF} \cdot 3.1\text{HF}$ and $\text{N}_2\text{O}_3\text{-HF}$, for the Nitrofluor process have compared favorably with the original solvent, $\text{N}_2\text{O}_4\text{-HF}$, in dissolution studies with several fuel element materials. Both alternate solvents are capable of dissolving Zr, Al, UO_2 ceramic fuel, and type 304 stainless steel at satisfactory rates. Thus, their versatility approaches that of $\text{N}_2\text{O}_4\text{-HF}$, the essential difference being that they do not disintegrate graphite or graphite-matrix fuels. This oxidizing action is related to the existence of NO_2^+ (nitryl) ions and/or molecular HNO_3 , neither of which is found in the alternate solvents. Pyrolytic graphite-coated UC_2 spheres and ZrC are also rapidly attacked by $\text{N}_2\text{O}_4\text{-HF}$.

During fuel dissolution the uranium and plutonium are converted to complex fluorides which can be appropriately fluorinated to their volatile hexafluorides by treatment with BrF_3 and F_2 , respectively. A significant improvement in uranium volatility was made by increasing the fluorination temperature from 100° to 150°C . After six treatments with BrF_3 , only 94 ppm uranium ($<0.2\%$) was retained in the salt residue containing excess zirconium. In a new volatilization reaction, the uranium in NOUF_6 salt was separated as a volatile compound by contacting the NOUF_6 with either gaseous $\text{NOF} \cdot 3.1\text{HF}$ or NOF rather than with BrF_3 . The reactions were performed in a tube furnace at 500°C , and UF_4 was identified as one of the solids that condensed in a downstream zone at 390° to 400°C . Although the reactions are not yet understood, the method appears to have interesting possibilities. Thus, $\text{NOF} \cdot 3.1\text{HF}$ could serve initially as the fuel solvent and subsequently as the volatilizing agent. A small alpha facility for determining plutonium behavior in the process is under construction and should be ready for use in September 1962.

Significant progress was made in solving the corrosion problem associated with handling the three solvents in nickel or Monel equipment. In loop tests, corrosion rates as low as ≈ 1 mil/month were achieved with certain combinations of solvent and container material for both zirconium-matrix fuel and UO_2 ceramic fuel. The exposure involved initial fuel dissolution followed by alternate evaporation and condensation cycles during a 1-week

Figure 4. Fused salt-liquid metal extraction apparatus.

period. The results showed that zirconium-matrix fuel was best handled in nickel equipment by using $\text{NOF} \cdot 3.1\text{HF}$ as the solvent. The best combinations for UO_2 fuel were nickel with $\text{N}_2\text{O}_4\text{-HF}$ and Monel with $\text{NOF} \cdot 3.1\text{HF}$. If both types of fuel are to be handled in the same equipment, nickel would be preferred to Monel.

Bench-scale equipment is being assembled for studies with UO_2 ceramic fuel. The results of this work will be applied to the engineering-scale equipment to be erected in the Volatility Studies Laboratory now under construction. This building will be ready for occupancy in September 1962 but will not be completed until early in 1963. The process equipment already fabricated will be installed during this interval. Some equipment lined with platinum may be used, since this material has exhibited complete resistance to attack under a wide variety of process conditions. A minor effort will be devoted to process studies with graphite-matrix fuel.

Aqueous Chemical Processing. A modified Sulfex process for dissolving stainless steel in a stainless steel vessel has been developed. In this process the vessel is kept passivated by the addition of 0.05 to 0.1 $M \text{HNO}_3$; the dissolution of the fuel element in a Ni-o-nel basket is initiated either by a mild steel wire wrapping the element or by the brazing material used in assembling the fuel element. The nitrate level is maintained by reoxidation of NO in a reflux condenser.

A steam-jacketed 309 SCb stainless-steel pilot-plant dissolver with a liquid capacity of ≈ 80 liters has been constructed for demonstration of this process. Several successful runs have been made in which stainless steel has been dissolved to a concentration of 25 g/l with no significant attack on the stainless steel dissolver vessel.

Fused Salt-Liquid Metal Extractions. An 18-month experimental study of rates of mixing and mass transfer in fused salt-liquid metal extraction columns was completed at the end of the fiscal year. The two phases used were the NaCl-KCl-MgCl_2 eutectic and Bi-Mg alloy. Cerium was the transferring solute. Experiments were carried out at 500°C . The apparatus (Figure 4) consisted of two pumped loops, one for the liquid metal and one for the salt, which interconnected at one or the other of two extraction columns. Superficial flow rate ranges were 11,000 to 57,000 lb Bi/hr-ft² and 1100 to 5700 lb salt/hr-ft².

Axial mixing in the continuous salt phase was studied by imposing a step change in tracer (Na^{24} as Na^{24}Cl) concentration on the influent salt and measuring the resulting tracer transient in the effluent salt. Interpretation of breakthrough curves so obtained for a 4-in.-i.d., 42-in.-long disk-and-doughnut column led to a value of 5 for the Peclet number for axial mixing, independent of the flow rates of the two phases. This indicated that there would be little effect of axial mixing on over-all rates of mass transfer.

Mass transfer experiments were carried out in the same column at several concentrations of magnesium in bismuth and with several different flow rate ratios of the two phases. Preliminary interpretation of the data led to over-all mass transfer coefficients in qualitative agreement with the predictions of a model [Handlos and Baron, *A.I.Ch.E. Journal* 3, 127 (1957)] which has been successfully used for aqueous-organic systems near room temperature. The height of a transfer unit

(HTU) was found to vary strongly with the concentration of magnesium in bismuth, as expected. The minimum HTU predicted for the column is ≈ 2 in.

Radiation Engineering

Radiation Development Center. Most of the heavy construction of building, cells, and laboratories for the High Intensity Radiation Development Laboratory has been completed (Figure 5). Yet to be done is the installation of cell doors, windows, and internal equipment. Completion of the entire laboratory is scheduled for December 1962.

Conceptual Design Studies. At the request of the Division of Isotopes Development of the AEC, conceptual studies were made in cooperation with Hughes Aircraft Company and Associated Nuclear, Inc., of transportable Co^{60} irradiators and transportable accelerator irradiators to be used as test units for the radiopasteurization of fruit in the California area.

Blood Irradiator. A design has been completed for the BNL Medical Department of a 750-curie Co^{60} source for the extracorporeal irradiation of blood.

Co^{60} Low-Level Dose Distribution Studies. Dose distribution studies in heterogeneous targets have been completed by utilizing rectangular arrays of

3-in.-diameter cylinders. Work has begun on arrays of spheres of various diameters.

Investigations have been completed on the dose distribution in rotating, semi-infinite, water-filled targets of 6-in. diameter utilizing semi-infinite, hollow cylindrical Co^{60} sources 3 in. in diameter. Experimental apparatus has been assembled to extend these investigations by using targets and sources ranging in diameter from 3 to 12 in. with air gaps ranging from 1 to 6 in.

Dosimetry. An extensive program has been carried out for the evaluation of all available commercial solar cells for possible use as radiation dosimeters. The work completed thus far indicates that the n on p type of silicon solar cell has wide application as a radiation dose rate dosimeter over a range of 10^4 to 10^7 rad/hr.

A low-level anthracene scintillation dosimeter has been developed utilizing $\frac{1}{4}$ -in. diam $\times$ $\frac{1}{4}$ -in. high crystal. The range of this instrument is 1 mrad/hr to 1 rad/hr.

Heat Generation. Studies to supplement the heat transfer measurements made on the ETR Co^{60} shipping container have been performed in a series of *in situ* thermal experiments employing up to 150,000 curies of Co^{60} . The experimental source container was designed to yield fundamental information on temperature distribution and the

Figure 5. The High Intensity Radiation Development Laboratory, May 1962.

basic mechanism of heat transfer within the source array. The data collected indicate that (1) the temperatures within an ordered array of radiation sources are largely dependent on source spacing, heat transfer being primarily conductive where sources are closely spaced; (2) small amounts of water in an air-filled source array lower maximum temperatures via an evaporation-condensation mechanism of heat removal; and (3) temperatures are significantly dependent on vertical positioning of the sources, particularly in a partially filled source container. These experiments, when complete, will provide a semiempirical or empirical relationship useful for designing future source shipping containers in which heat generation is a significant problem.

The occurrence of electrical discharge in lead shielding glass, resulting from gamma-ray irradiation, has been demonstrated in an initial investigation of this phenomenon. An $8 \times 9 \times 10$ -in. slab of 3.3-density lead glass exhibited electrical cracks, thermal cracks, and induced stress patterns subsequent to a Co^{60} radiation dose of 1.1×10^7 rad at a dose rate of $\approx 5 \times 10^5$ rad/hr. The study of radiation effects on large glass slabs will be continued, with glass thickness and cross section as major parameters. A fundamental study is also planned in which shallow electron "trapping," presumably related to the discharge phenomenon, will be investigated in smaller samples by means of measurements of color-center build-up, stress birefringence, and thermoluminescence in a variety of glasses.

Research Irradiator Project. In the last fiscal year an irradiator designed for radiopasteurization studies was built and installed at MIT, and two additional units were built and installed at the University of California (Davis), and the University of Washington, Seattle. These units employ $\approx 30,000$ curies of Co^{60} and can irradiate packages up to $14 \times 18 \times 6$ in. in size.

Foreign Exhibits. A second irradiator similar to the one used in Egypt and South America was fabricated and shipped to Beirut, Lebanon. Three stainless steel encapsulated cobalt plates containing 2300 curies are used as a source of gamma radiation. This unit was also used in Athens, Greece.

The gamma source strength of the original unit, used in exhibits at Cairo, Egypt; Buenos Aires, Argentina; Rio de Janeiro, Brazil; Lima, Peru; and Mexico City, Mexico, was supplemented by

Figure 6. Heat transfer to turbulent, in-line flow of mercury through a staggered rod bundle, under conditions of fully established flow.

Figure 7. Heat transfer to turbulent, in-line flow of mercury through a staggered rod bundle, at point near the flow inlet.

Figure 8. Heat transfer to turbulent, in-line flow of mercury through unbaffled rod bundle, showing entrance effect.

700 additional curies prior to the opening of the exhibit in Mexico City.

Liquid Metal Heat Transfer Research

Late in the fall of 1960, an improved mercury heat transfer loop was placed in operation for the purpose of measuring heat transfer coefficients for turbulent, in-line flow through unbaffled rod bundles. This loop has since been operated several times. Data were taken on the central rod of a 13-rod bundle. The rods were $\frac{1}{2}$ in. in o.d. and spaced in an equilateral triangular array, the distance between centers being 0.875 in.

For about a year, data were taken in the Peclet number range 900 to 4000; then placement of a smaller orifice in the flowmeter made it possible to study Peclet numbers as low as 200.

Some typical results are shown in Figures 6, 7, and 8. The meanings of the symbols in the figures are as follows:

- Ne = Nusselt number = hD_e/k ,
- h = heat transfer coefficient,
- D = outside diameter of rods,
- D_e = equivalent diameter of bundle,
- k = conductivity of liquid metal,
- Pe = Peclet number = $D_e v C_p \rho / k$,
- v = average linear velocity,
- L_f = upstream flow length through bundle,
- L_h = upstream heated length of rods,
- ρ = density of liquid metal,
- P = distance between rod centers,
- C_p = heat capacity of liquid metal,
- Pr = Prandtl number = $C_p \mu / k$, and
- μ = viscosity.

Figure 6 shows some typical results obtained under conditions of fully established flow at an L_f/D_e ratio of 41.1 and an L_h/D_e ratio of 38.7. The curve drawn through the data points was predicted on the basis of previous analytical and experimental results obtained at Brookhaven. Figure 7 shows the typical heat transfer relationship existing near the flow inlet. The curve is straight rather than "dog-leg" shaped, owing to the high degree of turbulence. At a Peclet number of 200, however, the Nusselt number appears to be independent of the L_f/D_e and the L_h/D_e ratios, presumably because the degree of turbulence is insufficient to produce a significant amount of eddy transport, and therefore the heat transfer coefficient remains essentially at the streamline value.

Figure 9. Alkali metal heat transfer loop during late stages of construction.

Figure 8 shows a set of typical curves depicting the effects of the L_f/D_e ratio and Peclet number on the Nusselt number. In this set of results, L_h was 2.0 in. less than L_f .

In the fall of 1961, the construction of an alkali metal heat transfer loop (Figure 9) was completed, but it was not placed in operation until January, and the first data were taken in February 1962. Initially, this loop will be used to obtain data on the in-line flow of NaK through rod bundles to see if the results agree with those obtained with the mercury loop. The present bundle consists of 19 0.500-in. rods spaced 0.875 in. apart in an equilateral triangular array. Thus far, data have been taken on two rods, each located in the central

position. The results from the first test element showed appreciable scatter, although they were in substantial agreement with the curve shown in Figure 6. The results from the second element fell low. The results from both elements were affected by rotation, which indicated that geometrical dissymmetry was present. It is suspected that there was insufficient clearance for the rods to slide freely through the bottom guide plate as they expanded on being heated. The six rods surrounding the test element were removed and straightened, and the clearances were increased in the guide plate. Data will soon be taken on a new nickel-plated test element.

A loop is under construction which will provide fundamental heat transfer information to be used in the design of boilers and condensers for auxiliary

Figure 11. New type of rotary seal recently developed for use with the rotary calciner.

power plants in space vehicles using alkali metals as working fluids. It should be ready for operation in the spring of 1963.

Along with the experimental studies, related analytical studies are being made. As an example of the latter, a successful attempt was recently made to reconcile the differences between theory and experiment that have always existed in liquid metal heat transfer. Evaluating $\bar{\psi}$ (for use in the standard calculational methods) by the following equation has been found to bring theory and experiment into substantial agreement for flow through tubes, annuli, and rod bundles:

$$\bar{\psi} = 1 - \frac{1.82}{\text{Pr}(\epsilon_M/\nu)^{1.4}_{\max}},$$

where

- $\bar{\psi}$ = average effective value of ϵ_H/ϵ_M ,
- ϵ_H = eddy diffusivity for heat transfer,
- ϵ_M = eddy diffusivity for momentum transfer,
- Pr = Prandtl number,
- ν = kinematic viscosity, and
- $(\epsilon_M/\nu)_{\max}$ = maximum value in flow cross section.

Ultimate Waste Disposal

Preparation of Phosphate Glass. During the past year major emphasis was placed on the formation of phosphate glass as a means of incorporating fission products in stable media. The phosphate system, in which the material is in the liquid phase throughout the process, lends itself readily to the continuous mode of processing. Continuous glass formation was carried out on a small scale at

Figure 10. Pilot plant for producing phosphate glass from simulated high level wastes.

1100°C by using a platinum crucible 2 in. in i.d. and 6 in. long, equipped with an overflow weir designed to provide maximum retention time in the crucible and with a bottom outlet jacketed tube to function as a freeze valve. Runs of 500 hr with simulated Purex and 125 hr with redox wastes were carried out.

To study operational problems and glass handling techniques on a more realistic scale a small pilot plant (Figure 10) for producing phosphate glass from various types of simulated high level wastes was constructed. Operation of the plant has been postponed pending the installation of a platinum-clad Inconel crucible, 8 in. in diameter and 24 in. long, to replace the high-density alumina crucible (shown in the figure) which was found to be incapable of withstanding the stresses imposed by high temperature service.

Calcination. A calciner of the rotary type for the treatment of high activity, aluminum-bearing wastes affords the combined advantages of continuous operation and effective dust control. At present the rotary calciner development work is directed toward perfecting the required mechanical equipment rather than process studies. In particular, a new type of rotary seal (see Figure 11) has been developed which operates on the principle of supplying steam under pressure to a very small gap between the rotating face and the stationary face, so that the small leakage of steam into the calciner with relatively high velocity will prevent the escape of radioactive gases or dust to the atmosphere. Other major mechanical improvements are being made in the calciner pilot plant, and a continuous 1000-hr run is planned during the latter months of 1962.

Applied Mathematics

As indicated in last year's report, the reorganization of the Applied Mathematics Division has resulted in its orientation to a large extent, but by no means exclusively, toward those areas of research in which large-scale digital computers serve as basic research instruments. The acquisition by the Division of a modern computer system has perforce meant that the major part of the work of the Division during the year has been directed toward acquainting the various departments of the Laboratory with this system. All members of the staff have devoted a major portion of their time to this effort, irrespective of the nature of their appointments, and in many cases with considerable sacrifice of their own personal research interests.

COMPUTER FACILITY, SYSTEMS, AND UTILITY PROGRAMS

During August 1961, an IBM 1401 system was installed as the first step in the provision for BNL of an advanced computation facility and was immediately put to use for the preliminary processing of input for the IBM 704 at NYU, which the Laboratory was then using, and the printing of output from that machine. During October, while the NYU machine was being replaced by an IBM 7090, the Division arranged for the commercial rental of time on both a 704 and a 7090 so that a smooth change-over of BNL's work to the new machine was achieved. In February the Laboratory acquired its own 7090 and was able to discontinue use of the machine at NYU a few hours after IBM's engineers had checked out the new BNL machine.

The high speed at which machines of the 7090 class operate, together with the fact that the only efficient means of getting information in and out is the use of magnetic tapes (the input is placed on them through the 1401, which also prints the output), renders them far less accessible to the originator of any given problem than computers of earlier design. For that reason much thought has been given to matters of organization and scheduling so that the solutions of problems that

do not require excessive printing and are of short duration on the 7090 will be available to the originator within an hour or two. Consequently, much of the printing for runs of longer duration must be done at night, with results available the next day.

Since many of the problems handled by the 7090-1401 complex at Brookhaven are of a special nature, and the machines themselves have many special features not included in the conventional system, an extensive effort was necessary to develop a monitor program suitable for BNL's requirements. A number of modifications and additions to the standard IBM FORTRAN monitoring system were made, and a BNL monitor program is now in satisfactory operation. Furthermore, extensive additions have been made to the libraries of utility programs available with the machines, as dictated by the Laboratory's needs.

The following table is indicative of the growing amount of work which this installation handles.

Month	Machine	No. of hours operated
June 1961	NYU 704	204
Aug. 1961	" "	250
Nov. 1961	" 7090	45
Jan. 1962	" "	71
Mar. 1962	BNL "	132
May 1962	" "	171
June 1962	" "	202

The 7090 operates at approximately five times the speed of the 704.

The Merlin computer continues to be available for those problems to which it is particularly suited, and additions are also being made to the library of programs for that machine.

PROGRAMMING AND THEORETICAL RESEARCH

As indicated initially, most of this work has been undertaken in collaboration with research workers in other fields, both at BNL and at universities, or motivated by their needs. Sometimes no more than the writing of programs for a well-

defined mathematical problem is involved, but more often than not the work calls for consultation and collaboration on the theoretical formulation of the problem, or independent work in this area. In some cases an assignment may be completed in a few days; in others, it may require months or even years and involve the efforts of several people. Here only a few examples will be discussed briefly.

A wide variety of problems in the physical sciences require, for their solution, the inversion of high-order matrices, the determination of their eigenvalues and eigenvectors, or simply the solution of linear systems of equations of high order. The large number of arithmetic operations involved in handling such matrix problems requires extreme care in the choice of method and careful attention to the problem of cumulative errors in arithmetic; failure to exercise such care can render formal arithmetic results meaningless. Therefore, over the past year considerable effort has been devoted to this field, and an extensive set of pro-

grams has been developed for the solution of such problems on the 7090, and to a lesser extent on Merlin.

Members of the Division have for some years now been working with the Solid State Group of the Physics Department, writing programs for the study of radiation damage in crystals. The mathematical model of the problem leads to systems of coupled differential equations of the order of 1000 or more, with a variety of boundary conditions. Since these programs were originally written for the 704, extensive rewriting was necessary to prepare them for the 7090. This was completed early in the year. Furthermore, the increased capabilities of the 7090 together with the development of more sophisticated programming techniques has made it possible to consider more complex problems, and a new set of programs has been written. Considerable interest in this work has developed outside the Laboratory, and a number of visitors have been supplied with the programs and instruction in their use.

Figure 1. Scene in the new Computer Center. The operator at the console (right) is setting up a problem. At the rear, magnetic tape is being installed for input to the problem. The printer is to the left.

In collaboration with members of the Physics Departments at BNL and Columbia University, the problem of two-body inelastic collision going into n particles was formulated mathematically for the calculation of invariant phase space and mass distribution, energy spectra, and angular correlations. The problem has been programmed for the 7090 and the program is now in use.

In collaboration with members of the staff of the Institute for Advanced Study, work has been started on a program for the calculation of W -particle production cross sections. Completion of this program awaits some modification of the mathematical model.

For the past several years members of the Division have been engaged with members of the Chemistry Department in the development of a more sophisticated and elaborate approach to the low energy intranuclear cascade problem and the concomitant preparation of the necessary machine program. This work was originally planned for Merlin, but in the spring of 1961, when it became obvious that the greater power of the 7090 should be utilized, those parts of the program that had been written for Merlin were rewritten for the 7090, and the rest of the program was completed for that machine. The program is currently in use by the Chemistry Departments at BNL and Columbia University, and plans are under way for its use at the University of Chicago.

A great deal of the Division's work has been related to problems arising in connection with the processing of bubble chamber data. First, since the measuring machines in use at BNL have paper tape output, a number of programs had to be written for transferring the data in appropriate format to magnetic tapes for various 7090 programs. Second, since it was decided to use, in addition to the BNL set of programs, the FOG-CLOUDY-FAIR set of 7090 programs written at the Lawrence Radiation Laboratory (LRL), in which the input format is quite different, an extensive set of both 1401 and 7090 programs had to be written to convert the output of the BNL measuring devices to a radically different form. Third, differences in the nature of the bubble chambers and the different energy ranges of the accelerators, among other considerations, raised questions with respect to FOG-CLOUDY-FAIR that were far from trivial. In collaboration with members of the Bubble Chamber Group and with the virtually indispensable

assistance of the authors of FOG-CLOUDY-FAIR, these problems are being resolved. Finally, again with members of the Bubble Chamber Group and the LRL staff, various programs are being prepared in connection with the use of flying-spot digitizers in conjunction with the FOG-CLOUDY-FAIR system.

The increasing volume of bubble chamber pictures being produced each year has led to a variety of attempts in several universities and laboratories to reduce the amount of human intervention necessary in the extraction of pertinent physical information from the photographs. The procedures now in use involve first the scanning of the film with the aid of such semiautomatic devices as the Frankenstein, the operator of the device punching the requisite data into cards or paper tape, and second the processing and analysis of these data with elaborate high-speed digital computer systems. The work indicated above is all directed toward new approaches to the first part of the process and envisages its incorporation into the second with only minor modification of the latter.

The Physics Department now has under construction a Hough-Powell mechanical flying-spot digitizer, which is to be finished within a few months. The input is the film itself, and through a direct data channel the digitizer feeds into the memory of the computer what it "sees" on the film. But since much of this information is of no interest, human intervention is still necessary to provide the computer with a set of instructions as to which areas of the film are of interest; this involves the use of a separate off-line scanning device.

Automatization of this latter procedure, whether by programmed instruction to the computer, by special-purpose hardware, or a combination of both, is obviously the next step. In the Applied Mathematics Division a group has for the past year been working on the first approach. Indications are that a reliable simulation of the scanning process can ultimately be achieved, although for purely economic reasons it may be better to have a portion of the work handled by special-purpose devices rather than the computer itself. It was not clear at the outset whether such a program could run fast enough to be economically competitive with "conventional" systems involving human scanners working in parallel. Considerable emphasis was therefore placed on achieving high data-handling rates, and several coding schemes have now been devised that are expected to settle

the question of speed affirmatively. In all the schemes completed or proposed to date, it must be emphasized, what has been or will be achieved is not automatic scanning but rather the speedy reduction of the original mass of data to a form (such as a table of track parameters) more amenable as input to a true, but as yet hypothetical, scanning program. The latter will inevitably have intrinsic logical complications, but, since it would handle relatively small quantities of data, it is reasonable to assume that it can be made relatively fast. Work on this phase of the problem is now under way. Apart from its obvious practical importance, this work is of interest for its possible wider ramifications in the area of pattern recognition generally.

Toward the end of the year, in conjunction with development by the Engineering Group of the Division (see below) of an automatic spark chamber film measuring projector, and in collaboration with members of the Physics Department, the preparation of a set of spark chamber data processing programs was begun.

A brief account is given below of work not programmatically connected with other Departments of the Laboratory.

Certain preliminary work on the use of a digital computer for performing the Feynman diagram calculations mentioned in last year's report has been completed and will be published shortly. Research on this problem continues.

A theoretical study of unstable particles based on a modified Lee model has been performed. For this purpose, a four-point interaction was superimposed on the conventional Lee Hamiltonian and was found to be completely soluble in the physical sector corresponding to a physical and/or bound state particle.

Research has been initiated in the application of operator theory to the study of boundary value

problems for elliptic differential equations. Toward the end of the year, with additions to the staff, research was begun in the field of parabolic differential equations and their applications.

ENGINEERING DEVELOPMENT

Since the efficient use of a modern high-speed computer system for problems of the sort encountered at BNL involves a variety of engineering problems not only in the design and construction of special-purpose peripheral equipment, both on- and off-line, but also frequently in modifications to the central system itself, a computer engineering group was added in 1961 to the staff of the Division. Mr. Jerome A.G. Russell was appointed Assistant Head of the Division for Engineering Research and Development in November, and a small group is now working in this field under his direction. This group first made a thorough and careful engineering study of the present and potential capabilities of the Merlin computer. In addition, the group has under consideration several useful expansions of the 7090-1401 system. As in the case of the theoretical staff, however, most of their work has been in support of other researchers in the Laboratory, for the most part in connection with the efficient use of the 7090-1401 system. Members of the group have designed and installed the electronic interface between the 7090 and the flying-spot digitizer and provided further substantial design aid for the construction of the latter. They have completed the preliminary design and begun construction of the spark chamber film measuring projector mentioned above, and have initiated the study of a cathode-ray-tube flying-spot digitizer. In addition, the group has provided assistance to other groups in the Laboratory on various electronic engineering problems not connected with computer problems.

LIFE
SCIENCES

Biology

The activities of the Biology Department center on the special facilities of the Laboratory and include studies of the biological effects of radiation and the use of isotopes for the elucidation of basic problems in biology. In general, problems are being investigated by the methods of molecular biology. The growth of the Department has been increasingly in the direction of this approach to biology and has involved studies on protein structure, enzyme kinetics, the molecular structure and function of antibodies, the molecular structure of chromosomes and its relation to the mutation process, etc. These concepts and techniques have found immediate application in such pressing problems as the nature of radiation-induced mutations, the intimate nature of the mechanism of radiation damage in plants and animals, and the details of the basic reactions involved in photosynthesis.

Each year the Department sponsors a symposium on a topic of current interest in biology. This year the symposium, entitled *Enzyme Models and Enzyme Structure*, was attended by 194 scientists in addition to members of the Brookhaven laboratory staff and will be published as the 15th volume in the Brookhaven Symposia series.

During the year a "gamma forest" was established. It consists of a large radioactive cesium source in the center of an oak-pine forest in the northeast corner of the Laboratory site. The source went into operation on November 20, 1961, and the immediately adjacent trees are showing marked radiation damage. A careful ecological study is being made of the region.

This report briefly indicates major areas of research and some findings made during the year.

ANIMAL PHYSIOLOGY

Radiation-Induced Aging

Radiation causes a decrease in the life expectancy of animals which closely resembles the natural aging process. A study of both natural and radiation-induced aging is being undertaken to determine the causes of each. It has been postu-

lated that aging results from the accumulation of viable mutations in the somatic cells of the body. This theory gains support from the fact that radiation both shortens the life span and causes mutations. However, nitrogen mustard, a chemical mutagen, has been found not to shorten the life span of mice, which argues strongly against the theory. This contradiction necessitated a direct comparison of the somatic mutations produced under various conditions, and a method was developed based on the assumption that chromosome aberrations are proportional to somatic mutations. Aberrations have been found to increase linearly with age in normal animals and to reach surprisingly high values while the animals are relatively young. Nitrogen mustard increases this rate only slightly, which shows that this agent is not mutagenic for most somatic cells in mammals. However, after a single dose of x rays the abnormalities increase strikingly and are very slowly eliminated. When animals are repeatedly subjected to gamma rays given at a low rate, the abnormalities increase faster than in controls but not as fast as computed for animals given a single equivalent dose at a high rate, which shows that there is some chromosome healing after small doses of radiation. Chromosome damage from a single dose of neutrons is very severe and seems to increase with time after exposure.

Tests for specific organ functions are being developed for mice and applied to mice of all ages. To date, kidney function tests have shown a decreasing functional capacity of this organ with advancing age, but this does not seem to be affected by radiation, which indicates that in the species of mouse used the kidney is probably not a "weak link" as far as aging is concerned.

Effects of Irradiation on the Central Nervous System

X-irradiation of the central nervous system with doses comparable to those used therapeutically evokes clinical signs and symptoms, after an unpredictable latent period of months or more, attributable to injury at the site of the irradiation.

A number of examinations are being made of monkeys, ranging in age from very young to adults, to determine first indications, progress, and consequences of the radiation injury and its nature. The work includes electron-microscopic, histochemical, and histological studies.

Deoxyribonucleic Acid Synthesis

Irradiation is known to affect the synthesis of deoxyribonucleic acid (DNA). Since the nucleic acid is the carrier of genetic information and many radiation effects are genetic in nature, any effects of radiation upon DNA and its metabolism are of potential importance. Although most experimental work has indicated depressed incorporation of labeled precursor into DNA as a consequence of radiation, results obtained with different systems and conditions as a whole appear discordant. Analysis of the problem has suggested possible roots of apparent discrepancies and experimental resolution. The necessary technical developments are under way.

Alterations in DNA metabolism may be important in carcinogenesis induced by irradiation and by other agents. Application of the carcinogen, 7,12-dimethylbenz(α)anthracene, to the plucked skin of mice under conditions known to be efficient for initiation of tumors has been found to depress incorporation of labeled thymidine into DNA in the growing hair follicle of mice $\approx 50\%$ within 24 hr. The carcinogen also delays passage of cells through the period of DNA synthesis, but does not appear to affect other phases of cell division. Whether the mechanisms of action of the carcinogen and radiation are the same or complementary is not yet known.

Action of Insulin on Liver

Increase in uptake of glucose by several isolated tissues (diaphragm, heart muscle, and adipose tissue) in response to added insulin is readily demonstrable and is caused by an increase in permeability of the cell membrane to glucose. The increase in permeability is two-way, i.e., glucose can leave as well as enter the cell more readily. Ordinarily, however, the glucose concentration inside the cell is vanishingly small because the free glucose is rapidly phosphorylated, an essentially irreversible reaction catalyzed by hexokinase. Intracellular release of free glucose by hydrolysis of glucose-6-PO₄ is restricted to a few types of cells,

notably liver cells, which contain the appropriate hydrolyzing enzyme for this purpose. In contrast to the ease with which insulin can be shown to increase glucose uptake by muscle and adipose tissue, it has proved difficult, and indeed may be impossible, to demonstrate an effect of insulin on glucose metabolism in isolated liver tissue. Since the liver continually supplies glucose to the blood, the net movement of glucose in the liver cell is from the inside out. Reasoning from the increase in permeability of the muscle cell membrane, insulin should increase the flow of glucose out of the liver (if anything) if its effect on the liver cell is analogous.

By the use of trace amounts of C¹⁴-glucose it is possible to measure simultaneously in the whole animal total glucose uptake and new (C¹²) glucose production by the liver. When an intravenous insulin infusion is accompanied by simultaneous glucose infusion to avoid lowering the blood glucose concentration, insulin not only increases glucose uptake but also decreases new glucose production by the liver. This effect was difficult to demonstrate because the same amount of insulin given without the glucose does not decrease or increase glucose production by the liver. This is due to a counterbalancing influence brought into play by the lowered blood glucose level; i.e., hypoglycemia in the absence of insulin infusion increases the rate of glucose production by the liver.

Thus in the whole animal insulin acts in a way not predictable on the basis of its best-documented effect at the cellular level, namely, it increases the permeability of certain cell membranes. This may mean that the cellular action of insulin is incorrectly understood. On the other hand, it may mean that the effect observed in the intact liver is not direct but is mediated by the cessation of signals for glucose production coming from a glucose receptor. Such a receptor would "see" more glucose in the presence of infused insulin (and in the absence of lowered blood glucose concentration) if it were separated from the blood glucose by a membrane whose permeability to glucose could be increased by insulin.

Spontaneous Disease in Laboratory Animals

A second attempt to set up a "disease-free" colony of mice is under way. The first colony, bred from animals obtained initially by aseptic Caesarean section and fostered on germ-free females, was

found to have a high incidence of cerebral encephalitozoonosis and was consequently discarded. The second colony was started in the same way from other stock. No infection has been observed in this colony, and mortality has been about one-third that of the first (7 vs 22%). However, a number of conceivably genetic abnormalities have been observed. One affects only male mice; death is sudden, and the essential lesion is massive bleeding into the pleural cavity. The nature of this "hemophilic" syndrome is under observation.

Antibody Specificity

The fundamental problem involved in antibody specificity is whether the specificity, i.e., the ability of each antibody to react only with its particular antigen, results from a genetic change or a rearrangement of the three-dimensional structure after the globulin molecule is synthesized. The genetic theory of formation requires that antibodies differ in their amino acid content or sequence, while the rearrangement theory does not. The large size of antibody molecules, polypeptide chains containing ≈ 1600 amino acids, precludes determination of their total sequence with present techniques. However, since the only measurable differences in antibodies lie in their immunological sites, which involve 10 to 20 amino acid residues, the problem can be attacked by finding marking agents that react at the active site of one but not another antibody.

A suitable method has been devised for labeling the active site of antibody. The method utilizes the fact that groups at the active site of the antibody can be protected against reactions with the labeling agent by prior addition of the specific antigenic group, the hapten. Thus, after treatment during which all groups outside the active site react with the specific reagent, the hapten is removed and the groups at the active site are allowed to react with a radioactive form of the reagent. The method has been applied successfully to incorporate I^{131} into the active site of antibody directed against the negatively charged phenylarsonic acid group. By a correlation of loss of activity following iodination with the appearance of modified residues, a tyrosyl residue has been identified as the reactive amino acid at the active site.

The labeling method has also been used to establish a suitable control, i.e., an antibody whose active site is not marked with radioactive iodine.

Iodination of antibody directed against the positively charged phenyltrimethylammonium group was shown to decrease the immunological activity by one-third the amount observed for the anti-phenylarsonic acid antibody under equivalent experimental conditions. Further, prior addition of the hapten did not protect against loss of activity by iodination. Clearly the active sites of the two antibodies differ in their chemical groups.

Based on these findings, an answer to the problem of antibody specificity appears obtainable from comparison of the iodine-labeled peptide or peptides from the active site of arsonic antibody with the iodinated peptides from the ammonium antibody, since the finding of a unique structure for the active-site peptide would strongly support the genetic theory. Methods have been worked out for quantitative rupture of the 22 S—S bonds in antibody, subsequent digestion of resulting fragments by trypsin, and recovery of peptides without significant loss of the iodine label. Results to date indicate that comparison of iodinated peptides from antibodies is experimentally feasible.

GENETICS

Genetic Plant Tumors

Three approaches are being taken in research on tumors that form in certain hybrids among species of *Nicotiana*. These are (1) factors affecting tumor induction in seedlings by applications of indoleacetic acid (IAA) and kinetin, (2) studies on IAA and kinetin response in tissue cultures of parental species and their tumor-forming hybrid, and (3) nonspecific stress and slow growth as factors in tumor development. In the seedling studies it was noted that added kinetin markedly increases tumor induction when applied to the stem meristem. Kinetin apparently interacts with endogenous IAA to upset normal morphogenetic processes. Tissue culture studies have revealed pronounced differences in response of the parental species and tumorous hybrid to kinetin in the presence of IAA. Parental tissues respond by increased growth whereas the hybrid tissues do not. It appears that the hybrid has sufficient endogenous growth factors, but that IAA and kinetin are required as additives for rapid growth of the parental species tissues. The effectiveness of IAA and kinetin is influenced by the level of inorganic salts in the culture medium. Stress with high and

low oxygen tension, crowding, inorganic ion imbalance, or radiation is effective in inducing tumor formation. Retarded growth is associated with enhanced tumor induction, which suggests that certain degradation products may be involved.

The results to date suggest several basic approaches to the plant tumor problem. Kinetin is a structural analogue of adenine. Thus, in effect, the system may be expected to lead to a study of differentiation and morphogenesis by direct or indirect alteration of the genetic potential of treated plants. The localization and site of action are being investigated with tritiated kinetin.

The Mutagenic Action of Coliphage X-373

Infection of bacterial cells by temperate bacteriophages may lead to permanent or transient changes in the genotype of the resultant lysogenic hosts through the well-known phenomena of phage conversion and phage-mediated transduction. A third type of phage-promoted host modification is being investigated.

Coliphage X-373, discovered as a contaminant in a stock of *Escherichia coli* K-12, induces mutations to nutritional auxotrophy in phage-sensitive sub-strains of K-12 at a frequency generally 2000 times greater than that of spontaneous mutations in parallel uninfected cultures. In sensitive cultures exposed to high multiplicities of phage, roughly 15% of cells survive to produce lysogenic clones; about 40% of the latter contain large numbers of mutants. The ratio of mutant to wild-type cells in individual clones may vary from 0.01 to 1.0, which suggests that mutation takes place soon after infection, a view supported by the finding that additional mutations cannot be induced by superinfection of stable lysogenic strains. Moreover, the spontaneous mutability of lysogenic strains is no greater than that of nonlysogenic strains. Thus, the mutagenic activity of X-373 is considered a consequence of one or more phage functions operative in newly infected cells, but inoperative in lysogenic cells.

Mutations induced by X-373 appear to be completely undirected. Over 300 mutants have been identified; the majority are deficient for one or another of 15 amino acids; others have requirements for purines, pyrimidines, or vitamins. It is also possible to induce mutations resistant to the virulent phages T1, T4, T5, and T6, and to sodium azide, but not streptomycin. All mutants are lysogenic

for X-373, but lysogeny *per se* is not required for expression of the mutant phenotypes because the two characters are separable by recombination in Hfr $\times$ F⁻ crosses. The mutants seem to be extraordinarily stable. No instances of spontaneous or diethyl sulfate-induced reversion to the wild type have been detected, and it is tentatively concluded that X-373 creates extensive lesions in the host genome, i.e., deletions or multisite alterations, which accords with the finding that X-373 cannot induce reverse mutations in nonlysogenic auxotrophs. Moreover, phage X-373 appears to be devoid of transducing activity.

Preliminary attempts to map the prophage locus of X-373 have failed to demonstrate close linkage to known markers of *E. coli*; X-373, like phage P1, may have no specific chromosomal attachment site. Attempts to induce mutations with phage P1 and with another unidentified coli-Shigella temperate phage were unsuccessful. Thus, at the moment, mutagenic activity seems to be a unique property of phage X-373.

Control of Cysteine Biosynthesis in Bacteria

In continuing analyses of genetic properties in mutants of *Salmonella typhimurium*, ≈ 460 cysteine-requiring auxotrophs have been classified in terms of recombination, complementation, and growth responses. Five cysteine-controlling regions have been recognized: *cysA*, *B*, *C*, *E*, and *G*. Available evidence suggests that each is located in a different part of the genome. *A* and *B* each contain three complementation units, *C* five units, and *E* two units; *G* has not yet been analyzed. Every single-site mutant tested has been placed unambiguously in one of these complementation groups, which, together with the difference in nutritional requirement detected among mutants belonging to different groups of the same region, makes it probable that each complementation unit represents a separate gene locus. Biochemical studies carried on at the Johns Hopkins University support this view. Assuming every complementation unit to be a different locus, at least 14 gene loci grouped in 5 clusters are involved in control of cysteine biosynthesis.

Extensive studies are being carried out with 276 single-site markers and 94 deletions located in the *cysC* region. This region is believed to comprise a cluster of five gene loci, arranged and designated as follows: *cysC*, *D*, *H*, *I*, and *J*. Single-site mutations are distributed among them fairly evenly.

Loci *C* and *D* appear to be adjacent, as do *I* and *J*. A "genetically silent" section, equal to at least half the length of the whole *cysC* region, is located between *D* and *H*, and a much shorter similar section between *H* and *I*. None of the known single-site mutations is located within these sections.

Of the 94 deletions, 81 are similar in extent ("ditto" deletions), their left ends falling in the left half of the long silent section and their right ends in the section beyond the *J* locus. In addition, 8 of the remaining 13 deletions have one end located in the same silent section. The proposed explanation of these unique findings assumes that the long silent section includes a duplication of the section where the right ends of the ditto deletions are located, and that the deletions originate because duplicated sections frequently synapse, which results in formation of a loop within the gene string and consequent omission of the loop from the newly formed genome during replication. The unusual properties of this silent section can be explained by assuming that it is composed, at least in part, of uncoded DNA.

Fate of Transforming Deoxyribonucleic Acid in *Pneumococcus*

A hereditary trait may be conferred upon certain cells by treatment with DNA from others bearing the trait. Study of such genetic transformation in pneumococcus using P^{32} -labeled DNA from a streptomycin-resistant strain indicates that the initially double-stranded, highly polymerized material was converted about equally into low molecular weight fragments and a single-stranded form during or immediately following entry of the DNA into the recipient cells. Material from both fractions was incorporated into the nucleic acid of the host, the fragments by normal synthetic processes, the single-stranded form by direct integration. The latter process represents the mechanism by which genetic information is transferred to the recipient cell. In agreement with other studies, about one-fifth of the labeled material was found to be so integrated. The breakdown of DNA on entry explains previous observations – that the extent of phenotypic expression, e.g., enzyme synthesis, of the introduced material depends upon the amount genetically integrated rather than the amount introduced, and upon the prompt eclipse, following entry, of material active in transformation.

Cooperative Radiation Mutation Program

The radiation facilities at Brookhaven are being used to irradiate plant parts such as seeds, cuttings, and similar material for eventual study here and elsewhere. The objective of this work is to study the usefulness of such irradiations in producing mutational changes that represent possible crop improvements. Irradiations have been performed for many individuals and groups in this and other countries.

Dr. H.T. Chang of the Joint Commission on Rural Reconstruction, Plant Industry Division, has described three new strains of peanut and 16 new strains of soybean which show more resistance to diseases and higher yields than the original varieties. Dr. Glen W. Burton of the U.S.D.A., Crops Research Division, Georgia Coastal Plain Experiment Station, has reported on a type of dwarf mutant of Pensacola bahiagrass with seed stalks only about 6 in. long. If this dwarf characteristic can be transferred into a normal genetic background it may be possible to use this new bahiagrass as turf.

CELL PHYSIOLOGY

Nuclear Variables and Radiosensitivity in Plants

Continuing study is being made of the manner in which nuclear variables control the radiosensitivity of plants. Tolerance data on many new species have been obtained, and from these and previous findings a new correlation has been developed. The data show that increasing nuclear volume (v) increases sensitivity and increasing chromosome number (n) decreases sensitivity. This suggests that sensitivity should be related to v/n , the chromosome volume, in some predictable fashion. Data for 10 species are shown in Figure 1. The relationship is clearly highly significant and thus can be taken as a reliable index of radiosensitivity. This correlation, considered in conjunction with relationships established earlier between nuclear volume, chromosome number, and radiosensitivity, makes possible the prediction of expected tolerances of species that have not been irradiated. From cytological data on several hundred species of plants it is predicted that gymnosperms as a group are, for plants, extremely sensitive. Many species can be expected to have a LD_{50} between 300 and 600 r for an acute gamma exposure. For these species continuous exposures of 2 to 5 r/day

will cause serious effects on growth, fertility, or viability which will manifest themselves in from 1 to 5 years. Angiosperms, including deciduous trees, will in general be much more resistant. It can be safely predicted, therefore, that gymnosperms, which include many important economic plants, would be among the first plants to show severe radiation damage in case of major nuclear accidents or war. It also appears that many of these plants may be extremely useful in radiobiological studies.

Differential Radiosensitivity of the Shoot Meristem in *Taxus media*

In *Taxus media* the pattern of damage to the shoot meristem produced by chronic irradiation is in order of decreasing sensitivity as follows: apical initial layer, lateral initial layer, control mother cell zone, transition zone, axillary cells, and rib meristem. The reasons for this particular order are being sought. In these chronically irradiated trees the relatively modest dose of 175 r given at 3.5 r

per 20-hr day produces total disruption of the apex hairs. In addition, a regenerative process occurs which results in new, abnormal, but apparently operative meristems.

Changes in Deoxyribonucleic Acid Content and Volume of Nuclei as Factors Affecting Radiosensitivity

The nuclear volume of meristem cells in the soybean increases after initiation of germination to a maximum at 48 hr and later decreases. The DNA content begins to increase at 48 hr, reaches a peak at 72 hr, and then decreases. The sensitivity of the plant (defined by dry weight at 30 days) to

Figure 1. Relationship between chromosome volume at interphase and acute lethal dose of x or gamma radiation in plants. Similar relationships also obtain for other measures of radiosensitivity. The relationships are useful in predicting the radiosensitivity of other plants from consideration of nuclear variables. The plants used are (1) *Trillium grandiflorum* (Michx.) Salisb. (white trillium); (2) *Lilium longiflorum* Thumb. (white trumpet lily); (3) *Podophyllum peltatum* L. (May apple); (4) *Hyacinthus* c.v. "Innocence"; (5) *Chlorophytum elatum* Ker. (spider plant); (6) *Zea mays* L. (corn); (7) *Muscari botryoides* (L.) Mill. (grape hyacinth); (8) *Crepis capillaris* (L.) Wallr. (hawk's-beard); (9) *Lycopersicum esculentum* Mill. (tomato); (10) *Gladiolus* c.v. "Friendship."

Figure 2. Estimation of cell population kinetics by double labeling with thymidine. By supplying thymidine- H^3 and $-C^{14}$ at appropriate intervals during cell replication and subsequently locating and identifying the isotopes by autoradiography, the duration of phases of the nuclear cycle may be determined. The autoradiograph is of chromosomes of the root tip of *Tradescantia* treated with labeled thymidine. The two nuclei with photographic grains localized over the nuclear area were labeled with tritium only. The nucleus with a spray of grains around it was labeled only with C^{14} .

x rays given acutely is least at 24 hr after germination and greatest at 72 hr. Later the sensitivity decreases, but not to the level at 24 hr. The findings illustrate the direct relationship between DNA content and volume of nuclei and the sensitivity of plants to radiation.

Cell Population Kinetics and Chromosome Duplication

Study of the kinetics of the growth of cell populations is facilitated by a recently developed double-labeling procedure employing thymidine- C^{14} and thymidine- H^3 . The cell population is supplied with one form of thymidine and then, after an appropriate interval, with the other. The labeled thymidine, which is stably incorporated into the DNA of the nucleus, then is detected and identified with respect to kind of label by autoradiography. The ratios of C^{14} -labeled to H^3 -labeled cells found in several labeling sequences define the duration of various subdivisions of the cell cycle relative to DNA synthesis and flow rates of cells through parts of the cycle. By means of this procedure it has been found that DNA synthesis is relatively insensitive to major environmental alterations, while the periods before and after synthesis are affected markedly. For example,

continued exposure of *Tradescantia* root tips to 30 r of gamma rays per day lengthens the periods before and after synthesis to several times the normal duration, but has practically no effect on the period of DNA synthesis. Extremes of temperature have a similar effect.

By use of thymidine- H^3 and autoradiography it has been found in *Tradescantia* root tips that DNA synthesis, which is confined to a definite period within the nuclear cycle, proceeds at a fairly constant rate until near the end of the synthetic period. At that time the rate decreases. The decrease reflects localization of DNA replication to specific sites on the chromosome arms. Generally the terminal portions of the chromosome arms continue replication after other parts cease DNA production.

Transfer of Genetic Information

Amino acids are believed to be represented in nucleic acid language by specific trinucleotide sequences. The simplest possible genetical change or mutational event should involve replacement of one nucleotide by another in DNA. This would set off a chain of reactions: replacement of one nucleotide by another in a messenger ribonucleic acid (RNA) molecule; replacement of one amino

Figure 3. DNA synthesis during chromosome replication. The autoradiographs are of chromosomes of the root tip of *Tradescantia* treated with thymidine- H^3 during the mid-portion (a and b) and near the end (c) of DNA synthesis. In a and b the isotope is distributed throughout the chromosome arms; in c it is confined mainly to the ends of the chromosome arms.

acid by another in a protein. Although little is known about the sequence of nucleotides in nucleic acids, a great deal is known about the sequence of amino acids in some proteins. Based on interrelationships indicated by available amino acid replacement data, on the assumption that amino acids showing a high degree of mutual replaceability have the majority of their coding nucleotides in common, and other considerations, a partial solution to the RNA-protein code has been proposed.

The sequence of most amino acids in proteins is believed to be determined by a nucleotide coding mechanism. However, there are indications that the mechanism does not operate on all amino acids. For example, proline is hydroxylated after incorporation into the polypeptide chain. In this case, the final form of the amino acid is determined after its incorporation into the polypeptide chain. Clearly hydroxyproline has no mutual representation in nucleic acids. Instead, the polypeptide chain is coded for proline, and some is converted into hydroxyproline. Although convincing experimental evidence is lacking, separate coding has been assumed for glutamate and its amide. It is known that amino acids form specific complexes with soluble RNA molecules which serve as immediate precursors in protein synthesis. In extracts prepared from the bacterium *Escherichia coli*, it has been found that glutamine is incorporated directly into the soluble RNA, whereas glutamic acid is first converted to glutamine. This suggests that of the two amino acids only glutamine is directly incorporated into the protein, and that the glutamic acid residues arise by hydrolysis of the glutamine residues.

Ribosomal particles exist in various states of aggregation. In *Escherichia coli* there are two fundamental ribonucleoprotein particles, a 50 S and a 30 S (ribonucleoprotein particles are named according to their approximate sedimentation constants; S refers to the Svedberg constant). These may combine to form a 70 S component which, in turn, may dimerize to form a 100 S component. It has been found that optimum amino acid incorporation *in vitro* requires the presence of both the 50 S and the 30 S ribosomal subunits, but not the 100 S ribosome. These results, combined with the findings of others, suggest that the 70 S ribosome, and only the 70 S ribosome, is essential for protein synthesis.

Synthesis of Messenger Ribonucleic Acid

The mechanism for transfer of genetic information from DNA to messenger RNA is obscure. Theoretical considerations suggest the formation, by hydrogen bonding, of a triple helical intermediate composed of double helical DNA as template, plus a newly synthesized single strand of RNA. Formation of a triple-strand intermediate would require minimum configurational change in DNA and allow ready dissociation of the RNA after synthesis, in keeping with the speed and selectivity with which DNA in the organism responds to inducers and repressors. Reasonable hydrogen bonding, bond lengths, and bond angles exist for two molecular models of a triple helical intermediate. The nucleotide-base ratios for the DNA and RNA in one model correspond to findings in animal nuclei. The ratios in the other model conform to findings in bacteria. At present, conflicting data are available to support either the triple-strand intermediate or the alternative double-strand complex proposed earlier.

Role of Nucleic Acids in the Formation of Chloroplasts

In green algae and plants, photosynthesis (the light-energized conversion of carbon dioxide into sugars and other compounds) is localized in cellular organelles known as chloroplasts. The formation of chloroplasts is also light dependent. Cells protected from light form plastid precursors, the proplastids. Upon exposure to light the proplastids grow and mature as chloroplasts. Since new proteins are synthesized during growth of chloroplasts, nucleic acids are expected to play an important role in the process. To determine whether new specific nucleic acids are required for plastid synthesis, inhibitors of nucleic acid synthesis were tested for effects on chloroplast development. *Euglena* cells were grown in the dark and then illuminated. The formation of chlorophyll, which is correlated with the appearance of specific chloroplastic proteins, during exposure to light was taken as an index of chloroplast development. 5-Fluorouracil, an inhibitor of DNA and RNA synthesis, at concentrations of 5 to 50×10^{-4} M, depressed chlorophyll formation. Equivalent concentrations of uracil but not thymine prevented the depression. Also, 5-fluorodeoxyuridine, an inhibitor of DNA synthesis, had no effect. Thus, the

Figure 4. Electron micrograph of *Anacystis nidulans*. In this photosynthetic bacterium the chlorophyll appears confined to the triple-layered lamellae visible in (1). Each lamella is separated from the next by ≈ 500 Å. As deduced from (2), the interlamellar space contains the pigment phycocyanin. Light absorbed by the phycocyanin presumably can migrate to either bounding lamella, which offers an explanation for the increased photosynthetic efficiency of light absorbed by this pigment.

findings indicate that synthesis of RNA rather than DNA is necessary for chloroplast development. Addition of the fluorouracil to the cells after chlorophyll formation has proceeded to $\approx 10\%$ of the maximum had little effect on subsequent pigment production. The effect of the fluorouracil was not prevented by delayed addition of uracil to illuminated cells. If, however, the cells were kept in the dark for one day before addition of uracil, chlorophyll synthesis in the light was only slightly depressed. These observations suggest not only that formation of chloroplasts requires RNA synthesis but also that the synthesis is initiated by light.

Structural Basis of Photosynthesis

Study of photosynthetic bacteria has indicated the level of structural organization needed to support the fundamental phenomena of photosynthesis (capture of light energy and transformation into chemical energy) to be much lower than hitherto supposed. However, the bacterial systems lack certain characteristics of photosynthesis in higher plants, such as photosynthetic oxygen evolution and pronounced effects on photosynthetic efficiency due to illumination with light of different colors. The unicellular blue-green alga, *Anacystis nidulans*,

is an obligate phototroph with high photosynthetic activity that possesses these characteristics; the rate of oxygen evolution is high, and light absorbed by the blue protein, C-phycocyanin, is more efficient in producing oxygen evolution than light absorbed by the primary photosynthetic pigment, chlorophyll *a*. When this microorganism is broken, the phycocyanin and the chlorophyll are recovered in different fractions. The chlorophyll sediments readily under centrifugation, whereas the blue protein does not. The chlorophyll-containing fraction consists of lamellae that correspond to the triple-layered lamellae observed in electron micrographs of the organism. These lamellae exhibited an unusual order; each is separated from the next by a distance of ≈ 500 Å. When procedures that reveal the phycocyanin in electron micrographs are employed, this 500-Å interlamellar region exhibits rather high electron density. It appears, therefore, that the phycocyanin is present in very high concentration between the chlorophyll-containing lamellae.

The 500-Å separation between lamellae is too great for significant energy transfer between an excited chlorophyll molecule in one lamella and another chlorophyll molecule in an adjacent lamella. Since the phycocyanin is interlamellar, it has been proposed that this structural organization can account for a higher photosynthetic effi-

ciency with light absorbed by phycocyanin than chlorophyll. Light energy absorbed by phycocyanin can migrate to either bounding lamella, whereas light absorbed by chlorophyll must be utilized in that lamella. According to this concept, the enhanced efficiency is the result not of events peculiar to the phycocyanin, but rather of a second-order effect due to structural organization of a rather high level.

Electron Transfer in Photosynthesis

The changes in light absorption of anaerobic suspensions of the green photosynthetic bacteria *Chlorobium thiosulfatophilum*, strain L, and *Chloropseudomonas ethylicum*, strain 2K, within the spectral interval 380 to 570 mμ during irradiation with monochromatic far-red light which is absorbed by

Figure 5. Change in light absorption spectrum in *Chlorobium thiosulfatophilum* L caused by irradiation with 800-mμ light. The change is attributable to oxidation of *c*-type cytochrome(s). Measurements were made with a double-beam spectrophotometer of the Chance type.

Figure 6. Absolute spectra of a preparation of chlorophyll-770 from (A) *Chlorobium thiosulfatophilum*, strain L, and (B) *Chloropseudomonas ethylicum*, strain 2K. The height of the red peak has arbitrarily been made equal to 1.0 for each curve. The solvents were tris buffer of pH 7.5 (---), 80% methanol (—), and wet ether (—). The pigment is involved in the absorption of light by these photosynthetic bacteria and subsequent oxidation of cytochrome.

the photosynthetic pigments indicates oxidation of *c*-type cytochrome(s) in these bacteria, a response previously observed in purple bacteria. Two wavelengths of far-red actinic light have been tested. Light of 725 m μ , which is within the main absorption band of chlorobium chlorophyll, appears to give the same effect as light of 800 m μ , which corresponds to a minor absorption band on the red side of the main band.

During an attempt to isolate the cytochromes from the green bacteria, the pigment responsible for the minor absorption band in the far red was separated from the major chlorophyll component. The far-red pigment remains in solution after centrifugation of crude cell extracts at 140,000 g for 90 min and is precipitable from the solution by ammonium sulfate. The spectrum of this pigment is completely different from that of the main chlorobium chlorophyll. It resembles bacteriochlorophyll spectroscopically, but is tentatively designated chlorophyll-770 by virtue of the position of the red peak in ether. It is planned to test the hypothesis that sensitization of the light-induced cytochrome oxidation involves transfer of electronic excitation energy from chlorobium chlorophyll to chlorophyll-770 within intact bacteria.

BIOCHEMISTRY

Enzyme Changes in Barley Leaves After Infection With Powdery Mildew

The increased respiration of plant cells characteristically accompanying infection by a wide range of pathogens continues to attract attention. In green barley leaves respiration is increased up to threefold after infection with *Erysiphe graminis* var. *hordei*. In contrast, the respiration of infected etiolated leaves changes but little. To gain understanding of the nature of the increased respiration, enzymic activities of noninfected and infected green and etiolated leaves were measured. In green leaves at ≈ 96 hr after inoculation concomitant increases occurred in respiration and in the activity of glucose-6-phosphate dehydrogenase, 6-phosphogluconate dehydrogenase, and enolase. However, the activities of isocitrate dehydrogenase, malate dehydrogenase, aconitase, and DPNH-cytochrome *c* reductase changed insignificantly. In etiolated leaves, on the other hand, there was no appreciable increase in respiration or enzyme activity. The increase in respiration thus ap-

pears to be due in part to an increase in the activity of the hexose monophosphate shunt system. Other experiments indicate that the increase in respiration may be related to a decrease in photosynthesis.

Protein Structure

Bovine pancreatic ribonuclease A is at present one of the most thoroughly investigated enzymes known: its covalent structure is defined, and extensive knowledge of its structural configuration is at hand. Knowledge of the relation between its structure and function has been provided by two types of investigation: modification studies, and the examination of biological variation. Among the former, substitution studies with the alkylating agent 2,4-dinitrofluorobenzene have been continued, and it is now possible to report the isolation and proof of structure of essentially homogeneous samples of the following lysine-substituted derivatives of ribonuclease A: (α)-1-dinitrophenylribonuclease; (α)-1,41-bis-dinitrophenylribonuclease; and 7,41-bis-dinitrophenylribonuclease. Quantitative studies have indicated that the latter derivative may be formed as a consequence of initial substitution at position 41 (affording the well-characterized 41-dinitrophenylribonuclease) with concomitant exposure of the lysine residue at position 7 and subsequent reaction at that position. The result provides direct proof that chemical modification at one side-chain in a protein molecule can alter the reactivity of a second side-chain in a remote position of the amino acid sequence.

Two types of biological variation have been investigated, the isozymes of bovine ribonuclease, and porcine ribonuclease. Extensive studies with bovine pancreatic zymogen granules have shown that multiple forms of ribonuclease are elaborated by the pancreas. The most prevalent are ribonuclease B and ribonuclease C. Ribonuclease B has been isolated from bovine pancreatic juice in highly purified form. It has been found to be a naturally occurring modified form of ribonuclease A; the modification is unprecedented and has been found to be due to the attachment in an as yet undefined manner of an oligosaccharide moiety composed of hexose and hexosamine to the protein molecule.

Mechanism of Muscular Contraction

The mechanism of muscular contraction is a problem of fundamental importance because it

represents a way of converting chemical energy immediately into mechanical work, and in man alone many diseases are directly related to failure of this basic function. To gain understanding of this process a continuing study is under way. The essential proteins, myosin and actin, which catalyze the decomposition of certain chemical compounds and are themselves shortened during muscular contraction have been identified. The chemical compound adenosine triphosphate (ATP), which provides the energy for the contraction, and its decomposition products adenosine diphosphate (ADP) and inorganic phosphate (P) are also known. However, the intermediate mechanism by which the energy of ATP is transferred to the protein structure is unknown. It has long been presumed that some intermediate complex must exist, but evidence for such a complex has been obtained only recently by means of experiments using water labeled with O^{18} . When ATP is split by water, the entering oxygen atom could finally appear in either ADP or P. In the presence of muscle proteins, which catalyze the hydrolysis of ATP, the O^{18} from the labeled water is intro-

duced only into the P. Thus, formation of a phosphoryl protein intermediate may be presumed. Prediction is that only one oxygen atom should be introduced; however, three appear instead of one. The extra oxygen atoms must be introduced at some point during the hydrolysis. Since the O^{18} of the water does not exchange with the oxygen of unhydrolyzed ATP or added P, the introduction of the extra oxygen into P derived enzymatically from ATP must occur at some intermediate stage in the hydrolysis.

The metal ion dependence and specificity with respect to nucleotide in the exchange reaction and over-all process of muscle contraction are similar. If indeed a phosphorylated protein explains the exchange reaction, it may also explain the change in shape of the myosin and actomyosin during contraction. The presumed new form of the proteins may well have a shorter structure than the original. Hence, the energy of ATP may lead to mechanical shortening by stabilizing a protein molecule in a geometric configuration differing from some original configuration.

Hemoglobin Synthesis

Various processes involved in hemoglobin synthesis by erythrocytes taken from the duck before and after x-irradiation and other treatment are being measured to determine whether radiation affects the processes directly or indirectly through inhibition of erythropoiesis. At present the latter action of radiation appears to be sufficient to explain the changes observed after irradiation of the animal. These changes – decreases in uptake of iron by the cells and conversion of this iron into heme, and decreases in formation of porphyrin, heme, and globin – presumably reflect maturation or aging of the erythrocyte. Inferences from the findings are that heme and globin formation are not connected in obligatory fashion, and that heme formation does not depend upon the uptake of iron by the cells, but is limited by porphyrin formation. With respect to time and degree, the iron uptake exhibits the most consistent pattern of change following the irradiation.

Figure 7. The formal representation of the simplest possible living system in which a spontaneous flow of electric current can occur. The broken lines indicate three equi-electrochemical potential surfaces (*eecp*). Note that an *eecp* surface is orthogonal to the lines of electric current flow. The system is represented in linear form as

external phase, membrane 1, internal phase,
membrane 2, external phase,

and has an emf expressed by

$$E = \frac{1}{F} \sum_j (t_j^1 - t_j^2) \Delta \mu_{i,j}$$

where the t 's are ionic transport members and $\Delta \mu_{i,j}$ is the chemical potential difference.

BIOPHYSICS

Origin of Electric Currents in Living Systems

The origin of electric currents in living systems has always been something of a mystery. It is possi-

ble however to give a general analysis of closed circuit systems in which spontaneous electric currents circulate. In terms of this analysis any isothermal closed circuit system that contains at least two barriers of different sets of ionic transport numbers interposed between homogeneous phases of different ionic compositions has an electromotive force. When such a circuit is opened through one of the homogeneous phases, the system in linear array is terminated by identical terminal phases, and an experimentally determinable emf appears between the identical terminal phases. Thus the simplest possible isothermal system exhibiting an emf is

phase α , barrier 1, phase β , barrier 2, phase α' , and its emf is given by

$$E = \frac{1}{F} \sum_j (t_j^1 - t_j^2) \Delta\mu_{ij}.$$

In this equation E represents the emf, F is the Faraday constant, t_j^1 and t_j^2 are the ionic transport numbers of the j th ion in the two barriers, and $\Delta\mu_{ij}$ is the chemical potential difference of an ion pair between phase β and phase α . The indicated summation is to be taken over all ions to which the barriers are individually permeable.

In the simplest possible model of a living cell the barrier becomes the limiting membrane of the cell. For a spontaneous electric current to be generated by the living cell, the limiting membrane of the cell must have at least two regions with different sets of ionic transport numbers. The electric current when it exists consists of anions and cations which circulate in opposite direction. There are no electrodes and hence no electrode reactions in living systems, and the energy to maintain the electric current is derived from some fraction of the free energy of dilution of interior phase with the external phase. When the cell is in a basal state, technically a stationary state, the emf in the system is related to the rate of metabolism, since μ_{ij} is then given by the flow of nutriment and end products through the limiting membrane of the cell. If the cell is not in a basal state then no simple relation exists between the emf of the biological source and the rate of metabolism.

Biological Effects of Cosmic Rays

Observations made several years ago on mice taken in balloons to the top of the atmosphere indicated that heavy cosmic-ray particles cause

Figure 8. Photograph of mouse irradiated with an 800-rad dose of deuterons in two places through a slit aperture 0.250 mm in width in the form of a cross. The photograph, taken 101 days after irradiation, shows graying only in the irradiated area.

Figure 9. Entrance to facility used to study ecological effects of radiation.

Figure 10. Radiation source in gamma forest. The source is maintained above ground 20 hr/day. The dose rate ranges from several thousand roentgens within a few meters to ≈ 2 r at 125 meters. The pine tree in the foreground was killed by radiation.

Figure 11. Predicted effects of chronic irradiation on a Long Island oak-pine forest. The radiation source is considered to be at the center and to deliver several thousand roentgens per day at a few meters. Prediction is based on the nuclear volumes and chromosome numbers of the species irradiated. The prediction will be compared with observation in a similar, actual forest.

graying of the hair, from which it has been inferred that these particles might have serious effects on man in such organs as the brain and the eye. Since no accelerators can produce such high energy particles, the problem was attacked experimentally by simulating a "thin-down" track from such a particle with a microbeam of deuterons 25 μ in diameter. Since an individual hair follicle is difficult to identify over long periods, for this part of the study the microbeam was formed as crossed slits 25- μ or 250- μ wide. This pattern could be identified on the mouse and showed that the graying response occurs with both slit widths at doses expected in cosmic-ray tracks. However, for both the brain and the eye, doses enormously higher than this are required to produce damage. It is concluded that graying of the hair may occur in astronauts, but this is probably the only damage that will result from cosmic-ray particles.

ENVIRONMENTAL BIOLOGY

Radiation Ecology

The interests of the Biology Department have been extended this year to include research in environmental biology, with emphasis on the effects of ionizing radiations on ecological systems and their components. The program extends previous work at the cellular and organismal levels to the population and community levels in an effort to

define more precisely the importance of ionizing radiation as an environmental factor. For this work a 9500-curie source of gamma radiation has been established in the center of a carefully selected forest stand on the Laboratory site. This source, which is exposed 20 hr/day, has been in operation since November 1961; it delivers several thousand roentgens per day within a few meters of the source and ≈ 2 r/day at 125 meters.

Several complementary research projects carried out in cooperation with other institutions are now under way as part of this program. Most of these projects utilize the new radiation facility and contribute to quantitative estimates of radiation effects on forest populations and to estimates of changes in rates of energy fixation and movement within the experimental ecological system.

A prediction of effects of the first year of irradiation on the populations of the various woody species was made on the basis of nuclear volume and chromosome number. Preliminary observations during the spring confirm the relative sensitivities indicated by this prediction. An ultimate goal of the program is development of the ability to predict quantitatively radiation effects on organisms, populations, and communities. The preliminary results of the first winter's irradiation lend new strong support to the hypothesis that nuclear volume and chromosome number alone provide a powerful basis for such a prediction.

Medical Research

The research program of the Medical Department, based on studies contributing to better understanding of biological processes in man, is aimed at increasing the knowledge of the effects of ionizing radiations upon the human being. Some clinical and laboratory investigations are directed toward enhancing beneficial applications of these radiations or improving measures for preventing or counteracting detrimental effects. Other investigations are centered on the biological processes of man in order that through the knowledge of normal function elucidations of disease states may be furthered, and improved methods of diagnosis and therapy may be developed. These approaches are complementary, for, whether at the level of the single cell or the entire body, there appear to be no effects of ionizing radiation that have not been observed in naturally occurring diseases or produced by other toxic agents. Thus, the study of radiation effects contributes to the understanding of diseases, and vice versa, and the pace of progress in both is controlled by the advances in knowledge of normal biological processes.

The diversified talents and extensive facilities uniquely concentrated at Brookhaven National Laboratory provide unusual advantages for pursuing these objectives. The broadly based program of the Department incorporates the experience and skills not only of those devoted to research in the field of medicine, but of many from the various disciplines in other departments of the Laboratory. Similarly, not only are the facilities of the hospital, laboratories, and reactor of the Medical Research Center utilized, but the instrumentation and special laboratories of other departments and the large nuclear devices may be used as well. Thus, the program is enriched through the productive interchange of ideas, information, facilities, and services, and the opportunities within its framework are limited only by the scope of the vision and interest of the individual investigator.

Specific examples of research under way are given in the following pages. The report is not comprehensive, and the reader may obtain additional information from publications of the Department.

EXTRACORPOREAL IRRADIATION OF THE CIRCULATING BLOOD OF MAMMALS

Radiation injury of the whole body confounds the interpretation of data collected on changes in the blood, since cells within the blood and the tissues that form new blood cells are injured simultaneously. It was reasoned that irradiation of the flowing blood without injury to the tissues that form blood cells would perform the dual service of assisting in the interpretation of changes observed after whole-body irradiation and providing a tool for study of the life span of blood cells and the regulation of new production. Surgical pumping and irradiation techniques developed in 1960 made possible the present studies. Preliminary studies performed upon various species resulted in the selection of the calf as the most desirable experimental animal. The current experimental setup is illustrated in Figure 1. Figure 2 shows the production of a pronounced depression of the blood lymphocytes, and Figure 3 presents an example of the slow recovery from the lymphocyte depression.

The method has opened up new avenues of attack on the ubiquitous lymphocyte with respect to quantification of its production rate, classification of lymphocytes by life span, and determination of the site of removal of aged or injured lymphocytes. Similar studies to be made on leukemic cattle will make it possible to determine the differences between normal and leukemic lymphocytes with respect to these parameters. It is hoped that these studies and perfection of the technique will lead to similar studies on human beings with leukemia.

EFFECT OF RADIATIONS OF DIFFERENT LINEAR ENERGY TRANSFER

In the past, evaluation of fast neutron effects usually has been handicapped by lack of neutron sources yielding narrow energy spectra. Apparatus recently constructed at this Laboratory now permits the exposure of mice to neutrons of selected discrete energies by locating groups of animals in sites at different angles to the beam-target axis of

a 3-Mev Van de Graaff generator employed to accelerate protons into a tritium target. Special equipment developed at Columbia University also is available to measure both depth-dose and quality of radiation, the latter by employing graded sizes of chambers representing microscopic volumes of tissue-equivalent material.

Initial studies of spleen-thymus weight loss indicated relative biological effectiveness (rbe) values ranging from ≈ 4.5 at a neutron energy of 0.43 Mev to 3.0 at 1.8 Mev (250-kvp x rays were used as the comparison radiation). Although slightly higher values were obtained for sperma-

togonia depletion, and slightly lower values for incorporation of iododeoxyuridine (a precursor of deoxyribonucleic acid), both showed the same relationship of rbe to neutron energy as that observed with spleen and thymus. Spermatogonia depletion after exposure to 14-Mev neutrons is being evaluated, and equipment has been built for similar study of the 2 to 6-Mev energy range.

The use of x radiation for comparison in the above work necessitated consideration of the quantitative effect of dose rate. An analysis of the data, including published rodent $LD_{50/30}$ dose-rate studies, yielded an expression indicating propor-

Figure 1. Extracorporeal irradiation of circulating blood of calf. Upper left, schematic diagram of experimental setup. Upper right, recording thermistors, pump, and water bath for heat exchanger. Lower left, calf in stanchion with blood flowing at 300 ml/min. Lower right, irradiation coil (1100-ml internal volume) receiving 300 rad/min.

tionality between biological effectiveness and the cube root of dose rate. Data from a study of human skin dose-threshold fitted this expression. Further, all sets of data indicated the same fractional change in 50% effect dose between any two given dose rates.

PATHOLOGICAL AND RADIOLOGICAL OBSERVATIONS RESULTING FROM EXPERIMENTAL USE OF HIGH ENERGY CHARGED PARTICLES

In 1946 Wilson published a discussion of the physical properties of fast protons and the concept that biological effects in tissue near the end of the range of the Bragg curve should be considerably enhanced by use of these particles because of greater specific ionization. Subsequently, Tobias and colleagues (1952 and 1958) and Falkner and associates (1959) reported their experimental observations on animals (rabbit ears with transplanted tumors and rat pituitary) and man (pituitary) following exposure to deuterons of 190 Mev,

protons of 340 Mev, and protons of 185 Mev. These investigators believe that both high energy deuterons and protons have future possibilities for precise irradiation of both deep-seated and superficial tumors.

The availability of the BNL 60-in. cyclotron and the proton synchrotron (Cosmotron) have made it possible to initiate a joint study with Drs. Baker and Moore of the Physics Department. Mice are being used in studies of (1) the histopathological effects of irradiation (penetration not exceeding 2 mm) with 20 to 40-Mev protons, deuterons, and alpha particles; (2) the effects of whole-body irradiation with 3-Bev protons; and (3) the indirect or "abscopal" effects of radiation delivered to the abdominal region. About 5000 mice have been irradiated to date. Irradiation of the heads of rabbits at an energy of 3 Bev is being investigated with respect to detectable histochemical alterations in the oxidative pathways in the brain. The Bragg effect of 50 to 90-Mev protons on the rectum and kidney of the dog is being established. Therapeutic effects of these various

Figure 2. Immediate production of depression of lymphocytes by extracorporeal irradiation of the blood for 12 hr.

Figure 3. An example of the prolonged lymphocyte depression produced by extracorporeal irradiation of the blood. Apparent recovery period, 64 days.

high energy protons are being investigated with reference to both the target (e.g., transplanted neoplasms in mice) and the host.

MEDICAL RESEARCH REACTOR STUDIES

The Medical Research Reactor (MRR) provides an intense source of neutrons and other forms of radiation that are useful in a variety of experiments.

Early in the year two patients with glioblastoma multiforme brain tumors were treated by neutron capture therapy; this completed a series begun during the previous year. In conjunction with the neutron capture therapy program and to broaden the radiobiological basis for the procedure, a number of experiments have been conducted in animals on aspects of the problem that cannot be effectively studied in patients. In mice, tumor implants were irradiated to determine the optimal time of exposure. In dogs the acute effects of high doses of thermal neutrons and of particles from the $B^{10}(n,\alpha)Li^7$ reaction on the brain and the effects of whole-body irradiation were studied. With pigs, the effects of fractionation of neutron dose on the skin were studied. In tissue cultures, HeLa cells have been used in efforts to determine autoradiographically the intracellular distribution of boron and to study the effects of heavy particle irradiation in a relatively simple cell system. Other animal experiments have included investigation of the effects of neutron irradiation on the eyes of rabbits, the retinal effects being studied by micro-electrode electroretinographic, biochemical, histochemical, and autoradiographic methods.

Work in mixed field dosimetry has included experiments on small graphite CO_2 -filled ionization chambers for measuring internal gamma-ray doses in animal studies. A gamma-insensitive continuous thermal neutron monitor that reads out the integral dose directly has been developed. Neutron spectrum studies have been carried out by using fission threshold foils and fission chambers (solid state detectors). Investigators from the Physics Department have used the MRR neutron beam in studies involving the fission of uranium.

DIFFERENTIAL MULTIDETECTOR

The use of radioactively labeled substances that concentrate in neoplasms is being investigated in

the development of diagnostic procedures. The localization of positron emitters depends upon the principle that annihilation of the positron produces two gamma rays which are emitted in opposite directions.

In the positron scanning method currently in use, only two detectors are used, a coincidence count between the two detectors being recorded as an event on the line connecting the detectors. The use of multiple detectors, with coincidence counts being recorded not only between directly opposing pairs but between all pairs, offers the possibility of shortening the counting time and of providing three-dimensional localization. The problems of data storage and data reduction are, however, much greater than when a single pair of detectors is used.

A working model using 32 one-inch-diameter crystals was assembled, the detectors being distributed in a circle (see Figure 4). Empirical tests using a point source at various points within a paraffin phantom and using a 2000-channel analyzer to store the data indicated that memory patterns could be correlated with tumor locations. Translation of the memory patterns to activity distribution patterns, however, requires processing by a large computer.

It is expected that clinical applications in the locating of tumors can be initiated with the existing 32 detectors.

MEDICAL STUDIES OF THE PEOPLE OF THE MARSHALL ISLANDS

Under the direction of the Medical Department of Brookhaven National Laboratory, a team of physicians and technicians from the United States and from the Trust Territory of the Pacific Islands conducts annual medical surveys of the Marshallese people accidentally exposed to fallout in 1954. Some 239 Marshallese people were exposed to sublethal doses of radiation with significant hematological effects. Many of them sustained beta burns of the skin and internal exposure to absorbed radionuclides. Considerable knowledge has been gained from these studies of acute and long-term effects of fallout on human beings.

The 1962 survey (8 years post exposure) was carried out in March at Rongelap Island, and the results are being processed. The 1961 survey report (7 years post exposure) indicates, in brief, that

the incidence of illness and disease was about the same in the exposed population as in the unexposed comparison population. Growth and development studies in children revealed a distinct tendency toward a slight lag in growth in exposed boys (but not girls) of ages <12 . Increasing frequency of pigmented lesions in skin areas previously involved with beta burns was found, but no evidence of malignant change was noted. Peripheral blood elements (particularly platelets) continued to show slightly lower levels in the more heavily exposed Rongelap group. No leukemia or other type of malignancy was detected, and no increased aging or similar late effect of radiation was seen in the exposed population.

Determination of body burdens of radionuclides was carried out by *in vivo* whole-body counting techniques with use of an 8-in. crystal in a "portable" 21-ton steel room and a transistorized 256-channel pulse-height analyzer, situated on board the survey ship, and by radiochemical analyses of urine samples. The IBM 704 computer was programmed to perform the analysis of complex gamma spectra obtained. No significant differences were noted between the body burdens of the people exposed to fallout in 1954 and those of the comparison population living in the same low-level radiation environment for the past four years. The Cs^{137} level ($14.7 \mu\text{C/kg}$) was about the same as that measured two years ago; the Zn^{65} level ($1.51 \mu\text{C/kg}$) had dropped to 17% of the mean level of two years ago; Co^{60} , in small amounts, was detected for the first time. The mean urinary excretion of Sr^{90} was $7.2 \mu\text{C/l}$, or 14% higher than that measured in 1959.

THE KINETICS OF RADIONUCLIDES IN ANIMALS AND MAN

The kinetics of the uptake and turnover of strontium in young rats were studied. The rate of calcium accretion and exchange capacity of various bones of the rat was determined following irradiation. Both these indices of bone metabolism were found to differ in irradiated rats from those in normal, aging rats. An attempt was made to alter the accretion and exchange in rat bones by dietary means. Stable strontium administered orally was found to decrease the accretion rate of Sr^{85} by bone, but it had no effect on the exchange or release of bone-fixed Sr^{85} .

Figure 4. Pilot model of differential multidetector for locating tumors by means of positron emitters.

Sr^{85} retention was measured in a number of patients with the whole-body counter. Retention over a one-year period could be described equally well by a series of exponential functions or by a power function, but it is not possible to extrapolate forward a series of exponentials. If the data are fitted with a power function and projected to the life span of man, then the biological half-life of the "equivalent" exponential, fitted to the curve from 20 to 40 years, is 110 years. Other studies attempted unsuccessfully to increase the turnover of bone-fixed Sr^{85} in patients by supplemental feeding of stable calcium and strontium salts.

Other long-term studies carried out with the use of the whole-body counter measured the retention of Cs^{137} , Zn^{65} , and Sc^{46} over the period of a year in a number of patients. Biological half-life values obtained were 52 to 114 days for Cs^{137} , 340 days for Zn^{65} , and >4 years for Sc^{46} .

In another group of patients studied over a 30-day period, the short-term kinetics of Sr^{85} and Ca^{47} were characterized in terms of accretion, exchange, and biological loss from bone (resorption and long-term exchange). A four-compartment

electronic model was developed with an analogue computer. The data suggest that identical roles in the skeletal metabolism for calcium and strontium over long periods of time cannot be postulated on the basis of the metabolic similarities observed in the short interval following injection of the tracers.

SKELETAL CELL PHYSIOLOGY

In order to study the effects of irradiation on the cellular complement of the skeletal system and the possibility of radiation-induced aging, it was first necessary to determine the changes that take place in the proliferative potential of cells during growth, aging, and the repair of trauma. Tritiated thymidine, a DNA precursor that is taken up by cells during DNA synthesis, was used throughout these studies. Salient features of bone growth development, repair, and aging uncovered during the latest in this series of investigations are as follows: (1) The proliferative potential of skeletal cells is highest at birth and within 8 wk diminishes to a very low level. (2) The role of the periosteum in bone formation has been extended to include participation in peripheral expansion of the epiphyseal plate and spherical growth of the ends of long bones (see Figure 5). (3) Aged osteogenic cells do not appear to lose their ability to undergo mitosis in spite of the considerable morphological change that takes place with increasing age. In addition, they can react to trauma within the same time period as that required for young cells to react. (4) A quantitative, rather than qualitative, change takes place in skeletal cells during the course of aging. (5) The initial proliferative response seen beyond 8 hr following trauma is a generalized reaction which returns to a normal level within 5 days. This basic response is not altered with increasing age and cytomorphological changes.

EFFECTS OF RADIATION ON BONE GROWTH OF RATS

Changes have been noted in the growth and development of children exposed to radiation resulting from atomic detonation. Since the doses received were lower than would be expected to cause direct effects on bone growth, studies were designed to investigate in animals possible indirect or "abscopal" action of radiation on bone growth. Tibial bone growth in rats was determined by

measuring x-ray pictures and noting changes in tibial length after exposure to different doses of x radiation. A special rat-holding cage was devised to permit roentgenography without anesthesia.

Direct and indirect effects of various doses of x radiation (200 to 800 r) have been measured at various times after exposure by comparing tibial growth rate in the exposed leg with that in the opposite leg, which was shielded with lead during exposure. Results on several hundred rats indicate that radiation-induced inhibition of bone growth at the various doses is due to indirect effects as well as to direct effects on the bone. "Abscopal" inhibition effects of exposure were found to be from 2 to 4 times greater than the direct effects on the bone. Both effects appeared to be dose dependent. Pair-feeding experiments showed that $\approx 65\%$ of the indirect effect was apparently nutritional due to the reduced food consumption associated with radiation exposure. Evaluation of partial-body shielding and other experiments are under way to determine the organ systems that might be responsible for "abscopal" effects other than those caused by reduced food consumption.

IMMUNE MECHANISMS IN HOMOLOGOUS AND HETEROLOGOUS (WASTING) DISEASE: THEIR SIGNIFICANCE IN POSTRADIATION TREATMENT

Postradiation transfusion of mouse and rat bone marrow protects recipient mice against lethal doses of whole-body radiation, yet recovery of the hematopoietic system of the irradiated host is frequently followed by wasting disease resulting in death during the second month after irradiation. It has been postulated that wasting disease and delayed deaths result from an immunological response of the donor marrow against host tissues. Experiments at Brookhaven have dealt with testing the capacity of transfused marrow cells in recipient mice to produce antibody to tetanus toxoid. The appearance of tetanus antitoxin in the serum of the host indicates antibody formation by immunologically competent donor cells. Antibody formation was obtained when isologous (same strain) and homologous (same species) mouse bone marrow was transplanted to recipient mice. Also, antitoxin formation was elicited from cell preparations obtained from spleen, thymus, and lymph-node tissues. Minimal titers of antibody were produced by spleen and thymus cell preparations.

SKELETAL CELL PHYSIOLOGY

OBJECTIVE:

To study the contribution of periosteal proliferation during growth and repair of long bones.

METHOD:

Autoradiographic analysis using tritiated thymidine.

RESULTS:

I. Normal bone

Periosteum contributes cells
to all growth parameters.

II. Traumatized bone

AUTORADIOGRAM
(YOUNG CELLS)

The biological response to fractures is
not altered by aging except quantitatively.

Figure 5.

The highest titers of antibody were elicited from bone marrow cells in mouse-to-mouse donor-host combinations. It has not been possible to demonstrate antibody formation by heterologous rat bone marrow in irradiated mice. The immunological competency of the donor rat marrow was limited in heterologous rat-to-mouse combinations.

STUDIES ON SYNTHESIS AND METABOLISM OF NUCLEIC ACIDS (RNA AND DNA) IN MAMMALIAN CELLS IN CULTURE

These studies are directed toward the understanding of the synthesis, metabolism, and function of nucleic acids in mammalian cells. Several cell lines, particularly the human cancer cell HeLa S3, are maintained in culture yielding a controlled homogeneous system. Tritium-labeled pyrimidine nucleosides used as precursors for nucleic acids are localized autoradiographically within subcellular structures. Nucleic acid fractions isolated chemically from the cells are analyzed quantitatively by spectrophotometry, and the amount of isotope is determined by liquid scintillation counting. The combination of methods yields information as to the site and time course of RNA synthesis, its time relationship to DNA synthesis and mitosis, and its metabolic interrelationship with nucleotides of low molecular weight and with DNA. By utilizing in addition the ability of certain cell lines in culture to produce specific proteins in the form of antibodies, functional characteristics of the nucleic acids in the interplay between genetic code in DNA and cell-specific proteins can be studied. The results obtained serve as references in the study of the effects of chemical and physical influences, such as irradiation, enzymes, toxic agents, and drugs.

H^3 -cytidine, H^3 -uridine and H^3 -thymidine were used in HeLa S3 cells to establish the site and time course of RNA synthesis in relation to the total cell cycle, DNA synthesis, and mitosis. Data highly indicative of a precursor role of nuclear RNA for cytoplasmic RNA was obtained. Once synthesized, RNA was found to be preserved, while nucleotides of small molecular weight turned over rapidly with release from the cell of breakdown products and nucleosides which were reincorporated into RNA and DNA. The existence of intermediate precursors for "delayed incorporation" into DNA was established. Chemical analysis of the distribu-

tion of H^3 between the pyrimidine bases in RNA, DNA, and precursor pool indicated and confirmed that most of the RNA remained stable and that cytoplasmic RNA was derived from nuclear RNA. The previously observed "delayed incorporation" of precursor (cytosine and thymine) into DNA was found to be related to a cytosine derivative, and the conversion of this compound from riboside form to deoxyriboside form could be partially inhibited. This suggests a function of an RNA intermediate in supplying precursors for DNA. Low concentration RNase inhibited partially the incorporation of H^3 -thymidine and H^3 -deoxyuridine into DNA, but not of H^3 -cytidine or H^3 -uridine into RNA and DNA. Autoradiography and special histochemical treatment of human cells in culture allowed the demonstration of chromosomes containing newly synthesized RNA, a further support for DNA-associated RNA synthesis in the intact cell. Indications of transfer of labeled RNA between homologous mammalian cell lines in culture have been observed in preliminary experiments.

DEOXYRIBONUCLEIC ACID METABOLISM

In previous years I^{131} -labeled iododeoxyuridine (IDU) has been shown, like thymidine, to be a specific label for DNA. It is utilized less efficiently than thymidine, being largely degraded into iodouracil and then to iodide. Once these catabolic products have been excreted, the radioactivity retained by the animal reflects the total DNA metabolism in the tissues. Subsequent loss of activity measures the amount of cell death, since DNA does not turn over in the living cell.

In continuation of these studies, synthesis of IDU by direct iodination of deoxyuridine has been shown to proceed via the addition of hypoiodous acid, with the formation of a stable intermediate which is converted by acid to IDU. Studies in mice of the simultaneous incorporation of two tracers (I^{131} DU and H^3 -thymidine) into DNA have shown a much greater sensitivity of I^{131} DU to physiological variations, including thymidine load. This finding may explain the recently reported sensitivity to radiation and cortisone in the uptake of IDU.

Cell death following radiation or following the injection of nitrogen mustard has been measured quantitatively *in vivo* by the use of two isotopes of iodine, the second to permit a correction for the iodide pool. Cell death from radiation begins with-

in an hour after exposure and is most pronounced among cells labeled shortly before irradiation.

Attempts to improve the utilization of IDU for studies in tumor metabolism have shown that the fluorine analogue FDU markedly potentiates the utilization of IDU. This effect is observed when as little as 10^{-10} moles of FDU are injected into a mouse and can reasonably be ascribed to a depression of the thymidine pool by stoichiometric combination with the enzyme synthesizing thymidine. Localized potentiation of IDU incorporation has been achieved by clamping the portal vein for 10 min upon injection of IDU. The resulting decreased catabolism of IDU by the liver increased the interval during which the compound was available for labeling the bone marrow and other peripheral tissues.

THE LETHAL EXPRESSION OF H^3 -THYMIDINE IN CLONAL POPULATIONS OF MAMMALIAN CELLS IN CULTURE

In the course of establishing a concentration of H^3 -thymidine that would prove adequate as a tracer for DNA synthesis and would not kill cells or in any way interfere with normal metabolism, it was noted that a small fraction of the cell population consistently survived and produced normal colonies of HeLa S3 cells after exposure to high concentrations of H^3 -thymidine (0.1 to 1 μ C/ml, specific activity 1.9 C/mM). Under the experimental conditions a 100% lethal effect could not be demonstrated, and it was possible to show that the persistent fraction were not progeny of radiation-resistant cells.

Further investigation revealed that the persistent fraction represented cells in the population that had a generative cycle considerably longer than 22 hr and did not enter DNA synthesis during the 24-hr growth period in the radioactive medium. The probability of colony development was greatly influenced by the size of the clone population at the time of the addition of the radioactive medium. Autoradiograms of HeLa S3 cells clearly showed that in young cultures having only 2 to 4 cells per clone the percentage of clones in which *all* cells were labeled after a 24-hr growth period in medium with 0.1 μ C/ml H^3 -thymidine was much greater than in older cultures having an average of ≈ 40 cells per clone. In the latter group $\approx 64\%$ of the original colonies exposed to H^3 -thymidine

gave rise to normal clones; this was in excellent agreement with the distribution of unlabeled cells. When the incubation time in the radioactive medium was extended to 48 hr, autoradiograms showed that all cells within the clones, regardless of the numbers in the clone population, had incorporated H^3 -thymidine and that all cells were equally susceptible to the lethal action of the radioisotope.

ORIGIN, RATE OF DIVISION, GROWTH, AND FATE OF BODY CELLS

These investigations are concerned with studying the origin, proliferation, and fate of cells under normal conditions and under abnormal conditions such as hemorrhage, hypertransfusion, exposure to radiation, and disease. The chief technique used involves tritiated nucleic acid precursors such as thymidine or cytidine as cell markers and high-resolution autoradiography with stripping or dipping film. The importance of knowledge of the origin and potentialities of cells is obvious. Characterization of the time parameters of cell proliferation in the normal steady state and in abnormal non-steady state conditions will serve to establish the groundwork for determining the mechanism(s) of regulation in the normal steady state and the basic defect in blood dyscrasias. Knowledge of the pathogenesis should make possible the development of a rational basis for therapy for the malignant and nonmalignant states.

The presence in normal peripheral blood and thoracic duct lymph of cells capable of synthesizing DNA, and thus presumably of proliferation, has been demonstrated. Blood and thoracic-duct lymphocytic cells from donor rats given tritiated thymidine were easily demonstrable in the bone marrow, spleen, and lymph nodes of normal and irradiated parabiotic or intact recipients, and some evidence of "cell transformation" was found; however, it could not be established that these cells could become erythrocytic or myelocytic precursors. The level of toxicity of tritiated thymidine for lymphocytes, bone marrow cells, and the very radiosensitive spermatocytes of the mouse was found to be well above the dose usually employed in tracer studies.

Labeled bone marrow cells were shown to be capable of differentiating into fibroblasts in artificially induced inflamed and wound areas in the

Figure 6. Autoradiograms showing concentration of I^{125} -labeled ribonuclease in the kidney. Upper left, cross section of kidney (unstained), $10\times$. Upper right, section of cortex (unstained), $50\times$. Bottom, high power view of glomerulus and surrounding tubules (stained), $500\times$. Note individual silver grains lying densely over tubular cells.

rat. The proliferative capacity of normal rat bone marrow cells and of marrow cells at various times after exposure to radiation was determined. A significant effect of radiation on the process of DNA synthesis in the individual cell was shown to exist. Detailed studies with dogs indicated that the rapid response in terms of reticulocyte and red cell output from the bone marrow after bleeding is not accomplished by a change in generation time or of DNA synthesis time of nucleated precursors.

Methods of evaluating in detail the time parameters of erythrocytic precursors have been worked out for the dog. Generation and maturation times are ≈ 10 hr, and the nuclei of erythrocytic precursors are extruded ≈ 10 hr after the final division. Initial studies in human beings have indicated somewhat longer times for comparable erythroblast parameters. The upper limit for blood transit time of human granulocytes was shown to be ≈ 22 hr. Lymphocyte turnover time was not

established but was shown to be of the order of weeks to months. Conceptual schemes for interpretation of tritiated thymidine autoradiographic data have been worked out. Acute and chronic leukemias were shown to proliferate relatively slowly as compared to normal steady state cell production rates. Multiple myeloma cells were shown to have a generation time of ≈ 100 hr, and it was shown that a marked death function of the abnormal cells must be present in order for myeloma patients to live as long as they do. Patterns of appearance of mature peripheral granulocytes in polycythemia, multiple myeloma, and leukemias are essentially unchanged from the normal steady state. Studies on proliferation in different types of leukemia have been carried out during and after therapy.

PLASMA PROTEIN METABOLISM

In previous years proteins have been labeled *in vitro*, usually with I^{131} , and the rates of their disappearance from the circulation and/or the rate of excretion of the catabolic product (iodide) have been determined. While catabolism of the larger proteins such as serum albumin appears to occur in the reticuloendothelial system, there has been a suggestion of partial catabolism in the kidney tubules as a result of glomerular filtration and tubular absorption. This effect has now been shown more clearly by the use of the small protein ribonuclease, with a molecular weight of only 14,000. More than half of an injected dose of iodine-labeled ribonuclease appeared in the kidneys of the mouse or rat within 5 min after intravenous injection, and autoradiography using I^{125} has clearly demonstrated that the radioactivity is concentrated in the cells bordering the lumen of the tubule (see Figure 6).

Radioiodine that rapidly leaves the kidney as inorganic iodide has been demonstrated *in vitro* by incubating slices of kidneys taken from rats shortly after injection of labeled ribonuclease. Catabolism in the kidney is depressed *in vivo* by mercuric chloride.

Ribonuclease labeled with Cr^{51} is rapidly concentrated in the kidneys, but this label differs from I^{131} in that it remains fixed in the kidney for some days. This cellular retention of chromium resembles the retention in red cells labeled with chromium and suggests that chromium carried into a

cell may remain there longer than iodine after the carrier molecule has been catabolized. This may thus provide a general approach to the elucidation of sites of protein catabolism.

STUDIES ON COLLAGEN

Collagen, which constitutes $\approx 1/3$ of the total proteins of the animal body, is the structural protein that holds the body together. It contains two unique amino acids, hydroxyproline and hydroxylysine. The latter has been a continuing subject of investigation in this Laboratory, particularly in regard to its additional reactive hydroxyl group. The hydroxylysine has been shown to originate entirely from lysine; during synthesis of collagen in the animal body part of the lysine involved has been shown to be changed to hydroxylysine. The mechanism by which the lysine is hydroxylated is being studied. In the formed collagen part of the $-\text{CH}(\text{OH}) \cdot \text{CH}_2(\text{NH}_2)$ groups of the hydroxylysine do not react with periodate, a behavior suggesting that such groups may be involved in covalent cross-linkages in the collagen molecule. The existence and nature of such linkages are also being investigated.

During the past year it has definitely been established that the hydroxylation cannot occur after collagen has been laid down in its permanent insoluble form, since the ratio of the radioactivities in lysine and hydroxylysine in rat skin collagen that had been labeled by injecting C^{14} -lysine remained constant throughout this period. In the course of these studies it has also been observed that collagen is metabolically inert, showing no evidence of replacement in rat skin during 15 weeks of study.

A similar experimental approach has revealed that this metabolic inertness also extends to the reactivity of the hydroxylysyl residue in collagen towards periodate. The same fraction reacts in both young and old collagen, which indicates that no additional cross-links involving this grouping are formed in the permanent structure.

TRYPTOPHAN METABOLISM IN DISEASE STATES

An area of tracer study that has received attention during the past year involves the metabolic rate of the amino acid tryptophan in neoplastic

disease. Clinical studies are in progress utilizing tryptophan-2- C^{12} , tryptophan-7a- C^{14} , 3-hydroxyanthranilic acid-7- C^{14} , and anthranilic acid-7- C^{14} . The apparatus used in these clinical studies is shown in Figures 7a and 7b. Although insufficient numbers of patients have been studied to permit comments concerning the influence of various diseases on the metabolic rate of the labeled tryptophan, other significant data on human beings have been obtained with the $C^{14}O_2$ apparatus mentioned above.

As shown in Figure 7c, with a constant level of C^{14} activity in an oral dose of tryptophan-2- C^{14} , the quantity of $C^{14}O_2$ expired by a patient may be elevated by increasing the dosage of unlabeled tryptophan administered. These data suggest that humans convert given quantities of tryptophan into protein, serotonin, niacin, coenzymes, and CO_2 , but, when the amount of tryptophan in the body spaces reaches a critical level, the excess is metabolized into $C^{14}O_2$ via the kynurenine-3-hydroxyanthranilic acid-acetate pathway or some other pathway to be elucidated.

In these studies proof has been obtained of the existence of a new metabolic product of tryptophan in the urine of a multiple myeloma patient. The aminophenols 3-hydroxyanthranilic acid and 3-hydroxykynurenine appear in elevated quantities in the urine of patients with bladder tumors, anemias, and leukemias. Another aminophenol, *o*-aminophenol, has been found in the urine of human beings, and it occurs in elevated quantities in the urine of bladder tumor patients. In studies of DL-tryptophan-7a- C^{14} metabolism, C^{14} -labeled *o*-aminophenol was successfully isolated from the urine of a multiple myeloma patient. The mechanism of formation of *o*-aminophenol appears to be the direct decarboxylation of 3-hydroxyanthranilic acid formed in the metabolism of tryptophan. Additional evidence for this pathway has been obtained in studies of the metabolism of ring-labeled 3-hydroxyanthranilic acid in mammary tumor mice. C^{14} -labeled *o*-aminophenol was isolated from the urine of these mice.

STUDIES OF CARBOHYDRATE AND FAT METABOLISM WITH CARBON AND HYDROGEN RADIOISOTOPES

The maxim that the body derives energy by conversion of sugar to carbon dioxide and water

has been scrutinized by means of a new approach in this Laboratory in the past year. Much is known of the various chemical steps in this over-all process. However, by the combined use of C^{14} as a carbon tracer and tritium (H^3) as a hydrogen label, more is being learned about the mechanism by which the energy inherent in carbon-carbon and carbon-hydrogen bonds is parceled, stored, and eventually converted to useful work by enzymes and coenzymes, and the efficiency of the process. Studies in this area of intermediary metabolism may yield information on the sites of action of some hormones and also on primary biochemical pathology in diabetes, obesity, hyperthyroidism, and similar abnormal metabolic conditions.

Hereditarily obese mice have been found to convert 10 times as much of the labeled hydrogen on the first carbon atom of glucose to liver fatty acids and 4 times as much to carcass fatty acids as do their lean littermate controls, while at the same time liberating equal amounts of C^{14} from the first carbon atom of glucose to carbon dioxide *in vivo*. This suggests superactive chemical reduction of fatty acid intermediates and excessively efficient use of glucose energy for fat synthesis in the obese animals. However, the amount of tritium found in total fatty acids is a very small percentage and the amount in body water a large percentage of the administered glucose tritium, even in the obese, which indicates other main sources of metabolic hydrogen for fatty acid synthesis. Increase in the ratio of C^{14} in carbon dioxide to tritium in body water further indicates more extensive synthetic metabolic activities in the obese compared to the lean mice. Preliminary study with lactic acid-2- C^{14} - H^3 suggests a similar difference in the mice with this substrate, which passes on its labeled hydrogen via a different coenzyme (diphosphopyridine nucleotide) than does glucose-1- H^3 (Figure 7d).

Comparable studies with patients (see Figures 7a and 7b) have so far included obese nondiabetic subjects and lean diabetic subjects. The examination of $C^{14}O_2$ in breath and H^3_2O in blood and urine further suggests differences in synthetic metabolic activity between these two groups and also an enhancing effect of insulin in the diabetic patient. For instance, insulin increases the amount of C^{14} from lactate in carbon dioxide but decreases the amount of H^3 from lactate in body water of the diabetic patient. The turnover of tritium in the

Figure 7. Studies of intermediary metabolism of carbohydrates and amino acids using radioactive carbon and tritium. (a) Collection of C¹⁴O₂ from patient. (b) Measurement of C¹⁴O₂ from patient. (c) Influence of tryptophan dose level on percent of C¹⁴O₂ expired. (d) Possible intermediate steps (in brief) for C¹⁴ and H³ transfer from glucose and lactate to fatty acids, carbon dioxide, and water. TPN and DPN = tri- and diphosphopyridine nucleotide coenzymes, respectively. CoA = coenzyme A.

body water over a period of several days is another aspect being studied in patients. Turnover time after administration of labeled glucose, but not of labeled lactate, is substantially longer than after administration of labeled water.

A brief description of the principal intermediate metabolic steps involved in these studies is given in Figure 7*d*. Figures 7*a* and 7*b* show the process and apparatus for continuous collection and analysis of $C^{14}O_2$ (and total carbon dioxide) in the breath of patients.

STUDIES ON METABOLISM OF LABELED VITAMIN B_{12} , RADIOIRON, AND RADIOPOTASSIUM IN HEALTHY HUMAN BEINGS AND IN PATIENTS

Blood cell production is dependent upon absorption and normal transport of vitamin B_{12} and iron to blood-forming tissues. The objective of this study is the measurement of absorption, excretion, transport, and turnover rates of these substances in healthy individuals and in patients with blood dyscrasias due to an aberration of metabolism of one or the other. A method for the detection of intrinsic factor defect (pernicious anemia) and malabsorption of vitamin B_{12} utilizing the whole-body counter has been developed. The biological half-time of Co^{60} -labeled B_{12} was shown to be ≈ 360 days. Plasma clearance of B_{12} was shown to be a multicomponent exponential with great prolongation in chronic granulocytic leukemia and polycythemia. The prolongation was correlated with an increased binding capacity of the plasma for vitamin B_{12} . Co^{60} -labeled B_{12} was shown to persist and represent unchanged B_{12} for over 90 days after injection or ingestion; hence Co^{60} activity for at least this time interval represents the behavior of unlabeled vitamin. Turnover rates of vitamin B_{12} in chronic granulocytic leukemia were found to be significantly slower than in normal individuals. Tissue distribution of labeled vitamin B_{12} has been determined in 12 patients following autopsy. The dose calculations show the infinity doses in organs of deposition to be less than "permissible" dosages.

Studies of Fe^{59} absorption and loss – the latter both by physiologic routes and by hemorrhagic loss of tagged red cells – have been under way for almost two years with the whole-body counter. The method results in a normal radioiron absorp-

tion of 5 to 25% and rate of loss of 0.0110 to 0.182% of the tracer per day. Malabsorption syndromes depress absorption well below the normal range, whereas iron deficiency results in increased absorptions as high as 97%. Quantitation of abnormal blood loss, notably that of menorrhagia, has also been possible with this technique.

In the course of therapy of many blood dyscrasias, corticosteroids are used in pharmacologic doses that may seriously deplete body potassium stores and endanger the life of the patient. Determination of total body potassium is only practical by the whole-body determination of K^{40} in a whole-body counter. Serial studies of K^{40} content are made on all individuals in whom steroid therapy is commenced. Computer analysis with both digital and analogue types is often required.

KINETICS OF SODIUM IN HUMAN AND EXPERIMENTAL HYPERTENSION

The operational hypothesis for this work is that a derangement in sodium metabolism plays an important role in the hypertensive process in man. Furthermore, excess dietary consumption of sodium (as ordinary table salt) is believed to be the most common manner in which this derangement is brought about. Evaluation of the hypothesis has involved the following:

1. The average salt intake of humans has been correlated with prevalence of hypertension in the respective communities of origin.
2. The long-term effects of chronic excess salt intake on several animal species are under continuous study.
3. With the aid of special whole-body counters for both man and small animals, abnormalities in handling Na^{22} have been recognized in hypertensive humans and animals.
4. The genetic factors known to be operating in human hypertension are being approached in animal studies by the development of two strains of rats. One strain does not develop hypertension, while the other does, although both have the same dietary salt intake.

The most important accomplishments were concerned with the demonstration that sodium is of fundamental importance in the hypertensive process. It was demonstrated that dietary treatment of hypertension depends primarily upon sodium restriction for its effectiveness. Furthermore, a paral-

lel increase in average salt intake and prevalence of hypertension was found to occur in the members of five communities studied throughout the world, which suggests a causal relationship between salt intake and hypertension. Finally, it was observed that long-term low salt diets are not harmful, which indicates that the high salt intakes common in western societies are not only probably injurious but also unnecessary. In 15 human subjects studied with the whole-body counter the rate of loss of Na^{22} from the body was found to be significantly slower in those with hypertension than in those without, because hypertensives have higher tissue sodium concentrations than normotensives. Chronic excess salt consumption by ≈ 200 rats and 17 dogs was observed to be associated with significant increases in plasma cholesterol concentrations in half of the animals. The importance attached to elevation of cholesterol in human atherosclerosis suggests that excess salt intake may play an accessory role in this disease also. Sea salt produced more hypertension in rats than did sodium chloride, which suggests that trace elements may aid in producing hypertension.

Explanation of the important observation of a decreased rate of Na^{22} loss in hypertensives was found. In 35 rats with hypertension induced only by salt-feeding, $\approx 2/3$ did not respond to subsequent salt withdrawal. This indicates that excess salt ingestion can cause permanent hypertension, at least experimentally. The development of two strains of rats, one that develops hypertension and one that does not, although both strains have identical salt intakes, suggests a strong genetic factor in the susceptibility to hypertension that results from salt-feeding.

STUDIES ON TRACE METAL METABOLISM

Investigation of manganese metabolism in some disease states has been continued using whole-body and localized external radioisotope counting techniques in association with neutron activation analysis. Incomplete studies suggest some possibility of altered radiomanganese kinetics in a subgroup of patients with Parkinsonism. A pilot investigation of hyperthyroidism revealed the perturbation of normal kinetics predicted by studies published previously. A different abnormal pattern, consistent with earlier findings in a case of reversible hydralazine-induced "collagen disease,"

has been observed in a few patients with collagen diseases. Concurrent animal experiments have confirmed an effect of hydralazine upon the organ distribution of injected Mn^{54} .

Additional aspects of normal manganese physiology also have been studied. The concentration of manganese in human blood, plasma, and spinal fluid under varying conditions was measured by activation analysis. Because of the hepatic regulation of body manganese, reported earlier, an extensive study of manganese (and other metals) in rat bile was undertaken. Preliminary results reveal marked but different effects of hemorrhagic shock and of anoxia. The incorporation of manganese into a metabolically stable erythrocyte porphyrin was further documented in human beings, and more complete chemical characterization of this compound is planned. The β_1 globulin that binds the labile manganese of human plasma was further defined by immunoelectrophoresis, and preparative procedures were begun using column chromatography and continuous electrophoresis. Finally, investigations in mice of the effects of acute whole-body gamma irradiation showed significant alterations in excretion and organ distribution of normal radiomanganese.

Animal experiments with radioisotopes have shown zinc metabolism to be under homeostatic control, but the zinc pathway through the body is less specific than that shown previously for manganese. Thus cadmium can affect the metabolism of zinc and can substitute for it incompletely, although cadmium itself is not homeostatically regulated. In the liver zinc, in further contrast to cadmium, becomes sequestered within normal macromolecular constituents.

The role of transition-group trace metals in the generation and reaction of organic free radicals was emphasized. Spectrophotometric, potentiometric, and electron paramagnetic resonance (EPR) methods were utilized to determine in detail the univalent redox reactions of metal ions with phenothiazine drugs. These results are providing a model for current EPR investigations of metal interactions with free radicals in biochemical systems.

The procedures developed previously for manganese determination by neutron activation analysis are being extended and generalized to provide feasible multielement analyses on single biological samples.

METABOLIC STUDIES IN NEOPLASTIC DISEASES UTILIZING RADIOACTIVELY LABELED PROTEINS

The present studies are a continuation of clinical investigations concerned with the physico-chemical properties and functions of the serum proteins in neoplastic diseases. In the patient with a known type of cancer the turnover rates of normal globulin, aberrant globulin, and other serum proteins labeled with I^{131} are determined. The turnover of I^{131} -labeled proteins, prepared by chemical fractionation, is determined not only by the conventional method of blood and urine sampling but also by the simpler approach of utilizing the whole-body gamma spectrometer. This device permits *in vivo* measurement of very low levels of an internally deposited gamma emitter. Should a differential metabolic recognition exist between the normal and aberrant proteins in a patient, it might serve as a basis for a diagnostic test. Serial turnovers carried out on a group of 50 women with cancer of the breast indicate a slower rate of catabolism of the normal gamma globulin as compared to the aberrant globulin (each patient received both types).

Multiple myeloma is a malignant neoplastic disease in which the globulins produced by the tumor plasma cells of the bone marrow rise to excessively high levels in the blood. This hyperproteinuria constitutes a fundamental protein metabolic disturbance which has been investigated by determining the biological half-life of (a) labeled normal gamma globulin, (b) autologous and homologous labeled gamma globulin in multiple myeloma, and (c) labeled beta globulin. The rate of catabolism of the gamma globulin is accelerated in patients with hypergammaglobulinemia, which indicates a very high rate of synthesis of the gamma globulin in these subjects. The studies now being completed with labeled beta globulin indicate that it is degraded faster than gamma globulin in patients with hyperbetaglobulinemia. The fundamental metabolic defect in these hyperproteinemic patients with multiple myeloma is therefore due to an accelerated synthesis of the abnormal proteins, which more than compensates for the accelerated rate of catabolism and results in excessively large amounts of these proteins in the blood.

ANALOGUE COMPUTER ANALYSIS OF TRACER KINETICS

The objectives of this project are the development of an analogue computer system and its application to the analysis of radioactive tracer data obtained in the Medical Department. The existing instrument uses a combination of direct and indirect analogue features giving maximum simplicity of operation without undue limitations on the scope of the problems that can be handled. The data display system used, unique in this kind of instrument, employs a 120-position mercury jet switch to generate points. The points may be moved to any position in the display to simulate data points, which are displayed on an oscilloscope simultaneously with the analogue curves.

During the year the computer was enlarged to provide increased capacity for handling problems involving unequal flow rates between compartments and for combining compartments. The chief applications involved analyses of data on creatinine metabolism, the production of CO_2 from labeled precursors, strontium and calcium metabolism, and the degradation of labeled proteins.

MEDICAL DEPARTMENT CONCLAVES

As part of its general educational endeavor the Department has sponsored jointly with the Division of Biology and Medicine of the US AEC a series of annual conclaves on nuclear science in medicine. Participation is by invitation only, and the objective is to provide two full days of discussion and demonstration dealing with a special field of medicine in which the visitors are qualified. Members of the Brookhaven Medical Department Staff give talks as a basis for discussion as well as demonstrations of various devices useful in atomic energy programs. The sixth such conclave, held October 23 and 24, 1961, was entitled "The Hazards of Uses of Atomic Radiation in Medical Practice, Teaching, and Research in Internal Medicine," and was attended by 46 chairmen of departments of internal medicine from 42 medical schools in the United States (including one from Puerto Rico) and from 4 in Canada.

SUPPORTING ACTIVITIES

Technical Operations and Services

The technical services and facilities essential to the Laboratory's research programs are provided by a number of organizational units whose operations are described below under appropriate headings.

REACTOR OPERATIONS

The Brookhaven Graphite Research Reactor (BGRR) operated throughout the year without a major delay or operating incident. Operation was at full power (20 Mw) during 84% of the year. In addition to 19 scheduled shutdowns for service work and changing experimental equipment, there were 9 unscheduled shutdowns of the reactor, of which 6 were caused by human error and 2 by failure of the Long Island Lighting Company's power supply. The remaining shutdown was occasioned by the need to inspect a fuel channel because of an abnormally high temperature indication. The unusually large number of human errors may be attributed primarily to the 5-year overhaul of the control rod drive units during the year and to the "fail safe" nature of many of the associated circuits, which led to shutdown of the reactor following minor errors.

To meet normal reactivity requirements it was necessary to increase the reactor loading from 575 channels, the design number, to 607 channels.

The fuel removed from the reactor averaged 41.7% burn-up of the original U^{235} . No serious difficulties were encountered in the behavior of the fuel or in its handling.

The graphite moderator structure of the BGRR was annealed for the 17th time in September 1961. The 4-month annealing cycle was then suspended pending development of a new annealing procedure. This procedure, which was approved by the Reactor and Critical Experiment Safety Committee on May 21, 1962, calls for more frequent annealings at lower temperatures (annealing interval, 3 weeks; maximum annealing temperature, 250°C; minimum temperature to achieve desired result, 150°C). The new procedure is expected to result in considerable recovery of the past growth of the structure.

Training activities were conducted which resulted in the qualification of all reactor supervisors and operators on a formal basis. This program involved classroom lectures in the subject areas of health physics, reactor instrumentation, and reactor technology followed by written examinations.

Difficulties experienced with the Fairbanks Morse electric motor drives for the three new primary cooling fans were corrected early in fiscal 1962, and the economies in cooling costs expected from operation of these fans were realized. Since the electrical costs associated with cooling the reactor are a major factor in the cost of its operation, these figures are plotted for several years in Figure 1.

Inspections by both the Laboratory and the AEC consumed a considerable amount of the time of the principal staff members. The reactor was inspected in great detail by an *ad hoc* committee appointed by the Laboratory's Director. It was inspected in somewhat less detail by two AEC committees, one from the Division of Operational Safety and the other from the Health and Safety Laboratory of the New York Operations Office, acting for the Brookhaven Office. In addition, the Brookhaven Office made weekly reactor inspections. The resulting reports were acted upon by internal committees, and minor changes in equipment and procedures have been made.

Utilization of the BGRR

Approximately 27 of the reactor's experimental holes are assigned to the Laboratory's research de-

Figure 1. Electrical costs associated with cooling the BGRR.

Table 1
Summary of Reactor Use Charges

	Use charges (in thousands of dollars)							Percent of total usage	Dollar amounts
	0	100	200	300	400	500	600	700	
<u>Fiscal 1962</u>									
Outside Organizations								5.7	\$ 65,156
Irradiation Service Group								4.6	51,739
BNL Research Departments									
Physics								63.2	724,096
Chemistry								9.7	111,870
Nuclear Engineering								13.0	149,008
Biology								3.8	43,500
								100.0	\$1,145,369
<u>Fiscal 1961</u>									
Outside Organizations								10.5	\$121,621
Irradiation Service Group								7.4	84,720
BNL Research Departments									
Physics								53.4	615,933
Chemistry								8.7	99,900
Nuclear Engineering								14.2	165,193
Medical								2.0	23,896
Biology								3.8	43,410
								100.0	\$1,154,673
<u>Fiscal 1960</u>									
Outside Organizations								11.4	\$149,119
Irradiation Service Group								7.5	97,920
BNL Research Departments									
Physics								47.4	621,827
Chemistry								6.9	90,480
Nuclear Engineering								16.1	211,960
Medical								6.0	79,200
Biology								4.7	62,240
								100.0	\$1,312,746

Table 2
Reactor Irradiation Service Charges
(Excluding Handling Charges)

	Fiscal 1962		Fiscal 1961		Fiscal 1960	
Outside Organizations	\$ 56,905		\$ 56,670*		\$ 88,971	
BNL Research Departments						
Physics	\$19,527		\$23,837		\$17,730	
Chemistry	10,077		7,715		8,534	
Nuclear Engineering	36,269		27,428		11,234	
Medical	8,643		6,170		3,934	
Biology	1,538		1,200		218	
Instrumentation and Health Physics	1,711	77,765	550	66,900	2,380	44,030
Total	\$134,670		\$123,570		\$133,001	

*The decrease in charges from 1960 to 1961 reflects the establishment of separate accounts for service irradiations, processed radioisotopes, and handling.

partments for projects that are described in the pertinent sections of this report. A number of reactor experimental facilities have been assigned to outside research organizations for use under various arrangements. Tables 1 and 2 indicate the relative use of the reactor by the research departments and outside organizations, as well as the charges applied to these programs to recover the operating expenses of the reactor. The outside organizations making significant use of the reactor, together with brief descriptions of their activities, are listed below.

Bell Telephone Laboratories. Studies of magnetic and crystal structure were carried out by using a neutron spectrometer to obtain neutron diffraction patterns from crystals. Measurements were made at room temperature and at 4.2° K. (Work done under a utilization agreement.)

Rensselaer Polytechnic Institute. Energy deposition in gaseous chemical systems has been studied. A loop experiment has been carried out to determine the effect of fission fragments on oxygen-nitrogen mixtures circulating through a stainless steel pipe containing quartz fibers with 15 wt % U_3O_8 (90% enriched in U^{235}). Sealed quartz capsules that contain various gaseous mixtures, with and without uranium, were also irradiated at different temperatures and pressures. (Work performed under Government Contract No. AT-30-4-321.)

General Electric Company. After several months of inactivity, work was resumed using a neutron

spectrometer. Studies of magnetic and crystal structure were carried out from neutron diffraction patterns obtained from various crystals and powders at different temperatures. (Work done under a utilization agreement.)

Sperry Gyroscope Company. Studies of radiation damage to electronic components and semiconductor devices were made during one reactor operating cycle. Electrical properties were measured while the irradiation was in progress to detect any changes due to the radiation field.

Naval Research Laboratory. Short-term irradiation of magnetic core materials was carried out to study changes in magnetic properties caused by reactor radiation damage. Long-term irradiations of steel specimens were carried out at reactor ambient temperatures to study the effects of radiation damage caused by the reactor radiation environment. (Work done under Government Contract No. NRL-AEC-11-61.)

Columbia University. A single-crystal neutron spectrometer effective in the neutron energy range from 0.0006 to 10 ev was used to measure total neutron cross sections of interest in the study of atomic motion in solids and liquids. The properties of solid-state detectors were studied by placing the devices in a collimated neutron beam. The detectors were also used to study alpha-particle distributions in ternary fission.

Wright Air Development Center. A three-NaI(Tl)-crystal pair spectrometer in coincidence with a single NaI(Tl) crystal was used to study

gamma-gamma coincidence following neutron capture.

Naval Ordnance Laboratory. Diffraction studies made jointly with the BNL Physics Department (Solid State) were carried out by using single crystals and a polarized beam of neutrons. In addition, powdered magnetic materials were studied with use of a powder diffractometer.

Procurement of Special Materials

The procurement for the scientific departments of all radioactive and stable isotopes as well as special materials controlled by the AEC is a responsibility of the Isotopes and Special Materials Group. In this connection, 282 purchase orders were placed for radioisotopes, 71 for stable isotopes, and 44 for special materials. Against these orders, ≈ 537 shipments were received and processed. These figures represent an increase over last fiscal year of 20% in orders placed and 30% in receipts. Of the orders placed, $\approx 26\%$ were for the Medical Department, 27% for Biology, 16% for Physics, 14% for Nuclear Engineering, 14% for Chemistry, and 3% for Instrumentation and Health Physics.

The Isotopes and Special Materials Group is also responsible for annual inventories of radium sources and purchased stable isotopes and for negotiations for the loan of valuable isotopes. The number of radium sources at Brookhaven remained unchanged at 39 with a total activity of 5.86 curies and a value of \$38,883. The inventory of purchased stable isotopes (\$34,494) was about the same as last year's, since additional purchases totaling \$15,380 were offset by consumption valued at \$14,962. During the year the number of loan agreements for stable isotopes increased from 35 to 43 with a value of \$140,000; in addition, two special samples had an estimated value of \$950,000.

The revised AEC policy for obtaining irradiation services in the Materials Testing Reactor (MTR) and the Engineering Test Reactor (ETR) operated by Phillips Petroleum Company at the National Reactor Testing Station, Idaho Falls, Idaho, requires semiannual forecasts of anticipated usage and permits direct arrangements with the Idaho Operations Office of the AEC. The Brookhaven irradiation programs in these reactors during fiscal 1962 are summarized below.

BNL-15: Production of 500,000 Curies of Co^{60} . The irradiation of cobalt in the ETR was com-

Table 3

Source and Special Nuclear Materials on Hand
at End of Year (in kilograms)

	Fiscal year		
	1962	1961	1960
<u>Station BRL</u>			
Natural U	2,043	2,688	2,895
Depleted U	2,968	2,950	14,538
$\text{U}^{235} > 75\%$	142	69	163
$\text{U}^{235} < 75\%$	6,410	7,815	11,850
U^{233}	0.047	0.051	0.051
Pu^{239}	3	2	7
Th	216	252	254
Heavy water	15,030	13,179	13,321
<u>Station BRG</u>			
$\text{U}^{235} > 75\%$	108	84	—
<u>Station BRM</u>			
$\text{U}^{235} > 75\%$	3	3	—

pleted in fiscal 1961, and delivery of the last 30,000 curies of Co^{60} was made this year.

BNL-18: Testing of Prototype Fuel Elements for the HFBR. This program was initiated in the ETR in May 1960 for the testing of one element. In fiscal 1961 the program was revised to include testing of three more elements, and in fiscal 1962 five additional elements, some to be irradiated in the MTR, were added to the program. Testing was completed on three of the nine fuel elements during this fiscal year and is in progress on four more. The expiration date of this program is expected to be March 31, 1963.

BNL-19: Radiation Damage in Carbon. This program involved short irradiations of five capsules of diamond dust in the MTR. These irradiations, a continuation of a series performed under programs BNL-3 and BNL-11, were completed early this year.

BNL-21: Radiation Damage in Pure Iron. Four capsules containing iron tensile test specimens are being irradiated in the MTR. Because of capsule fabrication problems, start of the irradiation was delayed until late this fiscal year. Completion is expected by September 1962.

BNL-22: Radiation Damage in Graphite. This program involves irradiation in the ETR of three instrumented capsules containing special graphite

samples. The irradiation was begun late this fiscal year and is expected to be completed by June 30, 1963.

BNL-23: Radiation Damage in Carbon. This program, a continuation of the BNL-19 series, calls for irradiation in the MTR of three capsules of diamond dust. Two capsules have been irradiated and returned, and the program is expected to be completed in October 1962.

BNL-24: Radiation Damage in Carbon. This program, which involves irradiation in the MTR of three capsules of diamond dust in continuation of the BNL-23 series, should be completed by December 1962.

BNL-25: Production of Special Research Isotopes. Under this program irradiation in the MTR of one capsule containing twelve samples of enriched stable isotopes was started late this fiscal year; completion is expected by September 1962.

Source and Special Nuclear Material Accountability

Table 3 presents a 3-year summary of the amounts of source and special (SS) nuclear materials on hand at Brookhaven at the end of each fiscal year. Based on dollar values published by the AEC, the current inventory of SS materials represents a value of $\approx \$5,600,000$. The net increase from last year's figure of $\approx \$4,000,000$ was due primarily to the additional $U^{235} > 75\%$ and heavy water received during the year.

The individual decreases in the amounts listed are the result of shipments of surplus items and irradiated natural uranium fuel containing plutonium. The increases are due to receipt of additional material. In the case of Station BRG, however, the increase in $U^{235} > 75\%$ is the net difference between the receipt of new fuel elements and the shipment of spent fuel elements.

A purchase order for 3000 new fuel elements for the BGRR was initiated late last year with Metals and Controls, Inc. Actual production began with the start of fiscal 1962, and much attention was given to checking the initial production efforts and eliminating various difficulties. When the first delivery was received from the fabricator, only about a 2-month supply of fuel elements for the reactor was on hand. Difficulties arising about halfway through the production program required BNL to test ≈ 900 delivered fuel elements. These tests resulted in the rejection of some 300 elements and

their return to the fabricator. Delivery of the 3000 fuel elements was completed in June 1962. It is anticipated that by April 1963 the inventory of fuel elements will be reduced to a 1-year supply and that a new order will be initiated at that time.

On June 1, 1962, the BGRR and the Medical Research Reactor (MRR) facilities became subject to inspection by the International Atomic Energy Agency (IAEA) for a period of two years under the Four Reactors Agreement with the U.S. Government. In preparation for this inspection, the accountability for SS nuclear materials at Brookhaven had been separated in March 1961 into three accounting stations, MGR for the graphite reactor, MMR for the medical reactor, and BNL for the remainder of the Laboratory. In October 1961, with a change in responsibility from the New York Area Office to the local Brookhaven Office of the AEC, these station letters were altered to the present BRG, BRM, and BRL.

For purposes of IAEA inspection, the accounting system was revised with the assistance of representatives of the AEC Division of Nuclear Materials Management. The SS accountability manuals for Stations BRG and BRM were revised, effective April 1, 1962, to conform to IAEA inspection requirements; however, they were not issued, pending clarification of some policies during the first official IAEA inspection. Two inspectors, Dr. Michael Hignatsberger and Mr. Carlos Buchler, arrived on May 30, 1962, the day preceding the effective date of the agreement, and discussed procedural arrangements and some policy agreements. On June 1 they verified the status of the nuclear materials inventories and the reactor facilities for both the MRR and the BGRR. They returned on June 14 to observe the charging and discharging of fuel elements during a routine shutdown of the BGRR.

Under the terms of the Four Reactors Agreement, a substitute must be provided for the reactor-produced plutonium removed from any of the four facilities, usually in spent fuel shipments. The AEC has designated Brookhaven National Laboratory as the "bank" for such replacement material and has supplied one kilogram of plutonium-239 to cover such removals from any of the four reactors involved.

Irradiation Services and Isotope Production

As shown in Table 4, the volume and income involved in providing irradiation services and

Table 4

Summary of Irradiation Services to Outside Users
(Handling and Other Charges Included)

	Fiscal 1962		Fiscal 1961		Fiscal 1960	
	Number	Volume, \$	Number	Volume, \$	Number	Volume, \$
Reactor irradiations	641	75,918	828	84,858	1,025	54,294
Processed radioisotopes	349	59,926	330	64,490	325	51,133
Research subsidy	—	—	—	(42,095)	—	(33,183)
Co ⁶⁰ sources	3	3,077	6	6,098	5	8,186
Cyclotron irradiations	12	2,342	62	9,885	22	2,854
Total	1,005	141,263	1,226	165,331	1,377	116,467

Table 5

Types of Customers Receiving Shipments

	Fiscal 1962		Fiscal 1961		Fiscal 1960	
	% of Customers	% of Shipments	% of Customers	% of Shipments	% of Customers	% of Shipments
Hospitals	22	51	20	39	18	40
Industrial	36	24	39	30	42	31
Universities	26	19	24	24	24	20
Government	8	3	7	4	7	5
Foreign	8	3	10	3	9	4

Table 6

Production of Special Isotopes by Hot Laboratory Division

	Fiscal 1962				Fiscal 1961				Fiscal 1960			
	No. of shipments		Activity, mC		No. of shipments		Activity, mC		No. of shipments		Activity, mC	
	On site	Off site	On site	Off site	On site	Off site	On site	Off site	On site	Off site	On site	Off site
Mg ²⁸	2	169	0.1	25	—	142	—	32	3	167	0.15	32.8
I ¹³²	—	71	—	1480	—	111	—	1830	7	70	70	1150
Y ⁹⁰ beads	—	60	—	2233 ^a	1	33	1 ^a	1014 ^a	—	15	—	431 ^a
Tc ^{99m}	—	20	—	2000	—	5	—	500	—	—	—	—
K ⁴³	19	1	9.1	0.1	—	—	—	—	—	—	—	—
Mo ⁹⁹	—	15	—	1010	—	3	—	30	—	—	—	—
I ¹²⁴	10	—	65.8	—	1	1	25	1	—	—	—	—
Y ⁹⁰ generators	—	9	—	1000	—	9	—	1100	—	—	—	—
Ar ³⁸	—	8	—	0.24 ^b	—	10	—	0.36 ^b	—	—	—	—
I ¹³³	—	1	—	20	8	27	150	540	7	69	46	2400
Te ^{129m}	—	1	—	1	—	—	—	—	—	—	—	—
F ¹⁸	—	—	—	—	—	2	—	64	2	6	16	48
Cu ⁶⁷	—	—	—	—	—	—	—	—	—	1	—	1

^aIndividual beads, not mC; activity per bead controlled by length of irradiation requested by customer.^bAr³⁸ is stable; unit is cc, not mC.

processed isotopes to outside organizations have shown a general decrease since last fiscal year. This decrease was probably the net result of several factors, namely, the cessation of the AEC research subsidy last year, the increased availability of similar services commercially, and the increase this year in BNL service rates. The distribution of these services by type of customer is given in Table 5.

Revised prices for BGRR rentals and service irradiations became effective for internal users on July 1, 1961, and for external users on August 10, 1961. The increases of $\approx 11\%$ in the rental rates and of 15% to 50% in the various service irradiation rates reflect the changes in anticipated costs and usage in the budget estimates for fiscal 1962 and 1963.

A system of costing for irradiation services was developed which related the cost assignment to volume of services performed. This system made it possible to follow fluctuations in the usage of the reactor for irradiation services, which is computed as a 12-month moving average. The over-all decrease for the year amounted to $\approx 20\%$.

Medical Research Reactor

The MRR was operated on 256 occasions during the year and 297,490 kw-hr of operation were accumulated. The total integrated energy to date is 1,141,470 kw-hr. The reactor is operated only upon written request from the experimenter to the Medical Department's Reactor Programmer, and each planned experiment is evaluated for reactor safety by the Reactor Operations Group and for personnel safety by the Medical Health Physics Group before permission is granted to conduct the experiment.

There were two unscheduled shutdowns of the MRR during the year. Water leaking into one of the control rod magnets caused one shutdown, and an inadvertent bump given to a mercury switch in the water-flow safety switch caused the second. The resultant delays were of minor duration.

No sticking or unsafe failure of the control rods occurred during the year. Semiannual inspections conducted in November 1961 and March 1962 revealed the rods to be in excellent condition. Some minor modifications were made in the rod coupling magnet enclosures to prevent accidental shutdowns due to leakage of water into these enclosures.

In April 1962 a Tracerlab, Inc., MWP-1A fission products water monitoring system was

installed in the primary coolant system of the MRR. Preliminary investigations indicate that the system will operate satisfactorily and that any significant increase in the fission product level in the primary coolant will be detected. Final evaluation of the system awaits permanent placement of indicators and recorders and additional system operating experience.

Installation of an electrically operated, remotely controlled target conveyor in the animal treatment facility was recently completed. This conveyor will accommodate a wide range of materials, including small animals, and will permit long-range programs to be carried out without unnecessary exposure of the experimenter while setting up and retrieving his target materials. Research programs utilizing this conveyor are in the planning stages.

HOT LABORATORY OPERATIONS

Radioisotope Production

The Hot Laboratory Division produces routinely certain isotopes that require chemical processing following irradiation. Noteworthy accomplishments this year include the first shipment of K^{43} of high ($>90\%$) radioisotopic purity, the first shipment of one of these processed isotopes (10 mC of I^{132}) to South America (specifically, to Rio de Janeiro), and attainment of a new high (40 mC/g) in the specific activity of Mg^{28} . In addition it is interesting to note that this year shipments were made of Ar^{38} to Australia, Switzerland, and France, of Mg^{28} to France and Canada, and of I^{133} to Sweden. Simultaneous irradiation of multiple targets is now being used routinely to reduce isotope production costs; this procedure was first used in an attempt to offset the temporary decrease in demand for isotopes caused by termination of the AEC research subsidy. Table 6 summarizes the routine production of special isotopes.

HEALTH PHYSICS

The over-all program of the Health Physics Division has been similar to that of recent years except for increased emphasis on monitoring at or beyond the site boundary and on dosimetry at the Alternating Gradient Synchrotron (AGS). The total staff, which includes those engaged in general safety and waste disposal and reclamation activi-

Figure 2. Depth dose in paraffin phantom due to high energy beams and stray radiation fields at the BNL 33-Bev Alternating Gradient Synchrotron; preliminary data, December 1961 to May 1962.

ties, has increased from 62 to 66. New facilities include a 400-channel gamma spectrometer for use primarily in bio-assay and area monitoring, new film processing equipment to replace that provided when the personnel monitoring program was initiated 14 years ago, and a shot-blasting room for the decontamination and reclamation center. A stronger Co^{60} source and improved arrangements have been provided for the high-level instrument calibration facility.

AGS Dosimetry

The greatly increased experimental activity and beam intensity at the AGS have required a continuation of dosage measuring and evaluation activities. The exposure situations encountered are extremely diverse, varying from familiar gamma-ray and neutron fluxes to multi-Bev fluxes of μ -mesons, protons, π -mesons, and exotic mixtures of these and other particles. Because of the diversity of hazards and the huge scale of operations, it was decided last fall that continuous coverage by health physics technicians should be provided for both the Cosmotron and the AGS, and the training of

a staff for this purpose has been a major undertaking.

In addition to the performance of routine monitoring and other services, active investigation was made of three dosimetry problems that are worthy of brief mention. The first concerns a region where a high energy flux of μ -mesons emerges from a 16-ft-thick wall of heavy concrete. The μ -mesons are accompanied by a complex spectrum of secondaries, some of which give tracks on the neutron monitoring (NTA) film used for personnel monitoring even though the μ -mesons themselves cause no tracks. Preliminary data indicate that the present system of interpreting personnel monitoring results does not underestimate the dose, but a continuing study in this area is planned.

Another problem of interest is the build-up of dose rate as high energy fluxes pass through the body. Considerable data, similar to the representative cases shown in Figure 2, have already been accumulated by using paraffin phantoms. In general, a substantial build-up is found for pure beams of high energy particles where lower energy components have been removed. Thus curves A and B represent 4.5 and 1.0-Bev/c beams of protons and π^+ -mesons. Curves C and D were taken at locations where high energy components can penetrate but where lower energy components have not been removed for experimental reasons. Curves E and F are for places where scattered radiations of lower energy predominate. Here any build-up due to high energy components is outweighed by absorption of those with lower energy. Knowledge of these build-up curves is essential for proper interpretation of the results of instrument surveys.

A third problem of major importance is the evaluation of dose rates resulting from activation of targets and machine components. Such activation is greatly increased by the strong-focusing feature of the AGS. Surface dose rates of the order of 1000 rad/hr have already been encountered, and even higher values are anticipated.

Waste Disposal and Reclamation

Use of the decontamination and reclamation center increased considerably during the year, with 375 batches of items ranging from hand tools to 2000-gal tanks being decontaminated. Also cleaned and reconditioned were 5000 lead bricks and 1230 heavy-concrete bricks. Three men are employed full time in this work, which has been quite profit-

able in terms of dollars saved. The facility is also used by many Laboratory groups as a "hot work" area for removal and processing of irradiated samples or for their own decontamination activities.

Solid and liquid wastes were packaged in 10-ton concrete vaults as in recent years. The processing of 61 such vaults resulted in an accumulation of 204 at the year's end; 5 vaults in poor condition were shipped to Oak Ridge for disposal. The process of solidifying the radioactive slurry from the evaporation plant was found to be greatly improved by the use of vermiculite. The amount of cement required per vault was reduced from 65 to 24 bags, and stresses in the concrete were reduced. By adding the slurry from the bottom of a dry cement-vermiculite mixture it has been possible to dispense completely with stirring.

Data relating to the liquid waste system are presented in Table 7. The amount of activity leaving the site in the stream to which the effluent is discharged totaled 175 mC during the fiscal year, and the average concentration was 1.24×10^{-7} $\mu\text{C/cc}$.

Area Monitoring

Monitoring of radiation levels at and beyond the site boundary was somewhat intensified as a result of the Russian weapons tests last fall. The average fallout rate in rain and settled dust was 42,990 dis/min/m² per week for the year ending June 30, 1962, or nearly 30 times that for the previous year. The background dose rates recorded at the two off-site monitoring stations increased gradually during the year from an initial value of 1.86 to a current value of 3.6 mr/week. Filterable activity in the air due to fallout averaged 32 $\mu\text{C/m}^3$ during the last half of September but has averaged 7.8 $\mu\text{C/m}^3$ since then, as compared with a barely measurable value of 0.1 $\mu\text{C/m}^3$ or less last year.

Because of the more stringent limits on off-site levels of I¹³¹ recommended by the Federal Radiation Council, increased attention was paid to iodine monitoring of the reactor stack effluent as well as milk and vegetation samples. About 10 mC/day of this isotope is discharged. Measurements of vegetation and milk samples in the vicinity and an analysis of the atmospheric dilution from the stack to the site boundary indicate that average levels at and beyond the boundary are <1% of the prescribed occupational limit and therefore do not

Table 7
Summary of Liquid Waste Data

	Fiscal 1962	Fiscal 1961
Input to filter beds, gal/day	612,000	519,600
Output from filter beds, gal/day	537,000	395,800
Net loss in filter beds, %	11.8	23.8
Stream above discharge point, gal/day	450,000	288,900
Stream at site boundary, gal/day	1,053,000	652,600
Rainfall, in./mo	3.6	3.6
Activity concentration at input to filter beds, C/cc	3.2×10^{-13}	7.9×10^{-13}
Activity concentration at output from filter beds, C/cc	2.4×10^{-13}	3.9×10^{-13}
Reduction in activity concentration, %	25	50
Activity at input to filter beds, mC/mo	22.6	45.1
Activity at output from filter beds, mC/mo	15.1	17.9

Figures are averages for each fiscal year.

require detailed monitoring. Such measurements were hampered last fall because I¹³¹ due to weapons tests was far in excess of that due to BNL operations.

Interest in gamma-ray propagation was stimulated by the necessity of stipulating the size of a Cs¹³⁷ source that could be used by the Biology Department for the new gamma forest. On the basis of field measurements and a theoretical treatment, a source of 9000 curies was set up. The radiation dose rate at the edge of the site, 760 meters distant, amounts to ≈ 150 mr/year or 30% of the off-site limit. Suitable fences and interlocks have been provided, and appropriate administrative procedures have been established to insure safe operation of the forest.

Personnel Monitoring

The workload of the personnel monitoring group continued to increase markedly, mostly because of the expanding programs of research at the synchrotrons. Film dosimeters were processed at the rate of 1214 per week during calendar 1961, an increase of 26% compared with that for 1960.

Table 8

Distribution of Exposures of Individuals Receiving
Regular Personnel Monitoring Service

Exposure index range*	Calendar 1961	Calendar 1960
0.0-0.4	2187	1959
0.5-0.9	106	77
1.0-1.4	28	23
1.5-1.9	20	25
2.0-2.4	14	9
2.5-2.9	6	5
3.0-3.4	9	4
3.5-3.9	11	4
4.0-4.9	15	11
5.0-5.9	2	15
6.0-6.9	0	8
7.0-7.9	0	4
8.0-9.9	0	3
10.0 or greater	0	0
Total	2398	2147

*Exposure index = $\frac{1}{2}$ of β exposure in rad + γ exposure in r + neutron exposure in rem.

Average exposure per individual decreased somewhat compared with that for the previous year, and, despite the increased scale of operations, there was a large drop in the number of exposures >5.0 . The distribution of exposures for the 2398 individuals regularly monitored is shown in Table 8.

General Safety

The Laboratory completed two periods of one million man-hours without an accident involving absence from work during calendar 1961. The accident frequency rate was 3.4 per million man-hours worked as compared with 2.9 for 1960. The accidents were considerably less serious, on the average.

MECHANICAL ENGINEERING

The level of activity in the past year in the Mechanical Engineering Division has increased slightly, but the rate of addition of personnel has decreased. The group showing the greatest increase is that performing work for the Physics Department. Replacements are expected to offset small decreases in the size of other groups. The present complement of the Division (138) is outlined in Table 9.

Division personnel assigned to the Bubble Chamber Group are concerned mainly with the largest construction project, the 80-in. liquid hydrogen bubble chamber. Design of this chamber and the fabrication of most of its major components have been completed this year. Very large components are involved; for example, the bubble chamber magnet will be a cube of steel columns roughly 25 feet on a side fabricated to accuracies of 0.005 in. and fastened with bolts of 3½-in. diameter. This assembly houses a vacuum chamber that is a stainless steel weldment fabricated to pressure vessel standards and finished to high tolerances — one of the largest to be fabricated to such high standards.

The coil to power the magnet has a 2×2-in. copper cross section (the largest cross section of hollow conductor copper used so far at Brookhaven) consolidated in a vacuum-impregnated epoxy casting to make a structure capable of withstanding the magnetic forces in the magnet.

Within the vacuum chamber is the liquid hydrogen container made of nonmagnetic stainless steel castings and weldments also of pressure vessel quality. The chamber surfaces are finished to very close tolerances for the liquid hydrogen seals.

The completed structure, which will weigh 500 tons, can be moved to different positions on a track 70 ft long; it also can be raised 2 ft and rotated to varying angles in azimuth. Design and fabrication necessitated the development and utilization of techniques at the limit of technology available for large-sized equipment.

The other components for the 80-in. bubble chamber are nearing completion. The expansion system has been in prototype operation to test the life of valves and components under operating conditions. The refrigerating system has been completed by the fabricator and is being shipped. The building for the chamber was occupied on April 1st, and installation of the auxiliary equipment such as compressors, controls, and finally the large magnet components is proceeding.

With the increase in number of photographic recording chambers, picture analysis has become a major problem. Design and development of components for the flying-spot digitizer has proceeded. In particular a film transport system for 35-mm film has been designed and completed, and work is in progress on a similar device for 70-mm film. The development of new devices for data reduc-

tion and analysis will continue at an increasing rate.

Other Division members have been active in operating phases of the BNL 20-in. liquid hydrogen bubble chamber, which has been in almost continuous operation at the Cosmotron and the AGS. Personnel assigned to the Columbia-BNL 30-in. chamber have successfully completed the erection and testing of this chamber and are preparing it for the first experimental run.

Members of the Division assigned to the Nuclear Engineering Department have rendered engineering assistance, in cooperation with the Department's engineers, on various departmental projects such as exponential and critical physics experiments for many kinds of uranium-moderator lattices and gamma-irradiation facilities for source development and for radiation experiments on foods and chemical substances. Work has also been done on liquid metal heat transfer research equipment for both heavy and light metals and on corrosion-testing facilities, particularly for liquid metals, for metallurgical research. In addition, work has been done on equipment and machinery for severe service in radioactive waste treatment and reactor fuel processing procedures constantly being developed for engineering evaluation of future trends.

Some Division members are connected with the HFBR Project staff, which is responsible to the AEC for technical administration and approval of all plans and specifications and for examination of field work. Engineering service has been rendered in the preparation of manuals and test programs for the Reactor Division, and assistance is being given in the areas of operator training and spare parts procurement. This has also involved the design and procurement of many items of auxiliary equipment needed for operation and fuel handling.

The Division's members have provided extensive assistance for the HFBR Project in administration, programming, preliminary and final design, and initiation of procurement. They have also been involved in the construction, inspection, and operation of the reactor, and in the development of components and selection of materials.

Division members maintain some activities in cooperation with ANS standards committees and ASME committees.

The Division's work at the Cosmotron has been divided amongst the areas of machine improve-

Table 9

Assignment of Mechanical Engineering Division
Personnel as of June 30, 1962

Department	Engi- neers	Designers and Draftsmen	Secretarial and Clerical
Accelerator			
AGS	21½	18	½
Cosmotron	11	8	—
Central Design	½	8	1½
Nuclear Engineering	6	18	1
Physics			
Bubble chambers	16	11	—
General design	4	11	—
Reactor	—	2	—
Total	59	76	3

ment, experiment support, and liquid hydrogen production and use. The major machine improvement was the installation, after completion of design and fabrication, of the new magnetically shielded straight sections during an extended shutdown involving the removal of major components from the Cosmotron. Other improvements made to the machine included strengthening the pole-face winding attachments and leveling the magnet. The three new straight sections have independent vacuum systems using 10-in.-diam PMC oil diffusion pumps. This has resulted in a greater uniformity in high vacuum around the Cosmotron ring.

Extensive modifications are being made in the Van de Graaff injection system. New ion sources, an electrostatic beam chopper, and beam energy modulator rf cavities were designed and built for future installation. A 14×17-in. spark chamber with an associated liquid hydrogen target to operate in the field of the 14-in. bubble chamber was completed, and a redesigned beryllium external beam monitor prototype has been built and tested. Five additional units are under construction. Six adjustable magnet stands to permit accurate positioning of large bending magnets were built.

Members of the Division installed facilities for fabricating Mylar liquid hydrogen targets, and several Mylar and stainless steel targets were designed and built for use at the AGS and the Cosmotron. An 18-in.-diam by 6-ft-long liquid hydrogen target for the AGS was procured and tested.

The hydrogen liquefier facility under the supervision of Division personnel produced 120,000 liters of liquid hydrogen despite two 6-week shut-downs for repairs. A deuterium hydrolysis plant from the Bureau of Standards at Boulder, Colo., is being installed at the liquefier facility. When this is completed shortly it will produce 30 ft³ of deuterium gas per hour.

A group from the Division continued to assist experimental physics groups in setting up and running experiments in the six beams of the Cosmotron. Two 2500-amp, water-cooled electrical cables for energizing bubble chamber magnets were installed along the East and North walls of the Target Area, and eight new air-conditioning units were installed inside the Cosmotron radiation shield to cool the machine area. The radiation shield was modified to provide an access labyrinth.

A major project has been the design and fabrication of six universal neutron spectrometers for the use of the Physics and Chemistry Departments at the HFBR. These spectrometers vary in some particulars, but standard designs and components serve as the core of the instrument construction. Design and construction were completed of a student research neutron spectrometer for the Puerto Rican Nuclear Center.

Two additional dual scanner units were constructed for use in scanning 35-mm bubble chamber film, and an instrument for determination of gap length distribution in bubble chamber tracks was designed and fabricated.

A high-pressure Čerenkov counter for identifying low velocity particles at the AGS was manufactured, and a 10-gap neon spark chamber was completed. An interesting feature of the design was the use of 0.005-in.-thick concentric cylinders of aluminum for energy loss plates.

Preliminary mathematical analysis and design were started on a 3-rotor, slow-neutron chopper system for the HFBR. This involved the determination of stresses and optimization in design of a high-speed composite rotor.

Other items worked on were cryostats, goniometers, an ultracentrifuge, a crystal growth furnace, and liquid Čerenkov counters.

With the AGS in operation for 15 shifts/week, much effort has gone into experiment support and machine operation, and Division members have worked with the experimenters and AGS liaison

personnel to set up and modify experiments in the experimental area.

Operating manuals have been written and instruction schedules devised for maintenance of the mechanical equipment at the AGS by Division personnel.

Additional details of the Division's extensive work at the AGS are given in the section on High Energy Accelerators.

Personnel assigned to the Central Design Group were involved in the design and installation of equipment for the Physics, Biology, Chemistry, and Medical Departments and the Health Physics Division. The vertical arm spectrometer was completed and installed on the roof of the BGRR, and several other spectrometers and auxiliary equipment, including a crystal rotation device, a crystal holder, a 500°C oven, and several cryostats, were also completed. A machine to grind up whole animals in seconds was completed and tested for the Biology Department, as well as devices and instruments needed for the ecology radiation field. A low-velocity Čerenkov counter for use at the AGS, several shielded chambers made from 30-in.-diam gun barrels, a 10 and 50-curie irradiation machine for the Health Physics Division, and a modification of the hydraulic press used in rupturing living cells were designed. Central Design personnel were involved in the development of the Hough-Powell automatic flying spot digitizer being constructed for use with the IBM 7090 computer.

MACHINE SHOPS

The total number of productive hours for the Machine Shops Division decreased for the second consecutive year despite continually increasing demands. Although additions to the staff have been made, the rate of increase has not been sufficient to keep pace with the workload. Furthermore, most of the additional personnel have been assigned to nonproductive clerical, technical, and administrative positions. Whenever the nature of the work has permitted, and with the requisite approval, jobs have been assigned to outside contractors. The contracting activity has increased considerably during the past year and is expected to continue at an accelerated rate in the coming year.

No significant change has been noted in the character or distribution of the workload within

the Division, although a somewhat larger number of individual jobs was handled than in previous years. Work processed through intralaboratory requisitions accounted for 90% of the total productive hours, the remaining 10% being used for jobs undertaken on a short-order basis. The total of 119,144 hours was $\approx 5\%$ less than in fiscal 1961. The average backlog was somewhat smaller than in fiscal 1961, but still was almost 18,000 hours per month, with the sheet metal and welding group continuing to show a higher proportion of backlog than of productive hours worked.

The contributions of the scientific departments to the Division's workload are indicated in Table 10. Of most significance is the continued drop in the proportion of work done for the Nuclear Engineering Department, from 36.6% in fiscal 1960 to 28.6% for fiscal 1962, and the marked increase in that done for the Physics Department, 30.4% in fiscal 1962 as against 11.2% in fiscal 1960. Much of the heavy workload for the Physics Department can be directly attributed to the 80-in. bubble chamber project. Over 75% of the Division's workload was contributed by the Nuclear Engineering and Physics Departments and the Cosmotron.

Considerable emphasis was placed on improving existing service activities. In cooperation with the Supply and Materiel Division a modern cutting shop was established at the metals warehouse to provide fast cutting service and, by cutting "to tolerance," to eliminate as nearly as possible the costly double-cutting operations previously necessary.

The work of inspection and quality control continued to grow during the year. Additional equipment acquired included Zyglo tanks and fixed and portable Magnaflux units. The only major nondestructive testing procedure not in current use is x-ray. The unified thread system was adopted as the BNL standard and a full range of Class 2 and 3 gauges purchased. The system of inspection reporting was completed in cooperation with the Mechanical Engineering and Purchasing Divisions, and work is continuing on other internal inspection systems and procedures.

In line with organizational changes planned in fiscal 1961 a Shop Services Section was established under the supervision of an assistant foreman as the first step in relieving the first-line foremen of as much paper work as possible and freeing them to devote more time to the scheduling of workloads and direct supervision of work in progress.

Table 10
Contributions to Machine Shops Workload

	% Man-hours		% Change in workload
	Fiscal 1961	Fiscal 1962	
Accelerators			
AGS	8.6	8.6	—
Cosmotron	14.3	16.5	+ 15.4
Biology	3.1	2.1	- 32.3
Chemistry	3.4	2.5	- 26.5
Instrumentation and			
Health Physics	0.7	0.6	- 14.3
Medical	3.1	3.3	+ 6.5
Nuclear Engineering	33.3	28.6	- 14.1
Physics	25.1	30.4	+ 21.1
Reactor	6.6	4.4	- 33.3
Miscellaneous	1.8	3.0	+ 66.7

Equipment worth $\approx \$175,000$ was acquired for replacements and additions. Of this amount $> \$67,000$ was acquired from government surplus, and the balance represents cash purchases of new equipment. Major items included a Thompson surface grinder, a Thompson broach grinder, a 36-in.-high $\times$ 18-ft-long Lodge & Shipley lathe, a new Kearney & Trecker Model 205S-12 horizontal universal milling machine, a Wysong & Miles Model 1252 shear, and a Lodge & Shipley Model 0810 shear with a $\frac{1}{2}$ -in. $\times$ 10-ft-wide capacity. Also included were a number of smaller pieces of equipment. Special emphasis was placed on new machines for the cutting shop and the close-tolerance machine shop.

Some of the work done during the year was of particular interest. For example, extremely close coordination between the Mechanical Engineering and Machine Shops Divisions was required in the development of a source assembly consisting of four encapsulated cobalt strips, two D-shaped filler bars, and an aluminum envelope; the maximum radial gap of 0.004 in. for the 12 interfaces was the critical requirement here. Also of interest were the fabrication of a vertical arm spectrometer, the critical assembly and welding of the bellows seal for the 80-in. bubble chamber, the fabrication of shipping casks for Mark I and Mark IA sources, and the continuing production of very precisely contoured tensile specimens for use in the study of neutron irradiation effects on body-centered-cubic

metals. The construction of a slow-neutron chopper and an in-pile high temperature furnace, the salvaging of quantities of sintered uranium oxide pellets, and the fabrication of an aluminum spark chamber and liquid hydrogen target for a *K*-meson experiment at the Cosmotron also exemplify the wide variety of fabrication problems encountered in more than 1650 jobs in fiscal 1961.

In addition, jobs were completed for Argonne National Laboratory, Columbia University, New York University, Rensselaer Polytechnic Institute, the University of Pennsylvania, and the University of Wisconsin.

TECHNICAL INFORMATION

Toward the end of the fiscal year, construction was completed on a wing at the west end of the Research Library Building. This additional space has been provided for the Information Division's functions in report processing, report distribution, and technical editing, which were formerly conducted in a separate, ex-Army building. The benefits accruing from housing under a single roof these various related activities in the field of technical information are already evident. In addition, the new wing provides the Library with a new micro-card reading room and a room with four study carrels.

The Library has recently given considerable attention to the improvement of staff services. An important organizational step in this direction was placement of the Medical Department Library under the direct supervision of the Research Library administration. This arrangement is expected to provide an improved book collection, a more efficient interlibrary loan service, and a coordinated reference service. A significant increase in the use of library services and materials has been noted. About 2000 more books, journals, and reports were borrowed this year than in the past year. The demand for reference services was also greater, particularly in the form of requests for bibliographies and literature searches. The need for translations of Russian scientific literature has placed a continuing demand on the time of the one member of the reference staff qualified to provide such service.

In the Classified Library, the receipt of classified reports has continued at about the same level. Of ≈ 700 documents accessioned, 199 are secret. No

classified reports were published by the Laboratory during the past year. An inventory conducted early in 1962 revealed holdings of 14,269 classified documents, of which 8059 are secret. A visit last winter by three members of the AEC's Declassification Branch resulted in the declassification of ≈ 2000 pieces of correspondence. It is hoped that subsequent reviews of this kind can be arranged, in order to reduce further the Laboratory's classified holdings.

Another activity in the field of technical information is the processing of all manuscripts originating from BNL-sponsored research for publication in the scientific and technical literature. As an index to the Laboratory's increasing output of such papers, the Information Division processed during the year a total of 710 items (including published abstracts of speeches), as compared with 708 in the previous year, 607 in fiscal 1960, and 486 in fiscal 1959.

Sixty-three BNL technical reports were processed by the Division and given the appropriate distribution. Most of these reports are edited and otherwise prepared for publication by the Editorial Section, which works closely with the Photography and Graphic Arts Division in ensuring that the Laboratory's formal technical publications meet the highest possible standards in format, style, and accuracy.

During the year, the BNL Patent Office was made a part of the Information Division. This is a logical arrangement, since the Patent Office must review all the scientific manuscripts originating from BNL-supported research in order to report to the AEC all discoveries and inventions contained in such documents, prior to their release for publication. During the current year 768 docu-

Table 11

	Fiscal year		
	1962	1961	1960
Photographs	11,062	8,758	9,133
Photomicrographs	2,768	2,167	1,898
Lantern slides	10,371	9,271	7,576
Prints	52,212	44,300	46,747
Photostats, Xerox	46,291	31,928	32,698
Ozalid prints	506,128	597,205	502,227
Film processed, ft	1,409,730	1,453,018	196,532
Charts and graphs drawn	3,982	3,581	3,058

ments were processed, which resulted in the preparation of 40 preliminary invention reports and 19 records of invention. Seven inventions became the subject of patent applications during this period. These statistics are representative of the normal workload over the past few years and are comparable with those for 1961, during which 10 patent applications were filed, 760 documents were processed, and 40 preliminary reports and 26 records of invention were submitted. In 1960, the processing of 645 documents resulted in 10 patent applications, although 72 preliminary invention reports and 22 records of invention were submitted.

PHOTOGRAPHY AND GRAPHIC ARTS

The Photography Group has experienced an increase in workload, as is evident from the production figures given in Table 11. The recording of component construction for the 80-in. liquid hydrogen bubble chamber at a variety of locations has added to the scheduling problem.

New equipment includes a fluorescence microscope and additional electronic flash sources for the studio. A microdensitometer for the evaluation of bubble chamber films and other microimages has been acquired. This device should prove useful in a number of fields.

The continuous film processor has operated well and was able to handle a load of 1.4 million feet which was concentrated almost entirely in the second half of the year. The sensitometric control procedures have ensured a uniform process, and the amount of footage lost (0.035%) was largely attributable to mechanical defects in the exposed film.

A motion picture on 16-mm color film describing aspects of research in high energy physics at the AGS is being produced. With the excellent cooperation of the Physics Department and the AGS

Table 12

	Fiscal year		
	1962	1961	1960
Offset pages	11,207,710	10,294,755	9,921,637
Film negatives and positives	10,610	9,896	9,050
Mimeo impressions	2,283,530	1,769,058	1,560,263
Stencils run	15,104	13,613	11,530
Sheets collated and bound	3,363,100	2,819,350	2,996,192
Report copies issued	100,725	92,745	115,473

the script has been written and footage has been completed, and the film is expected to be ready in the fall of 1962.

During the past year the Graphic Arts Group has experienced a continued increase in its printing and duplicating activities. Comparative figures for the past three years are presented in Table 12. In an effort to keep abreast of the requirements of the Laboratory, the Group has consolidated all its operations in one building and has added another person to the staff.

The Ektalith equipment and the new automated Multilith press have begun to function well, and some work is being diverted from the Ozalid process to this better and less expensive process.

The two Fotosetters continue to be operated for four of the possible six shifts, and a fifth will be added in the near future. This will effectively represent maximum utilization of the machines, since the remaining shift must be kept available for maintenance.

The Group has continued to investigate the possibilities of tape operation of the Intertype Fotosetter. Discussions have been held with the company and the prospects for installation of such equipment in the future seem good.

Administration and Operations

The management operations reviewed in this section have been carried out in a manner designed to facilitate and encourage the research activities of the Laboratory's scientific staff.

PERSONNEL

Scientific Staff and Students

During the 12 months ending May 31, 1962, the number of individuals holding appointments to the scientific staff remained relatively constant. Comparative year-end statistics shown in Table 1 do not reveal the characteristic high turnover; 42 scientists and engineers were appointed to the regular staff, and 40 terminated or were granted leaves of absence.

For the first time in seven years the number of salaried visitors decreased. This was due, in part,

to a temporary "leveling-off" of the number of postdoctoral Research Associates. Turnover in this category is always high because of the limited term of these appointments. During the year, 49 scientists began salaried appointments as Research Associates and 51 completed their terms. In addition, 21 scientists began nonsalaried postdoctoral appointments, and 8 terminated or were transferred to other categories. Of the 105 Research Associates (salaried and nonsalaried) holding appointments on May 31, 50 were foreign scientists.

Although the number of nonsalaried visitors continued to increase, the number who were participating at the Laboratory on a full-time basis remained the same as last year. Graduate and undergraduate students held 46 of the 73 additional nonsalaried appointments made this year.

Table 1
Scientific Staff and Students on May 31, 1961 and 1962

	Regular staff		Visitors			
			Salaried		Nonsalaried	
	1962	1961	1962	1961	1962	1961
By appointment category						
Staff						
Senior Scientist	52	52	1	0	9	11
Scientist	112	103	0	7	115	120
Associate Scientist	107	112	10	7	119	104
Assistant Scientist	65	67	6	10	93	87
Research Associate	—	—	76	78	29	16
Students						
Junior Research Associate	—	—	10	7	57	41
Research Assistant	—	—	0	0	57	27
Total	336	334	103	109	479*	406**
By academic degree						
Ph.D. or M.D.	204	203	92	99	317	287
Master	55	55	9	7	73	55
Bachelor	72	71	2	3	82	57
No degree	5	5	0	0	7	7

*65 of these appointees were at BNL on a full-time basis as of May 31, 1962.

**65 of these appointees were at BNL on a full-time basis as of May 31, 1961.

Table 2
Classification of Visiting Scientists and Students Participating in BNL Program
for One Month or More, June 1, 1961 – May 31, 1962

	Guests and salaried visitors				1961 Summer program		Total	
	More than 3 months		Less than 3 months					
	Salaried	Guest	Salaried	Guest	Salaried	Guest	Individuals	Institutions
University staff	15	34	6	33	45	47	180	88
Thesis students	10	18	0	14	0	3	45	21
Student Research Assistants	1	7	0	11	89	52	160	55
Subtotal	26	59	6	58	134	102	385	115 different
Industry	0	11	0	2	0	1	14	10
Other institutions	13	51	3	12	12	13	104	60
Total	39	121	9	72	146	116	503	185 different

Since May 31, 1957, the numbers of salaried and nonsalaried visitors have nearly doubled. During the same period the regular scientific staff has grown from 281 to 336, an increase of $\approx 20\%$.

May 31, instead of June 30, was selected as the date for Table 1 to exclude the large number of visitors who were at Brookhaven only during the summer. Similarly, the year ending May 31 was used in Table 2 to include participants in only one summer program.

Table 2 lists the number of visiting scientists (exclusive of salaried Research Associates) and students who participated at the Laboratory for a cumulative period of one month or more between June 1, 1961, and May 31, 1962. The total of 503 visitors is the largest in Brookhaven's history and compares with 450 during the previous year. Fifteen graduate students completed the research required for their doctoral theses during the year.

In addition to those included in Table 2, 219 scientists and students worked at Brookhaven for less than one month.

Table 3 shows the extent to which consultants' services have been employed during each of the past three fiscal years. The use of these specialists increased to the highest level since 1954.

Summer Program for 1962

Arrangements have been made for 270 visiting scientists and students to work at Brookhaven during the summer of 1962. Of this number, 132 are

staff members from colleges, universities, industrial organizations, and other institutions; and 138 are students.

Special groups include 91 students participating in the Laboratory's eleventh annual summer student program and 16 Health Physics Fellows.

Employment

During fiscal 1962 a total of 472 employees were hired; of these, 379 joined the nonscientific staff and 93, including Research Associates, were appointed to the scientific staff. With 267 separations, the net increase in employees was 205, to give a total for the Laboratory of 2593 employees, excluding temporary appointees, research collaborators, and guests. Employment statistics for fiscal 1962 are compared with those for fiscal 1961 in Table 4.

The need for specialized engineers, technicians, and related personnel continued to be the greatest employment responsibility. To meet it, recruiting efforts were increased at colleges and technical schools. An extensive campaign to employ a large number of Programmers and Mathematics Associates had gratifying results.

The employee orientation program was expanded beyond the area of explaining benefits programs and policies. It now covers a thorough indoctrination in the programs and facilities of the Laboratory. Since the Laboratory does not produce a specific product with which employees can

Table 3

Consultants' Services

	Fiscal year		
	1962	1961	1960
Total contracts in effect June 30	88	79	80
No. of consultants used	59	35	36
No. of man-days of service	577	372	336

identify themselves, it is hoped that this orientation will result in greater awareness of the Laboratory's aims and the individual's part in accomplishing these aims.

In summary, the demand for highly trained technical and professional personnel still exceeds the supply. However, except in a few critical areas, most openings are filled in a reasonable period of time.

Labor Relations

New contracts were negotiated with the Long Island Guards Union (LIGU) and the Oil, Chemical, and Atomic Workers (OCAW) International Union. The agreement reached with the LIGU provided an increase of $\approx 2.5\%$ each year under a two-year contract. Members of the Directly Affiliated Local Union (DALU) No. 24426, AFL-CIO, received an increase of 2.6% under the two-year contract negotiated in 1961 which will expire December 31, 1962.

The agreement reached with the OCAW Union covering 16 reactor operators and 6 Hot Laboratory employees was unique in several aspects. The new contract, effective April 1, 1962, specifies that all employees in this bargaining unit will be treated as nonbargaining-unit wage technicians with respect to matters of Laboratory policy, including technician wage ranges and merit reviews. The arbitration provision of the grievance procedure stipulated in this contract excludes the interpretation of Laboratory policy and the granting or withholding of wage increases. The question of whether or not the Laboratory's personnel policy has been applied consistently is subject to arbitration, as are other express provisions of the contract, including discharge for just cause. The contract has no expiration date, but it may be canceled by either party upon 90 days' notice. However, should the Union terminate the contract, the rates

Table 4

Employment Statistics*

	Fiscal 1962		Fiscal 1961	
	Number	Annual rate (%)	Number	Annual rate (%)
Scientific staff	439		437	
Nonscientific staff	2154		1951	
Total	2593		2388	
<u>Accessions</u>				
Scientific staff	93	21	126	31
Nonscientific staff	379	18	375	20
Total	472	19	501	22
<u>Separations</u>				
Scientific staff	91	21	74	18
Nonscientific staff	176	9	164	9
Total	267	11	238	11
<u>Net Accessions</u>				
Scientific staff	2	0.1	52	1.3
Nonscientific staff	203	9.9	211	11.4
Total	205	8.2	263	11.7

*Figures do not include 100 temporary summer non-student employees. Guests and temporary student employees are included in Table 2.

of pay of individual employees revert to the "standard" rate of the applicable wage range for the classification 60 days after such notification has been given.

An NLRB election was held for the purpose of determining whether the DALU would represent Cosmotron operations technicians. Of 51 eligible voters, 49 voted, 22 for joining the DALU and 27 against joining, and the Union's bid for representation was defeated.

Training and Communications

A revised Supervisors' Personnel Manual was distributed to all supervisors. The number of copies issued was increased to give greater coverage in this area. The program of supervisory seminars was expanded this year to include more than 250 participants, a considerable increase over previous seminars.

Employee Benefits Plans

Because of the amendment to the Social Security Law permitting men to retire at age 62, the Laboratory policy on Group Medical Insurance was altered. Under the new policy, an employee who terminates on or after his 62nd birthday may continue his Group Medical Insurance on the same basis as an employee retiring at age 65.

ARCHITECTURAL PLANNING

During the past year, the increasing pace of the Laboratory's activities was felt in all the areas in which the Architectural Planning Division participates. Eight major construction projects with a value in excess of \$11,000,000 were completed. New construction valued at \$15,000,000 was in various stages of design and engineering, and proposals for new research devices, facilities, and buildings in the amount of \$33,000,000 were prepared. The General Plant Projects program continued at the high level of activity reached the previous year. A number of studies relating to site utilities and services were completed, and to keep pace with the increased level of activity the organization was strengthened and modified.

The construction projects completed during the year were the Physics Building, the Radiation Development Center, quarters for visiting scientists, an addition to the Critical Assembly Facility, an addition to the Metallurgy Building, an addition to the AGS Experimental Area and Service Building, the 80-in. Bubble Chamber Building, and the Volatility Studies Laboratory.

The contracts for architect-engineer and construction work for these projects were held by the AEC and were administered and supervised by its Brookhaven Office. BNL was represented by the Architectural Planning Division, which provided the liaison required during the stages from early planning through construction.

Buildings and facilities under design by architect-engineers were the Chemistry Building, the Health Physics and Instrumentation Building, and a laboratory wing for the Cosmotron. Requirements for the Physics and Mathematics Addition to the Physics Building and the Controlled Environment Laboratory for the Biology Department were reviewed preparatory to development by architect-engineers.

Construction proposals including preliminary building layouts, cost estimates, and descriptive brochures were submitted for the following: Nuclear Engineering Building, Phase II; Hot Laboratory addition and modifications; AGS - North Experimental Area Building and an addition to the Service Building; Medical Research Center - animal laboratories and a self-contained Low Background Counting Facility; an addition to the Radiation Development Center; the High-Pressure Gamma Radiation Loop Facility; an addition to the Steam Plant; General Engineering and Services Center, Phase II; the Tandem Van de Graaff Accelerator Facility; and the Administrative Services Center, Lecture Hall, and Cafeteria.

The General Plant Projects program continued at the high level of activity achieved during the previous year. The projects initiated during the year amounted to \$1,200,000. This represented, however, a greater engineering effort than in previous years in that there were only four individual projects in excess of \$50,000, the balance consisting of miscellaneous modifications, additions, and adjustments to research facilities and supporting staff functions. Many of these projects presented unique engineering problems. As has been the case in prior years, the services of the Plant Maintenance Division were relied upon heavily to coordinate many of the projects.

Some of the projects accomplished as part of this program were an Animal Quarantine Building for the Medical Department, \$134,000; a building addition, storage barn, and miscellaneous facility modifications for the Biology Department, \$146,000; relocation and modification of temporary army buildings for interim use, \$80,000; projects in support of the research devices, including fire protection for the Graphite Research Reactor cooling towers, shielding doors, Cosmotron Equipment Assembly Addition, \$120,000; extension of fire protection in various buildings and implementation of safety recommendations, \$80,000; and installation of air-conditioning systems in buildings of the World War II type now modified as permanent research facilities (such as the Metallurgy Building, the Research Staff Lounge, and the Library), in addition to modifications to air-conditioning systems required for the changing use of various laboratory areas and operating procedures at the Cosmotron, Medical Research Center, Hot Laboratory, and Graphite Research Reactor, \$220,000.

During the year, detailed studies of the site's utilities and services, based on projections of the Laboratory's future requirements, were conducted by outside engineering consultants and were jointly coordinated by the Architectural Planning and Plant Maintenance Divisions. As a result of these studies, plans are being implemented for additional wells to meet the increasing demand for water and for a water treatment plant to eliminate or reduce the severe corrosion problems resulting from the present quality of the water. An addition to the Steam Plant is being requested to serve the Laboratory's needs through 1969. Studies involving sewage treatment, electric distribution, and electric capacity have also been conducted. To coordinate the electric power requirements of the research programs and the general plant requirements, a committee has been established consisting of three members representing the Architectural Planning Division, the Plant Maintenance Division, and the scientific departments.

To cope with the tremendous increase in major Laboratory construction projects, a project coordination group has been created within the organization. This group's prime responsibility is to assist the scientific departments in the development of their requirements and to act as coordinators between them, the Architect-Engineer, and the AEC.

PLANT MAINTENANCE

The functions of the Plant Maintenance Division include the operation and maintenance of all Laboratory utilities, including the Hot Laundry; the maintenance of all buildings, grounds, and vehicles; and the provision of services including communications, mail distribution, housing, and automotive or commercial carrier travel arrangements. In addition to these routine aspects of the Division's responsibilities, the specialized aspects such as facility modification projects, major maintenance programs, plant engineering, special staff services, and research support functions have continued to grow with the growth of the Laboratory.

During the past year, Plant Maintenance acted as coordinator for 116 facility modification projects, totaling almost \$500,000. These projects, funded as part of the General Plant Projects program, were performed by various contractors, with Plant Maintenance providing the engineering and contract supervision. A partial list includes the ecology

field, rehabilitation of an additional dormitory, completion of items connected with the Physics Building, quarters for visiting scientists, the addition to the Metallurgy Building, miscellaneous air-conditioning installations and modifications, relocation and modification of various barracks, and installation of fire detection systems.

During the past year, 122 major maintenance projects were initiated and completed. Again, the Plant Maintenance Division provided engineering and contract supervision for work conducted by outside contractors. Some of the projects undertaken were repainting a 350,000-gal elevated water storage tank; repairing and repainting Meteorology weather towers; refurbishing the interiors of the Biology, Cyclotron, and Graphite Research Reactor Buildings; repairing and resurfacing the road system and railroad tracks; and restoring site areas scarred by building removals and building construction.

The Laboratory's role as host to over 100 scientific seminars, conferences, and meetings annually is supported by the special staff services group which takes care of the transportation, travel arrangements, housing, and other accommodations required by the participants, who come from all parts of the world. While the increase in the number of such functions over the past few years has not been great, there has been a marked increase in the number of participants.

A very large portion of the Plant Maintenance Division's effort continues to be in the area of direct support to the scientific research effort. Increasing use was made of the various skills of craftsmen within the Division, such as electricians, steam fitters, and plumbers, in the setting-up and connecting of experimental equipment and apparatus.

Extensive use was made during the year of the services of consulting engineers in the examination of existing utilities and services to determine their condition and ability to meet the growing needs of the Laboratory. Among the utilities studied were the sewage treatment plant and the steam distributing system. The Division also initiated and participated with the Architectural Planning Division in the study of the water supply system and the attendant problems relating to the quality of the water.

The utilization of manpower and the distribution of costs within the Plant Maintenance Division

Table 5

Manpower Utilization

Type of work	Fiscal 1962		Fiscal 1961	
	Productive man-years	Percent of total work	Productive man-years	Percent of total work
Plant utilities operations	49	16	62	26
Building janitor service	54	18	35	14
Decontamination and hot laundry	14	5	10	4
Total plant operations	117	39	107	44
Communications service	14	7	13	5
Transportation, housing, and staff services	18	8	15	6
Total staff services	32	15	28	11
Special services for others	79	27	43	17
Facility improvements	13	4	5	2
Total special services	92	31	48	19
General repairs and maintenance	37	12	55	22
Special long-term maintenance	9	3	11	4
Total maintenance	46	15	66	26

Because of changes in reporting procedures, detailed comparisons between Fiscal 1961 and Fiscal 1962 are not possible.

Table 6

Costs of Supplies, Materials, and Purchased Labor

	Fiscal 1962	Fiscal 1961
Plant utilities operations		
Fuel oil	\$ 306,114	\$ 263,531
Electricity	1,047,375	975,190
Building janitor supplies	39,825	36,030
Decontamination and hot laundry supplies	17,481	21,860
Total plant operations	1,410,795	1,296,611
Staff services		
Telephone, teletype, and mail	295,176	247,242
Transportation (gasoline)	21,752	22,222
Housing and cafeteria material and supplies	71,867	56,180
Total staff services	388,795	325,644
Maintenance		
General materials	265,717	219,567
Special materials and contracts	383,421	424,579
Total maintenance	649,138	644,146
General plant projects	339,017	—
Grand total	\$2,787,745	\$2,266,401

sion for fiscal 1961 and 1962 are analyzed in Tables 5 and 6.

SECURITY AND PLANT PROTECTION

The Security and Plant Protection Division continued to perform its assigned functions during fiscal 1962. Although the scope of the security function at the Laboratory has contracted substantially since Brookhaven's early days, it has been necessary to maintain an extremely active AEC security clearance program. At the end of the fiscal year more than 1000 Laboratory employees had active "Q" clearance, while over 300 had active "L" clearance. These clearances are required to enable Brookhaven scientific and technical personnel to have full access to restricted data, both at the Laboratory and at other laboratories and institutions. A modified security clearance continues to be required of all Brookhaven employees upon employment. The increasing participation by foreign scientists and students in the Laboratory's research program has served to place greater responsibility upon the Security Office, since a special security clearance is prescribed in such cases.

The physical growth of the Laboratory and the expansion of its research programs have required a greater emphasis on the Laboratory's plant protection program. The increase in this aspect of the Division's functions has been met through the adaptation of industrial plant protection techniques and practices to the specialized requirements of Brookhaven. No additional personnel have been required for the accomplishment of these measures.

The Fire Group maintained its excellent record in meeting potentially dangerous situations at the Laboratory. An active fire inspection and fire prevention program has been instituted as a continuing and important aspect of the Fire Group's activities; close liaison with scientific and technical personnel engaged in research experiments and other activities representing potential fire hazards is regularly maintained, with gratifying results. Fire prevention inspections made by Fire Group personnel during the year totaled 1755, and 459 corrective actions were subsequently taken by operating personnel as a result of these inspections. Fortunately, there were no major fires on site during fiscal 1962; responses to fire alarms, however,

continue to increase in proportion to the Laboratory's growth.

	1962	1961	1960
Responses on site	165	132	103
Responses off site	7	5	5
Investigation - no response	49	34	39
Total	221	171	147

BUSINESS MANAGEMENT

During fiscal 1962 the Brookhaven Business Manager was appointed to another project of Associated Universities, Inc.; later in the year, the Assistant Business Manager was appointed Business Manager with responsibility for the Purchasing and the Supply and Materiel Divisions.

The total number of personnel increased slightly during the year, as shown below.

	Staff	Clerical	Fiscal 1962 total	Fiscal 1962 total
Business Office	3½	2	5½	5
Purchasing Division	14	14	28	28
Supply and Materiel Division	12½	57	69½	68
Total	30	73	103	101

The volume of business transacted this year is compared with that for fiscal 1961 in the following tabulation.

	Fiscal 1962	Fiscal 1961
Number of procurement transactions	32,500	28,000
Value of procurement actions	17,000,000	15,000,000
Number of receiving actions	37,000	35,000
Number of stores issues	130,000	122,000
Value of inventory turnover	2,152,000	1,974,000

In addition, there were 217 transactions by which excess material valued at \$2,393,800 was obtained for the Laboratory at the cost of transportation only.

The procurement forms and procedures instituted last year have worked well, and the 264 charge accounts utilized and the 117 offers of sale negotiated have proved effective in terms of economical buying and improved deliveries. The use of petty cash for purchases of low-cost items available from local sources has resulted in quick action on requisitions and a reduction in handling

and paper work. An evaluation of over-all systems and procedures is being conducted.

The main stockroom of the Supply and Materiel Division is being expanded. Two outlying stockrooms were added during the year, which brought the total of departmentally located, individually manned stockrooms to eight. The receiving section has been relocated in a larger building, and a materials-flow plan has been put into effect to improve the receiving functions, with use of a Xerox machine when additional copies of receiving documents are required. In conjunction with the Machine Shop Division, the Supply and Materiel Division has instituted

an improved metals-cutting operation that has resulted in greater efficiency of service.

FISCAL

During the past year the Fiscal Division completed its data processing study and selected a solid-state magnetic-tape on-line computer system to replace the existing electronic calculator and off-line punch card system, in use for the past 8½ years. The new equipment was received in June 1962, and the conversion program is now in process.

A comparative balance sheet for the Laboratory appears in the Introduction to this report.

Appendix A

PUBLICATIONS, JULY 1, 1961 – JUNE 30, 1962

This list includes official Laboratory publications, abstracts of papers which were or will be presented at scientific meetings, and publications by staff members, consultants, and guests. All these listings result from work done at the Laboratory; they were submitted during the review period.* Abstracts are indicated by (A); letters to the editor, (L); and notes, (N). Acceptance for future publications is designated by (in press).

GENERAL PUBLICATIONS

Annual Report, July 1, 1961. BNL 682 (AS-15).

Progress Reports, Nuclear Engineering Department:

May 1 – August 31, 1961. BNL 696 (S-59).

September 1 – December 31, 1961. BNL 705 (S-60).

Conference Reports:

Brookhaven Symposia in Biology No. 14. *Fundamental Aspects of Radiosensitivity*. BNL 675 (C-31).

Proceedings of the Eleventh Annual AEC Corrosion Symposium. BNL 728 (C-33).

Brookhaven Lecture Series:

1. *Radioastronomy and Communication Through Space*, EDWARD M. PURCELL. BNL 658 (T-214).
2. *Current Ideas on the Endocrine Regulation of Cellular Processes*, IRVING SCHWARTZ. BNL 685 (T-232).
3. *Inside the Protein Molecule*, WERNER HIRS. BNL 649 (T-208).
5. *Neutron Physics Of and With the High Flux Beam Research Reactor*, HERBERT KOUTS. BNL 664 (T-218).
9. *The Physics of Semiconductor Radiation Detectors*, G.L. MILLER. BNL 699 (T-241).
10. *Theory of the Gene*, MILISLAV DEMEREC. BNL 739 (T-267).
12. *Excessive Salt Intake and Hypertension: A Dietary and Genetic Interplay*, LEWIS K. DAHL. BNL 733 (T-263).
13. *Galaxies*, OTTO STRUVE. BNL 710 (T-247).
14. *A Computer Learns To See*, PAUL HOUGH. BNL 725 (T-259).
18. *Accelerators of the Future*, JOHN P. BLEWETT. BNL 741 (T-269).

Weekly Bulletin 14, No. 51-2; 15, No. 1-51.

Weekly Selected Reading List 14, No. 16-52; 15, No. 1-15.

Miscellaneous:

Bibliography on Gamma-Ray Radiography, 1950 to 1961 Inclusive, With Subject and Author Indices, J.C. AUSTIN. BNL 716 (L-326).

Three Lectures on Elementary Particle Resonances, R.H. DALITZ. BNL 735 (T-264).

*Also included are those listings from the last annual report [BNL 682 (AS-15)] for which complete reference information was not then available.

STAFF PUBLICATIONS AND ABSTRACTS

Accelerator Department

BAKER, W.F. Computations for AGS experimental beams. Description of computer program. Informal Report BNL 6431.

BARTON, M.Q. AND SACHARIDIS, E.J. Time dependence of space charge beam losses in the Cosmotron. Informal Report BNL 6430.

BARTON, M.Q., SNYDER, H.S. AND NIELSEN, C.E. Longitudinal instability and cluster formation in ion beams. Informal Report BNL 5758.

BARTON, M.Q. – See also HOCHMAN, E.H.

BLEWETT, J.P. *Accelerators of the Future, Brookhaven Lecture Series No. 18*, BNL 741 (T-269), June 13, 1962.

BLEWETT, J.P. Developments in ultra-high-energy accelerators. Presented at AIEE Meeting, Detroit, Oct. 1961.

BLEWETT, M.H. Cosmotron. *Encyclopedia of Electronics*, C. Susskind, Editor, pp. 153-4, Reinhold, New York, 1962.

COTTINGHAM, J.G. Servo system for magnetic controlled constant intensity flat top beam spill-out. Informal Report BNL 6424.

COURANT, E.D. Acceleration of polarized protons to relativistic energies. Informal Report BNL 6432.

COURANT, E.D. Efficiency of multiple traversal targets. Informal Report BNL 6433.

CRITTENDEN, R.R., MUSGRAVE, B. AND MARTIN, H.J. Pion-proton interactions in the neighborhood of the Λ^0 - K^0 threshold. (A) *Bull. Am. Phys. Soc.* 7, 468 (1962).

DANBY, G.T. Magnetic measurements of AGS experimental magnets. Informal Report BNL 6435.

DRYDEN, R.A. – See GOULD, C.L.

FELDMAN, P. – See SWARTZ, C.E.

FICKINGER, W.J., SANDWEISS, J. AND SANFORD, J.R. The Cosmotron parallel plate velocity spectrometer (beam separator). Informal Report BNL 5757.

GOULD, C.L. AND DRYDEN, R.A. One year of operating experience with 70 getter-ion pumps. Presented at 8th Intern. Symp. Am. Vacuum Soc., Washington, Oct. 1961.

HAHN, H. The canonical equations for synchrotron oscillations. Informal Report BNL 5752.

- HAHN, H. Nonlinear bunch motion at transition. Informal Report BNL 6425.
- HAHN, H. AND HALAMA, H.J. Modifications of the wide-band sum electrode system. Informal Report BNL 5749.
- HAHN, H. AND ORGASS, R.J. A dynamic RF phase meter.
- HAHN, H. - See also PLOTKIN, M.
- HALAMA, H.J. - See HAHN, H.; PLOTKIN, M.
- HOCHMAN, E.H. AND BARTON, M.Q. Synchrotron oscillations. Informal Report BNL 6420.
- MARTIN, H.J. - See CRITTENDEN, R.R.
- MASCHKE, A.W. Target efficiency measurements at the AGS. Informal Report BNL 6426.
- MUSGRAVE, B. - See CRITTENDEN, R.R.
- NIELSEN, C.E. - See BARTON, M.Q.
- ORGASS, R.J. - See HAHN, H.
- PLOTKIN, M. High level Q-meter. *Electron. Design* (in press).
- PLOTKIN, M. Magnet coil tester. *Electron. Design* (in press).
- PLOTKIN, M., RAKA, E.C., HALAMA, H. AND HAHN, H. Beam capture and acceleration in the Brookhaven Alternating Gradient Synchrotron. Presented at IRE Western Conv., San Francisco, Aug. 1961.
- RAKA, E.C. Some remarks on beam controlled acceleration. Informal Report BNL 5759.
- RAKA, E.C. - See also PLOTKIN, M.
- ROBERTSON, D.S. Internal bunching in the Alternating Gradient Synchrotron. Informal Report BNL 6434.
- SACHARIDIS, E.J. - See BARTON, M.Q.
- SANDWEISS, J. - See FICKINGER, W.J.
- SANFORD, J.R. - See FICKINGER, W.J.
- SMITH, L.W. Possible use of a betatron core on the Cosmotron. Informal Report BNL 6436.
- SNYDER, H.S. - See BARTON, M.Q.
- SOUKAS, A.V. Hall probe regulator action and Hall generator evaluation. Informal Report BNL 5750.
- SWARTZ, C.E. AND FELDMAN, P. Beryllium foil monitor for external proton beam. *Rev. Sci. Instr.* **33**, 565-6 (1962).
- TRANS, A. Drift tube accelerator. Informal Report BNL 5788.
- TURNER, C.M. Cosmotron injection system engineering and operation notes. Pulsed liner, angle corrector and inflector pulsing systems. Informal Report BNL 5751.
- WALTERS, J.L. The ball tuner change tables. Informal Report BNL 6427.
- WALTERS, J.L. Design of reflectometers for coaxial lines used at AGS linac. Informal Report BNL 6428.
- WALTERS, J.L. Flattening procedures for empty cylindrical cavities. Informal Report BNL 5753.
- WALTERS, J.L. Some properties of stub tuners which can be displayed on Smith charts. Informal Report BNL 6429.

Applied Mathematics Division

- SHIMAMOTO, Y. Graph theory and parametric representations of Feynman amplitudes. *Nuovo Cimento Suppl.* (in press).

Architectural Planning Division

- RUDDY, J.M. Design and fabrication of metal work for two high intensity radiation experimental cells. *J. Struct. Div. Am. Soc. Civil Engrs.* **88**, 23-32 (1962).

- RUDDY, J.M. How to select gamma shielding doors. *Nucleonics* **20**, No. 6, 94-5 (1962).
- RUDDY, J.M. Shield doors grow in size, cost. *Eng. News-Record* **167**, 50-1 (Nov. 23, 1961).
- RUDDY, J.M. Versatile dc distribution gives wide range of voltages and currents. *Power* **106**, No. 4, 68-9 (1962).

Biology Department

- ALSTON, R.E. AND SPARROW, A.H. Somatic mutation rates in double and triple heterozygotes of *Impatiens balsamina* following chronic gamma irradiation. *Radiation Botany* **1**, 229-32 (1962).
- ALTSZULER, N. - See STEELE, R.
- AUGENSTINE, L.G. The effects of ionizing radiation on enzymes. *Advan. Enzymol.* **24** (in press).
- AUGENSTINE, L.G. Radiobiological mechanisms: Comparative distribution and role of ionization, excitation, and energy and charge migration. Chapter in *Progress in Biophysics and Biophysical Chemistry*, Pergamon Press, New York (in press).
- AUGENSTINE, L.G. AND GHIRON, C.A. Inactivation of trypsin by ultraviolet light. I. The correlation of inactivation with the disruption of constituent cystine. *Proc. Natl. Acad. Sci. U.S.* **47**, 1530-47 (1961).
- AUGENSTINE, L.G., GHIRON, C.A., GRIST, K.L. AND MASON, R. The inactivation of trypsin by ultraviolet light. II. The involvement of intramolecular hydrogen bond disruption. *Proc. Natl. Acad. Sci. U.S.* **47**, 1733-41 (1961).
- AUGENSTINE, L.G. AND GRIST, K.L. Aspects of radiation-inactivation of enzymes. (A) Presented at IAEA Symp. Biological Effects of Ionizing Radiation at the Molecular Level, Br'no, Czechoslovakia, July 1962.
- AUGENSTINE, L.G. - See also RILEY, P.; WEINBERG, C.J.
- BARNES, H.W. - See ORDY, J.M.
- BERECH, J. JR. AND CURTIS, H.J. Renal function in the aged and x-irradiated mouse. (A) Presented at Radiation Res. Soc. Meeting, Colorado Springs, May 1962.
- BERGERON, J.A. AND SMILLIE, R.M. Relations between photosynthesis and the structure and enzymes of *Euglena* chloroplasts. (A) Presented at Am. Soc. Plant Physiologists Meeting, AIBS, Corvallis, Oreg., Aug. 1962.
- BERGERON, J.A. - See also GETTNER, M.E.
- BISHOP, J.S. - See STEELE, R.
- BORNER, G. - See INNES, J.R.M.
- BOURDEAU, P.F. - See WOODWELL, G.M.
- BREWBAKER, J.L. Cyanidin-red white clover, duplicate recessive mutant in *Trifolium repens* L. Submitted to *J. Heredity*.
- BREWBAKER, J.L. What can you expect from atomic-irradiated seed? Submitted to *Flower and Garden*.
- BREWBAKER, J.L. AND EMERY, G.C. Pollen radiobotany. *Radiation Botany* **1**, 101-54 (1962).
- BROWN, J.A.M. Effect of thymidine analogues on reproductive morphogenesis in *Arabidopsis thaliana* (L) Heynh. Submitted to *Nature*.
- BROWN, J.A.M. Effects of thymidine analogues on the apical meristem of *Arabidopsis thaliana* (L) Heynh. (A) Presented at Botan. Soc. Am. Meeting, AIBS, Corvallis, Oreg., Aug. 1962.
- CARLOUGH, M. - See QUASTLER, H.
- CARTER, J.G. - See WEINBERG, C.J.

- CONTI, S.F. AND GETTNER, M.E. Electron microscopy of cellular division in *Escherichia coli*. *J. Bacteriol.* **83**, 544-50 (1962).
- COX, E. - See LEVINE, M.
- CROWLEY, C. - See CURTIS, H.J.
- CURTIS, H.J. The biological effects of heavy cosmic ray particles. *Proc. 3rd Intern. Space Sci. Symp., Washington, April 30-May 5, 1962* (in press).
- CURTIS, H.J. The late effects of radiation. *Proc. Am. Phil. Soc.* (in press).
- CURTIS, H.J. The present status of the radiation hazard: Types of radiation and mechanisms of action on living organisms. (A) Presented at Ann. Meeting Am. Phil. Soc., Philadelphia, April 1962.
- CURTIS, H.J. AND CROWLEY, C. Chromosome aberrations in liver cells in relation to aging. Presented at IAEA/UNESCO Symp. Aetiology of Late Somatic Effects of Ionizing Radiations, London, March 1962.
- CURTIS, H.J. AND CROWLEY, C. Life span shortening from various tissue insults. Presented at IAEA/UNESCO Symp. Aetiology of Late Somatic Effects of Ionizing Radiations, London, March 1962.
- CURTIS, H.J. - See also BAKER, C.P. (Physics); BERECH, J. JR.; ORDY, J.M.; SMITH, H.H.; STEVENSON, K.G.; VON SALLMANN, L.; ZEMAN, W.
- CURTISS, R. - See LEVINE, M.
- DALMAS, R.L. - See SCHWINGHAMER, E.A.
- DEBODO, R.C. - See STEELE, R.
- DEMEREK, M. *Theory of the Gene, Brookhaven Lecture Series No. 10*, BNL 739 (T-267), Oct. 18, 1961.
- DORNFEST, B.S., LOBUE, J., HANDLER, E.S., GORDON, A.S. AND QUASTLER, H. Mechanisms of leukocyte production and release. I. Factors influencing leukocyte release from isolated perfused rat femora. *Acta Haematol.* **28**, 42-60 (1962).
- DORNFEST, B.S., LOBUE, J., HANDLER, E.S., GORDON, A.S. AND QUASTLER, H. Mechanisms of leukocyte production and release. II. Factors influencing leukocyte release from isolated perfused rat legs. *J. Lab. Clin. Med.* (in press).
- DORNFEST, B.S. - See also GORDON, A.S.
- DUNN, A. - See STEELE, R.
- EDMUNDSON, A.B. AND HIRS, C.H.W. On the structure of sperm whale myoglobin. I. The amino acid composition and terminal groups of the chromatographically purified protein. II. The tryptic hydrolysis of the denatured protein. III. The tryptic peptide derived from the amino-terminal portion of the molecule. IV. Amino acid sequences of the smaller tryptic peptides containing aromatic residues. Submitted to *J. Mol. Biol.*
- EMERY, G.C. - See BREWBAKER, J.L.
- ENGLBERGER, F.M. - See KOSHLAND, M.E.
- EVANS, H.J. Chromosome aberrations induced by ionizing radiations. *International Review of Cytology*, G.H. Bourne and J.F. Danielli, Editors, Chapter 13, pp. 221-321, Academic Press, New York, 1962.
- EVANS, H.J. AND SPARROW, A.H. Nuclear factors affecting radiosensitivity. II. Dependence on nuclear and chromosome structure and organization. *Brookhaven Symposia in Biology No 14*, pp. 101-27, BNL 675 (C-31), Nov. 1961.
- EVANS, H.J. - See also SPARROW, A.H.
- EVANS, W.R. AND SMILLIE, R.M. The role of nucleic acids in the formation of chloroplasts. (A) Presented at Am. Soc. Plant Physiologists Meeting, AIBS, Corvallis, Oreg., Aug. 1962.
- FRENKEL, J.K. - See INNES, J.R.M.
- FULHABER, E. - See PERROTTA, C.A.
- GADDONE, S.M. - See KOSHLAND, M.E.
- GAETJENS, E. - See KOSHLAND, D.E. JR.
- GAY, H. - See WOODS, P.S.
- GETTNER, M.E., SCHWINGHAMER, E.A. AND BERGERON, J. Electron microscopy of phage from *Rhizobium trifolii*. (N) *Biochim. Biophys. Acta* **55**, 209 (1962).
- GETTNER, M.E. - See also CONTI, S.F.
- GHIRON, C.A. - See AUGENSTINE, L.G.
- GOOLNICK, B.G. Techniques of autoradiography using tritiated thymidine and results in thymus and sarcoma 180. (A) *Radiation Res.* **16**, 560 (1962).
- GOOLNICK, B.G. - See also QUASTLER, H.
- GORDON, A.S., DORNFEST, B.S., LOBUE, J., HANDLER, E.S. AND QUASTLER, H. Mechanisms regulating leucocyte release from perfused femurs. (A) Presented at Am. Soc. Hematol. Meeting, Los Angeles, Nov. 1961.
- GORDON, A.S. - See also DORNFEST, B.S.
- GRIMES, P. - See VON SALLMANN, L.
- GRIST, K.L. - See AUGENSTINE, L.G.; RILEY, P.
- GUNCKEL, J.E. - See HAGEN, G.L.; MEISELMAN, N.
- HAGEN, G.L., GUNCKEL, J.E. AND SPARROW, A.H. Morphology and histology of tumor types induced by x, gamma, and beta irradiation of a tobacco hybrid. *Am. J. Botany* **48**, 691-9 (1961).
- HAMMOND, E.C. - See WOODWELL, G.M.
- HAMPTON, J.C. - See QUASTLER, H.
- HANDLER, E.S. - See DORNFEST, B.S.; GORDON, A.S.
- HEPPEL, L.A. - See KOSHLAND, D.E. JR.
- HILMOE, R. J. - See KOSHLAND, D.E. JR.
- HIRS, C.H.W. *Inside the Protein Molecule, Brookhaven Lecture Series No. 3*, BNL 649 (T-208), Jan. 11, 1961.
- HIRS, C.H.W. - See also EDMUNDSON, A.B.
- INNES, J.R.M. Comparative neuropathology: Diseases of the spinal cord of animals. *Livre Jubilaire Docteur Ludo van Bogaert, Institut Bunge*, pp. 390-6, Acta Medica Belgica, Brussels, 1962.
- INNES, J.R.M. AND CARSTEN, A. Delayed effects of localized x-irradiation of the nervous system of experimental rats and monkeys. *Brookhaven Symposia in Biology No. 14*, pp. 200-3, BNL 675 (C-31), Nov. 1961.
- INNES, J.R.M. AND SAUNDERS, L.Z. *Comparative Neuropathology*, 839 pp., 363 plates, 4000 ref., Academic Press, New York, 1962.
- INNES, J.R.M., ZEMAN, W., FRENKEL, J.K. AND BORNER, G. Occult endemic encephalitozoonosis of the central nervous system of mice (Swiss-Bagg-O'Grady strain). *J. Neuropathol. Exptl. Neurol.* (in press, Oct. 1962).
- JAKOWSKA, S., NIGRELLI, R.F. AND SPARROW, A.H. Additional radiobiological studies on normal and anemic newts, *Diemictylus viridescens*. (A) Presented at AAAS Meeting, Denver, Dec. 1961.
- KELLY, H. - See SMILLIE, R.M.
- KEMBER, N.F. Kinetics of population of bone forming cells in the normal and irradiated rat. *Proc. Utah Symp. Aspects of Internal Irradiation, Salt Lake City, May 1961*, Pergamon Press, New York (in press).

- KEMBER, N.F., QUASTLER, H. AND WIMBER, D.R. Adaptation to continuous irradiation. (N) *Brit. J. Radiol.* 35, 290 (1962).
- KLEIN, J.R. Depression of heme formation and the production of free porphyrin in duck erythrocytes. Submitted to *Am. J. Physiol.*
- KLEIN, J.R. Protoporphyrin and heme formation by erythrocytes from the x-irradiated and bled duck. *Am. J. Physiol.* 201, 663-7 (1961).
- KLEINFELD, R. - See ZEMAN, W.
- KOSHLAND, D.E. JR. The active site of chymotrypsin and methods for the study of active sites. (A) Presented at Biophys. Soc. Meeting, London, Dec. 1961.
- KOSHLAND, D.E. JR. Biological specificity in protein-small molecule interactions. *Proc. 1st Intern. Pharmacol. Meeting, Stockholm, 1961*, Pergamon Press, Oxford (in press).
- KOSHLAND, D.E. JR. Catalysis in life and in the test tube. In *Horizons in Biochemistry*, Academic Press, New York (in press).
- KOSHLAND, D.E. JR. The comparison of nonenzymic and enzymic reaction velocities. *J. Theoret. Biol.* 2, 75-86 (1962).
- KOSHLAND, D.E. JR. The nature of the catalytic site of enzymes. (A) Presented at N.Y. Acad. Sci. Symp. Antibody to Enzymes - A Three-Component System, New York, Sept. 1962.
- KOSHLAND, D.E. JR., GAETJENS, E. AND THOMA, J.A. Comparison of acid and enzyme catalyzed hydrolyses of acetals. (A) Presented at Am. Chem. Soc. Meeting, Atlantic City, Sept. 1962.
- KOSHLAND, D.E. JR., HILMOE, R.J., HEPPEL, L.A. AND SPRINGHORN, S.S. Cleavage point during hydrolyses catalyzed by ribonuclease and phosphodiesterase. *Biochim. Biophys. Acta* 53, 214-15 (1961).
- KOSHLAND, D.E. JR., YANKEELOV, J.A. AND THOMA, J.A. Specificity and catalytic power in enzyme action. Submitted to *Federation Proc.*
- KOSHLAND, D.E. JR. - See also LEVY, H.M.; RAY, W.J. JR.; STRUMEYER, D.H.
- KOSHLAND, M.E., ENGLBERGER, F.M. AND GADDONE, S.M. The effect of iodination on the binding of haptene to arsonic antibody. *J. Immunol.* (in press).
- KUO, J. - See NIMS, L.F.
- LACHANCE, L.E. The effect of temperature on the cytogenetic damage induced by fast neutrons in *Tradescantia* microspores. *Radiation Botany* 1, 203-8 (1962).
- LACKS, S. Molecular fate of DNA in genetic transformation of pneumococcus. *J. Mol. Biol.* 5, 119-31 (1962).
- LAMERTON, L.F., LORD, B.I. AND QUASTLER, H. Studies of cell population kinetics in normal and continuously irradiated animals. Submitted to *Strahlentherapie*.
- LAMERTON, L.F. - See also WIMBER, D.R.
- LEVINE, M. AND COX, E. The ultraviolet sensitivity of frozen phage. Submitted to *Radiation Res.*
- LEVINE, M. AND CURTIS, R. Genetic fine structure of the C region and the linkage map of phage P22. *Genetics* 46, 1573-80 (1961).
- LEVINE, R.P. AND SMILLIE, R.M. The pathway of triphosphopyridine nucleotide photoreduction in *Chlamydomonas reinhardtii*. *Proc. Natl. Acad. Sci. U.S.A.* 48, 417-21 (1962).
- LEVY, C.K. AND QUASTLER, H. Acute responses of the vestibular apparatus to x-irradiation. *Radiation Res.* 16, 189-200 (1962).
- LEVY, H.M., RYAN, E.M., SPRINGHORN, S.S. AND KOSHLAND, D.E. JR. Further evidence for oxygen exchange at an intermediate state in myosin hydrolysis. *J. Biol. Chem.* 237, 1730-1 (1962).
- LEVY, H.M., SHARON, N., RYAN, E.M. AND KOSHLAND, D.E. JR. Effect of temperature on the rate of hydrolysis of ATP and ITP by myosin with and without modifiers. *Biochim. Biophys. Acta* 56, 118-26 (1962).
- LIPKIN, M. AND QUASTLER, H. Cell population kinetics in the colon of the mouse. *J. Clin. Invest.* 41, 141-6 (1962).
- LIPKIN, M. AND QUASTLER, H. Cell retention and incidence of carcinoma in several portions of the gastrointestinal tract. (L) *Nature* 194, 1198-9 (1962).
- LIPKIN, M. AND QUASTLER, H. Studies of protein metabolism in intestinal epithelial cells. *J. Clin. Invest.* 41, 646-53 (1962).
- LIPKIN, M., QUASTLER, H. AND MUGGIA, F. Protein synthesis in the irradiated intestine of mice. (A) *Federation Proc.* 21, 256 (1962).
- LOBUE, J. - See DORNFEST, B.S.; GORDON, A.S.
- LORD, B.I. - See LAMERTON, L.F.
- MARIC, M.M. - See MATARAJAN, A.T.
- MASON, R. - See AUGENSTINE, L.G.
- MCCARTER, J.A. AND QUASTLER, H. Effect of dimethylbenzanthracene on cellular proliferation cycle. *Nature* 194, 873-4 (1962).
- MCCARTER, J.A. AND QUASTLER, H. Note on the effect of a carcinogenic hydrocarbon on synthesis of DNA. (N) *Biochem. Biophys. Res. Commun.* 55, 552-3 (1962).
- MEISELMAN, N., GUNCKEL, J.E. AND SPARROW, A.H. The general morphology and growth responses of two species of *Nicotiana* and their interspecific hybrid after chronic gamma irradiation. *Radiation Botany* 1, 69-79 (1961).
- MERICLE, L.W. AND MERICLE, R.P. Mutation induction by proembryo irradiation. *Radiation Botany* 1, 195-202 (1962).
- MERICLE, L.W. AND MERICLE, R.P. Radiosensitivity of the developing plant embryo. *Brookhaven Symposia in Biology No. 14*, pp. 262-86, BNL 675 (C-31), Nov. 1961.
- MERICLE, R.P. - See MERICLE, L.W.
- MIKSCH, J.P. Changes in radiosensitivity of *Glycine max* during early stages of growth. (A) Presented at Botan. Soc. Am. Meeting, AIBS, Corvallis, Oreg., Aug. 1962.
- MIKSCH, J.P. The influence of DNA and nuclear volume on sensitivity changes of *Glycine max* seedlings. (A) Presented at 2nd Intern. Congr. Radiation Res., Harrogate, England, Aug. 1962.
- MIKSCH, J.P., SPARROW, A.H. AND ROGERS, A.F. The effects of chronic gamma irradiation on the apical meristem and bud formation of *Taxus media*. Submitted to *Radiation Botany*.
- MILLER, L.N. - See WOODWELL, G.M.
- MUGGIA, F. - See LIPKIN, M.
- NATARAJAN, A.T. AND MARIC, M. The time-intensity factor in dry seed irradiation. *Radiation Botany* 1, 1-9 (1961).
- NELSON, D.R. - See WEINBERG, C.J.
- NIGRELLI, R.F. - See JAKOWSKA, S.

- NIMS, L.F. Origin of electric currents in living systems. Submitted to *Am. J. Physiol.*
- NIMS, L.F. Physiological unit of electrical energy. (A) Presented at Ann. Meeting Biophys. Soc., Washington, Feb. 1962.
- NIMS, L.F. Steady-state material transfer through biological barriers. *Am. J. Physiol.* **201**, 987-94 (1961).
- NIMS, L.F. Tracers, transfer through membranes, and coefficients of transfer. *Science* **137**, 130-2 (1962).
- NIMS, L.F. AND KUO, J. Electromotive force of transfer systems containing membranes. Submitted to *Am. J. Physiol.*
- NIMS, L.F. AND THURBER, R.E. Ion distribution patterns in stationary state systems. *Am. J. Physiol.* **201**, 995-8 (1961).
- NIMS, L.F. AND THURBER, R.E. Whole-body x-irradiation, nitrogen excretion, and the adrenal gland. *Endocrinology* **70**, 589-94 (1962).
- NIMS, L.F. - See also THURBER, R.E.
- OLSON, J.M. Light-induced cytochrome oxidation in green sulfur bacteria. (A) Presented at Am. Soc. Biol. Chem. Meeting, Atlantic City, April 1962.
- OLSON, J.M. AND ROMANO, C.A. A new chlorophyll from green bacteria. (N) *Biochim. Biophys. Acta* **59**, 726-8 (1962).
- ORDY, J.M., BARNES, H.W., SAMORAJSKI, T., CURTIS, H.J., WOLIN, L. AND ZEMAN, W. Pathologic and behavioral changes in mice following CNS deuteron irradiation. Submitted to *Radiation Res.*
- PERROTTA, C.A. Effects of irradiation on uterine and vaginal epithelium of the mouse. (A) Presented at Am. Assoc. Anatomists Meeting, Minneapolis, March 1962.
- PERROTTA, C.A. Initiation of cell proliferation in the vaginal and uterine epithelia of the mouse. *Am. J. Anat.* **111**, 195-204 (1962).
- PERROTTA, C.A., WOLFSBERG, M.F. AND FULHABER, E. Proliferation and DNA synthesis in the uterine and vaginal epithelium and in the regenerating liver of the mouse. (A) *Radiation Res.* **16**, 561 (1962).
- PRENSKY, W. Uptake of thymine-methyl- H^3 by pachytene chromosomes of *Lilium longiflorum*. (A) Submitted to *Records Genet. Soc. Am.*
- QUASTLER, H. Autoradiographic studies of DNA synthesis in irradiated mice. (A) Presented at 2nd Intern. Congr. Radiation Res., Harrogate, England, Aug. 1962.
- QUASTLER, H. Effects of irradiation on synthesis and loss of DNA. In *Chemical and Biological Action of Radiations*, Vol. 6, M. Haissinsky, Editor, Academic Press, New York, and Masson et Cie., Paris (in press, 1962).
- QUASTLER, H. X-irradiation and DNA synthesis: Competitive or noncompetitive inhibition? (A) Presented at Ann. Meeting Biophys. Soc., Washington, Feb. 1962.
- QUASTLER, H., CARLOUGH, M., GOOLNICK, B.G. AND WOLFSBERG, M.F. Effect of x-irradiation on DNA synthesis in some squamous cell epithelia. (A) *Radiation Res.* **16**, 561-2 (1962).
- QUASTLER, H. AND HAMPTON, J.C. Effects of ionizing radiation on the fine structure and function of the intestinal epithelium of the mouse. I. Villus epithelium. Submitted to *Radiation Res.*
- QUASTLER, H. AND ZUBAY, G. An RNA-protein code based on replacement data. II. Adjustment and extension. *J. Theoret. Biol.* (in press).
- QUASTLER, H. - See also DORNFEST, B.S.; GORDON, A.S.; KEMBER, N.F.; LAMERTON, L.F.; LEVY, C.K.; LIPKIN, M.; MCCARTER, J.A.; SHERMAN, F.G.; STEIN, O.L.; WIMBER, D.E.; WULFF, V.J.; ZUBAY, G.
- RAY, W.J. JR. AND KOSHLAND, D.E. JR. Identification of amino acids involved in phosphoglucomutase action. *J. Biol. Chem.* **237**, 2493-505 (1962).
- RAY, W.J. JR. AND KOSHLAND, D.E. JR. Role of amino acid residues on chymotrypsin action as indicated by an "all-or-none" assay. (A) Presented at Am. Chem. Soc. Meeting, Washington, March 1962.
- RIGOPOULOS, N. - See SMILLIE, R.M.
- RILEY, P., GRIST, K. AND AUGENSTINE, L. The dependence of enzyme activity on the integrity of specific disulfide bonds. (A) Presented at Ann. Meeting Biophys. Soc., Washington, Feb. 1962.
- ROGERS, A.F. - See MIKSCH, J.P.
- ROMANO, C.A. - See OLSON, J.M.
- RYAN, E.M. - See LEVY, H.M.
- SAMORAJSKI, T. - See ORDY, J.M.; ZEMAN, W.
- SCARPELLI, D.G. - See ZEMAN, W.
- SCHAEFFER, G.W. Tumor induction by an IAA-kinetin interaction in a *Nicotiana* hybrid. Submitted to *Science*.
- SCHAIER, L.A. - See SPARROW, A.H.
- SCHWINGHAMER, E.A. Studies on induced variation in the rhizobia. III. Host range modification of *Rhizobium trifolii* by spontaneous and radiation-induced mutation. *Am. J. Botany* **49**, 269-77 (1962).
- SCHWINGHAMER, E.A. AND DALMAS, R.L. Studies on induced variation in the rhizobia. II. Radiation sensitivity and induction of antibiotic-resistance markers. *Appl. Microbiol.* **9**, 410-14 (1961).
- SCHWINGHAMER, E.A. - See also GETTNER, M.E.
- SCOTT, K.J. AND SMILLIE, R.M. Enzyme changes in barley leaves after infection with powdery mildew. (A) Presented at Am. Soc. Plant Physiologists Meeting, AIBS, Corvallis, Oreg., Aug. 1962.
- SENGUN, A. - See WOODS, P.S.
- SHARON, N. - See LEVY, H.M.
- SHAVER, D.L. The effect of structural heterozygosity on the degree of preferential segregation in allotetraploids of *Zea*. (A) Presented at Genet. Soc. Am. Meeting, AIBS, Corvallis, Oreg., Aug. 1962.
- SHAVER, D.L. Five short articles. Submitted to *Maize Genetics Cooperative News Letter*.
- SHAVER, D.L. A study of aneuploidy in autotetraploid maize. Submitted to *Can. J. Genet. Cytol.*
- SHAVER, D.L. AND SPARROW, A.H. The relationship between nuclear or chromosome volume and rate of radiation-induced somatic mutation in higher plants. (A) Presented at Genet. Soc. Am. Meeting, AIBS, Corvallis, Oreg., Aug. 1962.
- SHERMAN, F.G., QUASTLER, H. AND WIMBER, D.R. Cell population kinetics in the ear epidermis of mice. *Exptl. Cell Res.* **25**, 114-19 (1961).
- SHERMAN, F.G. - See also WULFF, V.J.
- SMILLIE, R.M. Algal photosynthetic pyridine nucleotide reductases. (A) Presented at Am. Soc. Plant Physiologists Meeting, AIBS, Corvallis, Oreg., Aug. 1962.
- SMILLIE, R.M. Distribution and properties of plant fructose 1,6-diphosphatases and their role in photosynthesis and respiration. Submitted to *J. Biol. Chem.*

- SMILLIE, R.M. Formation and function of soluble proteins in chloroplasts. (A) Submitted to *Can. J. Botany*.
- SMILLIE, R.M. Photosynthetic and respiratory activities of growing pea leaves. Submitted to *Plant Physiol.*
- SMILLIE, R.M. AND RIGOPOULOS, N. Carotenoid production by streptomycin-bleached *Euglena*. *J. Protozool.* **9**, 149-51 (1962).
- SMILLIE, R.M., RIGOPOULOS, N. AND KELLY, H. Enzymes of the reductive pentose phosphate cycle in the purple and the green photosynthetic sulfur bacteria. *Biochim. Biophys. Acta.* **56**, 612-14 (1962).
- SMILLIE, R.M. - See also BERGERON, J.A.; EVANS, W.R.; LEVINE, R.P.; SCOTT, K.J.
- SMITH, H.H. Genetic control of *Nicotiana* plant tumors. Submitted to *Trans. N.Y. Acad. Sci.*
- SMITH, H.H. The reactor as a tool for research in plant sciences and agriculture. *Proc. IAEA Symp. Programming and Utilization of Res. Reactors, Vienna, Oct. 1961* (in press).
- SMITH, H.H. Studies on the origin, inheritance and mutation of genic-cytoplasmic male sterility in *Nicotiana*. (A) Submitted to *Records Genet. Soc. Am.*
- SMITH, H.H., CURTIS, H.J., WOODLEY, R.G. AND STEIN, O.L. The deuteron microbeam as a tool in botanical research. *Radiation Botany*, **1**, 255-68 (1962).
- SPARROW, A.H. Predictions of radiosensitivities of plant species. (A) Presented at 2nd Intern Congr. Radiation Res., Harrogate, England, Aug. 1962.
- SPARROW, A.H. AND EVANS, H.J. Nuclear factors affecting radiosensitivity. I. The influence of nuclear size and structure, chromosome complement, and DNA content. *Brookhaven Symposia in Biology No. 14*, pp. 76-100, BNL 675 (C-31), Nov. 1961.
- SPARROW, A.H. AND SCHAIRER, L.A. Plant radiosensitivities and potential damage from fallout radiation. (A) Presented at Ann. Meeting Radiation Res. Soc., Colorado Springs, May 1962.
- SPARROW, A.H. AND WOODWELL, G.M. Prediction of the sensitivity of plants to chronic gamma irradiation. *Radiation Botany* (in press).
- SPARROW, A.H. - See also ALSTON, R.E.; EVANS, H.J.; HAGEN, G.L.; JAKOWSKA, S.; MEISELMAN, N.; MIKSCH, J.P.; SHAVER, D.L.
- SPRINGHORN, S.S. - See KOSHLAND, D.E. JR.; LEVY, H.M.
- STEELE, R., ALTSZULER, N., DUNN, A., BISHOP, J.S. AND DEBODO, R.C. Effect of 2-deoxy-D-glucose on glucose production and uptake in intact dog. (A) *Federation Proc.* **21**, 81 (1962).
- STEELE, R., BISHOP, J.S., DUNN, A., ALTSZULER, N. AND DEBODO, R.C. Hepatic incorporation of the C¹⁴ of a C¹⁴-glucose load. (A) *Federation Proc.* **21**, 201 (1962).
- STEELE, R., DEBODO, R.C., ALTSZULER, N., DUNN, A. AND BISHOP, J.S. Effects of insulin on hepatic glucose metabolism and glucose utilization by tissues. Submitted to *J. Clin. Endocrinol. Metab.*
- STEELE, R., DUNN, A., ALTSZULER, N., BISHOP, J.S. AND DEBODO, R.C. Effects of insulin, a glucose load, and diet on glucose production and utilization. (A) *Federation Proc.* **21**, 205 (1962).
- STEELE, R., WINKLER, B., ALTSZULER, N., DUNN, A. AND DEBODO, R.C. Effects of growth hormone on FFA metabolism. (A) *Federation Proc.* **21**, 198 (1962).
- STEFFENSEN, D.M. Four short articles. Submitted to *Maize Genetics Cooperative News Letter*.
- STEFFENSEN, D.M. A variable distance for chromosome exchange dependent on cellular activity, temperature and other conditions. (A) Presented at Ann. Meeting Radiation Res. Soc., Colorado Springs, May 1962.
- STEIN, O.L. AND QUASTLER, H. The effect of tritiated thymidine on lateral root formation. Presented at Am. Soc. Cell. Biol. Meeting, Chicago, Nov. 1961.
- STEIN, O.L. AND QUASTLER, H. The effect of tritiated thymidine on the morphogenesis of lateral roots. *Tritium in the Physical and Biological Sciences*, Vol. 2, pp. 149-53, IAEA, Vienna, 1962.
- STEIN, O.L. - See also SMITH, H.H.
- STEVENSON, K.G. AND CURTIS, H.J. Chromosomal aberrations in irradiated and nitrogen-mustard treated mice. *Radiation Res.* **15**, 774-84 (1961).
- STRUMEYER, D.H. AND KOSHLAND, D.E. JR. Roles of the methionine residues in chymotrypsin action. (A) Presented at Am. Chem. Soc. Meeting, Atlantic City, Sept. 1962.
- THOMA, J.A. - See KOSHLAND, D.E. JR.
- THURBER, R.E. AND NIMS, L.F. Liver glycogen and blood sugar levels in x-irradiated newborn rats. *Endocrinology* **70**, 595-9 (1962).
- THURBER, R.E. - See also NIMS, L.F.
- VON SALLMANN, L., CURTIS, H.J., AND GRIMES, P. The effect of a deuteron microbeam on the mouse crystalline lens. *Arch. Ophthalmol.* **67**, 163-70 (1962).
- WEINBERG, C.J., NELSON, D.R., CARTER, J.G. AND AUGENSTINE, L.G. Thermoluminescence from gamma-irradiated biochemicals. Investigation of emission spectra. *J. Chem. Phys.* **36**, 2869-79 (1962).
- WIMBER, D.E. DNA synthesis and cell proliferation in chronically irradiated *Tradescantia* roots and mouse gut studied with a C¹⁴- and H³-thymidine double labeling technique. (A) Presented at 2nd Intern. Congr. Radiation Res., Harrogate, England, Aug. 1962.
- WIMBER, D.E. The effect of temperature on the duration of the nuclear cycle and its various subdivisions in *Tradescantia* root tips. (A) Presented at Botan. Soc. Am. Meeting, AIBS, Corvallis, Oreg., Aug. 1962.
- WIMBER, D.E. AND QUASTLER, H. A C¹⁴- and H³-thymidine double labeling technique in the study of cell proliferation in *Tradescantia* root tips. Submitted to *Exptl. Cell. Res.*
- WIMBER, D.R. Autoradiographic studies of DNA synthesis in the gastrointestinal epithelium and the bladder of the mouse. (A) *Radiation Res.* **16**, 561 (1962).
- WIMBER, D.R. AND LAMERTON, L.F. Cell population studies on the intestine of continuously irradiated rats. Submitted to *Radiation Res.*
- WIMBER, D.R. - See also KEMMER, N.F.; SHERMAN, F.G.
- WINKLER, B. - See STEELE, R.
- WOLFSBERG, M.F. Cell renewal in the squamous epithelium of the mouse forestomach. (A) Submitted to *Records Genet. Soc. Am.*
- WOLFSBERG, M.F. - See also PERROTTA, C.A.; QUASTLER, H.
- WOLIN, L. - See ORDY, J.M.
- WOODLEY, R.G. - See BAKER, C.P. (Physics); SMITH, H.H.
- WOODS, P.S. Autoradiographic studies of ribonucleic acid metabolism with tritium-labeled cytidine. *Tritium*

- in the *Physical and Biological Sciences*, Vol. 2, pp. 335-46, IAEA, Vienna, 1962.
- WOODS, P.S., GAY, H. AND SENGUN, A. Organization of the salivary-gland chromosome as revealed by the pattern of incorporation of H^3 -thymidine. *Proc. Natl. Acad. Sci. U.S.* **47**, 1486-93 (1961).
- WOODWELL, G.M. Effect of chronic gamma radiation on vegetation. (A) Presented at Ecol. Soc. Am. Meeting, AIBS, Corvallis, Oreg., Aug. 1962.
- WOODWELL, G.M. AND BOURDEAU, P.F. Measurement of dry matter production of the plant cover. *Proc. UNESCO Symp. Methodology of Plant Eco-Physiology, Institut de Botanique, France, April 1962* (in press).
- WOODWELL, G.M. AND HAMMOND, E.C. *A Descriptive Technique for Study of the Effects of Chronic Ionizing Radiation on a Forest Ecological System*. BNL 715 (T-251), March 1962.
- WOODWELL, G.M. AND MILLER, L.N. Effect of chronic gamma irradiation on the distribution of cambial activity in *Pinus rigida* Mill. (A) Presented at Ecol. Soc. Am. Meeting, AIBS, Corvallis, Oreg., Aug. 1962.
- WULFF, V.J., QUASTLER, H. AND SHERMAN, F.G. The incorporation of H^3 -cytidine in mice of different ages. *Arch. Biochem. Biophys.* **95**, 548-9 (1961).
- YANKEELOV, J.A. - See KOSHLAND, D.E. JR.
- ZEMAN, W. AND CURTIS, H.J. Metabolic and histochemical studies on direct radiation-induced nerve cell necrosis. (A) *Proc. 4th Intern. Congr. Neuropathol., Munich, Sept. 1961*, Vol. 1, pp. 141-7, Georg Thieme Verlag, Stuttgart, 1962.
- ZEMAN, W., CURTIS, H.J. AND BAKER, C.P. Histopathologic effect of high energy particle microbeams on the visual cortex of the mouse brain. *Radiation Res.* **15**, 496-514 (1961).
- ZEMAN, W., CURTIS, H.J., SCARPELLI, D.G. AND KLEINFELD, R. Chemical and enzymatic changes in nerve cells irradiated with high energy deuteron microbeams. *Response of the Nervous System to Ionizing Radiation*, T. J. Haley and R.S. Snider, Editors, pp. 429-39, Academic Press, New York, 1962.
- ZEMAN, W., SAMORAJSKI, T., AND CURTIS, H. J. Histochemical studies on mouse brains irradiated with high energy deuteron microbeams. *Effects of Ionizing Radiation on the Nervous System*, pp. 297-308, IAEA, Vienna, 1962.
- ZEMAN, W. - See also BAKER, C.P. (Physics); INNES, J.R.M.; ORDY, J.M.
- ZUBAY, G. The specificity of incorporation of glutamic acid and glutamine into protein. *Proc. Natl. Acad. Sci. U.S.* **48**, 894-6 (1962).
- ZUBAY, G. A theory on the mechanism of messenger-RNA synthesis. *Proc. Natl. Acad. Sci. U.S.* **48**, 456-61 (1962).
- ZUBAY, G. AND QUASTLER, H. An RNA-protein code based on replacement data. *Proc. Natl. Acad. Sci. U.S.* **48**, 461-71 (1962).
- ZUBAY, G. - See also QUASTLER, H.
- ALLEN, A.O. - See also CZAPSKI, G.; HAYON, E.; HUMMEL, A.
- BAKER, E.W. AND KATCOFF, S. Fission of Ag and Br in nuclear emulsion by 1.0-3.0-Bev protons. *Phys. Rev.* **126**, 729-34 (1962).
- BASSETT, W.A., KERR, P.F., SCHAEFFER, O.A. AND STOENNER, R.W. Potassium-argon ages of volcanics near Grants, New Mexico. *Bull. Geol. Soc. Am.* (in press).
- BASSETT, W.A., KERR, P.F., SCHAEFFER, O.A. AND STOENNER, R.W. Potassium-argon dating of the late tertiary volcanics and mineralization of Marysvale, Utah. *Bull. Geol. Soc. Am.* (in press).
- BELL, R.P. AND CROOKS, J.E. The secondary hydrogen isotope effect in the dissociation constant of formic acid. *Trans. Faraday Soc.* **58**, 1409-11 (1962).
- BELL, R.S. AND SUTIN, N. The kinetics of the formation and dissociation of mono-(1,10-phenanthroline)-iron (II). *Inorg. Chem.* **1**, 359-63 (1962).
- BELLON, P.L. - See SCATTURIN, V.
- BIELSKI, B.H.J. AND SAITO, E. The activation energy for the disproportionation of the HO_2 radical in 0.8 *N* sulfuric acid solution. (N) Submitted to *J. Phys. Chem.*
- BIELSKI, B.H.J. - See also BLACK, G.W.; CZAPSKI, G.; SAITO, E.
- BIGEISEN, J. Correlation of tritium and deuterium isotope effects. *Tritium in the Physical and Biological Sciences*, Vol. 1, pp. 161-8, IAEA, Vienna, 1962.
- BIGEISEN, J. Vapor pressures of isotopic molecules. (A) Presented at Ann. Meeting Soc. Chim. Phys., Paris, June 1962.
- BIGEISEN, J., HASCHEMEYER, R.H., WOLFSBERG, M. AND YANKWICH, P.E. Temperature dependence of the carbon isotope effect in the dehydration of formic acid by concentrated sulfuric acid. *J. Am. Chem. Soc.* **84**, 1813-15 (1962).
- BIGEISEN, J. AND RIBNIKAR, S.V. Structural effects in the vapor pressures of isotopic molecules. O^{18} and N^{15} substitution in N_2O . *J. Chem. Phys.* **35**, 1297-1305 (1961).
- BIGEISEN, J., RIBNIKAR, S.V., STERN, M.J. AND VAN HOOK, W.A. Molecular geometry and the vapor pressures of the isotopic ethylenes: Chemical and structural isomers. (A) Presented at Ann. Meeting Soc. Chim. Phys., Paris, June 1962.
- BIGEISEN, J., RIBNIKAR, S.V. AND VAN HOOK, W.A. Molecular geometry and the vapor pressure of isotopic molecules: The equivalent isomers *cis*-, *gem*-, and *trans*-dideuteroethylenes. Submitted to *J. Chem. Phys.*
- BIGEISEN, J., STERN, M.J. AND VAN HOOK, W.A. Molecular geometry and the vapor pressure of isotopic molecules: C_2H_3D and $C^{12}H_2=C^{13}H_2$. Submitted to *J. Chem. Phys.*
- BLACK, G.W. AND BIELSKI, B.H.J. The radiation-induced oxidation of 4-*t*-butylpyrocatechol in aqueous solutions. *J. Phys. Chem.* **66**, 1203-4 (1962).
- BONNER, F.T., ROTH, E.G., SCHAEFFER, O.A. AND THOMPSON, S.O. Chlorine-36 and deuterium study of Great Basin lake waters. *Geochim. Cosmochim. Acta* **25**, 261-6 (1961).
- CACACE, F. AND WOLF, A.P. The effect of radiation on the reactions of recoil carbon-11 in ammonia. (N) *J. Am. Chem. Soc.* **84**, 3202-3 (1962).

Chemistry Department

- ALLEN, A.O. Hugo Fricke and the development of radiation chemistry: A perspective view. *Radiation Res.* **17**, 255-61 (1962).

- CACACE, F. AND WOLF, A.P. Uso della reazione $N(p,\alpha)C^{11}$ nello studio della chimica di rinculo del carbonio. (N) Submitted to *Gazz. Chim. Ital.*
- CHEN, J.C.Y. Notes on the theory of electron scattering by molecules: The effects of negative-ion states. *J. Chem. Phys.* (in press).
- CHEN, K.L. AND MILLER, J.M. Comparison of nuclear reactions of α -particles with Sc^{45} and of deuterons with Ti^{47} . (A) *Bull. Am. Phys. Soc.* 7, 302 (1962).
- CHRISTMAN, D.R. - See DAWSON, R.F.; SLAVIN, M.; SOLT, M.L.
- COBB, J.C. Determination of lead in meteorites by alpha activation analysis. Submitted to *Geochim. Cosmochim. Acta*.
- COLLINS, K.E. AND HARBOTTLE, G. Effects of very low concentrations of radiation-produced defects on the thermal annealing of (n,γ) -produced Br^{82} in C_2Br_6 . (A) Presented at Am. Chem. Soc. Meeting, Atlantic City, Sept. 1962.
- COPPENS, P. Intramolecular repulsions in substituted aromatic molecules. (L) *J. Chem. Phys.* 36, 2523-5 (1962).
- CORLISS, L.M. AND HASTINGS, J.M. The magnetic structure of manganese chromite. (A) *J. Appl. Phys. Suppl.* 33, 1138 (1962).
- CORLISS, L.M. - See also HASTINGS, J.M.
- CROOKS, J.E. - See BELL, R.P.
- CUMMING, J.B., FRIEDLANDER, G., HUDIS, J. AND POSKANZER, A.M. Spallation of aluminum by 28-Gev protons. *Phys. Rev.* 127, 950-4 (1962).
- CUMMING, J.B., FRIEDLANDER, G. AND KATCOFF, S. $C^{12}(p,pn)C^{11}$ cross section at 28 Gev. *Phys. Rev.* 125, 2078-81 (1962).
- CZAPSKI, G. AND ALLEN, A.O. The reducing radicals produced in water radiolysis: Solutions of oxygen-hydrogen peroxide-hydrogen ion. *J. Phys. Chem.* 66, 262-6 (1962).
- CZAPSKI, G., BIELSKI, B.H.J. AND SUTIN, N. The kinetics of the oxidation of hydrogen peroxide by cerium(IV). (N) *J. Phys. Chem.* (in press).
- CZAPSKI, G. AND SCHWARZ, H.A. The nature of the reducing radical in water radiolysis. *J. Phys. Chem.* 66, 471-4 (1962).
- DAVIS, R. JR. AND STOENNER, R.W. A study of the production of Ar^{37} by 3-Bev protons in a thick sample of iron. (A) Presented at Ann. Meeting Am. Geophys. Union, Washington, April 1962.
- DAVIS, R. JR., STOENNER, R.W. AND SCHAEFFER, O.A. Cosmic ray produced Ar^{37} and Ar^{39} activities in recently fallen meteorites. (A) Presented at IAEA Symp. Radioactive Dating, Athens, Greece, Nov. 1962.
- DAVIS, R. JR. - See also STOENNER, R.W.
- DAWSON, R.F. AND CHRISTMAN, D.R. Biosynthesis of the pyridine ring of the *Nicotiana* alkaloids. Presented at Am. Chem. Soc. Symp. Biosynthesis of Natural Products, Chicago, Sept. 1961.
- DAWSON, R.F. - See also SOLT, M.L.
- DODSON, R.W. - See ROIG, E.
- ELLIOTT, N. Paramagnetism of tetraphenanthroline dihydroxyl diferric chloride. *J. Chem. Phys.* 35, 1273-4 (1961).
- FRIEDLANDER, G. AND YAFFE, L. Electron capture to positron ratios in the decay of Cs^{129} and Cs^{127} . (N) *Can. J. Phys.* 40, 1249-51 (1962).
- FRIEDLANDER, G. - See also CUMMING, J.B.
- FRIEDMAN, L. AND IRSA, A.P. The collision-induced dissociation of N_2O^+ . *J. Chem. Phys.* 36, 1082-3 (1962).
- GROVER, J.R. Effects of angular momentum and gamma-ray emission on excitation functions. *Phys. Rev.* (in press).
- GROVER, J.R. The nuclear reactions of tantalum with 5.7-Gev protons. *Phys. Rev.* 126, 1540-54 (1962).
- HAHN, R.L. AND MILLER, J.M. The reactions of alpha particles with tin-124. *Phys. Rev.* 124, 1879-87 (1961).
- HAMILTON, W.C. Bond distances and thermal motion in ferrous fluosilicate hexahydrate: A neutron diffraction study. *Acta Cryst.* 15, 353-60 (1962).
- HAMILTON, W.C. Comments on recent work on hydrogen atom location and thermal motion determination by neutron diffraction. *Proc. Intern. Conf. Magnetism and Cryst., Kyoto, Japan, Sept. 1961. J. Phys. Soc. Japan* 17, Suppl. B2, 374-5 (1962).
- HAMILTON, W.C. The structure of solids. *Ann. Rev. Phys. Chem.* 13 (1962) (in press).
- HAMILTON, W.C. Use of the minimum function in the solution of the crystal structure of orthorhombic acetamide. (A) Presented at Am. Crystallographic Assoc. Meeting, Villanova, Pa., June 1962.
- HAMILTON, W.C. AND IBERS, J.A. A neutron diffraction study of polycrystalline $HCrO_2$ and $DHCrO_2$. *Proc. Intern. Conf. Magnetism and Cryst., Kyoto, Japan, Sept. 1961. J. Phys. Soc. Japan* 17, Suppl. B2, 383-4 (1962).
- HARBOTTLE, G. - See COLLINS, K.E.; VELJKOVIC, S.R.; YOSHIHARA, K.
- HASCHEMEYER, R.H. - See BIGELEISEN, J.
- HASTINGS, J.M. AND CORLISS, L.M. The magnetic structure of manganese chromite. *Phys. Rev.* 126, 556-65 (1962).
- HASTINGS, J.M. - See also CORLISS, L.M.
- HAYON, E. The photochemistry of iodide ion in aqueous solution. *J. Phys. Chem.* 65, 1937-40 (1961).
- HAYON, E. AND ALLEN, A.O. Evidence for two kinds of "H atoms" in the radiation chemistry of water. *J. Phys. Chem.* 65, 2181-5 (1961).
- HEYMANN, D. AND SCHAEFFER, O.A. Constancy of cosmic rays in time. *Physica* (in press).
- HÖGFELDT, E. A simple averaging approach to surface heterogeneity. Submitted to *J. Phys. Chem.*
- HUDIS, J. - See CUMMING, J.B.
- HUMMELL, A. AND ALLEN, A.O. Radiation chemistry of aqueous solutions of ethanol and the nature of the oxidizing radical OH. *Radiation Res.* 17, 302-11 (1962).
- IBERS, J.A. - See HAMILTON, W.C.
- IRSA, A.P. - See FRIEDMAN, L.
- KATCOFF, S. - See BAKER, E.W.; CUMMING, J.B.
- KAUFMAN, S.B. Isobaric yields in the reactions of 2.9-Bev protons with arsenic-75. *Phys. Rev.* 126, 1189-94 (1962).
- KERR, P.F. - See BASSETT, W.A.
- MC EWEN, W.E. AND WOLF, A.P. Mechanism of the formation of benzylacetophenone and methylethylphenylphosphine from methylethylphenylbenzylidene-phosphorane and styrene oxide. (L) *J. Am. Chem. Soc.* 84, 676-7 (1962).
- MILLER, J.M. - See CHEN, K.L.; HAHN, R.L.
- MOLLENAUER, J.F. Gamma-ray emission from compound nucleus reactions of helium and carbon ions. *Phys. Rev.* 127, 867-79 (1962).

- MOLLENAUER, J.F., WAGNER, S. AND MILLER, G.L. *Silicon Surface Barrier $\Delta E/\Delta x$ Detectors for Nuclear Reaction Studies*. BNL 737 (T-266), May 15, 1962.
- MOLLENAUER, J.F. - See also DONOVAN, P.F. (Physics).
- MOORE, G.G. AND WOLF, A.P. The mechanism of the disproportionation of ethylbenzene-1- C^{14} . (A) Presented at Am. Chem. Soc. Meeting, Washington, March 1962.
- MULLEN, R.T. AND WOLF, A.P. The photolysis of carbon-2- C^{14} -suboxide in ethylene. (L) *J. Am. Chem. Soc.* **84**, 3214-16 (1962).
- NAUGHTON, J.J. Investigations of the potassium-argon method of age determination using solid isotopic dilution and gas removal by crushing. Submitted to *Science*.
- NAUGHTON, J.J. AND SCHAEFFER, O.A. Some preliminary age determinations on Hawaiian lavas. (A) Presented at Intern. Symp. Volcanology, Tokyo, Japan, May 1962.
- PORILE, N.T. Simple nuclear reactions of Ga^{69} and Ga^{71} with high energy protons. *Phys. Rev.* **125**, 1379-85 (1962).
- PORILE, N.T. A study of the $Al^{27}(\alpha, Be^7)Na^{24}$ reaction from threshold to 41 Mev. *Phys. Rev.* **127**, 224-30 (1962).
- PORILE, N.T., POSKANZER, A.M. AND RHO, M. Recoil studies of reactions of indium with medium energy protons and alpha particles. *Phys. Rev.* (in press).
- POSKANZER, A.M. - See CUMMING, J.B.; PORILE, N.T.
- RAPP, D. AND WESTON, R.E. JR. A comment on the calculation of rate constants from transition-state theory. (L) *J. Chem. Phys.* **36**, 2807-8 (1962).
- RHO, N. - See PORILE, N.T.
- RIBNIKAR, S.V. - See BIGELEISEN, J.
- ROIG, E. AND DODSON, R.W. The thallos-thallic exchange at various acidities in perchlorate media. *J. Phys. Chem.* **65**, 2175-81 (1961).
- ROTH, E.G. - See BONNER, F.T.
- RUBINSON, W. The Pb x-rays that accompany Po^{210} decay: Measurement of the M photon yield, and theory. Submitted to *Phys. Rev.*
- RUSH, J.J., TAYLOR, T.I. AND HAVENS, W.W. JR. Proton motions in solids by slow neutron scattering cross sections. (L) *J. Chem. Phys.* **35**, 2265-6 (1961).
- RUSH, J.J., TAYLOR, T.I. AND HAVENS, W.W. JR. Rotational freedom of ammonium ions and methyl groups by cross section measurements with slow neutrons. *J. Chem. Phys.* **37**, 234-8 (1962).
- RUSH, J.J. - See also SAFFORD, G.J. (Physics).
- SAITO, E. AND BIELSKI, B.H. J. The electron paramagnetic resonance spectrum of the HO_2 radical in aqueous solution. (L) *J. Am. Chem. Soc.* **83**, 4467 (1961).
- SAITO, E. - See also BIELSKI, B.H. J.
- SALKIND, A.J. - See SCATTURIN, V.
- SAYRE, E.V. The intentional use of antimony and manganese in ancient glasses. (A) Presented at 6th Intern. Glass Congr., Washington, July 1962.
- SAYRE, E.V. AND SMITH, R.W. Techniques of glass manufacture in antiquity. (A) Presented at 3rd Symp. Archaeol. Chem., Atlantic City, 1962.
- SCATTURIN, V., BELLON, P.L. AND SALKIND, A.J. Neutron diffraction analysis of AgO crystal structure. *J. Electrochem. Soc.* **108**, 819 (1961).
- SCHAEFFER, O.A. - See BASSETT, W.A.; BONNER, F.T.; DAVIS, R. JR.; HEYMANN, D.; NAUGHTON, J.J.
- SCHWARZ, H.A. Determination of some rate constants for the radical processes in the radiation chemistry of water. *J. Phys. Chem.* **66**, 255-62 (1962).
- SCHWARZ, H.A. - See also CZAPSKI, G.; STURM, J.
- SELTZER, S. The secondary α -deuterium isotope effect and the mechanism of the Diels-Alder reaction. (N) *Tetrahedron Letters* **11**, 457-60 (1962).
- SELTZER, S. The secondary α -deuterium isotope effect in the thermal decomposition of α -phenylethylazo-2-propane. Submitted to *J. Am. Chem. Soc.*
- SELTZER, S. - See also WESTON, R.E. JR.
- SLAVIN, M. Effect of slit width on background correction in photographic photometry. *Appl. Spectry.* **16**, 127-9 (1962).
- SLAVIN, M. AND CHRISTMAN, D.R. Dry ice cooling unit for constant-temperature baths between ambient and $-50^\circ C$. *Rev. Sci. Instr.* **33**, 382-3 (1962).
- SMITH, R.W. - See SAYRE, E.V.
- SOLT, M.L., DAWSON, R.F. AND CHRISTMAN, D.R. The gibberellic acid inhibition of nicotine biosynthesis. *Tobacco Science*, Vol. 5, 1961, pp. 95-8, Lockwood Trade Journal Co., New York, 1962.
- STERN, M.J. - See BIGELEISEN, J.
- STOENNER, R.W. AND DAVIS, R. JR. Tests of the spatial constancy of cosmic radiation with the Harleton meteorite. (A) Presented at Ann. Meeting Am. Geophys. Union, Washington, April 1962.
- STOENNER, R.W. - See also BASSETT, W.A.; DAVIS, R. JR.
- STURM, J. AND SCHWARZ, H.A. The helium-ion irradiation of liquid methyl iodide. *Radiation Res.* (in press).
- SUTIN, N. Electron exchange reactions. *Ann. Rev. Nucl. Sci.* (in press).
- SUTIN, N. Homogeneous electron transfer reactions. (A) Presented at Am. Chem. Soc. Meeting, Atlantic City, Sept. 1962.
- SUTIN, N. - See also BELL, R.S.; CZAPSKI, G.
- THOMPSON, S.O. A search for Ar^{42} in the atmosphere. (A) Presented at Ann. Meeting Am. Geophys. Union, Washington, April 1962.
- THOMPSON, S.O. - See also BONNER, F.T.
- UNSEREN, E. AND WOLF, A.P. The disproportionation of ethylbenzene-1- C^{14} , under the influence of aluminum bromide and hydrogen bromide. *J. Org. Chem.* **27**, 1509-13 (1962).
- VAN HOOK, W.A. - See BIGELEISEN, J.
- VELJKOVIC, S.R. AND HARBOTTLE, G. Low-temperature investigation of the Szilard-Chalmers reaction in potassium permanganate. (L) *J. Inorg. Nucl. Chem.* **23**, 159-60 (1961).
- VELJKOVIC, S.R. AND HARBOTTLE, G. Recoil-atom reactions at low temperature. *Nature* **191**, 1287-8 (1961).
- WEISSMAN, S.I. Relation between absorption and fluorescence spectra of triphenylmethyl. (N) *J. Chem. Phys.* (in press).
- WESTON, R.E. JR. Isotope effects in chemical reactions. *Ann. Rev. Nucl. Sci.* **11**, 439-60 (1961).
- WESTON, R.E. JR. AND SELTZER, S. The secondary deuterium isotope effect in the pyrolysis of dimethylmercury. Submitted to *J. Phys. Chem.*
- WESTON, R.E. JR. - See also RAPP, D.
- WITHNELL, R. A 4π proportional counter suitable for 4π β - γ coincidence counting and routine use. *Nucl. Instr. Methods* **14**, 279-80 (1961).

- WOLF, A.P. - See CACACE, F.; McEWEN, W.E.; MOORE, G.G.; MULLEN, R.T.; UNSEREN, E.
- WOLFSBERG, M. Isotope effects on intermolecular interactions and isotopic vapor pressure differences. Submitted to *J. Chim. Phys.*
- WOLFSBERG, M. Rate constants for the unimolecular decomposition of excited molecular species near threshold. *J. Chem. Phys.* **36**, 1072-7 (1962).
- WOLFSBERG, M. - See also BIGELEISEN, J.
- YAFFE, L. - See FRIEDLANDER, G.
- YANKWICH, P.E. - See BIGELEISEN, J.
- YOSHIHARA, K. AND HARBOTTLE, G. Recoil atom reactions and annealing processes in cobaltic hexamine bromide. Submitted to *Radiochimica Acta*.

Instrumentation and Health Physics Department

- CARSTEN, A. - See INNES, J.R.M. (Biology).
- CHASE, R.L. Integrator and digitizer - IH-78-1. Informal Report BNL 5619.
- CHASE, R.L. Recent developments in multichannel pulse-height analysis. *IRE Trans. Nucl. Sci.* **NS-9**, No. 3, 275-9 (1962).
- CHASE, R.L. A servo-stabilized analogue-to-digital converter for high resolution pulse-height analysis. *IRE Trans. Nucl. Sci.* **NS-9**, No. 1, 119-22 (1962).
- CHASE, R.L. Servo-stabilized RF vernier time-of-flight analyzer. Informal Report BNL 6439.
- CHASE, R.L. - See also FISCHER, J.
- COWAN, F.P. Certification of health physicists. *Health Phys.* **8**, 109-11 (1961).
- COWAN, F.P. A preliminary report on health physics problems at the Brookhaven Alternating Gradient Synchrotron (AGS). Presented at 1st Intern. Conf. Shielding Around High Energy Accelerators, Saclay, France, Jan. 1962.
- DEL CAMPO, R.G. - See SINGER, I.A.
- DISTENFELD, C.H. A description of the BNL research reactor environmental air control, alarm, and monitoring system. *7th AEC Air Cleaning Conf., Brookhaven National Laboratory, Oct. 1961*, pp. 8-22, TID-7627, March 1962.
- DONOVAN, P.F. - See GIBSON, W.M.
- FISCHER, J., HIGINBOTHAM, W.A. AND CHASE, R.L. A gated hodoscope readout system for several simultaneous inputs retaining pulse height information. Informal Report BNL 5620.
- FISCHER, J. AND ZORN, G.T. Reduction of delay between particle passage and spark chamber spark. *IRE Trans. Nucl. Sci.* **NS-9**, No. 3, 261-74 (1962).
- GIBSON, W.M., MILLER, G.L. AND DONOVAN, P.F. Semiconductor particle spectrometers. Chapter in *Alpha, Beta, Gamma Spectroscopy*, 2nd Edition, K. Siegbahn, Editor, North-Holland Publishing Co., Amsterdam (in press).
- GOSCINSKI, S.C. Drop counter. Informal Report BNL 5622.
- HIGINBOTHAM, W.A. Instrumentation for use with Bev accelerators. *IRE Trans. Nucl. Sci.* **NS-8**, No. 4, 34-9 (1961).
- HIGINBOTHAM, W.A. Recent advances in instrumentation. Chapter in *Symposia on Advances in Tracer Methodology*, Plenum Press, New York (in press).
- HIGINBOTHAM, W.A., SUGARMAN, R.M., POTTER, D.W. AND ROBERTSON, J.S. A direct analogue computer for multi-compartment systems. Submitted to *Ann. N.Y. Acad. Sci.*
- HIGINBOTHAM, W.A. - See also FISCHER, J.; PALMER, J.P.; RANKOWITZ, S.; SUGARMAN, R.
- JACOBS, J. Beam current integrator. Informal Report BNL 6438.
- KAZUHIKO, I. - See SINGER, I.A.
- LLACER, J. Low frequency amplifier IH-130-1. Informal Report BNL 6437.
- MERKLE, W. - See PALMER, J.P.
- MERRITT, F.C. - See SUGARMAN, R.
- MILLER, D.M. 6-Kilovolt capacitor charging supply. Informal Report BNL 6442.
- MILLER, D.M. 5-Kilovolt, 1.25-kilowatt capacitor charging supply. Informal Report BNL 5618.
- MILLER, D.M. 0 to 2-Kilovolt flash tube supplies. Informal Report BNL 6443.
- MILLER, D.M. 4-Kilovolt power supply. Informal Report BNL 5621.
- MILLER, D.M. Power supply for BHG lamp. Informal Report BNL 6444.
- MILLER, D.M. AND PALMER, J.P. 3-Kilojoule, 5-kilovolt flash tube capacitor charging supply. Informal Report BNL 5617.
- MILLER, G.L. *The Physics of Semiconductor Radiation Detectors, Brookhaven Lecture Series No. 9*, BNL 699 (T-241), Sept. 27, 1961.
- MILLER, G.L. - See also GIBSON, W.M.; MOLLENAUER, J.F. (Chemistry).
- NAGLE, C.M. - See SINGER, I.A.
- OGDEN, E.C. - See RAYNOR, G.S.
- PALMER, J.P., HIGINBOTHAM, W.A. AND MERKLE, W. Solid state precision current shunt for experimental magnets. Informal Report BNL 5662.
- POTTER, D.W. - See HIGINBOTHAM, W.A.
- RANKOWITZ, S., ROBERTSON, J.S., HIGINBOTHAM, W.A. AND ROSENBLUM, M. J. Positron scanner for locating brain tumors. *IRE Intern. Conv. Record* (in press).
- RANKOWITZ, S. - See also FINE, S. (Medical).
- RAYNOR, G.S., OGDEN, E.C., SMITH, M.E. AND SINGER, I.A. Pollen sampling and dispersion studies at Brookhaven National Laboratory. *J. Air Pollution Control Assoc.* **11**, 557-62 (1961).
- ROGERS, E.J. Van de Graaff proton source receives 110 kilovolt boosts. *Electronics* **35**, No. 13, 58-9 (1962).
- ROSENBLUM, M. J. - See RANKOWITZ, S.
- SINGER, I.A., KAZUHIKO, I. AND DEL CAMPO, R.G. Peak to mean pollutant concentration ratios for various terrain and vegetation cover. (A) *J. Air Pollution Control Assoc.* (in press).
- SINGER, I.A. AND NAGLE, C.M. *A Study of the Wind Profile in the Lowest 400 Feet of the Atmosphere. Progress Report No. 9*, BNL 697 (T-239), Sept. 1961; *Final Report*, BNL 718 (T-254), March 31, 1962.
- SINGER, I.A. - See also RAYNOR, G.S.
- SMITH, M.E. - See RAYNOR, G.S.
- SUGARMAN, R.M. Millimicronote No. 22: Evaluation of Hamner XN-690 coincidence circuit. Informal Report BNL 6441.
- SUGARMAN, R.M. Nanosecond logic and its applications to nuclear instrumentation. (A) Presented at IRE Pro-

- fess. Group Nucl. Sci. Meeting, Garden City, L.I., Feb. 1962.
- SUGARMAN, R.M. Random note No. 2 on HPD frame identification. Informal Report BNL 5624.
- SUGARMAN, R.M. Random notes on HPD frame identification problem. Informal Report BNL 5623.
- SUGARMAN, R., MERRITT, F.C. AND HIGINBOTHAM, W.A. *Nanosecond Counter Circuit Manual*. BNL 711 (T-248), Feb. 1962.
- SUGARMAN, R.M. - See also HIGINBOTHAM, W.A.
- WAGNER, S. - See MOLLENAUER, J.F. (Chemistry).
- WALL, G.N. Digital-to-analogue converter (punched tape to X-Y plotter). Informal Report BNL 6440.

Medical Department

- ADAMIK, E.A. - See CRONKITE, E.P.; JANSEN, C.R.
- APONTE, G.E. - See SHELLABARGER, C.J.
- ARIEL, I. Irradiation with interstitial radioactive palladium. (A) Presented at Soc. Nucl. Med. Meeting, Dallas, Texas, June 1962.
- BAKER, H. - See HOOGSTRAATON, B.
- BATEMAN, J.L. AND BOND, V.P. Dependence of short-term mammalian radiation effects on dose rate. (A) Presented at Radiological Soc. N. Am. Meeting, Chicago, Nov. 1961.
- BATEMAN, J.L., ROSSI, H.H., BOND, V.P. AND GILMARTIN, J. The dependence of RBE upon energy of fast neutrons. 2. Biological evaluation at discrete neutron energies in the range 0.43 to 1.80 Mev. *Radiation Res.* 15, 694-706 (1961).
- BERTINCHAMPS, A.J. - See PAPAVALIOU, P.S.
- BIERMAN, E.L. - See DREILING, D.A.
- BOND, V.P., FLIEDNER, T.M. AND USENIK, E. Early bone marrow hemorrhage in the irradiated dog. *Arch. Pathol.* 73, 13-29 (1962).
- BOND, V.P., ODARTCHENKO, N., COTTIER, H., FEINENDEGEN, L. AND CRONKITE, E.P. The kinetics of the more mature erythrocytic precursors studied with tritiated thymidine. Presented at Conf. Erythropoietin and Regulation of Erythropoiesis, San Francisco, Jan. 1962.
- BOND, V.P., RUBINI, J.R., FLIEDNER, T.M. AND CRONKITE, E.P. Study of the effect of therapy on cells in the blood of leukemic patients capable of incorporating H^3 -thymidine. *J. Lab. Clin. Med.* 59, 412-18 (1962).
- BOND, V.P. - See also BATEMAN, J.L.; FEINENDEGEN, L.E.; FLIEDNER, T.M.; KILLMANN, S.A.; KLIGERMAN, M.M.; PERMAN, V.; RUBINI, J.R.; SHELLABARGER, C.J.; SPRARAGEN, S.C.; STONER, R.D.; TSUYA, A.
- BORG, D.C. AND COTZIAS, G.C. Interaction of trace metals with phenothiazine drug derivatives: I. Structure-reactivity correlations; II. Formation of free radicals; III. Theoretical part. *Proc. Nat'l. Acad. Sci. U.S.* 48, 617-52 (1962).
- BORG, D.C. - See also COTZIAS, G.C.
- BRECHER, G. - See DEBONS, A.F.; KILLMANN, S.A.
- BROWN, R.R. - See HANKES, L.V.
- COHN, S.H., CONARD, R.A., GUSMANO, E.A. AND ROBERTSON, J.S. Use of a portable whole-body counter to measure internal contamination in a fallout-exposed population. *Health Phys. J.* (in press).
- COHN, S.H., LIPPINCOTT, S.W., GUSMANO, E.A., FINE, S. AND JANSEN, C. Short-term kinetics of Ca^{47} and Sr^{85} in man. (A) Presented at 2nd Intern. Congr. Radiation Res., Harrogate, England, Aug. 1962.
- COHN, S.H., LIPPINCOTT, S.W., GUSMANO, E.A. AND ROBERTSON, J.S. Comparative kinetics of Ca^{47} and Sr^{85} in man. Submitted to *Intern. J. Radiation Biol.*
- COHN, S.H., SPENCER, H., SAMACHSON, J., FELDSTEIN, A. AND GUSMANO, E.A. Influence of dietary stable strontium and calcium on the turnover of bone-fixed Sr^{85} in man. *Proc. Soc. Exptl. Biol. Med.* 110, 526-8 (1962).
- COHN, S.H., SPENCER, H., SAMACHSON, J. AND ROBERTSON, J.S. The long-term component of Sr turnover in man. Presented at Soc. Nucl. Med. Meeting, Dallas, Texas, June 1962.
- COHN, S.H., SPENCER, H., SAMACHSON, J. AND ROBERTSON, J.S. The turnover of strontium-85 in man as determined by whole-body counting. *Radiation Res.* 17, 173-85 (1962).
- COHN, S.H. - See also CONARD, R.A.; PRICE, D.C.; REIZENSTEIN, P.G.; ROBERTSON, J.S.
- CONARD, R.A., MACDONALD, H.E., MEYER, L.M., COHN, S., SUTOW, W.W., KARNOFSKY, D., JAFFE, A.A. AND RIKLON, E. *Medical Survey of Rongelap People Seven Years After Exposure to Fallout*. BNL 727 (T-260), May 1962.
- CONARD, R.A. - See also COHN, S.H.
- COTTIER, H., CRONKITE, E.P., TONNA, E.A. AND NIELSEN, N.O. Leukemogenic effect of whole-body Co^{60} - γ -irradiation compared with H^3 -thymidine and H^3 -cytidine: Preliminary report on the development of thymic lymphomas in $C^{57}BL/6J$ mice. Presented at IAEA/UNESCO Symp. Aetiology of Late Somatic Effects of Ionizing Radiations, London, March 1962.
- COTTIER, H., ROOS, B., RENTSCH, H. AND CRONKITE, E.P. An electron-microscopic study of *in vivo* primary thrombus formation. (A) Am. Soc. Hematol. Meeting, Los Angeles, Nov. 1961.
- COTTIER, H. - See also BOND, V.P.
- COTZIAS, G.C., BORG, D.C. AND SELLECK, B. Specificity of the zinc pathway through the body: The turnover of Zn^{65} in the mouse. *Am. J. Physiol.* 202, 359-63 (1962).
- COTZIAS, G.C., BORG, D.C. AND SELLECK, B. Virtual absence of turnover in cadmium metabolism: Cd studies in the mouse. *Am. J. Physiol.* 201, 927-30 (1961).
- COTZIAS, G.C. AND PAPAVALIOU, P.S. The state of binding of natural manganese in human cerebrospinal fluid, blood and plasma. *Nature* 195, 823-4 (1962).
- COTZIAS, G.C. - See also BORG, D.C.; HUGHES, E.R.; PAPAVALIOU, P.S.
- CRONKITE, E.P., JANSEN, C.R., MATHER, G., ADAMIK, E.A. AND SIPE, C.R. The induction of a prolonged lymphopenia by extracorporeal irradiation of the blood of the calf. (A) *Radiation Res.* 16, 586 (1962).
- CRONKITE, E.P., JANSEN, C.R., MATHER, G., NIELSEN, N., USENIK, E.A., ADAMIK, E. AND SIPE, C.R. Studies on lymphocytes. I. Lymphopenia by prolonged extracorporeal irradiation of circulating blood. *Blood* 20, 203-13 (1962).
- CRONKITE, E.P. - See also BOND, V.P.; COTTIER, H.; DEBONS, A.F.; FLIEDNER, T.M.; HUGHES, E.R.; HWANG, W.S.; JANSEN, C.R.; KILLMANN, S.A.; MEYER, L.M.; PERMAN, V.; PRICE, D.C.; REIZENSTEIN, P.G.; RUBINI,

- J.R.; SCHWARTZ, I.L.; SHELLABARGER, C.J.; SIPE, C.R.; TONNA, E.A.
- DAHL, L.K. *Excessive Salt Intake and Hypertension: A Dietary and Genetic Interplay, Brookhaven Lecture Series No. 12*, BNL 733 (T-263), Dec. 13, 1961.
- DAHL, L.K. Experimental hypercholesterolemia from chronic excess salt feeding. *Am. J. Cardiol.* **8**, 705-18 (1961).
- DAHL, L.K. The possible role of chronic excess salt consumption in the pathogenesis of essential hypertension. *Am. J. Cardiol.* **8**, 571-5, 587-92 (1961).
- DAHL, L.K. AND HEINE, M. The enhanced hypertensogenic effect of sea salt over sodium chloride. *Am. J. Cardiol.* **8**, 726-31, 754-6 (1961).
- DAHL, L.K., HEINE, M. AND TASSINARI, L. Effects of chronic excess salt ingestion. Evidence that genetic factors play an important role in susceptibility to experimental hypertension. *J. Exptl. Med.* **115**, 1173-90 (1962).
- DAHL, L.K., HEINE, M. AND TASSINARI, L. Role of genetic factors in susceptibility to experimental hypertension due to chronic excess salt ingestion. *Nature* **194**, 480-2 (1962).
- DAHL, L.K., SMILAY, M.G., SILVER, L. AND SPRARAGEN, S.C. Evidence for a prolonged biological half-life of Na^{22} in patients with hypertension. *Circulation Res.* **10**, Pt. 1, 313-20 (1962).
- DAHL, L.K., SMILAY, M.G., SILVER, L. AND SPRARAGEN, S.C. Prolonged biological half-life of sodium-22 in patients with essential hypertension. *Nature* **192**, 267-8 (1961).
- DAHL, L.K. - See also OMENN, G.S.; SPRARAGEN, S.C.
- DEBONS, A.F., SILVER, L., CRONKITE, E.P., JOHNSON, H.A., BRECHER, G., TENZER, D. AND SCHWARTZ, I.L. Localization of gold in the mouse brain in relation to gold-thioglucose obesity. *Am. J. Physiol.* **202**, 743-50 (1962).
- DEBONS, A.F. - See also DREILING, D.A.; SCHWARTZ, I.L.
- DE MEUTTER, R.C. - See SHREEVE, W.W.
- DREILING, D.A., BIERMAN, E.L., DEBONS, A.F., ELSBACH, P. AND SCHWARTZ, I.L. The effect of ACTH, hydrocortisone, and glucagon on the plasma non-esterified fatty acid concentration (NEFA) in normal subjects and in patients with liver disease. *Metabolism* **11**, 572-8 (1962).
- DREW, R.M. AND PAINTER, R.B. Further studies on the clonal growth of HeLa S3 cells treated with tritiated thymidine. *Radiation Res.* **16**, 303-11 (1962).
- DREW, R.M. - See also FEINENDEGEN, L.E.
- EASTERDAY, O.D. AND HAMEL, H. Acute intravenous and intraperitoneal toxicity studies on sodium tetraborate decahydrate. *Arch. Intern. Pharmacodyn.* (in press).
- EASTERDAY, O.D. - See also FARR, L.E.
- EISENTRAUT, A. - See RUBINI, J.R.
- ELSBACH, P. - See DREILING, D.A.
- FARR, L.E. Discussion of lecture by Dr. C.A. Tobias, "Biological effects of heavy radiation particles." Presented at World Federation Neurosurgical Societies Meeting, Washington, Oct. 1961.
- FARR, L.E. The use of radioisotopes in medicine. *Proc. Youth Conf., Chicago, Nov. 1961* (in press).
- FARR, L.E. AND EASTERDAY, O.D. Radioactive isotopes - metabolic, diagnostic, and therapeutic uses. Submitted to *J. Gerontol.*
- FARR, L.E. AND KONIKOWSKI, T. Absolute control of an experimental sarcoma in mice by boron-thermal neutron interaction. (A) Presented at Am. Soc. Exptl. Pathol. Meeting, Atlantic City, April 1962.
- FARR, L.E. AND KONIKOWSKI, T. Selective tumor response to neutron capture therapy of transplantable neoplasms in the mouse. (A) Presented at Soc. Nucl. Med. Meeting, Dallas, Texas, June 1962.
- FARR, L.E. - See also FINE, S.
- FEINENDEGEN, L.E. AND BOND, V.P. Differential uptake of H^3 -thymidine into the soluble fraction of single bone marrow cells, determined by autoradiography. *Exptl. Cell Res.* (in press).
- FEINENDEGEN, L.E., BOND, V.P. AND HUGHES, W.L. RNA mediation in DNA synthesis in human cancer cells in culture (HeLa) studied with tritium-labeled cytidine and thymidine. *Exptl. Cell Res.* **25**, 627-47 (1961).
- FEINENDEGEN, L.E. - See also BOND, V.P.
- FELDSTEIN, A. - See COHN, S.H.
- FINE, S., ROBERTSON, J.S., RANKOWITZ, S. AND FARR, L.E. Intravenous infusion droplet indicator. Submitted to *Electronics*.
- FINE, S., WOLINS, W., ROBERTSON, J.S. AND FARR, L.E. Electrode system for serial electrocardiographic tracings. Submitted to *J. Appl. Physiol.*
- FINE, S. - See also COHN, S.H.
- FLIEDNER, T.M., CRONKITE, E.P. AND BOND, V.P. Pathogenesis and regeneration of radiation-induced bone marrow injury and therapeutic implications. Submitted to *Strahlentherapie*.
- FLIEDNER, T.M., CRONKITE, E.P. AND BOND, V.P. Potentials and limitations of H^3 -thymidine labeling of hemopoietic cell systems in the study of their dynamics of proliferation. Presented at 8th Congr. European Soc. Haematol., Vienna, Aug. 28-Sept. 2, 1961.
- FLIEDNER, T.M., STODTMEISTER, R., MEYER, L. AND CRONKITE, E.P. Problems of bone marrow cell trans-fusions. Presented at Symp. Radioactive Isotopes in Hematology, Freiburg, West Germany, March 1962.
- FLIEDNER, T.M. - See also BOND, V.P.; KILLMANN, S.A.; RUBINI, J.R.; TSUYA, A.
- FORSYTH, E.M. - See PRICE, D.C.
- GILMARTIN, J. - See BATEMAN, J.L.
- GULIBON, J. - See PAPAVALIOU, P.S.
- GUSMANO, E.A. - See COHN, S.H.
- HALE, W.M. - See STONER, R.D.
- HAMEL, H. - See EASTERDAY, O.D.
- HANKES, L.V., BROWN, R.R. AND RAI, K. Metabolism of DL-tryptophan-2- C^{14} in human subjects. (A) Presented at Am. Soc. Biol. Chemists Meeting, Atlantic City, April 1962.
- HANKES, L.V., SCHMAELER, M. AND RAI, K. o-Aminophenol: A urinary product of tryptophan metabolism in the human. *Proc. Soc. Exptl. Biol. Med.* **110**, 420-2 (1962).
- HANKES, L.V., SCHMAELER, M. AND STONER, R.D. Metabolism of C^{14} -carboxyl-labeled 3-hydroxyanthranilic acid in normal and neoplastic mice. Submitted to *Proc. Soc. Exptl. Biol. Med.*
- HAYANO, I. - See TSUYA, A.
- HEINE, M. - See DAHL, L.K.

- HELD, B.T. - See KLIGERMAN, M.M.
- HOOGSTRATON, B., BAKER, H. AND REIZENSTEIN, P.G. Correlation between serum folic acid activity and response to antifolate therapy. (A) Presented at Am. Soc. Hematol. Meeting, Los Angeles, Nov. 1961.
- HUGHES, E.R. AND COTZIAS, G.C. Adrenocorticosteroid hormones and manganese metabolism. *Am. J. Physiol.* **201**, 1061-4 (1961).
- HUGHES, E.R., COTZIAS, G.C. AND CRONKITE, E.P. Total-body irradiation and manganese metabolism. *Nature* **193**, 792-3 (1962).
- HUGHES, W.L. - See FEINENDEGEN, L.E.; PAINTER, R.B.
- HWANG, W.S., TONNA, E.A. AND CRONKITE, E.P. The localization and distribution of tritiated histidine in growing mouse incisors. *Nature* **193**, 896 (1962).
- JANSEN, C.R., CRONKITE, E.P., MATHER, G., NIELSEN, N.O., RAI, K., ADAMIK, E.A. AND SIPE, C.R. Studies on lymphocytes. II. The production of lymphocytosis by intravenous heparin in calves. *Blood* **20**, 443-52 (1962).
- JANSEN, C.R. - See also COHN, S.H.; CRONKITE, E.P.
- JOHNSON, H.A. - See DEBONS, A.F.; SCHWARTZ, I.L.
- JONES, I. - See MEYER, L.M.
- KELLER, S. - See RUBINI, J.R.
- KILLMANN, S.A., CRONKITE, E.P., BOND, V.P. AND FLIEDNER, T.M. Kinetics of human leukemic cells studied with tritiated thymidine *in vivo*. (A) Presented at 8th Congr. European Soc. Haematol., Vienna, Aug. 28-Sept. 2, 1961.
- KILLMANN, S.A., CRONKITE, E.P., FLIEDNER, T.M. AND BOND, V.P. Cell proliferation in multiple myeloma studied with tritiated thymidine. *Lab. Invest.* (in press).
- KILLMANN, S.A., CRONKITE, E.P., FLIEDNER, T.M. AND BOND, V.P. Mitotic indices of human bone marrow cells. I. Number and cytologic distribution of mitoses. *Blood* **19**, 743-50 (1962).
- KILLMANN, S.A., CRONKITE, E.P., FLIEDNER, T.M., BOND, V.P. AND BRECHER, G. Mitotic indices of human bone marrow cells. II. The use of mitotic indices for estimation of the time parameters of the generative cycle particularly in serially connected multiplicative compartments. Submitted to *Blood*.
- KLIGERMAN, M.M., BOND, V.P., HELD, D.T., MATHER, G. AND NIELSEN, N. Anoxic protection of the small intestine of dogs by intra-arterial catheterization. *Radiology* (in press).
- KONIKOWSKI, T. - See FARR, L.E.
- LIPPINCOTT, S.W. - See COHN, S.H.
- MATHER, G. - See CRONKITE, E.P.; JANSEN, C.R.; KLIGERMAN, M.M.; OMENN, G.S.
- MAXFIELD, M. AND WOLINS, W. A molecular abnormality of urinary mucoprotein in cystic fibrosis of the pancreas. *J. Clin. Invest.* **41**, 455-62 (1962).
- MEYER, L.M., CRONKITE, E.P., MILLER, I.F., MULZAC, C.W. AND JONES, I. Co⁶⁰ vitamin B₁₂ binding capacity of human leukocytes. *Blood* **19**, 229-35 (1962).
- MEYER, L.M., REIZENSTEIN, P.G. AND CRONKITE, E.P. Serum binding of vitamin B₁₂ analogues: Identification of binding groups in the B₁₂ molecule. Submitted to *Blood*.
- MEYER, L.M. - See also CONARD, R.A.; FLIEDNER, T.M.
- MILLER, I.F. - See MEYER, L.M.; PAPAVALIOU, P.S.
- MULZAC, C.W. - See MEYER, L.M.
- NIELSEN, N.O. - See COTTIER, H.; CRONKITE, E.P.; JANSEN, C.R.; KLIGERMAN, M.M.
- ODARTCHENKO, N. - See BOND, V.P.
- OMENN, G.S., DAHL, L.K. AND MATHER, G.W. Effects of chronic excess salt feeding. Demonstration of normal thyroid function in salt-fed dogs with hypercholesterolemia: Thyroidal I¹³¹ uptake, plasma protein bound iodine, and histological studies. Submitted to *Circulation Res.*
- PAINTER, R.B. AND HUGHES, W.L. Nucleic acid metabolism and the lethal effect of radiation on cultured human cells (HeLa). *Ann. N.Y. Acad. Sci.* **95**, 960-8 (1961).
- PAINTER, R.B. - See also DREW, R.M.
- PAPAVALIOU, P.S., BERTINCHAMPS, A.J., MILLER, S.T., GULIBON, J. AND COTZIAS, G.C. Procedures for the study of bile physiology in the rat. Submitted to *J. Appl. Physiol.*
- PAPAVALIOU, P.S. - See also COTZIAS, G.C.
- PAVELEC, M. - See TONNA, E.A.
- PERMAN, V., CRONKITE, E.P., BOND, V.P. AND SORENSEN, D.K. The regenerative ability of hemopoietic tissue following lethal x-irradiation in dogs. *Blood* **19**, 724-37 (1962).
- PERMAN, V., SORENSEN, D.K., USENIK, E.A., BOND, V.P. AND CRONKITE, E.P. Hemopoietic regeneration in control and recovered heavily irradiated dogs following severe hemorrhage. *Blood* **19**, 738-42 (1962).
- PLAZIN, J. - See VAN SLYKE, D.D.
- PRICE, D.C., COHN, S.H., WASSERMAN, L.R., REIZENSTEIN, P.G. AND CRONKITE, E.P. The determination of iron absorption and loss by whole-body counting. *Blood* **20**, 517-31 (1962).
- PRICE, D.C., FORSYTH, E.M., COHN, S.H. AND CRONKITE, E.P. The study of menstrual blood loss and consequent iron deficiency by Fe⁵⁹ whole-body counting. (A) Presented at Can. Soc. Clin. Invest. Meeting, Jan. 1962.
- RAI, K. - See HANKES, L.V.; JANSEN, C.R.
- REIZENSTEIN, P.G., CRONKITE, E.P. AND COHN, S.H. Measurement of absorption of vitamin B₁₂ by whole-body gamma spectrometry. *Blood* **18**, 95-101 (1961).
- REIZENSTEIN, P.G., ROBERTSON, J.S. AND CRONKITE, E.P. Relations of the turnover of tracer vitamin B₁₂ to that of unlabeled B₁₂ in the body stores. Presented at 2nd European Symp. Vitamin B₁₂ and Intrinsic Factor, Hamburg, July 1961.
- REIZENSTEIN, P.G. - See also HOOGSTRATON, B.; MEYER, L.M.; PRICE, D.C.
- RENTSCH, H. - See COTTIER, H.
- ROBERTSON, J.S. Summary - Physiological Aspects (N.Y. Acad. Sci. Conf. Multicompartment Analysis of Tracer Experiments, New York City, May 1962). Submitted to *Ann. N.Y. Acad. Sci.*
- ROBERTSON, J.S. AND COHN, S.H. Use of analogue computers in studies of strontium and calcium metabolism. (A) Presented at N.Y. Acad. Sci. Conf. Multicompartment Analysis of Tracer Experiments, New York City, May 1962.
- ROBERTSON, J.S. - See also ARIEL, I.; COHN, S.H.; FINE, S.; HIGINBOTHAM, W.A. (Instr. & Health Physics); RANKOWITZ, S. (Instr. & Health Physics); REIZENSTEIN, P.

- ROOS, B. - See COTTIER, H.
- ROSSI, H.H. - See BATEMAN, J.L.
- RUBINI, J.R., BOND, V.P., KELLER, S., FLIEDNER, T.M. AND CRONKITE, E.P. DNA synthesis in circulating blood leukocytes labeled *in vitro* with H^3 -thymidine. *J. Lab. Clin. Med.* **58**, 751-62 (1961).
- RUBINI, J.R. AND CRONKITE, E.P. Blockade of H^3 -thymidine utilization *in vitro*. Presented at 8th Congr. European Soc. Haematol., Vienna, Aug. 28-Sept. 2, 1961.
- RUBINI, J.R., KELLER, S., EISENTRAUT, A. AND CRONKITE, E.P. *In-vitro* metabolism of H^3 -thymidine. *Tritium in the Physical and Biological Sciences*, Vol. 2, pp. 247-67, IAEA, Vienna, 1962.
- RUBINI, J.R. - See also BOND, V.P.
- SAMACHSON, J. - See COHN, S.H.
- SANEN, F.J. - See TONNA, E.A.
- SCHMAELER, M. - See HANKES, L.V.
- SCHWARTZ, I.L., CRONKITE, E.P., JOHNSON, H.A., SILVER, L., TENZER, D. AND DEBONS, A. Radioautographic localization of the "satiety center." *Trans. Assoc. Am. Physicians* **74**, 300-17 (1961).
- SCHWARTZ, I.L. - See also DEBONS, A.F.; DREILING, D.A.
- SELLECK, B. - See COTZIAS, G.C.
- SHELLABARGER, C.J., APONTE, G.E., CRONKITE, E.P. AND BOND, V.P. Studies on radiation-induced mammary gland neoplasia in the rat. VI. The effect of changes in thyroid function, ovarian function, and pregnancy. *Radiation Res.* (in press).
- SHELLABARGER, C.J., BOND, V.P. AND CRONKITE, E.P. Studies on radiation-induced mammary gland neoplasia in the rat. VII. The effects of fractionation and protraction of sublethal total-body radiation. *Radiation Res.* **17**, 101-9 (1962).
- SHELLABARGER, C.J. - See also TONNA, E.A.
- SHREEVE, W.W. Diabetic ketosis. Submitted to *Trans. N.Y. Acad. Sci.*
- SHREEVE, W.W. AND DE MEUTER, R.C. Pathways of carbohydrate formation in man. III. Asymmetric labeling of blood glucose by DL-lactate or pyruvate-2- or 3- C^{14} in diabetic and nondiabetic subjects. Submitted to *J. Biol. Chem.*
- SILVER, L. - See DAHL, L.K.; DEBONS, A.F.; SCHWARTZ, I.L.
- SIPLE, C.R. AND CRONKITE, E.P. Studies on the application of the Coulter electronic counter in enumeration of platelets. *Ann. N.Y. Acad. Sci.* **99**, 262-70 (1962).
- SIPLE, C.R. - See also CRONKITE, E.P.; JANSEN, C.R.
- SMILAY, M.G. - See DAHL, L.K.
- SORENSEN, D.K. - See PERMAN, V.
- SPENCER, H. - See COHN, S.H.
- SPRARAGEN, S.C., BOND, V.P. AND DAHL, L.K. DNA synthesizing cells in rabbit heart tissue after cholesterol feeding. *Circulation Res.* (in press).
- SPRARAGEN, S.C., BOND, V.P. AND DAHL, L.K. Role of hyperplasia in vascular lesions of cholesterol fed rabbits studied with thymidine- H^3 autoradiography. *Circulation Res.* **11**, 329-36 (1962).
- SPRARAGEN, S.C. - See also DAHL, L.K.
- STODTMEISTER, R. - See FLIEDNER, T.M.
- STONER, R.D. AND BOND, V.P. Antibody formation by transplanted bone marrow, spleen, thymus, and lymph node cells in isologous, homologous, and heterologous recipients. (A) Submitted to *Blood*.
- STONER, R.D. AND HALE, W.M. Radiation effects on primary and secondary antibody responses. *The Effects of Ionizing Radiation on Immune Processes*, Charles A. Leone, Editor, pp. 183-219, Gordon & Breach, New York, 1962.
- STONER, R.D. - See also HANKES, L.V.; TERRES, G.
- SUTOW, W.W. - See CONARD, R.A.
- TASSINARI, L. - See DAHL, L.K.
- TENZER, D. - See DEBONS, A.F.; SCHWARTZ, I.L.
- TERRES, G. AND STONER, R.D. Specificity of enhanced immunological sensitization of mice following injections of antigens and specific antisera. *Proc. Soc. Exptl. Biol. Med.* **109**, 881-91 (1962).
- TONNA, E.A. An autoradiographic examination of the utilization of tritiated histidine by cells of the skeletal system. *Nature* **193**, 1301-2 (1962).
- TONNA, E.A. Changes in the connective tissue framework of long bones during aging: A polarized study following metachromatic staining. (A) Presented at Ann. Meeting Gerontological Soc., Miami, Nov. 1962.
- TONNA, E.A. The connective tissue framework of the femur. *Anat. Record* (in press).
- TONNA, E.A. The localization and distribution of tritiated histidine in bone and changes concomitant with aging. (A) *J. Gerontol.* **16**, No. 4 (1961).
- TONNA, E.A. AND CRONKITE, E.P. An autoradiographic study of periosteal cell proliferation with tritiated thymidine. *Lab. Invest.* **11**, 455-62 (1962).
- TONNA, E.A. AND CRONKITE, E.P. Changes in the skeletal cell proliferative response to trauma concomitant with aging. *J. Bone Joint Surg.* **44A**, 144-64 (1962).
- TONNA, E.A. AND CRONKITE, E.P. Utilization of tritiated histidine (H^3 HIL) by skeletal cells of adult mice. *J. Gerontol.* (in press).
- TONNA, E.A., CRONKITE, E.P. AND PAVELEC, M. An autoradiographic study of the localization and distribution of tritiated histidine in bone. *J. Histochem. Cytochem.* (in press).
- TONNA, E.A., SANEN, F.J. AND CRONKITE, E.P. Fracture repair after treatment with deoxyribonucleic acid from fish. *Nature* **195**, 157-9 (1962).
- TONNA, E.A., SHELLABARGER, C.J. AND CRONKITE, E.P. An autoradiographic study of rat femora nine months after administration of tritiated thymidine. (A) *Anat. Record* **142**, 329 (1962).
- TONNA, E.A. - See also COTTIER, H.; HWANG, W.S.
- TSUYA, A., HAYANO, I., BOND, V.P. AND FLIEDNER, T.M. The effects of radiation on total and DNA synthesizing cells of the bone marrow and blood. Submitted to *Intern. J. Radiation Biol.*
- USENIK, E. - See BOND, V.P.; CRONKITE, E.P.; PERMAN, V.
- VAN SLYKE, D.D. Gasometric methods of analysis. Presented at Intern. Symp. Microchem. Techniques, Pennsylvania State University, Aug. 1961.
- VAN SLYKE, D.D. AND PLAZIN, J. Determination of carbon and its radioactivity. IV. Transfer of CO_2 to counting tubes without use of liquid nitrogen. *J. Biol. Chem.* (in press).
- WASSERMAN, L.R. - See PRICE, D.C.
- WOLINS, W. - See FINE, S.; MAXFIELD, M.

Nuclear Engineering Department

- ADLER, G. Radiation-induced solid state polymerization. *Proc. Symp. High Power Accelerator Applications, New York City, Oct. 1961*, pp. 69-78, Radiation Dynamics, Inc., Westbury, N.Y., 1962.
- ADLER, G. - See also KENEALLY, E.; STEINBERG, M.
- ARCIPIANI, B., RICABARRA, D. AND RICABARRA, G.H. A note on the measurement of the U^{238} cadmium ratio. (L) Submitted to *Nucl. Sci. Eng.*
- ARONSON, A. - See MICHAEL, P.; ROBBA, W.A.
- AUERBACH, C., FINSTON, H.L., KISSEL, G. AND GLICKSTEIN, J. An incremental approach to derivative polarography, *Anal. Chem.* 33, 1480-4 (1961).
- AUERBACH, E.H. - See MOORE, S.O.
- AUSKERN, A. AND OSAWA, Y. Xenon diffusion in uranium carbide powder. (N) *J. Nucl. Mater.* (in press).
- AUSTIN, J.C. *Bibliography on Gamma-Ray Radiography, 1950 to 1961 Inclusive, With Subject and Author Indices.* BNL 716 (L-326), March 1962.
- AVRUTIK, E.R. - See DWYER, O.E.
- BALLANTINE, D.S. - See HOIGNE, J.
- BANSLABEN, A.J. AND FINSTON, H.L. An alpha handling system of commercially available components. (A) Presented at Am. Nucl. Soc. Meeting, Washington, Nov. 1962.
- BARTLETT, C.B. - See HORN, F.L.
- BAUMANN, C.G. AND METZ, D.J. Gamma radiation-induced polymerization of alpha-methyl styrene. The effect of water on molecular weight distribution. (N) Submitted to *J. Polymer Sci.*
- BEZLER, P. - See TUCKER, W.
- BLOCH, F.W., MACKENZIE, D.R. AND WISWALL, R.H. JR. Preparation and properties of fluorocarbons of interest in reactor technology. *2nd Conf. Nuclear Reactor Chemistry, Gatlinburg, Tenn., Oct. 1961*, pp. 109-20, TID-7622, July 1962.
- BRACKER, J. - See EGAN, J.J.
- BUHL, P.A. - See KELSCH, J.J.
- CARLSON, C.M. A bond-cell method for complex molecules and metals. Submitted to *J. Chem. Phys.*
- CASTLEMAN, A.W. JR. The chemical and physical behavior of released fission products. (A) Presented at 3rd Conf. Nucl. Reactor Chem., Gatlinburg, Tenn., Oct. 1962.
- CHAPMAN, R.N. - See STEINBERG, M.
- CHEN, J.C. Boiling potassium heat transfer project loop design and development. (A) Presented at US AEC High-Temp. Liquid-Metal Heat Transfer Technol. Meeting, Brookhaven National Laboratory, May 1962.
- CHEN, J.C. Comparison of heat transfer characteristics of fuel elements in particulate and in solid forms. Presented at Am. Nucl. Soc. Meeting, Chicago, Nov. 1961.
- CHERNICK, J., HONECK, H.C., MICHAEL, P., MOORE, S.O. AND SRIKANTIAH, G. The correlation of integral experiments and high energy cross sections. *Physics of Fast and Intermediate Reactors*, Vol. 1, pp. 435-43, IAEA, Vienna, 1962; *Nucl. Sci. Eng.* 13, 205-14 (1962).
- CHERNICK, J. - See also MICHAEL, P.
- CHOW, J.G.Y., HARE, J., NIELSEN, A. AND PALLAS, F. Procedure for disassembling an uranium-bismuth loop in the BNL metallurgy hot cell. *Proc. 9th Hot Lab. Equip. Conf., Chicago, Nov. 1961*, pp. 188-94, American Nuclear Society, Chicago, 1962.
- CHOW, J.G.Y., MCRICKARD, S.B. AND GURINSKY, D.H. Effect of fast neutron irradiation on the mechanical properties of pure iron. *Proc. IAEA Symp. Radiation Damage in Solids and Reactor Materials, Vienna, Oct. 1962* (in press).
- CHOW, J.G.Y., MCRICKARD, S.B. AND GURINSKY, D.H. The mechanical properties of irradiated iron and iron alloys. (A) Presented at ASTM Symp. Radiation Effects on Materials and Neutron Dosimetry, Los Angeles, Sept. 1962.
- COLOMBO, P. - See STEINBERG, M.
- DE VILLIERS, J.W.L. - See STARR, E.
- DOERING, R.F., TUCKER, W.D. AND STANG, L.G. JR. A simple device for milking high-purity Y^{90} from Sr^{90} . *J. Nucl. Med.* (in press).
- DWYER, O.E., Eddy transport in liquid-metal heat transfer. (A) Presented at US AEC High-Temp. Liquid-Metal Heat Transfer Technol. Meeting, Brookhaven National Laboratory, May 1962.
- DWYER, O.E. Equations for bilateral heat transfer to fluids flowing in concentric annuli. Submitted to *Nucl. Sci. Eng.*
- DWYER, O.E. Wall and bulk temperatures for fluids flowing in concentric annuli. Informal Report BNL 6177.
- DWYER, O.E., HOWE, H.E. AND AVRUTIK, E.R. Removal of the more noble nonvolatile fission products from uranium-bismuth fuels. *Nucl. Sci. Eng.* 12, 15-22 (1962).
- DWYER, O.E. AND TU, P.S. Unilateral heat transfer to liquid metals flowing in concentric annuli. *Nucl. Sci. Eng.* (in press).
- EGAN, J.J., MCCOY, W. AND BRACKER, J. The standard molar free energy of formation of $MgCl_2$, $CeCl_3$, $ThCl_4$, and UCl_3 obtained by solid-state emf techniques. *Proc. IAEA Symp. Thermodynamics of Nuclear Materials, Vienna, May 1962* (in press).
- EGAN, J.J. - See also WISWALL, R.H. JR.
- EPSTEIN, S.G. - See WEEKS, J.R.
- ESHAYA, A.M. - See KENNEY, W.F.; SALZANO, F.J.
- FELBERBAUM, J. - See SHER, R.
- FINSTON, H.L. AND YELLIN, E. Determination of carbon-hydrogen ratios by neutron scattering. Submitted to *Anal. Chem.*
- FINSTON, H.L. - See also AUERBACH, C.; BANSLABEN, A.J.; JORDAN, J.
- FLEITMAN, A.H., ROMANO, A.J. AND KLAMUT, C.J. Corrosion of carbon steel by high temperature mercury. (A) Presented at Ann. Meeting Electrochem. Soc., Boston, Sept. 1962.
- FLEITMAN, A.H., ROMANO, A.J. AND KLAMUT, C.J. Corrosion of low carbon steel in boiling mercury. (A) Presented at AIME Symp. Corrosion by Metal Heat Transfer Fluids, New York City, Jan. 1962.
- GALANTER, L. - See KUHLE, O.A.; MULLER, A.; RIZZO, F.X.
- GARD, J. - See KENEALLY, E.
- GLICKSTEIN, J. - See AUERBACH, C.
- GREEN, L. Table of reactor coolant properties. *Nucleonics* 19, No. 11, 140-4 (1961).
- GREEN, L. - See also HATCH, L.P.; ROBBA, W.A.

- GREENE, M.W. AND TUCKER, W.D. An improved gallium-68 cow. *Intern. J. Appl. Radiation Isotopes* 12, 62-3 (1961).
- GREENE, M.W. - See also STERNLIEB, I.; TUCKER, W.D.
- GURINSKY, D.H. - See CHOW, J.G.Y.; ROMANO, A.J.; SCHWEITZER, D.G.
- HARE, J. - See CHOW, J.G.Y.
- HATCH, L.P., ROBBA, W.A., POWELL, J.R., REGAN, W.H., GREEN, L. AND HOFFMAN, K.C. Settled bed fuel reactors. Presented at Natl. Meeting Am. Inst. Chem. Engrs., Los Angeles, Feb. 1962.
- HATCH, L.P., WETH, G.C. AND TUTHILL, E.J. Ultimate disposal of high level radioactive wastes. Fixation in phosphate glass with emphasis on the continuous mode of plant operation. *Proc. IAEA Symp. Treatment and Storage of High Level Radioactive Wastes, Vienna, Oct. 1962* (in press).
- HATCH, L.P. - See also REGAN, W.H.; REILLY, J.J.; ROBBA, W.A.
- HELLSTEN, A. - See SHER, R.
- HILLMAN, M. Decay of Sc^{48} . (A) *Bull. Am. Phys. Soc.* 7, 462 (1962).
- HILLMAN, M. Formation cross sections for Ca^{47} using 14.5-Mev neutrons. *Nucl. Phys.* (in press).
- HOFFMAN, K.C. - See HATCH, L.P.; ROBBA, W.A.
- HOIGNE, J. AND BALLANTINE, D. Irradiation-induced effects in magnesium oxide catalysts. Submitted to *Intern. J. Appl. Radiation Isotopes*.
- HONECK, H.C. AND TAKAHASHI, H. The angular dependence of thermal neutron spectra in lattices. Submitted to *Nucl. Sci. Eng.*
- HONECK, H.C. - See also CHERNICK, J.
- HORN, F.L., BARTLETT, C.B. AND TIVERS, R.J. Chemistry of anhydrous systems using hydrogen fluoride and nitrogen oxides. *Proc. 2nd Intern. Symp. Fluorine Chem., Estes Park, Colo., July 1962* (in press).
- HORN, F.L. - See also JOHNSON, R.
- HOWE, H.E. - See DWYER, O.E.
- HRABAK, G.C. - See SCHWEITZER, D.G.
- JOHNSON, R., HORN, F.L. AND STRICKLAND, G. The "Phosphor" process for reprocessing ceramic fuels. Presented at Am. Nucl. Soc. Meeting, Boston, June 1962.
- JORDAN, J. AND FINSTON, H.L. Identification of active electrode sites by radioactive tracers. *J. Chim. Phys.* (in press).
- KAMMERER, O.F. - See KELSCH, J.J.
- KATZ, H.M. *Chemical Decladding of Stainless Steel Fuel Elements in a Stainless Steel Dissolver, Progress Report No. 1.* BNL 749 (T-275), July 1962.
- KATZ, H.M. *Some Methods of Preparing Tetravalent Uranium Solutions.* BNL 691 (T-236), Sept. 1961.
- KELSCH, J.J., GOLAND, A.N. AND KAMMERER, O.F. Observation of the $\text{Li}^6(n,\alpha)\text{H}^3$ reaction by electron microscopy of Li^6F films. (L) Submitted to *Appl. Phys. Letters*.
- KELSCH, J.J., KAMMERER, O.F. AND GOLAND, A.N. Electron microscopy of the effects of the passage of energetic particles through solids. (A) Presented at 5th Intern. Congr. Electron Microscopy, Philadelphia, Sept. 1962.
- KELSCH, J.J., KAMMERER, O.F., GOLAND, A.N. AND BUHL, P.A. Electron microscopy of metals exposed to fission fragments. (A) *Bull. Am. Phys. Soc.* 7, 172 (1962).
- KELSCH, J.J., KAMMERER, O.F., GOLAND, A.N. AND BUHL, P.A. Observation of fission fragment damage in thin films of metals. *J. Appl. Phys.* 33, 1475-82 (1962).
- KENEALLY, E., GARD, J. AND ADLER, G. The temperature dependence of birefringence in irradiated polyethylene. Submitted to *J. Polymer Sci.*
- KENNY, W.F. AND ESHAYA, A.M. *Adsorption of Xenon on Activated Charcoal.* BNL 689 (T-235), Jan. 1962.
- KISSEL, G. - See AUERBACH, C.
- KLAMUT, C.J. - See FLEITMAN, A.H.; ROMANO, A.J.; WEEKS, J.R.
- KLOTZ, P. - See NEWMAN, L.
- KOPPEL, J.U. Time dependent, space independent neutron thermalization. *Nucl. Sci. Eng.* 12, 532-41 (1962).
- KOUTS, H. The use of small reactors for reactor physics. *Proc. IAEA Symp. Programming and Utilization of Res. Reactors, Vienna, Oct. 1961* (in press).
- KRAMER, H., PRUZANSKY, J. AND STEINBERG, M. Formation of hydrazine from the radiolysis of ammonia in the presence of mineral solids. *Intern. J. Appl. Radiation Isotopes* (in press).
- KUHL, O.A. A high intensity radiation development laboratory at Brookhaven National Laboratory. Presented at Am. Nucl. Soc. Meeting, Chicago, Nov. 1961.
- KUHL, O.A., GALANTER, L., O'LEARY, V. AND MANOWITZ, B. *Conceptual Design Studies on Facilities for Radiation Pasteurization of Food, IN-267, Reports and Industrial News Branch, Division of Public Information, US AEC, Washington, Oct. 1961.*
- LELLOUCHE, G.S. Reactor size sufficient for stability against spatial xenon oscillations. (L) *Nucl. Sci. Eng.* 13, 60-1 (1962).
- LELLOUCHE, G.S. Space-dependent xenon oscillations. *Nucl. Sci. Eng.* 12, 482-9 (1962).
- LEVINE, M.M. Fission-product burn-up in fast reactors. *Physics of Fast and Intermediate Reactors, Vol. 2, p. 247, IAEA, Vienna, 1962.*
- LEVINE, M.M. AND STEINBERG, M. Design of radiochemical reactors with special reference to the polymerization of ethylene. *Nucl. Sci. Eng.* 12, 498-504 (1962).
- LEVINE, M.M. - See also ROBBA, W.A.
- LIU, C.H. Electroanalytical techniques in molten lithium sulfate-potassium sulfate eutectic. Submitted to *Anal. Chem.*
- LIU, C.H. Electrode potentials in molten lithium sulfate-potassium sulfate eutectic. *J. Phys. Chem.* 66, 164 (1962).
- LOEFFELHOLZ, M. - See STEINBERG, M.
- MACKENZIE, D.R. - See BLOCH, F.W.
- MANOWITZ, B. Applications of radiation to chemical processes. Presented at Am. Inst. Chem. Engrs. Symp., New York City, Oct. 1961.
- MANOWITZ, B. Let's be realistic about fuel reprocessing costs. *Nucleonics* 20, No. 2, 60-4 (1962).
- MANOWITZ, B. - See also KUHL, O.A.
- MASLAN, F. Xenon thermodynamic diagram. *Nucleonics* 20, No. 3, 53 (1962).
- MCCOY, W. - See EGAN, J.J.
- McNICHOLAS, J. - See ROBBA, W.A.
- McRICKARD, S.B. - See CHOW, J.G.Y.
- METZ, D.J. - See BAUMANN, C.G.

- MEYER, R.A. AND SCHWEITZER, D.G. Postirradiation measurements of thermal conductivity between 200 and 600°C for low permeability graphites. *Proc. 5th Conf. Carbon, Pennsylvania State University, June 1961*, Pergamon (in press).
- MICHAEL, P., CHERNICK, J. AND ARONSON, A. Theoretical analysis of the Snell experiment. Presented at Am. Nucl. Soc. Meeting, Chicago, Nov. 1961.
- MICHAEL, P. - See also CHERNICK, J.
- MOORE, S.O. *Neutron Cross Section Evaluation Group Newsletter No. 6*. BNL 732 (N-6), June 1962.
- MOORE, S.O., AUERBACH, E.H. AND PORTER, C.E. Analysis of U^{238} inelastic scattering, 0-1 Mev. (A) *Bull. Am. Phys. Soc.* 7, 357 (1962).
- MOORE, S.O. - See also CHERNICK, J.; SHER, R.
- MORELL, A.G. - See STERNLIEB, I.
- MULLER, A., RIZZO, F. AND GALANTER, L. *Solar Cell Dosimeter, Report No. 1*. BNL 721 (T-255), Feb. 1962.
- MURRENHOFF, A.P. - See TUCKER, W.D.
- NEWMAN, L. On the synergistic effect in solvent extraction. (N) Submitted to *J. Inorg. Nucl. Chem.*
- NEWMAN, L. AND KLOTZ, P. A secondary interaction of tri-*n*-octylamine with hydrochloric acid and thenoyltrifluoroacetone. Submitted to *J. Phys. Chem.*
- NEWMAN, L. AND KLOTZ, P. The solvent extraction of thorium by thenoyltrifluoroacetone and the influence of aqueous complexation by the chelating agent. Submitted to *J. Phys. Chem.*
- NEWMAN, L. AND KLOTZ, P. Synergistic effect of tri-*n*-octylamine on the solvent extraction of thorium by thenoyltrifluoroacetone. Submitted to *J. Am. Chem. Soc.*
- NIELSEN, A. - See CHOW, J.G.Y.
- O'LEARY, V. - See KUHL, O.A.
- OSAWA, Y. - See AUSKERN, A.
- PALLAS, F. - See CHOW, J.G.Y.
- PANCER, G. - See ROBBA, W.A.
- PORTER, C.E. Further remarks on energy level spacings. (A) *Bull. Am. Phys. Soc.* 7, 348 (1962).
- PORTER, C.E. - See also MOORE, S.O.
- POWELL, J.R. - See HATCH, L.P.
- PRICE, G.A. Measurements of reactor parameters in subcritical and critical assemblies: Lattices of slightly enriched uranium in water. (A) Presented at Am. Nucl. Soc. Meeting, Boston, June 1962.
- PRICE, G.A. - See also STARR, E.
- PRUZANSKY, J. - See KRAMER, H.
- PUROHIT, S.N. AND RAJAGOPAL, A.K. Scattering of thermal neutrons in the Doppler approximation. *Nucl. Sci. Eng.* 13, 250-60 (1962).
- RAJAGOPAL, A.K. - See PUROHIT, S.N.
- RAJAGOPAL, V. Determination of reactor transfer functions by statistical correlation methods. *Nucl. Sci. Eng.* 12, 218-24 (1962).
- REGAN, W.H., REILLY, J.J., WIRSING, E. AND HATCH, L.P. The application of inert fluidized beds to the volatility reprocessing of reactor fuels. Presented at Am. Inst. Chem. Engrs. Meeting, New York City, Dec. 1961.
- REGAN, W.H. - See also HATCH, L.P.; REILLY, J.J.
- REILLY, J.J., REGAN, W.H., WIRSING, E. AND HATCH, L.P. Studies on uranium recovery from unirradiated reactor fuel elements by volatile separation in inert fluidized beds. *Ind. Eng. Chem.* (in press).
- REILLY, J.J. - See also REGAN, W.H.
- RICABARRA, D. - See ARCIPIANI, B.
- RICABARRA, G.H. - See ARCIPIANI, B.
- RIZZO, F.X. AND GALANTER, L. Experimental build-up factors for semi-infinite water target and cobalt-60 slab sources. (A) Presented at Am. Nucl. Soc. Meeting, Chicago, Nov. 1961.
- RIZZO, F.X. - See also MULLER, A.
- ROBBA, W.A., ARONSON, A., GREEN, L., HATCH, L.P., HOFFMAN, K.C., LEVINE, M., McNICHOLAS, J. AND PANCER, G. *The Settled Bed Fast Reactor, Status Report*. BNL 713 (T-250), Jan. 1962.
- ROBBA, W.A. - See also HATCH, L.P.
- ROMANO, A.J., KLAMUT, C.J. AND GURINSKY, D.H. The investigation of container materials for Bi and Pb alloys. I. Thermal convection loops. (A) Presented at Ann. Meeting Electrochem. Soc., Boston, Sept. 1962.
- ROMANO, A.J. - See also FLEITMAN, A.H.
- SALZANO, F.J. Summary of fission product adsorption work at BNL. *7th AEC Air Cleaning Conf., Brookhaven National Laboratory, Oct. 1961*, pp. 224-41, TID-7627, March 1962.
- SALZANO, F.J. AND ESHAYA, A.M. Sorption of xenon in high density graphite at high temperatures. *Nucl. Sci. Eng.* 12, 1-3 (1962).
- SCHEINBERG, I.H. - See STERNLIEB, I.
- SCHWEITZER, D.G. The activation energy for annealing single interstitials in neutron-irradiated graphite and the absolute rate of formation of displaced atoms. *Phys. Rev.* (in press).
- SCHWEITZER, D.G. Oxidation and heat transfer studies in graphite channels. IV. Combined effects of temperature, flow rate, diameter, and reactivity on the length of channel cooled by air. *Nucl. Sci. Eng.* 12, 59-62 (1962).
- SCHWEITZER, D.G. Thermal properties of air-cooled graphite channels. *Nucl. Sci. Eng.* 13, 275-82 (1962).
- SCHWEITZER, D.G. AND GURINSKY, D.H. Oxidation and heat transfer studies in graphite channels. II. Thermal changes with time and distance in air-cooled graphite channels. *Nucl. Sci. Eng.* 12, 46-50 (1962).
- SCHWEITZER, D.G., HRABAK, G.C. AND SINGER, R.M. Oxidation and heat transfer studies in graphite channels. I. The effect of air flow rate on the C-O₂ and CO-O₂ reactions. *Nucl. Sci. Eng.* 12, 39-45 (1962).
- SCHWEITZER, D.G. AND SINGER, R.M. Oxidation and heat transfer studies in graphite channels. III. The reactivity of BNL graphite and its effect on the length of channel cooled by air. *Nucl. Sci. Eng.* 12, 51-8 (1962).
- SCHWEITZER, D.G. - See also MEYER, R.A.
- SEARS, J. AND STEINBERG, M. *Fission Fragment Energy Deposition for Formation of Chemical Products in Two-Phase Systems*. BNL 695 (T-238), Sept. 1961.
- SHER, R. AND FELBERBAUM, J. *Neutron Cross Section Evaluation Group, Least-Squares Analysis of the 2200-m/sec Parameters of U^{233} , U^{235} , and Pu^{239}* . BNL 722 (T-256-92-94-1), June 1962.
- SHER, R. AND MOORE, S. *Neutron Cross Section Evaluation Group Newsletter No. 5*. BNL 694 (N-5), Oct. 1961.
- SHER, R., TASSAN, S., WEINSTOCK, E.V. AND HELLSTEN, A. Low energy neutron cross sections of Dy¹⁶⁴. *Nucl. Sci. Eng.* 11, 369-76 (1961).

- SINGER, R.M. - See SCHWEITZER, D.G.
- SRIKANTIAH, G. Analysis of a beryllium-moderated critical experiment. (A) Presented at Am. Nucl. Soc. Meeting, Chicago, Nov. 1961.
- SRIKANTIAH, G. - See also CHERNICK, J.
- STANG, L.G. JR. Production of "special" radioisotopes with particular reference to the less developed countries. (A) Presented at IAEA Symp. Practical Applications of Short-Lived Radioisotopes Produced in Small Research Reactors, Vienna, Nov. 1962.
- STANG, L.G. JR. - See also DOERING, R.F.
- STARR, E. AND DE VILLIERS, J.W.L. Determination of diffusion cooling in graphite by measurement of the average neutron velocity. Submitted to *Nucl. Sci. Eng.*
- STARR, E. AND PRICE, G.A. Measurement of the diffusion parameters of graphite and graphite-bismuth by pulsed neutron methods. (A) Presented at Am. Nucl. Soc. Meeting, Boston, June 1962.
- STEINBERG, M. Chemonuclear processing. (A) Presented at Am. Nucl. Soc. Meeting, Boston, June 1962.
- STEINBERG, M. Fission fragment energy deposition efficiency. Submitted to *Nucleonics*.
- STEINBERG, M., COLOMBO, P., CHAPMAN, R.N. AND ADLER, G. Polymerization of ethylene by cobalt-60 gamma radiation. (A) Presented at 2nd Intern. Congr. Radiation Res., Harrogate, England, Aug. 1962.
- STEINBERG, M., COLOMBO, P., CHAPMAN, R.N. AND ADLER, G. *Radiation Processing, Report No. 4, Polymerization of Ethylene by Cobalt-60 Gamma Radiation*. BNL 740 (T-268), July 1962.
- STEINBERG, M. AND LOEFFELHOLZ, M. Energy deposition efficiency of fission fragment recoil radiation from thin uranium-containing elements. (A) Presented at 2nd Intern. Congr. Radiation Res., Harrogate, England, Aug. 1962.
- STEINBERG, M. - See also KRAMER, H.; LEVINE, M.M.; SEARS, J.; TUCKER, W.D.
- STERNLIEB, I., MORELL, A.G., TUCKER, W.D., GREENE, M.W. AND SCHEINBERG, I. The incorporation of copper into ceruloplasmin *in vivo*: Studies with copper-64 and copper-67. *J. Clin. Invest.* **40**, 1834-40 (1961).
- STRICKLAND, G. - See JOHNSON, R.
- TAKAHASHI, H. Generalization of the singular integral method for an anisotropic scattering medium. *J. Nucl. Energy Pt. A* **15**, 192-5 (1961).
- TAKAHASHI, H. - See also HONECK, H.C.
- TASSAN, S. Conversion ratios in slightly enriched uranium, water-moderated lattices. (L) Submitted to *Nucl. Sci. Eng.*
- TASSAN, S. - See also SHER, R.
- TIVERS, R.J. - See HORN, F.L.
- TU, P.S. - See DWYER, O.E.
- TUCKER, W.D., GREENE, M.W. AND MURRENHOF, A.P. Production of carrier-free Tc^{132} , I^{132} , Mo^{99} , and Tc^{99m} from neutron-irradiated uranium by fractionated sorption on alumina. *Atompraxis* **5**, 163-7 (1962).
- TUCKER, W.D., WAIDE, C., BEZLER, P. AND STEINBERG, M. The Brookhaven chemonuclear in-pile test loop. (A) Presented at Am. Nucl. Soc. Meeting, Boston, June 1962.
- TUCKER, W.D. - See also DOERING, R.F.; GREENE, M.W.; STERNLIEB, I.
- TUTHILL, E.J. - See HATCH, L.P.
- VAN NORMAN, J.D. Controlled potential coulometry in fused lithium chloride-potassium chloride eutectic. *Anal. Chem.* **34**, 594-6 (1962).
- WAIDE, C. - See TUCKER, W.D.
- WEEKS, J.R. Corrosion and mass transfer in alkali-liquid metal systems. In *Proc. 1962 Golden Gate Metals Conf. Materials, Science, and Technology for Advanced Applications*, D.R. Mash, Editor, Prentice-Hall, New York (in press).
- WEEKS, J.R. Corrosion in liquid metal cooled reactors. (A) Presented at Natl. Assoc. Corrosion Engrs. Meeting, Albany, N.Y., Oct. 1962.
- WEEKS, J.R. Possible correlations between thermoelectric potentials and liquid metal corrosion. Submitted to *Japan. Electrochem. Soc. J.*
- WEEKS, J.R. Reaction kinetics and thermodynamics in liquid metal corrosion. (A) Presented at AIME Symp. Corrosion by Metal Heat Transfer Fluids, New York City, Feb. 1962.
- WEEKS, J.R. AND EPSTEIN, S.G. Possible correlations between thermoelectric potentials and liquid metal corrosion. (A) Presented at Ann. Meeting Electrochem. Soc., Boston, Sept. 1962.
- WEEKS, J.R. AND KLAMUT, C.J. Liquid metal corrosion mechanisms. *Proc. IAEA Conf. Corrosion of Reactor Materials, Salzburg, Austria, June 1962* (in press).
- WEINSTOCK, E.V. - See SHER, R.
- WEISS, J. Radiolysis of polyaromatic compounds. (A) Presented at 2nd Intern. Congr. Radiation Res., Harrogate, England, Aug. 1962.
- WEISS, J. Radiolysis of toluene. (A) Presented at 2nd Intern. Congr. Radiation Res., Harrogate, England, Aug. 1962.
- WETH, G.C. - See HATCH, L.P.
- WIRSING, E. - See REGAN, W.H.; REILLY, J.J.
- WISWALL, R.H. JR. AND EGAN, J.J. Thermodynamics of solutions of actinides and fission products in bismuth. *Proc. IAEA Symp. Thermodynamics of Nuclear Materials, Vienna, May 1962* (in press).
- WISWALL, R.H. JR. - See also BLOCH, F.W.
- YELLIN, E. - See FINSTON, H.L.

Physics Department

- ADAIR, R.K. - See CRITTENDEN, R.R.
- ALBURGER, D.E. Gamma-ray decay of the 7.66-Mev level of C^{12} . *Phys. Rev.* **124**, 193-8 (1961).
- ALFRED, L.C.R. Shells and electron groups in the Thomas-Fermi atom. *Phys. Rev.* **125**, 214-19 (1962).
- ALPERIN, H.A., BROWN, P.J., NATHANS, R. AND PICKART, S.J. A polarized neutron study of antiferromagnetic domains in MnF_2 . (L) *Phys. Rev. Letters* **8**, 237-9 (1962).
- ALPERIN, H.A. - See also BROWN, P.J.; NATHANS, R.
- ARTMAN, R.A. AND GOLAND, A.N. Neutron irradiation of bismuth telluride. (A) *Bull. Am. Phys. Soc.* **7**, 186 (1962).
- AYER, F. AND BOLZE, E.M. A versatile bubble chamber track-measuring stage. Submitted to *Rev. Sci. Instr.*
- AYER, F., PICKUP, E., RAPPE, E.F. AND WALL, G. A bubble chamber track-centering device with digitized output. Submitted to *Rev. Sci. Instr.*
- BACHMAN, A.H. - See SAMIOS, N.P.

- BAKER, C.P., CURTIS, H. J., ZEMAN, W. AND WOODLEY, R.G. The design and calibration of a deuteron micro-beam for biological studies. *Radiation Res.* **15**, 489-95 (1961).
- BAKER, C.P. - See also DONOVAN, P.F.; ZEMAN, W. (Biology).
- BAKER, W.F. A computer program to optimize magnets in a beam transport system. Submitted to *Nucl. Instr. Methods* (special issue).
- BAKER, W.F., COOL, R.L., JENKINS, E.W., KYCIA, T.F., LINDENBAUM, S. J., LOVE, W.A., LUERS, D., NIEDERER, J.A., OZAKI, S., READ, A.L., RUSSELL, J. J. AND YUAN, L.C.L. Particle production by 10 to 30-Bev protons incident on Al and Be. (L) *Phys. Rev. Letters* (in press).
- BALTAY, C., SANDWEISS, J., SANFORD, J., BROWN, H., WEBSTER, M. AND YAMAMOTO, S. The separated beam at the AGS: Performance with antiprotons and π^+ -mesons. *Proc. CERN Intern. Conf. High Energy Physics, Geneva, July 1962* (in press).
- BALTAY, C. - See also BROWN, H.
- BARTON, G. AND KACSER, C. Some analytic properties of a decay amplitude with final state interactions: The single-loop four-point diagrams in $K \rightarrow 3\pi$ decay. *Nuovo Cimento* **21**, 988 (1961).
- BÉG, M.A.B. Correspondence between τ and η decays. (L) *Phys. Rev. Letters* **9**, 67-9 (1962).
- BÉG, M.A.B. AND DECELLES, P.C. Current conservation and $|\Delta T| = \frac{1}{2}$ rule in θ -decay. *Phys. Rev.* (in press).
- BÉG, M.A.B. AND DECELLES, P.C. Heavy pions and the πN vertex function. (L) *Phys. Rev. Letters* **9**, 30-1 (1962).
- BÉG, M.A.B. AND DECELLES, P.C. P -wave interactions in τ decay. (L) *Phys. Rev. Letters* **8**, 46-7 (1962).
- BÉG, M.A.B., DECELLES, P.C. AND MARR, R.B. Decay modes of K^* . *Phys. Rev.* **124**, 622-3 (1961).
- BERNABEI, A., BORST, L.B. AND SAILOR, V.L. Neutron resonance in samarium. *Nucl. Sci. Eng.* **12**, 63-7 (1962).
- BERNSTEIN, J. AND FEINBERG, G. Electromagnetic mixing effects in elementary particle physics. Submitted to *Nuovo Cimento*.
- BERNSTEIN, J. AND FEINBERG, G. A needle in a haystack. *Proc. CERN Intern. Conf. High Energy Physics, Geneva, July 1962* (in press).
- BERNSTEIN, J. AND FEINBERG, G. On the production and decay of intermediate vector mesons. *Phys. Rev.* **125**, 1741-5 (1962).
- BERTANZA, L., BRISSON, V., CONNOLLY, P., HART, E., HOUGH, P., MITTRA, I., MONETI, G., RAU, R., SAMIOS, N.P., SKILLICORN, I., YAMAMOTO, S., GOLDBERG, M., HARTH, E., LEITNER, J., LICHTMAN, S. AND WESTGARD, J. K^-p interactions at 2.3 BeV/c. (A) *Bull. Am. Phys. Soc.* **7**, 296 (1962).
- BERTANZA, L., BRISSON, V., CONNOLLY, P.L., HART, E.L., MITTRA, I.S., MONETI, G.C., RAU, R.R., SAMIOS, N.P., SKILLICORN, I.O., YAMAMOTO, S.S., GOLDBERG, M., GRAY, L., LEITNER, J., LICHTMAN, S. AND WESTGARD, J. K^-p interactions at 2.24 BeV/c. I. Effective mass distributions; II. Production properties. *Proc. CERN Intern. Conf. High Energy Physics, Geneva, July 1962* (in press).
- BERTANZA, L., BRISSON, V., CONNOLLY, P.L., HART, E.L., MITTRA, I.S., MONETI, G.C., RAU, R.R., SAMIOS, N.P., SKILLICORN, I.O., YAMAMOTO, S.S., GOLDBERG, M., GRAY, L., LEITNER, J., LICHTMAN, S. AND WESTGARD, J. Properties of the Ξ^- -hyperons. *Proc. CERN Intern. Conf. High Energy Physics, Geneva, July 1962* (in press).
- BERTANZA, L., BRISSON, V., CONNOLLY, P.L., HART, E.L., MITTRA, I.S., MONETI, G.C., RAU, R.R., SAMIOS, N.P., SKILLICORN, I.O., YAMAMOTO, S.S., LEITNER, J. AND WESTGARD, J. The decay of a negative hyperon into $\Lambda^0 + e^- + \bar{\nu}$. (L) *Phys. Rev. Letters* **9**, 19-21 (1962).
- BERTANZA, L., CONNOLLY, P.L., CULWICK, B.B., EISLER, F.R., MORRIS, T.W., PALMER, R., PRODELL, A. AND SAMIOS, N.P. $\Lambda^0 K^0$ production by pions on protons. (L) *Phys. Rev. Letters* **8**, 332-5 (1962).
- BERTANZA, L., CULWICK, B.B., MITTRA, I.S., SAMIOS, N.P., THORNDIKE, A.M., YAMAMOTO, S.S. AND LEA, R.M. Production of strange particles by 5 BeV/c π -mesons. (A) *Bull. Am. Phys. Soc.* **7**, 49 (1962).
- BOLOTIN, H.H., LI, A.C. AND SCHWARZSCHILD, A. Level structure of Sn^{118} and Sn^{120} from the decay of Sb isotopes. *Phys. Rev.* **124**, 213-23 (1961).
- BOLOTIN, H.H. - See also CHRIEN, R.E.
- BOLZE, E.M. - See AYER, F.
- BORST, L.B. - See BERNABEI, A.
- BOUTIN, H., SAFFORD, G. AND BRAJOVIC, V. A study of the liquid and solid phases of HF, HCl, and HBr by neutron inelastic scattering. (A) *Bull. Am. Phys. Soc.* **7**, 500 (1962).
- BOUTIN, H. - See also BRAJOVIC, V.; SAFFORD, G.
- BRAJOVIC, V., BOUTIN, H., SAFFORD, G. AND PALEVSKY, H. A study of rotational freedom in several ammonium salts by slow neutron inelastic scattering. (A) *Bull. Am. Phys. Soc.* **7**, 500 (1962).
- BRAJOVIC, V. AND PALEVSKY, H. Scattering of cold neutrons by crystalline water. (A) *Bull. Am. Phys. Soc.* **7**, 356 (1962).
- BRAJOVIC, V. - See also BOUTIN, H.; SAFFORD, G.
- BRISSON, V. - See BERTANZA, L.
- BROWN, H., CULWICK, B., FOWLER, W., GAILLOUD, M., KALOGEROPOULOS, T., KOPP, J., LEA, R., LOUITT, R., MORRIS, T., SHUTT, R., THORNDIKE, A., WEBSTER, M., BALTAY, C., FOWLER, E., SANDWEISS, J., SANFORD, J. AND TAFT, H. Observation of production of a $\Xi^- + \Xi^+$ pair. (L) *Phys. Rev. Letters* **8**, 255-7 (1962).
- BROWN, H. - See also BALTAY, C.
- BROWN, P. J., NATHANS, R., ALPERIN, H.A. AND PICKART, S. J. The effects of magnetic fields on the antiferromagnetic domain patterns in MnF_2 . (A) *Bull. Am. Phys. Soc.* **7**, 278 (1962).
- BROWN, P. J. - See also ALPERIN, H.A.; FRAZER, B.C.; KOCHER, C.W.
- BRUNHART, G., MARSHAK, H., REYNOLDS, C.A., SAILOR, V.L., SCHERMER, R.I. AND SHORE, F. J. Spin assignments of low energy neutron resonances in erbium and dysprosium. (A) *Bull. Am. Phys. Soc.* **7**, 305 (1962).
- CHADWICK, G.B. - See DUKE, P. J.; FUJII, T.; HIEN, N.C.; KEMP, M.A.R.
- CHRIEN, R.E., BOLOTIN, H.H. AND PALEVSKY, H. Resonance capture γ rays from platinum. *Phys. Rev.* **127**, 1680-7 (1962).
- CHRIEN, R.E., JAIN, A.P. AND PALEVSKY, H. Anomalous neutron scattering of Ar^{36} . *Phys. Rev.* **125**, 275-6 (1962).
- CHRIEN, R.E. - See also JAIN, A.
- CHURCH, E.L. AND WENESER, J. Penetration effects in magnetic-dipole L -shell internal conversion. (A) *Bull. Am. Phys. Soc.* **7**, 490 (1962).

- COHEN, V.W., MORAN, T. AND PENSELIN, S. Hyperfine structure of Cs^{134m} . *Phys. Rev.* **127**, 517-23 (1962).
- COHEN, V.W. - See also PENSELIN, S.
- COLLINS, G.B. - See DUKE, P.J.; FUJII, T.; HIEN, N.C.; KEMP, M.A.R.
- CONNOLLY, P. - See BERTANZA, L.
- COOL, R.L., JENKINS, E.W., KYCIA, T.F., HILL, D.A., MARSHALL, L. AND SCHLUTER, R.A. A measurement of the magnetic moment of the Λ^0 -hyperon. *Phys. Rev.* (in press).
- COOL, R.L. - See also BAKER, W.F.
- COTTER, M.J., MUETHER, H.R. AND PALEVSKY, H. Inelastic scattering of cold neutrons by liquid lead. (A) *Bull. Am. Phys. Soc.* **7**, 24 (1962).
- CRASEMANN, B., EMERY, G.T. AND KANE, W.R. A new isotope: 23-hour Re^{189} . (A) *Bull. Am. Phys. Soc.* **7**, 353 (1962).
- CRITTENDEN, R.R., LEIPUNER, L.B., ADAIR, R.K., MUSGRAVE, B. AND MARTIN, H.J. $\pi + p$ Interactions in the neighborhood of the Λ threshold. (A) *Bull. Am. Phys. Soc.* **7**, 281 (1962).
- CULWICK, B., GAILLOUD, M., MORRIS, T., THORNDIKE, A., LEA, R., KALOGEROPOULOS, T. AND FERBEL, T. Anti-proton elastic scattering. (A) *Bull. Am. Phys. Soc.* **7**, 298 (1962).
- CULWICK, B. - See also BERTANZA, L.; BROWN, H.
- DAMASK, A.C. Effects of neutron irradiation on nonfissionable alloys. Presented at IAEA Symp. Radiation Damage in Solids and Reactor Material, Venice, May 1962.
- DAMASK, A.C. Radiation and atomic rearrangement in alloys. Submitted to *Nucleonics*.
- DAMASK, A.C. AND DIENES, G.J. Kinetics of vacancy-interstitial annihilation. I. Impurity-interstitial trapping. *Phys. Rev.* **125**, 444-6 (1962).
- DAMASK, A.C. AND DIENES, G.J. Theory of point defect annealing in metals. (A) Presented at Intern. Conf. Crystal Lattice Defects, Kyoto, Japan, Sept. 1962.
- DAMASK, A.C. - See also DIENES, G.J.; FUJITA, F.E.; WAGENBLAST, H.
- DECELLES, P.C. - See BÉG, M.A.B.; DURAND, L. III.
- DIENES, G.J. AND DAMASK, A.C. Kinetics of vacancy-interstitial annihilation. II. Di-interstitial formation. *Phys. Rev.* **125**, 447-50 (1962).
- DIENES, G.J. - See also DAMASK, A.C.; HATCHER, R.D.
- DONOVAN, P.F., KANE, J.V., BAKER, C.P., MOLLENAUER, J.F. AND ZUPANCIC, C. Interactions in multibody nuclear disintegration processes. (A) *Bull. Am. Phys. Soc.* **7**, 287 (1962).
- DONOVAN, P.F. - See also ZUPANCIC, C.
- DUKE, P.J., CHADWICK, G.B., COLLINS, G.B., FUJII, T., HIEN, N.C. AND TURKOT, F. Proton-proton collisions and the one-pion exchange model. (A) Presented at Intern. Conf. Elementary Particles, Aix-en-Provence, France, Sept. 1961.
- DUKE, P.J. - See also FUJII, T.; HIEN, N.C.; KEMP, M.A.R.
- DURAND, L. III, DECELLES, P.C. AND MARR, R.B. Lorentz invariance and the kinematic structure of vertex functions. *Phys. Rev.* **126**, 1882-98 (1962).
- EISLER, F.R. - See BERTANZA, L.
- EL-WAHAB, M.A. AND KANE, J.V. An exact analytic treatment for scintillation counter time resolution functions. *Nucl. Instr. Methods* **15**, 15-22 (1962).
- EMERY, G.T., FIEBIGER, N.F. AND KANE, W.R. Evidence for an octopole excitation in U^{239} . (A) *Bull. Am. Phys. Soc.* **7**, 11 (1962).
- EMERY, G.T., KANE, W.R., MCKEOWN, M., PERLMAN, M.L. AND GOLDBABER, G.S. Properties of collective states in osmium-186. Presented at Conf. Magnetic Lifetimes and Properties of Nuclear States, Gatlinburg, Tenn., Oct. 1961.
- EMERY, G.T. - See also CRASEMANN, B.; SCHMORAK, M.
- FAZIO, G.G. - See MELISSINOS, A.C.
- FEINBERG, G. - See BERNSTEIN, J.
- FERBEL, T. - See CULWICK, B.
- FICKINGER, W.J., PICKUP, E., ROBINSON, D.K. AND SALANT, E.O. p - p Interactions at 2 Bev. I. Single-pion production. *Phys. Rev.* **125**, 2082-90 (1962).
- FIEBIGER, N.F. γ - γ Cascades from the reaction $\text{U}^{238}(n, \gamma)\text{U}^{239}$. (A) *Bull. Am. Phys. Soc.* **7**, 11 (1962).
- FIEBIGER, N.F., KANE, W.R. AND SEGEL, R.E. Thermal neutron capture gamma rays in iron. *Phys. Rev.* **125**, 2031-7 (1962).
- FIEBIGER, N.F. - See also EMERY, G.T.; KANE, W.R.; OLNESS, J.W.
- FIREMAN, E.L. AND ROWLAND, R.S. An additional measurement of the tritium content of atmospheric hydrogen of 1949. *J. Geophys. Res.* **66**, 4321 (1961).
- FLANAGAN, T.B. The effects of reactor irradiation upon the subsequent thermal decomposition of lead stypnate. *J. Phys. Chem.* **66**, 416-21 (1962).
- FLANAGAN, T.B. AND KIM, C.H. The effect of reactor irradiation upon the kinetics of an endothermic solid reaction. The dehydration of manganous oxalate dihydrate. *J. Phys. Chem.* **66**, 926-30 (1962).
- FOWLER, E. - See BROWN, H.
- FOWLER, W. - See BROWN, H.
- FOX, J.D. - See KANE, W.R.
- FRAZER, B.C. Neutron diffraction studies of ferroelectrics. *Proc. Intern. Conf. Magnetism and Cryst., Kyoto, Japan, Sept. 1961. J. Phys. Soc. Japan* **17**, Suppl. B2, 376-82 (1962).
- FRAZER, B.C. AND BROWN, P.J. The antiferromagnetic structure of CrVO_4 and the anhydrous sulfates of divalent Fe, Ni, and Co. *Phys. Rev.* **125**, 1283-91 (1962).
- FRAZER, B.C. - See also KAY, M.I.
- FUJII, T., CHADWICK, G.B., COLLINS, G.B., DUKE, P.J., HIEN, N.C., KEMP, M.A.R. AND TURKOT, F. Elastic proton-proton scattering at 1.35, 2.1, and 2.9 Bev. (A) *Bull. Am. Phys. Soc.* **7**, 45 (1962).
- FUJII, T. - See also DUKE, P.J.; HIEN, N.C.; KEMP, M.A.R.
- FUJITA, F.E. Oxidation and dislocation mechanism in fatigue crack formation. (A) Presented at Intern. Conf. on Fracture, Lake Wilderness, Maple Valley, Wash., Aug. 1962.
- FUJITA, F.E. AND DAMASK, A.C. Electrical resistivity changes in neutron-irradiated iron-carbon. (A) *Bull. Am. Phys. Soc.* **7**, 344 (1962).
- FUJITA, F.E. AND DAMASK, A.C. The influence of radiation-produced defects on the precipitation of carbon in iron. (A) Presented at Intern. Conf. Crystal Lattice Defects, Kyoto, Japan, Sept. 1962.
- GAILLOUD, M. - See BROWN, H.; CULWICK, B.
- GILLIAM, O.R. - See WYLIE, D.W.
- GOLAND, A.N. Atomic displacements in solids by nuclear radiation. Submitted to *Ann. Rev. Nucl. Sci.*

- GOLAND, A.N. Comparison of electron-microscope observations in neutron-irradiated copper and gold. (L) *J. Nucl. Mater.* (in press).
- GOLAND, A.N. - See also ARTMAN, R.A.; KELSCH, J.J. (Nuclear Engineering); VINEYARD, G.H.
- GOLDBERG, M. - See BERTANZA, L.
- GOLDHABER, G.S. - See EMERY, G.T.; SCHMORAK, M.
- GRAY, L. - See BERTANZA, L.
- GUPTA, S.N. Relativistic Schrödinger's equation for the two-nucleon system. *Phys. Rev.* **124**, 1638-40 (1961).
- HART, E.L., LOUITT, R.I., LUERS, D., MORRIS, T.W., WILLIS, W.J. AND YAMAMOTO, S.S. Multiple meson production in proton-proton collisions at 2.85 BeV. *Phys. Rev.* **126**, 747-56 (1962).
- HART, E.L., LOUITT, R.I. AND MORRIS, T.W. Proton-proton scattering at 1.5 BeV. (A) *Bull. Am. Phys. Soc.* **7**, 349 (1962).
- HART, E.L. - See also BERTANZA, L.
- HARTH, E. - See BERTANZA, L.
- HATCHER, R.D. AND DIENES, G.J. Relaxation and activation energies for an interstitial neutral defect in an alkali halide lattice. *Phys. Rev.* **124**, 726-35 (1961).
- HIE, N.C., CHADWICK, G.B., COLLINS, G.B., DUKE, P.J., FUJII, T., KEMP, M.A.R. AND TURKOT, F. Inelastic p - p scattering at 1.35 BeV. (A) *Bull. Am. Phys. Soc.* **7**, 45 (1962).
- HIE, N.C. - See also DUKE, P.J.; FUJII, T.; KEMP, M.A.R.
- HILL, D.A. - See COOL, R.L.
- HOUGH, P.V.C. *A Computer Learns To See, Brookhaven Lecture Series No. 14*, BNL 725 (T-259), Feb. 14, 1962.
- HOUGH, P.V.C. General purpose visual input for a computer. Submitted to *Ann. N.Y. Acad. Sci.*
- HOUGH, P.V.C. - See also BERTANZA, L.
- JACH, J. The thermal decomposition of irradiated α -lead azide. Submitted to *Trans. Faraday Soc.*
- JAIN, A.P., CHRIEN, R.E. AND PALEVSKY, H. Average neutron cross sections in the 10 to 100-kev region. (A) *Bull. Am. Phys. Soc.* **7**, 22 (1962).
- JAIN, A.P., CHRIEN, R.E. AND PALEVSKY, H. Relationship between average neutron cross sections and strength functions. (A) *Bull. Am. Phys. Soc.* **7**, 289 (1962).
- JAIN, A.P. - See also CHRIEN, R.E.
- JENKINS, E.W. - See BAKER, W.F.; COOL, R.L.
- KACSER, C. - See BARTON, G.
- KALOGEROPOULOS, T. - See BROWN, H.; CULWICK, B.; SAMIOS, N.P.
- KAN, H.K.A. - See TOWNSEND, P.D.
- KANE, J.V. - See DONOVAN, P.F.; EL-WAHAB, M.A.; LOWE, J.; ZUPANCIC, C.
- KANE, W.R., FIEBIGER, N.F. AND FOX, J.D. Coincidence studies of the thermal neutron capture gamma rays of chromium. *Phys. Rev.* **125**, 2037-48 (1962).
- KANE, W.R. - See also CRASEMANN, B.; EMERY, G.T.; FIEBIGER, N.F.
- KAY, M.I., FRAZER, B.C. AND UEDA, R. The disordered structure of NaNO_2 at 185°C . *Acta Cryst.* **15**, 506-8 (1962).
- KEATING, D.T. AND WALKER, C.B. Extinction and diffuse neutron intensity measurements. Submitted to *Acta Cryst.*
- KEMP, M.A.R., CHADWICK, G.B., COLLINS, G.B., DUKE, P.J., FUJII, T., HIE, N.C. AND TURKOT, F. Inelastic proton-proton scattering at 2.1 and 2.9 BeV. (A) *Bull. Am. Phys. Soc.* **7**, 46 (1962).
- KEMP, M.A.R. - See also FUJII, T.; HIE, N.C.
- KIM, C.H. - See FLANAGAN, T.B.
- KISTNER, O.C., KOCHER, C.W., MOZER, B. AND SWAN, J.B. Recoil-free nuclear resonance absorption spectrum of Fe^{57} impurity atoms in beryllium. (A) *Bull. Am. Phys. Soc.* **7**, 294 (1962).
- KISTNER, O.C. AND MOZER, B. Origin of the Mössbauer absorption spectra of iron impurity atoms in beryllium. (A) *Bull. Am. Phys. Soc.* **7**, 505 (1962).
- KISTNER, O.C. AND SUNYAR, A.W. Decay of long-lived Ag^{108m} . (A) *Bull. Am. Phys. Soc.* **7**, 342 (1962).
- KOCHER, C.W. AND BROWN, P.J. The atomic moments and hyperfine fields in Fe_2Ti and Fe_2Zr . *J. Appl. Phys. Suppl.* **33**, 1091-2 (1962).
- KOCHER, C.W. - See also KISTNER, O.C.
- KOPP, J. - See BROWN, H.
- KYCIA, T.F. - See BAKER, W.F.; COOL, R.L.
- LEA, R.M. AND VITTITOE, C. Bubble density measurements for particle identification. *Rev. Sci. Instr.* **33**, 243-4 (1962).
- LEA, R.M. - See also BERTANZA, L.; BROWN, H.; CULWICK, B.; SAMIOS, N.P.
- LEIPUNER, L.B. - See CRITTENDEN, R.R.
- LEITNER, J. - See BERTANZA, L.
- LEVY, P.W. Annealing of radiation-induced defects in fused silica. (A) Presented at 2nd Intern. Conf. Radiation Effects in Glass and Related Materials, Rochester, N.Y., April 1962.
- LEVY, P.W. Stages in the annealing of radiation-induced color centers and defects in quartz. (A) *Bull. Am. Phys. Soc.* **7**, 187 (1962).
- LEVY, P.W. - See also TOWNSEND, P.D.; WYLIE, D.W.
- LI, A.C. AND SCHWARZSCHILD, A. Lifetimes of the $4+$ and $2+$ rotational states in Dy^{162} and Er^{168} . (A) *Bull. Am. Phys. Soc.* **7**, 359 (1962).
- LI, A.C. AND SCHWARZSCHILD, A. Lifetimes of the $2+$ and $4+$ rotational states in Hf^{180} . Presented at Conf. Magnetic Lifetimes and Properties of Nuclear States, Gatlinburg, Tenn., Oct. 1961.
- LI, A.C. - See also BOLOTIN, H.H.
- LIGHTMAN, S. - See BERTANZA, L.
- LINDENBAUM, S.J. Shielding of high-energy accelerators. *Ann. Rev. Nucl. Sci.* **11**, 213-58 (1961).
- LINDENBAUM, S.J., LOVE, W.A., NIEDERER, J.A., OZAKI, S., RUSSELL, J.J. AND YUAN, L.C.L. The antiproton-proton and proton-proton total cross sections from 4 to 20 BeV/c. (L) *Phys. Rev. Letters* **7**, 185-8 (1961).
- LINDENBAUM, S.J., LOVE, W.A., NIEDERER, J.A., OZAKI, S., RUSSELL, J.J. AND YUAN, L.C.L. $\pi^+ + p$ and $\pi^- + p$ Total cross sections from 8 to 20 BeV/c. (L) *Phys. Rev. Letters* **7**, 352-4 (1961).
- LINDENBAUM, S.J. - See also BAKER, W.F.; MELISSINOS, A.C.
- LOUITT, R. - See BROWN, H.; HART, E.L.
- LOVE, W.A. - See BAKER, W.F.; LINDENBAUM, S.J.
- LOWE, J. AND MCCLELLAND, C.L. RF vernier time to pulse-height converter. (A) *Bull. Am. Phys. Soc.* **7**, 261 (1962).
- LOWE, J., MCCLELLAND, C.L. AND KANE, J.V. Lifetimes of the first excited states of C^{15} and B^{10} . *Phys. Rev.* **126**, 1811-18 (1962).

- LOWE, J. - See also McCLELLAND, C.L.
 LUERS, D. - See BAKER, W.F.; HART, E.L.
 MARR, R.B. - See BÉG, M.A.B.; DURAND, L. III.
 MARSHAK, H. Transmission of polarized neutrons through polarized Li^6 nuclei. (A) *Bull. Am. Phys. Soc.* 7, 305 (1962).
 MARSHAK, H., POSTMA, H., SAILOR, V.L., SHORE, F.J. AND REYNOLDS, C.A. Spin assignments of low-energy resonances using polarized neutrons and polarized Sm^{149} nuclei. Submitted to *Phys. Rev.*
 MARSHAK, H. - See also BRUNHART, G.; POSTMA, H.; SAILOR, V.L.
 MARSHALL, L. A magnetic lens system of unit magnification for high energy charged particles. Submitted to *Rev. Sci. Instr.*
 MARSHALL, L. - See also COOL, R.L.
 MARTIN, H.J. - See CRITTENDEN, R.R.
 MARTIN, J.P. AND ZUCKER, M.S. Differential cross section and polarization angular distributions for $\text{O}^{16}(n,n)\text{O}^{16*}$. (A) *Bull. Am. Phys. Soc.* 7, 72 (1962).
 McCLELLAND, C.L. AND LOWE, J. Upper limit on the F^{18} 940-kev level lifetime. (A) *Bull. Am. Phys. Soc.* 7, 340 (1962).
 McCLELLAND, C.L. - See also LOWE, J.
 McKEOWN, M. - See EMERY, G.T.
 MEETZ, K. Spectrum of the scattering integral operator in the physical energy sheet. *J. Math. Phys.* 3, 690-9 (1962).
 MELISSINOS, A.C., FAZIO, G.G., YAMANOUCHI, T., LINDENBAUM, S.J. AND YUAN, L.C.L. (L) *Phys. Rev. Letters* 7, 454-7 (1961).
 MERRIL, D.W. JR. - See PALEVSKY, H.
 MITTRA, I. - See BERTANZA, L.
 MONETI, G. - See BERTANZA, L.
 MOORE, J.A. - See PALEVSKY, H.
 MORAN, T.I. - See COHEN, V.W.; PENSELIN, S.
 MORRIS, T.W. - See BERTANZA, L.; BROWN, H.; CULWICK, B.; HART, E.L.
 MOZER, B., MYERS, V.W. AND OTNES, K. Observation of localized modes of vibration in low concentration alloys. (A) *Bull. Am. Phys. Soc.* 7, 16 (1962).
 MOZER, B. AND OTNES, K. Measurements of the vibrational frequency spectrum of nickel-palladium alloys. (A) Presented at IAEA Symp. Inelastic Scattering of Neutrons in Solids and Liquids, Chalk River, Canada, Sept. 1962.
 MOZER, B., OTNES, K. AND MYERS, V.W. Measurement of a simple defect mode of vibration. (L) *Phys. Rev. Letters* 8, 278-80 (1962).
 MOZER, B. - See also KISTNER, O.C.; VINEYARD, G.H.
 MUETHER, H.R. - See COTTER, M.J.
 MUNIR, B.A. - See SALANT, E.O.; ZORN, G.T.
 MUSGRAVE, B. - See CRITTENDEN, R.R.
 MYERS, V.W. - See MOZER, B.
 NATHANS, R., PICKART, S.J. AND ALPERIN, H.A. Spin density distributions and Hartree-Fock calculations for the iron group series. Submitted to *J. Phys. Soc. Japan*.
 NATHANS, R. - See also ALPERIN, H.A.; BROWN, P.J.; PICKART, S.J.
 NIEDERER, J.A. - See BAKER, W.F.; LINDENBAUM, S.J.
 OLNES, J.W. AND FIEBIGER, N.F. Levels of Se^{77} and Se^{78} from (n,γ) measurements. (A) *Bull. Am. Phys. Soc.* 7, 315 (1962).
 OTNES, K. AND PALEVSKY, H. A new apparatus for inelastic, quasi-elastic and elastic cold neutron measurements. (A) Presented at IAEA Symp. Inelastic Scattering of Neutrons in Solids and Liquids, Chalk River, Canada, Sept. 1962.
 OTNES, K. - See also MOZER, B.; PALEVSKY, H.; WAKUTA, Y.
 OZAKI, S. - See BAKER, W.F.; LINDENBAUM, S.J.
 PALEVSKY, H., MOORE, J.A., SUTTER, R.J., STEARNS, R.L., WAKUTA, Y. AND MERRIL, D.W. JR. Nucleon charge exchange at 3 Bev. (A) *Bull. Am. Phys. Soc.* 7, 46 (1962).
 PALEVSKY, H., OTNES, K. AND WAKUTA, Y. Low frequency hydrogen vibrations in potassium dihydrogen phosphate. (A) Presented at IAEA Symp. Inelastic Scattering of Neutrons in Solids and Liquids, Chalk River, Canada, Sept. 1962.
 PALEVSKY, H. - See also BRAJOVIC, V.; CHRIEN, R.E.; COTTER, M.J.; JAIN, A.; OTNES, K.; WAKUTA, Y.
 PALMER, R. - See BERTANZA, L.
 PASTERNAK, S. AND STERNHEIMER, R.M. An orthogonality property of hydrogenlike radial functions. Submitted to *J. Math. Phys.*
 PENSELIN, S., MORAN, T., COHEN, V.W. AND WINKLER, G. Hyperfine structure of the electronic ground states of Rb^{85} and Rb^{86} . *Phys. Rev.* 127, 524-8 (1962).
 PENSELIN, S. - See also COHEN, V.W.
 PERLMAN, M.L. - See EMERY, G.T.
 PICKART, S.J. Fourier projections of unpaired electron densities. Submitted to *Acta Cryst.*
 PICKART, S.J. AND NATHANS, R. Alloys of the first transition series with Pd and Pt. (A) *J. Appl. Phys. Suppl.* 33, 1336-8 (1962).
 PICKART, S.J. - See also ALPERIN, H.A.; BROWN, P.J.; NATHANS, R.
 PICKUP, E., ROBINSON, D.K. AND SALANT, E.O. π - π Resonance in π^-p interactions at 1.25 Bev. (L) *Phys. Rev. Letters* 7, 192-5 (1961).
 PICKUP, E., ROBINSON, D.K. AND SALANT, E.O. p - p Interactions at 2 Bev. II. Multiple-pion production. *Phys. Rev.* 125, 2091-101 (1962).
 PICKUP, E., ROBINSON, D.K. AND SALANT, E.O. Three-pion mass distributions and the η -meson. (L) *Phys. Rev. Letters*, 8, 329-32 (1962).
 PICKUP, E. - See also AYER, F.; FICKINGER, W.J.; ROBINSON, D.K.; SALANT, E.O.
 POSTMA, H., MARSHAK, H., SAILOR, V.L., SHORE, F.J. AND REYNOLDS, C.A. The transmission of polarized neutrons through samples of polarized Ho^{165} nuclei. *Phys. Rev.* 126, 979-85 (1962).
 POSTMA, H. - See also MARSHAK, H.; SAILOR, V.L.
 PRODELL, A. - See BERTANZA, L.
 RAHM, D.C. A rapid beam ejector for the Cosmotron. *Rev. Sci. Instr.* 32, 1116-19 (1961).
 RAPPE, E.F. - See AYER, F.
 RAU, R. - See BERTANZA, L.
 READ, A.L. - See BAKER, W.F.
 REYNOLDS, C.A. - See BRUNHART, G.; MARSHAK, H.; POSTMA, H.; SAILOR, V.L.
 ROBINSON, D.K., PICKUP, E. AND SALANT, E.O. Test of the one pion exchange model. (A) *Bull. Am. Phys. Soc.* 7, 265 (1962).

- ROBINSON, D.K. - See also FICKINGER, W.J.; PICKUP, E.; SALANT, E.O.
- ROSE, M.E. AND YANG, C.N. Eigenvalues and eigenvectors of a symmetric matrix of $6j$ symbols. *J. Math. Phys.* **3**, 106 (1962).
- ROWLAND, R.S. - See FIREMAN, E.L.
- RUSSELL, J.J. - See BAKER, W.F.; LINDENBAUM, S.J.
- SAFFORD, G., BRAJOVIC, V. AND BOUTIN, H. An investigation of the energy levels in alkaline-earth hydroxides by inelastic scattering of slow neutrons. (A) *Bull. Am. Phys. Soc.* **7**, 499 (1962).
- SAFFORD, G.J., RUSH, J.J., TAYLOR, T.I. AND HAVENS, W.W. JR. The effect of rotational freedom in several ammonium salts and dimethyl acetylene on the inelastic scattering of slow neutrons. Submitted to *Nucl. Sci. Eng.*
- SAFFORD, G. - See also BOUTIN, H.; BRAJOVIC, V.
- SAILOR, V.L., SCHERMER, R.I., SHORE, F.J., REYNOLDS, C.A., MARSHAK, H. AND POSTMA, H. Nuclear polarization in polycrystalline holmium-indium alloy. *Phys. Rev.* **127**, 1124-9 (1962).
- SAILOR, V.L. - See also BERNABEI, A.; BRUNHART, G.; MARSHAK, H.; POSTMA, H.
- SALANT, E.O., PICKUP, E., ROBINSON, D.K. AND MUNIR, B.A. Pion-pion interactions and pion production branching ratios at the third resonance. *Revs. Mod. Phys.* **33**, 435-6, (1961).
- SALANT, E.O. - See also FICKINGER, W.J.; PICKUP, E.; ROBINSON, D.K.
- SAMIOS, N.P., BACHMAN, A.H., LEA, R.M., KALOGEROPOULOS, T.E. AND SHEPHARD, W.D. Multiple-meson production in π^-p collisions at 4.7 BeV/c. *Proc. CERN Intern. Conf. High Energy Physics, Geneva, July 1962* (in press).
- SAMIOS, N.P., PLANO, R., PRODELL, A., SCHWARTZ, M. AND STEINBERGER, J. The parity of the neutral pion and the decay $\pi^0 \rightarrow 2e^+ + 2e^-$. *Phys. Rev.* **126**, 1844-9 (1962).
- SAMIOS, N.P. - See also BERTANZA, L.
- SANDWEISS, J. - See BALTAY, C.; BROWN, H.
- SANFORD, J. - See BALTAY, C.; BROWN, H.
- SCHERMER, R.I. - See BRUNHART, G.; SAILOR, V.L.
- SCHLUTER, R.A. - See COOL, R.L.
- SCHMORAK, M., EMERY, G.T. AND GOLDBABER, G.S. $M3$ isomeric transition in tin-113. *Phys. Rev.* **124**, 1186-9 (1961).
- SCHWARZSCHILD, A. Review of latest developments in delayed coincidence measurements. Presented at Conf. Electromagnetic Lifetimes and Properties of Nuclear States, Gatlinburg, Tenn., Oct. 1961.
- SCHWARZSCHILD, A. - See also BOLOTIN, H.H.; LI, A.C.
- SEGEL, R.E. - See FIEBIGER, N.F.
- SEHGAL, M.L. Decay of Sb^{115} . *Phys. Rev.* **125**, 968-71 (1962).
- SEHGAL, M.L. Isomeric cross section ratios in (n, γ) reactions. Submitted to *Phys. Rev.*
- SHEPHARD, W.D. - See SAMIOS, N.P.
- SHORE, F.J. - See BRUNHART, G.; MARSHAK, H.; POSTMA, H.; SAILOR, V.L.
- SHUSKUS, A.J. - See WYLIE, D.W.
- SHUTT, R. - See BROWN, H.
- SKILLICORN, I. - See BERTANZA, L.
- STEARNS, R.L. - See PALEVSKY, H.
- STERNHEIM, M.M. Second-order effects of nuclear magnetic fields. *Phys. Rev.* (in press).
- STERNHEIMER, R.M. Electronic polarizabilities of the alkali atoms. *Phys. Rev.* **127**, 1220-3 (1962).
- STERNHEIMER, R.M. Second-order quadrupole effect for the nuclear hexadecapole coupling in ions. *Phys. Rev.* **127**, 812-16 (1962).
- STERNHEIMER, R.M. - See also PASTERNAK, S.
- SUNYAR, A.W. - See KISTNER, O.C.
- SUTTER, R.J. - See PALEVSKY, H.
- SWAN, J.B. - See KISTNER, O.C.
- TAFT, H. - See BROWN, H.
- THORNDIKE, A.M. - See BERTANZA, L.; BROWN, H.; CULWICK, B.
- TOWNSEND, P.D., KAN, H.K.A. AND LEVY, P.W. Effects of irradiation on the photoconductivity of rutile. (A) *Bull. Am. Phys. Soc.* **7**, 187 (1962).
- TURKOT, F. - See DUKE, P.J.; FUJII, T.; HIEN, N.C.; KEMP, M.A.R.
- UEDA, R. - See KAY, M.I.
- VINEYARD, G.H., MOZER, B. AND GOLAND, A.N. Thermal spikes in an atomic model. (A) *Bull. Am. Phys. Soc.* **7**, 170 (1962).
- VITTITOE, C. - See LEA, R.M.
- WAGENBLAST, H. AND DAMASK, A.C. Discussion of the paper of Dr. Berry. (N) *Acta Met.* **10**, 333 (1962).
- WAGENBLAST, H. AND DAMASK, A.C. Kinetics of carbon precipitation in irradiated iron. *Phys. Chem. Solids* **23**, 221-7 (1962).
- WAKUTA, Y., OTNES, K. AND PALEVSKY, H. Low frequency hydrogen vibrations in potassium dihydrogen phosphate. (A) *Bull. Am. Phys. Soc.* **7**, 356 (1962).
- WAKUTA, Y. - See also PALEVSKY, H.
- WALKER, C.B. - See KEATING, D.T.
- WALL, G. - See AYER, F.
- WEBSTER, M. - See BALTAY, C.; BROWN, H.
- WENESER, J. - See CHURCH, E.L.
- WESTGARD, J. - See BERTANZA, L.
- WICK, G.C. Angular momentum states for three relativistic particles. *Ann. Phys. N.Y.* **18**, 65-80 (1962).
- WILKINSON, D.H. A self-consistent determination of the effective radii of heavy nuclei for alpha-particle emission. *Phys. Rev.* **126**, 648-52 (1962).
- WILLIS, W.J. - See HART, E.L.
- WINKLER, G. - See PENSELIN, S.
- WYLIE, D.W., SHUSKUS, A.J., YOUNG, C.G., GILLIAM, O.R. AND LEVY, P.W. Electron spin resonance of radiation-induced defects in potassium azide. *Phys. Rev.* **125**, 451-3 (1962).
- YAMAMOTO, S.S. - See BALTAY, C.; BERTANZA, L.; HART, E.L.
- YAMANOUCHI, T. - See MELISSINOS, A.C.
- YANG, C.N. - See ROSE, M.E.
- YOUNG, C.G. - See WYLIE, D.W.
- YUAN, L.C.L. - See BAKER, W.F.; LINDENBAUM, S.J.; MELISSINOS, A.C.
- ZORN, B.S. Deuterons in $p-p$ collision at 2 BeV. (A) *Bull. Am. Phys. Soc.* **7**, 349 (1962).
- ZORN, B.S. Pion-pion resonances in a pure $T=1$ state. (L) *Phys. Rev. Letters* **8**, 282-3 (1962); Erratum. *Ibid.* 386.
- ZORN, B.S. AND ZORN, G.T. Bubble density in a hydrogen bubble chamber. Submitted to *Nuovo Cimento*.

- ZORN, G.T. AND MUNIR, B.A. π - p Interactions at 5 BeV/c.
(A) *Bull. Am. Phys. Soc.* 7, 281 (1962).
ZORN, G.T. - See also FISCHER, J. (Instr. & Health Physics); ZORN, B.S.
ZUCKER, M.S. - See MARTIN, J.P.
ZUPANCIC, C., KANE, J.V. AND DONOVAN, P.F. A knock-out model for the reaction $O^{16}(\alpha, 2\alpha)C^{12}$ ground state.
(A) *Bull. Am. Phys. Soc.* 7, 287 (1962).
ZUPANCIC, C. - See DONOVAN, P.F.

Reactor Division

- OSBORNE, C.L. *The Seventeenth Annealing of the Graphite Moderator Structure in the BNL Reactor, Sept. 1961*. BNL 701 (T-242). For Official Use Only.
POWELL, R.W. AND OSBORNE, C.L. *Reactor Operations Monthly Report, July 1961*. BNL 688 (T-234). For Official Use Only.

THOMPSON, D. Facility Availability, Brookhaven Graphite Research Reactor, January 1951 to March 1962. *Nucleonics* (in press).

THOMPSON, D. *Reactor Operations Monthly Reports* (For Official Use Only):

Aug. 1961	BNL 693 (T-237)
Sept. 1961	BNL 698 (T-240)
Oct. 1961	BNL 704 (T-244)
Nov. 1961	BNL 707 (T-245)
Dec. 1961	BNL 708 (T-246)
Jan. 1962	BNL 712 (T-249)
Feb. 1962	BNL 717 (T-253)
Mar. 1962	BNL 723 (T-257)
Apr. 1962	BNL 729 (T-261)
May 1962	BNL 730 (T-262)
June 1962	BNL 744 (T-271)

Appendix B

OFFICERS AND SCIENTIFIC STAFF

Gerald F. Tape, *Acting Director* †
 Maurice Goldhaber, *Director*
 Clarke Williams, *Deputy Director*
 Charles E. Falk, *Associate Director*
 R. Christian Anderson, *Assistant Director*
 G. Norris Glasoe, *Assistant Director*
 Samuel M. Tucker, *Assistant Director*

G. Kenneth Green, *Chairman*, Accelerator Department
 Howard J. Curtis, *Chairman*, Biology Department
 Richard W. Dodson, *Chairman*, Chemistry Department
 Horner Kuper, *Chairman*, Instrumentation and Health Physics Department
 Lee E. Farr, *Chairman*, Medical Department *
 Victor P. Bond, *Chairman*, Medical Department
 Warren E. Winsche, *Chairman*, Nuclear Engineering Department
 George H. Vineyard, *Chairman*, Physics Department

John W. Calkin, *Head*, Applied Mathematics Division
 Irving J. Polk, *Head*, Mechanical Engineering Division
 Robert W. Powell, *Head*, Reactor Division

J. Georges Peter, *Director*, Architectural Planning
 H. Russell Cort, *Budget Officer*
 Maxwell M. Small, *Business Manager* *
 William J. Catacosinos, *Business Manager*
 Lewis R. Burchill, *Controller*
 Dennis Puleston, *Information Officer*
 Charles F. Dunbar, *Legal Counsel*
 Joseph S. Washburne, *Personnel Manager*
 Edward J. Burke, *Security Officer*

Accelerator Department

G. Kenneth Green, *Chairman*
 Roger R. Adams
 Mark Q. Barton
 Richard A. Beth
 John W. Bittner
 John P. Blewett
 M. Hildred Blewett
 Hugh N. Brown
 William Chinowsky*
 Theodore N. Constant
 James G. Cottingham
 Gordon T. Danby
 Donald A. Davis
 Edward W. Dexter
 Eric B. Forsyth

†On January 18, 1962, Dr. Tape was appointed Vice President of Associated Universities, Inc.

*Terminated before July 1, 1962.

Hyota Fujita
(on leave from Mitsubishi Atomic Power Industries, Inc., Japan)
 William Gefers
 Salvatore T. Giordano
 Harald Hahn
 Henry J. Halama
 Ralph R. Kassner
 Lawrence B. Leipuner
 William T. Link*
 Isador J. Livant
 Robert A. Loper
 Alfred W. Maschke
 Lowell McLean
 Walter W. Merkle
 William H. Moore
 Martin Plotkin
 Eugene C. Raka
 Raymond H. Rheaume
 David S. Robertson*
(on leave from Australian National Univ.)

Everett J. Rutan
 Eleftherios J. Sacharidis
(postdoctoral appointment)
 Edward E. Shelton
 Lyle W. Smith
 Julius Spiro
 Clifford E. Swartz
 Clarence M. Turner
 Arie van Steenberg
 John L. Walters*

Applied Mathematics Division

John W. Calkin, *Head*
 John R. Cannon
 Katherine T. Chen*
(postdoctoral appointment)
 Robert B. Marr
 Joel D. Pincus
 Stuart S. Rideout
 Jerome A.G. Russell
 Yoshio Shimamoto

Biology Department

Howard J. Curtis, *Chairman*
 Leroy G. Augenstein*
 Dai H. Bai*
(on leave from
Agricultural Experimental Station, Korea)
 John A. Bergeron
 John Berech, Jr.
 James L. Brewbaker*
 John A.M. Brown
(postdoctoral appointment)
 Milislav Demerec
 H. John Evans*
(assigned from
Medical Research Council, England)
 William R. Evans
(postdoctoral appointment)
 David L. Filmer
(postdoctoral appointment)
 Eric C. Gaetjens*
(postdoctoral appointment)
 Anne D. Gounaris
(postdoctoral appointment)
 Lewis J. Greene
 Ashraful U. Haque*
(on leave from Univ. of Dacca, Pakistan)
 William S. Hillman
 C.H.W. Hirs
 H. Robert Horton
(assigned from U.S. Public Health Service)
 James R. Innes
 Yashwant Karkhanis
(postdoctoral appointment)
 Norman F. Kember*
(postdoctoral appointment)
 Carl E. Kirchner*
(postdoctoral appointment)
 J. Raymond Klein
 Daniel E. Koshland, Jr.
 Marian E. Koshland
 Beyoung H. Kwack*
(IAEA Fellow)
 Sanford A. Lacks
 Eugene P. Lazzari
(postdoctoral appointment)
 Myron Levine*
 Harvard Lyman
(on leave from Brooklyn College)
 Leo W. Mericle*
(on leave from Michigan State Univ.)
 Rae P. Mericle*
(assigned from Michigan State Univ.)
 Jerome P. Miksche
 Samuel M. Mozersky
(assigned from National Inst. of Health)
 Leslie F. Nims
 John M. Olson
 Carmie A. Perrotta
(assigned from Yale Univ.)

*Terminated before July 1, 1962.

Andrew E. Petracco
(graduate student from Yale Univ.)
 Thomas H. Plummer, Jr.
(postdoctoral appointment)
 Wolf Prenskey
(assigned from U.S. Public Health Service)
 Henry Quastler
 Louis C.R. Sarda
(assigned from
Univ. of Sciences, Marseilles, France)
 Gideon W. Schaeffer
(assigned from U.S. Public Health Service)
 Erwin A. Schwinghamer*
 Kenneth Scott
(postdoctoral appointment)
 Seymour Shapiro*
 Donald L. Shaver
(postdoctoral appointment)
 Robert M. Smillie
 Harold H. Smith
 Arnold H. Sparrow
 Robert Steele
 David H. Strumeyer
(postdoctoral appointment)
 Christiaan Sybesma
(postdoctoral appointment)
 Austin L. Taylor
(postdoctoral appointment)
 Donald E. Wimber
 Marilyn F. Wolfsberg
(assigned from U.S. Public Health Service)
 Philip S. Woods*
 George M. Woodwell
(on leave from Univ. of Maine)
 Geoffrey Zubay

Chemistry Department

Richard W. Dodson, *Chairman*
 Augustine O. Allen
 Desire J. Apers
(on leave from Inst. Interuniversitaire
des Sciences Nucléaires, Belgium)
 Elizabeth W. Baker
 William A. Bassett*
(postdoctoral appointment)
 Benon H. Bielski
 Jacob Bigeleisen
(on leave to
Swiss Federal Inst. of Technology, Zurich)
 Ningondappa S. Biradar*
(on leave from Government of India)
 Tycho F. Burger*
(postdoctoral appointment)
 Fulvio Cacase
(on leave from Univ. of Rome, Italy)
 Thomas C. Castorina
(assigned from Picatinny Arsenal)
 Joseph C.Y. Chen
(postdoctoral appointment)
 Lung-Wen Chiao
(postdoctoral appointment)

David R. Christman
 Yung Yee Chu
 James C. Cobb
 Kenneth E. Collins
(postdoctoral appointment)
 Philip Coppens
(postdoctoral appointment)
 Lester M. Corliss
 William C. Corning
(graduate student from Univ. of Rochester)
 Edgard H. Crevecoeur*
(on leave from Inst. Interuniversitaire
des Sciences Nucléaires, Belgium)
 James B. Cumming
 Gidon Czapski
(postdoctoral appointment)
 Raymond Davis, Jr.
 Hartmut Diebler
(postdoctoral appointment)
 Bodo Diehn
(graduate student from Univ. of Kansas)
 Paul F. Donovan
(assigned from
Bell Telephone Laboratories, Inc.)
 Israel Dostrovsky
(on leave from
Weizmann Inst. of Science, Israel)
 Guenther Dulz
(graduate student from Columbia Univ.)
 Norman Elliott
 Harry R. Fickel*
(postdoctoral appointment)
 Michael H. Ford-Smith*
(postdoctoral appointment)
 Zeev Fraenkel*
(on leave from
Weizmann Inst. of Science, Israel)
 Simon Freed
 Gerhart Friedlander
 Lewis Friedman
 James R. Grover
 Richard L. Hahn*
(postdoctoral appointment)
 Masayuki Hamada
(on leave from Radiation Center
of Osaka Prefecture, Japan)
 Walter C. Hamilton
 Garman Harbottle
 Julius M. Hastings
 Jerome Hudis
 Andries Hummel
(assigned from
Ministry of Education, Arts, and Sciences,
Netherlands)
 James A. Ibers
 Adolph P. Irsa
 Karin Karlstrom
(assigned from Columbia Univ.)
 Demetrius Katakis*
(postdoctoral appointment)
 Seymour Katcoff

Priscilla Kaufman*
 Ann M. Kistner
(graduate student from Columbia Univ.)
 James F. Mollenauer
(postdoctoral appointment)
 Gordon G. Moore
(postdoctoral appointment)
 Robert T. Mullen
(postdoctoral appointment)
 John J. Naughton*
(on leave from Univ. of Hawaii)
 Elinor F. Norton
 Morris L. Perlman
 Norbert T. Porile
 Arthur M. Poskanzer
 Leon A. Rajbenbach*
(postdoctoral appointment)
 K. Narayano Rao*
(postdoctoral appointment)
 Carol S. Redvanly
 Bruno Reitzner
(assigned from Picatinny Arsenal)
 Louis P. Remsberg, Jr.
(postdoctoral appointment)
 Slobodan V. Ribnikar*
(postdoctoral appointment)
 J. Keith Rowley
 William Robinson
 John S. Rush*
(graduate student from Columbia Univ.)
 Eiichi Saito*
(on leave from Centre d'Études Nucléaires, Saclay, France)
 Edward V. Sayre
 Oliver A. Schaeffer
 Harold A. Schwarz
 Stanley Seltzer
 Morris Slavin
 Heinz I. Stangl*
(postdoctoral appointment)
 Marvin J. Stern
(postdoctoral appointment)
 Gerhard Stocklin
(postdoctoral appointment)
 Raymond W. Stoenner
 Norman Sutin
 Shigeo Tanaka
(on leave from Univ. of Tokyo, Japan)
 Yutaka Tanaka*
(on leave from Inst. of Physical and Chemical Research, Japan)
 Anthony J. Tench
(postdoctoral appointment)
 T. Darrah Thomas*
 Sydney O. Thompson
 W. Alexander Van Hook
(postdoctoral appointment)
 Slobodanka R. Veljkovic*
(on leave from Univ. of Belgrade, Yugoslavia)

*Terminated before July 1, 1962.

Richard Visser*
(on leave from Univ. of Amsterdam, Netherlands)
 Helge T. Von Koch
(postdoctoral appointment)
 Ralph E. Weston, Jr.
 Alfred P. Wolf
 Max Wolfsberg
 Armin Wyttenbach
(postdoctoral appointment)
 Kenji Yoshihara*
(on leave from Japan Atomic Energy Research Inst., Tokai-mura)

Instrumentation and Health Physics Department

Horner Kuper, *Chairman*
 Arland L. Carsten
 Robert L. Chase
 Frederick P. Cowan
 Carl H. Distenfeld
 Jack C. Faust
 Joachim Fischer
 Charles F. Foelix
 Lee Gemmell
 John S. Handloser
 William J. Hartin
 William A. Higinbotham
 Andrew P. Hull
 Kazuhiko Imai*
(on leave from Japan Atomic Energy Research Inst., Tokai-mura)
 Robert O. McClintock
 Charles B. Meinhold
 Gabriel L. Miller
 James P. Palmer
 Howard R. Pate
 Paulo P. Petry
(on leave from Univ. of Rio Grande do Sul, Brazil)
 Leigh F. Phillips
 David W. Potter
 Seymour Rankowitz
 Edwin J. Rogers
 Martin J. Rosenblum
 George E. Schwender
 Irving A. Singer
 Maynard E. Smith
 Robert J. Spinrad
 Raymond W. Stong
 Robert M. Sugarman*

Mechanical Engineering Division

Irving J. Polk, *Head*
 Robert D. Baldwin, Jr.
 Vernon J. Buchanan
 Basil De Vito
 Carl R. Flatau
 Carl L. Goodzeit
 Charles L. Gould
 John J. Grisoli

Rudolph S. Hodor
 Kenneth C. Hoffman
 Donald W. Huszagh
 Jack E. Jensen
 Michael B. Karelitz
 David A. Kassner
 Calman Lasky
 Robert J. McCracken
 Kurt F. Minati
 George Nugent
 Oliver S. Reading
 Albert P. Schlafke, Jr.
 Charles Theisen
 William G. Walker

Medical Department

Lee E. Farr, *Chairman**
 Victor P. Bond, *Chairman*
 Abbas Alaghemand*
(medical associate)
 Vicente Alcober
(on leave from Facultad de Medicina, Univ. de Valencia, Spain)
 John O. Archambeau
 Robert B. Aronson
(medical associate)
 John L. Bateman
(assigned from National Inst. of Health)
 Donald C. Borg
 Wenceslao Calvo
(on leave from Facultad de Medicina, Univ. de Valencia, Spain)
 Donald H. Clifford
(on leave from Univ. of Minnesota)
 William H. Clode*
(medical associate)
 Stanton H. Cohn
 Spencer L. Commerford
 Robert A. Conard, Jr.
 Hans Cottier
(on leave from Univ. of Bern, Switzerland)
 George C. Cotzias
 Eugene P. Cronkite
 Lewis K. Dahl
 Paul D. Doolan
(assigned from Univ. of California)
 Ruth M. Drew
 Otho D. Easterday*
 Ludwig E. Feinendegen
(Damon Runyon Memorial Fund Fellow)
 Samuel Fine*
(medical associate)
 Theodor M. Fliedner
(medical associate)
 Conrad T.O. Fong*
(assigned from New York Univ. School of Medicine)
 Edith M. Forsyth
 Richard L. Golden
(medical associate)

Richard A. Hammerstrom
(medical associate)
Lawrence V. Hanks
Max W. Hess
(medical associate)
Walter L. Hughes
Cornelius R. Jansen
(assigned from Atomic Energy Board,
Union of South Africa)
Hooney C. Kahng
(Worcester Foundation Fellow)
Henry J. Koch
(medical associate)
Louis C. Lax
(medical associate)
Stuart W. Lippincott
Robert A. Love
George C. Mather
(medical associate)
Nicholas Odartchenko
(medical associate)
Paul S. Papavasiliou
(medical associate)
Bengt H. Persson*
(medical associate)
Edwin A. Popenoe
David C. Price*
(medical associate)
Kanti R. Rai
(medical associate)
James S. Robertson
Eckart Schackow
(medical associate)
Lewis M. Schiffer
(medical associate)
Brigitte Schultze
(medical associate)
Irving L. Schwartz*
Yukio Shigeta
(medical associate)
Walton W. Shreeve
Sanford C. Spraragen*
(medical associate)
Elmer E. Stickley*
Richard D. Stoner
Edgar A. Tonna
Cornelius J. van der Merwe
(assigned from Atomic Energy Board,
Union of South Africa)
Donald D. Van Slyke
William Wolins*
Y. Lucas Yamamoto*
(medical associate)

Nuclear Engineering Department

Warren E. Winsche, *Chairman*
George Adler
Leonard B. Adler
Sidney J. Altschuler
(graduate student from Columbia Univ.)

*Terminated before July 1, 1962.

Biagio Arcipiani*
(postdoctoral appointment)
Seymour Aronson
Clemens Auerbach
Allan Auskern
David S. Ballantine
William N. Bishop
Fritz Bloch
Joseph S. Bryner
Charles Carlson*
Albert W. Castleman
John Chen
Jack Chernick
Joe G.Y. Chow
John T. Clarke
Carol H. Collins
(postdoctoral appointment)
Noel R. Corngold
Anita J. Court
Jacobus W.L. de Villiers*
(assigned from Atomic Energy Board,
Union of South Africa)
Russell N. Dietz
(graduate student from
Polytechnic Inst. of Brooklyn)
Robert F. Doering
Kenneth W. Downes
Orrington E. Dwyer
James J. Egan
José F. Evangelista*
(assigned from the Brazilian Army)
Stephen W. Feldberg
(postdoctoral appointment)
Harmon L. Finston
Albert H. Fleitman
Althea Glines
Murray D. Goldberg
David H. Gurinsky
Floris J. Haasbroek*
(on leave from
Dept. of Agricultural Technical Services,
Union of South Africa)
Loramus P. Hatch
Robert L. Hellens
Joseph M. Hendrie
John J. Herbst
(graduate student from New York Univ.)
Raymond J. Heus
Frank B. Hill
Manny Hillman
Henry C. Honeck
Frederick L. Horn
Robert J. Isler
Kazumi Iwamoto*
(on leave from Japan Atomic Energy
Research Inst., Tokai-mura)
Richard Johnson
Sheldon Kalish
Fumihiko Kamijo*
(assigned from Showa Deko K.K., Japan)
Otto F. Kammerer

Herbert M. Katz
John J. Kelsch
Carl J. Klamut
Juan U. Koppel*
(postdoctoral appointment)
Herbert J.C. Kouts
Theodore J. Krieger
Otto A. Kuhl
Gerald S. Lellouche
Melvin M. Levine
Marco Loeffelholz*
(assigned from Società Ricerche Impianti
Nucleare, Montecatini, Italy)
Donald R. MacKenzie
Bernard Manowitz
Michael W. Maresca*
Frank D. Maslan
Donald J. Metz
Paul A. Michael
Francis T. Miles
Sophie Moore
Leonard Newman
Yasutaka Osawa*
(assigned from Mitsubishi Atomic Power
Industries, Inc., Japan)
Guyon P. Pancer
James P. Phelps
Charles E. Porter, Jr.
James R. Powell, Jr.
Glenn A. Price
Surendra N. Purohit
(postdoctoral appointment)
Venkatesalu Rajagopal*
(postdoctoral appointment)
Rajiengar Ranganathan
(graduate student from New York Univ.)
Chad J. Raseman
James J. Reilly, Jr.
Guillermo H. Ricabarra*
(on leave from National Atomic Energy
Commission, Argentina)
María B. Ricabarra*
(on leave from National Atomic Energy
Commission, Argentina)
Powell Richards
Francis X. Rizzo
William A. Robba*
Ranier Rolle
(assigned from Atomic Energy Board,
Union of South Africa)
Jerome Sadofsky
Francis J. Salzano
César A. Sastre
Clifford H. Scarlett
Peter Schofield*
(assigned from AERE, Harwell, England)
Donald G. Schweitzer
Eric Sermet-Chadwick
(NATO Fellow)
Charles S. Shapiro
(graduate student from Syracuse Univ.)

Thomas V. Sheehan
 Rudolph Sher*
 Robert M. Singer
 Louis M. Slater
 John L. Speirs
 Govindasharama Srikantiah
(graduate student from Columbia Univ.)
 Louis G. Stang, Jr.
 Eugene Starr*
 John R. Stehn
 Meyer Steinberg
 Gerald Strickland
 John E. Suich
*(graduate student from
 Massachusetts Inst. of Technology)*
 Herbert Susskind
 Hiroshi Takahashi*
(postdoctoral appointment)
 Stelvio Tassan*
(on leave from New York Univ.)
 Yoel Tikochinsky
*(graduate student from
 the Hebrew Univ., Israel)*
 Walter D. Tucker
 Nazakat Ullah
(postdoctoral appointment)
 John D. Van Norman
 Derek Verdin
*(assigned from United Kingdom
 Atomic Energy Authority, England)*
 Jean I. Wagner
 Charles H. Waide
 John R. Weeks
 Eugene V. Weinstock
 Jerome Weiss
 Henry H. Windsor
 Edward Wirsing, Jr.
 Richard H. Wiswall, Jr.
 Emanuel Yellin*
(postdoctoral appointment)
 Wen-Shi Yu
*(graduate student from
 Polytechnic Inst. of Brooklyn)*

Physics Department

George H. Vineyard, *Chairman*
 Fernando Agullo-López
*(on leave from Junta de Energía Nuclear,
 Madrid, Spain)*
 Ian J.R. Aitchison
(postdoctoral appointment)
 David E. Alburger
 Louis C.R. Alfred*
(postdoctoral appointment)
 Ismael Almodovar
(assigned from Puerto Rico Nuclear Center)
 Harvey A. Alperin
*(assigned from
 U.S. Naval Ordnance Laboratory)*

Richard Arndt
(postdoctoral appointment)
 Frederick Ayer II*
 Charles P. Baker
 Winslow F. Baker
 Mirza A.P. Bég
(postdoctoral appointment)
 Brother Austin Bernabei
(graduate student from New York Univ.)
 Jeremy Bernstein
 Luciano Bertanza
*(on leave from
 Inst. di Fisica, Univ. di Pisa, Italy)*
 Martin Blume
 Henri Boutin
(on leave from Pennsylvania State Univ.)
 Veljko Brajovic*
*(on leave from
 Univ. of Belgrade, Yugoslavia)*
 Dermot J. Bredin
 Jean-Claude Brisson
*(on leave from
 Commissariat à l'Énergie Atomique,
 France)*
 Violette Brisson*
*(on leave from Centre National
 de la Recherche Scientifique, France)*
 P. Jane Brown
(postdoctoral appointment)
 Gunter Brunhart
(postdoctoral appointment)
 Robert E. Chrien
 Eugene L. Church
(assigned from Frankford Arsenal)
 Victor W. Cohen
 George B. Collins
 Philip L. Connolly
 Rodney Cool
 Maurice N. Cotter*
(postdoctoral appointment)
 Ernest D. Courant
 Bernd Crasemann
(on leave from Univ. of Oregon)
 Ray R. Crittenden
 Bernard B. Culwick
 Arthur C. Damask
(assigned from Frankford Arsenal)
 Horace R. Danner
 Paul C. DeCelles
(postdoctoral appointment)
 Edward der Mateosian
 George J. Dienes
 Achiel J. Deruytter
*(assigned from
 Centre d'Étude de l'Énergie Nucléaire,
 Belgium)*
 Philip J. Duke*
*(assigned from National Inst. for Research
 in Nuclear Science, England)*
 Loyal Durand III*
(postdoctoral appointment)

Frederick R. Eisler
 Guy T. Emery
 Anne Englert*
*(on leave from
 European Research Associates, Belgium)*
 Cavid Erginsoy
(on leave from IAEA, Vienna, Austria)
 Nikolaus F. Fiebiger*
*(on leave from
 Univ. of Frankfurt, West Germany)*
 Ted B. Flanagan*
 Kenneth J. Foley
(postdoctoral appointment)
 William B. Fowler
 B. Chalmers Frazer
 Tadao Fujii
 Francisco Eiichi Fujita
*(on leave from Japan Atomic Energy
 Research Inst., Tokai-mura)*
 Maurice Gailloud
*(assigned from
 Laboratoire de Recherches Nucléaires,
 École Polytechnique, Switzerland)*
 Arthur F. Garfinkel*
(graduate student from Columbia Univ.)
 Allen N. Goland
(assigned from Watertown Arsenal)
 Gertrude S. Goldhaber
 Samuel A. Goudsmit
 Konstantin Goulianos
(graduate student from Columbia Univ.)
 Vasken B. Hagopian
*(graduate student from
 Univ. of Pennsylvania)*
 Edward L. Hart
 Leland J. Haworth
(on leave to the AEC)
 Arthur Herschman
(on leave from Worcester Polytechnic Inst.)
 John W. Hess
(assigned from Picatinny Arsenal)
 Nguyen C. Hien
(assigned from Carnegie Inst. of Technology)
 John Hornbostel
 Paul V.C. Hough
 Joseph Jach
(assigned from Picatinny Arsenal)
 Adeshwar P. Jain
(postdoctoral appointment)
 Edgar W. Jenkins
 Gary L. Jensen
(graduate student from Univ. of Michigan)
 Jovan V. Jovanovic
(postdoctoral appointment)
 H.K. Alan Kan*
(assigned from Picatinny Arsenal)
 Walter R. Kane
 David T. Keating
 Michael A. Kemp
*(assigned from National Inst. for Research
 in Nuclear Science, England)*

*Terminated before July 1, 1962.

Ottmar C. Kistner
 Charles Kocher*
(postdoctoral appointment)
 Joshua K. Kopp
 Saul Krasner
(assigned from Picatinny Arsenal)
 Thaddeus F. Kycia
 Ronald E. Larsen
(assigned from Frankford Arsenal)
 Paul W. Levy
 Angela C. Li
(graduate student from New York Univ.)
 Seymour J. Lindenbaum
 Robert I. Louttit
 William A. Love
 James Lowe*
(postdoctoral appointment)
 Dierk Luers*
(on leave from Max Planck Inst., Munich, West Germany)
 Italo Mannelli
(assigned from Univ. of Pisa, Italy)
 James E. Mapes
(assigned from Picatinny Arsenal)
 Harvey Marshak
 Joseph P. Martin*
(postdoctoral appointment)
 Clyde L. McClelland
 Werner A.W. Mehlhop
(assigned from Univ. of California)
 Nariman B. Mistry
(graduate student from Columbia Univ.)
 Inder S. Mittra
(on leave from Punjab Univ., India)
 Sergio Monaro
(on leave from National Inst. of Nuclear Physics, Italy)
 Giancarlo Moneti
(on leave from Univ. of Rome, Italy)
 John A. Moore
 Thomas I. Moran
(postdoctoral appointment)
 Thomas W. Morris
 Bernard Mozer
 Basharat A. Munir*
(on leave from Florida State Univ.)
 Vernon W. Myers*
(on leave from Pennsylvania State Univ.)
 Robert Nathans
 James A. Niederer*
 Kazuyuki Ogawa
(assigned from Tokyo Shibaura Electric Co., Ltd., Japan)
 John W. Olness
(assigned from Wright-Patterson Air Force Base)

Carl M. Olsmats*
(postdoctoral appointment)
 Jean Oostens
(assigned from Inst. Interuniversitaire des Sciences Nucléaires, Belgium)
 Kare Ottes
(on leave from Aktiebolaget Atomenergi, Stockholm, Sweden)
 Satoshi Ozaki
 Harry Palevsky
 Robert B. Palmer
 Simon Pasternack
(Editor of The Physical Review)
 Siegfried Penselin*
(on leave from Univ. of Heidelberg, West Germany)
 Robert H. Phillips
 Stanley J. Pickart
(assigned from U.S. Naval Ordnance Laboratory)
 Eric Pickup*
(on leave from National Research Council, Canada)
 Hans Postma*
(on leave from Universities of Leiden and Groningen, Netherlands)
 Albert G. Prodell
 David C. Rahm
 Ralph R. Rau
 A. Lincoln Read
(postdoctoral appointment)
 Kenneth F. Riley
(postdoctoral appointment)
 Donald K. Robinson
 Joseph H. Rosolowski
(on leave from General Electric Co.)
 Henry Ruderman
(postdoctoral appointment)
 John J. Russell, Jr.
 George J. Safford
(assigned from Columbia Univ.)
 Vance L. Sailor
 Edward O. Salant
 Nicholas P. Samios
 James R. Sanford
(assigned from Yale Univ.)
 Robert I. Schermer
 Marcel R. Schmorak
(postdoctoral appointment)
 Ivan Schroder
(graduate student from Columbia Univ.)
 Melvin Schwartz
(assigned from Columbia Univ.)
 Arthur Z. Schwarzschild
 Madan L. Sehgal
(postdoctoral appointment)
 Ralph P. Shutt
 Ian O. Skillicorn
(postdoctoral appointment)

Joseph E. Smith
 Hartland S. Snyder
(deceased May 2, 1962)
 Morton M. Sternheim
(postdoctoral appointment)
 Rudolph Sternheimer
 Andrew W. Sunyar
 John B. Swan*
(on leave from Univ. of Western Australia, Nedlands)
 Michael J. Tannenbaum
(graduate student from Columbia Univ.)
 Alan Thorndike
 Peter D. Townsend
(assigned from Picatinny Arsenal)
 T. Laurence Trueman
(postdoctoral appointment)
 Frank Turkot
 Joseph B. Vise
(graduate student from Columbia Univ.)
 Yoshinisa Wakuta*
(on leave from Kyushu Univ., Japan)
 Ernest K. Warburton
 Medford S. Webster
 Joseph Weneser
 James Westgard
(graduate student from Syracuse Univ.)
 Gian Carlo Wick
 William J. Willis
(on leave to CERN, Geneva, Switzerland)
 Sanford E. Wolf
(on leave from Brandeis Univ.)
 Yehuda Wolfson
(on leave from Weizmann Inst. of Science, Israel)
 Sukeyasu S. Yamamoto
 Taiji Yamanouchi
(assigned from Univ. of Rochester)
 Thomas J. Ypsilantis
 Luke C.L. Yuan
 Gus T. Zorn
 Martin S. Zucker
 Crtomir Zupancic
(postdoctoral appointment)

Reactor Division

Robert W. Powell, *Head*
 John J. Floyd
 Gerald C. Kinne
 Frederick J. Morse
(graduate student from Rensselaer Polytechnic Inst.)
 Charles L. Osborne
 Jack E. Phillips
 dePuyster G. Pitcher
 Harry H. Steinhauer, Jr.*
(assigned from Rensselaer Polytechnic Inst.)
 Dudley Thompson

*Terminated before July 1, 1962.