TEST OF THE VALENCY MODEL OF NEUTRON CAPTURE IN MO-92 + NEUTRONO. A. Wasson * and G. G. Slaughter Oak Ridge National Laboratory **

Abstract: Resonant neutron capture ray spectra were measured in Mo-92 to test the Valency Model. The dominant strength of the primary rays feeding the $1/2^+$ first excited state suggests that $p \rightarrow s$ transitions are enhanced.

Renewed interest in the applicability of the single particle model to radiative neutron capture has occurred following the observation of the Brookhaven Group (1) that the ray spectra from p-wave neutron capture in Mo-98 and Mo-92 are consistent with this model. As a further test of the model, more detailed resonant neutron capture γ ray measurements on the target nucleus Mo-92 were done at the Oak Ridge Electron Accelerator. The γ -ray spectra were recorded with a Ge (Li) diode using the 10 m flight path and a 30 nsec beam burst. Useful spectra were obtained for neutron energies from 300 eV to 100 keV with a total of 23 resonances resolved for $E_n < 25$ keV. A total of 14 resonances are assigned to p-wave capture on the basis of either a strong 8066 keV γ ray transition to the $5/2^+$ ground state or no interference in the neutron transmission measurement (2). Similarly 5 resonances are assigned to S-wave capture with the parity of the remainder undetermined.

A total of 12 rays with energies exceeding 5.0 MeV were analysed and found to account for most of the total radiation width for most resonances. Absolute γ ray widths were determined using the 4.9 eV gold resonance as a standard. Nearly all of the low energy γ ray intensities result from two-step γ ray decays from the capturing state to the ground state. Eight primary γ rays are accompanied by a low energy transition to the ground state. The low energy γ ray intensity is correlated with the primary intensity which shows that the low energy γ ray intensities are not useful for determination of resonance spins. Also a new level at 2141 keV excitation was identified for both the primary and secondary γ rays.

The valency model (3) predicts the γ ray spectra to be dominated by transitions to the $5/2^+$ ground state and $1/2^+$ first excited state for p $3/2$ resonances and by transitions to the $1/2^+$ state for p $1/2$ resonances. This prediction is fulfilled for most of the resonances. The detailed

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

Fig. 1. Predicted and measured ray intensities

results for two $P\ 3/2$ resonances are shown in Fig. 1. The intensities shown are normalized to the summed intensities of the 12 γ rays. On the left are the experimental widths while the valency model predictions are on the right. For the remaining resonances the agreement is less although the 7129 keV transition to the $1/2^+$ first excited state remains dominant. Although the valency model components should be more important for the resonances with the largest reduced neutron widths, the 12.8 keV resonance has a small neutron width, while resonances with much larger widths show less agreement with the model. The large strength of the transitions to the ground and first excited states is indicated in Fig. 2. Here the average γ ray intensities for the three classes of resonances are shown. It is also of interest to note that the M1 γ ray intensities from the S-wave resonances are as large as the E1 intensities from the p-wave resonances. This near equality is also verified by the near equality in the total radiation widths for S- and p-wave resonances (2). The continuing strength of the highest energy γ rays for the un-resolved neutron energy region above 20 keV is shown in Fig. 3. The dominance of the 7129 keV ray is a violation of the statistical model predictions and leads support to the observation of Lane (4) that the $p \rightarrow s$ transitions are more single particle than the $p \rightarrow d$ transitions.

These preliminary results indicate that the single particle components are significant in the γ ray decay widths for resonant neutron capture in Mo-92.

Fig. 2. Average γ -ray intensities for three classes of resonances

Fig. 3. Average γ ray intensities for two neutron energy intervals

* Guest scientist from Brookhaven National Laboratory

** Operated by Union Carbide Corporation for the USAEC.

1. S. F. Mughabghab, R. E. Chrien, O. A. Wasson, G. W. Cole and M. R. Bhat Phys. Rev. Letters 26, 1118 (1971)
2. O. A. Wasson, B. J. Allen, R. R. Winters, R. L. Macklin and J. A. Harvey Bull. Amer. Phys. Soc. 17, 556 (1972)
3. J. E. Lynn, The Theory of Neutron Resonance Reactions, (Clarendon Press, Oxford, 1968)
4. A. M. Lane (Private communication to S. F. Mughabghab)