MAR 24 1965

83975

83975

MASTER

A. (d,p-f) ANGULAR CORRELATION STUDY OF FISSION BARRIER
TRANSITION STATES: ENERGY GAP OF A HIGHLY DEFORMED NUCLEUS*

by

Harold C. Britt, William R. Gibbs, James J. Griffin, and Richard H. Stokes

University of California
Los Alamos Scientific Laboratory
Los Alamos, New Mexico

NOT FOR PUBLIC RELEASE - OFFICIAL
DISTRIBUTION MAY BE MADE. OFFICIAL
REQUESTS MAY BE FILLED. REPORT
CONTAINS NOTHING OF PATENT INTEREST.
PROCEDURES ON FILE IN RECEIVING
SECTION.

Ed. N.B. Running Head:

ENERGY GAP IN TRANSITION STATE SPECTRUM

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

*Work done under the auspices of the U. S. Atomic Energy Commission.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

ABSTRACT

The (d,p) reaction was used to produce fissioning nuclei with excitations from the fission threshold to 5 MeV above. As a function of excitation the angular correlation of the fission fragments and the reaction protons has been measured. A theoretical correlation function is developed to permit values of the parameter K_0^2 to be derived from the observed angular anisotropy. For a Pu^{239} target the values obtained for K_0^2 show a stepwise increase with increasing excitation. The largest step occurs at ~ 2.6 MeV above the fission threshold, and is interpreted as the beginning of quasi-particle excitations of the highly deformed fissioning nucleus. On this basis the pairing energy gap appears to be significantly increased for large nuclear deformations. Also, K_0^2 is observed to change at ~ 0.7 and ~ 1.6 MeV above threshold. It is suggested that these energies are the onset of collective vibrational transition states through which fission occurs.

1. INTRODUCTION

The (d,p) reaction has been used in a previous experiment¹ to measure the probability of fission as a function of excitation energy. It was shown that fission thresholds can be measured for target nuclei which undergo thermal neutron-induced fission. For these nuclei, the (d,p) reaction produces even-even fissioning nuclei with excitation energies near the fission threshold; a region of excitation energy which is inaccessible to neutron-induced fission experiments.

Another property of the (d,p) reaction is that fissioning nuclei can be formed in states of moderately high angular momentum, so that it is possible to investigate the angular correlations of the fragments at excitation energies near the fission threshold. For high excitations, the results of anisotropy theory² for fission induced by particle capture, show that the anisotropy is an increasing function of the average angular momentum of the fissioning nucleus and a decreasing function of the amount of excitation above the fission threshold. Because the (d,p) reaction can transfer considerable angular momentum and at the same time induce fission with minimum excitation, large correlation effects may be anticipated. A measurement of these correlations as a function of excitation energy reveals information on the transition states of the nucleus as it passes over the fission barrier.² These transition states consist of collective and particle excitations of the fissioning nucleus at saddle point deformation. Of particular interest are measurements for fissioning nuclei throughout a band of several MeV above threshold, because in this region superfluidity

effects due to nuclear pairing³ are most important. At these low excitation energies it is possible to observe effects which distinguish the finite isolated nuclear superfluid⁴ from the infinite systems which are usually considered.^{5,6} In particular, it was expected that the angular correlation should show a discontinuity at the excitation energy corresponding to the onset of two quasiparticle excitations. In the threshold region, evidence for the opening of new transition states has been seen in the previous measurements¹ of fission barrier penetrability. In the penetrability data, because of competition from neutron emission and the effect of deuteron breakup, only a range of about 1.6 MeV was accessible to clear interpretation. The present anisotropy measurements are insensitive to the above two obscuring effects and permit unambiguous interpretation over a wider excitation range.

In the present experiment, measurements were made of the angular correlations of fragments emitted in the fission of Pu^{240} and U^{234} . The results clearly show the expected nuclear pairing effects, and also show evidence for low lying collective excitations near the fission threshold.

2. EXPERIMENTAL PROCEDURES

The experimental method was similar to that used in the previous determination of fission thresholds.¹ Semiconductor fission detectors were situated in a plane defined by the incoming 14.9-MeV deuteron beam and the observed reaction protons. The proton detector consisted of a semiconductor transmission ΔE detector followed by a semiconductor E detector in which the

protons stop. The pulses from the two detectors were used with an analog computer⁷ to distinguish protons from other emitted particles.

A fast-slow coincidence system was used to identify protons in coincidence with fission. A fast coincidence with $\tau = 50$ nsec was required between the fission detector and the E detector, and a slow coincidence was required with the proton identification system. A pulse from the coincidence system was used to gate a multichannel analyzer which recorded a pulse height proportional to the sum of the pulses in the AE and E detectors (i.e., proportional to the proton energy). In one set of measurements, three fission detectors were used simultaneously to determine the coincident counting rate for three different fission angles. In this case, three coincidence systems were used to gate three separate 100-channel blocks of a 400-channel analyzer.

During the measurements a simultaneous determination of the chance coincidence contribution to the spectra was made. This was done using another coincidence system which was identical to that used for the true spectra except that the signal from the E detector was delayed by about 160 ns (2 cyclotron periods) from true coincidence. The chance spectra were also recorded in a 100-channel block of the analyzer. In the measurement where three fission detectors were used, only one of the chance spectra was recorded and for the other two detectors the total number of chance counts was recorded on scalars. In the analysis of the data, it was assumed that the shape of the chance spectrum was the same for each fission angle. During the measurements, the chance rates

were typically 5% of the true rates.

The deuteron beam which was obtained from the Los Alamos variable energy cyclotron had a mean energy of 14.9 MeV. The beam energy was determined by slowing the deuterons with aluminum absorbers and comparing the solid state detector response of the degraded beam to the response of 5.1-MeV alpha particles. The proton energy scale was calibrated using the elastically scattered deuteron group from Pu and U, and proton groups from the (d,p) reaction on C^{12} and O^{16} . The overall width of the response in the energy measuring channel was 200 keV FWHM. This width includes deuteron beam energy spread, detector response, noise, and kinematic broadening. The targets were prepared by vacuum evaporation of $Pu^{239}O_2$ and $U^{233}O_2$ onto $60 \mu g/cm^2$ carbon backings. The Pu^{239} and U^{233} deposits had thicknesses of 175 and $360 \mu g/cm^2$, respectively.

The data were analyzed by first correcting each spectrum for the chance contribution. The results were transformed to the center-of-mass system of the fissioning nucleus. This transformation was performed using values for the recoil velocity and angle of the fissioning compound nucleus that were calculated from the deuteron energy, and the energy and angle of the proton. The transformation was performed assuming that all the fragments have a single mass and a single energy. The fragment mass was taken as one half the mass of the fissioning nucleus and the fragment energy was taken as the mean kinetic energy measured for thermal neutron-induced reactions forming the same fissioning nuclei.⁸

3. EXPERIMENTAL RESULTS

There were two types of measurements performed to obtain the results presented here. In the first measurement, a single fission detector was used with the Pu^{239} target and measurements were made for various angles of fission detector in order to determine the detailed shape of the angular correlation. This type of measurement is illustrated schematically in Fig. 1. In the second set of measurements, three detectors were placed at fixed laboratory angles of +20, +110, and +155 deg with respect to the beam. With this arrangement, a series of runs were taken to obtain the best possible statistical accuracies on the anisotropy measurements as a function of the proton energy.

The coincident proton energy spectra which were obtained in the measurements with the three fixed detectors are shown in Fig. 2. These spectra are the sum of the coincident spectra measured for the three fission detectors. The spectra shown in Fig. 2 have essentially the same shape as those obtained in the previous measurement.¹ The arrows shown in Fig. 2 represent the definition of the proton energy corresponding to the excitation energy at the fission threshold. These positions for the threshold energy were obtained⁹ by comparing the shapes of the coincident proton energy spectra with the spectra obtained in the previous threshold measurements.¹ As in the previous measurements¹ the threshold is defined as the energy at which the fission probability reached 50% of its first plateau value.¹⁰ These thresholds were then used to obtain the $E^* - E_{\text{th}}$ scale of excitation energy above the fission threshold used in the following figures.

Figure 3 shows the angular correlations obtained in the first measurement with the single fission detector for the Pu^{239} target. In this figure, results are shown for intervals of the excitation energy above the fission threshold of 0-1, 1-2, and 2-3 MeV for laboratory proton angles of 140° and 75° . The center-of-mass angles are measured relative to the classical recoil angle of the fissioning nucleus produced in the (d,p) reaction. The values of these recoil angles for the threshold excitation are indicated on the figure. Both positive and negative angles¹¹ are shown in Fig. 3 in order to demonstrate that within the statistical accuracies the classical recoil angle is the symmetry angle for the angular correlations. The solid lines shown in Fig. 3 represent least squares fits of the experimental results to the function

$$W(\theta) \propto 1 + g_2 P_2(\cos\theta). \quad (1)$$

It is seen that within the experimental uncertainties this function gives an adequate fit to the results. The results shown in Fig. 3 illustrate three general characteristics of the angular correlations: (1) the anisotropies $W(0^\circ)/W(90^\circ)$ decrease by approximately a factor of 2 for every 1 MeV of excitation energy above the threshold, (2) the maximum anisotropy is 2.8 for the 140° proton angle and 2.0 for the 75° proton angle, and (3) as discussed above, the symmetry angle for the angular correlation is close to the classical recoil angle of the fissioning nucleus.

Following the above experiment, measurements were made with much higher statistical accuracy using three fixed fission detectors and col-

lecting data at the three angles simultaneously. In this measurement the proton detector was placed at 140° because of the larger anisotropies and because the theoretical interpretation was more straightforward for a backward proton angle (see section 4). With this arrangement data were taken for both Pu^{239} and U^{233} targets.

The results were fitted to Eq. (1) and the values obtained for g_2 are shown in Fig. 4. In the energy interval 0-5 MeV above the threshold excitation, approximately 1×10^5 and 5×10^4 events were recorded for the Pu^{239} and U^{233} targets, respectively. The Pu^{239} data is therefore statistically superior to that for U^{233} . Figure 3 shows that near threshold the anisotropies for the U^{233} target are approximately a factor of 2 less than for the Pu^{239} target. For this reason also, the Pu^{239} data provides more precise information on the nuclear structure at the fission barrier. These data therefore provide the basis for most of the conclusions in this paper.

4. THEORY

To compute the angular distribution of fission fragments from the (d,pf) reaction, we need the amplitudes (a) for formation of a compound nucleus in a state (J, J_z, λ) when the proton is observed with momentum $\vec{K}_p$ and (b) for fission of the compound nucleus from this compound state with fragments emitted into the solid angle $\hat{\theta}_f$. Then

$$W(\hat{\theta}_f, \vec{K}_p) = \left| \sum_{J, J_z, \lambda} A(\vec{K}_p; J, J_z, \lambda) B(\hat{\theta}_f; J, J_z, \lambda) \right|^2$$

is the angular correlation of the fragments with the proton. A sum over final spin projections and an average over initial spin projections is implicit in all relevant expressions.

To compute the stripping amplitude, we assume that the neutron is captured into a compound state (J, J_z, λ) via a single particle state (l, j, m) coupled to the unexcited core whose angular momentum (I, I_z) is equal to the angular momentum of the target nucleus. Then

$$A(\vec{K}_p; J, J_z, \lambda) = \sum_{l, j, m} A(l, j, m; I, I_z; J, J_z; \vec{K}_p) \gamma_{lj}^{J\lambda} \quad (3)$$

where $\gamma_{lj}^{J\lambda}$ is the reduced width for the compound state (J, λ) and $A(l, j, m; I, I_z; J, J_z; \vec{K}_p)$ is to be computed from an appropriate stripping calculation.

To compute the fragment angular distribution from the compound state (J, J_z, λ) we assume that a fission through a transition state (J, J_z, K) emits fragments into the angle $\hat{\theta}_f$ with an amplitude given by the symmetric top wave function, $\vartheta_{J_z K}^J(\hat{\theta}_f)$, which describes the orientation of the fissioning nucleus as it passes over the fission barrier,² and that the amplitude for the compound state (J, λ) to decay through a transition state with projection K of angular momentum along the nuclear symmetry axis is $b_K^{J\lambda}$. Then

$$B(\hat{\theta}_f; J, J_z, \lambda) = \sum_K b_K^{J\lambda} \vartheta_{J_z K}^J(\hat{\theta}_f) \quad (4)$$

In the present experiment the sum over λ in equation (2) involves many states of a given J -value because of the experimental resolution

(~ 200 keV) and the narrow spacing of the relevant compound levels (~ 100 eV). One is therefore justified in the approximation

$$\sum_{\lambda\lambda'} b_K^{J\lambda*} b_{K'}^{J'\lambda'} \gamma_{\ell j}^{J\lambda*} \gamma_{\ell' j'}^{J'\lambda'} = \sum_{\lambda} \left| b_K^{J\lambda} \right|^2 \left| \gamma_{\ell j}^{J\lambda} \right|^2 \delta_{KK'} \delta_{\ell\ell'} \delta_{jj'} \delta_{JJ'} \quad (5)$$

$$= \overline{\left| b_K^J \right|^2} \overline{\gamma^2} \delta_{KK'} \delta_{\ell\ell'} \delta_{jj'} \delta_{JJ'} N(J, \ell, j).$$

Where $N(J, \ell, j)$ is the number of single particle states with quantum numbers (J, ℓ, j) and $\overline{\gamma^2}$ is the average reduced width for such single particle states.

We shall assume that from each single particle state (ℓ, j) every J state allowed by angular momentum coupling to the target spin is available. Then $N(J, \ell, j) = N_1(J, j) \cdot N_2(\ell, j)$ where $N_1(J, j) = \Delta(j, I, J)$ and $N_2(\ell, j)$ is still to be specified on physical grounds. $\Delta(abc)$ is one if a, b, c satisfy the triangular inequalities; zero, otherwise. The angular distribution is then a simple sum over products of density matrices:

$$W(\vec{K}_p, \hat{\theta}_f) \propto \sum_{J, \ell j, K} \overline{\left| b_K^J \right|^2} N_1(J, j) N_2(\ell, j) \left\{ \sum_{J_z J'_z} E_{(d, p)}^{J_z J'_z}(\vec{K}_p; J, \ell j) E_{fission}^{J_z J'_z}(\hat{\theta}_f; J, K) \right\} \quad (6)$$

where

$$E_{fission}^{J_z J'_z}(\hat{\theta}_f; J, K) = \mathcal{D}_{J_z, K}^{J*}(\hat{\theta}_f) \mathcal{D}_{J'_z, K}^J(\hat{\theta}_f) \quad (7)$$

and

$$E_{(d, p)}^{J_z J'_z}(\vec{K}_p; J) = \sum_{m m'} A^*(\ell j m; I, I_z; J, J_z; \vec{K}_p) A(\ell j m'; I, I_z; J, J'_z; \vec{K}_p) \quad (8)$$

is the usual density matrix for stripping a neutron (l, j) into a single particle state J .

Equation (6) states that the full angular correlation function is a sum over angular correlations for various possible j, l, J, K values weighted with the relative probability, $|b_K^J|^2 \cdot N(J, l, j)$, that the compound nucleus is formed through a state (j, l, J) and decays through a transition state, K .

To reduce equation (6) to easily calculable form we first expand

$E_{fission}^{J_z J'_z}$ in statistical tensors:

$$E_{fission}^{J_z J'_z} = \sum_{Lm} Y_L^m(\theta_f, \varphi_f) \sqrt{4\pi(2L+1)} (LJOK|LJJK) (LJM_z' | LJJ_z) \frac{1}{8\pi^2} \quad (9)$$

The sum over K is now quite simple, and if one assumes $\overline{|b_K^J|^2} = \overline{|b_{-K}^J|^2}$, it has the property:

$$\sum_K \overline{|b_K^J|^2} (LJOK|LJJK) = 0, \quad L \text{ odd} \quad (10)$$

so that only even L need be considered.

The analogous expansion of $E_{(d,p)}^{J_z J'_z} (l, j, J; \vec{K}_p)$ is given by Satchler¹² for plane waves and by Huby, Refai, and Satchler¹³ for distorted waves. The notation used in the present paper differs from that given by these authors.

$$E_{(d,p)}^{J_z J'_z} (l, j, J; \vec{K}_p) = \sum_{L'M'} (L'JM_z' | L'JJ_z) \eta_{L'}(l, j, J) \rho_{L', l}^{M'}(\vec{K}_p) \quad (11)$$

where,

$$\eta_L(I l j J) = W(L l j \frac{1}{2}; l j) W(L j J I; j J) [2J + 1] (2l + 1)^{\frac{1}{2}} (2j + 1) (2L + 1) \quad (12)$$

and

$$\rho_{Ll}^M(\vec{K}_p) = \sum_{m, m'} (L l M m | L l l m') \beta_l^m(\vec{K}_p) \beta_l^{m'}(\vec{K}_p)^* \quad (13)$$

when no spin-orbit interaction is included. The double sum over J_z, J'_z can be rearranged into the Clebsch-Gordan orthogonality relationship, proportional to $\delta_{LL'}, \delta_{MM'}$. Thus L' is equal to L and, therefore, assumes only even values. We note also that

$$\rho_{ol}^o = \sum_m |\beta_l^m(\vec{K}_p)|^2 = \sigma_l(\vec{K}_p) \quad (14)$$

is proportional to the stripping cross section for an l -wave neutron, and that, for even L ,

$$(L l o o | L l l o) W(L l j \frac{1}{2}; l j) = \Delta(l \frac{1}{2} j) \cdot \Delta(L l l) \cdot (L j o \frac{1}{2} | L j j \frac{1}{2}) [(2l + 1)(2j + 1)]^{-\frac{1}{2}} \quad (15)$$

Then by defining $\tilde{\rho}$ by means of

$$\rho_{Ll}^M(\vec{K}_p) = (L l o o | L l l o) \sigma_l(\vec{K}_p) \tilde{\rho}_{Ll}^M(\vec{K}_p) \quad (16)$$

we obtain

$$W(\vec{K}_p, \hat{\theta}_f) \propto \sum_{LM} \sum_{l j J K} (2L + 1)^{-\frac{1}{2}} g(j J K) \cdot \Delta(l \frac{1}{2} j) \sigma_l(\vec{K}_p) (2J + 1) \cdot \tilde{\rho}_{Ll}^M(\vec{K}_p) \overline{b_K^J}^2 \quad (17)$$

$$\cdot N_1(J, j) N_2(l, j) Y_L^{M*}(\hat{\theta}_f)$$

where

$$g_L(jJK) = (2L+1)(LJK|LJK)[(Lj0\frac{1}{2}|Lj\frac{1}{2})] W(LjJ; jJ)[(2j+1)(2J+1)]^{\frac{1}{2}} \quad (18)$$

Equation (17) can now be reduced by the following summations over l , J and K :

$$\frac{1}{\sum_{l,j} \sigma_l(\vec{K}_p) N_2(l,j)} \sum_l \sigma_l(\vec{K}_p) N_2(l,j) \tilde{\rho}_{Ll}^M(\vec{K}_p) = d_{Lj}^M(\vec{K}_p) \quad (19)$$

$$\frac{\sum_j N_1(J,j)(2J+1) \sum_K g_L(jJK) \overline{|b_K^J|^2}}{\sum_{j,j} N_1(J,j)(2J+1) \sum_K \overline{|b_K^J|^2}} = g_L(j;I,\alpha) \quad (20)$$

where α denotes one or more parameters describing the physical assumptions to be made concerning $\overline{|b_K^J|^2}$ and $N_1(J,j)$. Then the final result is

$$W(\vec{K}_p, \hat{\theta}_f) = (4\pi)^{\frac{1}{2}} \sum_{L,M,j} (2L+1)^{-\frac{1}{2}} g_L(j;I,\alpha) d_{Lj}^M(\vec{K}_p) Y_L^{M*}(\hat{\theta}_f). \quad (21)$$

To obtain numerical results, one has now to specify the physical assumptions to be made and carry through the computation of $g_L(j;I,\alpha)$ and $d_{Lj}^M(\vec{K}_p)$. We have made the simplest possible assumptions concerning the available single particle neutron states; namely, that all levels in the major oscillator shell, $N = 7$, are available with equal probability at every excitation energy. This assumption is reasonable (a) because of the fact that the stable quadrupole deformation of the target results in significant mixing of the spherical levels so that various spherical

eigenfunctions are already admixed into each deformed single particle state, and (b) more than one single particle state is being excited at any given experimental excitation energy, and finally (c) because at the excitation energies involved (~ 5 MeV) a given single particle state is mixed significantly into compound states⁽¹⁴⁾ within an energy $W \approx 2$ MeV, where W is the imaginary part of the neutron optical potential.⁽¹⁵⁾ On all three counts a given spherical single particle state will be excited over a wide range of experimental excitation energies, tending thereby to make applicable at every excitation energy the average properties of the whole collection of such states. In practice, this assumption implies that every odd l state ($1 \leq l \leq 7$) is available with equal probability, so that $N_2(l, j) = \Delta(l, j) \times \text{constant}$. Since l is assumed to be odd only, there is associated with each value of j a unique value of l . Thus we may define $\bar{\sigma}_j = \sigma_l(j)$. The results would not be significantly modified if some (or all) even l values were included. The results are therefore not as restrictive as the odd- l assumption might suggest.

The transition state probability, $\overline{|b_K^J|^2}$, is assumed to be proportional to the number of transition states available at the fission barrier at the specified excitation energy. For a Fermi-gas description of the transition state level density, one would have

$$\overline{|b_K^J|^2} \propto \exp - \frac{\hbar^2}{2I} \left\{ \frac{1}{J_{\perp}} [J(J+1) - K^2] + \frac{1}{J_{\parallel}} K^2 \right\} \quad (22)$$

For the nucleus highly deformed to the fission barrier, one expects

$\mathcal{F}_\perp \gg \mathcal{F}_\parallel$ so that the form

$$|\overline{b_K^J}|^2 \propto \exp - \frac{K^2}{2K_0^2} \quad (23)$$

is an adequate first approximation.⁽¹⁶⁾ Here K_0^2 is an energy dependent parameter which will be fit to experiment.

If the nuclear level density deviates from a simple Fermi gas picture,⁽⁶⁾ then K_0^2 is expected to show deviations from the Fermi gas prediction. It is in fact the dependence of K_0^2 on excitation energy which is the primary object of the present experiment.

With these assumptions the coefficient $g_L(j;I;K_0^2)$ can be computed and there remains the computation of $\tilde{p}_{Ll}^M(\vec{K}_p)$.

To compute $d_{Lj}^M(\vec{K}_p)$ we first consider the plane wave approximation for the stripping process in which only a single value of j is involved. Then apart from constant factors the Butler amplitude (using surface evaluation) is

$$\beta_l^m = j_l(\kappa R) Y_l^m(\theta_\kappa, \varphi_\kappa) (2l+1)^{-\frac{1}{2}} \quad (24)$$

where θ_κ and φ_κ are the angles of the recoil momentum vector $\vec{\kappa} = \vec{K}_D -$

$\frac{M_I}{M_F} \vec{K}_p$. Then

$$\rho_{Ll}^M = j_l^2(\kappa R) (Ll00|Ll10) (2L+1)^{-\frac{1}{2}} Y_L^M(\theta_\kappa, \varphi_\kappa) \quad (25)$$

and

$$d_{Lj}^M = \left(\frac{4\pi}{2L+1} \right)^{\frac{1}{2}} Y_L^M(\theta_\kappa, \varphi_\kappa) \quad (26)$$

Then

$$W(\theta_p, \varphi_p; \theta_f, \varphi_f) = \sum_L g_L(I; K_0^2) \cdot P_L(\cos \tilde{\theta}_f) \quad (27)$$

where $\tilde{\theta}_f$ is measured from the recoil direction. The quantity $g_L(I, K_0^2)$ is defined as

$$g_L(I, K_0^2) = \sum_j g_L(j; I, K_0^2) \bar{\sigma}_j / \sum_j \bar{\sigma}_j \quad (28)$$

One thus obtains a very simple angular correlation, exhibiting symmetry about the nuclear recoil direction.

In general, the plane wave approximation is a poor one for stripping to heavy nuclei. The exception is the angular correlations it predicts for proton angles, $140^\circ \leq \theta_p < 180^\circ$. For angular properties of such back-angle processes, the plane wave theory agrees quite well with the more complicated distorted wave theory, although the cross sections $\bar{\sigma}_l$ do not agree at all. Thus for comparison with the present data ($\theta_p = 140^\circ$) plane wave theory suffices for calculating $\bar{p}_{Ll}^M$, whereas more accurate DWBA values must be used for $\bar{\sigma}_j(K_p)$.

To demonstrate this we take $g_4(I, K_0^2) \ll g_2(I, K_0^2)$ for a given value of J , j , and l . The relative magnitude of g_4 and g_2 prevails in the present experimental data and is corroborated by calculation, which predicts $g_4 \approx 0.1 g_2$. The equation (21) becomes:

$$W(\theta_p, \varphi_p; \theta_f, \varphi_f) = 1 + g_2(j; I, K_0^2) \sum_{M=-2}^{+2} d_{2j}^M(K_p) Y_2^{M*}(\hat{\theta}_f) \quad (29)$$

For fixed values of θ_p, φ_p , each term in the sum expression can always be transformed by a simple rotation to the form

$$W(\theta_p, \varphi_p; \theta_f, \varphi_f = 0) = 1 + Q \cos 2(\theta_f - \theta_0) \quad (30)$$

where

$$Q(K_p) = \frac{3g_2 \sqrt{\gamma^2 + \beta^2}}{2 + g_2 \alpha} ;$$

$$\tan 2\theta_0 = -\frac{\gamma}{\beta}$$

and

$$\alpha = \frac{1}{2} d_{2j}^0 + \sqrt{\frac{3}{2}} d_{2j}^2$$

$$\beta = \frac{1}{2} d_{2j}^0 - \frac{1}{\sqrt{6}} d_{2j}^2 \quad (31)$$

$$\gamma = \sqrt{\frac{2}{3}} d_{2j}^1$$

Then by direct calculation and comparison with the plane wave theory, one finds that for each of the relevant j -values, Q and θ_0 approach the plane wave value for $\theta_p \geq \theta_c = 140^\circ$. Examples for $j = \frac{3}{2}$ and $l = 1, 2$ are shown in Figure 5. These cases were computed for $K_0^2 = 0$ which guarantees $g_L(j; l; K_0^2 = 0)$ equal to $g_L(j; l; 0)$ which is independent of J . The agreement in Figure 5 therefore justifies the use of plane wave theory in computing $\tilde{\rho}_{Ll}^M$ for the present case. For $\bar{\sigma}_j$, however, the DWBA values have been used throughout.

Using equations 28, 20 and 23 we have computed $g_2(I, K_0^2)$ for $I = 1/2$ and $I = 5/2$ corresponding to Pu^{239} and U^{233} . These functions are plotted in Figure 6. From the measured values of g_2 one may then infer values of K_0^2 as a function of excitation energy. These are shown in Figure 7. It may be noted that K_0^2 is near zero at the fission threshold as expected a priori for an even-even nucleus and that the values of K_0^2 at higher excitation energies are in reasonable agreement with those measured independently by means of neutron induced fission.

5. INTERPRETATION OF RESULTS

In a previous analysis of the energy dependence of the neutron and alpha-particle induced fission anisotropies, an estimate was obtained for the pairing-gap parameter, Δ_0 , in Pu^{240} distorted to its saddle point shape: $\Delta_0 = 1.16 \text{ Mev}$.⁶ This value is rather larger than that appropriate to Pu^{240} at its stable shape.¹⁷ According to the pairing model one would expect to find among the transition states no 2 quasi-particle excitations, until the available energy exceeds $2\Delta_0 \approx 2.30 \text{ Mev}$. Indeed, aside from some few collective vibrational excitations and their rotational bands, one expects to find only the rotational states of the ground state band (with $K = 0$) up to this minimum two quasi-particle energy. Above $2\Delta_0$, on the other hand, the two quasi-particle states should immediately become numerically dominant.⁴ This sudden change of character would reflect itself in a sudden increase from $K_0^2 \approx 0$ to $K_0^2 \approx \overline{2k_p^2} \approx 16$ to 22, where $\overline{k_p^2}$ is the mean square value K for the last filled single particle states. The steep rise in K_0^2 for Pu^{240} at

$E^* - E_f \approx 2.6$ Mev is quite consistent with this picture and should, we believe, be interpreted as direct evidence that $2\Delta_0 \approx 2.6$, a value somewhat larger than that mentioned above, but not inconsistently so.¹⁸

One might inquire why the energy gap should be larger at the highly deformed saddle point shape than at the ground state shape. One possible answer is that the pairing effect in nuclei is essentially a finite size effect, depending for a given nucleus upon the surface to volume ratio of the shape in question. Such an explanation is indicated by the work of Kennedy, et al¹⁹ who studied pairing effects as a function of thickness in an infinite (in two dimensions) slab of nuclear matter. It would also be consistent with the results of studies of pairing effects in infinite nuclear matter,^{20,21} which give a nuclear pairing effect much smaller than that observed in stable nuclei.

There also occurs in Figure 7 evidence for two smaller increases in K_0^2 at energies 0.7 and 1.6 above the fission threshold. The suggestion has been advanced²² that this behavior is qualitatively consistent with excited vibrational states analogous to the γ -vibrations of deformed nuclei about their stable shapes. The simplest γ -vibration has $K = \pm 2$. The value $K_0^2 \approx 3$ inferred for $0.7 < E^* - E_f < 1.6$ Mev, therefore corresponds well with a situation where fission is occurring through transition states of this kind, as well as through higher rotational states of the ground state ($K = 0$) band. Similarly, a double γ -vibrational excitation would be expected to occur at about twice that excitation energy. There would then occur a triplet of excited states with $K = \pm 4, 0$.

The value $K_0^2 \approx 7$ observed for $1.6 < E^* - E_f < 2.6$ Mev is consistent with fission through such barrier states as well as through higher rotational members of the $K = 0$ and $K = 2$ bands discussed above. Although only a speculation, this suggestion is interesting in pointing out the role of the larger pairing gap for the observation, and perhaps even for the existence, of "multi-phonon" vibrational states: in nuclei with equilibrium deformation, such states occur at energies where two quasi-particle states would normally be available to mix with the collective state or at least to obscure it for the experimentalist.

A comparison can be made between the behavior of K_0^2 from the present experiment and the previously measured variation of fission barrier penetrability.¹ In figure 8 the lines which have been drawn through the data points of the two experiments are plotted on the same scale of excitation. This figure emphasizes that both types of data show an increase at about 0.7 MeV excitation above the fission barrier, corresponding to the suggested onset of a vibrational band. A similar correlation at $E^* - E_f = 1.6$ and 2.6 MeV is obscured in the penetrability data by the beginning of both neutron emission and deuteron breakup.

The conclusions of the present paper may contribute to the interpretation of neutron fission cross sections. Specifically it is interesting to look for possible effects of the pairing energy gap in the well-determined data²⁵ for Pu^{239} . Indeed, these data show an inflection in cross section from a plateau value of 1.65 barns below 1 MeV, to 2.0 barns above. This change is rather well centered at 1 MeV, an energy which produces 2.6-MeV excitation energy in the

fissioning Pu^{240} nucleus. The corresponding large increase of K_0^2 at this energy suggests that this increase in cross section results from the onset of quasi-particle excitations.

As noted in section 3, the data for U^{234} fission is inferior statistically to that for Pu^{240} . In Figure 7 the same curve (dotted) has been drawn through the U^{234} data as was given by the Pu^{240} case to indicate that there is no evidence for a qualitative difference between the K_0^2 behavior for these two nuclides, and that the difference in the observed anisotropies is adequately accounted for by the difference in target spins.

REFERENCES

1. J. A. Northrop, R. H. Stokes, and K. Boyer, *Phys. Rev.* 115, 1277 (1959).
2. A. Bohr, Proceedings of the International Conference on the Peaceful Uses of Atomic Energy, Geneva, 1955 (United Nations, New York, 1956), Vol. 2; I. Halpern and V. M. Strutinskiĭ, Proceedings of the Second United Nations International Conference on the Peaceful Uses of Atomic Energy, Geneva, 1958 (United Nations, Geneva, 1958), Vol. 15. J. Griffin, Proceedings of the International Conference on Nuclear Physics, Paris, 1958 (Dunod, Paris, 1958); *Phys. Rev.* 116 107 (1959); 127, 1248 (1962); Proceedings of the International Conference on Nuclear Structure, Kingston, 1960, edited by D. A. Bromley and E. W. Vogt (University of Toronto Press, Toronto, Canada, 1960). V. M. Strutinskiĭ, *Nucl. Phys.* 27, 348 (1961).
3. A. Bohr, B. R. Mottelson, and D. Pines, *Phys. Rev.* 110, 936 (1958); S. T. Belyaev, *Kgl. Danske Videnskab. Selskab, Mat.-Fys. Medd.* 31, No. 11 (1959); The Many Body Problem (John Wiley and Sons, Inc., New York, 1959), p. 377; B. R. Mottelson, The Many Body Problem (John Wiley and Sons, Inc., New York, 1959), p. 283; J. Griffin and M. Rich, *Phys. Rev.* 118, 850 (1960); S. G. Nilsson and O. Prior, *Kgl. Danske Videnskab. Selskab, Mat.-Fys. Medd.* 32, No. 16 (1960); and other references too numerous to list.
4. M. Rich and J. Griffin, *Phys. Rev. Letters* 11, 19 (1963).
5. D. W. Lang, *Nucl. Phys.* 42, 353 (1963).

6. J. Griffin, Phys. Rev. 132, 2204 (1963).
7. R. H. Stokes, Rev. Sci. Instr. 31, 768 (1960).
8. J. C. D. Milton and J. S. Fraser, Can. J. Phys. 40, 1626 (1962).
9. The equivalent neutron energies at threshold as defined from the data of Fig. 2 are more negative by about 200 keV than the results of the previous measurement.¹ This discrepancy is within the uncertainties of the proton energy calibrations and does not affect the $E^* - E_p$ energy scale.
10. L. N. Usachev, V. A. Pavlinchuk, and N. S. Rabotnov, Soviet Physics, JETP, 44, 1950 (1963) point out an alternative prescription for the threshold. Their estimate of the difference between the two interpretations is, however, exaggerated by the large value of E_{curv} ($\approx .800$ MeV) they assume. A value $E_{\text{curv}} \leq 0.350$ MeV, as is indicated by the results of reference 1, reduces the change in threshold considerably from their estimate. Cf. also reference 18.
11. In the layout shown in Fig. 1, positive angles are measured in a clockwise direction from the recoil angle and negative angles in a counter-clockwise direction.
12. G. R. Satchler, Proc. Phys. Soc. (London) A66, 1081 (1953).
13. Huby, Refai, and Satchler, Nuclear Phys. 9, 94 (1958).
14. A. M. Lane, Proc. Int'l. Conf. on the Nuclear Optical Model, Florida State University, Tallahassee, Florida (1959).
15. A. E. S. Green, Proc. Int'l. Conf. on the Nuclear Optical Model, Florida State University, Tallahassee, Florida (1959).

16. J. Griffin, Phys. Rev. 127, 1248 (1962); J. Griffin and W. R. Gibbs, Phys. Rev., in press; J. Simmons et al., Phys. Rev., in press.
17. S. G. Nilsson and O. Prior, Mat. Fys. Medd. Dan. Vid. Selsk. 32 No. 16 (1960).
18. H. C. Britt (private communication) points out that a reinterpretation of the results of reference 1 might shift the measured threshold upward by perhaps 200 kev. This would reduce the value $2\Delta_0$ inferred here by approximately the same amount. Cf. also reference 10.
19. R. C. Kennedy, L. Wilets, and E. M. Henley, Phys. Rev. Letters 12, 36 (1964).
20. V. J. Emory and A. M. Sessler, Phys. Rev. 119, 248 (1960).
21. E. M. Henley and L. Wilets, Phys. Rev. Letters 11, 326 (1963).
22. J. J. Griffin and M. Rich, Bull. Amer. Phys. Soc. II, 8, 525 (1963).
23. J. E. Simmons, R. L. Henkel and R. B. Perkins, Phys. Rev. (in press).
24. J. Simmons and R. Henkel, Phys. Rev. 120, 198 (1960).
25. R. L. Henkel, Los Alamos Scientific Laboratory Report LA-2114 (1957) unpublished.

FIGURE CAPTIONS

1. Schematic drawing of the experimental arrangement. R designates the direction of the recoiling excited nucleus from which the fission fragments arise, and the ellipse suggests the fragment angular distribution for a fixed proton momentum.
2. Spectra of protons ($\theta_p = -140^\circ$) coincident with fission fragments. These data are the sum of the spectra obtained with $\theta_f = +20^\circ$, $+110^\circ$, and $+155^\circ$.
3. Angular correlations taken with a single fission detector for a Pu^{239} target. For laboratory proton scattering angles of 75° and 140° , the angular distribution of fission fragments is shown for various intervals of the excitation energy above the fission threshold. The lower two curves are for 0-1 MeV excitation, the middle two curves for 1-2 MeV excitation, and the upper two curves for 2-3 MeV excitation. The angular scale is taken relative to the recoil direction of the excited fissioning nucleus. θ_{lab} designates the value of the laboratory recoil angle for the two values of θ_p .
4. Measured values of the coefficient g_2 for U^{233} and Pu^{239} targets. These values were derived from data taken with three fixed fission detectors and with $\theta_p = 140^\circ$.
5. A comparison of distorted and plane wave calculations for $j = \frac{3}{2}$ and $K_0^2 = 0$. These curves illustrate the characteristic feature that for $\theta_p \geq 140^\circ$ the D.W.B.A. and P.W.B.A. give identical results for Q and φ .

6. K_0^2 vs g_2 calculated by means explained in the text. The difference between the curves results primarily from the difference in the target spins. This effect accounts adequately for the differences observed between the U^{233} and Pu^{239} data.
7. Values of K_0^2 (solid points) calculated from measured values of g_2 by means of the theory developed in the text. The triangles are from neutron data and its analysis for Pu^{239} can be found in Griffin⁶ and Simmons, et al²³; for U^{233} , see Simmons and Henkel.²⁴
8. Comparison between the variation of K_0^2 and the fission barrier penetrability, P.