Columbia University
Division of Nuclear Science and Engineering

PROGRESS REPORT FOR JANUARY 1, 1971, THROUGH JUNE 30, 1972
TO THE UNITED STATES ATOMIC ENERGY COMMISSION

MASTER

The research reported in this document was
made possible through the support extended
to Columbia University by the United States
Atomic Energy Commission, under

A.E.C. Contract AT (30-1) 73, Scope C
and

A.E.C. Contract AT (11-1) 2176

W. W. Havens, Jr., Director

Reproduction in whole or in part is permitted for any purpose
of the United States Government.

NOTICE

This report was prepared as an account of work
sponsored by the United States Government. Neither
the United States nor the United States Atomic Energy
Commission, nor any of their employees, nor any of
their contractors, subcontractors, or their employees,
makes any warranty, express or implied, or assumes any
legal liability or responsibility for the accuracy, com-
pleteness or usefulness of any information, apparatus,
product or process disclosed, or represents that its use
would not infringe privately owned rights.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

COLUMBIA UNIVERSITY

I. NEUTRON SPECTROSCOPY

A. Neutron Resonance Cross Section Measurements (J. Rainwater, F. Rahn, H. Liou, G. Hacken, M. Slagowitz, W.W. Havens, Jr., S. Wynchank)

Recently our attention has been concentrated on the analysis of the data obtained during the two preceding experimental runs. The data on the Eu and Sm isotopes have been published in the Physical Review (Phys. Rev. C 5, June (1972)), as well as the paper on the statistical tests for single level populations to neutron spectroscopy data (Phys. Rev. C 5, 1002 (1972)). Our results for the minima in the total cross section in Fe have been published in Nuclear Science and Engineering (N.S.&E. 47, 372 (1972)). The paper on resonance integrals of the rare earth isotopes Sm, Eu, Gd, Er, Yb, Lu, and W has been accepted by Nuclear Science and Engineering for publication and it will probably be published in June 1972.

A short catalogue of resonance energies suitable for the calibration of time of flight neutron spectrometers has been prepared. This work was done in cooperation with Dr. Rainwater's group. Table A1 gives resonance energies of some selected prominent sharp neutron resonances in readily available isotopes useful for this purpose. These resonance energies should be helpful in eliminating possible energy shifts in neutron spectroscopy data which may arise in neutron velocity selector results. The most common shift is due to difficulties in determining the zero point on the timing scale, in which case all calculated energies will be inaccurate by a quantity that varies monotonically with energy.

The work of the analysis of the $^{154},^{158},^{160}\text{Gd}$ data taken in 1968 and 1970 is nearing completion. Figure A1 shows examples of our data for these isotopes. We obtained individual parameter results up to 1 keV in ^{154}Gd and 10 keV in $^{158},^{160}\text{Gd}$. Table A2 lists the parameters E_0 , Γ_n , and in favorable cases Γ_γ for these isotopes, along with

preliminary assignments. These values have been revised from the ones we presented in our last report. Figure A2 shows the curves of $N(E)$ versus E and the histogram of adjacent level spacings for ^{154}Gd . A good fit to a straight line is obtained to 300 eV, giving a $\Delta_{\text{exp}}=0.22$, compared with theoretical $\Delta_{\text{DM}}=0.28\pm0.11$ expected in the Dyson-Mehta theory of adjacent level spacings. The Wigner distribution gives good agreement with the histogram of adjacent spacings in this energy interval, with a correlation coefficient of adjacent spacings of $\rho(S_i S_{i+1}) = -0.6\pm0.3$ versus the $\rho = -0.27$ predicted by the Orthogonal Ensemble (O.E.) theory. From Figure A3 we determine our value of the strength function. The slope of the line drawn to $\Sigma \Gamma_n^0$ versus E gives our "best choice" value of $S_0 = (2.0\pm0.4)\times 10^{-4}$ for this isotope. We found $\langle \Gamma_\gamma \rangle = 88$ meV for 25 levels in ^{154}Gd , with few values of Γ_γ more than 20% from the average. A χ^2 variance analysis gives $\nu = 151$ degrees of freedom, mainly due to experimental uncertainties in the individual Γ_γ values.

Examples of our ^{158}Gd and ^{160}Gd results are presented in Figure A4. Figure A5 shows the level densities and Wigner plots for the $^{158},^{160}\text{Gd}$ isotopes. Good fits to a straight line for the level spacing are obtained up to ~ 4 keV for these isotopes. In ^{158}Gd , we obtain $\Delta_{\text{exp}} = 0.29$ (versus $\Delta_{\text{DM}} = 0.38\pm0.11$) and a $\rho(S_i S_{i+1}) = -0.14\pm0.15$; while for ^{160}Gd we obtain $\Delta_{\text{exp}} = 0.32$ (versus $\Delta_{\text{DM}} = 0.30\pm0.11$) and a $\rho(S_i S_{i+1}) = -0.33\pm0.17$. The experimental average spacing of s levels is $\langle D \rangle = 86\pm4$ eV and 202 ± 27 eV respectively. The $\sigma(k)$ of the spacings distributions for k spacings is much lower than implied by Bohigas and Flores' Two Body Random Matrix Theory, and is in general agreement with Wigner Gaussian Orthogonal Ensemble. Figures A6 and A7 show determinations of S_0 for ^{158}Gd and ^{160}Gd respectively. We find $S_0 = (1.5\pm0.2)\times 10^{-4}$ and $(1.8\pm0.4)\times 10^{-4}$ in these isotopes, with $\langle \Gamma_\gamma \rangle = 105$ meV ($n = 27$) and 111 meV ($n = 4$).

The work on the total cross section of natural iron at points of relative minima is complete and has been submitted for publication. Figure A-8 shows our new values, which represent our best experimental results to date. The knowledge of σ_T at the points presented in this figure is especially important, in that these energy regions represent "holes" in the total cross section which allow the transport and deep penetration of neutrons in several important applications, such as shielding and fast reactor design. Our results in Figure A-8 are lower than previous experimental results. The values of σ_T in this figure still do not represent our best experimental resolution, and we will redo our Fe measurements during our next experimental series.

The resonance capture integrals for the rare earth isotopes measured during 1968: ^{152}Sm , ^{154}Sm , ^{151}Eu , ^{153}Eu , ^{154}Gd , ^{158}Gd , ^{160}Gd , ^{166}Er , ^{167}Er , ^{168}Er , ^{170}Er , ^{168}Yb , ^{170}Yb , ^{171}Yb , ^{172}Yb , ^{174}Yb , ^{176}Yb , ^{175}Lu , ^{182}W , ^{183}W , ^{184}W , and ^{186}W have been evaluated. These isotopes lie in the region of the periodic table that exhibits large nuclear deformation and the split "4S" giant resonance in the strength function. Their relatively high level density and fractionally large Γ_γ (compared to Γ) make these isotopes suitable for application as control poisons in nuclear reactors. Table A-3 shows our values of the resonance capture integral as well as $\langle \Gamma_\gamma \rangle$ as determined from our data.

B. Cross Section and Resonance Parameters $A < 80$ (F. Rahn, J. Rainwater, W.W. Havens, Jr., M. Slagowitz, U. Singh)

Transmission measurements of the total neutron cross section of natural sodium samples were run during the 1968 and 1970 experimental series. The measurements covered the energy region of 10 eV to greater than 350 keV. The full width half maximum resolution of the data was approximately 0.2 nsec/meter. Our previous results on sodium were limited by resolution and background effects to energies below 100 keV. Our present data is an attempt to increase the energy interval so that it extends into the region where other recent measurements have been made at higher energies.

Our samples had a range of inverse thickness ($1/n$) values between 4.8 and 550 barns/atom. The background associated with our data presented special problems above 20 keV, where it was found to be especially dependent on the neutron energy. In order to assure an accurate normalization of the data, we used Fe and Co in the filter technique, which directly yields the total cross section of sodium at points in the energy spectrum where these materials have sharp resonance structure. In this manner we have determined the sodium cross section near the 376 Fe resonance to be 3.46 barns.

Figure B1 shows our results in the 20 to 360 keV region. The measured total cross section as shown in this figure and the "true" cross section differ by an amount which depends on the thickness of our samples and resolution effects. The resolution effects, important where the cross section is rapidly varying, is a function of the transmission, $T(E)$. The resonance parameters of the first two levels, obtained by an R-matrix analysis, are as follows: the 2.850 keV level is assigned $J=1$, $\ell=0$ with $\Gamma=388$ eV, and the 53.15 keV level can be fit with $J=2$, $\ell=1$ and $\Gamma=1228$ eV. These R-matrix parameters are consistent with peak cross sections of $\sigma_0=390$ and 37 barns respectively, and the shapes of these resonances. There is general agreement of our total cross section values with the Karlsruhe data in the region where the data sets overlap, 280 to 360 keV. Our values are slightly higher overall, and do not go as deep in the minimum close to 296 keV, which may be due to the lower resolution of our data in this region.

Measurements of the total cross section of natural copper and ^{63}Cu were also made during 1968 and 1970, for a series of thicknesses ($1/n$)=4.6 to 458 barns/atom. Figure B2 shows our results from 60 to 2 keV. The filter materials Co, Fe, Ta, Cl, and U were used to assure a proper normalization of the data. Work is beginning on the resonance parameter analysis, which will include area and shape formalisms, to determine the energies and neutron widths of the observed levels, isotopic identification, and attempt to assign their ℓ and J values.

C. Cross Section and Resonance Parameters of ^{232}Th and ^{238}U
(F. Rahn, H. Liou, J. Rainwater, W.W. Havens, Jr., M. Slagowitz,
S. Wynchank)

The analysis of the ^{238}U data from the 1970 run is complete. Preliminary total neutron cross section plots and resonance parameters appeared in our last report. Our work during this report period concentrated on the spin assignment of the observed levels. We have applied the following statistical tests in the s-population selection: Wigner distribution, Porter-Thomas distribution, Dyson-Mehta Δ statistic and $\text{Cov}(S_i, S_{i+1})$. In addition we used the F test, due to Dyson, for final level spin assignment. The F test (or F statistic) involves the evaluation at each resonance position E_i of a parameter F_i , defined as

$$F_i = \sum_{j \neq i} f(x_{ji}),$$

where $x_{ji} = (E_j - E_i)/L$, and $f(x) = \frac{1}{2} \ln\{[1+(1-x^2)^{1/2}]/[1-(1-x^2)^{1/2}]\}$. The summation is taken over all resonances between $(E_i - L)$ and $(E_i + L)$, where $L = M\langle D \rangle$ is the largest interval characterized by a properly chosen integer M times the average level spacing $\langle D \rangle$. Dyson finds that $\langle F_i \rangle = n - \ln(n) - 0.656$, with a standard deviation of $[\ln(n)]^{1/2}$ where $n = \pi M$. If a spurious level is present $\langle F_i \rangle = n$ at this spurious level. Thus a large positive fluctuation occurs at a "spurious" level, while a low value of F_i results near a missed level.

The problem in ^{238}U is eliminating as p wave or spurious the large number of excess levels seen in the data. Nearly all s wave levels were observed. We assumed that the stronger levels were s wave and therefore the analysis concentrated on the assignment of the weaker levels. The Porter-Thomas distribution was used to determine the expected number of small resonances. Table C-1 presents our spin assignments for ^{238}U . Figure C-1 and C-2 show the comparison of our data with the theoretical Porter-Thomas and Wigner distributions. For this set of s wave energy levels, we obtain a $\Delta_{\text{exp}} = 0.42$ (versus $\Delta_{\text{DM}} = 0.49 \pm 0.11$) and a $\text{Cov}(S_i, S_{i+1}) = -0.26 \pm 0.08$ (versus an expected $\text{Cov}(S_i, S_{i+1}) = -0.27$). Our best choice for $\langle D \rangle$ is 20.8 eV.

The preliminary transmission, self-indication and Moxon-Rae analysis of the data from the 1970 run has been completed. The ^{232}Th samples had thicknesses of $(1/n) = 10.72, 31.9, 36.0, 19)$ and 885 barns/atom. The transmission measurements were carried out at 200 meters with a nominal resolution of 0.1 nsec/meter. The self-indication detector was at 40 meters and the Moxon-Rae measurements were carried out at 33 meters. A large number of reference samples such as Fe, Ta, Co, Cu, etc., were used as beam filters to help in the absolute normalization of the data. The sharp structure in the cross section of these reference samples determines the transmission of Th at the energies where the structure occurs in the reference samples, independent of the details of the background which is energy and sample dependent. The knowledge of the Th transmission at these points establishes $T=0$ and $T=1$, and allows a self-consistent determination of the (T, σ) values. Moxon-Rae measurements

were performed simultaneously to give capture yields which could be related to the resonance parameters. Self-indication "D only" and "D+T" experiments were performed with a high efficiency plastic scintillation detector to give additional resonance information. There was good agreement between our "D only" and Moxon-Rae results.

The (T, σ) values of ^{232}Th were obtained by the transmission of the difference technique. A plot of the total neutron cross section appeared in our last progress report. We ran extra thick samples in the transmission measurement to accurately determine the total cross section between levels, where the cross section dips are due to potential-level interference.

To determine the resonance parameters, area analysis was mainly used and was supplemented by shape information. The area data gave a series of curves in the $(\Gamma_n, \Gamma_\gamma)$ plane, and the intersection of these curves gave the value of the parameters Γ_n and Γ_γ . The shape fits used values of σ_T in the wings of the stronger, isolated resonances, in places where σ_T was not rapidly varying, so that resolution effects were relatively small. We then obtain a series of curves for R' versus Γ or Γ_n , where R' is the (local) potential scattering length. An example of this analysis is in Figures C-3 and C-4. The combination of the area results and the shape results determines our values of the resonance parameters.

Table C-2 lists the resonance parameters $[E_0, g\Gamma_n \text{ and } \Gamma_\gamma]$ for 305 levels that we observed between 21 eV and 4 keV. In Figure C-5 we show the value of Γ_γ as a function of E for 84 levels up to 2400 eV. We found an average value of $\langle \Gamma_\gamma \rangle = [21.2 \pm 0.3 \text{ (statistical)} \pm 0.9 \text{ (systematic)} \text{ meV}]$ to be consistent with our data. We have assigned a systematic uncertainty of $\pm 0.9 \text{ meV}$ to account for the relative normalization of our data. We find no evidence for a quasiperiodic variation of Γ_γ with neutron energy. Figure C-6 shows the distribution of our Γ_γ values, with the histogram taken in intervals of 2 meV. The histogram can be suitably fit by a chi-squared function with 87 degrees of freedom, generated about a mean value of 21.2 meV. Our average value of 21.2 meV (0-2400 eV) compares with a value of $\langle \Gamma_\gamma \rangle = 20.5 \pm 3 \text{ meV}$ obtained by Forman at Los Alamos, 20.9 meV obtained by Asghar (0-350 eV) and 21.6 meV (0-350 eV) obtained by Ribon at Saclay.

Our values of the total neutron width Γ_n in Table D-1 tend to be somewhat higher than the previous Columbia results of Garg et al. in the energy region above 1 keV. Figure C-7 shows $\Sigma g\Gamma_n^0$ versus E for ^{232}Th . The energy interval used in obtaining S_0 affords relatively complete level resolution. Our best choice of the strength function is $S_0 = (0.84 \pm 0.08) \times 10^{-4}$. Missing or spurious levels of small Γ_n have small effect on S_0 . We are in good agreement with the recent Saclay results of Ribon [$S_0 = (0.87 \pm 0.10)$] and Harwell results of Asghar (0.8 ± 0.17) but are higher than the results of Garg (0.69 ± 0.07).

By the application of the various statistical tests in a manner similar to ^{238}U , we attempted to assign spins to the observed

resonances. The resulting s-wave population to 3 keV is in good agreement with all the statistical tests (Wigner distribution, Porter-Thomas distribution, Dyson-Mehta Δ , $\text{Cov}(S_i, S_{i+1})$, and F statistic). The stronger levels provided the framework for the choice of the weaker s-wave levels. The probability of another choice of the s-wave level population in ^{232}Th satisfying all the statistical tests is small. Figures C-8 and C-9 respectively show our comparison with the Porter-Thomas and Wigner distribution for our choice of the s-wave population in ^{232}Th .

Tables C3 and C4 give resonance parameters for levels we identify as p levels in ^{232}Th and ^{238}U . The calculation of the $g\Gamma_n^1$ values requires that an effective nuclear radius R be chosen. This gives a "barrier energy" E_1 where the neutron wave length λ equals R. We use $R=1.41 A^{1/3}$ fermi, giving $E_1=276$ keV and 270 keV for ^{232}Th and ^{238}U respectively. The statistical spin factor g is included, and is unity except for $P_{3/2}$ levels where it is equal to 2. The information from the $g\Gamma_n^1$ values for these levels permits us to make rough estimates for the p wave strength functions S_1 for the two isotopes. In making such estimates it must be noted that (a) most of the p levels are too weak for us to observe, (b) the fractional uncertainties in these $g\Gamma_n^1$ values are relatively large, (c) these weak levels are mainly seen using our thickest samples where a significant fraction of the energy range is not satisfactory for observing weak p levels due to the effects of the strong s levels, and (d) an examination of our data shows that we have a greatly reduced efficiency for detecting p levels above 400 eV in ^{232}Th and 1200 eV in ^{238}U .

We assume that the "true" p level density is three times that for s levels (due to statistical factors) and the $\langle g\Gamma_n^1 \rangle$ is the same for $P_{1/2}$ and $P_{3/2}$ levels. An estimate is made of the effective energy interval ΔE_1 , available not too near the strong s levels, and the strength function implied by the number of levels observed having $g\Gamma_n^1$ values greater than a given $g\Gamma_n^1$. We found $S_1=(0.9^{+0.5}_{-0.3})\times 10^{-4}$ for ^{232}Th and $(1.43^{+0.30}_{-0.27})\times 10^{-4}$ for ^{238}U . The uncertainties in these values were determined by using distinctly too few and too many levels to be consistent with our data.

From our analysis of total cross section data, including shape fitting of various well isolated s resonances, we were able to arrive at values of R' , the optical model potential scattering radius. Interference contributions from more than one resonance are significant in these isotopes, except in the vicinity of a few single isolated resonances, notably the 221 eV level in ^{232}Th and 347 eV level in ^{238}U . At these energy regions, we examined the behavior of σ_1 , particularly at the interference minima and in the wings where σ is sensitive to R' , and resolution effects are relatively unimportant. We found that $R'=9.1\pm 0.3$ f. for ^{232}Th and 9.6 ± 0.3 f. for ^{238}U .

D. Fission Cross Section of ^{233}U (J.P. Felvinci and E. Melkonian)

Increasing interest in thermal breeder reactors warrants a second (and critical) look at the ^{233}U cross sections. We reported earlier¹ that when the fission fragment energy distribution was subdivided into groups, the resulting partial fission cross sections showed significant variation. Suggestions were made that some of the differences observed in fission cross sections among several experimenters could be due to the different bias in the detection of the fission fragments.

In this report we present $\sigma_0\Gamma_F$ values for the fission resonances. The experiment which we report on was performed in the Spring of 1970 at the Nevis (NVS) synchrocyclotron. The details of the experiment were reported earlier¹. Both time-of-flights and fission fragment energies were measured simultaneously and recorded event-by-event on magnetic tape. The energy resolution was good enough to subdivide the fission cross section into partial fission cross sections corresponding to different energy groups.

In this report we present in Table D1 the $\sigma_0\Gamma_F$ values calculated from fitting the experimental data to a single level Breit-Wigner formula. The results are compared with values given in BNL-325. In Table D2 the G and H values calculated from the Adler-Adler formalism are given and are compared with other published results. The errors quoted include both statistical and possible fitting errors. Table D3 gives the fission resonance integrals

$$\int_{E_1}^{E_2} \sigma_F \frac{dE}{E}$$

over GAM (1/4 lethargy) intervals and they are compared with existing data. Table D4 summarizes the fission resonance integrals for different energy intervals corresponding to other experimenters' values.

We analyzed resonances in detail up to 30 eV and will continue to analyze them at higher energies. Since the fits to resonances above 30 eV involve matching at the boundary to the fits below 30 eV, it is possible that some resonance parameters just below 30 eV will be slightly changed. We believe that the change will be within the errors quoted.

The $\sigma_0\Gamma_F$ values from the fit to the Doppler and resolution broadened single-level Breit-Wigner formula show reasonable agreement with Nifenecker's results for the large resonances, but are quite different

for wide or very close resonances. In these cases naturally the exact resonance energy assumed and the width used is quite critical. Several minor changes in the resonance parameters kept the values within the quoted errors.

It might be pointed out that in the single level formalism a level at 2.16 eV is necessary for a good fit, but this can be removed in the Adler-Adler form. The additional level at 6.44 eV and 12.22 eV cannot be removed, as no interference term in the multilevel fit will give a good result. Inspection of the Weston et al.³ results and the calculation of de Saussure⁷ shows that they have also some excess area in these regions, though these are accentuated in our experiment. We believe again that these are wide levels having wider mass and energy distribution and thus slightly biased against in ionization chamber experiments. The level at 11.69 eV has an especially high ratio of low energy to medium energy fragments¹ and also a higher proportion of high energy γ -rays². We also note the changing ratio of areas between the 21.75 and 22.28 eV level in different pulse height cuts.

Inspection of the resonance integrals in different intervals in Table D3 reveals some discrepancies between our results and past experiments. We seem to be on the high side in the energy interval of 4 - 6.5 eV which is a region of small, wide levels and again in the region of 8.3 - 25 eV. At higher energies our results seem to be close to those of Weston et al.³ but are above the results of Moore et al.⁴ and Cao et al.⁵

The cross sections were calculated by assuming the energy dependence of the flux to be $\phi(t) = E^{0.60} dt$. This was confirmed by comparison with other experiments in the same run and also by Monte-Carlo calculations. The background was assumed to consist of a part which varies as the neutron flux and a fairly constant one. Normalization was obtained by comparing the cross sections at 1 eV, 20 keV, and the peak cross sections at several resonances with those of Weston et al.¹ Discrepancies between peak values at several resonances were noted and attributed to possible pulse height effects in the fission chamber they used.

We have observed in the past that our "background" is higher for the low energy fission fragments. We have ruled out admixture of α -particles as our bias was set around 35 MeV above the α -particles but below the fission fragments. Our conclusion is that some wide levels which contribute more to the "background" have a larger proportion of low energy fission fragments (also very high energy fragments) and thus are possibly discriminated against in other experiments.

References

1. Felvinci, J.P. and Melkonian, E. Proc. Third Conf. on Neutron Cross Sections and Technology, Vol. II, 855. Knoxville, Tennessee. 1971
2. Felvinci, J.P. and Melkonian, E. NYO-72-227, p.12. 1969.
3. Weston, L.W., et al. ORNL-TM-2140 and Nucl. Sci. Eng. 39, 1 (1968).
4. Moore, M.S., et al. Phys. Rev. 118, 714 (1960).
5. Cao, M.G., et al. Nuclear Data for Reactors, Vol. I, 419 (1970). IAEA, Vienna.
6. Nifenecker, H. Journ. de Physique 25, 877 (1964).
7. de Saussure, G., et al. Nuclear Data for Reactors, Vol. II, 757 (1970). IAEA, Vienna.

E. Delayed Fission Following Gamma-Ray Emission in $^{235}\text{U}+n$
 (M. Derengowski, J. Felvinci, and E. Melkonian)

This experiment, which was performed in the spring of 1970 and which used the Nevis synchrocyclotron as a pulsed neutron source, has been described in previous reports to the NCSAC. The data have been analyzed and an isomeric fission state with a half-life of 7.0 ± 2.2 nsec has been identified. This short-lived isomeric state has not been observed previously, although an isomeric state (or states) with half-lives ranging from 67 to 130 nsec has been reported. A longer lived state of at least 40 nsec is also required to fit our data, but the data are insensitive to the exact value.

We estimate that at least 2% of all fissions go through the 7 nsec isomeric state. While this contribution is much higher than observed previously, it is consistent with the trend noted by others that the relative proportion of isomeric states increases with decreasing neutron energy. In our case, fissions were induced primarily by neutrons of a few eV, while other measurements were performed with neutrons in the keV and MeV ranges.

Figure E-1 shows the data used in this analysis together with the best fit curve obtained by a least squares fit using five functions, namely 1) a gaussian representing gammas associated with prompt fission and having a standard deviation of 3.0 nsec, corresponding to the time resolution; 2) a short lifetime exponential broadened by the gaussian resolution function, the actual lifetime being adjusted for best fit; 3) a similarly broadened long lifetime exponential arbitrarily set at 80 nsec; 4) a function representing the effect of detecting gamma rays produced by the capture or inelastic scattering of prompt fission neutrons; and 5) a constant background arising from chance coincidences. All but number 5 are shown in Figure E-2.

A paper is being submitted to Physical Review Letters.

F. Fission Cross Section of ^{235}U (J. Felvinci and E. Melkonian)

It is known that different sets of cross section data frequently differ by considerable amounts (up to 15%). This could be due to incorrect flux dependence, inadequate background corrections, and improper normalization. The peak cross sections in the major resonances are considerably different in the works of Michaudon et al.¹, Bowman et al.², Brown et al.³, de Saussure et al.⁴, and Brooks et al.⁵. We decided to undertake a critical examination of these discrepancies and of the possibility that experimental problems could cause them. This is quite important as possible differences in the width and height of the resonances will certainly affect the resonance parameters and thus the quantities calculated from them. Subtle effects due to selective pulse height discrimination in fission chambers or underestimating scattering contributions could all contribute to the discrepancies.

Areas for the resonances were calculated by means of shape analysis and are given in Table F-1. Comparison with some other existing sets of data is given. The analysis performed was of a single level nature and no multilevel fits were attempted at this time. It was noticed though that our data agreed rather well with other experimenters who applied single level analysis, but was rather different from equivalent multilevel values.

The errors given in Table F-1 include both statistical and fitting errors. Some results quoted in the literature appear not to include the errors due to fitting and thus are smaller than they should be.

Table F-2 shows a comparison between the fission integrals,

$$\int_{E_1}^{E_2} \sigma_f dE,$$

for our data and those of de Saussure et al.⁶ Both sets are normalized to Deruytter's value of 237.1 eV b from 7.8 to 11.0 eV. These areas were calculated with the assumption that our flux varies as $\phi(t)dt = E^{0.60}dt$ from 1 eV to 10 keV. This variation was confirmed by other measurements during the same run and also by a Monte Carlo calculation performed to determine the flux emerging from the moderator. The background was taken to consist of two parts; one behaving as the flux and the other a constant.

Further work on the ^{235}U cross section is continuing; resonance integrals are going to be reported soon.

1. Michaudon, Derrier, Ribon, & Sanche, Nucl. Phys. 69, 545 (1965).
2. Bowman, Auchampough, Fultz, Moore, & Simpson, Conf. on Neutron Cross Sections & Technology, 1966, Washington DC, Vol. 2, p. 1004.
3. Brown, Berger, & Cramer, LA-3586, 1966.
4. de Saussure, Gwin, Weston, Ingle, Fullwood, & Hockenbury, ORNL TM-1804, 1967.
5. Brooks, Jolly, Schomberg, & Sowerby, AERE-M1670, 1966.
6. de Saussure et al., op. cit.

TABLE A1

Isotope	E_n (KeV)	ΔE_n (KeV)	Reference
^{23}Na	53.15	0.03	17
	296.5	1.0	17
^{27}Al	5.907	0.003	12
	119.8	0.3	12
	222.9	0.4	12
	255.7	0.5	12
	365.5	0.7	12
	435.5	1.0	12
	524.9	1.3	12
	646.9	1.7	12
	202.2	2.4	12
^{32}S	380.3	3.0	12
	591.7	8.0	12
	14.81	0.02	12
Cl (unassigned)	26.67	0.03	12
	113.8	0.2	12
	20.17	0.02	12
^{40}Ca	89.40	0.13	12
	133.8	0.3	16
	146.4	0.3	12
	170.9	0.4	16
	295.1	0.8	12
	640.9	1.0	15
	694.6	1.0	15
	758.0	1.0	15

TABLE A1

Isotope	E_n (KeV)	E_n (KeV)	Reference
Ti ⁴⁷	3.074	0.007	11
	10.510	0.014	11
Ti (unassigned)	11.040	0.016	11
	11.460	0.017	11
Mn ⁵⁵	17.80	0.02	12
	30.30	0.04	12
	41.09	0.05	12
	53.40	0.08	12
	81.59	0.08	12
Fe ⁵⁶	27.91	0.02	12
	73.96	0.07	12
	83.63	0.09	12
	129.53	0.14	12
	168.96	0.26	12
	188.0	0.3	12
	318.1	0.6	12
	435.5	1.0	12
	8.049	0.005	12
	10.692	0.010	12
Co ⁵⁹	13.268	0.010	12
	31.34	0.03	12
	41.49	0.06	12
	56.39	0.04	12
	66.28	0.06	12
	84.29	0.09	12
	142.3	0.1	12

TABLE A2

Neutron Resonance Parameters of ^{154}Gd

E_0 (eV)	ΔE_0	Γ_n (meV)	$\Delta \Gamma_n$	Γ_γ (meV)	$\Delta \Gamma_\gamma$
11.58	.02	.39	.06		
22.33	.06	11.5	1.0		
47.07	.07	2.8	.4		
49.50	.07	1.6	.2		
65.06	.05	21.	3.		
76.12	.08	1.1	.3		
100.68	.11	29.	4.0	80.	15.
105.61	.12	4.5	.3		
124.00	.15	115.	17.	85.	10.
139.18	.18	124.	16.	91.	10.
148.43	.20	33.	5.		
164.48	.23	105.	7.	73.	9.
170.35	.12	5.0	1.5		
201.61	.15	11.7	2.5		
211.02	.17	35.	5.		
224.01	.18	18.	3.		
252.77	.22	12.	2.		
257.46	.23	34.	5.		
269.16	.24	28.	4.		
331.71	.34	14.	2.		
333.76	.33	13.5	3.		
364.83	.38	7.	4.		
396.53	.43	95.	8.	65.	10.
407.56	.45	45.	9.		
444.73	.26	120.	15.	76.	11.
447.05	.27	94.	8.	69.	9.
467.95	.28	125.	15.	86.	9.
486.20	.31	69.	10.	75.	15.
511.52	.34	91.	12.	83.	13.
515.98	.32	50.	7.	80.	15.
551.76	.35	55.	7.	85.	15.
589.83	.43	152.	20.	115.	30.
625.50	.42	240.	35.	80.	10.
640.32	.44	61.	14.	90.	15.
684.65	.49	55.	10.		
687.11	.49				
773.97	.59	148.	12.	105.	15.
795.02	.61	35.	8.		
808.70	.63	195.	30.	95.	15.
811.86	.64	75.	20.		

TABLE A2 (cont.)

Neutron Resonance Parameters of ^{154}Gd

E_0 (eV)	ΔE_0	Γ_n (meV)	$\Delta\Gamma_n$	Γ_γ (meV)	$\Delta\Gamma_\gamma$
836.31	.33	540.	70.	104.	15.
845.32	.34	370.	50.	96.	15.
858.95	.35	265.	40.	97.	15.
897.42	.36	310.	40.	92.	12.
910.82	.37	63.	9.	84.	13.
916.87	.37	405.	50.	98.	10.
966.71	.41	275.	40.	105.	10.
985.50	.43	157.	28.	83.	14.

TABLE A2 (cont.)

Neutron Resonance Parameters of ^{150}Gd

E_0 (eV)	ΔE_0	Γ_n (meV)	$\Delta \Gamma_n$	Γ_γ (meV)	$\Delta \Gamma_\gamma$
22.30	.02	6.0	.7	96.	10.
101.08	.09	1.0	.2		
242.73	.17	46.	7.	105.	12.
277.15	.20	16.	2.		
344.76	.28	155.	15.	93.	8.
409.07	.36	265.	20.	110.	10.
503.31	.25	270.	20.	115.	10.
588.49	.31	56.	6.	112.	14.
692.86	.40	790.	100.	98.	10.
847.30	.54	1550.	200.	105.	12.
869.3	.5	3.3	1.6		
917.13	.31	460.	30.	88.	10.
1076.97	.37	270.	20.	70.	15.
1219.90	.47	850.	100.	110.	14.
1294.70	.25	430.	60.	112.	12.
1351.34	.28	25.	15.		
1427.61	.30	660.	50.	108.	14.
1458.84	.31	1070.	80.	135.	30.
1584.35	.33	360.	40.	93.	10.
1740.	1.	3.0	1.5		
1776.55	.41	480.	50.	128.	14.
1876.10	.44	265.	55.	102.	10.
*1911.	1.	1.5	1.0		
1944.68	.47	650.	120.	99.	10.
1990.60	.49	610.	120.	107.	13.
2113.	2.	8.9	4.0		
2243.40	.59	165.	35.	85.	17.
2317.33	.61	650.	80.	155.	25.
2361.	2.	7.6	4.0		
2386.66	.64	1115.	90.	117.	20.
2564.25	.71	350.	100.	135.	18.
2585.36	.72	80.	15.	70.	20.
2654.39	.75	225.	30.	95.	17.
*2706.	2.	2.4	1.5		
2795.73	.81	605.	30.	130.	25.
2868.78	.84	4200.	600.		
2933.82	.87	790.	40.	91.	13.
3071.	3.	42.	15.		
3094.15	.94	65.	25.		
3165.10	.98	220.	30.		

*p wave or spurious level

TABLE A2 (cont.)

Neutron Resonance Parameters of ^{158}Gd

E_0 (eV)	ΔE_0	Γ_n (meV)	$\Delta \Gamma_n$	Γ_γ (meV)	$\Delta \Gamma_\gamma$
3325.	3.	16.	8.		
3479.28	1.13	3400.	600.		
3535.	3.	21.	12.		
3586.21	1.18	770.	100.		
3624.14	1.20	40.	20.		
3684.	3.	24.	12.		
3768.10	1.26	2950.	300.		
3936.99	1.35	1500.	240.		
3982.70	1.38	4000.	800.		
4180.42	1.48	570.	40.		
4223.	4.	30.	15.		
4299.86	1.54	445.	80.		
4468.	4.	55.	20.		
4501.24	1.65	360.	70.		
4621.00	1.73	2050.	300.		
4796.15	1.81	1650.	200.		
4894.03	1.88	420.	80.		
4985.29	1.93	1100.	250.		
5029.93	1.97	100.	50.		
5113.00	2.00	50.	30.		
5196.10	2.06	30.	20.		
5334.18	2.16	360.	160.		
5414.01	2.18	910.	230.		
5611.23	2.30	6200.	1500.		
5680.93	2.35	1900.	500.		
5759.11	2.40	1600.	400.		
5993.42	2.56	1500.	300.		
6112.05	2.62	125.	50.		
6193.99	2.67	770.	195.		
6343.48	2.76	2050.	500.		
6376.81	2.79	2300.	600.		
6782.87	3.05	700.	200.		
6807.42	3.08	350.	150.		
6878.78	3.13	100.	50.		
7266.17	3.39	100.	50.		
7445.83	3.52	250.	70.		
7548.94	3.60	270.	80.		
7610.37	3.64	2950.	750.		
7676.25	3.76	5500.	1400.		
7854.23	3.84	1650.	450.		

TABLE A2 (cont.)

Neutron Resonance Parameters of ^{158}Gd

E_0 (eV)	ΔE_0	Γ_n (meV)	$\Delta\Gamma_n$	Γ_γ (meV)	$\Delta\Gamma_\gamma$
7936.85	3.87	600.	200.		
8010.94	3.93	300.	200.		
8235.46	4.07	1550.	400.		
8288.92	4.13	650.	200.		
8368.01	4.20	3700.	900.		
8660.54	4.72	850.	210.		
8740.54	4.48	6200.	1550.		
8867.21	4.57	1250.	300.		
9362.64	4.96	5800.	1500.		
9437.50	5.00	135.	80.		
9513.27	5.10	2100.	500.		
9626.00	5.18	2500.	630.		
9767.26	5.28	550.	250.		
9949.51	5.44	580.	260.		
9979.75	5.59	140.	80.		

TABLE A2 (cont.)

Neutron Resonance Parameters of ^{160}Gd

E_0 (eV)	ΔE_0	Γ_n (meV)	$\Delta \Gamma_n$	Γ_γ (meV)	$\Delta \Gamma_\gamma$
221.96	.14	49.	5.	120.	17.
*421.9	.5	1.5	.7		
447.9	.5	15.	6.		
478.89	.46	320.	40.	125.	15.
*571.81	.30	6.	4.		
*707.5	1.	3.8	2.		
752.59	.45	8.	4.		
904.93	.30	3100.	360.		
**1025.	1.	2.8	1.5		
1240.94	.48	3050.	360.	85.	18.
*1291.	1.	5.	2.5		
1417.91	.30	880.	100.	115.	15.
**1632.	1.	3.3	1.6		
*1695.	1.	6.	3.		
1808.35	.42	6500.	450.		
1828.9	1.2	200.	100.		
*1874.	1.	5.	3.		
1959.79	.48	250.	40.		
2268.92	.60	1450.	100.		
2396.29	.65	5500.	750.		
2516.92	.69	4500.	1000.		
*2555.	2.	16.5	8.		
2640.21	.75	3200.	800.		
2813.22	.82	510.	120.		
*2899.32	0.85	50.	30.		
**3174.	3.	27.5	13.		
3343.	3.	25.	12.		
3454.68	1.11	200.	100.		
*3563.	3.	40.	15.		
*3598.	3.	45.	20.		
3717.93	1.24	1400.	300.		
3870.20	1.32	1300.	300.		
3931.58	1.35	550.	130.		
4517.82	1.66	6100.	1500.		
4639.	4.	20.	10.		
**4667.	4.	15.	10.		
4794.	4.	50.	20.		
5018.23	2.05	2000.	500.		
5251.92	2.09	1200.	300.		
5482.27	2.23				

*p wave or spurious level

**probably spurious level

TABLE A2 (cont.)

Neutron Resonance Parameters of ^{160}Gd

E_0 (eV)	ΔE_0	Γ_n (meV)	$\Delta\Gamma_n$	Γ_γ (meV)	$\Delta\Gamma_\gamma$
5836.48	2.44	1950.	500.		
5927.89	2.50	3500.			
6390.78	2.80	2550.	650.		
6504.19	2.88	11500.	2500.		
6792.07	3.06	6300.	1600.		
6832.10	3.10	1480.	500.		
6900.65	3.20	400.	250.		
7303.64	3.43	1950.	500.		
7379.42	3.47	5600.	1400.		
7691.00	3.69	4300.	1100.		
7840.91	3.81	3100.	750.		
8305.48	4.15	3200.	750.		
8465.27	4.27	2500.	600.		
8693.70	4.44	2700.	6500.		
9095.60	4.77	3700.	900.		
9518.37	5.11	2400.	600.		
9662.38	5.20	2200.	550.		
9852.	6.				

TABLE A-3
(page 1)

Isotope	Mass % in natural element	$I_C^{0.414}$ (barns)	$\langle \Gamma_Y \rangle$ meV*	other values of I_C (barns)
^{152}Sm	26.7	2644 ± 604	65(9)	3100 ± 200^a , 3162 ± 104^b 2850 ± 300^c , 2920^f
^{154}Sm	22.7	31 ± 6	79(3)	
^{151}Eu	47.8	$5729 \pm 650^+$	90.0(70)	3741^d , $<3000^e$, 3420^f
^{153}Eu	52.2	$1745 \pm 195^+$	94.8(44)	1833^d , 1280 ± 100^e
^{154}Gd	2.1	179 ± 17	100(24)	303^g
^{158}Gd	24.9	60.5 ± 6	106(27)	72^h
^{160}Gd	21.9	6.9 ± 1.0	129(5)	4.8^h
^{166}Er	33.4	123 ± 13	92.4(10)	56.5 ± 11.3^g
^{167}Er	22.9	4899 ± 605	91(54)	
^{168}Er	27.1	35.6 ± 7	85(2)	
^{170}Er	14.9	45 ± 7		32.2^f
^{168}Yb	0.1	$34350 \pm 4500^{++}$		$30,950^i$
^{170}Yb	3.0	212 ± 20		326^i
^{171}Yb	14.3	345 ± 39	76.5(44)	313^i
^{172}Yb	21.8	26.2 ± 6	72(4)	18 ± 7^g , 23.5^i
^{174}Yb	31.8	27.1 ± 6	80(3)	33.8^i
^{176}Yb	12.7	8 ± 2	82(1)	7.6^i
^{175}Lu	97.4	1158 ± 280	60(8)	886^d
^{182}W	26.4	592 ± 60	75(2)	591^j
^{183}W	14.4	371 ± 42		387^j
^{184}W	30.6	14.1 ± 1.5		13.5^j , 7.5 ± 1.0^k , 14.9 ± 1.6^l
^{186}W	28.4	487 ± 50	47(3)	345^f , 551^j , 490 ± 80^k 441 ± 22^l , 318^m , 450 ± 36^n

*followed by the number of levels from which $\langle \Gamma_Y \rangle$ was obtained.

+the resonance integrals in Eu are complicated by low energy resonances near our cut-off energy of 0.414 eV. We have obtained the value of $I_C^{0.5} = 3265$ barns for ^{151}Eu and 1632 barns for ^{153}Eu .

++ $I^{0.5} = 31,900$ barns.

- a J. Walker, Can. J. of Phys. 39, 1184 (1961).
- b M. Cabell, J. Inorg. Nucl. Chem. 24, 749 (1962).
- c R. Tattersall, J. of Nucl. Energy A12, 32 (1960).
- d J. Rogers, Trans. Am. Nucl. Soc. 10, 259 (1967).
- e R. Tattersall, H. Rose, S. Pattenden, D. Jowitt, J. Nucl. Energy 12, 32 (1960).
- f Report: "Certain Accounts on the Utilization of the THAI Research Reactor" Bangkok Conf. THAI-AEC-10 (1967).
- g R. Dobrozemsky, A. Edlmair, F. Pichlmayer, F. Viehbock, "Pile Neutron Capture Cross Sections for Rare Earth Isotopes" EANDC (JR)-68 "L".
- h S. Mughubghub and R. Chrien, Phys. Rev. 180, 1131 (1969).
- i S. Mughubghub and R. Chrien, Phys. Rev. 174, 1400 (1968).
- j G. Joanou, C. Stevens, "Neutron Cross Sections for the W Isotopes" GA report 5885 (1964).
- k J. Scoville, Trans. Am. Nucl. Soc. 5, 377 (1962).
- l C. Pierce, D. Shook, NSE 31, 431 (1968).
- m J. Gillette, "Review of the Radioisotope Prog." ORNL-4013 (1965).
- n L. Beller, D. Latham, J. Otter, "Measurement of the Resonance Integral in ^{186}W " NAA-SR-12500 (1967).

Fig. A1. Transmission of ^{154}Gd sample with $(1/n)=344$ barns/atom. This sample was $\sim 45\%$ enriched in ^{154}Gd , $\sim 30\%$ in ^{155}Gd , and had significant amounts of ^{156}Gd and ^{157}Gd present. This figure shows the isotopic identification of the prominent resonances, with their energy parameter in eV. Less prominent resonances, not indicated for reasons of clarity, are due mainly to ^{155}Gd .

Fig. A2. (a) Number of levels versus E observed in ^{154}Gd . The sharp decrease in $N(E)$ above 300 eV is due mainly to our inability to make isotopic assignments of many weaker levels above this energy due to the presence of significant amounts of other Gd isotopes.

(b) The observed histogram of nearest neighbor spacings in ^{154}Gd and the theoretical Wigner distribution in the energy interval 0-300 eV over which we believe all, or nearly all, s-wave resonances have been identified.

Fig. A3. (a) Histogram of reduced neutron widths for ^{154}Gd observed in energy interval 0-1 keV. The Porter-Thomas curve is also shown for $n^*=70$ levels indicated from the level density in the lower energy interval 0-300 eV over which a good Δ_{exp} was obtained.

(b) $\Sigma \Gamma_n^0$ versus E . The slope of the straight line determines the s strength function, S_0 .

Fig. A4. Examples of our ^{158}Gd (lower plot) and ^{160}Gd (upper plot) results. These plots show transmission histograms for $(1/n)=240$ and 184 barns/atom respectively, with isotopic enrichments of 99% and 98%. As seen in this figure, almost no "showing through" of even the strongest resonances from the other Gd isotopes occurs.

Fig. A5. (a) Number of observed levels versus E for ^{158}Gd and ^{160}Gd . Only S levels are shown. Few, if any, levels are thought to have been missed below 4 keV. Two weak levels in ^{158}Gd and 13 weak levels in ^{160}Gd were observed in the "D only" data, which we tentatively assign as p wave.

(b) Histograms of observed level spacings & Wigner distributions for ^{158}Gd and ^{160}Gd in the energy interval 0-4 keV. Relatively good fits to the data are obtained.

Fig. A6 (a) Porter-Thomas distribution compared to the observed distribution of reduced neutron widths. The P-T distribution has been normalized to S_0 and the level density in the 0-4 keV energy region.

(b) $\Sigma \Gamma_n^0$ versus E for ^{158}Gd . The slope gives $S_0 = (1.5 \pm 0.2) \times 10^{-4}$ as the experimental s strength function.

Fig. A7 (a) Porter-Thomas distribution and observed reduced neutron widths in the interval 0-4 keV. There are 13 additional "very weak" levels observed which appear in the first histogram box which are likely to be p waves or spurious levels.

(b) $\Sigma \Gamma_n^0$ versus E for ^{160}Gd . The two lines shown give nearly extreme possibilities of S that are consistent with our data. Our best choice is $S_0 = (1.8 \pm 0.4) \times 10^{-4}$, which gives a somewhat stronger weighting to the 0-4 keV energy region.

Figure A-8. The measured total neutron cross section for natural iron in selected energy regions. The σ_T shown in this figure is influenced by resolution effects (effective resolution of the experiment = 0.6 nsec/meter). These resolution effects are greater where σ_T shows a rapid recovery from σ_{min} in terms of our timing channel widths.

Figure A-8. The measured total neutron cross section for natural iron in selected energy regions. The σ_T shown in this figure is influenced by resolution effects (effective resolution of the experiment = 0.6 nsec/meter). These resolution effects are greater where σ_T shows a rapid recovery from $\sigma_{\min}$ in terms of our timing channel widths.

(page 2)

Figure A-8. The measured total neutron cross section for natural iron in selected energy regions. The σ_T shown in this figure is influenced by resolution effects (effective resolution of the experiment = 0.6 nsec/meter). These resolution effects are greater where σ_T shows a rapid recovery from $\sigma_{\min}$ in terms of our timing channel widths.

(page 3)

Figure A-8. The measured total neutron cross section for natural iron in selected energy regions. The σ_T shown in this figure is influenced by resolution effects (effective resolution of the experiment ≈ 0.6 nsec/meter). These resolution effects are greater where σ_T shows a rapid recovery from $\sigma_{\min}$ in terms of our timing channel widths.

(page 4)

Fig. B1. Total cross section of natural sodium from 20 to 360 keV. This experimental σ_T is subject to effects which are dependent on the thickness of our samples and the resolution of the NVS time of flight spectrometer.

Fig. B1 (continued). Total cross section of natural sodium, 20-360 keV.

Fig. B2. The total cross section of natural copper in the energy interval 60 - 2.5 keV.

Fig. B2 (continued). Total cross section of natural copper in the energy interval 60 to 2.5 keV.

TABLE C-1
(page 1)G Values of U^{238} Calculated with $M = 12$

E(eV)	$\sqrt{\Gamma_n^0} \text{ (meV)}^{\frac{1}{2}}$	G (Old)	G (New)
6.68	0.77	4.36*	0.90*
Q 13.20P	0.02	4.00*	
21.00	1.38	2.77*	0.83*
36.70	2.27	1.60*	0.54*
66.30	1.76	1.12*	0.44*
80.77	0.48	1.44*	0.40*
Q 92.00P	0.09	1.44*	
102.78	2.55	0.80*	0.15*
116.93	1.82	0.01*	-0.14*
145.80	0.26	-1.24*	-0.70*
165.54	0.52	-1.67*	-0.79*
190.34	3.30	-2.18*	-0.94*
208.65	1.97	-2.52*	-1.05*
237.40	1.34	-2.03	-1.04*
Q242.88P	0.10	-2.04	
Q263.94	0.12	-2.13	-0.70
273.74	1.23	-2.32	-0.63
291.11	0.95	-2.86	-0.98
311.12	0.24	-3.64	-1.37
347.92	2.10	-3.84	-1.68
376.92	0.24	-3.36	-1.18
397.56	0.55	-2.73	-0.54
410.23	0.97	-2.61	-0.45
434.19	0.63	-2.09	-0.41
Q454.17	0.14	-1.59	-0.03
463.31	0.49	-1.32	-0.01
478.70	0.37	-1.42	-0.46
Q488.89P	0.14	-1.60	
518.59	1.38	-1.80	-0.80
535.49	1.26	-1.18	-0.54
Q556.05	0.14	-0.84	-0.39
580.20	1.06	-0.05	-0.09
595.15	1.83	0.46	-0.08
619.94	1.07	1.90	0.32
Q623.53P	0.13	2.20	
628.67	0.40	1.79	0.20
661.18	2.12	1.16	-0.56
Q677.00P	0.14	1.73	
693.23	1.14	2.22	0.15
708.46	0.84	3.10	0.56
721.80	0.22	4.04	0.82
Q730.10P	0.17	4.99	
732.26	0.22	5.02	0.54
Q742.95P	0.14	3.94	
765.05	0.49	3.32	0.43
779.14	0.24	3.23	0.71
790.88	0.42	2.82	0.57
821.58	1.43	2.81	0.19

E (eV)	$\sqrt{\Gamma_n^0}$ (meV) ^{1/2}	G (Old)	G (New)
Q845.62P	0.14	4.90	
851.02	1.38	5.46	0.98
855.15	1.66	5.38	1.14
865.52	0.37	4.83	0.61
Q891.29P	0.17	5.08	
905.11	1.22	6.33	0.04
909.90P	0.17	6.53	
925.18	0.53	6.56	0.44
Q932.50P	0.10	6.84	
935.87	2.19	6.54	0.50
958.43	2.26	5.37	-0.00
991.78	3.32	6.65	0.18
1000.3P	0.20	7.29	
1011.2	0.24	7.91	0.87
1023.0	0.45	8.61	1.24
1029.1	0.32	8.86	1.10
Q1033.2P	0.14	8.61	
1053.9	1.52	7.66	0.05
1058.1P	0.14	8.62	
Q1070.5P	0.10	8.71	
Q1081.1P	0.14	8.10	
Q1094.8P	0.14	8.52	
1098.3	0.67	8.65	0.56
Q1102.3P	0.14	8.23	
1108.9	0.95	7.03	0.65
1131.4	0.24	4.69	1.09
1140.4	2.55	4.02	1.09
1167.5	1.53	2.44	1.15
1177.6	1.36	1.99	1.22
1195.0	1.63	0.80	1.09
1210.9	0.51	-0.26	0.77
1245.1	2.55	-1.10	0.56
1267.0	0.87	-0.91	0.96
1273.2	0.89	-1.05	0.81
1298.4	0.28	-2.32	0.33
1317.2	0.33	-3.00	0.42
1335.7	0.17	-3.20	0.23
1353.0N			-0.23
1393.0	1.92	-1.01	0.25
1405.1	1.43	0.59	0.62
Q1410.0P	0.17	1.09	
Q1417.0P	0.17	1.42	
1419.6	0.50	1.40	0.72
1427.7	0.89	0.35	0.45
1444.1	0.75	-0.93	-0.44
1473.0	1.43	-2.32	-1.43
1523.1	2.35	-1.04	-1.14
1532.0	0.22	-0.46	-0.95

E(eV)	$\sqrt{\Gamma_n^0} \text{ (meV)}^{\frac{1}{2}}$	G (Old)	G (New)
Q1546.0P	0.14	0.24	
1550.0	0.17	0.23	-1.08
1565.0	0.22	-0.56	-1.25
1598.2	2.83	-1.19	-1.69
1622.9	1.45	-1.01	-1.38
1638.2	1.00	-0.46	-1.32
Q1545.4P	0.14	-0.32	
1662.1	2.00	-0.75	-1.69
1688.3	1.38	-0.72	-1.63
Q1700.7F	0.14	-0.34	
1709.4	1.16	-0.46	-1.35
1723.0	0.57	-1.05	-1.39
1744.0P	0.20	-2.06	
1755.8	1.22	-2.79	-1.36
1782.3	3.32	-3.50	-1.12
1797.7	0.22	-3.82	-0.74
1808.3	0.63	-4.56	-0.99
1845.6	0.56	-6.80	-1.23
1868.0N			0.10
1870.0N			0.06
1902.3	0.69	-7.93	-1.12
1917.1	0.71	-8.24	-1.15
1953.0N			-1.05
1968.7	3.61	-7.56	-0.58
1974.6	3.24	-7.45	-0.81
2023.6	2.12	-6.61	-0.76
2031.1	1.05	-6.29	-0.73
2070.0N			-0.55
2088.6	0.55	-3.29	0.25
2096.5	0.47	-2.79	0.22
2124.3	0.32	-1.54	0.12
2145.9	0.87	0.41	0.84
2152.8	1.95	1.00	0.74
2172.0P	0.22	1.77	
2186.0	2.79	2.57	0.70
Q2194.0F	0.22	3.12	
2201.4	1.55	3.03	0.92
2230.0	0.32	3.98	1.79
2235.7	0.32	3.41	1.92
Q2241.5F	0.17	4.31	
2259.1	1.17	4.45	2.10
2266.4	1.75	4.48	2.05
2281.3	1.52	4.23	1.24
2298.7P	0.22	4.07	
Q2302.0F	0.14	3.14	
231 .9	0.55	2.41	0.87
2337.4	0.32	1.55	1.20
2352.0	1.14	1.43	1.72

TABLE C-2

 ^{232}Th Resonance Parameters

(page 1)

$E_0(\text{eV})$	ΔE	$\Gamma_n^0(\text{meV})$	$\Delta\Gamma_n^0$	Γ_γ (meV)	$\Delta\Gamma_\gamma$	$E_0(\text{eV})$	ΔE	$\Gamma_n^0(\text{meV})$	$\Delta\Gamma_n^0$	Γ_γ (meV)	$\Delta\Gamma_\gamma$
21.78	.02	.41	.02	20. 2.		369.43	.31	1.30	.08	22. 2.	
23.43	.02	.67	.05	25. 2.		380.40	.33*	.006	.002		
58.84	.07*	.001	.001			391.53	.34	.006	.002		
59.48	.08	.51	.03	25. 2.		400.85	.35	.52	.05	18. 2.	
69.17	.10	5.29	.24	25. 2.		402.62	.36*	.004	.002		
90.08	.08*	.001	.001			411.62	.36	.007	.002		
112.93	.11	1.27	.09	20. 2.		420.92	.38	.02	.005		
120.78	.11	2.18	.14	22. 2.		454.1	.4	.05	.01		
128.21	.12*	.004	.001			462.3	.4	3.02	.23	22. 2.	
129.10	.13	.30	.02	18. 2.		476.3	.4	.01	.005		
5.72	.15*	.007	.002			488.6	.4	2.71	.18	18. 2.	
4.24	.15	.01	.002			510.31	.25	.17	.04		
170.34	.19	4.98	.38	26. 2.		528.46	.26	.52	.04	20. 3.	
178.62	.21*	.004	.001			533.55	.27*	.010	.003		
192.57	.23	1.33	.07	17. 2.		535.45	.27*	.01	.003		
196.13	.24	.005	.001			540.09	.28	.04	.004		
199.30	.24	.89	.07	18. 2.		569.87	.30	1.21	.08	19. 2.	
202.41	.25*	.002	.001			573.45	.30	.03	.01		
210.87	.25*	.001	.001			578.00	.30	.08	.02		
219.30	.28*	.001	.001			598.16	.32	.39	.04	19. 2.	
221.15	.29	2.08	.13	22. 2.		617.84	.33	.18	.02		
242.23	.16*	.002	.001			656.41	.35	1.99	.16	20. 2.	
251.48	.18	2.02	.13	24. 2.		660.66	.37*	.008	.004		
262.96	.19	1.48	.09	19. 2.		665.15	.38	.97	.06	18. 2.	
285.74	.21	1.77	.10	20. 2.		675.22	.39	7.89	.58	19. 2.	
290.12	.21*	.002	.001			687.3	.4	1.72	.15	23. 2.	
302.30	.23*	.006	.002			700.9	.4	.64	.08	17. 4.	
305.43	.24	1.49	.11	20. 2.		712.8	.4	.82	.11	19. 3.	
309.20	.24*	.003	.001			740.9	.4	6.98	.55	23. 2.	
321.47	.25*	.002	.001			764.7	.4	.03	.01		
328.92	.26	4.19	.33	26. 2.		771.7	.4*	.005	.003		
336.26	.27*	.003	.001			778.5	.4	.39	.03	26. 3.	
341.83	.28	1.95	.11	19. 2.		804.1	.5	6.35	.46	20. 2.	
361.47	.31*	.004	.002			820.9	.5	.03	.01		
365.11	.31	1.47	.08	21. 2.		836.6	.5	.04	.01		

*p levels or spurious

TABLE C-2 (CONTINUED)

(page 2)

E_0 (eV)	ΔE	Γ_n^0 (meV)	$\Delta\Gamma_n^0$	Γ_γ (meV)	$\Delta\Gamma_\gamma$	E_0 (eV)	ΔE	Γ_n^0 (meV)	$\Delta\Gamma_n^0$	Γ_γ (meV)	$\Delta\Gamma_\gamma$
842.2	.5	.96	.10	19. 2.		1233.3	.4 *	.02	.01		
846.7	.5 *	.005	.002			1243.1	.4 *	.54	.06	20. 3.	
850.5	.5	.03	.01			1248.6	.4	3.31	.45	15. 3.	
866.2	.5	.44	.07	23. 3.		1260.8	.5 *	.02	.01		
869.3	.5 *	.02	.01			1261.7	.5 *	.03	.01		
884.7	.5	.006	.003			1269.4	.5	.55	.04	20. 3.	
889.91	.29	1.27	.10	21. 2.		1287.84	.25 *	.01	.01		
905.97	.30	.05	.01			1291.90	.25	2.89	.39	25. 2.	
919.26	.30 *	.010	.005			1301.61	.26	1.44	.14	21. 2.	
926.47	.31	.007	.003			1335.71	.27	.10	.02		
934.24	.31 *	.007	.003			1345.65	.27 *	.03	.002		
943.22	.31	1.47	.16	23. 2.		1349.44	.27 *	.02	.01		
962.55	.33	.17	.02			1354.89	.27	2.58	.30	26. 2.	
982.65	.33	1.02	.10	21. 2.		1359.82	.28	.28	.02		
990.46	.34	2.22	.16	25. 4.		1372.54	.28	.02	.01		
1010.39	.35	3.55	.47	20. 3.		1377.85	.28	1.43	.19	24. 2.	
1021.36	.35 *	.01	.003			1384.59	.29 *	.008	.005		
1039.13	.37	.45	.05			1387.70	.29	.05	.01		
1043.74	.37 *	.02	.005			1397.95	.29	3.61	.29	19. 2.	
1064.54	.38	.13	.02			1426.43	.30	2.70	.24	21. 3.	
1073.73	.39 *	.006	.003			1433.84	.30	.92	.11	18. 3.	
1077.20	.39	.37	.06			1441.30	.30 *	.03	.01		
1093.26	.39	.05	.01			1461.28	.31	.03	.01		
1109.6	.4	.69	.05	17. 3.		1469.27	.31 *	.008	.005		
1114.9	.4	.05	.01			1476.15	.31	.05	.01		
1115.9	.4 *	.03	.01			1508.60	.32	.02	.01		
1120.2	.4	.08	.03			1509.56	.32 *	.09	.03		
1132.7	.4 *	.009	.003			1518.60	.33	4.62	.64	24. 3.	
1138.7	.4	.46	.03	19. 3.		1524.45	.33	4.99	.51	20. 3.	
1150.2	.4	.59	.05	22. 2.		1555.56	.34	.17	.02		
1194.4	.4	.24	.03			1581.41	.35	.53	.04	20. 4.	
1204.1	.4	.02	.01			1589.53	.35	7.27	1.00	24. 2.	
1217.3	.4 *	.02	.01			1602.63	.35	1.22	.15	24. 2.	
1223.8	.4 *	.01	.01			1610.47	.35 *	.02	.004		
1227.8	.4	.84	.05	23. 3.		1630.94	.35	12.63	1.24	19. 5.	

*p levels or spurious

TABLE C-2 (CONTINUED)

(page 3)

E_0 (eV)	ΔE	Γ_n^0 (meV)	$\Delta\Gamma_n^0$	Γ_γ (meV)	$\Delta\Gamma_\gamma$	E_0 (eV)	ΔE	Γ_n^0 (meV)	$\Delta\Gamma_n^0$	Γ_γ (meV)	$\Delta\Gamma_\gamma$
1640.35	.37	1.09	.12	25. 2.		2147.6	.5	1.92	.15	14. 4.	
1661.48	.37	2.80	.29	25. 4.		2158.5	.5*	.04	.01		
1677.73	.39	.61	.07			2162.9	.5	1.87	.15	26. 4.	
1689.65	.38*	.04	.01			2170.1	.5*	.06	.02		
1696.90	.38	.04	.01			2177.9	.5	1.59	.13	17. 3.	
1704.97	.38	.06	.01			2196.9	.5	1.00	.11	17. 3.	
1719.72	.39	.82	.07	17. 3.		2206.8	.5*	.04	.01		
1729.13	.39	.03	.01			2216.2	.5	.49	.04		
1740.2	.4	.14	.02			2222.0	.5	1.80	.19	27. 5.	
1746.5	.4	.72	.07	23. 4.		2233.5	.5	.03	.01		
1762.6	.4	2.43	.40	27. 4.		2270.9	.5	.59	.08		
1767.1	.4*	.04	.02			2276.3	.6	.96	.10	24. 3.	
1785.4	.4	.05	.02			2286.4	.6	5.44	.63	19. 5.	
1803.3	.4	2.00	.16	21. 3.		2307.2	.6*	.06	.02		
1811.9	.4	1.06	.14	20. 3.		2320.9	.6	.09	.03		
1824.0	.4	1.94	.16	18. 3.		2329.5	.6*	.04	.01		
1848.6	.4	.13	.02			2336.0	.6	2.28	.37	20. 4.	
1854.4	.4	1.04	.16	25. 3.		2344.4	.6	.09	.04		
1861.9	.4	.95	.14	22. 3.		2352.5	.6	.31	.06		
1897.1	.4*	.05	.02			2353.7	.6	.29	.06		
1899.9	.4	2.87	.32	29. 5.		2375.2	.6	2.42	.37	18. 3.	
1928.3	.4	.15	.02			2382.8	.6	.04	.02		
1931.1	.4	.22	.03			2391.1	.6	.07	.02		
1950.3	.4	2.88	.38	30. 4.		2418.5	.6	1.75	.18		
1971.2	.4	5.52	.52	25. 4.		2427.4	.6*	.05	.04		
1987.8	.4	.96	.11	18. 3.		2434.6	.6*	.06	.04		
2004.9	.4	.51	.07	22. 4.		2439.8	.6	.21	.03		
2015.4	.4*	.02	.01			2455.8	.6	3.53	.44		
2051.7	.5	.42	.07			2462.5	.6*	.07	.03		
2055.5	.5*	.01	.01			2474.9	.6	.04	.02		
2061.9	.5	1.43	.18	17. 3.		2491.8	.6	.13	.03		
2073.8	.5	.15	.04			2509.3	.6	6.29	.70		
2079.0	.5	.20	.04			2527.3	.6	.93	.12		
2097.3	.5	.02	.01			2557.1	.7	.07	.02		
2116.9	.5	1.63	.13	20. 3.		2563.1	.7	6.12	.59		

*p levels or spurious

TABLE C-2 (CONTINUED)

(page 4)

E_0 (eV)	ΔE	Γ_n^0 (meV)	$\Delta\Gamma_n^0$	E_0 (meV)	ΔE	Γ_n^0 (meV)	$\Delta\Gamma_n^0$
2569.6	.7	1.18	.20	3039.7	.9	1.07	.09
2604.0	.7 *	.03	.02	3050.2	.9	.09	.05
2612.5	.7	1.70	.20	3061.4	.9	.52	.05
2624.3	.7	.14	.04	3082.8	.9	.99	.11
2635.0	.7	3.12	.35	3103.6	.9	.03	.01
2655.5	.7	.09	.04	3109.2	.9	.39	.05
2663.4	.7	3.88	.39	3120.7	.9	.13	.04
2677.0	.7	.23	.04	3148.5	.9	3.30	.36
2688.4	.7	3.80	.39	3153.4	.9	3.47	.36
2713.0	.7	1.84	.23	3163.0	.9	.37	.07
2722.7	.7	.22	.04	3188.6	.9	1.33	.12
2733.2	.7	6.89	.57	3194.5	.9	.07	.05
2746.1	.7	.29	.06	3207.4	.9	1.59	.18
2775.6	.8	1.52	.25	3229.4	1.0	.30	.05
2792.9	.8	3.12	.36	3242.5	1.0	.25	.05
2803.4	.8 *	.08	.02	3252.7	1.0	1.46	.14
2815.6	.8	.55	.08	3270.0	1.0	.45	.07
2832.9	.8	.70	.09	3295.7	1.0	7.11	.78
2838.7	.8	.03	.03	3307.2	1.0	.07	.07
2843.2	.8 *	.03	.03	3317.6	1.0	.09	.05
2852.0	.8	3.93	.56	3331.8	1.0	.71	.10
2861.2	.8 *	.15	.06	3342.9	1.0	2.94	.31
2870.4	.8 *	.03	.01	3351.4	1.0	.31	.07
2883.9	.8	.08	.03	3383.5	1.0	1.20	.15
2895.9	.8	.04	.01	3409.5	1.0	.08	.03
2915.6	.8	.06	.02	3442.9	1.1	.34	.05
2932.0	.8 *	.03	.02	3471.9	1.1	.31	.05
2948.6	.8	1.77	.18	3510.0	1.1	.08	.05
2956.5	.8	.81	.11	3521.8	1.1	1.79	.30
2966.2	.8	.26	.06	3544.2	1.1	.12	.05
2980.0	.8	.16	.04	3574.4	1.1	.28	.05
2988.0	.9	.68	.11	3594.4	1.1	.35	.07
3006.5	.9	.03	.01	3611.6	1.1	2.00	.25
3017.3	.9	.51	.05	3625.5	1.1	.20	.05
3027.3	.9	4.18	.45	3636.7	1.2	.13	.05

*p levels or spurious

TABLE C-2 (CONTINUED)

(page 5)

$E_0(\text{meV})$	ΔE	$\Gamma_n^0(\text{meV})$	$\Delta\Gamma_n^0$
3651.7	1.2	1.11	.13
3674.2	1.2	.20	.05
3692.7	1.2	.10	.07
3707.6	1.2	.08	.05
3716.0	1.2	.43	.07
3722.9	1.2	1.61	.33
3732.8	1.2	.79	.11
3745.2	1.2	.08	.05
3759.2	1.2	.13	.05
3786.4	1.2	.47	.08
3799.4	1.2	.08	.05
3820.5	1.2	.65	.10
3827.0	1.3	1.76	.27
3848.5	1.3	.19	.06
3868.8	1.3	1.05	.13
3883.4	1.3	.18	.05
3906.0	1.3	3.52	.48
3923.4	1.3	.14	.05
3931.5	1.3	.59	.10
3951.1	1.3	.16	.06
3961.4	1.3	.65	.10
3970.3	1.3	1.14	.14
3976.5	1.3	1.90	.27
3994.4	1.3	.57	.09
4007.6	1.4	.11	.06

TABLE C-3. Neutron Resonance parameters for assigned p levels in ^{232}Th .

E_0 (eV)	ΔE_0	$g\Gamma_n^1$ (meV)	$\Delta g\Gamma_n^1$	E_0 (eV)	ΔE_0	$g\Gamma_n^1$ (meV)	$\Delta g\Gamma_n^1$	E_0 (eV)	ΔE_0	$g\Gamma_n^1$ (meV)	$\Delta g\Gamma_n^1$	E_0 (eV)	ΔE_0	$g\Gamma_n^1$ (meV)	$\Delta g\Gamma_n^1$
58.84	0.07	3.6	1.8	380.40	0.33	4.0	1.3	1217.3	0.4	3.9	1.4	1897.1	0.4	6.5	2.6
90.08	0.08	3.2	1.6	402.62	0.36	3.0	1.5	1223.8	0.4	2.6	1.2	2015.4	0.4	2.4	1.3
128.21	0.12	8.5	2.0	533.55	0.27	4.9	1.7	1233.3	0.4	4.8	1.7	2055.5	0.5	1.5	0.8
145.72	0.15	14.1	3.0	535.45	0.27	7.1	2.0	1260.8	0.5	3.7	1.3	2158.5	0.5	5.5	2.3
178.62	0.21	6.9	2.2	660.66	0.37	3.2	1.3	1261.7	0.5	5.6	1.8	2170.1	0.5	7.9	2.9
202.41	0.25	2.8	1.4	771.7	0.4	1.9	1.0	1287.84	0.25	3.0	1.3	2206.8	0.5	5.6	2.4
210.87	0.26	1.8	0.9	846.7	0.5	1.6	0.8	1345.65	0.27	5.3	2.0	2307.2	0.5	7.6	2.9
219.30	0.28	1.7	0.9	869.3	0.5	7.7	2.8	1349.44	0.27	4.4	1.6	2329.5	0.6	4.9	2.1
242.23	0.16	2.1	1.1	919.26	0.30	2.9	1.4	1384.59	0.29	1.6	0.9	2427.4	0.6	5.5	2.4
290.12	0.21	2.2	1.1	934.24	0.31	1.9	0.9	1441.30	0.30	6.1	2.5	2434.6	0.6	6.8	3.0
302.30	0.23	5.2	2.2	1021.38	0.35	2.9	1.3	1469.27	0.31	1.4	0.8	2462.5	0.6	8.0	3.2
309.20	0.24	3.0	1.4	1043.74	0.37	5.3	2.0	1509.56	0.32	16.6	3.4	2604.0	0.7	3.3	1.7
321.47	0.25	1.9	1.0	1073.73	0.39	1.6	0.8	1610.47	0.36	3.5	1.5	2803.4	0.8	8.5	3.5
338.26	0.26	2.6	1.3	1115.9	0.4	8.0	2.4	1689.65	0.38	5.9	2.1	2843.2	0.8	2.7	1.4
361.47	0.31	3.2	1.6	1132.7	0.4	2.2	1.1	1767.1	0.4	5.5	2.2	2861.2	0.8	14.5	5.0
												2870.4	0.8	2.7	1.5
												2932.0	0.8	2.7	1.5

TABLE C-4. Neutron resonance parameters for assigned p levels in ^{238}U .

E_0 (eV)	ΔE_0	$g\Gamma_n^1$ (meV)	$\Delta g\Gamma_n^1$	E_0 (eV)	ΔE_0	$g\Gamma_n^1$ (meV)	$\Delta g\Gamma_n^1$	E_0 (eV)	ΔE_0	$g\Gamma_n^1$ (meV)	$\Delta g\Gamma_n^1$	E_0 (eV)	ΔE_0	$g\Gamma_n^1$ (meV)	$\Delta g\Gamma_n^1$
10.22	0.01	14.6	3.0	337.19	0.27	2.2	1.1	846.9	0.4	11.0	3.3	1256.5	0.4	1.2	0.9
19.50	0.02	4.3	1.8	351.75	0.30	3.3	1.5	890.6	0.3	8.0	2.7	1371.6	0.3	3.2	1.7
45.19	0.07	1.8	0.9	354.66	0.31	1.2	0.8	909.5	0.3	12.7	3.5	1381.6	0.3	3.1	1.6
63.54	0.08	3.2	1.4	407.64	0.36	2.7	1.4	932.3	0.3	2.9	1.6	1410.5	0.3	2.0	1.2
83.57	0.08	1.4	0.7	439.71	0.39	4.7	2.1	940.1	0.3	2.9	1.5	1416.3	0.3	8.9	2.7
89.19	0.08	28.8	3.5	488.20	0.40	11.2	3.4	964.9	0.3	1.8	1.1	1549.8	0.3	5.3	2.3
121.61	0.11	1.2	0.7	498.93	0.25	2.0	1.1	976.8	0.3	5.3	1.9	1646.1	0.3	3.9	2.1
124.30	0.12	2.3	1.2	523.21	0.26	4.5	2.0	985.6	0.3	2.6	1.4	1744.9	0.4	7.3	2.6
152.42	0.16	5.7	2.2	542.34	0.28	1.1	0.8	1014.4	0.3	13.2	3.0	1912.6	0.4	1.6	1.0
158.89	0.17	0.9	0.7	606.12	0.33	4.5	2.2	1031.1	0.3	8.1	2.7	2175.2	0.5	3.9	2.0
173.11	0.19	3.0	1.3	624.80	0.34	17.3	3.8	1062.3	0.3	5.4	2.5	2288.9	0.6	2.2	1.2
202.30	0.25	3.7	1.5	668.40	0.37	3.9	1.8	1067.6	0.3	7.7	2.1	2798.1	0.8	3.7	1.9
214.97	0.27	3.4	1.4	677.50	0.39	10.6	3.1	1071.0	0.3	2.4	1.3	2907.1	0.8	2.6	1.5
242.60	0.16	10.6	2.4	712.49	0.40	3.6	1.5	1081.0	0.3	5.3	2.5	2922.1	0.8	6.8	2.7
253.88	0.18	6.6	2.0	729.4	0.4	9.4	2.9	1094.4	0.4	9.6	2.9				
255.37	0.19	4.0	1.7	743.2	0.4	3.9	1.6	1102.7	0.4	14.6	3.5				
257.10	0.19	1.3	0.8	756.0	0.4	5.7	2.6	1147.0	0.4	2.1	1.1				
275.76	0.20	4.7	2.1	808.2	0.4	4.7	2.3	1154.8	0.4	2.8	1.4				
282.29	0.21	3.5	1.5	815.3	0.4	2.3	1.2	1217.9	0.4	2.5	1.4				
294.96	0.22	1.6	0.9	832.4	0.4	2.9	1.5	1237.9	0.4	2.5	1.3

Figure C-1. Histogram of the reduced widths $\sqrt{g\Gamma_n^0}$ for the selected $\ell=0$ population for ^{238}U . The Porter-Thomas theory is given by the smooth curve.

Figure C-2. Histogram of adjacent level spacings for the selected ^{238}U $l=0$ population. The smooth curve represents the Wigner distribution.

Fig. C-3

Example of the area analysis for the $E_0 = 675 \text{ eV}$ level of ^{232}Th . The common intersection of the curves gives the values of the parameters $\Gamma_n = 205 \pm 15 \text{ meV}$ and $\Gamma_\gamma = 19 \pm 2 \text{ meV}$.

Fig. C-4

Example of the shape analysis for the $E_0 = 675$ eV level in ^{232}Th . The series of curves are derived from the sum (R^+) and difference (R^-) of the cross section for points equidistant in energy from E_0 . The cross section points are chosen in the wings of the resonance where the cross section is slowly varying and where resolution effects are less pronounced. The intersection of the points give further information on $\Gamma_n = 205 \pm 15$ meV and the (local) potential scattering radius R .

Figure C-5. The values of Γ_γ versus energy for 84 levels up to 2400 eV.

Figure C-6. The distribution of observed Γ_γ values in ^{232}Th taken in histogram intervals of 2 meV. The average $\langle \Gamma_\gamma \rangle$ is 21.2 meV. The curve is a chi-squared distribution with $\nu=87$ degrees of freedom.

Figure C-7. The $\Sigma g \Gamma_n^0$ versus E for ²³²Th. The slope of the curve determines the s wave strength function. Our best choice is $S_0 = (0.84 \pm 0.08) \times 10^{-4}$.

Figure C-8. Histogram of the reduced neutron widths $\sqrt{\Gamma_n^0}$ for our selection of the $\ell=0$ population in ^{232}Th . The Porter-Thomas theory is represented by the smooth curve.

Figure C-9. Histogram of adjacent level spacings for the $l=0$ population in ^{232}Th . The Wigner distribution is given by the curve.

TABLE D1

Resonance Parameters of ^{233}U (single level fits)

E_0	Γ_T	NVS $\sigma_0\Gamma_F(\text{eVb})$	Saclay ¹
1.55	600	127.95 $\pm$ 2.55	
1.79	240	161.07 $\pm$ 1.68	301
2.16	180	22.69 $\pm$ 1.07	
2.28	70	39.98 $\pm$ 0.84	54
3.30	550	20.73 $\pm$ 1.58	
3.60	180	27.07 $\pm$ 0.95	45
4.75	800	55.61 $\pm$ 3.35	63.5
5.88	300	25.29 $\pm$ 1.37	43.5
6.44	500	44.72 $\pm$ 2.17	
6.79	150	104.85 $\pm$ 1.81	130
7.46	200	1.56 $\pm$ 0.87	8.3
8.69	650	4.86 $\pm$ 1.83	5.3
9.33	350	19.66 $\pm$ 1.44	13.8
10.35	350	169.79 $\pm$ 2.71	163
11.20	350	18.04 $\pm$ 1.63	
11.69	200	5.79 $\pm$ 1.18	
12.22	500	18.13 $\pm$ 2.19	16.9
12.73	300	113.41 $\pm$ 2.62	122
13.66	320	31.02 $\pm$ 1.88	37
15.34	240	61.03 $\pm$ 2.15	47
15.84	250	10.57 $\pm$ 2.98	
16.13	200	25.18 $\pm$ 2.18	107
16.50	300	61.32 $\pm$ 2.40	12.8
17.93	200	14.52 $\pm$ 1.45	14.7
18.47	250	16.90 $\pm$ 1.81	9.1
18.91	300	99.49 $\pm$ 2.78	97
20.52	400	51.99 $\pm$ 2.45	51.5
21.75	200	51.10 $\pm$ 2.73	32.4
22.28	480	180.90 $\pm$ 4.92	199
22.89	760	18.16 $\pm$ 4.09	
23.67	500	29.72 $\pm$ 2.77	33
25.20	380	38.24 $\pm$ 2.63	43.7
25.86	360	7.33 $\pm$ 2.07	5.1
26.54	480	21.35 $\pm$ 2.68	8.3
27.55	400	5.26 $\pm$ 2.25	26
28.17	220	10.69 $\pm$ 1.89	27
29.00	480	56.25 $\pm$ 3.43	64
29.54	200		

1. Nifenecker, H. Jour. de Physique 25, 877 (1964).

TABLE D2

Resonance Parameters of ^{233}U (multilevel fits)

μ (eV)		ν (eV)		G_f (beV $^{3/2}$)		H_f (beV $^{3/2}$)	
NVS	ORNL ¹	NVS	ORNL ¹	NVS	ORNL ¹	NVS	ORNL ¹
1.55	1.43	0.30	0.29	78.15 $\pm$ 2.71	50.86	4.33 $\pm$ 2.87	-18.87
1.79	1.78	0.12	0.12	104.05 $\pm$ 1.93	114.26	-7.30 $\pm$ 1.47	-0.23
2.28	2.29	0.05	0.05	39.60 $\pm$ 0.71	42.74	14.76 $\pm$ 0.56	18.48
3.30	3.37	0.275	0.35	18.98 $\pm$ 2.31	28.12	1.01 $\pm$ 2.24	-13.61
3.60	3.63	0.09	0.08	24.32 $\pm$ 1.22	22.02	1.30 $\pm$ 1.15	6.80
4.75	4.61	0.40	0.40	62.28 $\pm$ 4.09	62.53	1.51 $\pm$ 3.76	-35.08
5.88	5.80	0.15	0.15	28.83 $\pm$ 7.21	20.14	-1.34 $\pm$ 2.35	-17.06
6.44		0.25		52.28 $\pm$ 4.61		6.70 $\pm$ 4.86	
6.79	6.80	0.075	0.09	140.36 $\pm$ 2.91	176.62	1.72 $\pm$ 2.51	31.33
7.46	7.47	0.10	0.10	3.57 $\pm$ 1.39	6.78	-1.17 $\pm$ 1.38	-0.67
8.68	8.69	0.32	0.28	6.19 $\pm$ 3.97	22.67	-2.85 $\pm$ 3.68	4.82
9.33	9.14	0.175	0.15	23.62 $\pm$ 3.07	17.48	-3.51 $\pm$ 3.09	-20.48
10.35	10.35	0.175	0.16	266.03 $\pm$ 5.37	275.89	14.40 $\pm$ 4.18	0.42
11.20	11.30	0.175	0.20	31.42 $\pm$ 4.27	33.41	5.06 $\pm$ 4.02	8.67
11.69		0.10		10.15 $\pm$ 3.18		0.63 $\pm$ 3.02	
12.22		0.25		30.74 $\pm$ 7.45		5.78 $\pm$ 6.97	
12.73	12.76	0.15	0.17	206.74 $\pm$ 6.16	238.89	3.22 $\pm$ 6.50	49.43
	13.45		0.20		41.96		14.53
13.66	13.65	0.16	0.14	59.34 $\pm$ 3.97	44.99	0.89 $\pm$ 3.34	-26.72
15.34	15.28	0.12	0.12	115.92 $\pm$ 5.17	119.77	-5.06 $\pm$ 4.79	-35.47
15.84		0.125		19.24 $\pm$ 7.99		16.10 $\pm$ 8.34	
16.13	16.13	0.10	0.22	58.63 $\pm$ 8.86	101.25	-7.61 $\pm$ 8.07	-4.79
16.50	16.52	0.15	0.13	127.05 $\pm$ 7.47	94.54	3.09 $\pm$ 6.76	26.62
17.93	17.93	0.10	0.12	35.81 $\pm$ 3.96	43.16	-1.46 $\pm$ 3.69	4.22
18.47	18.42	0.125	0.21	42.07 $\pm$ 5.79	58.68	-15.12 $\pm$ 6.49	1.83
18.91	18.86	0.15	0.15	204.74 $\pm$ 7.65	216.24	-15.29 $\pm$ 6.49	-46.49
20.52	20.53	0.20	0.20	100.30 $\pm$ 6.66	111.51	9.41 $\pm$ 5.93	6.53
21.75	21.85	0.10	0.13	102.85 $\pm$ 8.62	130.22	16.71 $\pm$ 8.23	86.08
22.28	22.23	0.24	0.24	453.40 $\pm$ 19.88	532.22	31.96 $\pm$ 17.96	-77.59
22.89	22.94	0.38	0.38	78.15 $\pm$ 19.98	96.58	5.27 $\pm$ 21.07	3.64
23.67	23.54	0.25	0.32	80.43 $\pm$ 9.67	65.49	-11.39 $\pm$ 9.73	-94.87
25.20	25.13	0.19	0.19	98.97 $\pm$ 9.05	80.77	-1.65 $\pm$ 8.14	-38.01
25.86	26.19	0.18	0.18	16.57 $\pm$ 9.07	-0.73	-1.80 $\pm$ 8.59	16.46
26.54	26.57	0.24	0.24	58.96 $\pm$ 10.49	70.76	6.28 $\pm$ 9.76	19.32
27.55	27.18	0.20	0.77	12.01 $\pm$ 8.73	14.32	-12.52 $\pm$ 7.94	-42.57
28.17	28.31	0.11	0.11	16.63 $\pm$ 7.45	13.54	2.84 $\pm$ 6.77	14.45
29.00	29.05	0.24	0.27	140.35 $\pm$ 15.08	176.58	30.93 $\pm$ 12.61	48.21
29.54	29.60	0.10	0.08	14.79 $\pm$ 7.74	12.87	13.62 $\pm$ 6.07	10.80

1. de Saussure, G., et al. Nuclear Data for Reactors, Vol. II, 757 (1970). IAEA, Vienna.

TABLE D3

Comparison of Resonance Integrals for ^{233}U

Energy Interval (eV)	Weston et al. ¹	Brooks ²	Nifen-ecker ³	Moore et al. ⁴	Cao et al. ⁵	NVS
0.414-0.532	33.34	35.14		34.20	34.77	35.05
0.532-0.683	30.54	32.29		31.47	32.14	30.75
0.683-0.876	28.84	30.84		29.53	30.61	30.15
0.876-1.126	34.76	37.17		35.54	36.99	36.00
1.126-1.445	56.61	53.87		54.63	59.48	59.00
1.445-1.855	138.23	139.95		140.54	137.32	136.00
1.855-2.382	97.56	95.55	97.63	92.02	99.64	95.15
2.382-3.059	14.55	15.22	13.18	17.19	14.70	16.55
3.059-3.927	26.14	25.54	23.11	25.82	24.73	26.80
3.927-5.043	20.41	20.07	17.92	19.82	18.94	21.85
5.043-6.475	22.10	22.07	19.92	20.83	20.57	25.80
6.475-8.313	35.09	30.98	33.44	31.55	33.56	34.70
8.313-10.67	30.48	28.59	28.43	25.85	28.00	32.10
10.67-13.70	27.61		26.98	23.17	26.55	28.30
13.70-17.59	18.99		19.30	16.36	19.62	20.40
17.59-22.59	32.47		30.87	23.21	30.57	33.80
22.59-29.07	16.76		17.53	15.12	17.50	16.96
29.07-37.03	15.18		15.54	11.77	15.08	15.00
37.03-47.83	6.76		7.27	5.90	8.46	6.86
47.83-61.42	11.29		10.91	8.23	13.17	11.40
61.42-78.86	9.77			7.14	11.44	9.83
78.86-101.3	8.74			6.51	7.83	8.80
101.3-130.0				8.09		8.73
130.0-166.9				4.70		5.60
166.9-214.4	5.82			4.07	5.54	5.61
214.4-275.2	5.51			4.59	5.57	5.87
275.2-353.4	5.42			4.51	4.73	5.05
353.4-453.8	3.65			3.77	3.50	3.50
453.8-582.6	3.54			2.80	3.31	3.30
582.6-748.1	3.84			3.17	3.40	3.80
748.1-960.5	3.26				2.72	3.00
960.5-1223	2.44				2.00	2.15

1. Weston, L.W., et al. ORNL-TN-2140 and Nucl.Sci.Eng. 39, 1 (1968).
2. Brooks, F.D., et al. AERE M-1709 (1966).
3. Nifenecker, H. Journ. de Physique 25, 877 (1964).
4. Moore, M.S., et al. Phys. Rev. 118, 714 (1960).
5. Cao, M.G., et al. Nuclear Data for Reactors, Vol. I, 419 (1970), IAEA, Vienna.

TABLE D4

Comparison of Resonance Integrals for ^{233}U

Energy Interval (eV)	Weston et al. ¹	Brooks ²	Nifenecker ³	Moore et al. ⁴	Cao et al. ⁵	NVS
0.414-10.67	568.65	567.28		559.00	571.45	579.90
1.85-61.42	375.41		362.03		371.9	385.67
0.414-101.2	716.24			676.4	721.17	731.25
166.9-748	27.78			22.91	25.87	27.13
0.414-1223	762.5					777.86

1. Weston, L.W., et al. ORNL-TM-2140 and Nucl.Sci.Eng. 39, 1 (1968).
2. Brooks, F.D., et al. AERE M-1709 (1966).
3. Nifenecker, H. Journ. de Physique 25, 877 (1964).
4. Moore, M.S., et al. Phys. Rev. 118, 714 (1960).
5. Cao, M.G., et al. Nuclear Data for Reactors, Vol. I, 419 (1970), IAEA, Vienna.

Figure E-1

Figure E-2

TABLE F-1

Resonance Parameters of ^{235}U

$E(\text{eV})^*$	$\Gamma(\text{mV})$	Present Results $\sigma_0\Gamma_f(\text{eV b})$	Michaudon ^a $\sigma_0\Gamma_f(\text{eV b})$	Blons ^b $\sigma_0\Gamma_f(\text{eV b})$	Cao ^c $\sigma_0\Gamma_f(\text{eV b})$
1.13	140	14.5±0.4			
2.03	50	1.0±0.2	1.03±0.07		
2.92	150	1.2±0.3			
3.15	190	8.7±0.4	8.1±0.80		
3.61	100	9.6±0.3	8.65±0.90		
4.85	40	1.0±0.2	1.51±0.15		
5.52	300	3.4±0.5			
5.90	300	2.8±0.5			
6.20	200	9.3±0.6			
6.39	50	10.2±0.4	14±1.5		10.1±0.2
7.08	100	10.7±0.5	10±1		9.8±0.1
8.46	40	0.5±0.4			
8.79	150	115.0±1.4	105.5±5		109.9±0.8
9.30	200	18.8±0.8	17±4	20±2	19.9±0.4
9.70	70	1.9±0.4		3	
10.21	95	4.1±0.5	5±1	4.7±0.5	3.9±0.1
11.67	43	7.1±0.5	5.95±0.50	7.5±1	7.0±0.1
12.40	80	50.5±1.1	47±3	50.0±3	54.6±0.7
12.90	83	1.4±0.4	2.9±0.5	2.7±0.5	3.5±0.1
13.31	40	1.0±0.4	3.3±0.6	2.6±1	
13.73	40	2.3±0.6	2.4	3	1.6±0.5
14.03	500	38.1±1.6	27±5	31±6	
14.51	45	2.0±0.5	5±1	2.5±0.5	2.9±0.1

* ±0.02 eV (throughout Table F-1)

a. Michaudon et al., op. cit.

b. Blons, Derrier, Michaudon, Proc. of 3rd Conf. on Neutron Cross Sections & Technology, Knoxville 1971. Conf-710301, Vol. 2, p. 829.

c. Cao, Migneco, Theobald, Wartena, & Winter, Neutron Cross Sections & Technology, NBS-229, Washington DC 1968, p. 481.

TABLE F-1 (CONTINUED)
(page 2)

E(eV)	r(mV)	Present Results $\sigma_0\Gamma_f$ (eV b)	Michaudon $\sigma_0\Gamma_f$ (eV b)	Blons $\sigma_0\Gamma_f$ (eV b)	Cao $\sigma_0\Gamma_f$ (eV b)
15.42	80	8.0±0.6	10.5±1	10.0±1	11.4±0.2
16.09	55	7.9±0.6	10±1	8.7±1	9.8±0.2
16.69	138	12.8±0.8	13.6±1.5	14.2±1	15.2±0.3
17.06	40	0.6±0.4			
18.07	160	20.2±1.0	14.5±2.5	18.5±2	18.6±1.6
18.66	80	2.5±0.6			
18.97	80	8.4±0.9		2±0.5	
19.31	105	117.2±2.0	103±5	121.3±3	118.8±1.0
20.16	40	3.6±0.5	5±1.5	5.2±1	
20.63	90	6.7±0.7	4.35±0.3	6.3±0.5	6.9±0.2
21.07	70	34.0±1.0	29±1.5	34.6±1	34.7±0.4
22.54	40	0.5±0.5			
22.97	90	13.5±0.9	11.7±0.6	13.6±0.5	14.2±0.2
23.44	80	15.3±1.0	8±1	7.0±0.5	6.0±0.2
23.69	60	19.3±1.1	22±4	30±5	
24.32	200	12.3±1.5	8±1.5	7.0±1	
24.47	50	0.9±1.0		5	
24.89	600	18.8±2.0		8	
25.28	100	5.6±1.1		44	
25.59	250	22.0±1.4			
26.50	150	16.1±1.1		22.0±2	
26.83	100	2.6±0.8		4.6	
27.18	50	1.9±0.6	4±1.5	3.4±1	
27.83	110	18.3±1.1	16.5±1	22.4±2	17.8±0.1
28.37	70	3.4±0.7	5±1	5.6±0.5	5.8±0.1
29.67	73	2.0±0.6	3±0.6	3.55±0.5	3.3±0.1
30.63	200	6.2±1.1	4.6±0.9	6.5±1	6.0±0.2
30.87	60	5.8±0.9	6.2±0.7	7.7±0.6	7.1±0.2
32.09	130	40.7±1.6	36±4	45±2	41.8±0.3
33.54	80	28.2±1.4	24.5±3.7	28.2±2	27.1±0.7

TABLE F-1 (CONTINUED)
(page 3)

$E(\text{eV})$	$\Gamma(\text{mV})$	Present Results $\sigma_0\Gamma_f(\text{eV b})$	Michaudon $\sigma_0\Gamma_f(\text{eV b})$	Blons $\sigma_0\Gamma_f(\text{eV b})$	Cao $\sigma_0\Gamma_f(\text{eV b})$
34.40	130	43.4 ± 1.8	33 ± 5	39.0 ± 2	37.3 ± 0.6
34.85	110	26.6 ± 2.0	15 ± 5	24.4 ± 2	25.5 ± 1.2
35.21	170	105.1 ± 2.9	102 ± 20	124.0 ± 4	118.3 ± 2.0
35.77	250	7.3 ± 1.5			
38.28	80	3.9 ± 0.9	12 ± 4	10.7 ± 2	
39.45	95	40.0 ± 1.7	41 ± 4	47.4 ± 1	47.3 ± 0.3
39.88	40	3.2 ± 1.0	6	6.7 ± 0.5	
40.65	52	5.4 ± 1.0	10	11.5 ± 1	
41.30	120	4.6 ± 1.4	8	7	
41.51	60	4.8 ± 1.4	4	5.0	
41.91	90	10.7 ± 1.3	19 ± 2	13.3 ± 1	10.5 ± 0.2
42.28	40	3.2 ± 1.0		7.1	
42.76	64	0.8 ± 1.0	3 ± 0.5	2.5 ± 0.2	2.3 ± 0.1
43.47	66	5.1 ± 1.1	8	7.2 ± 0.5	6.5 ± 0.1
44.10	90	7.5 ± 1.3	10 ± 3	11.0	
44.66	220	18.8 ± 1.9		21.0	
45.04	40	5.3 ± 1.3		13.	

TABLE F-2

 ^{235}U Fission Integrals (normalized to Deruytter)

Energy Interval (eV)	de Saussure ^a	Our Results
0.50-0.70	13.47	13.2
0.70-1.0	17.05	17.1
1.0 - 1.8	29.19	28.2
1.8 - 5.0	54.70	45.2
5.0 - 7.4	62.30	61.6
7.4 - 10.0	219.6	221.2
10.0 - 15.0	216.3	216.5
15.0 - 20.5	316.3	329
20.5 - 33.0	447.8	481
33.0 - 41.0	497.1	514
41.0 - 60.0	918.4	970
60.0 - 100.0	975.5	1044
100.0 - 200.0	2097.4	2325
200.0 - 300.0	2085.3	2150
300.0 - 1000.0	8138.5	8560

a. de Saussure et al., op. cit.