000-1733-2

MASTER

Robert L. Watters and James F. McInroy
Department of Radiology and Radiation Biology
Colorado State University, Fort Collins, Colorado

AT(11-1)- 1733

^{210}Po has been proposed for use in radionuclide electrical generators in space missions. Accidental release of the quantities of this material required for such an application could lead to the contamination of large areas. One of the hazards of such an accident could be the transfer of ^{210}Po from contaminated dairy pasture to man through the milk supply. This study is an attempt to estimate the magnitude of this problem. With the use of assumptions regarding the feeding of cattle and the intake of milk by man an estimate of the surface deposition to produce a maximum permissible body burden is made.

LEGAL NOTICE

The report was prepared as an outcome of developmental agreement work under the United States and the Commission, and may contain nothing on behalf of the Commission.

If there are any omissions or misstatements, approved or disapproved by respect to the accuracy, completeness, or timeliness of the information contained in this report, or that the use of any language, expression, definition, or process disclosed in this report may not fully represent the views, or

B. Address any limitations with respect to the use of, or for damages resulting from the use of any information, processes, methods, or products disclosed in this report.

C. As used in the above, "person acting on behalf of the Commission" includes any employee or representative of the Commission, or employee of such government, the agency, institution or organization, or individual, who has been authorized to participate in the program of development or research, or who has been authorized to receive, store, handle, transmit, disseminate, or otherwise use, any information generated in the employment or relationship with the Government, or the United States, with such person.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

With the advent of radionuclide electrical generators for space mission applications, a new potential source of radioactive fallout is possible. Small radionuclide sources have been used on space vehicles for a number of years (1), but with increasing mission requirements for electrical power these sources will become larger. Megacurie quantities may be used in the future, and ^{210}Po is one of the candidates for these SNAP systems when short term power requirements are indicated (2).

Before such a radionuclide system may be launched, a thorough radiological hazard evaluation must be made (3). This evaluation considers the probabilities of accident and loss of containment of the radionuclide at the various phases of launch, orbit attainment and re-entry. Although aerospace engineers design radioactive source containers to minimize the risk of atmospheric dispersal under any condition, there are still finite probabilities of loss of containment and contamination of the biosphere (4). The probabilities of the various modes of dispersal are determined from reliability data and the radiological doses are estimated from past experience and experimental data. Finally, the radiological hazard estimate is made from the evaluation of the potential radiation dose to the population and its probability of occurrence.

One of the important routes by which an acutely dispersed and deposited radionuclide may find its way into the diet of man is through

the milk from cows which graze on contaminated pasture. The research described in this paper is an attempt to appraise the magnitude of the intake of ^{210}Po through the milk supply after acute deposition and the body burdens which could result in man.

Experimental Methods.

A Holstein cow with a fistula to the rumen was used for the experiment. A dose of 110 μCi of $^{210}\text{PoCl}_4$ in 10 ml of 0.5N HCl was administered to the rumen from a plastic vial which was washed several times in rumen fluid. The administration was made after the morning milking and one liter samples of each subsequent milking were obtained periodically thereafter to determine the long-lived component of the excretion curve. Hand milking was employed throughout the experiment.

The milk samples were placed in three liter beakers with 250 ml of concentrated nitric acid and 250 ml of 70% perchloric acid. The initial reaction, which was quite vigorous, was induced by heating the mixture on a hot plate at the 200° F thermostat setting. After foaming subsided, the samples were allowed to digest at a thermostat setting of 400° F. The appearance of the liquid at this stage was that of a yellow to amber solution with a liquid fat layer floating on the surface. When the volume had reduced to approximately 500 ml, the fat began to oxidize and required additional amounts of nitric acid to prevent eruptions due to rapid oxidation by the perchloric acid. Digestion was

continued until all evidence of organic material disappeared and the solution was devoid of nitric acid. This occurred after the perchloric acid had fumed for some time and the volume had been reduced to about 50 ml of colorless solution. Complete dissolution and oxidation required about 24 hours.

The clear liquid was transferred to a 500 ml polyethylene bottle with hot water and the beaker was washed with hot, concentrated hydrochloric acid followed by two washings with hot water. Approximately 0.1 gm of ascorbic acid was added to the solution, which was then neutralized with 10N sodium hydroxide. The addition of 21 ml of concentrated hydrochloric acid and dilution to 500 ml produced a solution 0.5 N in HCl.

A 200 ml aliquot of the sample was assayed by spontaneous deposition of ^{210}Po onto nickel (5). The plating cell was a pyrex maternity nursing bottle from which the bottom had been cut. A clean nickel disc was placed in the plastic screwcap and the bottle was inverted and the sample introduced. Spontaneous deposition of polonium onto the nickel proceeded for three hours with stirring at a temperature of 85°C .

The plated discs were alpha counted in a Nuclear Measurements Corporation, PC-3B internal flow gas counter with a counting yield of 48% as determined with a standard ^{238}U source. The radiochemical yield for this procedure was $70 \pm 1\%$. The data was analyzed by a

computer program which determined the best regression lines by sequential least squares methods.

Results and Discussion

The milk secretion curve for the acute ingestion of ^{210}Po shown in Figure 1 indicates an initial buildup of ^{210}Po concentration in milk with the peak concentration occurring one day after ingestion. The subsequent reduction in concentration involves an apparent two component decay, which can be described by the equation:

$$C_m = D_o (3.0 \times 10^{-5} e^{-0.19T} + 2.8 \times 10^{-6} e^{-0.021T}) \quad (1)$$

where

D_o = the intake of polonium-210 on the first day in μCi

C_m = concentration of polonium-210 in milk ($\mu\text{Ci/liter}$)

T = time of secretion after initial intake

and the two coefficients, 3.0×10^{-5} and 2.8×10^{-6} are fractions of the initial ingested dose which are contributed to one liter of milk by the respective exponential secretion components after 24 hours.

One method for relating the intake of fallout radionuclides to concentration in milk is by expressing the transfer coefficient (6). This coefficient is the percent of the daily radionuclide intake per liter of milk at equilibrium with constant daily intake. If the above equation for ^{210}Po is integrated assuming constant daily intake, then the transfer coefficient will be given by:

$$\frac{100 C_m}{D_o} = 100 \left[1.6 \times 10^{-4}(1-e^{-0.19T}) + 1.3 \times 10^{-4}(1-e^{-0.021T}) \right] \quad (2)$$

When T is very long ($T \rightarrow \infty$) in relation to the decay constant, the transfer coefficient is estimated to be 0.029 by this method. This value may be compared to an estimated transfer coefficient for fallout ^{137}Cs of 0.40 under the same conditions.

A more realistic expression to describe the concentration of ^{210}Po in milk from cows grazing continuously on a pasture contaminated by an acute radionuclide dispersal can be developed from the single dose secretion curve, (7). From the equation so developed it is possible to derive an estimate of the body burden obtained by people consuming milk from these cows.

The assumptions used for the polonium body burden estimates are essentially those of Davis, et al (8):

1. The deposition time on the pasture is short (hours);
2. The chemical and physical properties are similar to the laboratory form of polonium;
3. The deposition is on pasture supplying all of the feed for the dairy cows;
4. Background and cumulative soil concentrations are negligible;
5. Human consumption of milk is one liter per day and occurs on the day of milking; and
6. The removal time of polonium from the pasture is similar to that of other radionuclides.

A further assumption which is implicit is that the distribution fractions for the body burden of ^{210}Po which are used by the ICRP Committee II are valid (9).

Two factors would affect the ^{210}Po concentration in milk where continued grazing follows acute pasture contamination:

1. The contribution of ^{210}Po to milk from each day of intake, and
2. The decreasing rate of polonium intake caused by pasture losses and radioactive decay.

If equation (1) represents the milk concentration, C_m , on any day, T , after a single intake, then for chronic ingestion the intake on day, t , after initial ingestion combined with all previous intakes will lead to a concentration, M , on any day, T , after the initial ingestion. The ^{210}Po concentration, M , will be the sum of the excretion components for the individual daily intakes and the daily intakes will be decreasing according to the effective half time for the ^{210}Po on the pasture, T_{eff} . This leads to the expression:

$$M = \int_{t=0}^{t=T} D_0 e^{-\frac{0.693t}{T_{\text{eff}}}} \left[3.0 \times 10^{-5} e^{-0.19(T-t)} + 2.8 \times 10^{-6} e^{-0.021(T-t)} \right] dt \quad (3)$$

where $(T-t)$ is the time between consumption of polonium by the cow and secretion into the milk.

Thompson (10) has estimated that the most probable effective half life for any radionuclide on vegetation can be calculated by the equation:

$$T_{\text{eff}} = \frac{13 T_r}{13 + T_r} \quad (4)$$

where T_r = radioactive half life.

For ^{210}Po with a half life of 138 days, the equation estimates the effective half life on vegetation to be twelve days. With this pasture half removal time

$$M = D_0 \quad 2.3 \times 10^{-4} (e^{-0.058T} - e^{-0.19T}) - 7.6 \times 10^{-5} (e^{-0.058T} - e^{-0.021T}) \quad (5)$$

A normalized curve for ^{210}Po milk concentrations at any time after the start of intake by the cow is shown in Figure 2. The concentration is expressed as the fraction of ^{210}Po intake on the first day per liter of milk. An estimate of the milk concentration on day, T , can be made with this curve for various acute contamination levels. If continuous grazing is assumed, the average utilized area factor (UAF) per cow is reported to be 42 m^2/day (11). The surface concentration, ($\mu\text{Ci}/\text{m}^2$), after an acute deposition, multiplied by the (UAF) will give the ^{210}Po intake, D_0 , on the first day.

The minimum polonium surface deposition which would produce the maximum permissible body burden (MPBB) from the daily intake of one liter of contaminated milk may be calculated from the ICRP exponential model (9):

$$\frac{d(qf_2)}{df} + \lambda(qf_2) = P$$

where:

q = maximum permissible body burden (MPBB) of
polonium-210 = 0.00 μCi for the general population.

f_2 = fraction of body burden in critical organ (spleen) = 0.07.

P = product of the radionuclide intake I and the fraction, f_w ,
reaching the spleen.

f_w = fraction reaching critical organ by ingestion = 0.002.

λ = $\frac{.693}{T_{\text{eff}}} = 0.017 \text{ day}^{-1}$

T_{eff} = effective half life of ^{210}Po in the spleen = 42 days.

For the case of acute deposition followed by continuous pasture removal, the intake term, I , would be replaced by M the concentration of polonium in milk on any day, T , after the initial ingestion by the cow, and the differential equation is:

$$\frac{d(qf_2)}{dt} + \lambda(qf_2) = f_w D_0 \left[2.3 \times 10^{-4} (e^{-0.058T} - e^{-0.19T}) - 7.6 \times 10^{-5} (e^{-0.058T} - e^{-0.021T}) \right] \quad (6)$$

A solution of this equation

$$qf_2 = f_w D_0 \left[0.19e^{-0.017T} - 0.0037e^{-0.058T} - 0.0013e^{-0.19T} - 0.017e^{-0.021T} \right] \quad (7)$$

may be solved for the value of D_0 which produces the MPBB. This initial intake is calculated to be 17 μCi and would correspond to a pasture surface concentration of 0.4 $\mu\text{Ci}/\text{m}^2$ if the previously stated assumptions about grazing are valid.

The estimate presented for the minimum deposition to produce the MPBB for ^{210}Po is no doubt conservative. The assumptions used are chosen to give the maximum effect. Any change in the feeding practices such as dry feed supplements or storage of hay for a period of time prior to feeding would increase the maximum deposition required. A less soluble form of the polonium would reduce the transfer from the cows digestive tract to the blood stream. Reduction in the quantity of milk consumed or appreciable delay in consumption would lead to a higher initial concentration to achieve the MPBB. Finally, the MPBB would be of short duration (approximately 20 days) so that higher body burdens may be justified in this type of analysis.

REFERENCES

1. D. G. Harvey, P. J. Dick and C. R. Fink, *Nucleonics* 21, No. 4, p. 56 (1963).
2. H. L. Davis, *Nucleonics* 21, No. 3, p. 61 (1963).
3. P. G. Lafyates, *Nuclear Safety* 1, No. 4, p. 1 (1960).
4. W. T. Harper, *Transactions American Nuclear Society* 9, No. 1, p. 161 (1966).
5. R. L. Blanchard, *Analytical Chemistry* 38, p. 189 (1966).
6. G. M. Ward, J. E. Johnson and L. B. Sasser, *Journal of Dairy Science* 50, No. 7, p. 1092 (1967).
7. F. W. Lengemann, R. A. Wentworth and F. L. Hiltz, *Health Physics* 14, p. 101 (1968).
8. J. J. Davis, F. W. Lengemann and G. F. Fries, *Radiation Health Data* 6, No. 7, p. 387 (1965).
9. Recommendation of the International Commission on Radiological Protection. Report of Committee II on Permissible Dose for Internal Radiation, ICRP, Publication 2, Pergamon Press, London (1959).
10. S. E. Thompson, Effective Half-Lives of Fallout Radionuclides on Plants with Special Emphasis on Iodine-131, UCRL 12388, University of California, Ernest O. Lawrence, Livermore, California (1965).

11. J. J. Koranda, Agricultural Factors Affecting the Daily Intake of Fresh Fallout by Dairy Cows, UCRL 12479, University of California, Ernest O. Lawrence Radiation Laboratory, Livermore, California (1965).

List of Figures

- Figure 1 A graph of the ^{210}Po milk secretion curve for an acute ingestion dose of 110 μCi of PoCl_4 to a Holstein cow.
- Figure 2 A graph of the predicted fraction of the first day ^{210}Po ingestion dose when continuous grazing on contaminated pasture occurs. The estimated pasture removal half time is 12 days.

FRACTION OF INGESTED DOSE OF ^{210}Po ON FIRST DAY FOUND
IN ONE LITER OF MILK