JUN 9 1965

MASTER

CAR 882-8

Carnegie Institute of Technology

Department of Physics

$\pi^- + p$ INTERACTIONS AT 646 MeV.

by

John D. Oliver

January, 1965

PATENT CLEARANCE OBTAINED. RELEASE TO
THE PUBLIC IS APPROVED. PROCEDURES
ARE ON FILE IN THE RECEIVING SECTION.

Contract AT(30-1)-882

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

UNCLASSIFIED

Car 882-8.

$\pi^- + p$ INTERACTIONS AT 646 MeV.

by

John D. Oliver

January, 1965

Carnegie Institute of Technology

Pittsburgh, Pennsylvania

Contract AT(30-1)-882.

A thesis, based upon the work reported here, has been submitted to the Department of Physics, Carnegie Institute of Technology, in partial fulfillment of the requirements for the degree of Doctor of Philosophy.

TABLE OF CONTENTS

	page
LIST OF TABLES.....	iv
I. ABSTRACT.....	v
II. INTRODUCTION.....	1
III. DESIGN OF THE EXPERIMENT.....	5
A. Beam Production and Design.....	5
B. Beam Composition.....	6
C. Bubble Chamber.....	7
IV. DATA ANALYSIS.....	9
A. Event Types.....	9
B. Scanning Procedure.....	9
C. Stop Scan Procedure.....	11
D. Track Count.....	13
E. Measuring Procedure.....	14
V. COMPUTING.....	16
A. Geometrical Reconstruction.....	16
a. Reconstructed variables.....	16
b. Tests of accuracy.....	17
c. Remeasurements.....	18
B. Kinematic Fitting.....	19
a. Theory.....	19
b. Error analysis.....	22
c. Determination of momentum.....	23
d. Fitting procedure.....	27
e. Test functions.....	27

V.	Computing (continued)	
	C. Editor.....	29
	D. Plotter.....	30
	E. Prirej.....	32
VI.	RAW RESULTS AND CORRECTIONS.....	34
	A. Events.....	34
	a. Introduction.....	34
	b. Events examined by Editor.....	34
	c. Corrections to events examined by Editor.....	36
	d. Corrections for events not examined by Editor...	44
	e. Correction to the differential cross section....	48
	B. Neutrals.....	51
	a. Measurement.....	51
	b. Corrections to neutrals.....	51
	C. Scanning Efficiency.....	54
VII.	CROSS SECTIONS.....	56
	A. Definition.....	56
	B. Total Track Length.....	57
	C. Density of Hydrogen.....	60
	D. Cross Sections.....	60
VIII.	THEORY AND DISCUSSION.....	64
	A. Partial Wave Analysis of Elastic Scattering.....	64
	a. Theory.....	64
	b. Small angle correction.....	68
	c. Fit to differential cross section.....	68

VIII. Theory and discussion (continued)

B. Inelastic Scattering.....	74
a. Theory.....	74
1. The Statistical Model.....	75
2. The Isobar Model.....	75
b. Single pion exchange.....	78
c. Results for effective two-body mass.....	78
d. Angular distributions.....	80
e. Multiple pion production.....	81
IX. CONCLUSION.....	85
X. ACKNOWLEDGEMENTS.....	87
APPENDIX I	
Derivation of Error Formulas.....	88
APPENDIX II	
The Olsson-Yodh Isobar Model.....	92
BIBLIOGRAPHY.....	96
FIGURE CAPTIONS.....	100

LIST OF TABLES

	page
I. Quantum Numbers for Peaks in $\sigma_T(\pi^- p)$	4
II. Uncorrected Results for Measured and Computed Events	36
IIIa. Reasons for Rejection by Plotter or Prirej	37
IIIb. Fractional Rejections for az, z or dip for events passing x and y	38
IV. Corrections for Large dip	44
V. Events not computed by Guts	45
VI. Events which failed Guts	45
VII. Corrected Number of Good Events	47
VIII. θ_{23} Corrections to Differential Cross Section	49
IX. Angular Distribution of Elastic Events	50
Xa. Results of the Experiment	61
Xb. Differential Cross Section	62
XI. Coefficients for $a_n = \sum_{ij} b_{ijn} A_i A_j^*$	67
XII. Coefficients a_n for 5th order curve	71
XIII. Coefficients a_n for all fits	72
XIV. Cross sections for $\pi^- + p \rightarrow \pi^- + \pi^+ + n + x^0$, where x^0 is π^0 or an energetic γ ray	83

ABSTRACT

A study of π^-p reactions at 646 MeV was made using a 14 inch liquid hydrogen bubble chamber. Approximately 6600 good events were analyzed and the following cross sections were obtained:

$$\begin{aligned}\sigma_{(\text{elastic})} &= 17.56 \pm 0.43 \text{ mb}, & \sigma(\pi^0\pi^-p) &= 4.65 \pm 0.17 \text{ mb}, \\ \sigma(\pi^-\pi^+n) &= 7.14 \pm 0.23 \text{ mb}, & \sigma_{\text{neut.}} &= 11.78 \pm 0.43 \text{ mb}, \\ \sigma(\pi^-\pi^+\pi^0n) &= 0.33 \pm 0.04 \text{ mb} & \text{and } \sigma(\pi^-p\pi^-\pi^+) &= 0.08 \pm 0.02 \text{ mb}.\end{aligned}$$

This gives a total cross section of 41.54 ± 0.82 mb. The elastic differential cross section was fitted to a 5th order polynomial in $\cos\theta^*$ with coefficients $a_0 = 0.27 \pm 0.02$, $a_1 = 1.48 \pm 0.11$, $a_2 = 3.86 \pm 0.22$, $a_3 = -0.29 \pm 0.53$, $a_4 = -0.65 \pm 0.28$, and $a_5 = 1.69 \pm 0.52$ (mb/ster). The fact that a 5th order curve was needed indicates that partial waves up to $F_{5/2}$ are present. The inelastic reactions are dominated by production of the $N^*(1238)$ isobar and were fitted to the model of Olsson and Yodh. The model failed to explain the angular distributions of the reaction products and also failed to explain the $M_{\pi^-\pi^+}$ plot. However, it was successful with the other two-body effective mass plots.

II. INTRODUCTION

The study of pion physics has been one of the most active and fruitful branches of modern physics. Intensive studies of the interaction of positive and negative pions with protons, as a function of pion energy, have revealed marked structures in the total cross sections and in the differential cross sections for elastic scattering.

Figure 1 shows the total and neutral cross sections for scattering of positive and negative pions by protons in the energy range 0-1200 MeV.¹⁻⁶ The $\pi^- p$ total cross section has three prominent peaks. The first of these is the now well established 3-3 resonance and will be discussed below. The second peak, which occurs at an incident π^- kinetic energy of about 600 MeV, is smaller and less is known about it. This second peak forms the subject of this work.

When this experiment was planned, data were available on the total and neutral cross sections and there was some knowledge of the elastic cross section in the region of the 600 MeV peak. However, many of the quoted elastic cross sections had large errors and very little was known about the inelastic cross sections. It was intended to make detailed studies of the $\pi^- + p$ reactions at 550, 600, and 646 MeV. These energies were chosen so as to bracket the second peak in order to make comparisons among the results at the maximum and at both sides of the peak. The experiment was performed in cooperation with groups from the University of Maryland⁷ and the University of Kentucky⁸ who analyzed the results at 550 and 600 MeV respectively.

One of the most useful concepts for interpreting our type of data is that of isotopic spin. If $|I, I_z\rangle$ represents an isospin wave

function, one can express the π^+p and π^-p isospin function as⁹

$$|\pi^+p\rangle = \left| \frac{3}{2}, \frac{3}{2} \right\rangle \quad (1a)$$

$$|\pi^-p\rangle = \sqrt{\frac{1}{3}} \left| \frac{3}{2}, -\frac{1}{2} \right\rangle - \sqrt{\frac{2}{3}} \left| \frac{1}{2}, -\frac{1}{2} \right\rangle \quad (1b)$$

These expressions mean that the π^+p reaction is a pure $I = \frac{3}{2}$ whereas the π^-p reaction is a combination of $I = \frac{3}{2}$ and $I = \frac{1}{2}$. We can give the expressions (1a) and (1b) a physical meaning if we recall that isospin was originally introduced into nuclear physics to explain the fact that n-n, p-p, and n-p forces are equal if electromagnetic interactions are neglected. Formally, this is expressed by the concept of charge independence which states that nuclear forces depend only on I , the total isospin, but not on I_z , the z component. If we extend this concept to meson-nucleon forces, we see that (1a) and (1b) imply that any structure in the π^+p total cross section should be reflected in the π^-p cross section because the latter contains an $I = \frac{3}{2}$ component. On the other hand, features in the π^-p total cross section may come solely from the $I = \frac{1}{2}$ state and need not appear in the π^+p cross section. Examining Figure 1 with this in mind, we see that $\sigma_T(\pi^-p) = \frac{1}{3} \sigma_T(\pi^+p)$ at the 200 MeV peak and that there are no peaks in $\sigma_T(\pi^+p)$ at 600 or 900 MeV. Consequently, we can assign $I = \frac{3}{2}$ to the 200 MeV peak and $I = \frac{1}{2}$ to the peaks at 600 and 900 MeV in the π^-p total cross section.

Customary usage reserves the term "resonance" for the case of a phase shift passing through 90° (cf. Ch. VIII). However, it is also possible that two partial waves could add in such a way as to produce a peak in a cross section. Indeed, several authors^{10,11} have

proposed inelastic mechanisms which could account for the peak at 600 MeV. Thus, a peak in total cross section need not indicate a resonance. It is well established that the peak at 200 MeV is due to a phase shift passing through 90° ; however, there is no clear evidence to establish either the 600 or 900 MeV peaks as resonances. Consequently, we shall refer to them as "peaks" in this thesis.

Although it has been relatively simple to assign isospin to the peaks, it has proved to be much more difficult to assign angular momentum and parity. In theory, one can make a phase shift analysis using the data from elastic scattering, polarization measurements, and photoproduction experiments.^{9,12-14} This has been done successfully at 200 MeV with the result that this resonance is known to be $J = \frac{3}{2}$, even parity. However, in this case the analysis is greatly simplified by a number of facts:^{1,9}

- 1) The relative sizes of the π^+p and π^-p peaks indicate that only the $I = \frac{3}{2}$ state need be considered. This follows from (1a) and (1b) because if no $I = \frac{1}{2}$ state is present, the π^-p cross section should be $\frac{1}{3}$ of $\sigma(\pi^+p)$, and this is indeed the case.
- 2) The scattering is purely elastic so that all the phase shifts are real.
- 3) The differential cross section for elastic scattering shows that only $\ell = 0$ and $\ell = 1$ waves are present in the partial wave expansion. That is, we need consider only s and p waves in the analysis.

The situation at the higher peaks is much more complex. One cannot ignore the $I = \frac{3}{2}$ state because $\sigma_T(\pi^+p)$ is about $\frac{1}{3}$ of $\sigma_T(\pi^-p)$. Furthermore, complex phase shifts are needed because the scattering is very inelastic. Finally, it is known that partial waves up to $F(\ell = 3)$ are present.¹⁵⁻²¹ A number of authors^{12,22-24} have reviewed the evidence and have concluded that the most probable assignments for the quantum numbers associated with these states are those shown in Table I, provided that the two higher peaks are true resonances.

TABLE I
Quantum Numbers for Peaks in $\sigma_T(\pi^-p)$

ENERGY	J	I	Parity
200	$\frac{3}{2}$	$\frac{3}{2}$	+
600	$\frac{3}{2}$	$\frac{1}{2}$	-
900	$\frac{5}{2}$	$\frac{1}{2}$	+

Chapter III of this thesis discusses the design of the experiment; Chapter IV is a discussion of the scanning and measuring procedures; Chapter V describes the computer programs used to analyze the results; Chapter VI and VII give the results and necessary corrections, and Chapter VIII is a discussion of the results.

III. DESIGN OF THE EXPERIMENT

A. Beam Production and Design

The experiment was performed using the Cosmotron at Brookhaven National Laboratory with the Adair 14 inch liquid hydrogen bubble chamber. Figure 2 is a plan drawing of the beam layout. The first pair of quadrupoles furnished horizontal and vertical focusing, the bending magnet provided momentum resolution and some additional vertical focusing, and the last quadrupole furnished additional horizontal focusing. The C magnet was used to compensate for the fringing field of the bubble chamber magnet; i.e., it deflected the beam so that it entered the chamber nearly parallel to a diameter. The beam optical system was designed with the aid of the Beam II program of Brookhaven National Laboratory²⁵ and was planned to give an average momentum of 775 MeV/c with a spread of $1\frac{1}{2}\%$ at the beam window of the bubble chamber. It was calculated that the quadrupoles increased the intensity of the beam by a factor of 13.5 at the chamber. This was necessary because the experiment was designed as a parasite experiment in which only part of the internal proton beam of the Cosmotron was used to produce the meson beam. Details of the electronics system used to control the internal proton beam are given by Vittitoe⁸. Final adjustments of the magnet currents were made with the wire orbit technique. The tension and current of the wire were set to represent π^- mesons with a momentum of 775 MeV/c and the current in the bending magnet was adjusted to bring the wire to a good focus. After the correct bending magnet current was determined, the current in the wire was varied. It was found that a $\frac{1}{2}\%$ change in the current in the wire caused the wire to be noticeably off-center. This indicated that the system had good momentum resolution.

B. Beam Composition

A measurement of the muon and electron contamination of the beam was made in order to determine how many pions were actually entering the bubble chamber. It was not possible to do this with differential range measurements because, at this energy, the range of the pion is large compared to its mean collision length. Instead, a gas (CO_2) filled Cerenkov counter in coincidence with two scintillation counters was used. The arrangement is shown in Fig. 3a. This method is based on the fact that the threshold for Cerenkov radiation is given by $\beta = \frac{1}{n}$ where n is the index of refraction and β is the velocity of the particle in units of the speed of light. At 775 MeV/c, the velocities of e , μ and π are respectively $\beta_e = 0.9999+$, $\beta_\mu = 0.991$ and $\beta_\pi = 0.985$. The index of refraction of CO_2 increases with pressure. Hence, as the pressure was increased, we detected successively in the Cerenkov counter, electrons, electrons and muons, and finally electrons, muons and pions. Counters 1 and 2 (Fig. 3a) detected all charged particles passing through the counter telescope. However, the Cerenkov counter detected only those particles with velocities above the threshold. Thus, the ratio $\frac{12^Y_C}{12}$, at any pressure, gives the fraction of the beam with velocities greater than the threshold velocity at that pressure. Figure 3b shows the curve of $\frac{12^Y_C}{12}$ vs. pressure obtained for a beam momentum of 775 MeV/c. The first plateau corresponds to muons and electrons and the second plateau corresponds to the addition of the pions. A plateau for electrons alone was not found because the measurement did not go to a low enough pressure. Using $\beta_\pi = .985$ and noting that at 0°C the index of refraction of CO_2 is given by²⁶

$$n = 1 + .0045 P$$

where P is in atmospheres, we find that the threshold pressure should be approximately 33.8 atmospheres or 34.9 kg/cm^2 which is in reasonable agreement with the 37 kg/cm^2 shown in Fig. 3b.

The results for the ratio $\epsilon_\mu \equiv \frac{\mu + e}{\pi + \mu + e}$ as a function of momentum are shown in Fig. 3c. It was found that $\epsilon_\mu = 0.145 \pm 0.005$ at 775 MeV/c.

C. Bubble Chamber

The bubble chamber (Fig. 4) is a copper cylinder 8 inches deep and 14 inches in inside diameter. It was mounted in a magnet in such a way that the plane of the windows was perpendicular to the floor. The beam direction, the magnetic field direction and the vertical direction form a trident of lines which are roughly mutually perpendicular. Three cameras were mounted 36.5 inches from the front window with their optical axes perpendicular to the plane of the windows. The optical system has a demagnification factor of about 12. The front window (Fig. 4b) has two fiducial marks, 10 inches apart, etched on the inside surface. These are used to define the coordinate system. The back window has a single fiducial mark, etched at its center, which helps to distinguish the three views but is not used to define the coordinate system.

The coordinate system (4c) is a left handed system with z increasing from front to back, y up, and x toward the incoming beam, i.e., the beam enters the chamber at the high x end and leaves it at the low x end. Two angles, azimuth and dip, (Fig. 4c) are used to define track directions. Let p be any momentum. Then the components of p are expressed in terms of the above angles as follows:

$$p_x = p \cos(\text{dip})\cos(\text{az})$$

$$p_y = p \cos(\text{dip})\sin(\text{az})$$

$$p_z = p \sin(\text{dip})$$

Dip ranges from -90° to $+90^{\circ}$ and ϕ has a range of 0° to 360° relative to the positive x axis. Thus a positive dip means the track is going in the positive z direction.

The bubble chamber magnet was designed to produce a central field of approximately 17.5 kilogauss with a current of 5000 amperes. The field was measured by the staff of Brookhaven National Laboratory and was found to obey the equation:

$$B(\text{Kilogauss}) = 17.486 + 0.0487 (z-4)^2 + 0.0338 r^2 + 0.00045 (z-4)^2 r^2 \quad (2)$$

In this formula, r is the radial distance from the central axis of the chamber (coordinates $x = 7''$, $y = 7''$) and all distances are measured in inches. The field was strong enough to produce a radius of curvature of about 60 inches for a particle with a momentum of 775 MeV/c.

IV. DATA ANALYSIS

A. Event types

At 650 MeV, a rather large number of reactions can take place. These are listed below with the notation (np or n prong) to indicate the number of visible outgoing tracks. They are labeled with the terms used to refer to them in this thesis. A typical 2p event is shown in Fig. 5.

$\pi^- + p \rightarrow \pi^- + p$	(2p)	elastic scattering	(3a)
$\rightarrow \pi^- + p + \pi^0$	(2p)	π^0 production	(3b)
$\rightarrow \pi^- + n + \pi^+$	(2p)	π^+ production	(3c)
$\rightarrow \pi^- + \pi^+ + \pi^- + p$	(4p)	multiple pion production	(3d)
$\rightarrow \pi^- + \pi^+ + n + \pi^0$	(2p)		(3e)
$\rightarrow \pi^- + \pi^0 + \pi^0 + p$	(2p)		(3f)
$\rightarrow \pi^0 + n$	(0p)	charge exchange	(3g)
$\rightarrow \pi^0 + \pi^0 + n$	(0p)	or stops	(3h)
$\rightarrow \pi^0 + \pi^0 + \pi^0 + n$	(0p)		(3i)

The dominant reactions are 3a, 3b and 3c which have a combined cross section of about 30 millibarns or about 3/4 of the total cross section at this energy. The stops have a cross section of about 11 mb and measurements at Saclay⁴ indicate that charge exchange is the dominant neutral reaction. Finally, the multiple pion production events are rare, having a cross section of less than 0.6 mb.

B. Scanning procedure

Forty-two rolls of film with about 26,000 frames were scanned for events. As shown on Fig. 4, each camera has a number assigned to it which is used to identify the view. That is, the picture taken by

camera 1 is called View 1 etc. In order to eliminate chance coincidences of tracks, at least two views must be used in scanning. The scanners were instructed to use View 1 and View 3 because the frame numbers on View 2 were almost illegible. The film was projected onto the screen of the scanning machine at approximately lifesize. All frames with less than 40 tracks were scanned for 2 prong, 4 prong or 1 prong events. 1 prong events were recorded only if the scattering angle was greater than 10° . In order to be considered suitable for measurement, the events had to satisfy certain criteria. These were defined by constructing a special template as shown in Fig. 6a. The template was divided into ten equal zones that were used to identify the rough location of an event for the measurer. When using the template, it was placed on View 1 with line AA directly over the right hand fiducial mark and the central horizontal line connecting the right and left hand fiducial marks. That is, the positive zones lay above the line connecting the left and right fiducials and the negative zones lay below the line. View 1 was chosen for use with the template because camera 1 was almost directly over the beam entrance window. Therefore, relatively little distortion of the entrance angle of the incident beam track is produced. This is illustrated in Fig. 6b which shows a section of the chamber in the plane defined by the two front fiducial marks and the camera. The heavy lines enclose the region which, as seen by camera 1, will project into the template. The dotted lines enclose the region of the chamber actually used (the final fiducial volume) by the computer programs. It can be seen that at the downstream end of the chamber there is a large region in which events fall inside the scanning template but outside the final fiducial volume. These events have short outgoing tracks in the downstream direction and,

indeed, some tracks which in reality enter the chamber wall will give the appearance of stopping within the chamber. Consequently, events in this region were rejected by the computer programs because they could not be accurately measured.

Referring again to Fig. 6a, the scanners were told to define an acceptable event as one meeting all of the following criteria:

1. The vertex was inside the template.
2. The incident track crossed line AA of the template.
3. The entrance angle of the incident track was within $\pm 5^\circ$ of the average beam direction.

The entrance angle was defined as the angle between the tangent to the incident track and the line between the left and right fiducials. A positive entrance angle meant that the track was going in the negative y direction. It was always measured at the point where the track crossed line AA of the template. The average beam direction was not rigorously defined and hence varied somewhat from scanner to scanner, roll to roll, and even frame to frame. However, it will be seen later that the final criteria were much stricter so that the variation in acceptance angle did not effect the final results.

All events which met the above criteria were recorded on special scanning sheets by roll, frame, and zone number. The scanner also indicated the type, i.e. 2 prong, 4 prong or 1 prong, and noted any special information such as second scatterings or $\pi - \mu - e$ decay.

C. Stop scan procedure

A separate scan of 39 rolls was made for stops. This differed from the regular event scan in two ways:

1. Only frames with less than 25 tracks were used.
2. Only zones 1 through 4 of the template were used.

These changes were necessary in order to provide good scanning efficiency. In the first place, it was found that the scanners would miss stopping tracks when there were more than 25 tracks on a frame. In the second place, as mentioned previously, a large fraction of the incident tracks only give the appearance of stopping in zone 5. A study of Fig. 6b will show that this is a result of the projection of tracks near the back window into zone 5 of the template even though the track actually left the chamber.

Another difficulty came from tracks which passed into the front or back window, and would therefore appear to stop. A special procedure was used to distinguish between true stops and tracks which went out a window. It was based on the fact that the end point of a track which goes out a window will be the same distance from the fiducial marks on that window no matter which view is used. The scanner measured the directed distance from the end point to a line drawn between the left and right fiducials. Calling this distance y_1 on View 1 and y_3 on View 3, we define $\zeta = y_1 - y_3$. This will be 0 if the track is out the front window and some constant number $\zeta_{\max}$ if it is out the back window. A track which really ends in the chamber will have a ζ someplace between these two limits. The scanner recorded the ζ for all stops. Figure 7 shows the ζ distribution for stops on several rolls. Nearly 55% of the stops lie in the peaks $\zeta \leq 0.1$ or $\zeta > 2.3$. These are considered to be out the front and back window respectively. There is no direct connection between ζ and z except for tracks which go out the windows.

D. Track Count

A measurement of absolute cross sections requires a knowledge of total number of tracks crossing the fiducial volume. In order to determine this, the scanners were told to count all the tracks on every tenth frame satisfying the following rules:

1. The track crossed AA on the template.

2. The entrance angle lay between 1.8° and 6.6° . The entrance angle limits were based on the azimuth distribution of the incident tracks of events. This will be discussed later (Chapter V) in the section on beam acceptance criteria. The angles were measured with a template which had the angular limits drawn on it. Recounts of a number of rolls showed that the track count was reproducible to one percent. One cannot reasonably expect better reproducibility because some tracks were obscure and others were close to the angular limits. This track count was combined with a calculation (Chapter VII) of average incident track length to give the total track length of incident pions.

A special track count of one-half of the rolls was done to check the curvature of the tracks by comparing the tracks with a circular template having a 55 inch radius. This curvature corresponds to the average beam momentum of 775 MeV/c (cf. Chapter V-C). It was found that the average 10 inch long track would fit this template and had a sagitta of 6 mm., but that 6% of the tracks had sagittas of 7 mm. or more. This larger sagitta could be due to a very large dip, a track close to the back window, or a momentum lower than the average. In the latter case, the momentum would have to be at least 16% lower than the average momentum.

It was not feasible to check the incident track of events by template in the manner described above because the average length of

these tracks is only 5 inches. This reduces the average sagitta from 6 mm to 1.5 mm and a deviation of 1 mm for a 10 inch track becomes only 0.25 mm for the 5 inch track. This difference is too small to be measured reliably by template. Consequently, any events corresponding to the large sagitta beam tracks are included in the total number of events analyzed. Therefore, the tracks with large sagittas were included in the final track count.

E. Measuring procedure

The events found were measured on either a Hermes or a Hydel measuring machine and the coordinates plus identifying remarks automatically punched into IBM cards. These measuring machines are standard commercial machines which are capable of measuring a point to an accuracy of five microns on the film. This corresponds to about 0.060 mm in the bubble chamber. The reliability of the machines was tested several times a week with a standard test pattern and a computer program. Whenever the test showed that a machine was not working properly, the difficulty was fixed and the appropriate events were remeasured.

The measurers assigned a twelve digit identification number (ID) to each event. It consisted of the roll and frame number, an event number, and a type number. The event number was 1 if there was only one event on the frame. Otherwise it distinguished the event being measured from others on that same frame. The type number identified the event as being 2-prong, 4-prong, $\pi - \mu - e$ decay or a second scattering. In addition to the ID, the measurer also assigned a measurement number indicating the number of times the event had been measured and indicated whether any positive track appeared to stop in the chamber. The ID number and the remarks were punched on all cards for an event. On each view, the

measurer measured the left and right fiducial marks and the three (one incident and two outgoing) tracks belonging to the event. Five roughly equally spaced points were measured on each track starting near, but not at, the vertex. In case a track appeared to stop in the chamber, one of the five points was taken to be the end point. The views and tracks were identified by numbers punched on each card.

Approximately 8850 events were measured at least once. 350 events were not measured at all. The most common reasons for not measuring an event were that the tracks were faint or obscure. About 50 events were not measured because they involved 2 or more secondary scatterings and it would have required major changes in the program before they could have been analyzed. Finally, 50 events were not measured because the outgoing tracks were so short that the scanner could not tell the positive track from the negative track. These were really outside the fiducial volume and should be subtracted from the 350 unmeasured events. The treatment of the remaining 300 unmeasured events will be described later. A complete flow chart for the data processing is shown in Fig. 16.

V. COMPUTING

A. Geometric Reconstruction

a. Reconstructed variables. - The geometric reconstruction of events was done with a computer program called Datpro²⁷. This program took the results of the measuring machine and converted them into appropriate three dimensional quantities describing the event. These quantities are the position of the vertex, the azimuth, dip, and radius of curvature of each track and the approximate lengths of the tracks. If the remarks indicated that a track might have stopped in the chamber, the exact length from the vertex to the end was computed. The program also computed the momenta of the tracks and errors in all these quantities.

The basic assumption in Datpro is that all tracks project onto the film as circles. This would be rigorously true only if the magnetic field were constant and the dip angles were all zero. The measurements made with a circular template (radius 55") indicate that this is a very good assumption for the incident tracks. However, one cannot expect it to be good for tracks with large dips and, indeed, there is considerable evidence that it is unreliable for dips greater than 65° . This limit is not as restrictive as it appears because phase space factors limit the number of events with large dips.

Datpro treats each track and view independently. For a given track on a view, it uses three points to determine a radius and a center of curvature. It then selects the pair of tracks which are most nearly perpendicular and determines their intersection from the known centers of curvature and radii. This defines the projected coordinates of the vertex for the view being used. It is relatively simple to find the three dimensional vertex coordinates once the projected vertex coordinates are known

for each view. This can be done with any two views and Datpro uses all three possible pairs of views and averages the results. Once all of the above quantities are known, Datpro constructs a tangent to each track and calculates the azimuth and dip angles. As the final step in the reconstruction, Datpro corrects each measured radius for the effects of position and dip, and then averages the three measured radii for each track to find the true radius in the chamber.

The momentum is related to the radius and dip by the formula $p \cos(\text{dip}) = kHR$ where H is the magnetic field and k is a numerical factor depending only on the system of units. The term $\cos(\text{dip})$ is needed because only the component of $\vec{p}$ which is perpendicular to $\vec{H}$ is affected by the field.

The only remaining variable of interest is the length of each track. If the track does not stop, this is defined as the distance between the first and last measured points on view 1 divided by $\cos(\text{dip})$. This definition does not give an accurate measure of the length but is useful in identifying the track if trouble occurs later in the analysis. A more rigorous method was used if the track stopped in the chamber. In that case, Datpro reconstructed the three dimensional coordinates of the endpoint and checked that it was actually in the chamber and not at one of the windows. If the track passed this test, the distance between the vertex and the endpoint was calculated using a formula which allowed for dip and curvature. Those tracks which were found to enter one of the windows were treated as non-stopping tracks.

b. Tests of Accuracy. - A number of tests were incorporated into Datpro to check the accuracy of measurement. The most important of these tests were those for radius and vertex fits. The radius test compares the sagittas of a track as computed for all three views. The

sagitta is defined as the distance from the midpoint of a track to the straight line joining the endpoints. It is related to the radius by the formula $S = \frac{L^2}{8R}$ where L is the straight line distance between the endpoints. Let S_i be the sagitta of a track computed for the i^{th} view and let $\bar{S}$ be the average of the three calculated sagittas for the track. The test requires that $|S_i - S_j| < 0.25 \times \bar{S}$ where i and j are any two of the views. The program tests all three possible distinct pairs of views and rejects the event if any pair fails the test.

It was mentioned earlier that the three dimensional vertex coordinates can be determined from any pair of views. This fact is used to define a test for vertex fit. Let (x_i, y_i) be the x and y coordinates of the vertex as determined by some pair of views. (The index i indicates which pair of views was used.) The test for vertex fit defines $S_{ij} = \sqrt{(x_i - x_j)^2 + (y_i - y_j)^2}$ and requires that S_{ij} be less than 0.045 inch for all three possible distinct combinations. This test was needed because inaccuracy in the vertex would cause large errors in the azimuth and dip. The rather arbitrary limits in these two tests are the results of experience with the program at Brookhaven.

A binary record was written on magnetic tape for use with the next program if the event passed all of the tests. The binary record contained the ID of the event and the dip, azimuth, momentum, length and radius of each track. The same information was written (in BCD) on another tape. This data was printed and examined by the physicist in order to check the program and examine individual events.

c. Remeasurements. - Those events which failed Datpro were listed for remeasurement. About 22% of all events failed the first measurement and 25% of these failed a second time. A sample of the events which failed twice was measured a third time and 67% of them failed again.

A detailed examination showed that two-thirds of the events which failed three times had been rejected by the radius test and that the offending tracks had $|\text{dip}|$ greater than 65° . The most reasonable explanation is that the circle approximation failed for tracks with large dips. Most of the remaining failures were poor vertex fits and all of these events had small opening angles between all three tracks. In such cases, the three tracks of the event overlapped for part of their lengths and the vertex was not well defined. Some of these were successfully measured by measuring the vertex directly in 2 of the views and using only those views in the reconstruction. Unfortunately, many of the events had vertexes which were badly defined in two views and could not be measured.

It was felt that the 67% rejection rate for third measures was too high to justify remeasuring events which had failed twice unless they could be handled by the special measuring procedure just described. This was particularly true because the difficulties appeared to be in the computer program rather than in the measuring. Any attempt to improve the results would have required either extensive rewriting of the program or widening the rejection criteria and accepting large errors in the measured quantities. It was decided to leave the events unmeasured and make corrections for them which will be described later.

B. Kinematic Fitting

a. Theory. - All events which passed Datpro were put through a kinematic fitting routine called Guts. Guts is a well-known program and has been described in detail in the literature.²⁸ Its operation is based on the fact that due to errors intrinsic to the measuring process, the measured values of the momenta and energies of the particles rarely, if ever, satisfy conservation of energy and momentum. Consequently, it

should be possible to get more accurate estimates of the true values by adjusting the measured momenta and energies to conform to the conservation laws. The amount of adjustment allowed should depend on the errors in the measured variables so that more freedom is given to quantities which have larger errors. Guts makes this adjustment by minimizing the test quantity

$$\chi^2 = \sum_{ij} (x_i - x_i^m) G_{ij}^{-1} (x_j - x_j^m) \quad (4)$$

subject to the constraints of conservation of energy and momentum. That is, the final fitted values of the variables always obey the conservation laws exactly even though the initial values did not. In the definition of χ^2 , x_i^m is the measured value of a kinematic quantity and x_i is the adjusted value. G_{ij} is an error matrix defined by $G_{ij} = \overline{\sigma_i \sigma_j}$ where σ_i is the error in x_i^m . The kinematic quantities used are the curvature ($k = \frac{1}{p \cos(\text{dip})}$), $\tan(\text{dip})$, and azimuth for each track. These were chosen because they are closest to what was actually measured and one hopes that the errors will be normally distributed.

In the process of choosing adjusted values for this experiment, Guts considered only the three hypotheses: elastic scattering, π^0 production and π^+ production. Every event was put through each of these fits and the measured quantities compared to fitted quantities using χ^2 as a goodness of fit criterion. The values of χ^2 resulting from the minimizing process will fall into two groups depending upon whether the event is elastic or inelastic; for elastic events the distribution of χ^2 will be one of 4 degrees of freedom, whereas for inelastic events it will have only 1 degree of freedom. It can be shown²⁸ that the number of degrees of freedom for the χ^2 function is equal to the number of equations of constraint used in the fitting process. For elastic events all four

equations are applied, whereas for inelastic events three equations are needed to determine the kinematics of the missing particle, leaving only one equation to be used in the fitting process. In general, if there are n pieces of data and the fitting procedure (least squares) is used to determine $m < n$ parameters, then the resulting distribution of χ^2 has $r = n - m$ degrees of freedom. In the case of elastic events, there are 9 pieces (3 for each particle) of experimental data. Since there are four equations of constraint, the least squares fitting procedure is really determining $9 - 4 = 5$ quantities. Therefore the χ^2 distribution will have $9 - 5 = 4$ degrees of freedom. For the inelastic events there are also 9 pieces of data and one is attempting to determine $12 - 4 = 8$ quantities (there are really 4 particles involved in an inelastic process and so therefore there are $3 \times 4 = 12$ kinematic quantities for each event). Therefore the χ^2 distribution will have only $9 - 8 = 1$ degree of freedom.

Strictly speaking, the error matrix is not diagonal because all the calculated quantities are related via the initial measurements.

However, Datpro involves such a large number of calculations, that one expects the connection between variables to be small. Consequently, we chose to ignore the correlation terms and used a diagonal error matrix, $G_{ij} = \sigma_i \sigma_j \delta_{ij}$. In this form, G_{ij}^{-1} becomes $G_{ij}^{-1} = \frac{1}{\sigma_i \sigma_j} \delta_{ij}$ and we can write (4) as:

$$\chi^2 = \sum_i \frac{(x_i - x_i^m)^2}{\sigma_i^2} \quad (5)$$

Equation 5 shows that underestimating an error will result in an χ^2 which is too large. Conversely, an overestimate of an error gives too small an χ^2 . Thus any systematic mistake in estimating errors will give the wrong χ^2 distribution when the events are processed.

b. Error analysis. - The error formulas used for this experiment were of the general form $\left(\frac{\Delta x}{x}\right)^2 = \frac{a}{(L \cos \text{dip})^n}$. These were derived semi-empirically on the basis that the largest errors should be associated with short projected track lengths (cf. Appendix I). The formulas used were

$$(\Delta \text{az})^2 = \frac{11.6}{(L \cos \text{dip})^{1.79}} \quad (\text{radians})^2 \quad (6a)$$

$$(\Delta \text{dip})^2 = \frac{1360}{(L \cos \text{dip})^{1.93}} \quad (\text{radians})^2 \quad (6b)$$

$$\left(\frac{\Delta P}{PR}\right)^2 = \frac{167}{(L \cos \text{dip})^{4.03}} \quad (\text{mils})^{-2} \quad (6c)$$

The functional form of these equations was suggested by theory, e.g., the R in (6c) reflects the fact that a track with a large radius of curvature will have a small sagitta and be difficult to measure. The exact values of the coefficients and exponents were determined by repeated remeasurements of a group of events. Five events were remeasured 40 times and 4 events were remeasured 10 times. A standard deviation was obtained for each track and variable and the left hand sides of 6a-6c were plotted vs L cos dip on log-log paper. The points were found to lie on straight lines and least square fits gave the final coefficients and exponents.

The formulae 6a-6c represent only measuring errors. The complete formulae used in Guts also include terms for multiple scattering. It was often true that the multiple scattering error in the azimuth was greater than the measuring error, whereas the measuring error was much more important for dip.

Guts produced a set of variables called the stretch function which were used to test the error assignments. These are defined by

$$S_i = \frac{x_i^f - x_i^m}{\sigma_i} \quad (7)$$

where x_i^f is the final fitted value of the variable and x_i^m is the initial measured value. It is clear that the stretch functions will be Gaussian with a mean of zero and a variance of one if the errors are properly assigned. It was found that the error formulae 6a-6c produced stretch functions which were reasonably close to the ideal distribution (Fig. 9).

c. Determination of momentum.- A constant value was used for the momentum of the incident track rather than the value computed by Datpro. This was felt to be reasonable because the measuring errors for the incident track were often much larger than the known momentum spread of the beam. The fixed incident momentum was adjusted for energy loss by using the formula $p = p_0 - \left. \frac{dp}{dx} \right|_{\min} xL$ to give a momentum spread across the chamber. Here p_0 is the constant input momentum, L is the length of the incident track, and $\left. \frac{dp}{dx} \right|_{\min}$ is the rate of momentum loss at minimum ionization. The errors in the incident momentum was always taken to be 3% of p .

The best value of p_0 to use was determined by studying elastic events. Elastic events were used for the following reasons:

1. All four constraints of conservation and momentum are available, and it was expected that this would make the fitting procedure more reliable than it would have been if inelastic events had been used.
2. They could be identified by coplanarity and angular correlation so the identification did not require χ^2 . This was useful because χ^2 was a function of p_0 .

When p_0 was set at 786 MeV/c, the average fitted momentum was 778 MeV/c and the stretch function for K_{inc} had a mean of 0.5. If we recall that $k_{inc} = \frac{1}{p_{inc} \cos(dip)} = \frac{1}{p_{inc}}$ (since incident tracks have dip = 0) we see that (7) gives

$$S_{k_{inc}} = \frac{\frac{1}{p_{inc}^f} - \frac{1}{p_{inc}^m}}{\sigma\left(\frac{1}{p_{inc}}\right)}$$

Thus a positive value of $S_{k_{inc}}$ indicates that the fitted momentum is smaller than the measured momentum. A trial of $p_0 = 777$ MeV/c gave a stretch function with a mean of 0.2 which showed that the momentum was still too large. The final value of 775 MeV/c for p_0 gave a mean of -0.028 for $S_{k_{inc}}$. The minus sign indicates that the momentum was a little too small but it was not considered worthwhile to try again.

It should be noted that the final value of 775 MeV/c for p_0 agrees with the wire orbit measurements. However, this was an independent determination since the value of p_0 was forced on us by the computer. Fig. 10a and 10b show the momentum distribution for the incident track of elastic events in the center of mass and the laboratory. The final average laboratory momentum for all events was 772 MeV/c. This differs somewhat from p_0 because of the energy loss mentioned previously. The average momentum represents the incident momentum evaluated at the center of the chamber whereas p_0 is the momentum at the beginning of the incident track.

Guts was originally set up to use the incident momentum calculated by Datpro. However, it was found that the average momentum from Datpro was 812 MeV/c and that Guts reduced this to 786 MeV/c during the fitting process. The fact that the average incident momentum calculated

by Datpro was found to be too large suggested that the magnetic field (Eq. 2) was incorrectly normalized. An attempt was made to test this hypothesis by examining events with long tracks. The test was restricted to such events because it was expected that measuring errors would be small for long tracks. It was again found that Guts lowered the momentum when making a fit. Consequently, the momenta of outgoing tracks were multiplied by 0.965 in order to make an approximate correction. The correction factor of 0.965 is the ratio $\frac{p_o}{p_{\text{Datpro}}}$ where p_o was taken as 786 MeV/c and p_{Datpro} was taken as 812 MeV/c. 786 MeV/c was used instead of the correct incident momentum of 775 MeV/c because the correction procedure was written before the correct momentum was learned. Changing the correction factor after the true momentum was learned would have been uneconomical because it would have been necessary to recompute several thousand events. In any case, the effect of using 786 MeV/c rather than 775 MeV/c is very small because of the parameter adjustment procedure inherent in the Guts program.

A correction was also applied to the outgoing tracks in order to allow for energy loss along the track. It was mentioned previously that Datpro assumes that a track can be fitted by a circle of constant radius. Actually, the energy loss would cause the radius to decrease as one moves away from the vertex and it is reasonable to assume that the measured momentum is the momentum about half way along the track rather than the momentum at the vertex. This correction was not difficult to make for pions since they were almost always minimum ionizing. The program used $p = p_{\text{meas}} + \left. \frac{dp}{dx} \right|_{\text{min}} \times F \times L$ where F is a factor that determines what fraction of the length is used. The problem was simulated on a computer and it was found that F depended slightly on the momentum,

position and length of the track but that 0.5 was a reasonable choice which was good for most cases.

The problem was more difficult with protons because the energy loss changes rapidly with energy. It was apparent from the graphs in the Berkeley range-energy tables²⁹ that one could write $p = cR^\alpha$ to describe the range momentum relations. Let P_m and R_m be the measured momentum and corresponding range. One can write

$$P_{\text{True}} = c(R_m + \Delta R)^\alpha = cR_m^\alpha \left(1 + \frac{\Delta R}{R_m}\right)^\alpha = P_m \left(1 + \frac{L \times F}{R_m}\right)^\alpha$$

as the correction for energy loss. The equations used were ($P = 142R^{.278}$, $F = .6$) for P_m less than 500 MeV/c and ($P = 110R^{.333}$, $F = .5$) for P_m greater than 500 MeV/c where R is in centimeters and P is in MeV/c. Here again, F was determined from a computer program. The net effect of these corrections was usually a decrease in momentum for pions and an increase for protons. The change for low momentum protons was often several times as large as the measuring error. The positive track was treated as a proton when attempting to fit the elastic or π^0 production hypothesis whereas it was treated as a pion for the π^+ production fit.

The corrections discussed above were not needed for tracks which ended in the chamber because the momentum of these tracks was determined from their range. No serious error was made if a stopping track was treated as a regular track, but treating a track which did not stop as a true stop resulted in poor or wrong fits by Guts. These were found and corrected later when the results of Guts were examined in detail.

d. Fitting procedure.- Guts tried to fit events only after making the above error calculations and momentum adjustments. It began by assuming the positive track was a proton and attempting fits to elastic scattering and π^0 production. It then treated the positive track as a π^+ and tried the π^+ production fit.

Guts contained a number of tests which were used to check the quality of a fit. The most important of these were: too large a χ^2 or too many iterations needed to make a fit. These tests were designed to be very liberal so that the experimenter would have freedom to examine the details of poor fits. As an example, occasional inelastic fits were found with χ^2 as large as 500 even though χ^2 of 6.64 represents a confidence level of less than 0.01 for such fits. The broadness of the tests made it very reasonable to assume that a fit which failed Guts could be definitely eliminated.

All events were put through all three hypotheses. All satisfactory fits for an event were written as binary records on magnetic tape. A record contained the input data, the fitted momenta and angles, the stretch functions and a number of functions for testing the fit. An event could have up to three records since successful fits were sometimes made to all three hypotheses. Events for which no fits could be made were listed and examined later.

e. Test functions.- The test functions provided by Guts were: χ^2 , the missing mass, and coplanarity and correlation functions. χ^2 has already been explained. The missing mass was defined as $(E_0 - E_1 - E_2)^2 - (\vec{P}_0 - \vec{P}_1 - \vec{P}_2)^2 = M^2$. In this formula, $(E_0, \vec{P}_0)$ are the measured incident energy and momentum and $(E_1, \vec{P}_1)$, $(E_2, \vec{P}_2)$ are the measured outgoing

energies and momenta. E_1 is defined by $E_1 = \sqrt{p_1^2 + m_1^2}$ ($c=1$) and is thus the total energy. The terms $E_0 - E_1 - E_2$ and $\vec{P}_0 - \vec{P}_1 - \vec{P}_2$ represent the energy and momentum which must be carried by some missing particle in order to conserve energy and momentum. Thus M^2 is the squared mass of a hypothetical neutral particle. It is possible for M^2 to be negative and so M is defined as

$$M = \sqrt{M^2} \quad \text{if } M^2 \geq 0$$

$$M = -\sqrt{|M^2|} \quad \text{if } M^2 < 0 .$$

In this terminology, a negative missing mass really means an imaginary mass. It is clear that M should be 0 for elastic scattering, 135 MeV for π^0 production and 939 MeV for π^+ production. One cannot expect perfect agreement since measuring errors spread out the values of M , but this does give some indication of the goodness of fit.

The coplanarity and correlation tests were designed to help separate elastic and inelastic events. The coplanarity was defined by:

$$\text{COP 1} = \theta \quad \text{where } \cos\theta = \hat{P}_0 \cdot \frac{\vec{P}_1 \times \vec{P}_2}{|\vec{P}_1 \times \vec{P}_2|} .$$

In this equation, $\vec{P}_1 \times \vec{P}_2$ is the normal vector to the plane of the outgoing tracks and $\hat{P}_0$ is a unit vector in the direction of the incident track. For elastic events, all three tracks lie in the same plane and $\theta = 90^\circ$. We defined $\text{COP 2} = \frac{90 - |\text{COP 1}|}{\Delta \text{COP 1}}$ where $\Delta \text{COP 1}$ is the measuring error in COP 1. COP 2 is the relative error and is more meaningful than COP 1 since measuring errors will keep COP 1 from equalling 90° .

The correlation test used the fact that knowledge of the

incident momentum and the angle between one outgoing track and the incident direction is sufficient to predict the other angle for an elastic event. One thus has a kinematic line relating θ_π and θ_p for elastic scattering. The correlation function, CORR, was defined to be 0 for events which lay exactly on the kinematic line and essentially was a measure of the distance from the measured point (θ_π, θ_p) to the kinematic line. Here again we used the relative measure defined as $\text{CORR } 2 = \frac{|\text{CORR}|}{\Delta \text{CORR}}$ as a criterion.

C. Editor

A special program examined the binary output of Guts and attempted to classify the events. Each fit for an event was examined and tested separately and a record kept of the results. The tests were:

$$\text{Elastic} \quad \chi^2 \leq 11.67 \quad \text{and} \quad \text{COP } 2 \leq 2.3 \quad \text{and} \quad \text{CORR } 2 \leq 2.3$$

$$\text{Inelastic} \quad \chi^2 \leq 5.41$$

These limits were chosen as representing the 2% confidence level of ideal χ^2 and normal distributions. An event was considered elastic only if it passed all three tests.

An event was considered to be classified only if it passed the tests for one fit and failed the other fits. Events which failed all the fits or passed more than one fit were considered unclassified. The correct fit for a classified event was copied onto a new tape with the elastic, π^0 and π^+ events each going onto separate tapes. Thus a tape of successful fits for each of the three hypotheses was created. The χ^2 , COP 2 and CORR 2, and missing mass distributions are shown in Figures 11-13.

Approximately 28% of all events failed Editor. All the records for events which failed were written on a fourth tape called the rejection tape, for later examination.

D. Plotter

The set of programs Datpro, Guts, Editor, was designed to process all events without any spatial restrictions. It was felt that events should be retained for as long as possible in order to give the experimenter as much flexibility as possible.

Plotter was written to examine the fits accepted by Editor. It essentially consisted of two parts:

1. A rejection routine which rejected events for spatial or kinematic reasons.
2. A plotting program which could be set up to calculate any desired quantities for each event and to plot the results.

It was thus used to make the final choice of events and provide the necessary plots for analysis.

The plotter rejection criteria fell into three categories applied in the following order: those defining the fiducial volume (x-y limits), those concerning the az, dip and z of the incident track, and those which required good measurements on the outgoing tracks. Figure 14 shows top view of the final fiducial volume. The sides $y = 5''$ and $y = 9''$ correspond to the limits of the original scanning template. The right hand edge at $x = 12''$ also corresponds to the original template. The left hand end is an arc of a circle with a 5 inch radius. The circle is concentric with the chamber and thus guarantees that any track

emerging from a vertex lying at the edge of the circle will be at least 2 inches long before it enters the chamber wall. This restriction was intended to assure good measuring accuracy. The broken line at $x = 4''$ is the limit used for stops, and will be discussed below.

The z distribution of events is shown in Fig. 8. This was well defined and events were accepted only if $1.6'' \leq z \leq 6.2''$.

Fig. 15 a and b show the dip and azimuth distribution for the incident track. It was clear that the beam was well defined by these two plots and accordingly the following restrictions were placed on the incident track:

1. $-3^\circ < \text{dip} < +2^\circ$
2. azimuth to lie between the lines $\text{az} = 13.8 - x$ and $18.6 - x$
where x is the x coordinate of the vertex in inches.

The x coordinate of the right hand fiducial was $x = 12''$ so requirement 2 gave $1.8^\circ < \text{az} < 6.6^\circ$ for the acceptable entrance angle of events. These limits were also used in the final track count discussed earlier. The azimuth increases as the beam crosses the chamber because the tracks curved downwards. It is easy to show that if the track is a circle and the azimuth is small, then there is a linear relation between x and azimuth. This is the source of requirement 2. Approximately 2% of the classified events failed on azimuth, 5% failed on dip, and $1\frac{1}{2}\%$ failed on both azimuth and dip. These three categories are mutually exclusive, i.e. if an event failed on both az and dip it was not counted in either az or dip .

The third group of rejection criteria applied to the outgoing tracks. All non-stopping tracks had to be at least 1.5 inches long, no event was accepted if one of its tracks had a momentum error greater

than one-half of the momentum. In addition, no $|\text{dip}|$ was allowed to be greater than 67.5° . This last requirement recognized the trouble with the bad radius test discussed earlier and was designed to treat all events with large dips on the same footing. About 26% of the events entering Plotter failed to pass its criteria. These are classified according to the reason for failure in Table III.

E. Prirej

The events which were not classified by the Editor program could not simply be thrown away. They were examined and classified one at a time by a physicist at a scanning table. However, in order to reduce the number of events to be examined, they were first analyzed by a computer program called Prirej. This program rejected all those events which failed to pass the tests contained in Plotter and gave a detailed printed output for the remaining events. This method reduced the number of events to be examined by hand from 2228 to 1053.

The print out from Prirej for events which passed the Plotter criteria contained the ID, CORR 2, COP 2, and the input momenta, azimuth and dips. The program also printed the fitted momenta, azimuths, and dips, the missing mass, and the χ^2 for each fit along with an estimate of the apparent bubble densities for the outgoing tracks. Upon examination, it was possible to classify 782 of these events. About half of the remainder were remeasured and 50% of them failed Editor again. A detailed discussion of the unclassified events will be found in Chapter VI.

Many decisions could be made on the basis of the test quantities alone without needing much recourse to bubble density. The apparent bubble density of a track varied with position in the chamber and with

the view. Consequently, it was used either to confirm a decision already made on the basis of the test function or to make a decision when the difference between π^+ and proton bubble densities was clear. Those cases for which no decision could be made from the test functions and for which the bubble density was not clearly in favor of one or another hypothesis were treated as ambiguous and not listed as classified.

A test was made by examining Guts output from which χ^2 , missing mass, COP 2 and CORR 2 were omitted. Thus only bubble density was available for classifying the events. This test showed that bubble density alone was accurate to 88%. That is, 12% of the time, decisions made on bubble density alone disagreed with decisions made by Editor. When one considers that the physicist examining the unclassified events had much more information available and that doubtful cases were not classified, it seems reasonable to assume that the classified events were correct to 5%.

The appropriate records of those events which were successfully hand-classified were copied onto the corresponding input tapes for plotter. The final rejection tape thus contained only the unclassified events which failed the plotter tests and those that passed the plotter tests but could not be classified.

VI. RAW RESULTS AND CORRECTIONS

A. Events

a) Introduction. - The purpose of this chapter is to discuss the bookkeeping and corrections to the data which were necessary in order to obtain the final cross sections. The overall flow of events through the programs and analysis is shown in Fig. 16. It is convenient to divide the events into two distinct groups for discussion. The first of these is the group of 7897 events which were put into Editor. Table II gives a breakdown of these events and Tables IIIa and IIIb give an explanation of why events were rejected by Plotter and Prirej.

The second group of events (1252) consists of 973 events which were never successfully put through Datpro plus the 279 events rejected by Guts. These are itemized in Tables V and VI and will be discussed after the discussion and corrections for the first group.

b) Events examined by Editor. - Table II and Figure 16 show that 7897 events were examined by Editor. Of these events, 2684 failed the Plotter or Prirej tests and can be temporarily disregarded. A total of 4160 events were classified by Editor and passed the Plotter tests. The 2228 events which failed Editor were put through Prirej and 1053 passed the Plotter criteria and were examined by hand. 782 of these events were easily classified and were added to the Plotter tapes. The remaining 271 events which were examined by hand fall into a number of separate categories:

1. 71 events classified after the Plotter tapes were made. These were not included in the plots because it was inconvenient to do so.

2. 40 events called ambiguous because the physicist was unable to decide between two or more fits. These were split proportionally among the different fits. In making this split, only the relevant numbers were used, i.e. events which were ambiguous between π^0 and π^+ production were split by $R = \frac{n_+}{n_0 + n_+}$ where n_+ is the number of classified π^+ events and n_0 is the number of classified π^0 events.
3. 41 events had the characteristics that Guts made a fit only to π^+ production, the χ^2 was large, the missing mass was about 1100 MeV, and both outgoing particles had small momenta and were heavily ionizing. These characteristics were also shared by 33 events that were completely rejected by Guts. The most likely explanation of these events is that they are multiple pion production from the process $\pi^- + p \rightarrow \pi^- + \pi^+ + \pi^0 + n$ or $\pi^- + \pi^+ + \gamma + n$.
4. 35 events had only one reasonable fit but for some reason, e.g. bubble density, the physicist was not satisfied with it. 17 of these were added to the ambiguous events discussed in (2) and 18 of them were called unclassifiable.
5. Finally, 84 events were considered unclassifiable. A typical event had $\chi_{el}^2 = 59.6$, $\chi_{inel}^2 = 69.2$ and $p_{inc}(el) = 707$ MeV/c and $p_{inc}(inel) = 957$ MeV/c. There was no justification for accepting either fit and no evidence that remeasuring the event would improve the fit. Indeed, some of them were remeasured, and no improvement was found. These events may represent cases for which the incident particle had a different momentum from the beam, i.e. the

Table II. Uncorrected Results for Measured and Computed Events

	Elastic	π^0	π^+	$\pi^+ \pi^- \pi^0 n$	Total
Classified by Editor	3310	983	1376	0	5669
Fail Plotter	834	254	421	0	1509
Balance	2476	729	955	0	4160
Classified by Hand					
Plotted	454	99	229	0	782
Unplotted	62	27	39	41	169
Good events	2992	855	1223	41	5111
Pass Plotter but Unclassifiable					102
Fail Plotter or Prirej					2684
Total Events					7897

incident particle may not have been a true beam particle.

These events were treated as being unclassifiable and the track count reduced proportionally.

The net result of these considerations is that out of 7897 events entering Editor 5111 were classified, 2684 were rejected on the basis of "Plotter" tests (in Plotter or Prirej) and 102 were unclassifiable.

c) Corrections to events examined by Editor.-Study of Table IIIa will show that events rejected by the Plotter criteria fall into two distinct groups. The first of these is those events rejected on beam tests (x,y,z, az or dip). These are not considered to be true events because the incident track was not part of the acceptable beam. They can be eliminated from the events used in calculating the cross sections,

Table IIIa. Reasons for Rejection by Plotter or Prirej

	Classified Events			Total Classified	Prirej	Total
	Elastic	π^0	π^+			
1. x or y	233	31	104	368	291	659
2. z only	23	4	3	30	9	39
3. az only	79	31	31	141	112	253
4. Both z and az	2	1	0	3	4	7
5. dip only	195	108	151	454	379	833
6. Both z and dip	11	5	3	20	26	46
7. Both az and dip	57	31	30	118	163	281
8. All of z, az, dip	5	6	10	21	26	47
<hr/>						
TOTAL BEAM REJECTIONS*	605	217	333	1155	1010	2165
<hr/>						
9. Length < 1.5 inch	75	14	35	124	106	230
10. $\Delta p > 1/2p$	23	0	0	23	32	55
11. $ \text{dip} > 67.5^\circ$	131	23	53	207	27	234
<hr/>						
TOTAL REJECTIONS**	834	254	421	1509	1175	2684

* These totals are the result of summing only rejections 1 through 8 which pertain to the incident track only.

** These totals include all the rejections (1 through 11).

Table IIIb. Fractional Rejections for az, z, or dip for Events
passing x and y

	Plotter			Prirej
	Elastic	π^0	π^+	
Total Events	3310	983	1376	2228
Failed x or y	233	31	104	291
Total pass x and y	3077	952	1272	1937
Percent Failed				
z only	0.75	0.42	0.24	0.46
az only	2.56	3.26	2.44	5.78
z and az	0.07	0.10	0.00	0.21
dip only	6.32	11.32	11.90	19.55
z and dip	0.36	0.53	0.32	1.34
az and dip	1.85	3.26	2.36	8.40
z, az and dip	0.16	0.63	0.79	1.34
% pass x and y but fail az, z, or dip	12.07	19.52	18.05	37.08

but they must be considered when the total track length is calculated. This point will be discussed in detail in the chapter on cross sections.

It may be of some interest to compare the percentage of events which passed x and y but failed on z, az, or dip. In making such a comparison, one must remember that the group of 1179 events rejected by both Editor and Prirej could never be classified as to event type. Therefore, the (elastic and inelastic) events shown in Table IIIa as rejected come only from the events which passed Editor and were subsequently rejected in Plotter. Table IIIb shows the percentage of failure for az, dip, or z with the 782 events which were hand classified included with the unclassified events. The most interesting result is the large difference between the elastic, inelastic and unclassified events which were rejected on dip. A possible reason for this difference is that events with bad dip may have the wrong momentum. If this were true, one would expect that Guts would find it easier to make an inelastic fit rather than an elastic fit. Consequently, these events would tend to concentrate in the inelastic or unclassified groups.

Another interesting point is that the overall rejection rate is much higher for unclassified events than for those classified by Editor. This suggests a correlation between the Plotter criteria and the ease of making a fit. This is as it should be if the Plotter criteria were correctly chosen to define the beam.

The second group of rejections shown in Table IIIa are events which passed the beam tests but were deliberately left out of the plots for one of the following reasons:

1. The length of an outgoing track was less than 1.5 inch.

2. The $|\text{dip}|$ of an outgoing track was greater than 67.5° .
3. The error in a momentum was greater than one half of the momentum.

These events are true events and must be included when calculating the cross sections. They were omitted from the plots because their characteristics implied large measuring errors which might distort the plotted quantities. The remainder of this section will discuss the way in which the number of events was corrected to allow for them.

The simplest correction to make is the 124 events shown in Table IIIa as classified but rejected on length. There is really no reason to question their classification and they can simply be added to the appropriate event types. The 106 events rejected in Prirej because of their length were not examined by hand. They can be most conveniently dealt with by adding them to the unmeasured events to be discussed in the next section.

The next correction to be discussed involves those events with large dip or large errors in momentum. All of the events which failed the bad radius test in Datpro had large dips. The events rejected by Plotter on the basis of $|\text{dip}| > 67.5^\circ$ should therefore be included with the bad radius events when making corrections to the raw results. Moreover, the error formula for momentum (6c) indicates that a large error in momentum implies either a short track length or a large dip. Events with short outgoing tracks were rejected by Plotter before the momentum test was made. This implies that many of those events which were rejected because $\Delta p > \frac{1}{2}p$ can be considered to have large dips. Thus the events with either bad radius, $|\text{dip}| > 67.5^\circ$, or $\Delta p > \frac{1}{2}p$ form a single group with the property of having an outgoing track with large

dip. It was possible to make detailed corrections for this group of events by examining a number of angular distributions in a system in which the incident track is taken as the x-axis. This system was defined by making two successive rotations of the coordinate system. The first was a rotation around the laboratory z-axis by an angle equal to the azimuth of the incident track. This produces an (x', y', z') system in which the x' axis lies in the plane defined by the incident track and the laboratory z-axis. The angle between the x' -axis and incident track is the dip of the track. We now rotate around y' through an angle equal to the dip of the incident track to produce the (x'', y'', z'') system in which the x'' -axis coincides with the incident track and the angle between the z and z'' axes is the dip of the track. Since the dip of most events was only a few degrees the z'' axis is really quite close to the chamber z axis. The angles used to make the corrections in this rotated system were defined by:

$$\cos\theta_{23} = \frac{(\vec{p}_- \times \vec{p}_+)_z}{|\vec{p}_- \times \vec{p}_+|}$$

$$\theta_{23} = \tan^{-1} \left(\sqrt{1 - \cos^2\theta_{23}} / \cos\theta_{23} \right)$$

$$\text{and } \phi_{\pm} = \tan^{-1} \left(\frac{p_{y_{\pm}}}{p_{z_{\pm}}} \right)$$

where $p_{\pm}$ refers to the momentum of the positive or negative outgoing track in the rotated laboratory system.

θ_{23} defines the angle between the normal to the plane of the outgoing particles and the new z axis. θ is the angle between the new z axis and the projection of a track on the new y-z plane. The angle θ_{23} was chosen as a useful variable for elastic scattering because in this case all three tracks lie in the same plane and the system should be independent of rotation of the plane around the incident track. Thus a plot of the number of events in an interval $\Delta \theta_{23}$ should in principle be independent of θ_{23} . Two effects were expected to alter this uniformity in the case of the observed elastic events.

1. Events for which the production plane was perpendicular to the windows ($\theta_{23} = 90^\circ$) were hard to see.
2. Elastic events with large dips had large θ_{23} since for elastic events one has $\theta_{23} = \tan^{-1} \left(\frac{p_z}{p_y} \right) = \tan^{-1} \left(\frac{\sin(\text{dip})}{\cos(\text{dip}) \sin(\text{az})} \right)$ or $\theta_{23} = \tan^{-1} \left(\frac{\tan(\text{dip})}{\sin(\text{az})} \right)$.

This meant that the events missing from the elastic plots because of large dips would be revealed as an asymmetry in the θ_{23} plots.

θ_{23} was not a convenient variable for inelastic scattering because the incident track does not in general lie in the plane defined by the two outgoing charged particles. This is because inelastic events contain an unseen neutral particle which carries off some of the momentum. This introduces kinematic effects which can prevent the θ_{23} distribution from being uniform. For example, $\theta_{23} = 90^\circ$ indicates that the plane of the outgoing charged particles is perpendicular to the window. This plane must contain the incident track for elastic scattering but can make an angle of up to 90° with the incident track for inelastic scattering. Accordingly, there is no clear reason for θ_{23} to have a

uniform distribution for inelastic events. The ϕ variables were more appropriate for inelastic scattering because of the fact that the distribution of outgoing tracks should be independent of rotation around the incident track direction for an unpolarized beam and target. $\phi = \tan^{-1} \left(\frac{p_y}{p_z} \right)$ measured the projection of the momentum on a plane perpendicular to the incident track and the number of events in the range $\Delta\phi$ should be independent of ϕ . One can see the relation between ϕ and dip by writing $p_y = p_o \cos(\text{dip}) \sin(\text{az})$, $p_z = p_o \sin(\text{dip})$ and $\phi = \tan^{-1} \left(\frac{\cos(\text{dip}) \sin(\text{az})}{\sin(\text{dip})} \right) = \tan^{-1} (\tan(90-\text{dip}) \sin \text{az})$. Events with large dip will have small ϕ and the missing events will show up as an asymmetry in the ϕ distribution.

For elastic events, conservation of momentum required that $\phi_+ = \phi_-$. Moreover, since we had $\phi = \tan^{-1} \left(\frac{p_y}{p_z} \right)$ and $\theta_{23} \text{ (elastic)} = \tan^{-1} \left(\frac{|p_z|}{p_y} \right)$, the three corrections were equivalent for elastic events and were expected to give the same results. The corrections for elastic events thus formed a convenient check on the computer program used to calculate θ_{23} and $\phi_{\pm}$ and showed that it worked correctly.

Figure 17 shows the $\phi_{\pm}$ and θ_{23} plots used to make the corrections for large dips. There are clear asymmetries in all of the plots except $\phi_+(\pi^0)$. A study of the kinematics of π^0 production showed that the maximum laboratory angle between the incident track and the proton was approximately 65° . This implied that very few proton tracks in π^0 production would be rejected on bad radius or large dip. Hence, it was reasonable for $\phi_+(\pi^0)$ to have no asymmetry. A plot was also made of ϕ_+ vs ϕ_- for π^+ production and it was found that the two corrections were independent, i.e., one had to add the corrections from ϕ_+ to that from ϕ_- . (It was estimated that no more than 13 events were common to

ϕ_+ and ϕ_- .) This was considered reasonable because there was no simple relationship between $\vec{p}_-$ and $\vec{p}_+$ for inelastic scattering.

The corrections were calculated by taking the average number of events per bin from 0 to 70° for θ_{23} and 20 to 90° for ϕ_+ and extrapolating to get the missing events. The results are shown in Table IV below and in Figure 17.

Table IV. Corrections for large dip

	θ_{23}	ϕ_+	ϕ_-	common	Total
Elastic	351	--	--	--	351
π^0 prod.	---	0	68	--	68
π^+ prod.	---	108	85	-13	180
					<hr/> 599

This total of 599 events missing because of large dips should be compared to the events known to be missing. From Tables V and IIIa we have 308 bad radius events expected to pass Plotter and 289 events missing because of $\Delta p > \frac{1}{2} p$ or large |dip|. This gives a total of 597 which compares very well with the calculated correction of 599. It was therefore decided that the above corrections accounted for all the events missing because of large dips, bad radius or poor momentum.

d) Corrections for Events not examined by Editor.- We can now discuss the events which are shown in Figure 16 as having never been measured, rejected by Datpro, or failed Guts. These are itemized in detail in Tables V and VI below.

Table V. Events not computed by Guts

Reason	Total	Expected to fail Plotter	Expected to pass Plotter
Bad Radius	424	116	308
Poor Vertex	131	36	95
Miscellaneous	118	32	86
Never measured	300	82	218
Total	973	266	707

Table VI. Events which failed Guts.

Type	Number
2p - failed Plotter	167
2p - passed Plotter	
$\pi^- - \pi^+ - \pi^0 - n$	33
Dalitz pair	6
4p (not checked by Plotter)	
Dalitz pair	10
$\pi^- \pi^+ \pi^- p$	17 *
1p (not checked by Plotter)	46
Total	279

* Some of these may be from $\pi^0 \rightarrow \gamma + e^+ + e^-$ where the e^+ and e^- have high momenta.

The columns of Table V which are labeled as expected to fail or pass. Plotter are an attempt to estimate how many of the uncomputed events would have passed or failed the x, y, z , az , or dip criteria for the incident track. Table IIIa shows that 2165 events failed for one of these reasons. This is 27.4% of the total of 7897 events which entered the Editor routine. Consequently the totals shown in Table V were multiplied by 0.274 to correct for the beam criteria. The 308 expected good events which failed because of bad radius have already been discussed in the previous section and are included in the large dip correction. The remaining 399 events may be lumped with 106 unclassified events rejected on length (mentioned in the previous section) to give a total of 505 unclassified events. It will be shown in the chapter on cross section that these may be handled by splitting them proportionally among the events. For convenience the results of this split have been included in Table VII as a miscellaneous correction. It should be noted that the split is made using the number of events shown in the subtotal row of Table VII. That is, all explicit corrections had already been made before the proportional split was done.

We now turn to the events which failed Guts and are listed in Table VI. Of the two prong events which passed Plotter, 33 have the characteristics previously associated with $\pi^- \pi^+ \pi^0 n$ or $\pi^- \pi^+ \gamma n$. The other 6 had spiral tracks typical of low energy electrons and are probably Dalitz pairs from π^0 decay. The 10 events shown as Dalitz pairs under 4p events also have one or two spiraling tracks and appear to be $\pi^- p \rightarrow \pi^- p \pi^0$ followed by Dalitz decay of the π^0 . There were 17 four-track events which were not clearly Dalitz pairs and were accepted as

cases of $\pi^- \pi^+ \pi^- p$. This is an upper limit for the number of cases of $\pi^- \pi^+ \pi^- p$ because some of them may be high energy Dalitz pairs. Finally, 46 events were found with only 1 visible prong. Presumably, these are small angle elastic scattering and are accounted for by the small angle correction discussed later.

Table VII below itemizes all the corrections which have been discussed above and shows the grand total of events. The small angle correction for elastic scattering will be discussed in detail in Section VIII on elastic scattering. It allows for events missed because of small scattering angles and was calculated by extrapolating the differential cross section to 0° .

Table VII. Corrected Number of Good Events

	Elastic	π^0	π^+	$\pi^- \pi^+ \pi^0 n$	$\pi^- \pi^+ \pi^- p$	Total
classified, plotted	2930	828	1184	-	-	4942
large dip	351	68	180	-	-	599
classified, unplotted	62	27	39	74	17	219
length	75	14	35	-	-	124
Dalitz pair	-	10	-	-	-	10
Subtotal	3418	947	1438	74	17	5894
miscellaneous	298	82	125	-	-	505
correction to large dip	52	10	30	-	-	92
small angle	153	-	-	-	-	153
Total	3921	1039	1593	74	17	6644
	± 80	± 34	± 45	± 9	± 4	± 118

Another small correction has not yet been discussed. It is shown in Table VII as "correction to large dip". It is based on the fact that the elastic, π^0 and π^+ events shown as classified but un-plotted, length or miscellaneous were not included in the θ_{23} or $\phi_{\pm}$ plots. If they had been included, we would expect that they would have a distribution similar to the events which were plotted. For example, there are 435 elastic events in the three categories just mentioned. If a θ_{23} plot were made for these events, we would expect a correction of $435 \times 351 / 2930$ or 52 events as shown in Table VII.

e) Correction to differential cross section.- All of the above corrections are total corrections, i.e. they apply to the calculation of cross sections for the various reactions listed in Chapter IV. It is necessary to split the correction to elastic scattering into a set of corrections for various ranges of $\cos\theta^*$ (where θ^* is the center of mass scattering angle) before calculating the differential cross section. These ranges and corrections are shown below in Table VIII. It should be noted that some ranges overlap. This makes it possible to interpolate and get more detailed corrections than would be possible with one set of ranges. The rather peculiar choice of ranges was dictated by the need to have enough events in a bin to provide good statistics. It also took into account the fact that the kinematics of elastic scattering restrict events with large dips to the range $\cos\theta^* > .7$ and $-.1 > \cos\theta^* > -.8$, so that most of the corrections should be in these ranges.

Table VIII. θ_{23} Corrections to Differential Cross Section

Range of $x = \cos\theta^*$	correction	Range	correction
$-1 \leq x < -0.5$	54	$-1 \leq x < -0.7$	22
$-0.5 \leq x < +0.2$	24	$-0.7 \leq x < -0.2$	53
$0.2 \leq x < 0.5$	6	$-0.2 \leq x < 0.4$	0
$0.5 \leq x < 0.6$	1	$0.4 \leq x < 0.6$	10
$0.6 \leq x < 0.7$	14		
$0.7 \leq x < 0.8$	50		
$0.8 \leq x < 0.9$	93		
$0.9 \leq x$	109		

Table IX shows the bins of $\cos\theta^*$ used in calculating the differential cross section and the corresponding uncorrected numbers and corrections. In order to make the detailed corrections shown, it was sometimes necessary to split the θ_{23} correction into smaller bins, e.g. the correction for $\cos\theta^* < -0.7$ must be subdivided into three parts. This was done by dividing the corrections proportionally among the bins, i.e., $n_c = \frac{n_i}{\sum n_i} c$ where n_i is the number of uncorrected events in the bin and the sum is over the bins included in the correction c . The correction for miscellaneous events was always done in this manner using the sum of the uncorrected events and the θ_{23} correction for n_i ; i.e. $\sum n_i = 3281$ events. The term miscellaneous events refers to all the corrections listed for elastic events in Table VII except the large dip and small angle corrections. The small angle correction applies only to the bin $\cos\theta^* \geq 0.9$ and is not included here because this bin is not needed for the analysis of angular distribution. This will be explained in detail in Chapter VIII.

Table IX. Angular distribution of Elastic Events

$\cos\theta^*$	uncorrected number	θ_{23} correction	miscellaneous correction ^a	total corrected number
1.0 to 0.9	391	109	72	572
0.9 to 0.8	415	93	76	584
0.8 to 0.7	426	50	71	547
0.7 to 0.6	364	14	56	434
0.6 to 0.5	256	1	38	295
0.5 to 0.4	189	9	30	228
0.4 to 0.3	137	0	21	158
0.3 to 0.2	127	0	19	146
0.2 to 0.1	71	0	11	82
0.1 to 0.0	48	0	7	55
0.0 to -0.1	22	0	3	25
-0.1 to -0.2	15	0	2	17
-0.2 to -0.3	15	4	3	22
-0.3 to -0.4	20	6	4	30
-0.4 to -0.5	38	11	7	56
-0.5 to -0.6	48	13	9	70
-0.6 to -0.7	69	19	13	101
-0.7 to -0.8	92	7	15	114
-0.8 to -0.9	101	8	16	125
-0.9 - 1.0	86	7	14	107
Total	2930	351	487	3768

^a miscellaneous: 487 events missing because they were classified but not put on plotter tape, ambiguous, not plotted because one track was short, assigned unmeasured, or correction to large dip.

B. Neutrals

a) Measurement. - The events discussed thus far have been those with charged outgoing tracks. These correspond to reactions 3a-3f. In order to quote a total cross section, it is necessary to know the total number of stops, i.e. reactions yielding only neutral particles. It was mentioned earlier that 39 rolls were scanned for stops. Three of these rolls could not be used for computing cross sections because no track counts were available for them. A test sample of 21 of the remaining 36 rolls were measured with the restriction that no stop was measured unless its ξ satisfied the requirement $0.1 < \xi < 2.4$. The Plotter criteria for stops were the same as for events except that the left-hand limit was taken as the straight line $x = 4''$ (see Fig. 14) to allow for the difference in scanning templates. The test sample showed that 54% of the stops passed the amended Plotter criteria. It also showed that stops for which the scanner found $\xi = 2.3$ were out the back window and should not have been measured. The 36 usable rolls contain 1898 stops satisfying the requirement $0.1 < \xi < 2.3$ and these were used when computing the cross section.

b) Corrections to neutrals. - A sample of about 1000 of the 1898 usable stops found was measured and it was found that Datpro rejected 30% of them. This was considerably higher than one would expect on the basis of measurements of events. A check showed that many of the rejections were on poor vertex fit. It is possible that this may be due to the fact that one has to measure the vertex of stops directly, whereas for events one can calculate it from the intersection of two tracks and should be able to find the vertex more accurately.

When the 678 measured stops which passed Datpro were put through Plotter, it was found that only 54% passed the criteria for stops. This is much lower than expected on the basis of results for events where the rejection rate was about 27%. A check showed that rejections on dip for stops were twice as large as for events and that the dip distribution for stops was much broader than the distribution for events.

It was conceivable that this large rejection rate for dip might be due to the reduced x limit used for stops, ^{so} a special run of the π^+ events was made using the reduced x -limit that was imposed for stops. It was found that the fraction of events which passed the x -y criteria and then failed on dip was independent of the x -limit used. It is very unlikely that the dip distribution for stops actually differs from that of events. The most probable explanation of the observed broadening is that it is due to the difficulty encountered in measuring.

The above discussion indicates that we cannot find the number of stops which would pass the x, y, z , az and dip criteria by using the 54% that passed the criteria for stops. However, the x -y test should not be sensitive to measuring error and it was found that 86% of the measured stops passed the x -y test. This gives 1632 stops on the 36 rolls which should pass the x -y test. Referring to Tables IIIa and IIIb, we find that 7238 events pass the x -y test for events and that 1506 of these fail on az , z , or dip. This means that 79.2% of all events passing x -y also pass az , z or dip. If we assume that this is independent of the x -y limit, then we have $N_s = 1293 \pm 43$ stops on 36 rolls which pass x, y, z , az , and dip. This is 68.1% of the original number rather than 54%. The error of 43 stops is the statistical error in

the original number.

We will need several other numbers in addition to N_s when we calculate the track length. The first of these is the number (N'_s) of stops which would have been found if all frames scanned for events had also been scanned for stops. In order to do this, we first note that there were 3.436×10^5 tracks on the 42 rolls that were scanned for events. The 36 rolls scanned for stops had 2.584×10^5 tracks. However, this number is too high because the track count for stops used a wider azimuth limit than the track count for events even though the scanning criteria for stops and events were identical. A recount of ten rolls of stops, using the final azimuth limits used for events, showed that the original stop track count should be multiplied by 0.794 in order to reduce it to the limits used for events. This gave a corrected total of 2.052×10^5 tracks on the frames scanned for stops. We define ϵ_T as $\epsilon_T \equiv \frac{3.436}{2.052} = 1.67$ as the correction for scanning fewer frames for stops than for events. The second correction needed allows for the fact that the average track length on frames scanned for stops was 20.3 cm but was 24.6 cm on frames scanned for events. This gives $\epsilon_L \equiv \frac{24.6}{20.3} = 1.21$ as the correction for the difference in average track length. Then $N'_s \equiv \epsilon_L \epsilon_T N_s = 2613$ stops if all frames scanned for events had been scanned for stops.

The second number of interest is the reverse of N'_s . It is the number of events (N'_e) which would have been found if only frames scanned for stops had been scanned for events. If we let N_e be the total of 6664 events listed in Table VII, then $\frac{N'_e}{N_e} = \frac{N_s}{N'_s}$, or $N'_e = 3288$ events. These two numbers (N'_s , N'_e) are not used directly for computing the cross

sections, but they are used to make a small correction in the total track length.

C. Scanning Efficiency

A correction must be applied to the data in order to allow for events missed in scanning. The scanning efficiency was computed by double scanning 23 rolls for events. Assume that there are actually N_0 events on a roll and that scanner A finds N_A events and scanner B finds N_B events. We define the efficiencies by $N_A = \epsilon_A N_0$ and $N_B = \epsilon_B N_0$. The simplest assumption is that the efficiencies are independent, i.e. A has as much chance of finding events found by B as he has of finding events not found by B. Under this assumption, the number of events found by both scanners is $N_C = \epsilon_A \epsilon_B N_0 = \epsilon_A N_B = \epsilon_B N_A$ and we can write $N_0 = \frac{N_A N_B}{N_C} = \frac{N_A}{\epsilon_A} = \frac{N_B}{\epsilon_B}$. This technique can be expected to overestimate the efficiency because some events will be much more likely to be missed than the majority of events and our assumption will be wrong. However, it does set a lower limit to the efficiency correction. It was necessary to use an average efficiency for the experiment because all of the rolls were not double scanned. Let ϵ_i be the average efficiency for scanner i computed from the rolls scanned by him that were later double scanned, and let n_i be the total number of rolls scanned by i. Then the average efficiency was defined by

$$\epsilon = \frac{\sum n_i \epsilon_i}{\sum n_i}.$$

On the basis of 23 rolls double scanned for events and 10 rolls double scanned for stops we have $\epsilon(\text{events}) = .977$ and $\epsilon(\text{stops}) = .968$. It is reasonable for the efficiency for events to be higher than the

for stops because stops are much less distinctive and do not catch the eye as well as events. However, one must remember that the chance of an event being missed by both scanners during a double scan is negligible if each scanner has an efficiency of 98%. Since half of the rolls were double scanned for events, it is reasonable to take $\epsilon(\text{events}) = 0.99$ instead of 0.977.

VII. CROSS SECTIONS

A. Definition

The total cross section is rigorously defined by the relation

$$N_T = N_0 \left(1 - \exp(-\sigma_T n \langle L \rangle) \right) \quad (8)$$

where N_T is the number of events of all kinds, N_0 is the number of particles crossing the target, n is the number of target atoms per cm^3 , σ_T is the total cross section per scatterer, and $\langle L \rangle$ is the average track length for an incident particle. The argument of the exponential is small (on the order of 0.03 for this experiment) and we can expand equation (8) to give

$$N_T = N_0 \sigma_T n \langle L \rangle \left(1 - \frac{1}{2} \sigma_T n \langle L \rangle + \dots \right)$$

We can take the first term as a very good first approximation to σ_T and use it to evaluate the correction term, $\sigma_T^{(1)} n \langle L \rangle = N_T / N_0$, and therefore we have $N_T = N_0 \sigma_T n \langle L \rangle \left(1 - \frac{1}{2} \frac{N_T}{N_0} \right) = \sigma_T n \langle L \rangle (N_0 - \frac{1}{2} N_T)$. This form gives us a physical meaning for the second term. $\langle L \rangle$ is the average track length and is calculated for tracks which go through the chamber.

However, N_0 is the total number of tracks including those which yield events. The term $N_0 \langle L \rangle$ therefore overestimates the total track length because tracks which end in events have an average length of $\frac{1}{2} \langle L \rangle$.

The second term $(-\frac{1}{2} N_T \langle L \rangle)$ corrects for this.

The partial cross sections are related to the total cross section by $\frac{\sigma_i}{\sigma_T} = \frac{N_i}{N_T}$ where σ_i and N_i are the cross section and number of events of type i . This can be written $N_i = \frac{N_T}{\sigma_T} \sigma_i$ or $N_i = \sigma_i n \langle L \rangle (N_0 - \frac{1}{2} N_T)$.

Thus the general expression $\sigma_1 = \frac{N_1}{nL}$, holds for either partial or total cross sections, where we have written L' for $\langle L \rangle (N_0 - \frac{1}{2} N_T)$ as the total effective track length.

B. Total Track Length

All classified events which passed the Plotter criteria were put through an orbit program which traced the path of the incident particle back to the entrance of the fiducial volume and forward to the point where it would have left the fiducial volume. The program assumed that the incident track was a circle and used the vertex coordinates, fitted azimuth and the radius of curvature to determine the coordinates of the center of the circle. The radius of curvature was found from the equation $P = kHR$ where P was the fitted momentum and H was the corrected average field for the track computed by Datpro. The fitted values of azimuth and momentum were used because they should be more accurate than the measured values. This method should give an unbiased average track length because the events are a random sample of all the pions entering the chamber. The result is an average track length of $\langle L \rangle_{\text{events}} = 24.62 \pm 0.05$ cm.

The same program and events were rerun using the fiducial volume for stops rather than events. This gave an average track length for stops of $\langle L \rangle_{\text{stop}} = 20.27 \pm 0.05$ cm. This method should be safe because there is no reason to believe that tracks ending in stops are different in any manner from tracks ending in events. The ratio $\frac{\langle L \rangle_{\text{event}}}{\langle L \rangle_{\text{stop}}} = 1.215$ was used earlier to normalize the number of stops to the event fiducial volume.

The orbit program also gave the distributions of the azimuth,

z, and y of the incident tracks at the entrance to the fiducial volume. These distributions were used in a Monte Carlo program to calculate the track length. The length distribution calculated in this manner agreed with the distribution found by the orbit program. Moreover, rerunning the Monte Carlo program several times showed that the average track length was insensitive to statistical fluctuations in the z, azimuth, and y distributions and that small systematic changes in these distributions did not effect the result. The errors quoted for track length are based on the observed statistical fluctuation in the Monte Carlo calculation.

The total track length is given by

$$L' = \langle L \rangle \left(N_0 (1 - \epsilon_\mu) (1 - \epsilon_{z,D}) - \frac{1}{2} (N_E + N'_S) \right). \quad (8)$$

N_0 is the number of tracks counted with azimuth limits corresponding to the Plotter criteria and ϵ_μ is the ratio $\frac{\mu + e}{\pi + \mu + e}$. Thus $N_0 (1 - \epsilon_\mu)$ is the number of incident pions which pass the azimuth limits. $\epsilon_{z,D}$ is defined by

$$\epsilon_{z,D} = \frac{\text{No. of events which pass x, y, az but fail z or dip}}{\text{No. of events which pass x, y and az}}$$

The term $(1 - \epsilon_{z,D})$ corrects the total number of pions to the number which would pass z and dip. It should be noted that azimuth has been accounted for in the track count and x, y limits are included in the calculation of $\langle L \rangle$. Finally, the term $\frac{1}{2} (N_E + N'_S)$, where $N_E + N'_S$ is the number of good events plus the corrected number of stops, is the correction to the track length previously discussed. Substituting the values:

$$\langle N_0 \rangle = \text{ave. number of tracks/frame} = 13.123 \pm .013$$

No. of frames scanned for events = 26,181

$$N_0 = (3.436 \pm 0.03) \times 10^5 \text{ tracks}$$

$$\epsilon_\mu = 0.145 \pm 0.005$$

$$\epsilon_{z,D} = 0.138 \pm 0.005$$

$$N_e + N'_s = 6644 + 2613 = 9257$$

$$\langle L \rangle = 24.62 \pm .05 \text{ cm for events}$$

we find the total track length for events is $L'_{\text{event}} = (6.118 \pm .09) \times 10^6 \text{ cm.}$

We also need the total track length for stops. This is calculated using

$$N_0 = (2.052 \pm .02) \times 10^5 \text{ tracks}$$

$$N'_e + N_s = 3288 + 1293 = 4581$$

$$\langle L \rangle_{\text{stop}} = 20.27 \pm .05 \text{ cm for stops}$$

and we find

$$L'_{\text{stop}} = (3.030 \pm .045) \times 10^6 \text{ cm.}$$

A problem which must be discussed at this point is what to do with events that were omitted from the plots. Suppose that N_G events were plotted and that N_B events were not plotted but passed the beam criteria. There are two equivalent methods of dealing with the N_B events.

1. Assign them proportionally:

$$N'_i = N_i + \frac{N_i}{N_G} N_B = N_i \left(\frac{N_G + N_B}{N_G} \right)$$

2. Reduce the track length and write:

$$L'' = \frac{N_G}{N_G + N_B} L'$$

$$\sigma_i = \frac{N_i}{nL''} = \frac{N_i}{nL'} \left(\frac{N_G + N_B}{N_G} \right)$$

The first method was used in Chapter VI to assign the 505 events shown in Table VII as miscellaneous. However, Table II shows 102 events abandoned as unclassifiable. It was suggested in Chapter VI that these might be events with a wrong incident momentum. It seems appropriate to use the second method to deal with these. Taking $N_B = 102$ and $N_G = 6644$ (Table VII), we define $\epsilon_m \equiv \frac{N_G}{N_G + N_B} = 0.985$ and $L''_{\text{events}} = \epsilon_m L'_{\text{events}} = (6.026 \pm .09) \times 10^6 \text{ cm}$.

C. Density of Hydrogen

The density was determined by two independent methods.

1. The vapor pressure of a liquid neon vapor pressure thermometer in contact with the chamber was $51 \pm 1 \text{ cm of Hg}$. This corresponds³⁰ to a liquid hydrogen density of $0.0626 \pm 0.0001 \text{ gm/cm}^3$.
2. The average range of muons from $\pi^+ \rightarrow \mu^+ \rightarrow e^+$ decay in the chamber was $1.029 \pm .025 \text{ cm}$. The Berkeley tables give a range of $1.103 \pm 0.003 \text{ cm}$ at a density of $0.0586 \pm 0.0006 \text{ gm/cm}^3$.

Thus our range corresponds to a density of $0.0628 \pm 0.0016 \text{ gm/cm}^3$. The two methods are in good agreement. Taking the value of $0.0626 \pm 0.0001 \text{ gm/cm}^3$, 1.008 as the atomic weight of hydrogen and 6.0248×10^{23} as Avogadro's number, we have $n = (3.742 \pm .006) \times 10^{22} \text{ atoms/cm}^3$ and

$$\begin{aligned} nL''_{\text{event}} &= (225.5 \pm 3.3) \times 10^{27} \text{ cm}^{-2} \\ nL'_{\text{stop}} &= (113.4 \pm 1.7) \times 10^{27} \text{ cm}^{-2} \end{aligned}$$

D. Cross Sections

Tables Xa and Xb below show the result of this experiment. The errors are a combination of the statistical errors and an estimated 5%

uncertainty in the corrections and the classification of Editor rejections. The number of events comes from Table VII corrected for a scanning efficiency of 99%. The number of neutrals is the N_g of Chapter VI corrected for a scanning efficiency of 96.8%. Note that the number of neutrals does not correspond to the criteria used for events and that the events and neutrals are separately normalized.

Table Xa. Results of the Experiment

Type of event	No. of events	absolute cross section (mb.)
elastic	3960 ± 81	17.56 ± 0.43
π^0 production	1049 ± 34	4.65 ± 0.17
π^+ production	1609 ± 45	7.14 ± 0.23
$\pi^- \pi^+ \pi^0 n$	74 ± 9	0.33 ± 0.04
$\pi^- \pi^+ \pi^- p$	17 ± 4	0.08 ± 0.02
all charged events	6711 ± 118	$\sigma_{ch} = 29.76 \pm 0.69$
neutrals	1336 ± 44	$\sigma_{neut} = 11.78 \pm 0.43$
Total cross section		$\sigma_T = 41.54 \pm 0.82$

Our results for the total and neutral cross sections are compared to other experiments in Fig. 18a and 18b. Figures 19a-19c show our elastic, π^0 , and π^+ production cross sections. The elastic cross section has a peak at 600 MeV and the results of this experiment are in good agreement with the other measurements shown.

The π^+ production cross section (Fig. 19b) is in good agreement with the results from Maryland⁷ and Kentucky⁸ but all three measurements

Table Xb. Differential Cross Section

$\cos\theta^*$	Experimental (mb/ster)	5th order fit (mb/ster)
1	6.49 ± 0.25	6.36 ± 0.22
.85	4.16 ± 0.19	4.55 ± 0.10
.75	3.90 ± 0.18	3.62 ± 0.09
.65	3.09 ± 0.16	2.86 ± 0.07
.55	2.10 ± 0.13	2.22 ± 0.06
.45	1.63 ± 0.11	1.69 ± 0.05
.35	1.13 ± 0.09	1.24 ± 0.04
.25	1.04 ± 0.09	0.87 ± 0.03
.15	0.58 ± 0.07	0.58 ± 0.03
.05	0.39 ± 0.05	0.35 ± 0.02
-.05	0.18 ± 0.04	0.20 ± 0.02
-.15	0.12 ± 0.03	0.13 ± 0.02
-.25	0.16 ± 0.03	0.14 ± 0.02
-.35	0.21 ± 0.04	0.22 ± 0.02
-.45	0.40 ± 0.05	0.35 ± 0.02
-.55	0.50 ± 0.06	0.53 ± 0.03
-.65	0.72 ± 0.07	0.71 ± 0.04
-.75	0.81 ± 0.08	0.85 ± 0.04
-.85	0.89 ± 0.08	0.89 ± 0.04
-.95	0.76 ± 0.08	0.76 ± 0.07

are systematically higher than the results of Kirz.³¹ It is possible that the results of Kirz may be too low because he measured and analyzed only events which scanners classified as probably π^+ on the basis of bubble density. The discrepancy between the two sets of results makes it difficult to decide whether or not a peak is present in the cross section for π^+ production.

The π^0 production cross section (Fig. 19c) is also in close agreement with the Kentucky and Maryland results. Figure 19c shows bubble chamber results^{7,8,44} which may indicate a small peak in the cross section near 600 MeV. Also shown are results of a counter experiment by Detoeuf et al³² which measured $\pi^- + p \rightarrow \pi^0 + (\text{charged particles})$. This cross section is much higher than the bubble chamber results and the disagreement is very difficult to explain. Some of the discrepancy may be from the reactions $\pi^- + p \rightarrow \pi^- + \pi^+ + \pi^0 + n$ or $\pi^- + p \rightarrow \pi^- + p + \pi^0 + \pi^0$ but this is less than 0.5 mb at the energy of this experiment.

VIII. THEORY AND DISCUSSION

A. Partial Wave Analysis of Elastic Scattering

a. Theory.- We shall briefly review the usual treatment of elastic scattering in terms of the partial wave analysis. We consider first a beam of spinless particles scattered from a different spinless particle as seen in the center of mass system. For this case, it is well known that the elastic scattering amplitude is given by:

$$f(\theta^*) = \frac{1}{2ik} \sum_{\ell} (2\ell + 1) (\exp(2i\delta'_{\ell}) - 1) P_{\ell}(\cos\theta^*) \quad (9)$$

Here k is defined by $p = \hbar k$ and δ'_{ℓ} is a complex phase shift. We can write $\delta'_{\ell} = \delta_{\ell} - \frac{1}{2}i \ln \eta_{\ell}$ where δ and η_{ℓ} are real and $0 \leq \eta_{\ell} \leq 1$. We then have

$$f(\theta^*) = \frac{1}{2ik} \sum_{\ell} (2\ell + 1) (\eta_{\ell} \exp 2i\delta_{\ell} - 1) P_{\ell}(\cos\theta^*) \quad (10)$$

η_{ℓ} is called the elasticity parameter and can be given a physical meaning by noting that we can write

$$\sigma_{\text{elastic}} = \frac{\pi}{k^2} \sum_{\ell} (2\ell + 1) |1 - \eta_{\ell} e^{2i\delta_{\ell}}|^2 \quad (11a)$$

$$\sigma_{\text{inelastic}} = \frac{\pi}{k^2} \sum_{\ell} (2\ell + 1) (1 - \eta_{\ell}^2) \quad (11b)$$

$\eta_{\ell} = 1$ corresponds to completely elastic scattering and $\eta_{\ell} = 0$ corresponds to the maximum possible inelastic scattering.

$f(\theta^*)$ is given a physical meaning by the relation $\frac{d\sigma}{d\Omega^*} = |f(\theta^*)|^2$.

It follows from the properties of the $P_\ell(\cos\theta^*)$ that we can write

$$\frac{d\sigma}{d\Omega^*} = \sum_n a_n \cos^n \theta^* \quad \text{and that the maximum } n \text{ in the power series is}$$

given by $N_{\max} = 2\ell_{\max}$ where $\ell_{\max}$ is the highest value of ℓ which is effective.

We have to extend this to π^-p scattering where we have $J = \ell \pm \frac{1}{2}$.²³

Each value of ℓ other than 0 will have two angular momentum amplitudes

$$A_{\ell, J} = A_{\ell, \ell \pm \frac{1}{2}} \equiv A_{\ell \pm} \equiv (\eta_{\ell \pm} \exp[2i\delta_{\ell \pm}] - 1)/2i.$$

The $A_{\ell \pm}$ can be further decomposed into isotopic spin terms by writing

$$A_{\ell \pm} = \frac{1}{3} (A_{\ell \pm} [T = \frac{3}{2}] + 2A_{\ell \pm} [T = \frac{1}{2}])$$

Moreover, the fact that the proton can undergo spin flip adds terms $P_\ell^1(\cos\theta^*)\exp(i\phi)$ to the partial wave expansion. The non-spin flip and spin-flip amplitudes are respectively

$$f(\theta^*) = \frac{1}{k} \sum [(L+1)A_{L+} + LA_{L-}] P_L(\cos\theta^*) \quad (12a)$$

$$g(\theta^*, \phi) = \frac{1}{k} \sum (A_{L+} - A_{L-}) P_L^1(\cos\theta^*) \exp(i\phi) \quad (12b)$$

$$\text{and} \quad \frac{d\sigma}{d\Omega^*} = |f(\theta^*)|^2 + |g(\theta^*, \phi)|^2 \quad (13)$$

The associated Legendre polynomials $P_L^1(\cos\theta^*)$ can be written as

$\sin\theta^*$ (polynomial in $\cos\theta^*$). It follows from this that we can still write

$$\frac{d\sigma}{d\Omega^*} = \sum_n a_n \cos^n \theta^* \text{ where } N_{\max} = \text{Min}(2J_{\max}, 2\ell_{\max}).$$

It is customary to use this power series expansion rather than attempt a phase shift analysis because of the large number of coefficients needed in a phase shift analysis. The a_n are related to the angular momentum amplitudes by

$$a_n = \frac{1}{k^2} \text{Re} \sum_{i \leq j} b_{i j n} A_i A_j^*$$

where i or j stand for $L_{\pm}$ and the $b_{i j n}$ are given in Table XI below.

As will be shown below, we will need to know the differential cross section at zero degrees (forward scattering). This is given by

$$\left(\frac{d\sigma}{d\Omega^*}\right)_0 = |f(0)|^2 = |\text{Re}f(0)|^2 + |\text{Im}f(0)|^2.$$

The spin flip amplitude does not contribute since $P_{\ell}^1(\cos\theta = 1) = 0$. The optical theorem³³ relates the imaginary part of $f(0)$ to the total cross section: $\sigma_{\text{total}}(\pi^- p) = \frac{4\pi}{k} \text{Im}f(0)$. The real part of $f(0)$ has been calculated by Cronin³⁴ from dispersion relations. It is therefore possible to have a value for $\left(\frac{d\sigma}{d\Omega^*}\right)_0$ which is independent of the angular distribution observed in this experiment. This is useful since, as mentioned before, it is difficult to see events where the scattered pion emerges near the incident beam.

Table XI. Coefficients for $a_n = \sum_{ij} b_{ijn} A_i A_j^*$

		n						
i	j	0	1	2	3	4	5	6
0	0	1	--	--	--	--	--	--
0	1-	--	2	--	--	--	--	--
0	1+	--	4	--	--	--	--	--
0	2-	-2	--	6	--	--	--	--
0	2+	-3	--	9	--	--	--	--
0	3-	--	-9	--	15	--	--	--
0	3+	--	-12	--	20	--	--	--
1-	1-	1	--	--	--	--	--	--
1-	1+	-2	--	6	--	--	--	--
1-	2-	--	4	--	--	--	--	--
1-	2+	--	-9	--	15	--	--	--
1-	3-	-3	--	9	--	--	--	--
1-	3+	3	--	-30	--	35	--	--
1*	1+	1	--	3	--	--	--	--
1+	2-	--	-10	--	18	--	--	--
1+	2+	--	--	--	12	--	--	--
1+	3-	3	--	-36	--	45	--	--
1+	3+	-3	--	-6	--	25	--	--
2-	2-	1	--	3	--	--	--	--
2-	2+	3	--	-36	--	45	--	--
2-	3-	--	--	--	12	--	--	--
2-	3+	--	21	--	-110	--	105	--
2+	3+	9/4	--	-9/2	--	45/4	--	--
2+	3-	--	45/2	--	-117	--	225/2	--
2+	3+	--	9	--	-30	--	45	--
3-	3-	9/4	--	-9/2	--	45/4	--	--
3-	3+	-9/2	--	207/2	--	-675/2	--	525/2
3+	3+	9/4	--	45/4	--	-165/4	--	175/4

b. Small angle correction.- In order to use the optical theorem, it is first necessary to determine the total cross section and this requires determining the small angle correction. This was done by fitting the number of events to a polynomial of the form $n_i = \sum_j a_j \cos^j(x_i)$ where n_i is the number of events in the i^{th} bin and x_i is the midpoint of the bin. The fitting was done by the least square method discussed by Orear³⁵ which consists of minimizing the quantity $\chi^2 = \sum_i \left(\frac{n_i - \sum_m a_m \cos^m(x_i)}{\delta n_i} \right)^2$ where δn_i is the error in the number of events in the i^{th} bin. The bin $0.9 < \cos\theta^*$ was omitted from the fit because we were attempting to determine the small angle correction to this bin. The remaining 19 points were fitted to 1st through 7th order curves. The first satisfactory fit was a 4th order curve and the best fit was a 5th order curve. Table IX lists 572 events in the bin $0.9 < \cos\theta^*$. The 4th order curve predicted 700 events in this bin and the 5th order curve gave 750 events. The number was finally taken as 725 events so the small angle correction was 153 events.

c. Fit to differential cross-section.- There are two methods of determining the coefficients a_j in the relation $\frac{d\sigma}{d\Omega^*} = \sum_j a_j \cos^j\theta^*$ after the small angle correction has been made and the optical theorem used to evaluate $\left. \frac{d\sigma}{d\Omega^*} \right|_{\theta^*=0}$. The first of these is a "lumped" method in which we write

$$n_i = 2\pi nL'' \int_{x_i-\Delta}^{x_i+\Delta} \frac{d\sigma}{d\Omega^*} dx \quad \text{where } x \text{ is } \cos\theta^* \text{ and get}$$

$$n_i = 2\pi nL'' \left. \frac{d\sigma}{d\Omega^*} \right|_{x_i} x 2\Delta. \quad \text{This yields a value for } \left. \frac{d\sigma}{d\Omega^*} \right|_{x_i} \quad \text{which represents}$$

the average in the range of $\cos\theta^*$ between $x_i - \Delta$ and $x_i + \Delta$ and we then write

$\left. \frac{d\sigma}{d\Omega^*} \right|_{x_i} = \sum_j a_j x_i^j$. This method is simple but has the disadvantage that the value of $\left. \frac{d\sigma}{d\Omega^*} \right|_{x_i}$ may not be the true value at x_i and will depend on the bin size. The second method, which we will call the "integral" method, uses

$$n_i = 2\pi nL'' \int_{x_i - \Delta}^{x_i + \Delta} \frac{d\sigma}{d\Omega^*} dx = 2\pi nL'' \int_{x_i - \Delta}^{x_i + \Delta} \sum_j a_j x^j dx$$

$$= 2\pi nL'' \sum_j \frac{a_j}{j+1} \left([x_i + \Delta]^{j+1} - [x_i - \Delta]^{j+1} \right)$$

and does not require evaluating $\frac{d\sigma}{d\Omega^*}$ at x_i . The term nL'' is the same as that used in Chapter VII on cross-sections.

The values of the a_n were determined by the same least square method used to determine the small angle correction. For the lumped method, the quantity minimized was

$$\chi^2_L = \sum_i \left(\frac{\left. \frac{d\sigma}{d\Omega^*} \right|_{x_i} - \sum_j a_j x_i^j}{\delta_i} \right)^2 + \left(\frac{\left. \frac{d\sigma}{d\Omega^*} \right|_{x_o=1} - \sum_j a_j}{\delta_o} \right)^2 \quad (14)$$

The quantity minimized for the integral method was

$$\chi^2_I = \sum_i \left(\frac{\frac{n_i}{2\pi nL''} - \sum_j \frac{a_j}{j+1} \left[(x_i + \Delta)^{j+1} - (x_i - \Delta)^{j+1} \right]}{\delta_i} \right)^2 + \left(\frac{\left. \frac{d\sigma}{d\Omega^*} \right|_{x_o=1} - \sum_j a_j}{\delta_o} \right)^2 \quad (15)$$

In both these equations $\left. \frac{d\sigma}{d\Omega^*} \right|_{x_i=0}$ is the differential cross section at 0° evaluated from the optical theorem and dispersion relations and δ_i represents the error in $\left. \frac{d\sigma}{d\Omega^*} \right|_{x_i}$ or $\frac{n_i}{2\pi nL''}$.

The minimum value of $n_{\max}$ needed to provide a good fit to the experimental results with

$$\frac{d\sigma}{d\Omega^*} = \sum_{n=0}^{n_{\max}} a_n \cos^n \theta^*$$

can be determined by plotting the confidence level for the fit vs the order of the fit. It has become customary to use instead a "goodness of fit" parameter defined as $\left(\chi^2/d \right)^{1/2}$ where d is the number of degrees of freedom available for the fit. This is based on the fact that the average χ^2 equals the number of degrees of freedom for a good fit. Thus, the "goodness of fit" parameter should approach 1 as the fit improves.

Both the "lumped" and "integral" methods were used to make fits to 1st through 7th order curves of $\frac{d\sigma}{d\Omega^*}$. In both cases, the bin $\cos\theta^* > 0.9$ was omitted and the fits were made to the remaining 19 points and the optical theorem point at $\theta^* = 0^\circ$. It was found that both methods gave nearly identical χ^2 and that the difference between the coefficients determined by the two methods were well within the errors. Figure 20 shows the differential cross section and the goodness of fit vs n . The first curve which gives a satisfactory fit when the optical theorem is used is the 5th order. The 6th and 7th order also give good fits but the 7th order can be eliminated because it gives $\left. \frac{d\sigma}{d\Omega^*} \right|_{\theta^*=0} = 4.47 \text{ mb/ster}$

when the fit is made without using the optical theorem. This is far below the expected value of 6.49 mb/ster. The choice between 5th and 6th order is more difficult because both have satisfactory confidence levels. However, when the fit is made without using the optical theorem point at 0° , the 5th order curve gives $\left. \frac{d\sigma}{d\Omega^*} \right|_{\theta=0} = 5.85 \pm 0.50$ mb/ster whereas the 6th order gives 5.45 ± 0.80 . Thus, the 5th order curve gives better agreement with the calculated value. Moreover, when the fitting is done with the forward scattering point included, the 6th order curve gives $a_6 = 0.48 \pm 1.03$ which is consistent with 0. One can thus conclude that the 5th order curve is the lowest order which will give a good fit and that higher orders are not essential.

Table XII shows the coefficients for the 5th order curve determined by both the "lumped" and "integral" methods. This table also shows the preliminary results of a recent Berkeley¹⁸ counter measurement at 650 MeV. The results of the two experiments appear to be in satisfactory agreement.

Table XII. Coefficients for 5th Order Curve

	Lumped Method	Integral Method	Berkeley
a_0	0.27 ± 0.02	0.27 ± 0.02	0.27 ± 0.03
a_1	1.48 ± 0.11	1.48 ± 0.11	1.37 ± 0.12
a_2	3.86 ± 0.22	3.86 ± 0.22	3.91 ± 0.26
a_3	-0.29 ± 0.53	-0.30 ± 0.54	0.33 ± 0.47
a_4	-0.65 ± 0.28	-0.64 ± 0.28	-1.06 ± 0.47
a_5	1.69 ± 0.52	1.70 ± 0.52	0.85 ± 0.61

The coefficients for all fits attempted are shown in Table XIII.

Table XIII. Coefficients a_n for all fits

order coeff	1	2	3	4	5	6	7
a_0	0.61 ± 0.02	0.37 ± 0.02	0.26 ± 0.02	0.24 ± 0.02	0.27 ± 0.02	0.27 ± 0.02	0.27 ± 0.02
a_1	0.75 ± 0.04	1.78 ± 0.05	1.10 ± 0.08	1.23 ± 0.08	1.48 ± 0.11	1.50 ± 0.12	1.52 ± 0.16
a_2		2.95 ± 0.07	3.47 ± 0.08	4.17 ± 0.20	3.86 ± 0.22	4.01 ± 0.40	3.99 ± 0.44
a_3			1.64 ± 0.14	1.37 ± 0.15	-0.29 ± 0.53	-0.39 ± 0.57	-0.62 ± 1.39
a_4				-1.01 ± 0.26	-0.65 ± 0.28	-1.24 ± 1.31	-1.16 ± 1.39
a_5					1.69 ± 0.52	1.77 ± 0.55	2.37 ± 3.29
a_6						0.48 ± 1.03	0.43 ± 1.07
a_7							-0.40 ± 2.16

The fact that a 5th order curve is needed indicates that partial waves up to $F_{5/2}$ are involved in the scattering. Moreover, the 5th order term must be produced by interference between $D_{5/2}$ and $F_{5/2}$ waves because a $D_{5/2}$ wave alone would give terms no higher than 4th order. Any $F_{7/2}$ wave which may be present must have a very small effect since otherwise we would expect to need a 6th order curve. These results are consistent with other workers who have found that 5th order is sufficient up to at least 900 MeV.¹⁷⁻²¹

Figure 21 shows the coefficients as a function of energy. There is some evidence for a peak in a_2 at 600 MeV but it is not pronounced. The fact that the peak in elastic scattering evident in Figure 19a does

not appear as a large peak in one of the a_n need not be surprising.

Table XI relating the a_n to the partial waves shows that the effects are likely to be spread out through many of the coefficients.

The shape of the differential cross section (Fig. 20) is also in agreement with the results of other experiments. The backward hump is known¹⁹ to become more pronounced between 720 and 1020 MeV and to disappear by 1300 MeV. The results at higher energies (720-1020 MeV) indicate that the cross section is essentially 0 at $\theta^* = 180^\circ$. The peak in the forward direction is typical of diffraction scattering but diffraction scattering would predict a flat tail rather than a hump in the backward hemisphere. The hump is usually interpreted as due to the spin-flip amplitude $\langle g(\theta^*, \phi) \rangle$ since $|g(\theta^*, \phi)|^2 = \sin^2 \theta^* \times (\text{polynomial in } \cos \theta^*)$ goes to 0 at $\theta^* = 180^\circ$ and can give a hump in the backward hemisphere.

B. Inelastic Scattering

a. Theory.- One of the difficulties in interpreting inelastic scattering is that there are a large number of variables which can be used and one must decide which variables are the important ones. In recent years, it has been common to study the production of "resonance" particles, i.e. short-lived ($\sim 10^{-23}$ sec.) particles which decay into two or more longer lived particles. For example, the reaction $\pi + N \rightarrow \pi_1 + \pi_2 + N$ might actually proceed through the mechanism $\pi + N \rightarrow \pi_2 + N^*$ accompanied by the subsequent decay of $N^* \rightarrow \pi_1 + N$. The "particle" N^* is thought of as a resonance between π_1 and N . If this were the mechanism, then the initial step is really a two body problem and one would expect N^* to have a definite mass. One looks for a resonance between two particles by studying the variable $M_{12}^2 = (E_1 + E_2)^2 - (\vec{p}_1 + \vec{p}_2)^2$. M_{12} is Lorentz-invariant and has the physical significance of being the mass of the hypothetical resonance. If the two particles actually form a resonance, then one would expect to find an enhanced number of events in a region around a definite value of the invariant mass. Due to the fact that the resonance is unstable, its mass is not well defined and this will be reflected in the spread of events in the region of enhancement. The width of the peak will be governed by the uncertainty relation: $\Delta m \Delta t \sim \hbar$ where Δt is the mean life of the resonance and Δm is the spread in the invariant mass. There are also quantum mechanical interference effects which have a tendency to wash out a possible resonance and so make identification difficult. A number of models have been introduced to aid the study of inelastic scattering. Of these, the simplest is phase space or the statistical model.

1. The Statistical Model

Phase space is essentially the statistical model introduced by Fermi.³⁶ It assumes that in the formula for transition probability, $\omega = \frac{2\pi}{\hbar} \frac{d\rho}{dE} |H_{if}|^2$, the matrix element is a constant independent of the momenta and energies of the outgoing particles. Then the phase space factor $\frac{d\rho}{dE}$ gives the distribution of whatever variables are being studied.

The statistical model is useful as a null hypothesis because any deviation from it gives information about the matrix element. Figure 22 shows the effective mass plots for this experiment. The solid line shows the phase space distribution calculated from the formulae given by Bloch³⁷ (non-Lorentz invariant 3 body phase space). It can be seen that all of the $M_{\pi N}$ plots show large deviations from phase spaced centered around 1230 MeV. These deviations can be interpreted as examples of the N_{33}^* isobar which has been observed by many experimenters.

2. The Isobar Model

The isobar model was first introduced by Lindenbaum and Sternheimer³⁸⁻⁴¹ who pointed out that the (3,3) resonance at 190 MeV in the lab (1230 MeV in the center of mass system) could be interpreted as an excited isobaric state of the nucleon. They suggested that single pion production could be considered as proceeding through this isobar as an intermediate state. Their model is based on the diagram below.

They computed the effect of this intermediate state on the assumption that the cross section for production of an isobar with $J = I \neq 3/2$ and mass m_1 was equal to the $\pi^+ + p$ total cross section at center of mass energy m_1 . That is, they wrote $\sigma_{\pi^+ + p}(m_1) = \text{const} \times |M_{\text{isobar}}|_{m_1}^2 \times \rho(m_1) = \sigma_I(m_1)$ where $|M_{\text{isobar}}|_{m_1}$ is the matrix element for formation of an isobar of mass m_1 and $\rho(m_1)$ is the phase space factor. They also assumed that the production and decay of the isobar was isotropic.

Bergia, Bonsignori and Stanghellini⁴² extended the isobar model to include interference between the two diagrams below. They pointed out

that we must treat these diagrams as coherent because the isobar has a finite lifetime and we do not observe it directly. Bergia et al used a modified form of the Breit-Wigner resonance formula⁴⁷ to represent the matrix element instead of calculating it from the $\pi^+ + p$ cross section. However, they kept the assumption of isotropic production and decay.

The most recent and complete version of the isobar model is that of Olsson and Yodh.⁴³ These authors took into account

1. The interference terms
2. P-wave decay of the isobar
3. S-wave production of the isobar
4. Requirements of Bose statistics

The results of the calculations for the reactions

$$\pi^+ + p \rightarrow \pi^+ + \pi^+ + n$$

$$\pi^+ + p \rightarrow \pi^+ + \pi^0 + p$$

$$\pi^- + p \rightarrow \pi^- + \pi^+ + n$$

$$\pi^- + p \rightarrow \pi^- + \pi^0 + p$$

are

$$M(\pi^+, \pi^+, n) = -\frac{a_3 \vec{\sigma} \cdot \hat{q}}{\sqrt{60}} \left[\frac{s(\hat{p}_2')}{D(2)} + \frac{s(\hat{p}_1')}{D(2)} \right] \quad (16a)$$

$$M(\pi^+, \pi^0, p) = \frac{a_3 \vec{\sigma} \cdot \hat{q}}{\sqrt{15}} \left[3/2 \frac{s(\hat{p}_1')}{D(1)} - \frac{s(\hat{p}_2')}{D(2)} \right] \quad (16b)$$

$$M(\pi^-, \pi^+, n) = \frac{a_3 \vec{\sigma} \cdot \hat{q}}{\sqrt{30}} \left[2/3 \frac{s(\hat{p}_2')}{D(2)} - \frac{s(\hat{p}_1')}{D(1)} \right] - \frac{a_1 \vec{\sigma} \cdot \hat{q}}{2\sqrt{3}} \left[1/3 \frac{s(\hat{p}_2')}{D(2)} + \frac{s(\hat{p}_1')}{D(1)} \right] \quad (16c)$$

$$M(\pi^-, \pi^0, p) = \frac{a_3 \vec{\sigma} \cdot \hat{q}}{3\sqrt{15}} \left[2 \frac{s(\hat{p}_2')}{D(2)} - 1/2 \frac{s(\hat{p}_1')}{D(1)} \right] + \frac{a_1 \vec{\sigma} \cdot \hat{q}}{3\sqrt{6}} \left[\frac{s(\hat{p}_1')}{D(1)} - \frac{s(\hat{p}_2')}{D(2)} \right] \quad (16d)$$

The formulae are derived in Appendix II.

The factors $\frac{a_3 \vec{\sigma} \cdot \hat{q}}{D(1)}$ denotes S wave production of an isobar formed by pion No. 1 and the nucleon from the $T = 3/2$ state of the initial pion-nucleon system. The terms in a_1 for the $\pi^- p$ system represent isobar formation from the $T = 1/2$ state of the initial system.

$D(1)$ represents the resonance denominator for the $(3,3)$ resonance and is given by $\frac{1}{D(1)} = \frac{(\Gamma(1)/q)^{1/2}}{W(1N) - W_R + i\Gamma(1)}$ where $\Gamma(1)$ is the width, W_R

the (3,3) resonance energy, and $W(1N)$ the energy of pion No. 1 and the nucleon in their center of mass. The incoming pion direction in the overall center of mass is given by the unit vector $\hat{q}$. The P-wave decay of the isobar is given by the function $s(\hat{p}_1') = (2\hat{q} \cdot \hat{p}_1' - i\vec{\sigma} \cdot (\hat{q} \times \hat{p}_1'))$ where $\hat{p}_1'$ is the direction of motion of pion no. 1 in the isobar rest frame, $\vec{\sigma}$ is the nucleon spin, and the isobar is formed from pion no. 1 and the nucleon. The only undetermined parameters are the magnitudes of a_1 and a_3 and their relative phase. These authors claim that their model accounts for the ratio of $\sigma(\pi^+ + p \rightarrow \pi^+ + \pi^+ + n)$ to $\sigma(\pi^+ + p \rightarrow \pi^0 + \pi^+ + p)$ and also accounts for all the effective mass plots except $M_{\pi\pi}^-$ from the reaction $\pi^- + p \rightarrow \pi^- + \pi^+ + n$.

b. Single pion exchange.- Another model often considered is the single pion exchange model diagrammed below.

This was not considered in this experiment because the mass plots showed that isobar production dominated the inelastic scattering. Moreover, Vittitoe⁸ at 600 MeV and Pickup et al⁴⁴ at 905 and 960 MeV also found that isobar formation dominated the reactions.

c. Results for effective two body mass.- The interpretation of the $M_{\pi N}$ plots (Fig. 22) in this experiment is complicated by the kinematics. The reaction $\pi^- + p \rightarrow \pi_1 + N_{\pi_2 N}^*$ is a two body reaction with the momenta of the two outgoing particles determined by the total available energy and M_{N^*} . The decay of $N_{\pi_2 N}^*$ into π_2 and N is also a two body reaction

with completely determined momenta. Therefore, when we calculate the effective mass of pion no. 1 and the nucleon, ($M_{\pi_1 N}$), it will be limited to a fairly narrow range. A specific calculation for this experiment using $M_{\pi_2 N} = 1240$ MeV showed that $M_{\pi_1 N}$ would be between 1200 and 1300 MeV. Since this is the range in which the isobar is found, we cannot tell whether we are looking at $\pi^- + p \rightarrow \pi_1 + N^*_{\pi_2 N}$ or $\pi^- + p \rightarrow \pi_2 + N^*_{\pi_1 N}$ simply by looking at the plots of this experiment.

Kirz³¹ has made a study of the reaction $\pi^- + p \rightarrow \pi^- + \pi^+ + n$ in the energy range 360-780 MeV. He finds that the peak in $M_{\pi^- n}$ stays centered about 1240 MeV whereas the peak in $M_{\pi^+ n}$ shifts position as the energy of the incident beam is changed. Therefore the peak in $M_{\pi^- n}$ probably represents a true isobar and the $M_{\pi^+ n}$ peak is a kinematic effect. This is consistent with the Olsson-Yodh model because $N^*_{\pi^- n}$ is a pure $T = 3/2$ state whereas $N^*_{\pi^+ n}$ is a mixture of $T = 3/2$ and $T = 1/2$. Consequently, the Olsson-Yodh model predicts that $N^*_{\pi^- n}$ should be much more common than $N^*_{\pi^+ n}$.

Unfortunately, there is not enough experimental data to allow us to decide whether both isobars are present in the reaction $\pi + p \rightarrow \pi^- + \pi^0 + p$ or whether one of the peaks is a kinematic effect. If we assume that $|a_1| > |a_3|$ because of the peak in the $T = 1/2$ cross section, the Yodh model predicts that $N^*_{\pi^- p}$ and $N^*_{\pi^0 p}$ should be present in equal amounts.

The $M_{\pi\pi}$ plots also have a number of points of interest. The peaking at the high mass end of $M_{\pi^- \pi^0}$ is predicted by the Olsson-Yodh model because the matrix element for $\pi^- + p \rightarrow \pi^- + \pi^0 + p$ is anti-symmetric in the two isobars. This means that it will be largest when the π^- and π^0 go away from each other. $M_{\pi\pi}$ is defined by $(M_{\pi\pi})^2 = (E_{\pi_1} + E_{\pi_2})^2 - (\vec{p}_{\pi_1} + \vec{p}_{\pi_2})^2$. If the pions are going away from each other, the second

term will be small and the $M_{\pi\pi}$ will be near its maximum.

On the other hand, the peaking at the high mass end of the $M_{\pi^-\pi^+}$ plot is not predicted by the Olsson-Yodh model. Kirz has found that this enhancement is present at other incident beam energies but that it shifts so that it always remains at the high end of mass spectrum. Therefore, one cannot interpret it as a π - π resonance. Olsson and Yodh have suggested that one might be able to explain the enhancement in terms of a $T = 0$, $J = 0$ $\pi\pi$ interaction. If the explanation is that there is a $\pi\pi$ interaction, then it must be $T = 0$ because the lack of any effect in $\pi^-\pi^0$ or $\pi^+\pi^+$ rules out $T = 1$ and $T = 2$.

d. Angular distributions. - The angular distributions in the overall center of mass are shown in Fig. 22. It will be observed that the angular distributions for π^+ and π^0 production are qualitatively different. In particular, the π^- and π^+ in π^+ production show a strong forward-backward asymmetry not present in the π^0 production plots. This asymmetry is consistent with the peaking in $M_{\pi^-\pi^+}$ because the pions going away from each other will cause $M_{\pi^-\pi^+}$ to be large as previously explained. Kirz has found that the asymmetry reverses below 600 MeV, with the π^+ going forward and the π^- going backwards.

The observed asymmetry does not seem to be consistent with the Olsson-Yodh assumption of S wave production. Fig. 24 shows the center of mass angular distribution of the π^+ for three ranges of the effective π^- -neutron mass. If the reaction occurred via the intermediate reaction $\pi^- + p \rightarrow \pi^+ + N^*$ followed by $N^* \rightarrow \pi^- + n$ and the isobar was produced in an S state relative to the π^+ , then the π^+ distribution should be isotropic. Moreover, the isobar will go opposite from the π^+ by conservation of

momentum. As shown by Fig. 24, the result is strongly dependent on M_{π^-N} . In particular, for $1190 \text{ MeV} < M_{\pi^-n} < 1290 \text{ MeV}$, the ratio $\frac{F - B}{F + B} = 0.17 \pm 0.04$ where F is the number of isobars with $\cos\theta^* > 0$ and B is the number of isobars with $\cos\theta^* < 0$.

e. Multiple pion production.- There was a special class of events which have the following characteristics:

1. Either no fit was made by Guts or only a fit to production was made and this fit was very poor.
2. The momenta of both outgoing tracks were low.
 $(|\vec{p}_1| + |\vec{p}_2|) \leq 400 \text{ MeV/c}.$
3. The missing mass for π^+ production was greater than 1050 MeV.
4. Both tracks were heavily ionizing.

These events cannot be Dalitz pairs because the bubble density was too great and they cannot have a proton for the positive track because protons with such low momenta would have a range less than the length of the tracks.

The most reasonable explanation is the reaction $\pi^- + p \rightarrow \pi^- + \pi^+ + n + \pi^0$ or $\pi^- + \pi^+ + n + \gamma$. The missing mass for this reaction is given by

$$M^2 = m_1^2 + m_2^2 + 2E_1E_2(1 - \beta_1\beta_2 \cos\theta_{12})$$

where 1 and 2 are the two neutral particles and θ_{12} is the angle between them. The missing mass cannot be less than $m_1 + m_2$ and the range of mass is limited by the fact that only 200 MeV is available to be shared among the four particles in the overall center of mass system. The kinematics of the reaction show that both the π^+ and π^- will have low momenta in the laboratory system. Thus this reaction would have the

desired characteristics.

It was stated above that the missing mass cannot be less than $m_1 + m_2$. This lower limit is 1075 MeV if m_1 and m_2 are a π^0 and neutron. The 74 reported events of this type actually include 10 events with missing mass between 1040 and 1075 MeV and χ^2 for π^+ production greater than 45. These were accepted because the χ^2 was so large and also because one would expect measuring errors to give some missing masses less than 1075 MeV. It is possible that some of these events may be the tail end of the π^+ production events. However, a study of the χ^2 and stretch distribution for π^+ events with one or both pions having momenta less than 200 MeV/c did not show any indication of systematic error connected with low momenta. Thus it is more reasonable to connect these 10 events to $\pi^+ + \pi^- + \pi^0 + n$ rather than to $\pi^+ + \pi^- + n$.

Kirz³¹ has found the same type of events at other energies and as shown in Table XIV there is a sudden increase in cross section at about 550 MeV. Our cross section is in good agreement with his values at 600 and 673 MeV. Pickup et al⁴⁴ has found a cross section of 0.8 mb at 905 MeV.

A search for the reaction $\pi^- + p \rightarrow \pi^- + p + \pi^0 + \pi^0$ was unsuccessful. The missing mass for such events would have to be greater than 270 MeV and one would expect $\chi^2_{\pi^0}$ to be large. Only 2 or 3 events met these requirements. Pickup et al report that at 905 MeV the cross section for $\pi^- + p \rightarrow \pi^- + p + (2 \text{ or more neutrals})$ is 0.2 mb which is a factor of 4 lower than the cross section for $\pi^- + p \rightarrow \pi^+ + \pi + (2 \text{ or more neutrals})$. This is consistent with our failure to find such events.

Finally, the reaction $\pi^- + p \rightarrow \pi^- + p \rightarrow \pi^- + \pi^+$ is easy to locate and we have less than 17 events which are known to be of this type.

Table XIV. Cross sections for $\pi^- + p \rightarrow \pi^- + \pi^+ + n + x^0$, where x^0 is π^0 or energetic γ ray.*

T_{beam} (MeV)	p_{beam} (MeV/c)	$\pi^- \pi^+ n x^0 / \pi^- \pi^+ n$	$\sigma_{\pi^- \pi^+ n x^0}$ (mb)
360	480	0.000 ± 0.002	0.000 ± 0.004
430	552	0.004 ± 0.002	0.015 ± 0.008
460	583	0.005 ± 0.002	0.020 ± 0.008
480	604	0.002 ± 0.002	0.010 ± 0.010
555	680	0.018 ± 0.007	0.10 ± 0.04
605	731	0.080 ± 0.014	0.49 ± 0.09
646	772	0.046 ± 0.006	0.33 ± 0.04
673	800	0.055 ± 0.012	0.34 ± 0.08
780	909	0.061 ± 0.012	_____

*All results are from Kirz except this experiment.

The question of why the reaction $\pi^- + \pi^+ + \pi^0 + n$ should be so much more common than the others is difficult to answer. Kirz has pointed out that the jump in cross section occurs near the threshold for n production and suggests that the reaction may be $\pi^- + p \rightarrow n + \eta$ followed by $\eta \rightarrow \pi^+ + \pi^0 + \pi^-$. The suggestion is supported by some recent spark chamber experiments^{45,46} which indicate that the cross section for η production is several millibarns at this energy. This would account for the fact that the cross section for $\pi^- + p \rightarrow \pi^- + \pi^+ + \pi^0 + n$ is

so much larger than the other multiple pion reaction cross sections. Unfortunately, it is not possible to test this the hypothesis because we cannot calculate $M_{\pi^+\pi^-\pi^0}$ from our data.

IX. CONCLUSION

The elastic, inelastic, and neutral production cross sections for negative pions on protons at 646 MeV were studied in this experiment. The results are in good agreement with other experiments. It was found that the differential cross section for elastic scattering could be fitted to a 5th order polynomial in $\cos\theta^*$. This showed that waves up to at least $F_{5/2}$ are present and that the effects of any higher partial waves are small.

The reactions $\pi^- + p \rightarrow \pi^- + \pi^+ + n$ is dominated by the formation of the $N_{3,3}^*$ isobar in its negative charge states. The Olsson-Yodh model accounts for the shape of the M_{π^-n} and M_{π^+n} effective mass plots but fails to account for the $M_{\pi^-\pi^+}$ plot. Moreover, the angular distributions of the reaction products in the center of mass are in poor agreement with the Olsson-Yodh model. In particular, the distribution of the $N_{3,3}^{*-}$ in the center of mass does not appear to be consistent with the Olsson-Yodh assumption of s-wave production.

The reaction $\pi^- + p \rightarrow \pi^- + \pi^0 + p$ also appears to be dominated by isobar production but the effect is not as strong as for the π^+ production reaction. This is due to fact that $N^*(\pi^-n)$ is a pure $I = 3/2$ state whereas the isobars involved in π^0 production are mixtures of $I = 1/2$ and $I = 3/2$. It is not clear whether both the $N^*(\pi^0p)$ and $N^*(\pi^-p)$ are present or whether only one is present. The Olsson-Yodh model gives a good fit to the effective mass plots for π^0 production but again fails to predict the angular distributions.

The cross section for $\pi^- + p \rightarrow \pi^- + \pi^+ + \pi^0 + n$ appears to be at least a factor of four greater than the reactions $\pi^- + p \rightarrow \pi^- + \pi^+ + \pi^- + p$

or $\pi^- + p + \pi^0 + \pi^0$. This difference may be due to η production, but it was not possible to test this idea.

X. ACKNOWLEDGEMENTS

I wish to thank the many people whose unfailing help and cooperation made this experiment possible. I especially wish to thank Dr. I. Nadelhaft for his constant advice and guidance, Drs. J. Ashkin and G. B. Yodh for planning the bubble chamber run, and the staff of Brookhaven National Laboratory for operating the chamber and providing Guts and Datpro.

I must also acknowledge the help of Mr. J. Crump, Mrs. J. Fierst, and Miss M. E. Lipchak who helped with the computer programs, and the staffs of the New York University Computation Center and the Gulf Oil Research and Development Center who provided time on their 7090's and helped with the running of the programs.

Finally, I must thank the following scanners who worked under the direction of Mrs. P. Simpson: M. Godich, A. Fine, D. Stasak, L. Stember, P. Bizub, H. Scheller, S. Engles, M. Lentz, J. Griffith, and L. McGraw.

APPENDIX I

Derivation of Error Formulas

In the following derivations, we treat all variables as independent and use the general formula:

$$Q = f(a, b)$$

$$(\delta Q)^2 = \left(\frac{\partial f}{\partial a}\right)^2 (\delta a)^2 + \left(\frac{\partial f}{\partial b}\right)^2 (\delta b)^2$$

1. The error in dip

We define $\sin(\text{dip}) = \frac{z_2 - z_1}{L}$ where z_1 and z_2 are the z coordinates of two points a distance L apart. Write $z_2 - z_1 = z$ and

$$\sin(\text{dip}) = \frac{z}{L} \quad (1)$$

$$\cos^2 \text{dip} (\Delta \text{dip})^2 = \frac{(\Delta z)^2}{L^2} + \frac{z^2}{L^2} \left(\frac{\Delta L}{L}\right)^2 \quad (2)$$

or

$$(\Delta \text{dip})^2 = \left(\frac{\Delta z}{L \cos \text{dip}}\right)^2 + \frac{z^2}{L^2} \left(\frac{\Delta L}{L \cos \text{dip}}\right)^2 \quad (3)$$

We drop the second term because the additional L^2 term makes it small.

This leaves us with

$$(\Delta(\text{dip}))^2 = \left(\frac{\Delta z}{L \cos(\text{dip})}\right)^2 \quad (4)$$

which is present for all dip and increases as $\text{dip} \rightarrow 90^\circ$. This is a physically reasonable behavior so we expect the error in dip to behave like (4). The term Δz comes from the measuring error in z .

2. The error in azimuth

We write $\tan az = \frac{y_2 - y_1}{x_2 - x_1} = \frac{y}{x}$ and treat x and y as independent variables. Then

$$\sec^4(az) (\Delta az)^2 = \tan^2 az \left[\left(\frac{\Delta y}{y} \right)^2 + \left(\frac{\Delta x}{x} \right)^2 \right] \quad (5)$$

We now write $x = L_p \cos az$ and $y = L_p \sin az$ where L_p is the projected length, i.e., $L_p = L \cos \text{dip}$.

$$\begin{aligned} (\Delta x)^2 &= \cos^2 az (\Delta L_p)^2 + L_p^2 \sin^2 az (\Delta az)^2 \\ (\Delta y)^2 &= \sin^2 az (\Delta L_p)^2 + L_p^2 \cos^2 az (\Delta az)^2 \\ \sec^4(az) (\Delta az)^2 &= \tan^2 az \left[2 \left(\frac{\Delta L_p}{L_p} \right)^2 + \left(\frac{1}{\tan^2 az} + \tan^2 az \right) (\Delta az)^2 \right] \\ \left[\frac{1}{\sin^2 az \cos^2 az} - \frac{\cos^2 az}{\sin^2 az} - \frac{\sin^2 az}{\cos^2 az} \right] (\Delta az)^2 &= 2 \left(\frac{\Delta L_p}{L_p} \right)^2 \end{aligned} \quad (6)$$

The left hand side of (6) reduces to $2(\Delta az)^2$ after some algebra so we have

$$(\Delta az)^2 = \left(\frac{\Delta L_p}{L_p} \right)^2 = \left(\frac{\Delta L_p}{L \cos \text{dip}} \right)^2 \quad (7)$$

3. The error in momentum

We use $P = \frac{HR}{\cos \text{dip}}$

$$(\Delta P)^2 = P^2 \left(\frac{\Delta R}{R} \right)^2 + P^2 \tan^2 \text{dip} (\Delta \text{dip})^2 \quad (8)$$

Now R is defined by $R = \frac{L_p^2}{8S}$

$$\frac{(\Delta R)^2}{R^2} = 4 \left(\frac{\Delta L_p}{L_p} \right)^2 + \left(\frac{\Delta S}{S} \right)^2 \quad (9)$$

But $1/S = \frac{8R}{L_p^2}$

Therefore (9) becomes

$$\left(\frac{\Delta R}{R} \right)^2 = 4 \left(\frac{\Delta L_p}{L_p} \right)^2 + \frac{64R^2}{L_p^4} (\Delta S)^2 \quad (10)$$

and we have

$$\left(\frac{\Delta P}{P} \right)^2 = 4 \left(\frac{\Delta L_p}{L_p} \right)^2 + \frac{64R^2}{L_p^4} (\Delta S)^2 + \tan^2 \text{dip} (\Delta \text{dip})^2 \quad (11)$$

The last term vanishes for small dip and the first term is expected to be smaller than the second. Therefore we use only the middle term and write

$$\left(\frac{\Delta P}{P} \right)^2 = \frac{64 (\Delta S)^2}{(L \cos \text{dip})^4} \quad (12)$$

The experimental and theoretical formulas are given below for comparison.

	Theoretical	Experimental
$(\Delta az)^2$	$= \frac{(\Delta L_p)^2}{(Lcosdip)^2}$	$\frac{11.6}{(Lcosdip)} \quad 1.79$
$(\Delta dip)^2$	$= \frac{(\Delta z)^2}{(Lcosdip)^2}$	$\frac{1360}{(Lcosdip)} \quad 1.93$
$\frac{\Delta P}{PR}^2$	$= \frac{64(\Delta S)^2}{(Lcosdip)^4}$	$\frac{167}{(Lcosdip)} \quad 4.03$

The agreement is good except for Δaz . This probably reflects the fact that x and y are not truly independent variables.

APPENDIX II

The Olsson-Yodh Isobar Model

We make the following assumptions:

1. Single pion production occurs only via the production of an isobar and a pion ($\pi + p \rightarrow N^* + \pi$)
2. The N^* and π are produced in a relative S state
3. The N^* decays as a $P_{3/2}$ state in its own center of mass system
4. The amplitudes do not depend on T_3 of the isobar, i.e. the amplitudes are charge independent.

We write $|\pi^+ p\rangle = |3/2, 3/2\rangle$

$$|\pi^- p\rangle = \sqrt{1/3} |3/2, -1/2\rangle - \sqrt{2/3} |1/2, -1/2\rangle$$

and we will need two amplitudes: a_3 to denote that the $T = 3/2$ component of the incoming wave produces the isobar and a_1 to denote that the $T = 1/2$ component of the incoming wave produces the isobar.

The general form of any term in the matrix element will be

$$M = a_1 \vec{\sigma} \cdot \hat{q} \frac{S}{D}$$

where $\hat{q}$ is a unit vector in the direction of the incoming pion in the overall center of mass and $\vec{\sigma}$ is the nucleon spin. The term $\vec{\sigma} \cdot \hat{q}$ is a pseudoscalar representing S wave production of a single pion.

Let pion number 1 be the pion which forms the isobar and let $\hat{p}_1^i$ be its momentum in the isobar rest frame. We write $S_1 = 2\hat{q} \cdot \hat{p}_1^i - i\vec{\sigma} \cdot (\hat{q} \times \hat{p}_1^i)$ as the projection operator which picks out the $P_{3/2}$ decay. We also write $\frac{1}{D_1} \frac{(\Gamma(1)/q)^{1/2}}{W_{(IN)} - W_R - i\Gamma(1)}$ to represent the $(3,3)$ resonance. $\Gamma(1)$ is the width, W_R the $(3,3)$ resonance energy, and $W_{(IN)}$ is the energy of pion number 1 and the nucleon in their center of mass.

We will derive the matrix element for the reaction $\pi^- + p \rightarrow \pi^- + \pi^+ + n$. The other matrix elements are derived in a corresponding way. The diagrams involved are:

We use the Clebsch-Gordan coefficients to decompose the incoming wave as follows:

$$\begin{aligned}
 \sqrt{\frac{1}{3}} \left| \frac{3}{2}, -\frac{1}{2} \right\rangle &= \sqrt{\frac{8}{45}} \left| \frac{3}{2}, \frac{1}{2} \right\rangle |1, -1\rangle - \sqrt{\frac{1}{45}} \left| \frac{3}{2}, -\frac{1}{2} \right\rangle |1, 0\rangle \\
 &\quad - \sqrt{\frac{2}{15}} \left| \frac{3}{2}, -\frac{3}{2} \right\rangle |1, 1\rangle \\
 -\sqrt{\frac{2}{3}} \left| \frac{1}{2}, -\frac{1}{2} \right\rangle &= -\sqrt{\frac{1}{9}} \left| \frac{3}{2}, \frac{1}{2} \right\rangle |1, -1\rangle + \sqrt{\frac{2}{9}} \left| \frac{3}{2}, -\frac{1}{2} \right\rangle |1, 0\rangle \\
 &\quad - \sqrt{\frac{1}{3}} \left| \frac{3}{2}, -\frac{3}{2} \right\rangle |1, 1\rangle.
 \end{aligned}$$

The terms in $|\frac{3}{2}, -\frac{1}{2}\rangle|1,0\rangle$ represent particles not present in the diagram and must be dropped. Then the amplitude becomes

$$A = \frac{a_3 \vec{\sigma} \cdot \hat{q}}{\sqrt{15}} \left[\sqrt{\frac{8}{3}} \frac{S_2}{D_2} |\frac{3}{2}, \frac{1}{2}\rangle|1, -1\rangle - \sqrt{2} \frac{S_1}{D_1} |\frac{3}{2}, -\frac{1}{2}\rangle|1,1\rangle \right] \\ - \frac{a_1 \vec{\sigma} \cdot \hat{q}}{\sqrt{3}} \left[\frac{1}{\sqrt{3}} \frac{S_2}{D_2} |\frac{3}{2}, \frac{1}{2}\rangle|1, -1\rangle + \frac{S_1}{D_1} |\frac{3}{2}, -\frac{3}{2}\rangle|1,1\rangle \right] \quad (1)$$

Expand once more: $|\frac{3}{2}, -\frac{3}{2}\rangle = |\frac{1}{2}, -\frac{1}{2}\rangle|1, -1\rangle$

$$|\frac{3}{2}, \frac{1}{2}\rangle = \sqrt{\frac{1}{3}} |\frac{1}{2}, -\frac{1}{2}\rangle|1, +1\rangle + \sqrt{\frac{2}{3}} |\frac{1}{2}, \frac{1}{2}\rangle|1,0\rangle$$

Again the term in $|1,0\rangle$ represents a π^0 and must be dropped and (1) becomes

$$A = \frac{a_3 \vec{\sigma} \cdot \hat{q}}{\sqrt{15}} \left[\frac{\sqrt{8}}{3} |1,1\rangle|1, -1\rangle \frac{S_2}{D_2} - \sqrt{2} \frac{S_1}{D_1} |1, -1\rangle|1,1\rangle \right] |\frac{1}{2} - \frac{1}{2}\rangle \\ - \frac{a_1 \vec{\sigma} \cdot \hat{q}}{\sqrt{3}} \left[\frac{1}{3} |1,1\rangle|1, -1\rangle \frac{S_2}{D_2} + \frac{S_1}{D_1} |1, -1\rangle|1,1\rangle \right] |\frac{1}{2}, \frac{1}{2}\rangle \quad (2)$$

However, (2) is not symmetric in the pion wave functions. We break (2) into symmetric and antisymmetric parts, and keep only the symmetric part as required by Bose statistics. This gives

$$M(\pi^- \pi^+ n) = \frac{a_3 \vec{\sigma} \cdot \hat{q}}{\sqrt{30}} \left[\frac{2}{3} \frac{S_2}{D_2} - \frac{S_1}{D_1} \right] - \frac{a_1 \vec{\sigma} \cdot \hat{q}}{2\sqrt{3}} \left[\frac{1}{3} \frac{S_2}{D_2} + \frac{S_1}{D_1} \right] \quad (3)$$

Similarly

$$M(\pi^-, \pi^0, p) = \frac{a_3 \hat{\sigma} \cdot \hat{q}}{3 \sqrt{15}} \left(2 \frac{s_2}{D_2} - \frac{1}{2} \frac{s_1}{D_1} \right) + \frac{a_1 \vec{\sigma} \cdot \hat{q}}{3 \sqrt{6}} \left(\frac{s_1}{D_1} - \frac{s_2}{D_2} \right)$$

$$M(\pi^+, \pi^0, p) = \frac{a_3 \vec{\sigma} \cdot \hat{q}}{\sqrt{15}} \left(\frac{3}{2} \frac{s_1}{D_1} - \frac{s_2}{D_2} \right)$$

$$M(\pi^+ \pi^+ n) = -\frac{a_3 \hat{\sigma} \cdot \vec{q}}{\sqrt{60}} \left(\frac{s_2}{D_2} + \frac{s_1}{D_1} \right)$$

where the ordering and subscripts are related as shown

i.e. $\frac{s_1}{D_1}$ means pion number 1 and the nucleon form the isobar.

BIBLIOGRAPHY

1. H. Bethe and F. de Hoffman, Mesons and Fields, (Row, Peterson and Company, Evanston, 1955), Vol. II, pp. 111-131.
2. J. C. Brisson, J. F. Detoeuf, P. Falk-Variant, L. Van Rossum, and G. Valladas, Nuovo Cimento 19, 211 (1961).
3. T. J. Devlin, B. J. Moyer and V. Perez-Mendez, Phys. Rev. 125, 690 (1962).
4. J. C. Brisson, P. Falk-Variant, J. P. Merlo, P. Sonderegger, R. Turley and G. Valladas, The Aix-en-Provence International Conference on Elementary Particles, Vol. I, pp. 44-55 (1961).
5. P. Bareyre, C. Bricman, G. Valladas, G. Villet, J. Bizard, and J. Segainot, Phys. Letters 8, 137 (1964).
6. C. B. Chiu, R. J. Cence, R. D. Eandi, R. W. Kenney, B. J. Moyer, J. A. Poirier, V. Z. Peterson, and W. B. Richards, Bull. Am. Phys. Soc. 8, 603 (1963), and private communication.
7. G. B. Yodh, T. B. Day, G. Quarenì, A. Quarenì-Vignudelli, R. A. Burnstein and I. Nadelhaft, (to be published in the Physical Review).
8. C. N. Vittitoe, B. R. Riley, W. J. Fickinger, V. P. Kenney, J. G. Mowat, and W. D. Shephard, Phys. Rev. 135, B232 (1964).
9. W. S. C. Williams, An Introduction to Elementary Particles, (Academic Press, New York, 1961).
10. R. F. Peierls, Phys. Rev. Letters 6, 641 (1961).
11. J. S. Ball and W. R. Frazer, Phys. Rev. Letters 7, 204 (1961).
12. R. F. Peierls, Phys. Rev. 118, 325 (1960).
13. R. R. Wilson, Phys. Rev. 110, 1212 (1958).
14. F. P. Dixon and R. L. Walker, Phys. Rev. Letters 1, 142 (1958).

15. L. K. Goodwin, R. W. Kenney, and V. Perez-Mendez, Phys. Rev. 122, 655 (1961).
16. C. D. Wood, T. O. Devlin, J. A. Helland, M. O. Longo, B. J. Moyer, and V. Perez-Mendez, Phys. Rev. Letters 6, 481 (1961).
17. J. A. Helland, C. D. Wood, T. J. Devlin, D. E. Hagge, M. O. Longo, B. J. Moyer, and V. Perez-Mendez, Phys. Rev. 134, B1079 (1964).
18. P. M. Ogden, D. E. Hagge, J. A. Helland, M. Banner, J. F. Detoeuf, J. Teiger, Bull. Am. Phys. Soc. 8, 618 (1963) and private communication.
19. B. J. Moyer, Rev. Mod. Phys. 33, 367 (1961).
20. F. Grard, G. Macleod, L. Montanet, M. Cresti, R. Barloutand, C. Choquet, J. M. Gaillard, J. Heughebaert, A. Leveque, P. Lehmann, J. Meyer, and D. Revel, Nuovo Cimento 22, 193 (1961).
21. J. I. Shonle, Phys. Rev. Letters 5, 156 (1961).
22. P. Falk-Variant and G. Valladas, Rev. Mod. Phys. 33, 362 (1961).
23. W. M. Layson, Nuovo Cimento 27, 724 (1963).
24. P. Auvid and C. Lovelace, Nuovo Cimento 33, 473 (1964).
25. J. Sandweiss and J. R. Sanford, "Beam II Program", BNL Cosmotron Report, November 1959.
26. American Institute of Physics Handbook, (McGraw-Hill, New York, 1963), Second Ed., Chap. 6, p. 95.
27. Datpro was written by R. K. Adair and was modified for this experiment.
28. J. P. Berge, F. T. Solmitz, and H. D. Taft, Rev. Sci. Instr. 32, 538 (1961).
29. J. H. Atkinson, Jr., and B. H. Willis, High Energy Particle Data, Vol. II, UCRL-2426 (rev.) 1957.
30. E. L. Hart, "Revised Optical Constants for the BNL 20 inch Chamber", BNL memorandum, June, 1962.

31. J. Kirz, J. Schwartz, and R. D. Tripp, Phys. Rev. 130, 2481 (1963).
32. J. F. Detoeuf, Y. Ducros, J. P. Merlo, A. Stirling, B. Thevenet, L. Van Rossum, and J. Zsebery, Phys. Letters 8, 74 (1964).

Additional references are included in this paper.

33. L. I. Schiff, Quantum Mechanics (McGraw-Hill, New York, 1955), p. 105.
34. J. W. Cronin, Phys. Rev. 118, 824 (1960).
35. J. Orear, Notes on Statistics for Physicists, UCRL#8417, August 1958.
36. E. Fermi, Progr. Theoret. Phys. (Japan) 5, 570 (1950).
37. M. M. Block, Phys. Rev. 101, 796 (1956).
38. R. M. Sternheimer and S. J. Lindenbaum, Phys. Rev. 105, 1874 (1957).
39. R. M. Sternheimer and S. J. Lindenbaum, Phys. Rev. 109, 1723 (1958).
40. R. M. Sternheimer and S. J. Lindenbaum, Phys. Rev. Letters 5, 24 (1960).
41. R. M. Sternheimer and S. J. Lindenbaum, Phys. Rev. 123, 333 (1961).
42. S. Bergia, F. Bonsignori, and A. Stanghellini, Nuovo Cimento 15, 1073 (1960).
43. M. Olsson and G. B. Yodh, Phys. Rev. Letters 10, 353 (1963).
44. E. Pickup, D. K. Robinson, E. O. Salant, F. Ayer, and B. A. Munir, Phys. Rev. 132, 1819 (1963).
45. F. Bulos, R. E. Lanou, A. E. Pifer, A. M. Shapiro, M. Widgoff, R. Panvini, A. E. Brenner, C. A. Bordner, M. E. Law, E. E. Ronat, K. Strauck, J. J. Szymanski, P. Bastien, B. B. Brabson, Y. Eisenberg, B. T. Feld, V. K. Fischer, I. A. Pless, L. Rosenson, R. K. Yamamoto, G. Calvelli, L. Guerriero, G. A. Salandin, A. Tomasin, L. Ventura, C. Voci, and F. Waldner, Phys. Rev. Letters 13, 486 (1964).

46. R. C. Cence, V. Z. Peterson, V. J. Stenger, C. B. Chiu, R. D. Eandi, R. W. Kenney, B. J. Moyer, J. A. Poirier, and W. B. Richards, Bull. Am. Phys. Soc. 9, 909 (1964).
47. J. M. Blatt and V. F. Weisskopf, Theoretical Nuclear Physics, (John Wiley and Sons, New York, 1952), Chap. 8, p. 392.

FIGURE CAPTIONS

1. Total and neutral cross sections for π^+p and π^-p .
2. Beam diagram.
- 3a. Counter arrangement.
- 3b. Coincidence curve for $p_{inc} = 775 \text{ MeV/c}$.
- 3c. u-e contamination of beam.
- 4a. Schematic sketch of chamber. The cameras are in a plane 36" from the front window.
- 4b. Front view of chamber.
- 4c. Definition of azimuth and dip.
5. A typical two-prong event.
- 6a. Front view of chamber showing scanning template.
- 6b. Section view of chamber. The dotted rectangle is the final fiducial volume.
7. ξ distribution of neutrals.
8. z distribution of events.
9. Stretch distributions for the incident track of elastic events. The solid line is the theoretical curve.
- 10a. Incident momentum distribution of elastic events in the center of mass.
- 10b. Incident momentum distribution of elastic events in the laboratory.

- 11a. χ^2 distribution for inelastic events.
- 11b. χ^2 distribution for elastic events. The solid lines give the theoretical distributions.
- 12a. COP2 distributions for elastic and inelastic events.
- 12b. CORR2 distributions for elastic and inelastic events.
- 13a. Elastic missing mass.
- 13b. π^0 missing mass.
- 13c. π^+ missing mass.
- 14. Top view of chamber showing final fiducial volume. The broken line is the left hand limit of the fiducial volume used for neutrals.
- 15a. Dip distribution of events.
- 15b. Azimuth distribution of events.
- 16. Flow chart showing processing of events.
- 17a. θ_{23} distribution for elastic events.
- 17b. ϕ_- and ϕ_+ for π^+ events.
- 17c. ϕ_- and ϕ_+ for π^0 events.
- 18a. Total cross section for π^-p from 500 to 700 MeV.
- 18b. Neutral cross section from 500 to 700 MeV.
The numbers correspond to the references.
- 19a. Elastic cross section.
- 19b. π^+ production cross section.
- 19c. π^0 production cross section.

- 20a. Differential cross section.
- 20b. Goodness of fit parameters.
- 21. a_n as a function of energy.
- 22. Effective mass plots. The solid lines are phase space. The broken lines are the prediction of the Olsson-Yodh models.
- 23. Angular distributions for inelastic events, θ^* is the center of mass angle between the incident pion direction and the outgoing particle. The broken lines are the Olsson-Yodh model.
- 24. Angular distribution for π^+ production as a function of effective π^- -neutron mass.

BEAM DIAGRAM

a

b

c

FIG. 3

(a) SCHEMATIC SKETCH OF CHAMBER

(b) FRONT VIEW OF CHAMBER

(c) DEFINITION OF AZIMUTH AND DIP

FIG. 5

FIG. 6a

FIG. 6b

FIG. 7

FIG. 8

FIG. 9

FIG. 10

FIG. II

FIG. 12

FIG. 13

FIG. 14

FIG. 15d

FIG. 15 b

FIG. 16

a

b

c

FIG. 17

FIG. 18

FIG 19 a

FIG. 20

b

FIG 21

FIG. 22

FIG. 23

FIG. 24