C/375/5.17.63

ORNL-3440
UC-34 - Physics
TID-4500 (19th ed., Rev.)

MASTER

RECURSION RELATIONS AND SIMPLIFIED
EXPRESSIONS FOR THE $6-j$ SYMBOLS

C. D. Zerby

OAK RIDGE NATIONAL LABORATORY

operated by

UNION CARBIDE CORPORATION

for the

U.S. ATOMIC ENERGY COMMISSION

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

Printed in USA. Price: \$0.50 Available from the
Office of Technical Services
U. S. Department of Commerce
Washington 25, D. C.

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

ORNL-3440

Contract No. W-7405-eng-26

Neutron Physics Division

RECURSION RELATIONS AND SIMPLIFIED EXPRESSIONS
FOR THE $6-j$ SYMBOLS

C. D. Zerby

Date Issued

MAY 16 1963

OAK RIDGE NATIONAL LABORATORY
Oak Ridge, Tennessee
operated by
UNION CARBIDE CORPORATION
for the
U.S. ATOMIC ENERGY COMMISSION

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

Abstract

It is shown that the 144 symmetry properties of the $6-j$ symbols which were discovered by Regge can be displayed with a 4×4 array which is invariant to the ordering of three of the rows and all columns. The elements of the array are linear combinations of the quantum numbers. In addition, it is shown that a large set of three-term recursion relations exist. The recursion relations are used to deduce simplified expressions for the $6-j$ symbols in special cases.

The Wigner 6-j symbol,¹ which is related to the Racah W-coefficient² by

$$\left\{ \begin{array}{ccc} a & b & c \\ d & e & f \end{array} \right\} = (-1)^{a+b+e+d} W(abed, cf), \quad (1)$$

can be represented by the notation

$$\left\{ \begin{array}{ccc} a & b & c \\ d & e & f \end{array} \right\} = \begin{bmatrix} a+b-c & a+e-f & b+d-f & -c+d+e \\ a-e+f & a-b+c & c+d-e & -b+d+f \\ b-d+f & c-d+e & -a+b+c & -a+e+f \\ c+d+e+1 & b+d+f+1 & a+e+f+1 & a+b+c+1 \end{bmatrix} = \begin{bmatrix} B_{11} & B_{12} & B_{13} & B_{14} \\ B_{21} & B_{22} & B_{23} & B_{24} \\ B_{31} & B_{32} & B_{33} & B_{34} \\ B_{41} & B_{42} & B_{43} & B_{44} \end{bmatrix}, \quad (2)$$

where every element B_{ij} is a positive integer or zero.

The advantage of the square array notation is that it efficiently displays the 144 symmetry properties of the 6-j symbol that were discovered by Regge³ since it is invariant to the ordering of the four columns and to the ordering of the first three rows. This notation is similar to that used by Regge⁴ to display the symmetry properties of the 3-j symbols. The invariance of the square array can be proved with the help of Racah's formula for the W-coefficient² modified according to Eq. 1 and expressed in terms of the elements B_{ij} :

-
1. E. P. Wigner, "On the Matrices Which Reduce the Kronecker Products of Representations of Simple Reducible Groups," unpublished.
 2. G. Racah, Phys. Rev. 62, 438 (1942); Phys. Rev. 63, 367 (1943).
 3. T. Regge, Nuovo cimento 11, 116 (1959).
 4. T. Regge, Nuovo cimento 10, 544 (1958).

$$\begin{bmatrix} B_{11} & B_{12} & B_{13} & B_{14} \\ B_{21} & B_{22} & B_{23} & B_{24} \\ B_{31} & B_{32} & B_{33} & B_{34} \\ B_{41} & B_{42} & B_{43} & B_{44} \end{bmatrix} = (-1)^{B_{41}+1} \left[\frac{B_{22}! B_{23}! B_{24}! B_{32}! B_{33}! B_{34}! B_{41}!}{B_{11}! B_{12}! B_{13}! B_{14}! B_{21}! B_{31}! B_{42}! B_{43}! B_{44}!} \right]^{\frac{1}{2}}$$

$$\times \begin{Bmatrix} B_{11} & B_{12} & B_{13} & B_{14} \\ B_{21} & B_{22} & B_{23} & B_{24} \\ B_{31} & B_{32} & B_{33} & B_{34} \\ B_{41} & B_{42} & B_{43} & B_{44} \end{Bmatrix}, \quad (3)$$

where

$$\{ \} = \sum_k \frac{(-1)^{k+B_{11}} B_{11}! B_{12}! B_{13}! B_{14}! B_{21}! B_{31}! (B_{11}+B_{41}-k)!}{k! (B_{11}-k)! (B_{12}-k)! (B_{13}-k)! (B_{14}-k)! (B_{21}-B_{11}+k)! (B_{31}-B_{11}+k)! B_{41}!} \quad (3a)$$

With the aid of the new notation it is possible to display (144)³ three-term recursion relations by means of the single expression

$$\begin{bmatrix} B_{11} & B_{12} & B_{13} & B_{14} \\ B_{21} & B_{22} & B_{23} & B_{24} \\ B_{31} & B_{32} & B_{33} & B_{34} \\ B_{41} & B_{42} & B_{43} & B_{44} \end{bmatrix} = - \left[\frac{B_{12} B_{13} B_{14} B_{41}}{B_{11} B_{42} B_{43} B_{44}} \right]^{\frac{1}{2}} \begin{bmatrix} B_{11}-1 & B_{12}-1 & B_{13}-1 & B_{14}-1 \\ B_{21} & B_{22} & B_{23} & B_{24} \\ B_{31} & B_{32} & B_{33} & B_{34} \\ B_{41}-1 & B_{42}-1 & B_{43}-1 & B_{44}-1 \end{bmatrix}$$

$$- \left[\frac{(B_{11}+B_{41})^2 B_{21} B_{31}}{B_{11} B_{42} B_{43} B_{44}} \right]^{\frac{1}{2}} \begin{bmatrix} B_{11}-1 & B_{12} & B_{13} & B_{14} \\ B_{21}-1 & B_{22} & B_{23} & B_{24} \\ B_{31}-1 & B_{32} & B_{33} & B_{34} \\ B_{41} & B_{42}-1 & B_{43}-1 & B_{44}-1 \end{bmatrix}, \quad (4)$$

which may be verified with Eq. 3.

In terms of Wigner's 6-j symbols, Eq. 4 becomes

$$\begin{aligned}
& (a+b+d+e+2) \left[(a-e+f+1)(b-d+f+1) \right]^{\frac{1}{2}} \left\{ \begin{array}{ccc} a & b & c \\ d & e & f \end{array} \right\} \\
& + \left[(a+e-f)(b+d-f)(-c+d+e)(c+d+e+1) \right]^{\frac{1}{2}} \left\{ \begin{array}{ccc} a & b & c \\ d - \frac{1}{2} & e - \frac{1}{2} & f + \frac{1}{2} \end{array} \right\} \\
& + \left[(a+b-c+1)(b+d+f+2)(a+e+f+2)(a+b+c+2) \right]^{\frac{1}{2}} \left\{ \begin{array}{ccc} a + \frac{1}{2} & b + \frac{1}{2} & c \\ d & e & f + \frac{1}{2} \end{array} \right\} = 0. \quad (4a)
\end{aligned}$$

By combining sets of two equations of the type given in Eq. 4 it is possible to obtain only four independent relations which are not related through the symmetry properties. Examples of the four relations are

$$\begin{aligned}
& \left[(-c+d+e)(a-e+f+1)(b-d+f+1)(c+d+e+1) \right]^{\frac{1}{2}} \left\{ \begin{array}{ccc} a & b & c \\ d & e & f \end{array} \right\} \\
& + \left[\frac{(-c+d+e)(c+d+e+1) - (a+b-c+1)(a+b+c+2)}{(a+b+d+e+2)} \right] \left[(a+e-f)(b+d+f) \right]^{\frac{1}{2}} \left\{ \begin{array}{ccc} a & b & c \\ d - \frac{1}{2} & e - \frac{1}{2} & f + \frac{1}{2} \end{array} \right\} \\
& - \left[(a+b-c+1)(-b+d+f)(-a+e+f)(a+b+c+2) \right]^{\frac{1}{2}} \left\{ \begin{array}{ccc} a + \frac{1}{2} & b + \frac{1}{2} & c \\ d - \frac{1}{2} & e - \frac{1}{2} & f \end{array} \right\} = 0, \quad (5)
\end{aligned}$$

$$\begin{aligned}
& (-a+c-d+f) \left[(a-e+f+1)(c+d+e+1) \right]^{\frac{1}{2}} \left\{ \begin{array}{ccc} a & b & c \\ d & e & f \end{array} \right\} \\
& + \left[(b-d+f+1)(a+e-f)(b+d-f)(-c+d+e) \right]^{\frac{1}{2}} \left\{ \begin{array}{ccc} a & b & c \\ d - \frac{1}{2} & e - \frac{1}{2} & f + \frac{1}{2} \end{array} \right\} \\
& - \left[(a+b-c+1)(-a+b+c)(c-d+e)(-a+e+f) \right]^{\frac{1}{2}} \left\{ \begin{array}{ccc} a + \frac{1}{2} & b & c - \frac{1}{2} \\ d & e - \frac{1}{2} & f \end{array} \right\} = 0, \quad (5a)
\end{aligned}$$

$$\begin{aligned}
& \left[\frac{(a+b-c)(a+b+c+1) - (-c+d+e)(c+d+e+1)}{(a+b+d+e+1)} \right] \left[(b+d+f+1)(a+e+f+1) \right]^{\frac{1}{2}} \begin{Bmatrix} a & b & c \\ d & e & f \end{Bmatrix} \\
& + \left[(a-e+f)(b-d+f)(a+b-c)(a+b+c+1) \right]^{\frac{1}{2}} \begin{Bmatrix} a - \frac{1}{2} & b - \frac{1}{2} & c \\ d & e & f - \frac{1}{2} \end{Bmatrix} \\
& - \left[(-a+e+f)(-b+d+f)(-c+d+e)(c+d+e+1) \right]^{\frac{1}{2}} \begin{Bmatrix} a & b & c \\ d - \frac{1}{2} & e - \frac{1}{2} & f - \frac{1}{2} \end{Bmatrix} = 0, \quad (5b)
\end{aligned}$$

and

$$\begin{aligned}
& (a-c+d-f) \left[(c+d+e+1)(b+d+f+1)(a+e+f+1)(a+b+c+1) \right]^{\frac{1}{2}} \begin{Bmatrix} a & b & c \\ d & e & f \end{Bmatrix} \\
& (b+c+e+f+1) \left[(a+b-c)(a+e-f)(b+d-f)(-c+d+e) \right]^{\frac{1}{2}} \begin{Bmatrix} a - \frac{1}{2} & b - \frac{1}{2} & c \\ d - \frac{1}{2} & e - \frac{1}{2} & f \end{Bmatrix} \\
& - (a+b+d+e+1) \left[(b-d+f)(c-d+e)(-a+b+c)(-a+e+f) \right]^{\frac{1}{2}} \begin{Bmatrix} a & b - \frac{1}{2} & c - \frac{1}{2} \\ d & e - \frac{1}{2} & f - \frac{1}{2} \end{Bmatrix} = 0. \quad (5c)
\end{aligned}$$

With the equations that have been presented it is possible to deduce some simplified expressions for the 6-j symbols in special cases in a manner similar to that described previously for the 3-j symbols.⁵ It should be noted, for example, that the sum given in Eq. 3a is always an integer or zero since each term in the sum is an integer. Hence, when $B_{11} = 0$, the summation index can only assume the value zero and the sum is equal to unity. This fact leads to the simplified expression

$$\begin{bmatrix} 0 & B_{12} & B_{13} & B_{14} \\ B_{21} & B_{22} & B_{23} & B_{24} \\ B_{31} & B_{32} & B_{33} & B_{34} \\ B_{41} & B_{42} & B_{43} & B_{44} \end{bmatrix} = (-1)^{B_{41}+1} \left[\frac{B_{22}! B_{23}! B_{24}! B_{32}! B_{33}! B_{34}! B_{41}!}{B_{12}! B_{13}! B_{14}! B_{21}! B_{31}! B_{42}! B_{43}! B_{44}!} \right]^{\frac{1}{2}}, \quad (6)$$

5. C. D. Zerby and R. R. Coveyou, Some Simplified Expressions for the 3-j Symbols, ORNL-TM-514 (to be published).

which applies to the case in which any B_{ij} for $i \leq 3$ is equal to zero since that element can be transposed to the B_{11} position using the symmetry properties. Thus a simplified equation for the 6-j symbol is obtain whenever $j_k = j_l + j_m$ in any set of quantum numbers that form a triangle relation in the 6-j symbol.

If the minimum of the B_{ij} is greater than zero, then a form of the recursion relation given in Eq. 4 can be used to relate the symbol to symbols with the value of the element reduced by one. In this way the symbol can finally be reduced to the form given in Eq. 6. Alternatively, the element with the minimum value can be transposed to the B_{11} position and Eq. 4 can be used in the form it is presented. In the latter case there is an advantage to using the recursion relation for the symbol defined in Eq. 3a since it is always an integer. The substitution of Eq. 3 into Eq. 4 yields

$$\left\{ \begin{matrix} B_{11} & B_{12} & B_{13} & B_{14} \\ B_{21} & B_{22} & B_{23} & B_{24} \\ B_{31} & B_{32} & B_{33} & B_{34} \\ B_{41} & B_{42} & B_{43} & B_{44} \end{matrix} \right\} = \left[\begin{matrix} B_{12} & B_{13} & B_{14} \end{matrix} \right] \left\{ \begin{matrix} B_{11}-1 & B_{12}-1 & B_{13}-1 & B_{14}-1 \\ B_{21} & B_{22} & B_{23} & B_{24} \\ B_{31} & B_{32} & B_{33} & B_{34} \\ B_{41}-1 & B_{42}-1 & B_{43}-1 & B_{44}-1 \end{matrix} \right\} \\ - \left[\begin{matrix} B_{21} & B_{31} & (B_{11}+B_{41}) \end{matrix} \right] \left\{ \begin{matrix} B_{11}-1 & B_{12} & B_{13} & B_{14} \\ B_{21}-1 & B_{22} & B_{23} & B_{24} \\ B_{31}-1 & B_{32} & B_{33} & B_{34} \\ B_{41} & B_{42}-1 & B_{43}-1 & B_{44}-1 \end{matrix} \right\}. \quad (7)$$

From Eqs. 3 and 7 and the fact that the symbol with braces is equal to unity when the element in the B_{11} position is equal to zero, one can quickly deduce that

$$\begin{bmatrix} 1 & B_{12} & B_{13} & B_{14} \\ B_{21} & B_{22} & B_{23} & B_{24} \\ B_{31} & B_{32} & B_{33} & B_{34} \\ B_{41} & B_{42} & B_{43} & B_{44} \end{bmatrix} = (-1)^{B_{41}+1} \left[\frac{B_{22}! B_{23}! B_{24}! B_{32}! B_{33}! B_{34}! B_{41}!}{B_{12}! B_{13}! B_{14}! B_{21}! B_{31}! B_{42}! B_{43}! B_{44}!} \right]^{\frac{1}{2}}$$

$$\times \left\{ \left[B_{12} B_{13} B_{14} \right] - \left[B_{21} B_{31} (B_{41}+1) \right] \right\} \quad (8)$$

$$\begin{bmatrix} 2 & B_{12} & B_{13} & B_{14} \\ B_{21} & B_{22} & B_{23} & B_{24} \\ B_{31} & B_{32} & B_{33} & B_{34} \\ B_{41} & B_{42} & B_{43} & B_{44} \end{bmatrix} = (-1)^{B_{41}+1} \left[\frac{B_{22}! B_{23}! B_{24}! B_{32}! B_{33}! B_{34}! B_{41}!}{2 B_{12}! B_{13}! B_{14}! B_{21}! B_{31}! B_{42}! B_{43}! B_{44}!} \right]^{\frac{1}{2}}$$

$$\times \left\{ \left[B_{12} B_{13} B_{14} \right] \left\{ (B_{12}-1)(B_{13}-1)(B_{14}-1) - B_{21} B_{31} B_{41} \right\} \right.$$

$$\left. - \left[B_{21} B_{31} (B_{41}+2) \right] \left\{ B_{12} B_{13} B_{14} - \left[(B_{21}-1)(B_{31}-1)(B_{41}+1) \right] \right\} \right\}, \quad (9)$$

etc. Equation 8 provides expressions for the symbols when $j_k = j_\ell + j_m - 1$ in any of the four triangle relations and Eq. 9 provides similar expressions for $j_k = j_\ell + j_m - 2$.

ORNL-3440
UC-34 - Physics
TID-4500 (19th ed., Rev.)

INTERNAL DISTRIBUTION

1. Biology Library	69. T. S. Kress
2-3. Central Research Library	70. C. E. Larson
4. Reactor Division Library	71. M. E. LaVerne
5-6. ORNL - Y-12 Technical Library	72. R. V. Miskell
Document Reference Section	73. S. K. Penny
7-56. Laboratory Records Department	74. C. W. Nestor, Jr.
57. Laboratory Records, ORNL R.C.	75. Frances Perey
58. Donald Arnurius	76. R. H. Ritchie
59. J. B. Ball	77. H. C. Schweinler
60. E. P. Blizard	78. M. J. Skinner
61. Brian Buck	79. J. A. Swartout
62. J. L. Clark (K-25)	80. A. M. Weinberg
63. J. W. T. Dabbs	81-85. C. D. Zerby
64. C. D. Goodman	86. R. A. Charpie (consultant)
65. A. A. Grau	87. P. F. Gast (consultant)
66. W. H. Jordan	88. R. F. Taschek (consultant)
67. H. W. Joy	89. T. J. Thompson (consultant)
68. M. Kastenbaum	

EXTERNAL DISTRIBUTION

90. Research and Development Division, AEC, ORO
91-727. Given distribution as shown in TID-4500 (19th ed., Rev.) under
Physics category (75 copies - OTS)