MAR 24 1965

LA-DC-6969

83975

NOT FOR PUBLIC RELEASE - OFFICIAL
DISTRIBUTION MAY BE MADE. OFFICIAL
REQUESTS MAY BE FILLED. REPORT
CONTAINS NOTHING OF PATENT INTEREST.
PROCEDURES ON FILE IN RECEIVING
SECTION.

MASTER

Internal Symmetry and Relativistic Wavefunctions*

C. L. Critchfield

Los Alamos Scientific Laboratory, Los Alamos, New Mexico

It is known that states of single particles having spin may be represented by symmetric polynomials of Dirac wavefunctions.⁽¹⁾ For spins $\frac{1}{2}$, 1, $\frac{3}{2}$, 2, ... the number of components is 4, 10, 20, 35 ..., respectively. When the spin is zero a five component wavefunction is required in order to obtain first order, coupled differential equations.

The generators of the homogeneous Lorentz group form an anti-symmetrical tensor (six-vector) and may be expressed as a commutator of the four-vector components γ_k of Dirac's theory

$$S_{mn} = \frac{1}{4}i[\gamma_m, \gamma_n]. \quad (1)$$

We also have

$$[S_{mn}, \gamma_k] = i g_{nk} \gamma_m, \quad (2)$$

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:
A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.
As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

where $\underline{g}_{nk}$ is the metric tensor of special relativity. Equations (1) and (2) plus commutators between $\underline{S}_{mn}$'s form the basis of an algebra of order 10 and rank 2 in which the elements $\underline{\sigma}_z$ and $\underline{\gamma}_4$ may be diagonal in their matrix form. The algebra is therefore $\underline{C}_2$ (or $\underline{B}_2$).⁽²⁾

We may then expect that supermultiplets, in representations of $\underline{SU}_4$ for nuclear states,⁽³⁾ and $\underline{SU}_6$ for particle states,⁽⁴⁾ have similar relationships to larger groups which include Lorentz transforms. For wavefunctions in nuclei one would require a rank of four since spin is generalized to three quantum numbers which may be designated as spin, isospin and seniority. The most likely algebras are then $\underline{B}_4$ and $\underline{C}_4$. Irreducible representations may be expressed as collections of sets of integers with either sign, four to a set. Each distinct set is a weight the multiplicity of which can be calculated from Freudenthal's recursion formula.⁽⁵⁾ In such a representation of $\underline{C}_4$ the sum of the four numbers is either even or odd for all sets whereas for $\underline{B}_4$ both occur. Representations of the exceptional algebra $\underline{F}_4$ involve half-integers as well as even and odd sums. In application to nucleons therefore $\underline{C}_4$ is indicated. Similar expressions for the representations of $\underline{A}_3$ (the algebra of $\underline{SU}_4$) contain only positive integers and their sum is a constant for a given irreducible representation. For a common highest weight, $\{\underline{N}_1, \underline{N}_2, \underline{N}_3, \underline{N}_4\}$, all weights of $\underline{A}_3$ will occur in $\underline{C}_4$ and it remains only to consider the multiplicities.

Let $\underline{D(X)}$ be the dimension of a particular representation of the algebra $\underline{X}$. Each representation of $\underline{SU}_4$ may be expressed as a sum of

terms of specified dimensions in spin and isospin space: $(D(A_1), D(A_1)) = (2T + 1, 2S + 1)$. In going over to relativistic waves we have the correspondence between the dimension of the ordinary spin part and that in C_2 which is shown in the following array

$$\begin{array}{l} \text{If} \quad D(A_1) = 1 \quad 2 \quad 3 \quad 4 \quad 5 \quad \dots \\ \text{then} \quad D(C_2) = 5 \quad 4 \quad 10 \quad 20 \quad 35 \quad \dots \end{array} \quad (3)$$

Equation (3) provides the basis for comparison of a given expansion for SU_4 with that for the same highest weight in Sp_8 which will be in terms of $(D(A_1), D(C_2))$. As is to be expected, the multiplicity of these latter terms may be greater than necessary to include those of SU_4 and also representations of C_2 occur which do not appear in Eq. (3), e.g., of dimensions 1, 14, 16, another 35, and so on. Calculations of all representations having a sum of N 's equal to four and less show that those of A_2 are indeed contained in those of C_4 in the sense of the correspondence in Eq. (3).

We may now generalize the procedure suggested for SU_4 to the internal symmetry group of baryons and mesons, including spin, viz. SU_6 . Instead of nucleons we have "quarks"⁽⁶⁾ and we look for the decomposition of the representation of C_6 with highest weight {300000} in terms of the dimensions in SU_3 and Sp_4 : $(D(A_2), D(C_2))$,

$$\{300000\} = (10,20) + \underline{(8,16)} + (8,4) + (1,4) . \quad (4)$$

This is of 364 dimensions. Ignoring $D(\underline{C}_2) = 16$, as before, and comparing with Eq. (3) we see that $(10,20)$ corresponds to the $\frac{3}{2}$ decuplet and $(8,4)$ to the $\frac{1}{2}$ octet of baryons. In addition there is a singlet of spin $\frac{1}{2}$ and the same parity. Applying this procedure to two twelve dimensional representations of $\underline{C}_6$, one for a quark and one for an anti-quark

$$\{100000\} \otimes \{00000\bar{1}\} = (9,10) + (8,5) + (1,5) + \underline{(8,1)} + \underline{(1,1)} . \quad (5)$$

Again ignoring the underlined $\underline{C}_2$ singlets we are left with a vector meson nonet and a pseudoscalar $(8,5)$ just as in the regular representation of $\underline{SU}_6$, plus a pseudoscalar singlet $(1,5)$. So far as Eqs. (4) and (5) are concerned we could have disregarded the five dimensional representations of $\underline{C}_2$ and kept the singlets with the same result. This is not done because there is no Dirac equation for a wavefunction of one component and moreover, under such a procedure, the representations of $\underline{Sp}_8$ that were calculated would not contain all those of $\underline{SU}_4$.

Thus the symplectic group of rank six, $\underline{Sp}_{12}$, permits the same quantum numbers as $\underline{SU}_6$ plus that of $\underline{\gamma}_4$ which when considered to be an infinitesimal operator, as it is here, is a form of particle gauge operator since the sign of its eigenvalues is that of the energy in hole-theory.

REFERENCES

*Work performed under the auspices of the U. S. Atomic Energy Commission.

¹V. Bargmann and E. P. Wigner, Proc. Natl. Acad. Sci. U.S. 34, 211 (1948).

²E. B. Dynkin, Amer. Math Soc. Trans. II, Vol. 6, p. 319.

³E. P. Wigner, Phys. Rev. 51, 106 (1937).

⁴F. Gürsey and L. Radicati, Phys. Rev. Letters 13, 173 (1964),
A. Pais, Phys. Rev. Letters 13, 175 (1964).

⁵M. Konuma, K. Shima and M. Wada, Suppl. Progr. Theoret. Phys. 28, 44 (1963).

⁶M. Gell-Mann, Phys. Letters 8, 214 (1964).