RECEIVED BY DTIC OCT 26 1971

UCLA - 12 - 8 40

MASTER

UNIVERSITY OF CALIFORNIA, LOS ANGELES

LABORATORY OF NUCLEAR MEDICINE AND RADIATION BIOLOGY

CONTRACT NO. AT (04-1) GEN-12

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

STABILITY OF UNSATURATED METHYL ESTERS OF FATTY ACIDS ON SURFACES*

by

Vida Slawson and James F. Mead

LABORATORY OF NUCLEAR MEDICINE AND RADIATION BIOLOGY

900 VETERAN AVENUE

UNIVERSITY OF CALIFORNIA, LOS ANGELES, CALIFORNIA 90024

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

*These studies were supported in part by Contract AT (04-1) GEN-12 between the Atomic Energy Commission and the University of California and by USPHS research career award GM-K6-19,177 from the Division of General Medical Sciences, National Institutes of Health, U.S.A.

SUMMARY The stability of unsaturated methyl esters is greater absorbed on silica gel than on a glass surface. Storage of small samples absorbed on silica gel may be a convenient addition to conventional methods of protecting labile fats against autoxidation.

SUPPLEMENTARY KEY WORDS

autoxidation

• silica gel storage

Preservation of samples before analysis is important to biochemists handling materials containing polyunsaturated fats. When storage is necessary, it should be at low temperature, under an inert gas, possibly with added antioxidant. These measures decrease but do not always prevent autoxidative and enzymatic changes.

In the course of investigating the increased stability to autoxidation found for unsaturated methyl esters chromatographed on silver nitrate-silicic acid (1), the effects of adsorption of these esters on different surfaces was studied. The type of surface can be important from several considerations. It may contain atoms of transition metals, causing ordinary metal catalysis (2). Very small amounts of certain metals can have profound effects on autoxidation rates (3-5). The orientation of molecules on the surface can also influence oxidations. When cottonseed oil is spread on a gelatin surface there is less oxidation than on a glass surface (6). Since the pi-bonding of unsaturated esters to silver ions is strong enough to permit chromatographic separation of compounds with different degrees of unsaturation, we hoped that the specific binding of the unsaturated bonds of methyl esters from biological materials to AgNO_3 -impregnated silica gel might be strong enough to afford significant protection against autoxidation. Hopefully, this would be a useful adjunct to methods commonly used to protect small quantities of unsaturated fats.

Materials. Pentane, hexane, cyclohexane, diethyl ether and silver nitrate were reagent grade and used as received. Cyclohexene, b.p. 83°C , was distilled to remove the p-methoxyphenol (b.p. 243°C) added as a

stabilizer by the manufacturer. It was then passed through a silica gel column and stored in the dark at 4°C. Silica gel (Baker 3405, lot 34110) was used as received for one part of the experiment. It was impregnated with 9.0% AgNO₃ by addition of an acetonitrile solution of the reagent to an acetonitrile slurry of the adsorbent, followed by removal of the solvent with vacuum and heat in a rotating evaporator. A mixture of methyl esters with 0-6 methylene interrupted double bonds was made from commercial esters of >90% purity. The ester mixture was stored at -20°C until purified by chromatography. A 2 x 14 cm column of silica gel (Baker, 3405) was washed with 50 ml of pentane. A cyclohexane solution of the esters was poured on the top of the column. 50 ml of pentane eluted 0.0024 g (I). 100 ml of 4% diethyl ether in pentane eluted 0.2375 g (II). 75 ml diethyl ether eluted 0.0176 g with typical autoxidation odor (III). I and III were discarded. II was diluted with 50 ml of hexane to give a concentration of 0.00475 g/ml. The flask was stoppered and kept at -20°C except when in use. GLC composition of the mixture was palmitate, 3.29%; oleate, 9.4%; linoleate, 12.1%; linolenate, 15.3%; arachidonate, 15.4%; eicosapentaenoate, 18.9%; and docosahexaenoate, 25.5%.

Procedure. A 1 ml aliquot of hexane containing 0.00475 g of mixed esters was pipetted into an empty 125-ml Erlenmeyer flask to serve as a control. 1 ml aliquots were also pipetted into similar flasks containing 0.1 g of AgNO₃-silica gel, 0.5 g of AgNO₃-silica gel, 0.1 g of silica gel, and 0.5 g of silica gel. The flasks were swirled to mix the contents. They were loosely covered with beakers to exclude dust but not air, and then placed in a 4°C refrigerator. The refrigerator was dark except for

occasional openings to remove and return contents. Sampling initially consisted of the addition of cyclohexene-diethyl ether (1:1), mixing, removal of an appropriate volume with a microsyringe, and direct injection on a polar column. The flask was immediately replaced in the refrigerator for storage until the next sampling. Because the broad solvent peaks interfered with accurate measurement of early peaks, the sampling technique was modified. A larger amount of cyclohexene was added and the flask swirled. A portion was withdrawn and filtered through a small plug of pyrex wool in a disposable Pasteur pipette. The cyclohexene was evaporated with nitrogen at room temperature. The sample was then dissolved in pentane and injected immediately on to the GLC column.

Gas Liquid Chromatography (GLC). A Barber-Colman Model 10 with a flame ionization detector was used for analyses. The column was 12.2% (w/w) ethylene glycol succinate on acid washed, siliconized Chromosorb G, 6 mm x 41", 80-100 mesh. At 185-200°C and 15-20 psig argon, methyl palmitate eluted in 2-3 minutes. Accuracy was within $\pm 5\%$ of major components.

Calculation. Methyl palmitate was presumed to be unchanged under conditions of the experiment. The area ratios of the unsaturated esters to methyl palmitate were calculated, divided by the initial area ratios for the individual esters, and multiplied by 100 to give percent of the original ester remaining.

Results. There was a considerable difference between the stability of the unsaturated esters on glass and the two adsorbents. After 16 days at 4°C there were no esters with more than three double bonds found on

the glass surfaces (3 trials). After 42 days at 4° C there were varying amounts of all the original compounds on the adsorbents. The differences between the two adsorbents are shown in Table 1. Although the silver nitrate-treated silica gel preserved the unsaturated esters longer than the glass surface, the unaltered silica gel was even better. It is possible that crystallization of silver nitrate in the pores of the silica gel reduced the surface area. Oxygen uptake by soybean oil on silica gel was an inverse function of the surface area when the amount of oil was less than required for a monomolecular layer (7). It is equally possible that Ag^+ promotes some oxidation.

← TABLE 1

Comparison of Oxidation on Different Surfaces. A very crude estimate of the difference in stability on a glass surface with that on the silica gel may be obtained by comparing the time required by a given ester to reach about 50% of its original value. Fortunately, the values for the 63-day silica gel sample and the 5-day glass control sample are similar for most of the esters. This is shown in Table 2. The esters stored on silica gel might be considered $\frac{63}{5}$, or 12.6 times more stable than the control. A more satisfying approach is to assume that methyl oleate does not autoxidize until the majority of the polyunsaturated materials are gone. Inspection of the GLC analyses for samples containing 3 or more double bonds indicated that the value for 18:1 was as likely to be above 100% as below (7 vs. 5), and the 18:1/16:0 ratios did not decrease. Chromatograms in which methyl oleate deviates from 100% by more than 5% are presumed to err by inaccuracy in the palmitate peak. They are corrected by multiplying each component by the factor necessary to adjust oleate

← TABLE 2

to 100%. From the original percent composition and percent remaining at a given time, the actual amount of each ester oxidized is calculated. A plot of μ moles of each ester oxidized vs. the number of double bonds in the ester is roughly linear. Figure 1 shows typical plots. The data from Table 2 have been included. The slope of a line gives the average μ moles of ester oxidized per double bond. Division by the days of storage gives the average oxidation rate in μ moles per bond per day. The comparison made from the data in Table 2 can also be made from Figure 1. The slope of the 5 day glass sample is 0.283; the oxidation rate is 0.057 μ moles/bond/day. The slope of the 63 day, 0.5 g silica gel sample is 0.231; the oxidation rate 0.0037 μ moles/bond/day. The ratio of the oxidation rates is $\frac{0.057}{0.0037}$. The esters are 15 times less stable on glass than on silica gel. For practical purposes it is sufficient to say that storage on silica gel has a significant protective effect.

Discussion. There is considerable evidence that polyunsaturated esters of fatty acids are much more susceptible to autoxidation than monounsaturated esters (8-10). The linear correlation of susceptibility to autoxidation with number of double bonds is not unreasonable, since there has been no oxidation of methyl oleate. However, this does not coincide with the relative reactivities of 18:2 and 18:3 shown by a recent literature survey (11). It is possible that our results have been influenced by the unequal composition of the starting mixture, or that decomposition products of the highly unsaturated esters have changed the rates of 18:2 and 18:3. Determination of the oxidation rates of single esters would answer these questions.

← FIGURE 1

There are several possible explanations, not mutually exclusive, for the effect of silica gel. A primary factor may be differences in trace metal content of this silica gel and glass. The importance of surface geometry of the adsorbent to adsorption (12, 13), and to chromatographic separation has been mentioned (14). Optimum pore size and surface area for chromatographic separations have been related to the size of the molecules separated (15, 16). It is possible that moisture content of the silica gel as well as its surface had an effect on the stability of the esters. This was demonstrated with methyl linoleate on cellulose (17, 18) where oxidation was markedly reduced by humidification. Water has clearly been shown to increase the binding of some compounds to silica gel (13). There was a substantial increase in the heat of adsorption of benzene and hexane after hydration of the silica gel. A logical presumption is that with a stronger bond to silica gel, an unsaturated molecule becomes less reactive with atmospheric oxygen.

In view of the general observations that oxidation rates for unsaturated materials are connected with surface area and moisture content, it is not difficult to accept our results. Determining basic reasons for the observed differences in oxidation rates would be of practical value because a better understanding of the most relevant factors would facilitate choice of a better support.

Conclusion. Highly unsaturated fatty acid esters when stored on silica gel were 12 to 15 times more stable than when stored on a glass surface. Therefore, in addition to usual precautions of low temperature and protection from oxygen, the use of silica gel to provide extra stability

is recommended. Deposition of fatty acid esters onto silica gel by evaporation of solvent and subsequent removal with a polar solvent are simple and rapid procedures. Since the unsaturated methyl esters are stable on silica gel for days, the collection and temporary storage of fractions from GLC separations would be facilitated by use of silica gel. If the technique is used for other lipid compounds, caution should be exercised, since undesirable changes may be accelerated by silica gel (19, 20).

Notice of grant support. These studies were supported in part by Contract AT(04-1)GEN-12 between the Atomic Energy Commission and the University of California and by USPHS research career award GM-K6-19,177 from the Division of General Medical Sciences, National Institutes of Health, U.S.A..

REFERENCES

1. Slawson, V., and R. Stein. 1970. Comparative autoxidative susceptibility of fatty esters with 0-6 methylene-interrupted double bonds. Lipids 5: 713-717.
2. Holman, R. T. 1954. Autoxidation of fats and related substances. Prog. Chem. Fats Other Lipids 2: 51-98.
3. Ingold, K. U. 1968. Principles of metal catalyzed lipid oxidation. In SIK Symposium. R. Marcuse, editor. SIK-Rapport No. 240. Kallebäck, Göteborg, Sweden. 11-41.
4. Uri, N. 1956. Metal ion catalysis and polarity of environment in the aerobic oxidation of unsaturated fatty acids. Nature (London) 177: 1177-1178.
5. Heaton, F. W., and N. Uri. 1961. The aerobic oxidation of unsaturated fatty acids and their esters: cobalt stearate-catalyzed oxidation of linoleic acid. J. Lipid Res. 2: 152-160.
6. Togashi, H. J., A. S. Henick, and R. B. Koch. 1961. The oxidation of lipids in thin films. J. Food Sci. 26: 186-191.
7. Honn, F. J., I. I. Bezman, and B. F. Daubert. 1951. Autoxidation of drying oils adsorbed on porous solids. J. Amer. Oil Chem. Soc. 28: 129-133.
8. Holman, R. T. 1966. General introduction to polyunsaturated acids. Prog. Chem. Fats Other Lipids 9: 3-12.
9. Holman, R. T., and O. C. Elmer. 1947. The rates of oxidation of unsaturated fatty acids and esters. J. Amer. Oil Chem. Soc. 24: 127-129.
10. Gunstone, F. D. 1958. An Introduction to the Chemistry of Fats and Fatty Acids. Wiley, New York. 114-115.

11. Ackman, P. G., J. C. Sipos, and P. M. Jangaard. 1967. A quantitation problem in the open tubular gas chromatography of fatty acid esters from cod liver lipids. Lipids 2: 251-257.
12. Adamson, A. W. 1967. Physical Chemistry of Surfaces, 2nd ed. Interscience, New York, London and Sydney. 406.
13. Kiselev, A. V. 1961. Surface chemistry, adsorption energy and adsorption equilibria. Quart. Rev. Chem. Soc. 15: 99-124.
14. Sporer, A. H., and K. N. Trueblood. 1959. A quantitative study of the relation between molecular structure and adsorption affinity on silicic-acid-celite columns. J. Chromatogr. 2: 499-512.
15. Klein, P. D. 1961. Silica gel structure and the chromatographic process. Effect of pore diameter on the adsorption and differential migration of sterol acetates. Anal. Chem. 33: 1737-1741.
16. Waksmundzki, I. A., J. Oscik, J. Matusiewicz, R. Nasuto, and J. Rozylo. 1961. The structure of silica gels adsorbing specifically pyridine, quinoline, and acridine. Przem. Chem. 40: 387-390. Chem. Abstr. 56: 20181 (1962).
17. Labuza, T. P., H. Tsuyuki, and M. Karel. 1969. Kinetics of linoleate oxidation in model systems. J. Amer. Oil Chem. Soc. 46: 409-416.
18. Labuza, T. P., N. D. Heidelbaugh, M. Silver, and M. Karel. 1971. Oxidation at intermediate moisture contents. J. Amer. Oil Chem. Soc. 48: 86-92.
19. Smith, I. I., and T. Foell. 1962. Thin-layer chromatography of steroids on starch-bound silica gel chromatoplates. J. Chromatogr. 9: 339-344.
20. Marinetti, G. V., J. Erbland, and E. Stotz. 1959. The structure of beef heart plasmalogens. J. Amer. Chem. Soc. 81: 861-864.

TABLE 1 VARIATION OF STABILITY OF UNSATURATED METHYL ESTERS
WITH SURFACE AFTER 42 DAYS*

Methyl Ester	AgNO ₃ -SiO ₂	SiO ₂
	% Remaining [†]	% Remaining [†]
18:1	97.6	103.1
18:2	41.7	106.2
18:3	24.7	79.6
20:4	10.4	83.5
20:5	8.7	76.3
22:6	7.9	59.9

* 0.00475 g of ester mixture on 0.1 g adsorbent, 4° C, dark

† Relative to 16:0

TABLE 2 · COMPARISON ON SILICA GEL AND GLASS SURFACE*

Methyl Ester	63 Days Silica Gel†	5 Days Glass
	% Remaining†	% Remaining†
18:1	104.8	101.5
18:2	85.2	84.0
18:3	75.9	74.2
20:4	75.0	65.7
20:5	67.7	57.1
22:6	64.7	60.2

* 0.00475 g of mixed methyl esters, 4° C, dark

† 0.5 g

‡ Relative to 16:0

FIG. 1 Relation of Oxidation to Amount of Unsaturation*

X	X	Glass, 5 days, uncorrected (TABLE 2)
.	.	Silica gel, 0.1 g, 63 days, corrected
O	O	Silica gel, 0.5 g, 63 days, uncorrected (TABLE 2)
△	△	AgNO ₃ -silica gel, 0.1 g, 42 days, uncorrected
△	△	AgNO ₃ -silica gel, 0.5 g, 42 days, corrected

*15.2 μ moles (0.00475 g) mixed esters, 4° C dark