PORTIONS OF THIS REPORT ARE ILLEGIBLE. It
has been reproduced from the best available
copy to permit the broadest possible avail-
ability.

i

Crystal Structures of Diketone Complexes of Alkaline Earth Metals and Other Compounds.

Table of Contents

Table of Contents	i
Abstract	v
Introduction	xiii
Acknowledgements and Dedication	xi
I. The Crystal and Molecular Structure of $((C_6H_5)_3PCH_3)^+$ $((B_6C_2H_8)Mn(CO)_3)^-$, etc.	
A. Introduction	1
B. Experimental Procedure	2
C. Determination and Refinement	5
D. Results and Discussion	12
E. References	23
II. Determination of the Crystal Structure of $XeF_5^+ AsF_6^-$.	
A. Introduction	29
B. Experimental Procedure	30
C. Solution and Refinement of the Structure	34
D. Results and Discussion	41
E. References	48
III. The Crystal and Molecular Structure of 1,3-diphenyl- 1,3-propanedione enol (HDPP)	
A. Introduction	49
B. Experimental Procedure	50
C. Determination and Refinement	54
D. Results and Discussion	61
E. References	71

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, agents, contractors, or subcontractors, by their conduct, warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

IV. The Crystal and Molecular Structure of bis(dimethyl- formamido)-bis(1,3-diphenyl-1,3-propanedionato)magnesium. $\text{Mg}(\text{DPP})_2(\text{DMF})_2$.	
A. Experimental Procedure	72
B. Determination and Refinement	77
C. Results and Discussion	87
D. References	98
V. The Crystal and Molecular Structure of bis(1,3-diphenyl- 1,3-propanedionato)calcium hemiethanolate. $\text{Ca}(\text{DPP})_2$ - $(\text{C}_2\text{H}_5\text{OH})_{\frac{1}{2}}$.	
A. Experimental Procedure	99
B. Determination and Refinement	103
C. Results and Discussion	115
D. References	131
VI. The Crystal and Molecular Structure of bis(1,3-diphenyl- 1,3-propanedionato)strontium hemiacetate. $\text{Sr}(\text{DPP})_2$ - $((\text{CH}_3)_2\text{CO})_{\frac{1}{2}}$.	
A. Experimental Procedure	132
B. Determination and Refinement	136
C. Results and Discussion	143
D. References	156

VII. Summary and Comparison

A. Introduction	157
B. $Mg(DPP)_2(DMF)_2$ and Other Octahedra	158
C. $Ca(DPP)_2(C_2H_5OH)_{\frac{1}{2}}$, $Sr(DPP)_2((CH_3)_2CO)_{\frac{1}{2}}$ and Other Polymers	160
D. A Note on the DPP^- Ligand	164
E. References	166

Crystal Structures of Diketone Complexes of Alkaline Earth Metals and Other Compounds

Abstract

Frederick John Hollander

The structures of six compounds were determined by means of x-ray crystallographic techniques applied to single crystals of $((C_6H_5)_3PCH_3)^+((B_6C_2H_8)Mn(CO)_3)^-$, $XeF_5^+AsF_6^-$, 1,3-diphenyl-1,3-propanedione enol (HDPP), $Mg(DPP)_2(dimethylformamide)_2$, $Ca(DPP)_2(C_2H_5OH)_2$, and $Sr(DPP)_2((CH_3)_2CO)_2$.

$((C_6H_5)_3PCH_3)^+((B_6C_2H_8)Mn(CO)_3)^-$ crystallizes in the triclinic space group $\overline{P}1$ with cell dimensions $a = 11.216(5)$ Å, $b = 11.285(4)$ Å, $c = 11.464(5)$ Å, $\alpha = 105.79(1)^\circ$, $\beta = 104.46(1)^\circ$, $\gamma = 98.04(1)^\circ$. The observed density is 1.25 gm/cm^3 , compared to 1.29 gm/cm^3 calculated for one molecule in the asymmetric unit. The structure was solved by Patterson methods and refined to an R value of 0.041 for 2862 reflections. The cation has a propeller shape as expected. The anion has a non-crystallographic mirror plane and confirms the structure expected on the basis of spectroscopic work. Comparisons are made to some other dicarbollide fragments.

$XeF_5^+AsF_6^-$ crystallizes in the monoclinic space group $P2_1/c$ with cell dimensions $a = 5.886(3)$ Å, $b = 16.054(10)$ Å, $c = 9.051(4)$ Å, $\beta = 91.564(35)^\circ$. Volume criteria were used to determine that there were four formula units per unit cell. The structure was solved by Patterson methods and refined to an R value of 0.116 for 925 reflections. The salt forms dimer rings containing two cations and two anions connected by bridging fluorines around

a center of symmetry. The forms of the cation and anion are respectively square pyramidal and octahedral. Deviations from ideal forms for both the cation and the anion are discussed and compared to results for other structures.

HDPP crystallizes in the orthorhombic space group $Fbca$ with cell dimensions $a = 10.853(1) \text{ \AA}$, $b = 24.441(1) \text{ \AA}$, $c = 8.755(1) \text{ \AA}$. The measured density of 1.22 gm/cm^3 compares to the calculated density of 1.28 gm/cm^3 for eight molecules in the unit cell. The structure was solved by direct methods and refinement proceeded to an R value of 0.027 for 1070 reflections. HDPP exists entirely in the enolic form and there is a strong internal hydrogen bond involving the enolic hydrogen. The structure is described and compared to the results of a previous study.

$Mg(DPP)_2(DMF)_2$ crystallizes in a monoclinic unit cell with dimensions $a = 16.893(3) \text{ \AA}$, $b = 12.853(2) \text{ \AA}$, $c = 16.927(3) \text{ \AA}$, $\beta = 117.085(5)^\circ$. The centric space group $C2/c$ was chosen over the non-centric group Cc on the basis of refinement in the former. The structure was determined by direct methods and refined to an R value of 0.067 on 1817 reflections. The coordination of the Mg atom is octahedral, with the Mg on a crystallographic two-fold axis. The two DMF molecules are cis to one another on the coordination octahedron.

$Ca(DPP)_2(C_2H_5OH)_{\frac{1}{2}}$ crystallizes in the triclinic space group $\bar{P}1$ with cell dimensions $a = 15.247(1) \text{ \AA}$, $b = 13.555(1) \text{ \AA}$, $c = 14.097(1) \text{ \AA}$, $\alpha = 74.64(1)^\circ$, $\beta = 95.97(1)^\circ$, $\gamma = 113.59(1)^\circ$. The density calculated with four formula units in the cell is $d_c = 1.32 \text{ gm/cm}^3$. The structure was solved by direct methods and refined

to an R value of 0.040 for 4503 reflections. The complex forms novel polymeric clusters around a center of symmetry. There are two Ca atom environments, one six- and one seven-fold coordinated. The form of the polymer is described in detail.

$\text{Sr}(\text{DPP})_2((\text{CH}_3)_2\text{CO})_{1/2}$ crystallizes in the triclinic space group $\bar{P}1$ with cell dimensions $a = 13.759(10) \text{ \AA}$, $b = 14.182(9) \text{ \AA}$, $c = 16.031(9) \text{ \AA}$, $\alpha = 107.57(3)^\circ$, $\beta = 90.36(3)^\circ$, $\gamma = 113.07(3)^\circ$. The data crystal was poor, with data out only to an effective limit of $2\theta = 25^\circ$ in molybdenum radiation. The structure was solved by Patterson methods and refined to an R value of 0.088 on 1561 reflections. The hydrogen atoms were not located. The structure shows similar novel polymerization to the Ca complex, though differences between the two structures are enough to make them non-isomorphous. The polymerization is described in detail.

Comparison is made between the DPP complexes studied in this thesis and other β -diketone complexes. The $\text{Mg}(\text{DPP})_2(\text{DMF})_2$ structure is compared with other octahedral β -diketone complex structures, and the Sr and Ca complexes are compared with polymeric acetylacetonates and with each other. Some notes are made on the structure of the DPP^- ligand.

INTRODUCTION

Six determinations of molecular and crystal structure by means of x-ray diffraction are presented in this thesis. The nature of structural determination is such that the most convenient way to present the investigative work and description is in the form of separate studies. The first six sections are the six such studies I have undertaken in my graduate career. The closing section is a comparison of the results of the four related structures to each other and to similar published structures. Tables and Figures are labeled independently in each of the seven sections and the references for each section are collected at the end of that section.

The first two structure determinations are totally unrelated, and were performed to confirm predicted molecular structures for other workers, specifically M. Frederick Hawthorne (Section I) and Prof. Neil Bartlett (Section II).

The remaining four structure determinations have in common the molecule 1,3-diphenyl-1,3-propanedione (HDPP). Section III gives the determination of the structure of HDPP, and sections IV - VI give the determinations of the structures of the complexes of the alkaline-earth metal ions Mg^{++} , Ca^{++} and Sr^{++} with DPP^- and various neutral solvation molecules.

The choice of subject for these four structure determinations was brought about by a chance reading of an article on the complexes of β -diketones with metals (Fackler, 1966). One of the facts noted in passing was that the series $M(DPM)_2$ ($HDPM = 2,2,6,6$ -tetramethyl-3,5-heptanedione = dipivaloylmethane), where

M was an alkaline-earth metal, had the following melting points: Be, 93-97°C; Mg, 94°C; Ca, 204°C; Sr, 200°C; Ba, 172°C. Intrigued by the anomaly in the melting point series between Mg and Ca, I investigated further to find that no work had been done in the area of crystal structures of β -diketone complexes of the alkaline-earth metals with the exception of two structures of alkaline-earth acetylacetonates (Amirthalingam, *et al.*, 1960; Morosin, 1967). Since the effect on the melting points was observed with β -diketones having more bulky side groups than $-\text{CH}_3$, I decided to investigate the alkaline-earth metal complexes of HDEM and HDPP (actually DEM^- and DPP^-). Accidents of procedure, rather than any plan, led to the study of the compounds herein rather than others in the two series.

A drastic change in molecular structure of the DPP complexes does take place between Mg and Ca, while the Sr complex has a structure similar to the Ca complex. A similar change may take place in the series of DEM complexes, or there may be an entirely different structural change which accounts for the originally observed melting point differences. A recent paper by Schwarberg, Sievers and Moshier (1970) indicates strongly that the $\text{Mg}(\text{DEM})_2$ complex is monomeric in solution and in the vapor phase, while the Ca, Sr and Ba DEM complexes are dimeric. Since the results found for the DPP series ranged from monomer to tetramer in terms of the basic $\text{M}(\text{DPP})_2$ unit, it is quite possible that the structural change is different in the DEM series.

It is obvious that the opportunities for further investigation in this area are far from exhausted. I hope to be able to

continue to investigate these structures, and not have this thesis become a dead end.

References

- Amirthalingam, V., Padmanabhan, V.M. and Shankar, J. (1960). Acta Cryst., 13, 201.
- Fackler, J.P. (1966). Progress in Inorganic Chemistry, 7, 361.
- Morosin, B. (1967). Acta Cryst., 22, 316.
- Schwarberg, J.E., Sievers, R.E. and Moshier, R.W. (1970). Anal. Chem., 42(14), 1828.

Acknowledgements

I would like to thank Dr. Allan Zalkin and Mrs. Helena Ruben for their guidance and aid, Dr. Mark Fisher and Dr. David St. Clair for their early inspiration, Dr. Siddhartha Ray for his example of careful workmanship, Art Olson and Fran Jurnak for keeping me from climbing the walls, and the Graphic Arts Dept. at LBL for their friendliness and efficiency on short notice.

I would most especially like to thank Dr. Richard E. Marsh, who started me on this whole trail, and Prof. David H. Templeton, my adviser, whose careful thought and thorough reasoning were invaluable. I would also like to thank the other two members of my thesis committee, Prof. Ken Raymond and Prof. Hans-Rudolf Wenk for their time and effort.

Much thanks to the Lawrence Berkeley Laboratory and to the United States Atomic Energy Commission for their generous support.

Dedication

This work is dedicated to my wife, Lynn, who puts up with me.

I. The Crystal and Molecular Structure of $(\text{C}_6\text{H}_5)_3\text{PCH}_3^+$
 $(\text{B}_6\text{C}_2\text{H}_8\text{Mn}(\text{CO})_3)^-$, the triphenylmethylphosphonium salt
 of the manganese tricarbonyl derivative of the $\text{B}_6\text{C}_2\text{H}_8^{2-}$
 dicarbollide ion.

A. Introduction

The $(\text{B}_6\text{C}_2\text{H}_8\text{Mn}(\text{CO})_3)^-$ ion was synthesized by Hawthorne and Pitts (1967) and a tentative structure assigned to it on the basis of IR and NMR spectroscopic data (Hawthorne and George, 1969). This study of the triphenylmethylphosphonium (TMP) salt was undertaken to confirm the proposed structure by x-ray crystallographic means.

B. Experimental Procedure

Well-formed crystals of the compound were supplied to us by M. Frederick Hawthorne as prepared by A. Denise Pitts. They were sealed in a vial under hexane, but proved to be air-stable during the data collection. The crystals were needlelike and a clear orange-red. Optical extinction between crossed polarizers was very sharp and the crystals were used as supplied to us with no further recrystallization. Precession photographs indicated that the unit cell was triclinic, and preliminary cell dimensions were measured from hk0 and h0l photographs. A crystal of approximate dimensions 0.18 mm x 0.14 mm x 0.19 mm was attached to a glass fiber and mounted on a G.E. XRD-5 single crystal diffractometer equipped with a manual quarter-circle Eulerian cradle goniostat. The orientation of the crystal was determined by centering reflections identified from the precession photographs, and a strong reflection, $(\bar{1} \bar{2} 1)$, found near $\chi = 90^\circ$ was moved to $\chi = 90^\circ$ to facilitate checking of the orientation of the crystal. Twenty-seven high-angle reflections were then accurately centered in χ , ϕ , and 2θ using molybdenum radiation ($K\alpha_1 = 0.709261 \text{ \AA}$, $K\alpha_2 = 0.713543 \text{ \AA}$, $K\beta = 0.632253 \text{ \AA}$), and a least-squares analysis performed to calculate accurate cell dimensions and crystal orientation matrix. The derived cell dimensions (and their standard deviations) are $a = 11.216(5) \text{ \AA}$, $b = 11.285(4) \text{ \AA}$, $c = 11.464(5) \text{ \AA}$, $\alpha = 105.79(1)^\circ$, $\beta = 104.46(1)^\circ$, $\gamma = 98.04(1)^\circ$. The observed density was 1.25 gm/cm^3 , measured by flotation in a mixture of n-hexane and carbon tetrachloride. This compares to a calculated density of 1.29 gm/cm^3 based on two formula units in a

unit cell volume of $1318 \text{ \AA}^3$. The measured density is low due to difficulties in getting the solution to wet the crystals properly.

Intensity data was collected on the same crystal using filtered Mo K α radiation (0.001" Zr in the diffracted beam) with the distances from the source and receiving apertures to the crystal 14.5 and 17.8 cm respectively. All reflections lying in a hemisphere in reciprocal space $\chi = 0^\circ$ - 90° , $\phi = 0^\circ$ - 360° out to $2\theta = 45^\circ$ ($\sin\theta/\lambda = 0.5385$) were counted for 10 sec with both crystal and counter stationary at an x-ray tube take-off angle of 4° . Background counts were taken from a plot of background vs. 2θ averaged over several values of χ and ϕ . Individual background intensities were measured for 10 sec off-set from the peak by 1° in 2θ for those reflections at $2\theta < 5^\circ$ and for those reflections affected by streaking from lower orders. A standard background correction of 1 count was applied to all reflections with $2\theta > 25^\circ$. Periodic checks of three standard reflections showed only small ($\pm 1\%$) random fluctuations in intensity over the data collection period. The data were then punched on cards and put into our standard data reduction programs. 3445 data were collected, of which 607 were measured as less than or equal to the background and another 5 of which had $I < \sigma(I)$.

Inspection of the intensity of the $(\bar{1} \bar{2} 1)$ as a function of ϕ showed no significant variation, and the values of μ and μ_R were low (5.9 cm^{-1} and 0.11), so no absorption correction was made to the intensities. Lorentz and polarization factors were applied to the data to convert them to F^2 values.

Atomic scattering factors of Cromer and Waber (1965) for the non-hydrogen atoms and those of Stewart, Davidson and Simpson (1965) for the hydrogen atoms were used. Real and imaginary dispersion corrections after Cromer (1965) were applied to the manganese and the phosphorus.

The function minimized by least-squares was $\Sigma w(\Delta F)^2 / \Sigma w F_o^2$. The weighting scheme used throughout the refinement was $w = 0.0$ for $F^2 < \sigma(F^2)$ and $w = 1/\sigma^2(F)$ for all other reflections. $\sigma(F^2)$ was adjusted to weight down intense reflections by adding a fraction, p , of F^2 . $\sigma(F)$ was calculated from $\sigma^2(F^2) = \sigma_o^2(F^2) + (pF^2)^2$; $\sigma_o(F^2) = (Lp)^{-1} \sigma(I)$; $\sigma(F) = F - (F^2 - \sigma(F^2))^{\frac{1}{2}}$, where p for this structure was 0.04.

The following programs for the CDC 6600 and CDC 7600 were used in this structure analysis and interpretation: PICK3, a cell dimension refinement and orientation calculation program; GONIO, a goniometric settings program; INCOR, ORDER, and SORT, general data reduction programs; FORDAP, A. Zalkin's Fourier analysis program; LSLONG and LESQ, our unpublished modifications of the Ganzel-Sparks-Trueblood least-squares program; DISTAN and DISMAT, crystallographic bond distance and angle calculation programs; DATLOK, D.J. St. Clair's unpublished weighting scheme analysis program; LSPLAN, our modified version of the least-squares planes program from the University of Pittsburgh; ORTEP, C. Johnson's (1965) molecular crystallographic plotting program; and LIST1 and LISTAP, data presentation programs.

17. B. B. Cunningham, J. Chem. Educ., 36, 31 (1959).
18. L. H. Bates, Modern Magnetism, 4th Edn., (Cambridge University Press, London, 1961).
19. L. Stein, Inorg. Chem., 3, 995 (1964).
20. P. Handler and C. A. Hutchinson, J. Chem. Phys., 37, 1210 (1956).
21. C. A. Hutchinson, Tung Tsang and B. Weinstock, J. Chem. Phys., 37, 555 (1962).
22. M. E. Hendricks, E. R. Jones, Jr., J. A. Stone and D. G. Karraker, J. Chem. Phys. 55, 2993 (1971).
23. J. L. Green, (Ph.D. Thesis), University of California, Lawrence Radiation Laboratory Report, UCRL-16516 (1965).
24. J. R. Peterson, (Ph.D. Thesis), University of California, Lawrence Radiation Laboratory Report, UCRL-17875 (1967).
25. J. C. Copeland, (M.S. Thesis), University of California, Lawrence Radiation Laboratory Report, UCRL-17718 (1967).
26. D. K. Fujita, (Ph.D. Thesis), University of California, Lawrence Radiation Laboratory Report, UCRL-19507 (1969).
27. T. C. Parsons, private communication (1969).
28. D. K. Smith, University of California, Lawrence Radiation Laboratory Report, UCRL-7196 (1963).
29. A. C. Larson, R. B. Roof, Jr., and D. T. Cromer, Los Alamos Scientific Laboratory Report, LA-3335 (1965).
30. D. E. Williams, Ames Laboratory Report, IS-1052 (1964).
31. L. E. J. Roberts and A. J. Walker, Physico-Chimie du Protactinium, (Editions du Centre National de la Recherche Scientifique, Paris, 1966), p. 51.

no peaks on the map large enough to be unidentified non-hydrogen atoms. Two cycles of full-matrix least-squares refining coordinates of all atoms, anisotropic thermal parameters for phosphorus and manganese, isotropic thermal parameters for all other non-hydrogen atoms, and three sets of isotropic thermal parameters (cage, methyl, phenyl) for the hydrogens, gave an $R_1 = 0.078$.

Comparisons of F_{obs} and F_{calc} at this time showed 55 data with large discrepancies. These reflections were remeasured using the same method as before and were incorporated into the data set after scaling to allow for a slight deterioration of the crystal over the refinement period.

All non-hydrogen atoms were then allowed to refine with anisotropic thermal parameters, and the individual isotropic thermal parameters of the hydrogens were allowed to refine. The refinement proceeded in two blocks, with the anion refined in one block and the cation in the other, while the scale factor was refined in both. Finally, two cycles of full-matrix least-squares on all atoms brought the shifts of all parameters to less than two percent of their calculated standard deviations and the refinement was considered complete. The final R_1 was 0.042 for 2862 reflections, the weighted R-value ($R_2 = \Sigma w(\Delta F)^2 / \Sigma w F_o^2$) was also 0.042, and the standard deviation of an observation of unit weight was 1.12. A difference Fourier map showed no peak greater than $0.21 \text{ e}^-/\text{\AA}^3$. The observed F values and the final differences are given in Table 1. The final atomic parameters are given in Table 2 with their standard deviations as calculated by the least-squares program.

Table 1 (2 pages)

Observed Structure Factors, Standard Deviations and
Final Differences for $(\text{C}_6\text{H}_5)_3\text{PCH}_3^+$ $(\text{B}_6\text{C}_2\text{H}_8)\text{Mn}(\text{CO})_3^-$.

All values multiplied by 10.0. An * indicates that the
reflection was given zero weight in least-squares.

Table 2

Coordinates and Thermal Parameters.^{a,b}

(a) Standard deviations of the least significant digit(s) are given in parentheses.

(b) The form of the temperature factor (B's in Å²) is:

$$T = \exp(-.25(B_{11}h^2a^{*2} + B_{22}k^2b^{*2} + B_{33}l^2c^{*2} + 2B_{12}hka^{*}b^{*} + 2B_{13}hla^{*}c^{*} + 2B_{23}klb^{*}c^{*}))$$

for the anisotropic atoms and
 $T = \exp(-B \sin^2 \theta / \lambda^2)$ for the isotropic atoms.

ATOM	K	Y	Z	811	822	833	812	813	823
HN	.30036(5)	.44937(5)	.17203(5)	4.395(3)	4.51(3)	5.97(3)	.73(2)	1.06(2)	.68(2)
H(1)	.3772(5)	.3177(5)	.2897(5)	5.0(3)	9.1(4)	5.0(3)	2.7(3)	2.2(2)	3.8(3)
H(2)	.3231(5)	.2421(4)	.0886(5)	5.5(2)	4.7(2)	5.2(2)	1.2(2)	.5(2)	.2(2)
H(3)	.4255(5)	.2026(6)	.2001(6)	6.7(3)	6.4(3)	6.2(3)	2.4(2)	1.5(3)	2.9(3)
H(4)	.5329(5)	.3362(5)	.2857(5)	5.0(2)	9.1(3)	5.0(2)	2.1(2)	.3(2)	2.4(2)
H(5)	.4865(5)	.2671(5)	.1072(5)	5.9(3)	7.9(3)	5.3(3)	3.0(2)	1.4(2)	1.3(2)
H(6)	.5015(4)	.4474(5)	.1832(4)	4.4(2)	6.5(3)	5.2(2)	.6(2)	1.4(2)	1.8(2)
C(1)	.4608(4)	.4503(4)	.3048(4)	4.7(2)	6.0(2)	3.6(2)	.7(2)	.5(2)	.7(2)
C(2)	.3959(4)	.3546(4)	.0604(4)	5.5(2)	6.7(2)	3.8(2)	2.2(2)	1.4(2)	1.4(2)
C(11)	.0498(3)	.3310(2)	.1695(3)	4.8(1)	5.4(1)	9.0(2)	.3(1)	2.3(1)	.6(1)
C(4)	.1486(4)	.3785(3)	.1895(3)	5.0(2)	4.0(2)	5.1(2)	1.0(2)	1.1(2)	.1(1)
C(22)	.2914(3)	.0865(3)	.3520(3)	6.5(2)	5.7(2)	8.8(2)	~1.1(1)	3.4(1)	~1.2(1)
C(5)	.2952(3)	.5924(4)	.2812(4)	4.5(2)	5.4(2)	5.5(2)	~2(2)	2.0(2)	.5(2)
C(3)	.1931(4)	.5405(3)	~.0393(3)	13.1(3)	10.5(2)	7.3(2)	6.1(2)	3.1(2)	4.5(2)
C(6)	.2330(4)	.5039(4)	.0435(4)	7.6(3)	5.9(2)	5.1(2)	2.6(2)	1.8(2)	1.4(2)
P	~.3372(10)	.0804(10)	.2972(8)	3.71(4)	4.19(4)	3.07(4)	.70(3)	.96(3)	.69(3)
C(3)	~.2272(5)	.1414(5)	.1748(5)	5.0(2)	4.5(3)	4.8(2)	2.1(2)	1.3(2)	1.4(2)
C(7)	~.0363(3)	.2194(3)	.4275(3)	4.0(2)	3.7(2)	3.7(2)	.7(1)	1.5(1)	1.1(1)
C(8)	.0931(3)	.2334(4)	.4674(4)	4.4(2)	4.5(2)	4.4(2)	.7(2)	1.1(2)	.6(2)
C(9)	.1670(4)	.3359(4)	.5687(4)	4.4(2)	5.6(2)	5.3(2)	~3(2)	.8(2)	1.0(2)
C(10)	.1122(5)	.4242(4)	.6300(4)	6.9(3)	4.3(2)	4.4(2)	~1.0(2)	1.4(2)	.8(2)
C(11)	~.0143(5)	.4125(4)	.5916(4)	7.0(2)	3.8(2)	5.0(2)	.9(2)	2.8(2)	.5(2)
C(12)	~.0894(4)	.3103(3)	.4898(4)	4.7(2)	4.7(2)	5.1(2)	.9(2)	2.0(2)	1.3(2)
C(13)	~.0441(3)	~.0176(3)	.2357(3)	3.8(2)	3.9(2)	3.9(2)	.4(1)	.7(1)	.6(1)
C(14)	~.0564(4)	.1386(4)	.2431(3)	4.7(2)	4.7(2)	3.6(2)	.6(2)	1.3(2)	1.3(2)
C(15)	.0159(4)	~.2164(4)	.1922(4)	6.0(3)	4.0(2)	4.0(2)	1.4(2)	1.1(2)	1.2(2)
C(16)	.1016(4)	~.1727(4)	.1386(4)	6.7(2)	5.3(2)	4.8(2)	2.6(2)	1.8(2)	.8(2)
C(17)	.1149(4)	~.0529(4)	.1326(4)	5.4(2)	5.7(2)	5.1(2)	1.6(2)	2.7(2)	1.2(2)
C(18)	.0425(4)	.0251(4)	.1805(3)	5.6(2)	3.7(2)	4.4(2)	1.1(2)	1.7(2)	1.1(2)
C(19)	~.2502(3)	.0020(3)	.3499(3)	5.5(2)	4.1(2)	3.9(2)	.4(2)	.5(1)	.3(1)
C(20)	~.2393(4)	.0277(4)	.4776(4)	3.6(2)	4.6(2)	4.4(2)	.2(2)	.7(2)	.7(2)
C(21)	~.3266(4)	~.0390(4)	.5131(5)	4.6(2)	5.8(2)	5.4(2)	.5(2)	2.1(2)	1.6(2)
C(22)	~.4359(4)	~.1260(4)	.4268(5)	4.9(2)	6.5(2)	7.4(3)	~2(2)	2.9(2)	2.1(2)
C(23)	~.4392(4)	~.1538(4)	.3000(5)	4.6(2)	6.8(3)	7.1(3)	~1.5(2)	~1.5(2)	~0(2)
C(24)	~.3521(4)	~.0873(4)	.2611(5)	5.2(2)	7.0(3)	4.8(2)	~2(2)	1.5(2)	.3(2)
H(1)	.338(3)	.309(3)	.356(4)	7.0(10)					
H(2)	.237(3)	.172(3)	.024(3)	6.2(8)					
H(3)	.427(3)	.111(4)	.203(4)	7.3(10)					
H(4)	.625(3)	.349(3)	.359(4)	7.4(9)					
H(5)	.538(3)	.225(3)	.047(4)	7.7(10)					
H(6)	.574(3)	.320(3)	.183(3)	5.7(8)					
H(7)	.468(3)	.516(3)	.380(3)	5.4(9)					
H(8)	.377(3)	.304(3)	~.016(4)	6.6(11)					
H(9)	.167(4)	.188(4)	.143(4)	7.3(12)					
H(10)	~.263(3)	.189(3)	.205(3)	5.5(10)					
H(11)	~.276(3)	.084(3)	.116(3)	4.5(10)					
H(12)	.129(3)	.172(3)	.426(3)	3.9(8)					
H(13)	.254(4)	.344(3)	.596(4)	6.9(11)					
H(14)	.164(3)	.489(3)	.694(4)	5.9(10)					
H(15)	~.054(3)	.468(3)	.622(3)	4.5(9)					
H(16)	~.169(2)	.303(2)	.466(2)	2.1(7)					
H(17)	~.111(3)	~.105(2)	.274(3)	2.4(9)					
H(18)	.002(3)	~.294(3)	.191(3)	5.0(9)					
H(19)	.149(3)	~.224(3)	.109(3)	5.1(9)					
H(20)	.176(3)	~.024(3)	.097(3)	4.4(8)					
H(21)	.046(3)	.100(3)	.170(3)	3.1(7)					
H(22)	~.174(3)	.084(3)	.537(3)	2.6(7)					
H(23)	~.319(3)	~.024(3)	.597(4)	5.9(10)					
H(24)	~.483(3)	~.175(3)	.456(3)	5.8(9)					
H(25)	~.506(4)	~.210(4)	.236(4)	7.4(11)					
H(26)	~.356(3)	~.056(3)	.183(3)	4.4(9)					

D. Results and Discussion

The crystal structure shows that the compound consists of discrete cationic and anionic species. A listing of the intramolecular bond distances in both the anion and the cation is given in Table 3.

1. Triphenylmethylphosphonium (TMP) ion.

The cation exhibits a pseudo-threefold axis of symmetry along the P-C(3) bond, the phenyl groups being tilted so as to form a propeller (Figure 1). The environment of the phosphorus is very closely tetrahedral, the range of the C-P-C bond angles being from 108.9° to 109.9° . The P-C(3) bond distance is shorter by three standard deviations than the average of the other three P-C bonds ($1.779(5)$ Å vs. $1.794(3)$ Å), a difference that is barely significant. The hydrogens on the methyl group are staggered with respect to the carbons on the phosphorus as expected.

The phenyl groups are very closely planar. Least-squares planes through the six carbons in each phenyl gave distances to these carbons less than the standard deviation in the atomic positions. Two of the planes have the P-C bond bent slightly ($d_{\text{P-plane}} = 0.025$ Å) but significantly out of the plane.

All C-H distances are normal for x-ray crystallographic studies, the average C-H bond length uncorrected for thermal motion being 0.89 Å. The average C-C bond length for the phenyl groups is $1.374(2)$ Å, in good agreement with published values. There is a measurable discrepancy between the length of the bonds from the meta- to the para-carbons (e.g. C(9)-C(10), Figure 1) which average to 1.363 Å (range: $1.358 - 1.368$ Å)

Table 3

Intra-molecular Bond Distances (Å).^a

Anion			Cation		
Atoms	Distance		Atoms	Distance	
Mn - C(1)	2.043(4)		P - C(3)	1.779(5)	
Mn - C(2)	2.047(4)		P - C(7)	1.796(3)	
Mn - B(1)	2.356(5)		P - C(13)	1.797(3)	
Mn - B(2)	2.349(5)		P - C(19)	1.788(3)	
Mn - B(6)	2.227(5)		C(7) - C(8)	1.380(5)	
Mn - C(4)	1.765(4)		C(8) - C(9)	1.376(5)	
Mn - C(5)	1.771(4)		C(9) - C(10)	1.364(6)	
Mn - C(6)	1.780(5)		C(10) - C(11)	1.358(6)	
C(1) - B(1)	1.573(7)		C(11) - C(12)	1.381(5)	
C(1) - B(4)	1.607(6)		C(12) - C(7)	1.379(5)	
C(1) - B(6)	1.571(6)		C(13) - C(14)	1.373(5)	
C(2) - B(2)	1.565(6)		C(14) - C(15)	1.381(5)	
C(2) - B(5)	1.609(6)		C(15) - C(16)	1.368(6)	
C(2) - B(6)	1.589(6)		C(16) - C(17)	1.362(5)	
B(1) - B(2)	2.122(7)		C(17) - C(18)	1.378(5)	
B(1) - B(3)	1.683(8)		C(18) - C(13)	1.383(5)	
B(1) - B(4)	1.743(7)		C(19) - C(20)	1.385(5)	
B(2) - B(3)	1.695(7)		C(20) - C(21)	1.376(5)	
B(2) - B(5)	1.764(7)		C(21) - C(22)	1.358(6)	
B(3) - B(4)	1.668(8)		C(22) - C(23)	1.367(6)	
B(3) - B(5)	1.667(8)		C(23) - C(24)	1.382(6)	
B(4) - B(5)	1.885(7)		C(24) - C(19)	1.385(5)	
B(4) - B(6)	1.971(7)		H(9) - C(3)	0.94(4)	
B(5) - B(6)	1.971(7)		H(10) - C(3)	0.86(4)	
C(4) - O(1)	1.162(4)		H(11) - C(3)	0.83(3)	
C(5) - O(2)	1.161(4)		H(12) - C(8)	0.92(3)	
C(6) - O(3)	1.157(4)		H(13) - C(9)	0.93(4)	
H(1) - B(1)	1.00(4)		H(14) - C(10)	0.89(4)	
H(2) - B(2)	1.10(4)		H(15) - C(11)	0.87(3)	
H(3) - B(3)	1.05(4)		H(16) - C(12)	0.85(3)	
H(4) - B(4)	1.12(4)		H(17) - C(14)	0.85(3)	

Table 3 (continued)

Anion		Cation	
Atoms	Distance	Atoms	Distance
H(5) - B(5)	1.07(4)	H(18) - C(15)	0.86(3)
H(6) - B(6)	1.06(3)	H(19) - C(16)	0.85(3)
H(7) - C(1)	0.92(3)	H(20) - C(17)	0.94(3)
H(8) - C(2)	0.90(4)	H(21) - C(18)	0.88(3)
		H(22) - C(20)	0.88(3)
		H(23) - C(21)	0.89(4)
		H(24) - C(22)	0.96(4)
		H(25) - C(23)	0.92(4)
		H(26) - C(24)	0.86(3)

 (a) Standard deviations appear in parentheses.

Figure 1

Conformation and labling of the cation. Perspective view roughly down the pseudo-threefold axis. Thermal ellipsoids have been scaled to enclose 50% probability. Hydrogen atoms have been given an artificial isotropic thermal parameter of $2.0 \text{ \AA}^2$.

XBL 724-666

and the other carbon-carbon bonds which average to 1.380 Å (range: 1.373 - 1.385 Å). The C-C-C angles also show a variation dependent on position, the angle adjacent to the phosphorus being reduced by about 1.5° with respect to the other angles. (Average for adjacent angles is 118.8° , range 118.4° to 119° ; average for all other angles 120.2° , range 119.8° to 120.6° , average for all angles is 120.0° .)

The results from this study compare well to the results of Fritchie (1966) on another TMP salt. The average values for all chemically different bonds are within the sum of their standard deviations of each other, though Fritchie's study is not well enough refined to confirm or deny the small differences noted here. The primary difference between the two TMP ions in the two studies is that in this study the ion assumes the expected propeller configuration, whereas packing forces distort it considerably in Fritchie's study.

2. The $(B_6C_2H_8)Mn(CO)_3$ anion.

This work confirms the structure proposed for both the $B_6C_2H_8^{2-}$ ligand and for the anion as a whole (George and Hawthorne, 1969; Hawthorne and Pitts, 1967).

The anion possesses an almost exact non-crystallographic mirror plane running through the Mn, B(3) and P(6) (Figure 2). Variations from true mirror symmetry for the carborane cage atoms, including hydrogens, are all less than twice the standard deviations of their positions as calculated by least-squares. The carbons of the carbonyl groups also conform well to the mirror

Figure 2

Conformation and labling of the anion. Perspective view parallel to the non-crystallographic mirror plane.

XBL 724-661

plane, but the associated oxygen atoms deviate from mirror symmetry by as much as five times the standard deviations of their positions. In addition, the carbonyl groups are very closely radial from the manganese. These observations confirm Hawthorne's spectroscopic conclusions (George and Hawthorne, 1969), and the slight deviations of the oxygens can undoubtedly be laid at the feet of crystal packing forces.

Hawthorne's description of the manganese + cage system is a tricapped trigonal prism, where the capping atoms are the two carbons and the unique boron. In the numbering system used here, the capping atoms are C(1), C(2) and B(3). The prism can best be seen in Figure 2, with the three parallel edges of the prism coming out of the paper between B(1) and B(4), B(2) and B(5) and Mn and B(6). The cage, excluding the Mn, can be described another way, by defining two planes, plane 1 consisting of B(3), B(4) and B(5), and plane 2 containing C(1), C(2), B(1) and B(2). Plane 1 and plane 2 are tipped with respect to one another and plane 1 is slipped back in order to accommodate B(6), causing the "front" end of plane 1 to rise, and B(3) to hang over the "back", giving the whole cage the appearance of a sled (Figure 3). This view of the $B_6C_2H_8^{2-}$ cage can most easily be related to the geometry of the $B_7C_2H_9^{2-}$ dicarbollide ion (St. Clair, 1969), while Hawthorne's view is more closely related to the $B_7H_7C_2(CH_3)_2^{2-}$ polyhedral species (Koetzle, et.al., 1968).

As shown in Figure 3, the manganese ion lies closest to the two carbon atoms. It is slightly displaced towards B(1) and

Figure 3

Stereoscopic view of the anion. View direction is approximately perpendicular to the non-crystallographic mirror plane through the anion. Temperature ellipsoids have been scaled to enclose 20% probability. Hydrogens have been given artificial thermal parameters of $2.0 \text{ \AA}^2$.

XBL 724-664

B(2) but its position is such that the Mn-B(6) distance is shorter by 0.12 Å than the Mn-B(1) and the Mn-B(2) distances even though B(6) is offset from plane 2 by 0.43 Å in the direction opposite the manganese. The distance from the manganese to plane 2 is 1.578 Å. The Mn-C and Mn-B distances are well within the ranges reported for other metallocarboranes, the Mn-C distances being about average and the spread between the Mn-C and Mn-B bond lengths being much greater than normal (St. Clair, 1969) due to the fact that the base of the ligand is highly non-planar when all atoms bonded to the manganese are considered.

The interatomic distances between borons and carbons are summarized in Table 3, and show a great deal of variation, both among themselves and with respect to average values reported for other carborane complexes. Note that the mirror plane is again in evidence, as all related bond distances are within two standard deviations of each other. The average B-C distance, 1.59 Å, is more than 0.10 Å shorter than the averages reported for icosahedral fragments (St. Clair, 1969), but agrees reasonably with the averages for the $B_7C_2H_9^{2-}$ fragment to which this is related by the removal of one atom. It is also in good agreement with the average B-C distances in non-icosahedral carboranes. (Hart and Lipscomb, 1968; Koetzle, *et.al.*, 1968) The average B-B distance in this structure is 1.82 Å, in good agreement with average values in other structures. This is misleading, however, due to the fact that the average includes three anomalously long distances; B(1)-B(2), B(4)-B(6) and B(5)-B(6). When these are excluded from the average, it drops to 1.73 Å,

which is significantly shorter than the average B-B distance reported for both the polyhedral carboranes and dicarbollide fragments but is close to the apical B-B distances reported by Koetzle, et.al. (1968).

The B(1)-B(2) distance of 2.122 Å is longer than any B-B distance I have been able to reference. However, the species referenced above ($B_7H_7C_2(CH_3)_2^{2-}$) does contain long B-B bonds of length around 2.0 Å. Koetzle's anion is related to this one by substitution of the $Mn(CO)_3$ for B2 (his notation) and the removal of the CH_3 groups from his carbons and their replacement by hydrogens. The long bond lengths observed in his study correspond in location to the long bonds observed here.

The C-H and B-H bond distances are normal for x-ray studies, and all bonds to hydrogen are directed approximately radially, as expected.

3. Packing

The packing of the cations and the anions can be seen in Figure 4. The cations pack in rows on the c axis with the phenyl groups of one cation interleaving with the phenyl groups of the cation related by a center of symmetry at $0, 0, \frac{1}{2}$ such that each phenyl of one cation is inclined at a relatively high angle to a phenyl ring of the other in a staggered configuration. There is a relatively close approach of parallel phenyl groups across the center of symmetry at $\frac{1}{2}, 0, \frac{1}{2}$, and a very close approach of parallel phenyl groups across the center of symmetry at $0, \frac{1}{2}, \frac{1}{2}$, with a distance C(11)-C(11)' of only 3.29 Å (not shown in Figure

4). Other inter-molecular contacts of less than the sum of the van der Waals radii are: O(1)-C(3) (methyl carbon) 3.45 Å, O(2)-C(12) 3.24 Å, and O(2)-C(22) 3.31 Å. The carbonyl-methyl contact can be seen near the origin in Figure 4 and the other two carbonyl contacts can be drawn from the anion that is closest to the front in Figure 4 and the carbonyl that is directed down and away from that manganese to the two phenyls involved. (One is hidden behind the carbonyl group and the other is attached to the phosphorus on the upper portion of the marked c axis.)

The cations thus can be considered to form sheets in both the ac and the bc planes, with close contacts near $\frac{1}{2}, 0, \frac{1}{2}$ and $0, \frac{1}{2}, \frac{1}{2}$, but with relatively open spaces near $\frac{1}{2}$ on the b axis and the a axis. The anions form zig-zag chains through the openings in this lattice of cation sheets. As can be seen from Figure 4, the carbonyl ends of the anion are opposed across the center of symmetry at $0, \frac{1}{2}, 0$ and the dicarbollide ends of the anion are opposed across the centers at $\frac{1}{2}, 0, 0$, $\frac{1}{2}, \frac{1}{2}, 0$ and $\frac{1}{2}, \frac{1}{2}, \frac{1}{2}$. None of these oppositions force close contact between atoms from one anion and atoms from another.

Figure 4

Stereoscopic view of the contents of the unit cell. The methyl carbons and the carbons on the dicarbollide ion are solid black. The phosphorus atoms have been stippled. Labeled axes are positive from the origin. Hydrogen atoms and the cations in the b direction have been omitted for clarity, as have the phenyl groups of the foremost cation.

XBL 724-723

E. References

- Cromer, D.T. (1965). Acta Cryst., 18, 17.
- Cromer, D.T. and Weber, J.T. (1965). Acta Cryst., 18, 104.
- Fritchie, C.J., Jr. (1966). Acta Cryst., 20, 107.
- George, A.D. and Hawthorne, M.F. (1969). Inorg. Chem., 8(8), 1801.
- Hart, H. and Lipscomb, W.N. (1968). Inorg. Chem., 7(6), 1070.
- Hawthorne, M.F. and Pitts, A.D. (1967). J. Am. Chem. Soc., 89, 7115.
- Johnson, C.K. (1965) Oak Ridge National Report-3794, Revised, oak Ridge, Tenn.
- Koetzle, T.F., Scarbrough, F.E. and Lipscomb, W.N. (1968). Inorg. Chem., 7(6), 1076.
- St. Clair, D.J. (1969). Ph.D. Thesis, University of California, Berkeley, Calif. UCRL-18910.
- Stewart, R.F., Davidson, E.R. and Simpson, W.T. (1965). J. Chem. Phys., 42, 3175.

II. Determination of the Crystal Structure of $\text{XeF}_5^+ \text{AsF}_6^-$.

A. Introduction

Several compounds of the general empirical formula F_{11}XeM have been reported. Crystal structures have been determined for $\text{M} = \text{Pt}, \text{Ru}$ (Bartlett, et.al., 1967; Morrell, 1971) and involve the XeF_5^+ cation and an octahedrally coordinated MF_6^- anion. This study was undertaken to determine the structure of F_{11}XeAs , and shows a structure consistent with previous work in being formulable as $\text{XeF}_5^+ \text{AsF}_6^-$.

B. Experimental Procedure

The compound $\text{XeF}_5^+ \text{AsF}_6^-$ was prepared by Manfred Wechsberg according to the method previously described (Bartlett and Wechsberg, 1971). Single crystals suitable for crystallographic study were obtained with great difficulty by sublimation of the material in quartz capillaries. Precession photographs taken of several crystals indicated that they were monoclinic, and the systematic absences ($h0\ell$, $\ell \neq 2n$, $0k0$, $k \neq 2n$) indicated uniquely space group $P2_1/c$ (C_{2h}^5 , No. 14).

A crystal of approximate dimensions 0.18 mm x 0.16 mm x 0.04 mm in the form of a thin plate perpendicular to b and bounded by (0 1 0), (0 $\bar{1}$ 0), ($\bar{1}$ 0 1), ($\bar{1}$ 0 $\bar{1}$) and the capillary wall was mounted on a Picker/Nuclear four-circle diffractometer automated by a DEC PDP 8I computer. Accurate cell dimensions were obtained by refinement on the measured positions of 12 reflections carefully centered in χ , ω , and 2θ ($\lambda = 0.709261 \text{ \AA}$) using the FACS-1 least-squares program on the PDP 8I. The crystal data are summarized in Table 1.

Table 1.

Crystal Data

$a = 5.886(3) \text{ \AA}$	Space Group - $P2_1/c$
$b = 16.564(10) \text{ \AA}$	$Z = 4$
$c = 8.051(4) \text{ \AA}$	$d_c = 3.51 \text{ gm/cm}^3$
$\beta = 91.564(35)^\circ$	$\mu = 90.8 \text{ cm}^{-1}$
$V = 784.6 \text{ \AA}^3$	

Intensity data were collected using a θ - 2θ scan technique with Mo K α radiation monochromatized with a graphite bent-crystal monochrometer ($2\theta_m = 12.02^\circ$). A base width for the scan of 1.1°

was used which was centered on the predicted Mo $K\alpha_1$ position and was augmented on the high 2θ side to allow for the α_1 - α_2 separation. The peak was scanned at a speed of $1^\circ/\text{min}$ and background measurements were counted for 10 seconds at each end of the scan (all angles are degrees 2θ). Attenuators were automatically inserted in the diffracted beam any time the count rate exceeded 7000 cps and the peak was rescanned and backgrounds taken with the attenuators in place. Reflections were collected in the hemisphere $+h, +k, \pm l$ for $2\theta = 0^\circ$ - 50° ($\sin\theta/\lambda = 0.596$). Three reflections, (3 0 0), (0 10 0) and (0 0 4), were monitored periodically during the data collection and showed anisotropic deterioration to between 43% and 63% of their original values. Due to the extreme difficulty of obtaining crystals suitable for x-ray crystallographic work, it was decided to use the data set anyway and a single set of factors was applied to all reflections to correct for the decay in the absence of any detailed knowledge of the decay anisotropy. 3071 reflections were recorded, not including multiple records of the standards. Of the 3071, 367 were measured with zero intensity and 713 in all had $I < \sigma(I)$. I and $\sigma(I)$ were calculated using the formulae:

$$I = C - \frac{t_c}{2t_b} (B_1 + B_2) \quad \sigma^2(I) = C + \frac{t_c^2}{4t_b^2} (B_1 + B_2)$$

where C is the total counts taken in scan time t_c , and B_1 and B_2 are the background counts, each taken for time t_b . Due to the high absorption coefficient (Table 1.), an absorption correction was made to the data. The faces named above were well-defined and

easily measurable, and the curved face of the crystal against the capillary wall was approximated by five faces ((1 0 1), (2 0 1), (1 0 0), (2 0 $\bar{1}$), (1 0 $\bar{1}$)). The correction calculated and then applied to the data varied by a factor of 2.5 from minimum to maximum. After application of the absorption correction the intensities were averaged to give a set of 1382 unique reflections. The standard deviations of the intensities were calculated by $\sigma(I) = (\sigma^2(I_{\text{ave}}) + p^2 I^2)^{\frac{1}{2}}$, where p (here = 0.05) is a factor introduced to reduce the weights given to intense reflections, which are more subject to systematic errors, and $\sigma(I_{\text{ave}})$ was calculated from counting statistics or from the deviation of the individual reflections from the average, whichever was larger. Lorentz and polarization factors were then applied to convert the measurements to F^2 values. Of the final 1382 reflections, 926 had $F^2 > \sigma(F^2)$.

Atomic scattering factors of Cromer and Waber (1965) were used for the preliminary refinements, using the dispersion terms of Cromer (1965). Towards the end of the refinement the more recent scattering factors of Doyle and Turner (1968) and the dispersion terms of Cromer and Liberman (1970) were used ($\Delta f'_{\text{Xe}} = -.684 \text{ e}^-$, $\Delta f''_{\text{Xe}} = 1.958 \text{ e}^-$, $\Delta f'_{\text{As}} = -.03 \text{ e}^-$, $\Delta f''_{\text{As}} = 2.007 \text{ e}^-$). Our least-squares program minimizes the function $\Sigma w(\Delta F)^2 / \Sigma w F_o^2$.

All calculations were performed on either our CDC 6600 or CDC 7600 computer. The following programs were used in the solution of this structure: RLFACS, a program for reduction of raw data from the Picker automatic diffractometer. HORSE, our modification of Hamilton's GONO9 program for absorption correction; INCOR and SORT, general data reduction programs; FORDAP,

A. Zalkin's Fourier analysis program; LSLONG and LESQ, modifications of the Ganzel-Sparks-Trueblood least-squares program; DATLOK, D.J. St. Clair's unpublished weighting scheme analysis program; DISTAN and DISMAT, crystallographic bond distance and angle programs, the latter of which calculates standard deviations using the correlation matrix from least-squares; ORTEP, C. Johnson's (1965) thermal ellipsoid plotting program; LSPLAN, our modification of the least-squares planes program from the University of Pittsburgh; and LISTI and LISTAP, data presentation programs.

C. Solution and Refinement of the Structure

The volume of the unit cell is $784.6 \text{ \AA}^3$. Zachariassen's (1949) criterion of $18 \text{ \AA}^3$ per fluorine yielded 43.6 fluorines per unit cell, which was consistent with the formulation of F_{11}XeAs and four formula units in the monoclinic unit cell. The assumption that all atoms were in general positions in the cell ($4c: x, y, z; -x, -y, -z; -x, \frac{1}{2}+y, \frac{1}{2}-z; x, \frac{1}{2}-y, \frac{1}{2}+z$) was subsequently confirmed by the refinement.

A three-dimensional Patterson map was constructed, and Xe-Xe, As-As and Xe-As peaks were identified, giving the positions of the two heavy atoms in the asymmetric unit. Four cycles of least-squares refinement of their coordinates and isotropic thermal parameters and a scale factor brought the conventional R value ($R_1 = \Sigma |\Delta F| / \Sigma |F_o|$) to 0.29. A Fourier map phased on the F values exhibited 11 major peaks other than those which appeared at the Xe and As positions. These peaks were arranged octahedrally around the As peak and in a square pyramid around the Xe peak. The 11 peaks were labeled as fluorines and refined with isotropic thermal parameters in least squares, while xenon and arsenic were allowed anisotropic thermal parameters. Then two cycles of full-matrix least squares during which the fluorines were allowed to refine anisotropically brought R_1 to 0.114 for 811 reflections, using the weighting scheme $w = 0.0$ for $I < 2\sigma(I)$, $w = 1/\sigma^2(F)$ for all others.

A difference Fourier map taken at this point showed several large peaks at unreasonable positions to be atoms. The standard deviation of an observation of unit weight, S, was equal to 3.00

and an analysis of the weighted ΔF 's showed that the intense reflections were given too much weight. An option in LESQ was used that allows for modification of $\sigma(F^2)$ (and hence of w) by adding a fraction, p , of F^2 to $\sigma(F^2)$. Several values of p were input and least-squares refinements run to check the effect and to determine what fraction should be used. Values from 0.05 to 0.20 were tried, and analyses of the weighted differences run for each value. (Note that this fraction p is in addition to the 0.05 added during the data reduction.) The weighting scheme used for these least-squares runs and for the final refinement was: $w = 0.0$ for $F^2 < \sigma(F^2)$ rather than $F^2 < 2\sigma(F^2)$, increasing the number of reflections included in the refinement from 811 to 925. R_1 decreased from $p = 0.0$ to $p = 0.10$ and then increased gradually. The weighted sum of differences decreased continuously of course, as did the weighting of the intense reflections. A general point of diminishing returns was reached around $p = 0.10$, and this value of p was used in the final refinement.

The final cycle of least-squares, refining all atoms anisotropically, gave $R_1 = 0.116$ for 925 reflections, an R_1 including zero weighted reflections of 0.143, a weighted R value ($R_2 = \Sigma w(\Delta F)^2 / \Sigma w F_o^2$) of 0.128 and $S = 1.57$. The shifts of all parameters were less than 2% of their estimated standard deviations and the refinement was considered complete. The difference Fourier map phased from the results of the last cycle of least-squares still showed many peaks greater than $1 \text{ e}^-/\text{\AA}^3$, the largest being associated with smearing of electron density in the b direction around the xenon and the arsenic. These peaks and

the high value of p are indications of the fact that the intensity decay noted during data collection could not be accurately accounted for in an isotropic manner.

The structure factors, their standard deviations and the final differences are given in Table 2. The final coordinates and thermal parameters for the atoms are given in Table 3.

Table 2.

Observed Structure Factors, Standard Deviations and Final Differences for $\text{XeF}_5^+ \text{AsF}_6^-$. An * indicates that the value was given zero weight in least-squares.

Table 3.

Coordinates and Thermal Parameters for $\text{XeF}_5^+ \text{AsF}_6^-$.^{a,b}

(a) Estimated standard deviations of the least significant digit(s) are given in parentheses.

(b) The form of the temperature factor (B_{ij} in units of $\text{\AA}^2$) is:

$$T = \exp(-.25(B_{11}\underline{h}^2\underline{a}^2 + B_{22}\underline{k}^2\underline{b}^2 + B_{33}\underline{l}^2\underline{c}^2 + 2B_{12}\underline{hk}\underline{a}*\underline{b}* + 2B_{13}\underline{hl}\underline{a}*\underline{c}* + 2B_{23}\underline{kl}\underline{b}*\underline{c}*).$$

ATOM	X	Y	Z	B11	B22	B33	B12	B13	B23
XE	-1957(3)	.1101(1)	.2501(2)	5.76(9)	8.41(11)	5.05(8)	.22(9)	.01(5)	.22(6)
AS	.6068(6)	.1067(2)	.7009(3)	4.16(15)	5.47(15)	3.03(12)	.07(15)	-.31(11)	-.43(10)
F1	.081(4)	.1888(10)	.126(2)	6.3(11)	6.6(9)	4.7(8)	.5(9)	.0(7)	1.2(7)
F2	.456(4)	.1590(11)	.187(2)	6.3(12)	7.6(11)	6.4(10)	-3.1(10)	1.6(9)	1.9(8)
F3	-.179(4)	.0607(11)	.050(2)	6.4(11)	8.1(10)	3.6(7)	1.1(9)	-1.5(7)	-.8(7)
F4	-.109(3)	.0830(16)	.269(2)	3.5(9)	14.7(18)	5.3(9)	-2.5(11)	.4(7)	1.3(10)
F5	.161(4)	.1892(11)	.405(2)	7.4(13)	7.9(10)	4.4(8)	1.7(10)	-2.5(8)	-1.3(7)
F6	.474(4)	.0869(10)	.509(2)	7.7(12)	7.6(9)	2.9(6)	.2(9)	-2.7(7)	-.7(6)
F7	.377(3)	.1625(12)	.756(2)	3.9(9)	7.4(11)	7.1(10)	2.1(9)	1.2(8)	-1.3(7)
F8	.486(3)	.0176(11)	.772(2)	4.6(9)	7.0(9)	3.9(7)	-1.2(8)	-.0(6)	1.2(6)
F9	.834(3)	.0457(9)	.643(2)	3.8(8)	5.8(8)	5.7(9)	1.3(7)	1.5(7)	.6(7)
F10	.734(4)	.1875(10)	.620(3)	7.2(13)	5.9(9)	7.4(11)	-1.5(9)	1.0(10)	-.6(7)
F11	.741(4)	.1226(11)	.888(2)	8.1(13)	9.8(12)	2.8(7)	.7(10)	-2.0(8)	-1.9(7)

D. Results and Discussion

The structure of $F_{11}XeAs$, like the reported structures of $F_{11}XePt$ (Bartlett, *et.al.*, 1967) and $F_{11}XeRu$ (Morrell, 1971), is well divided into cations (XeF_5^+) and anions (AsF_6^-).

In the cation, the xenon has five fluorines closely bound to it in a distorted square pyramid (Figure 1). The xenon is displaced from the plane of the basal fluorines by 0.31 Å in the direction of the lone pair electrons. The distances from the xenon to the fluorines are also shown in Figure 1; the standard deviation for all distances being ± 0.02 Å. The average of the $Xe-F_{\text{basal}}$ bond distances is 1.82(1) Å, as compared to the $Xe-F_{\text{ap}}$ distance of 1.76(2) Å, indicating that the shortening of the apical $Xe-F$ bond is significant in terms of the standard deviations of the distances. The average $F_{\text{ap}}-Xe-F_{\text{bas}}$ angle is 80.4° and ranges from 79.2° to 82.4° . The $F_{\text{bas}}-Xe-F_{\text{bas}}$ angles range from 86° to 90° (Table 4).

In addition to the five closely bound fluorines, the xenon is coordinated to three fluorines belonging to the AsF_6^- anion. These bridging fluorines show $Xe\cdots F$ distances from 2.65 Å to 2.83 Å (Figure 1). The angles $F_{\text{ap}}-Xe\cdots F_{\text{br}}$ range from 139° to 148° avoiding the central non-bonded electron pair. The conformation of the cation and the bridging fluorines found in this structure agrees well with the results of other studies involving the XeF_5^+ group (*loc. cit.* ; Bartlett and Leary, 1972; Bartlett and Sladky, 1970; Burbank and Jones, 1970).

The AsF_6^- anion shows a slight trigonal distortion from octahedral symmetry. Most of the distortion can be described

Figure 1.

Configuration, labeling and bond distances for $\text{XeF}_6^+ \text{AsF}_6^-$.
 Perspective view roughly perpendicular to the bc plane showing
 the environment around the center of symmetry at $\frac{1}{2}, 0, \frac{1}{2}$.
 Standard deviation of all bond lengths shown is $\pm 0.02 \text{ \AA}$.

XBL 724-665

Table 4.

Selected Interatomic Bond Angles (deg) for $\text{XeF}_5^+ \text{AsF}_6^-$.^a

Atoms	Angle	Atoms	Angle
F1 - Xe - F2	80(1)	F6 - As - F7	90(1)
F1 - Xe - F3	80(1)	F6 - As - F8	87(1)
F1 - Xe - F4	82(1)	F6 - As - F9	89(1)
F1 - Xe - F5	79(1)	F6 - As - F10	90(1)
F1 - Xe - F6	141(1)	F7 - As - F8	92(1)
F1 - Xe - F8 ^b	139(1)	F7 - As - F10	92(1)
F1 - Xe - F9 ^b	148(1)	F7 - As - F11	92(1)
F2 - Xe - F3	89(1)	F8 - As - F9	85(1)
F2 - Xe - F5	89(1)	F8 - As - F11	91(1)
F3 - Xe - F4	86(1)	F9 - As - F10	89(1)
F4 - Xe - F5	90(1)	F9 - As - F11	89(1)
		F10 - As - F11	91(1)

(a) Estimated standard deviations in parentheses.

(b) Atom related by center of symmetry at $\frac{1}{2}, 0, \frac{1}{2}$.

by a displacement of the As atom from the center of gravity of the F atoms. The bond distances to F6, F8 and F9 average to 1.744 Å (range: 1.740 - 1.751 Å) while those to F7, F10 and F11 average 1.694 Å (range: 1.676 - 1.707 Å). Similar distortions of bond lengths in the anion for bridging fluorines has been observed in $\text{XeF}_5^+ \text{RuF}_6^-$ (Morrell, 1971). The primary distortion of the fluorines themselves from an ideal octahedron can be described by a displacement of F8 towards F9.

Pairs of cations and pairs of anions are linked into a ring by the bridging fluorines, each xenon accepting one fluorine from one arsenic and two fluorines from the arsenic related by a center of symmetry (Figure 1).

The packing of the rings in the unit cell is shown in Figure 2. They stack in well-separated sheets perpendicular to the b axis and stack in herringbone fashion in the sheets. There are no inter-ionic contacts of less than the expected F-F van der Waals distance of 2.70 Å except in the cases of the bridging fluorines.

It should be noted that an earlier version of Figure 1 is reproduced in Bartlett and Sladky's (1970) review. Due to further refinements after the time that particular illustration was drawn, some of the bond distances have changed significantly.

Figure 2.

Stereoscopic view of the contents of the unit cell, showing packing of the rings. Axes labeled ^e are positive from the origin.

XBL 724-724

F. References

- Bartlett, N., Einstein, F., Stewart, D.F. and Trotter, J. (1967) J. Chem. Soc. (A), 1190.
- Bartlett, N. and Leary, K. (1972). Unpublished communication.
- Bartlett, N. and Sladky, F.O. (1970). "The Chemistry of Krypton, Xenon and Radon" UCRL-19658, Lawrence Berkeley Laboratory, Berkeley, Calif.
- Bartlett, N. and Wechsberg, M. (1971). Z. anorg. u. allgem. Chemie, 385, 5-17.
- Burbank, R.D. and Jones, G.R. (1970). Science, 168, 248.
- Cromer, D.T. (1965). Acta Cryst., 18, 17.
- Cromer, D.T. and Liberman, D. (1970). J. Chem. Phys., 53(5), 1891.
- Cromer, D.T. and Waber, J.T. (1965). Acta Cryst., 18, 104.
- Doyle, P.A. and Turner, P.S. (1968). Acta Cryst., A24, 390.
- Johnson, C.K. (1965). Oak Ridge National Report-3794, Revised, Oak Ridge, Tenn.
- Morrell, B.K. (1971). M.S. Thesis, University of California, Berkeley, Calif. LBL-156.
- Zachariasen, W.H. (1949). Acta Cryst., 2, 390.

III. The Crystal and Molecular Structure of 1,3-diphenyl-1,3-propanedione enol (HDPP).

A Introduction

As a preface to a study of the complexes of the alkaline-earth metals with β -diketones, the structure of 1,3-diphenyl-1,3-propanedione (HDPP), one of the ligands to be used in the study, was determined to provide a basis for comparison of the complexed and uncomplexed molecules. HDPP could have occurred as the keto form (I) but was found to exist solely as the enol (II) in the crystalline state.

Knowledge of the crystal form was also necessary to allow identification of the material when there were suspicions about the synthesis of the chelates. The HDPP frequently crystallized out by itself and did not react properly, and the knowledge of the crystal habits and space group enabled quick determination of whether or not this had happened.

(I) keto

(II) enol

B. Experimental Procedure

The HDPP was supplied to us by Eastman Kodak Co. (Stock No. 2197) in the form of a pink crystalline powder. The material was recrystallized from various organic solvents, giving clear colorless or orange crystals which were otherwise indistinguishable. The usual habit was a very thin, flexible blade, but crystals suitable for x-ray work were obtained by crystallization from aqueous ethanol. Comparison of powder, precession and Weissenberg photographs of various crystals indicated that there was no dependence of crystal structure on the solvent used for recrystallization.

Weissenberg and precession photographs established the systematic absences $Ok\bar{l}$, $k \neq 2n$; $h0\bar{l}$, $l \neq 2n$; $hk0$, $h \neq 2n$ and an orthorhombic unit cell. These absences uniquely determine the space group $Pbca$ (P_{2h}^{15} , No. 61). Preliminary cell dimensions were determined by measurement of the Weissenberg and powder patterns.

A crystal of approximate dimensions 0.19 mm x 0.23 mm x 0.12 mm was then mounted and aligned on a Picker/Nuclear four-circle diffractometer automated with a DEC PDP 8I computer and disc. Twelve reflections at high 2θ were carefully centered in X , ω and 2θ using $Cu K\alpha_1$ ($\lambda = 1.54051 \text{ \AA}$) radiation and used to refine the cell dimensions and orientation matrix for the crystal using the FACS-1 least-squares program on the PDP 8I. The final cell dimensions and their standard deviations are $a = 10.853(1) \text{ \AA}$, $b = 24.441(1) \text{ \AA}$, $c = 8.755(1) \text{ \AA}$, $\alpha = \beta = \gamma = 90^\circ$. The density was measured by flotation in aqueous zinc bromide solution and

gave the result $d_o = 1.22 \text{ gm/cm}^3$ which is low compared to the calculated density based on the refined structure of 1.28 gm/cm^3 . The error is due to the difficulty in wetting the crystals with an aqueous solution.

Intensity data were collected using graphite monochromatized ($2\theta_m = 26.45^\circ$) copper radiation and a θ - 2θ scan technique. The x-ray tube takeoff angle was 2° . Peaks were scanned at a speed of $1^\circ/\text{min}$ from a position 0.6° below the calculated $K\alpha_1$ peak to a position 0.6° above the predicted $K\alpha_2$ peak. Backgrounds were counted for 10 seconds with both crystal and detector stationary at positions offset from each end of the scan range by 0.7° (all degrees 2θ). All data in the quadrant $+\underline{h}, +\underline{k}, \pm \underline{l}$ were collected for two ranges of 2θ , the first range from 0° - 100° and the second from 100° - 120° . Aluminum attenuators were automatically inserted in the diffracted beam whenever the counting rate exceeded 7000 cps and the reflection and backgrounds remeasured with the attenuators in place. Three reflections, the $(0\ 4\ 0)$, $(0\ 2\ 3)$ and $(3\ 3\ 1)$, were monitored frequently during the data collection and exhibited a gradual decrease in intensity amounting to $3\frac{1}{2}\%$ over the entire data collection period. Random short-term variations in the intensities of the standards were of the same order of magnitude.

The data were originally punched onto paper tape by the PDP 8I, and was transferred to magnetic tape and converted to net intensities by program PAPER written for the CDC 6600. Net intensities and their standard deviations were calculated according to the following formulae:

$$I = C - \frac{t_c}{2t_b}(B_1 + B_2) \quad \sigma(I) = C + \frac{t_c^2}{4t_b^2}(B_1 + B_2)$$

where C is the total count over scan time t_c , and B_1 and B_2 are the background counts, each taken for time t_b . PAPER also applied a decay correction to the intensities.

3775 reflections were measured, not counting multiple measurements of the standards. Of these, 449 were space group extinctions, 320 had $I = 0.0$ (i.e. C less than the adjusted background) and another 481 had $I < \sigma(I)$.

The absorption coefficient, μ , was 6.9 cm^{-1} , and the average μR was 0.10. No absorption correction was applied to the intensities.

Lorentz and polarization corrections were applied to the data and equivalent reflections were averaged to produce $1/24$ unique reflections. A fraction, p , of the intensity was added to the standard deviations of all reflections to reduce the weighting of the more intense reflections, which tend to be affected by systematic errors. The formula used was:

$$\sigma^2(I) = \sigma^2(I_{\text{ave}}) + p^2 I^2$$

where $\sigma(I_{\text{ave}})$ was calculated from counting statistics or from scatter around the mean, whichever was larger. The fraction p was originally set to 0.02, but later increased to 0.04 to improve the weighting.

Our least-squares program minimizes the function $\sum w(\Delta F)^2 / \sum w F_o^2$. The weighting scheme used gave $w = 0.0$ to those reflections with $I < \sigma(I)$ and $w = 1/\sigma^2(F)$ to all other reflections.

Scattering factor tables used for the non-hydrogen atoms were those of Cromer and Waber (1965) and the hydrogen atom scattering factor tables used were those of Stewart, Davidson and Simpson (1965). Anomalous dispersion terms for C and O were ignored.

Programs written for the CDC 6600, other than those already mentioned, used in the solution, refinement and presentation of this structure were: INCOR, EDIT and SORT, general data reduction programs; WILSON, a program for applying Wilson's statistics to data and calculation of normalized structure factors; REL, R.E. Long's (1965) program for application of Sayre's relationships for centric structures; FORDAP, A. Zalkin's Fourier analysis program; LSLONG, our unpublished modification of the Ganzel-Sparks-Trueblood least-squares program; HFINDR, a program for prediction of hydrogen positions in a structure; LSPLAN, our modification of the least-squares planes program from the University of Pittsburgh; DATLOK, D.J. St. Clair's unpublished weighting scheme analysis program; ORTEP, Johnson's (1965) thermal ellipsoid plotting program; and LIST1 and LISTAP, data presentation programs.

C. Determination and Refinement

Normalized structure factors, E_h , were calculated using Wilson's (1942) method. The phases of the highest 209 E values ($E > 1.50$) were determined using Long's (1965) sign determination program, which iteratively applies the equation $\text{sign}(E_h) = \text{sign}(\sum_k E_k E_{h-k})$. The fixed positive phases of the (6 1 7), (3 6 4) and (1 3 2) defined the origin, and the phases of an additional four reflections, (10 10 1), (0 16 4), (7 14 1) and (8 6 3), were held fixed for each of the 16 computer runs in which they were allowed to have all combinations of positive and negative phases. A consistency index defined as:

$$C = \frac{\sum_h \sum_k E_h E_k E_{h-k}}{\sum_h \sum_k |E_h E_k E_{h-k}|}$$

was calculated for each combination. Of the 16 possibilities, one had $C = 0.75$, three others which differed only in the sign of one of the starting reflections had $C = 0.71$ - 0.72 , and all others had $C = 0.36$ - 0.56 .

A Fourier map was calculated using the E's phased from the run with the highest consistency index. The top seventeen peaks on the Fourier plot gave an unambiguous picture of the expected geometry of the HDPP molecule. Positions for the atoms were taken from the Fourier plot; they were labeled and assigned atomic scattering factors on the basis of the known molecular structure. The positions of all 17 atoms were then refined by least-squares with individual isotropic thermal parameters, using all the data. After five iterations of least-squares the R value ($R_1 = \sum |\Delta F| /$

$E|F_o|$) was 0.12. Inspection of the reflections with high E values showed that none of the phases of these reflections had changed from those predicted.

All atoms were then allowed to refine with anisotropic thermal parameters, and three cycles of least-squares gave $R_1 = 0.091$. A Difference Fourier showed peaks located near all the predicted positions for hydrogens in the enolic form of the β -diketone. The positions of the peaks were put into least squares and allowed to refine with isotropic thermal parameters. Three cycles of refinement brought R_1 to 0.049, but several of the hydrogen atoms refined to chemically unreasonable positions. The worst offender, H(2), was placed in its calculated position and three more cycles of least-squares performed during which it refined to its previous poor position and the R_1 value remained unchanged.

Analysis of the observed and calculated F's showed that F_o/F_c was consistently less than 1.0 for high intensity data. A correction factor for secondary extinction of the form $F = F_o(1.0 + kI)$, where I is the raw intensity, was applied, with $k = 2.08 \times 10^{-7}$. Three least-squares cycles reduced R_1 to 0.042, but the hydrogens still did not improve. The factor p mentioned above was also varied and its final value of 0.04 arrived at without correcting the misplaced hydrogens.

All hydrogens except H(7) (Figure 1) were then placed in positions calculated on the basis of perfect phenyl geometry for all three rings and a C-H distance of 1.0 Å. In the next few cycles of least-squares the positions of the hydrogens were held fixed, and the coordinates and thermal parameters of the non-hy-

drogen atoms and the thermal parameters of the hydrogens allowed to refine. Three cycles of least-squares reduced R_1 drastically to 0.029, with a weighted R value ($R_2 = \Sigma w(\Delta F)^2 / \Sigma w F_o^2$) of 0.037. The positional parameters of the hydrogens were then allowed to refine along with all the other parameters in least-squares. R_1 dropped to 0.027 and R_2 to 0.033 and all hydrogens refined to chemically reasonable positions. The standard deviation of an observation of unit weight was 1.04 and all parameter shifts were minute fractions of their standard deviations. A difference Fourier phased after the final cycle showed no peaks greater than $0.20 \text{ e}^-/\text{\AA}^3$ in either the positive or negative sense and no discernable correlations between the peaks and the structure. The refinement was regarded as complete at this point.

The observed structure factors and the final calculated structure factors are given in Table 1. The final coordinates and thermal parameters of the atoms are given in Table 2.

Table 1.

Comparison of F_{obs} and F_{calc} . An * indicates that the reflection was given zero weight in least-squares.

Table 2.

Final coordinates and thermal parameters for HDPP.^{a,b}

- (a) Standard deviations of the least significant digit(s) are given in parentheses.
- (b) The form of the temperature factor (B 's in units of $\text{\AA}^2$) is:
- $$T = \exp(-.25(B_{11}h^2a^{*2} + B_{22}k^2b^{*2} + B_{33}l^2c^{*2} + 2B_{12}hka^*b^* + 2B_{13}hla^*c^* + 2B_{23}k\ell b^*c^*))$$
- for anisotropic, and
- $$T = \exp(-B \sin^2\theta/\lambda^2)$$
- for isotropic thermal parameters.

ATOM	X	Y	Z	B11	B22	B33	B12	B13	B23
C(1)	.1032(2)	.29449(8)	.5250(3)	5.80(11)	4.24(9)	5.26(11)	1.01(8)	-.68(10)	-.50(9)
C(2)	.0515(3)	.25721(9)	.6239(3)	7.90(15)	4.31(10)	5.85(13)	1.19(10)	-1.23(12)	.22(10)
C(3)	-.0694(3)	.26209(8)	.6673(3)	7.84(16)	4.48(11)	6.35(15)	-.30(11)	-.30(12)	.67(10)
C(4)	-.1394(2)	.30493(8)	.6143(3)	5.53(12)	4.88(10)	6.93(14)	-.26(9)	-.23(11)	.92(10)
C(5)	-.0885(2)	.34302(8)	.5167(2)	4.70(10)	4.06(8)	5.73(12)	.11(7)	-.87(9)	.55(9)
C(6)	.0332(2)	.33030(6)	.4698(2)	4.56(9)	3.56(8)	4.42(10)	.07(7)	-.97(8)	-.42(7)
C(7)	.0918(2)	.37756(7)	.3641(2)	3.79(8)	4.46(8)	4.58(10)	.13(7)	-.72(8)	-.89(8)
C(8)	.0456(2)	.42909(7)	.3336(2)	4.05(9)	4.39(9)	4.54(11)	-.04(7)	.31(8)	-.10(8)
C(9)	.1056(2)	.46456(7)	.2309(2)	3.91(8)	4.17(9)	4.30(10)	-.73(7)	-.46(8)	-.79(8)
C(10)	.0568(2)	.51892(7)	.1881(2)	3.78(8)	4.24(8)	3.78(10)	-.99(7)	-.41(7)	-.26(8)
C(11)	-.0549(2)	.53778(7)	.2435(3)	4.97(10)	4.21(9)	4.87(11)	-.86(8)	.54(9)	.19(8)
C(12)	-.1001(2)	.58819(8)	.1993(3)	5.79(11)	4.31(9)	6.11(12)	-.24(9)	.86(11)	.72(9)
C(13)	-.0348(2)	.62036(9)	.0991(3)	6.16(12)	4.75(10)	5.62(13)	-.76(9)	-.28(10)	.78(10)
C(14)	.0756(2)	.66221(9)	.0438(3)	5.96(12)	5.74(12)	5.32(13)	-1.77(10)	-.01(10)	1.22(10)
C(15)	.1214(2)	.55183(9)	.0862(2)	4.42(9)	5.90(10)	4.60(11)	-1.16(8)	.22(9)	.07(9)
O(1)	.1927(1)	.36110(5)	.2998(2)	4.46(6)	5.38(6)	7.24(9)	.67(5)	.50(6)	-.34(7)
O(2)	.2081(1)	.49000(6)	.1691(2)	4.27(6)	5.74(7)	6.85(9)	-.35(6)	.95(6)	-.45(6)
H(1)	.1908(18)	.2930(7)	.4012(23)	5.8(5)					
H(2)	.1033(17)	.2275(8)	.6619(24)	7.3(6)					
H(3)	-.1084(18)	.2378(9)	.7315(27)	7.3(6)					
H(4)	-.2281(19)	.3080(8)	.6438(24)	6.9(6)					
H(5)	-.1362(17)	.3743(8)	.4622(22)	5.3(5)					
H(6)	-.0284(16)	.4409(6)	.3826(20)	4.3(4)					
H(7)	.2197(25)	.4019(11)	.2259(32)	11.1(8)					
H(8)	-.1048(15)	.5154(7)	.3075(20)	4.6(4)					
H(9)	-.1827(20)	.5996(8)	.2379(24)	6.6(6)					
H(10)	-.0643(19)	.0589(9)	.0860(28)	7.9(6)					
H(11)	.1250(19)	.6236(8)	-.0240(26)	6.8(6)					
H(12)	.1963(16)	.5362(7)	.0451(22)	5.0(4)					

XBL 707-1308

D. Results and Discussion

HDPP exists entirely in the enolic form of the β -diketone in the crystal. The enolic hydrogen, H(7), refined well and there were no indications of tetrahedrally bound hydrogens on the α -carbon, C(8). Inspection of the distances from the two oxygens to H(7) and of the C-C and C-O distances in the diketone ring (Figure 1) indicate that while the hydrogen is bonded almost symmetrically, and the ring is almost entirely delocalized, there is a slight but significant preference for the O(1)-H(7)---O(2) form of the diketone.

The two phenyl rings are planar to within two standard deviations of their positional parameters. The diketone ring, however, is significantly distorted in such a way that the bending of the phenyl rings from the plane of the diketone is taken up in part by the distortion. The angles between the three planes as calculated by the least-squares planes program are -17.0° between the first phenyl (C(1)-C(6)) and the diketone, -4.0° between the second phenyl ring and the diketone, and 13.1° between the two phenyl rings.

The average bond distances in the phenyl rings (C-C = $1.330(2)$ Å, C-H = 0.98 Å) are in good agreement with literature values. The C-C bonds adjacent to the diketone ring are significantly longer, and the C-C bonds furthest from the diketone significantly shorter than the average, and the interior angle nearest the diketone is significantly less than the theoretical 120° , presumably due to steric interaction. The effects of the intra-molecular close contacts also show up in the C-C-C angles

Figure 1.

Perspective view of one molecule of HDPP, showing labeling and bond distances. View is approximately perpendicular to the ab plane, positive b to the right and a down. Standard deviations of the C-C and C-O bonds are ± 0.002 - 0.003 Å. Standard deviations of distances to the hydrogens are ± 0.02 Å. Thermal ellipsoids have been scaled to include 50% probability and H atoms have been given an artificial thermal parameter of 1.0 Å^2 .

XBL 7CV-1346

Figure 2.

Interatomic bond angles in HDPP. Standard deviations of angles involving carbon and oxygen only is $\pm 0.2^\circ$. Standard deviations of angles involving hydrogen is $\pm 1^\circ$.

centered on C(7) and C(9) (Figure 2), which are greater than the expected 120° due to close contacts between phenyl hydrogens H(5) and H(8) and the diketone hydrogen H(6).

The bonds between the diketone system and the phenyl rings show normal shortening due to conjugation between the π -bonded rings.

The HDPP molecules are packed parallel to one another in interleaving sheets. Sheets made up of tilted HDPP molecules lying parallel to one another run in the z direction alternating inclination of the tilt with sheets to either side in the x and y directions. There are close ($< 3.50 \text{ \AA}$) contacts between HDPP molecules in both the x and the z directions (Figure 3; Table 3). There are no close contacts in the y direction. Some of the contacts in the x direction might be considered as hydrogen bonds from the phenyl carbons to the carbonyl oxygens, but the indications are not that strong given the equally close contacts between atoms which lack the possibility of hydrogen-bond formation.

Just before writing began on this thesis a previous study of the crystal structure of HDPP was brought to my attention through an article on the crystal structure of ω -(p-toluoyl)-acetophenone enol (related to HDPP by a para methyl on one of the phenyl rings) (Kato, 1971) which referred to a structure done by D.E. Williams (1966). The literature search was rechecked and it was found that while the paper was listed in the Author Index for Chem. Abstracts, it was not listed under any of the chemical names for HDPP, nor under its empirical formula, nor under the general

Figure 3.

Stereoscopic view of the contents of one unit cell. Labeled axes are positive from the origin. Oxygen atoms have been shaded and hydrogen atoms given artificial thermal parameters for clarity. Close inter-molecular contacts between heavy atoms are shown.

JCL 708-1106

Table 3.

Intermolecular contacts of less than 3.50 Å between heavy atoms.

Atom 1	Atom 2	Distance	Molecule of Atom 2 related to molecule of Atom 1 by:
C(6)	C(12)	3.48 Å	$\underline{x}$, $1-\underline{y}$, $1-\underline{z}$
O(1)	C(14)	3.41 Å	$\frac{1}{2}-\underline{x}$, $1-\underline{y}$, $\frac{1}{2}+\underline{z}$
O(2)	C(11)	3.44 Å	$\frac{1}{2}+\underline{x}$, $\underline{y}$, $\frac{1}{2}-\underline{z}$
O(2)	C(13)	3.47 Å	$-\underline{x}$, $1-\underline{y}$, $-\underline{z}$

heading "Structure Determination".

Williams' results are substantially identical to those found in this study, though comparison is difficult due to lack of reported standard deviations for his parameters. The present study comes to a better agreement ($R_1 = 0.027$ vs. Williams' $R_1 = 0.059$), and the refinement of this structure included the hydrogen thermal parameters where Williams' refinement did not. Discrepancies in positional parameters as great as ten times the standard deviations of the parameters in the present work are observed between the two studies, but consideration of the distances and angles (for which Williams does give standard deviations) reveals no differences between the two structures that are greater than the sum of the standard deviations for each.

The largest absolute difference between the two structures is in the symmetry of the intra-molecular hydrogen bond. Whereas this study found O-H distances of 1.22(3) Å and 1.28(3) Å from O(1) and O(2) to H(7), Williams finds corresponding distances of 1.18(6) Å and 1.34(6) Å, which would indicate a less delocalized form of the enol. His distances for C-O and C-C bonds within the diketone are not consistently different from those of this study, however, and it is more likely that this difference reflects differences in the refinement techniques and small differences in the data sets than any chemical difference between the structures.

E. References

- Cromer, D.T. and Weber, J.T. (1965). Acta Cryst., 18, 104.
- Johnson, C.K. (1965). Oak Ridge National Report-3794, Revised, Oak Ridge, Tenn.
- Kato, V.K. (1971). Acta Cryst., B27, 2028.
- Long, R.E. (1965). Ph.D. Thesis, University of California, Los Angeles, Calif.
- Stewart, R.F., Davidson, E.R. and Simpson, W.T. (1965). J. Chem. Phys., 42, 3175.
- Williams, D.E. (1966). Acta Cryst., 21, 340.
- Wilson, A.J.C. (1942). Nature (London), 150, 152.

IV. The Crystal and Molecular Structure of bis(dimethylformamido)-bis(1,3-diphenyl-1,3-propanedionato)magnesium. $\text{Mg}(\text{DPP})_2(\text{DMF})_2$

A. Experimental Procedure

The compound was synthesized by combining an aqueous solution of magnesium acetate ($\text{Mg}(\text{OAc})_2$) and an ethanolic solution of 1,3-diphenyl-1,3-propanedione (HDPP) in a strong $\text{NH}_3/\text{NH}_4\text{Cl}$, pH 10, buffer. All materials used were of reagent grade and were not further purified before use. A pale yellow precipitate formed immediately and the mixture was stirred for several hours before the precipitate was filtered and dried in air. The filtrate, on standing, yielded another precipitate which was filtered and shown to be identical to the first. The formal reaction is $\text{Mg}(\text{OAc})_2 + 2\text{HDPP} \xrightarrow{\text{pH } 10} \text{Mg}(\text{DPP})_2 + 2\text{HOAc}$.

Attempts were made to recrystallize the compound from various organic solvents and mixtures. Attempts at recrystallization from ethanol were marginally successful, yielding crystals which decomposed rapidly, even in sealed capillaries. Preliminary measurements on these crystals indicated a triclinic cell, but conflicting results were obtained. These crystals were not used in this structure determination due to their poor properties.

Evaporation in air of dimethylformamide (DMF) solutions of the compound yielded very good crystals which were used throughout the rest of this investigation. These crystals still tended to decompose on exposure to air, but were stable in sealed capillaries and in a desiccator over Drierite. The crystals were also stable in air during one period of very low humidity. This

indicates that moisture in the air is most likely responsible for the decomposition of the crystals. A powder pattern taken of the recrystallized material was different from that of the original precipitate, but over a period of exposure to air, the diffraction lines of the original material slowly reappeared.

Several crystals were mounted in sealed quartz capillaries for study. Weissenberg and precession photographs indicated a monoclinic unit cell with absences hkl , $h + k \neq 2n$ and $h0l$, $l \neq 2n$. These absences are consistent with space groups Cc or $C2/c$, with b as the unique axis.

A crystal was then mounted on a G.E. XRD-5 manual three-circle diffractometer and twelve high-angle reflections were carefully centered using $Cu K\alpha_1$ radiation ($\lambda = 1.54051 \text{ \AA}$). The cell dimensions were refined on the 2 θ measurements only, using program TTHCEL, a modification of a least-squares program supplied to us by H. Hope. TTHCEL refines the cell parameters and 2 θ origin, and minimizes the function $\sum w(d_o^{*2} - d_c^{*2})^2$, where the relative weights are assigned subjectively by the user. The refined cell parameters and their standard deviations were determined to be $a = 16.893(3) \text{ \AA}$, $b = 12.853(2) \text{ \AA}$, $c = 16.927(3) \text{ \AA}$, $\alpha = 90^\circ$, $\beta = 117.085(5)^\circ$. The density was not measured due to difficulty in finding a suitable flotation medium. The calculated density for an empirical formula $Mg(DPP)_2$ gave $Z = 4$, $d_c = 0.96 \text{ gm/cm}^3$ as the most reasonable result. The density based on the refined composition of $Mg(DPP)_2(DMF)_2$ with the observed Z of 4 is $d_c = 1.10 \text{ gm/cm}^3$.

Intensity data were collected from a crystal of approximate dimensions 0.25 mm x 0.25 mm x 0.30 mm sealed in a quartz capillary and mounted on a Picker/Nuclear four-circle diffractometer automated by a DEC PDP 8I computer and disc, using graphite monochromatized ($2\theta_m = 26.36^\circ$) Cu K α radiation. Intensities were collected using a θ - 2θ scan technique at an x-ray tube takeoff angle of 3° . Peaks were scanned at a rate of $1^\circ/\text{min}$ from 0.9° below the predicted K α_1 position to 0.9° above the predicted K α_2 position. Backgrounds were counted for 10 seconds at positions offset 0.6° from each end of the scan interval (all angles in 2θ). Aluminum attenuators were automatically inserted in the diffracted beam whenever the count rate exceeded 7000 cps, and the peak and backgrounds were remeasured with the attenuators in place. The reflections (4 0 0), ($\bar{2}$ 0 2) and (0 4 $\bar{4}$) were monitored periodically during the data collection and exhibited no decay in intensity. All reflections in the quadrant of reciprocal space $+h, +k, \pm l$ were measured out to a 2θ angle of 120° ($\sin\theta/\lambda < 0.562$). 2433 unique reflections were measured, of which 614 had $I < \sigma(I)$ (including $I = 0.0$). Net intensities and their standard deviations were calculated by the formulae:

$$I = C - \frac{t_c}{2t_b} (B_1 + B_2) \quad \sigma(I) = C + \frac{t_c^2}{4t_b^2} (B_1 + B_2)$$

where C is the total counts measured over scan time t_c and B_1 and B_2 are the background counts, each measured for time t_b . No absorption correction was performed ($\mu = 7.7 \text{ cm}^{-1}$). Lorentz and polarization factors were applied to the data to convert them to F^2 values.

The scattering factors of Doyle and Turner (1968) were used for neutral Mg, C and O, together with the real and imaginary dispersion terms of Cromer and Liberman (1970). The hydrogen scattering factors of Stewart, Davidson and Simpson (1965) were used.

Our least-squares program minimizes the function $\sum w(\Delta F)^2 / \sum w F_o^2$. The weighting scheme used throughout the refinement gave zero weight to those reflections where $F^2 < \sigma(F^2)$, and $w = 1/\sigma(F)$ for all others. $\sigma(F)$ was calculated by finite differences from $\sigma(F^2)$ and F^2 by

$$\sigma(F) = F - (F^2 - \sigma(F^2))^{\frac{1}{2}} ; \sigma^2(F^2) = \sigma_o^2(F^2) + (pF^2)^2$$

where p is a factor (here = 0.06) used to reduce the weights of intense reflections, which are more prone to undetected systematic errors.

The following programs, written for our CDC 6600 computer, were also used in the solution and refinement of this structure: MAGPIK, a program for interpretation of raw data from the Picker/DEC system; INCOR, EDIT, and ORDER, general data reduction programs; WILSON, an unpublished program written by Maddox and Maddox for applying Wilson's (1942) statistics to data and calculation of normalized structure factors; REL, R.E. Long's (1965) program for direct determination of centric phases; FORDAP, A. Zalkin's Fourier analysis program; LSLONG, our modification of the Ganzel-Sparks-Trueblood least-squares program; DISMAT, a crystallographic distances and angles program which calculates standard deviations using the correlation matrix from least-squares; ORTEP, C. John-

son's (1965) thermal ellipsoid plotting program; LSPLAN, our modification of the least-squares planes program from the University of Pittsburgh; and LIST1 and LISTAP, data presentation programs.

B. Determination and Refinement

Normalized structure factors, E_h , were calculated using Wilson's (1942) method. Analysis of the average values of E and E^2-1 strongly indicated the centric space group $C2/c$ ($\text{ave}(E) = 0.792$ vs. 0.792 (0.886) and $\text{ave}(E^2-1) = 0.973$ vs. 0.968 (0.736) theoretical values for the centric and (acentric) case). Since the most probable number of molecules in the unit cell was four according to density considerations, this implied that the Mg atom had to lie either on the two-fold axis at $x = 0, z = \frac{1}{4}$ or on a center of symmetry ($0, 0, 0$ or $\frac{1}{2}, \frac{1}{2}, \frac{1}{2}$) in the eight-fold space group. Since the Mg atom, however, was not expected to be "heavy" enough to show up well on a Patterson map, it was decided to use direct methods to solve the structure.

REL, R.E. Long's (1965) sign determination program was used in the attempt to determine the phases of the 316 reflections with $E > 1.50$. The instructions for REL stated that the indices of the reflections in the centered space group had to be converted to indices in a primitive unit cell and gave a matrix for doing so. After discovering the correct matrix for this transformation, a small program was written to convert the hkl of the reflections from the centered cell to a primitive cell and back again. (Later it was shown that the indices did not have to be transformed for input into the program and use of the transformation was stopped.) Two reflections were given arbitrary positive phases to fix the origin of the cell. Four other reflections had their phases fixed during each of the 16 runs in which they were allowed to have all combinations of positive and negative

phases. The program iteratively applies the equation $\text{sign}(E_h) = \text{sign}(\sum_k E_k E_{h-k})$ to determine as many of the remaining phases as possible.

Several early sets of starting reflections gave runs with poor agreement and reflections with undetermined phases. Later sets of starting reflections gave convergence to self-consistent phases for the high-E reflections. However, Fourier maps phased on the E's of the solutions with the highest consistency C,

$$C = \frac{\sum_h \sum_k E_h E_k E_{h-k}}{\sum_h \sum_k |E_h E_k E_{h-k}|}$$

and the least number of iterations to convergence showed numerous peaks, but no sign of a magnesium on a special position or any other portion of a chemically reasonable structure.

The attempt to solve the structure by direct methods was abandoned temporarily and attempts made to fit a model of the structure into the unit cell. The magnesium was forced to lie on one of the special positions, and the DPP ligands were assumed to be planar in concordance with the results of the study of its structure (Section III). The formula $\text{Mg}(\text{DPP})_2$ for the complex was assumed and the magnesium was allowed to be coordinated in a planar or near-tetrahedral manner, depending on the symmetry of the special position. While several models were found that would fit into the cell without having portions of the DPP's intersect point symmetry operations, none of them refined well when put into least-squares. The reason for the failure of the model-fitting attempt was, of course, the fact that the model was poor, as shown by the refinement of the actual structure.

In a return to direct methods, it was decided to choose a starting set with the lowest possible hkl for reasonably high E values rather than those reflections with the highest E's or the most number of interactions. Six starting reflections were chosen where no index was higher than 5 and all E's were greater than 2.20. The solution with the third highest consistency ($C = 0.63$) but a much lower number of iterations than the second highest set, phased an E-map which had a large peak in a general position which was surrounded by smaller peaks in a portion of an octahedron overlaid with further peaks that looked like a portion of a benzene ring which made sense if it was rotated through a nearby two-fold axis. A portion of a DPP was also identifiable. Since the centric solution to a basically non-centric structure can give such overlap, it was decided to attempt refinement of the positions of the peaks thus found in the non-centric space group Cc. A magnesium, five oxygen and four carbon atoms were labeled and allowed to refine for three cycles of full-matrix least-squares, giving an R value ($R_1 = \Sigma |\Delta F| / \Sigma |F_o|$) of 0.49. The x and z coordinates of the Mg atoms were held fixed to determine the origin of the cell. The results of the least-squares were used to phase a non-centric Fourier map which showed the original 10 peaks and 27 others in chemically reasonable positions. These were labeled and allowed to refine with the others to an R_1 of 0.53. Some carbons had thermal parameters refine to negative values (all atoms had isotropic thermal parameters). Another Fourier was phased, which showed 42 major peaks. These

indicated two DPP molecules with the exception of one missing phenyl carbon, and two four-atom "straight" chains. All 42 atoms were allowed to refine for five cycles of least-squares, with more problems with the thermal parameters, and a last Fourier map showed the missing phenyl carbon and two atoms on the chains which confirmed the suspicion that they were molecules of dimethylformamide (DMF). Refinement of all 45 atoms reduced R_1 drastically, to 0.16, and a difference Fourier showed no peaks greater than $0.5 \text{ e}^-/\text{\AA}^3$. Refinement still proceeded slowly.

At this point it was noticed that the $\text{Mg}(\text{DPP})_2(\text{DMF})_2$ complex possessed a non-crystallographic (in Cc) two-fold axis in the y direction and that the z of the Mg atom was very close to 0.25. A shift in the x coordinate of all the atoms was all that was necessary to bring the complex into the centric space group. The poor behavior of the refinement was thus explained, since supposedly unrelated parameters were actually heavily correlated. The structure was fitted to the centric group by changing the x coordinates of all the atoms and averaging the two-fold related positions, and an attempt was made to refine the structure in the centric group. Three cycles of centric full-matrix least-squares refined quickly and with only small shifts to an R_1 of 0.154. The results of the Wilson statistics and the excellent refinement of the structure in the centric space group confirm that choice.

All of the atoms were then allowed to refine with anisotropic thermal parameters, and the R_1 reduced to 0.141. A difference Fourier was phased and the hydrogen atoms of the DPP ligand were

found, but not those of the DMF. The 11 hydrogens were included in least-squares with isotropic thermal parameters and R_1 dropped to 0.078. One of the hydrogens "wandered" and was found again on the difference Fourier along with most of the hydrogens of the DMF molecule. Further refinements showed that the hydrogens of one methyl carbon, C(17), were refining poorly. They were removed and a difference Fourier calculated which showed two peaks around C(17) in reasonable positions. A third hydrogen position was calculated to complete a tetrahedron around the carbon. All three hydrogens were given an isotropic thermal parameter of $12 \text{ \AA}^2$, and they were held fixed during the last three cycles of least-squares, during which all parameter shifts dropped to less than 10% of their standard deviations. The final R_1 was 0.067 for 1817 reflections. The final R_2 ($\sum w(\Delta F)^2 / \sum w F_o^2$) was 0.065 and the standard deviation of an observation of unit weight was 1.30. The data were checked for secondary extinction with negative results.

Values of F_o and the final differences are given in Table 1. The final coordinates and thermal parameters of the atoms are given in Table 2.

Table 1. (2 pages)

F_0 , $\sigma(F_0)$ and final differences for $\text{Mg}(\text{DPP})_2(\text{DMF})_2$ ($\times 10.0$).

Entries marked with an * were given zero weight in least-squares.

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85															

Table 2.

Final coordinates and thermal parameters for $\text{Mg}(\text{DPP})_2(\text{DMF})_2$.^{a,b}

(a) Standard deviations of the least significant digit(s) are given in parentheses.

(b) The form of the temperature factor (B in units of $\text{\AA}^2$) is

$$T = \exp(-.25(B_{11}h^2a^{*2} + B_{22}k^2b^{*2} + B_{33}l^2c^{*2} + 2B_{12}hka^*b^* +$$

$$2B_{13}hla^*c^* + 2B_{23}k lb^*c^*)) \text{ for anisotropic, and}$$

$$T = \exp(-B \sin^2 \theta / \lambda^2) \text{ for isotropic thermal parameters.}$$

ATOM	X	Y	Z	B11	B22	B33	B12	B13	B23
RG	0	.3373(1)	.250	3.68(8)	3.44(8)	4.85(9)	0	2.14(7)	0
O(1)	-.0439(2)	.4524(2)	.1553(2)	3.8(1)	4.1(1)	5.3(1)	-.14(1)	2.0(1)	.5(1)
O(2)	.1176(1)	.3437(2)	.2419(2)	4.0(1)	3.9(1)	5.7(1)	.08(1)	2.6(1)	.3(1)
C(01)	-.0231(3)	.6479(3)	.0215(3)	5.1(2)	4.5(2)	4.8(2)	.3(2)	2.3(2)	-.1(2)
C(02)	-.0672(3)	.7344(4)	-.0270(3)	6.4(3)	5.4(2)	4.7(2)	-.2(2)	2.3(2)	.0(2)
C(03)	-.1258(3)	.7868(4)	-.0080(4)	6.2(3)	5.2(2)	6.4(3)	.9(2)	2.0(2)	1.4(2)
C(04)	-.1402(3)	.7543(4)	.0634(3)	4.7(2)	5.2(2)	7.1(3)	1.4(2)	2.4(2)	.6(2)
C(05)	-.1006(2)	.6647(3)	.1085(3)	4.3(2)	5.2(2)	5.1(2)	.5(2)	2.2(2)	.7(2)
C(06)	-.0398(2)	.6115(3)	.0890(2)	3.5(2)	4.0(2)	4.2(2)	-.2(1)	1.5(1)	.0(1)
C(07)	.0047(2)	.5173(3)	.1419(2)	4.1(2)	3.9(2)	4.1(2)	-.1(1)	1.9(1)	-.3(1)
C(08)	.0974(2)	.5087(3)	.1741(3)	3.7(2)	3.5(2)	5.5(2)	-.1(1)	2.3(2)	.4(2)
C(09)	.1479(2)	.4250(3)	.2216(2)	4.2(2)	3.5(2)	4.9(2)	.1(1)	2.7(2)	-.5(1)
C(10)	.2463(2)	.4253(3)	.2503(2)	3.6(2)	3.7(2)	4.4(2)	.1(1)	2.0(1)	-.1(1)
C(11)	.2917(3)	.3328(3)	.2601(3)	4.3(2)	4.2(2)	6.4(2)	.1(2)	2.4(2)	-.5(2)
C(12)	.3813(3)	.2321(5)	.2873(3)	5.0(2)	6.3(3)	7.5(3)	2.2(2)	3.0(2)	.6(2)
C(13)	.4285(3)	.4226(5)	.3074(3)	3.0(2)	9.2(4)	6.3(3)	.5(2)	2.4(2)	1.0(2)
C(14)	.3860(3)	.5152(5)	.3003(3)	4.3(2)	6.2(3)	7.8(3)	-.1.0(2)	2.0(2)	1.0(2)
C(15)	.2956(3)	.5167(4)	.2719(3)	4.4(2)	4.2(2)	7.1(3)	.1(2)	2.0(2)	.5(2)
O(3)	.0579(2)	.2222(2)	.3473(2)	4.5(1)	4.3(1)	5.5(1)	.7(1)	2.4(1)	1.0(1)
C(16)	.1330(3)	.1852(3)	.3767(3)	5.1(2)	5.5(2)	4.9(2)	1.0(2)	2.5(2)	.7(2)
N(1)	.1725(2)	.1250(3)	.4443(2)	5.6(2)	5.4(2)	4.6(2)	1.5(1)	1.6(1)	.8(1)
C(17)	.1281(4)	.0975(5)	.4965(4)	11.1(4)	10.5(4)	9.5(4)	3.2(3)	5.9(3)	4.9(3)
C(18)	.2608(6)	.082(1)	.4699(7)	7.3(4)	11.3(7)	10.4(6)	4.5(5)	3.0(4)	3.3(5)
H(01)	.016(3)	.615(3)	.008(2)	5.1(10)					
H(02)	-.057(2)	.754(3)	-.075(3)	6.5(11)					
H(03)	-.153(3)	.838(4)	-.044(3)	7.8(13)					
H(04)	-.187(4)	.788(4)	.069(3)	9.5(15)					
H(05)	-.109(2)	.641(3)	.161(2)	5.1(9)					
H(06)	.127(2)	.560(3)	.158(2)	4.7(9)					
H(07)	.258(2)	.275(3)	.238(2)	4.8(9)					
H(08)	.406(3)	.272(4)	.290(3)	7.9(14)					
H(09)	.491(3)	.427(3)	.331(3)	7.1(11)					
H(10)	.416(3)	.573(3)	.314(3)	5.7(11)					
H(11)	.264(2)	.573(3)	.267(2)	4.3(9)					
H(12)	.162(3)	.197(3)	.347(3)	5.6(11)					
H(13)	.291(4)	.100(5)	.533(5)	11.8(23)					
H(14)	.251(4)	.021(5)	.462(5)	9.4(25)					
H(15)	.298(5)	.125(6)	.449(5)	13.9(28)					
H(16)	.181	.095	.573	12.000					
H(17)	.078	.140	.487	12.000					
H(18)	.1088	.0205	.4882	12.000					

C. Results and Discussion

The structure is separated into well-defined complex groups with composition $\text{Mg}(\text{DPP})_2(\text{DMF})_2$. The magnesium has a formal charge of +2 and each of the DPP ligands has a formal charge of -1; the DMF molecules are formally neutral. The magnesium is in a special position on the two-fold axis, and is coordinated by two symmetry-related DPP and two symmetry-related DMF molecules. The two DPP ligands lie above and to either side of the magnesium in such a way that their mean planes form a propeller around the axis. The planes of the two DMF molecules are so oriented that there is no indication of a propeller in their configuration (Figure 1).

The six oxygen atoms coordinating the magnesium lie on the corners of a slightly distorted octahedron. The distances from the magnesium to the oxygens in the DPP ligand are identical to within a standard deviation, at 2.055(2) Å and 2.057(3) Å, while the Mg-O distance to the DMF is slightly, but significantly, longer at 2.095(3) Å. Both of these distances fall well within the range for typical Mg-O distances in the literature (2.0 Å - 2.15 Å), with the Mg-O(DPP) distance being shorter than the Mg-O(DMF) as expected from electrostatic interactions. The angles of the octahedron are also distorted (Table 3). The angles O(2)-Mg-O(2)' is concave in the positive y direction and the O(1)-Mg-O(1)' and O(3)-Mg-O(3)' planes are twisted out of perpendicularity to this plane (Figure 2).

The distances between oxygens coordinating the Mg range from 2.81 Å to 3.07 Å, with the minimum distance being between the

Figure 1.

Stereoscopic view of the complex, showing configuration and labelling. The two-fold crystallographic axis runs vertically in the plane of the drawing. Hydrogens have been omitted for clarity. Thermal ellipsoids are scaled to include 50% probability.

XBL 722-136

Figure 2.

Oxygen coordination of the Mg, showing distortion of the octahedron. Projection down the two-fold axis onto the ac plane.

Table 3.

Bond angles (deg) around magnesium in $\text{Mg}(\text{DPP})_2(\text{DMF})_2$.^a

Atom(1) -Mg- Atom(2)	Angle
O(1)	O(1) ^b 87.9
O(1)	O(3) 174.0
O(1)	O(3) ^b 91.3
O(2)	O(1) 86.3
O(2)	O(1) ^b 90.4
O(2)	O(2) ^b 175.4
O(2)	O(3) 87.8
O(2)	O(3) ^b 95.5
O(3)	O(3) ^b 90.2

(a) Standard deviation of all angles is $\pm 0.1^\circ$.

(b) Related by two-fold axis through Mg.

two oxygens in the same DPP ligand. There are several close C--O and C--C contacts within the complex, specifically across the two-fold axis, due to the close bonding of the oxygens to the magnesium. Analysis of the least-squares planes through portions of the molecules involved show that the DPP is pushed away from the two-fold axis so that the Mg atom is 0.58 Å above the least-squares plane of the diketone and that the DMF molecules are spread away from the axis so that Mg is above that molecular plane by 0.29 Å. (The distances given are the perpendicular distances to the plane involved. "Above" implies that the distance from the plane to the Mg has a component in the positive y direction.)

The two phenyl groups, the diketone ring and the DMF molecule are each planar to within five standard deviations of the coordinates of the atoms involved. The two phenyl rings are twisted with respect to the diketone plane by 47.6° and 30.9° for the first (C(1)-C(6)) and second (C(10)-C(15)) rings respectively. The first phenyl ring is also bent so that C(6) is out of the diketone plane.

The average C-C distance in the phenyl rings is 1.376 Å. Other bond distance in the DPP and DMF molecules are as expected (Section III, Sutton, et.al., 1958). The O-C-C and C-C-C angles in the diketone are spread (ave. 125°) as expected, to give the ligand a bigger bite. In the phenyl groups the interior angle nearest the diketone is significantly less than the ideal 120° (118.8° and 117.5° for the first and second phenyls respectively) adjusting to the steric interaction between the phenyls and the

diketone. The phenyl C-C bond distances to the para carbon atoms are systematically shorter than the C-C bond distances to the other carbons, also possibly due to steric interactions (Table 4).

The complexes themselves are packed in an interlocking manner shown in Figure 3. The phenyl rings extending away from the two-fold axis (C(10)-C(15)) project into the open space between the other two phenyl rings of a C centering related complex and the DMF molecules of the complex related to the second by a translation in y. The phenyl rings directed up the two-fold axis with respect to the Mg project into the area just below the DMF molecules of the complex above it. The phenyls also project into the relatively open area to the side of a complex related to it by a center of symmetry, and the DMF ligands into the open side of a complex related by the n glide at y = $\frac{1}{4}$.

There are only three short ($< 3.50 \text{ \AA}$) inter-complex contacts between non-hydrogen atoms. They are: from O(1) to a C(18) of the complex related to the first by the n glide at y = $\frac{1}{4}$, O(1)-C(18) distance $3.39 \text{ \AA}$, from O(2) to a C(2) of the complex related by the center of symmetry at O, $\frac{1}{2}, \frac{1}{2}$, O(2)-C(2) distance $3.48 \text{ \AA}$, and from C(4) to a C(4) of the complex related by the n glide at y = $\frac{3}{4}$, C(4)-C(4) distance $3.34 \text{ \AA}$. None of these contacts can be considered as hydrogen bonds.

Table 4.

Intra-molecular and intra-complex distances (Å).^a Mg(DPP)₂(DMF)₂

Atom 1	Atom 2	Distance	Atom 1	Atom 2	Distance
C(01)	C(02)	1.381(6)	C(01)	H(01)	0.90(4)
C(02)	C(03)	1.372(7)	C(02)	H(02)	0.94(4)
C(03)	C(04)	1.369(7)	C(03)	H(03)	0.88(5)
C(04)	C(05)	1.376(6)	C(04)	H(04)	0.94(5)
C(05)	C(06)	1.393(5)	C(05)	H(05)	1.01(4)
C(06)	C(01)	1.377(5)	C(08)	H(06)	0.94(4)
C(06)	C(07)	1.489(5)	C(11)	H(07)	0.90(4)
C(07)	O(1)	1.261(4)	C(12)	H(08)	0.87(5)
C(07)	C(08)	1.408(5)	C(13)	H(09)	0.95(5)
C(08)	C(09)	1.381(5)	C(14)	H(10)	0.87(4)
C(09)	O(2)	1.277(4)	C(15)	H(11)	0.88(3)
C(09)	C(10)	1.504(5)	C(16)	H(12)	0.86(4)
C(10)	C(11)	1.384(5)	C(17)	H(16) ^c	1.19
C(11)	C(12)	1.367(6)	C(17)	H(17) ^c	0.96
C(12)	C(13)	1.363(7)	C(17)	H(18) ^c	1.03
C(13)	C(14)	1.366(7)	C(18)	H(13)	0.98(6)
C(14)	C(15)	1.377(6)	C(18)	H(14)	0.79(6)
C(15)	C(10)	1.389(5)	C(18)	H(15)	1.02(8)
O(3)	C(16)	1.228(5)			
C(16)	N(1)	1.288(5)	Mg	O(1)	2.057(3)
N(1)	C(17)	1.441(6)	Mg	O(2)	2.055(2)
N(1)	C(18)	1.461(7)	Mg	O(3)	2.095(3)
O(1)	O(1) ^b	2.855(5)	O(2)	O(3)	2.878(3)
O(1)	O(2)	2.811(3)	O(2)	O(3) ^b	3.072(3)
O(1)	O(2) ^b	2.917(3)	O(3)	O(3) ^b	2.968(5)
O(1)	O(3) ^b	2.967(3)			

(a) Standard deviations given in parentheses.

(b) Related by two-fold axis through Mg.

(c) Positions not refined by least-squares.

Figure 3.

Stereoscopic view of the unit cell, showing packing of complexes. Labeled axes are positive from the origin. Hydrogens have been omitted for clarity.

MSL 722-280

D. References

- Cromer, D.T. and Liberman, D. (1970). J. Chem. Phys., 53(5), 1891.
- Doyle, P.A. and Turner, P.S. (1968). Acta Cryst., A24, 390.
- Johnson, C.K. (1965). Oak Ridge National Report-3794, Revised, Oak Ridge, Tenn.
- Long, R.E. (1965). Ph.D. Thesis, University of California, Los Angeles, Calif.
- Stewart, R.F., Davidson, E.R. and Simpson, W.T. (1965). J. Chem. Phys., 42, 3175.
- Sutton, L.E., Ed. (1958). "Tables of Interatomic Distances," The Chemical Society, London, Burlington House, W.1.
- Wilson, A.J.C. (1942). Nature (London), 150, 152.

V. The Crystal and Molecular Structure of bis(1,3-diphenyl-1,3-propanedionato)calcium hemiethanolate. $\text{Ca}(\text{DPP})_2(\text{C}_2\text{H}_5\text{OH})_{\frac{1}{2}}$

A. Experimental Procedure

The complex was synthesized by mixing a solution of calcium acetate in aqueous ethanol with an ethanolic solution of 1,3-diphenyl-1,3-propanedione (HDPP) and adding concentrated $\text{NH}_3/\text{NH}_4\text{Cl}$, pH 10, buffer. A lemon yellow precipitate appeared immediately upon addition of the buffer solution. The mixture was heated to reflux for one hour to allow the reaction to go to completion. The precipitate dissolved on heating, but reappeared as the mixture was cooled. It was filtered and dried in air and stored in a desiccator over Drierite. The formal reaction is $\text{Mg}(\text{OAc})_2 + 2 \text{HDPP} \xrightarrow{\text{pH } 10} \text{Mg}(\text{DPP})_2 + 2 \text{HOAc}$.

Attempts to recrystallize the complex failed, generally yielding a viscous orange oil, until well-formed clear yellow crystals were obtained from a slightly warmed solution of the compound in chloroform, which was allowed to evaporate very slowly. The oil obtained from other attempts at re-crystallization dissolved in many organic solvents, but formed a cloudy solid on addition of ethanol.

Crystals obtained from the chloroform solution were mounted on glass fibers. Precession photographs indicated a triclinic unit cell, and preliminary cell dimensions were determined. Another crystal was mounted on a Picker/Nuclear four-circle diffractometer automated with a DEC PDP 8I computer and disc. Twelve high-angle reflections were carefully centered using

monochromatized Cu $K\alpha_1$ radiation ($\lambda = 1.54051 \text{ \AA}$). The cell dimensions were refined on the 2θ measurements only, using program TTHCEL, a modified version of a program supplied to us by H. Hope. The program refines the real cell dimensions and minimizes $\sum w(d_o^* - d_c^*)^2$, where the relative weights are assigned subjectively by the user. The cell parameters obtained from this refinement were $a = 15.247(1) \text{ \AA}$, $b = 13.555(1) \text{ \AA}$, $c = 14.097(3) \text{ \AA}$, $\alpha = 74.64(1)^\circ$, $\beta = 95.97(1)^\circ$, $\gamma = 113.59(1)^\circ$. Calculation of the density of the compound for a cell volume of $2576 \text{ \AA}^3$ gave, for an empirical formula $\text{Ca}(\text{DPP})_2$, $d_c = 0.314 \cdot Z$. Attempts to measure the density failed, due to the fact that a solution could not be found which did not either dissolve the crystals or cause them to decompose, but observations did indicate that the most reasonable density would be for a Z of 4, giving $d_c = 1.26 \text{ gm/cm}^3$. The assumption of $Z = 4$ was confirmed by the refinement of the structure and the density based on the actual composition is $d_c = 1.32 \text{ gm/cm}^3$.

Data were collected using monochromatized ($2\theta_m = 26.42^\circ$) Cu $K\alpha$ radiation and a θ - 2θ scan technique. The peaks were scanned at a rate of $1^\circ/\text{min}$ from 1.0° below the calculated $K\alpha_1$ position to 1.0° above the calculated $K\alpha_2$ position. Backgrounds were counted for 10 seconds at positions offset by 0.5° from each end of the scan (all angles 2θ). Aluminum attenuators were automatically inserted into the diffracted beam whenever the count rate exceeded 7000 cps and the peak was rescanned and backgrounds re-measured with the attenuators in place. The hemisphere of data $+\underline{h}, \pm \underline{k}, \pm \underline{l}$ was collected for $2\theta = 0^\circ - 100^\circ$ ($\sin\theta/\lambda < 0.500$). Three

reflections, the (4 0 0), (0 3 0) and (0 0 3), were measured after every 100 data to monitor crystal decay. 5824 reflections were measured over the entire data collection period, including remeasurement of some reflections in connection with the decay correction. The standards showed a gradual fall in relative intensity from 1.00 to 0.942 over the first 3973 reflections, then a sharp drop to a relative intensity of 0.820 between reflections 3973 and 3974 caused by a malfunction in the equipment. The exact location of the change was determined by taking the data in that region again, after the data set was completed. The proper scaling was applied to all reflections, then the Lorentz and polarization factors were applied to the data. After the few equivalent reflections were averaged, there were 5294 unique reflections, of which 4504 had $F^2 > \sigma(F^2)$. Intensities and their standard deviations were calculated according to the formulae:

$$I = C - \frac{t_c}{2t_b} (B_1 + B_2) ; \sigma(I) = C + \frac{t_c^2}{4t_b^2} (B_1 + B_2)$$

where C is the total count in the peak, taken for scan time t_c , and B_1 and B_2 are the background counts, each taken for time t_b .

The atomic scattering factors of Doyle and Turner (1968) were used for Ca^{++} , O and C, along with the real and imaginary dispersion terms of Cromer and Liberman (1970). The scattering factors of Stewart, Davidson and Simpson (1965) were used for the hydrogen atoms.

Our least-squares program minimizes the function $\Sigma w(\Delta F)^2 / \Sigma w F_o^2$. The weighting scheme used throughout the refinement gave

zero weight to those reflections with $F^2 < \sigma(F^2)$ and gave $w = 1/\sigma^2(F)$ for all others. $\sigma(F)$ was calculated from $\sigma(F^2)$ by the method of finite differences.

The following programs written for our CDC 6600 computer were used during the solution and refinement of this structure. MAGPIK, INCOR and ORDER, general data reduction programs; WILSON, an unpublished Wilson-plot program by Maddox and Maddox; REL, R.E. Long's (1965) phase determination program; FORDAP, A. Zalkin's Fourier analysis program; LSLONG, our unpublished version of the Ganzel-Sparks-Trueblood least-squares program; DISMAT, a crystallographic distances and angles program which uses the correlation matrix from least-squares in calculating standard deviations; ORTEP, C. Johnson's (1965) thermal ellipsoid plotting program; HFINDR, a program for calculation of expected hydrogen atom positions; LSPLAN, our modified version of the least-squares planes program from the University of Pittsburgh; and LISTI and LISTAP, data presentation programs.

B. Determination and Refinement

The assumption of the centric space group $\overline{P}1$ was made throughout, and is confirmed by the solution and successful refinement of the structure.

A Patterson map was calculated which showed some peaks that could have been the interrelations of two Ca atoms, but when the derived positions of the Ca's were refined by least-squares one thermal parameter became negative and the R value ($R_1 = \Sigma |\Delta F| / \Sigma |F_o|$) was 0.56. This lack of any positive indication of a correct structure, plus the fact that the Ca atom possesses only 5% of the electrons in the structure, led to an attempt to solve the structure by direct methods.

Wilson's (1942) statistics were performed, and normalized structure factors, E_h , were generated in the usual manner. Phases were assigned to the 333 E values greater than 2.00 using R.E. Long's (1965) program, which iteratively applies the relationship $\text{sign}(E_h) = \text{sign}(\Sigma_k E_k E_{h-k})$. The fixed positive phases of the $(3 \ 2 \ 2)$, $(1 \ 3 \ 1)$ and $(2 \ 3 \ 2)$ determined the origin, and the signs of four other reflections, the $(5 \ 2 \ 5)$, $(2 \ 3 \ 4)$, $(0 \ 3 \ 3)$ and $(1 \ 5 \ 2)$ were held fixed during each of the 16 computer runs during which they were allowed to have all possible combinations of positive and negative phases. A consistency index defined as:

$$C = \frac{\Sigma_h \Sigma_k E_h E_k E_{h-k}}{\Sigma_h \Sigma_k |E_h E_k E_{h-k}|}$$

was calculated for the generated set of phases for each combination. Fourier maps using the phased E values as coefficients

were calculated for the three sets with the largest (0.94, 0.86, 0.77) values of C and the least number of iterations to convergence, but each showed only one large peak instead of the two expected for $Z = 4$ in the centric space group. A new set of starting reflections was chosen, with $(1 \bar{3} \bar{1})$, $(2 \ 3 \ 2)$ and $(3 \ \bar{4} \ \bar{2})$ positive to determine the origin and $(3 \ 1 \ \bar{5})$, $(1 \ \bar{5} \ \bar{7})$, $(5 \ \bar{4} \ 3)$ and $(2 \ \bar{3} \ \bar{1})$ used for the combinations, and the program was run again. Three sets of reflections were generated which were identical to the top three of the first trial, along with two other combinations with a high value of C (0.90). Of the two new solutions, the one which converged in the least number of cycles was used to phase an E-map. This map revealed two large peaks and a number of smaller peaks in configurations identifiable as portions of a DPP molecule. The 41 atoms found on the E-map were then refined by full-matrix least-squares with isotropic thermal parameters. Two cycles of refinement brought R_1 to 0.41, and the generated signs of the F's were used to phase a Fourier map of F_o . This Fourier showed 69 peaks and one shoulder in chemically reasonable positions for the expected formulation of $\text{Ca}(\text{DPP})_2$ with two Ca atoms in the asymmetric unit. The seventy atoms were refined as before for three cycles, and for three cycles with the two Ca atoms and eight O atoms allowed anisotropic thermal parameters, to an R_1 of 0.21. A difference Fourier showed three major peaks with heights of $6 \text{ e}^-/\text{\AA}^3$, $3 \text{ e}^-/\text{\AA}^3$ and $3 \text{ e}^-/\text{\AA}^3$, and no other peaks as high as $1 \text{ e}^-/\text{\AA}^3$. The three peaks were linked together with the largest peak at a distance of 2.41 Å from Ca(1). The only chemically reasonable explanation,

which was subsequently confirmed by refinement, was that the peaks represented an ethanol coordinated to the Ca. The atoms of the ethanol were put into least-squares with isotropic thermal parameters, together with the calculated positions of the 44 hydrogens for the DPP ligands. The hydrogens were all given an isotropic thermal parameter equal to the average thermal parameter of the ring they were bonded to, and neither their positions nor their thermal parameters were allowed to refine. R_1 reduced to 0.09 and the ethanol atoms refined well.

Memory limitations of the computer being used forced the further refinement of the structure to proceed in blocks. Refining the hydrogens with isotropic thermal parameters and all other atoms with anisotropic thermal parameters, it was necessary to divide the structure into three blocks. The two Ca atoms and the overall scale factor were refined in each block. The three blocks consisted of 1) the heavy atoms of the two "inner" DPP ligands, 2) the heavy atoms of the two outer DPP ligands and the ethanol and 3) the hydrogen atoms. After two cycles of 1), two cycles of 2), three cycles of 3) and two more cycles of 1), R_1 was reduced to 0.057 and a difference Fourier map was calculated. This showed that one of the hydrogens had refined to a position far from its proper location, and the hydrogen was repositioned correctly. The enolic hydrogen was located on the difference Fourier, the expected positions of the methylene and methyl hydrogens were calculated, and all six were placed in least-squares and allowed to refine. New blocks were formed so

that the hydrogen atoms could refine in the same block as the heavy atoms to which they were attached, and again the two Ca atoms and scale factor were refined with each group. After two cycles of all groups, R_1 was 0.048, but the methylene and methyl hydrogens of the ethanol were refining poorly. The expected positions were recalculated and were included in least-squares but not allowed to refine.

Refinement then continued on the larger CDC7600 computer, allowing the refinement of the structure in only two blocks, one containing the two Ca atoms and the "inner" DPP ligands including hydrogens, the other containing the two Ca's, the "outer" DPP ligands and the ethanol. Further attempts to refine the ethanol hydrogens gave three hydrogens which refined to good angles but poor distances, H(46), H(47) and H(48), while the other three refined well. The three poor hydrogens were allowed to keep their directions from their carbons, but were reset to the proper distances and allowed to refine. They returned to their previous positions and the thermal parameters of H(47) and H(48) became very large. Removal of H(47) and H(48) and calculation of a difference Fourier showed no recognizable peaks, so their theoretical positions were calculated for a C-H distance of 0.90 Å and fixed thermal parameters of 12.0 Å^2 and 14.0 Å^2 respectively, were assigned to them. They were forced by least-squares to retain their C-H distance.

Examination of the data at this point showed that those reflections with large F values also showed large values of $w\Delta F$, and a factor p ($= 0.05$) of F^2 was added to $\sigma(F^2)$ according

to the equation $\sigma^2(F^2) = \sigma_0^2(F^2) + (pF^2)^2$ to reduce the weights of the high-F reflections.

Refinement of the hydrogens on the ethanol as described above caused H(49) to refine to a large C-H distance, and the fixed C-H distances to H(47) and H(48) were increased to 1.0 Å. H(49) refined to a reasonable position, but its thermal parameter was ill-conditioned and was fixed at 15 Å^2 . Refinement of the full structure for four more cycles reduced all parameter shifts to less than 20% of their standard deviations, and most shifts were less than 10% of their standard deviations. The refinement was considered complete at this time. R_1 was 0.040 for 4503 reflections, and R_2 ($\sum w(\Delta F)^2 / \sum wF_0^2$) was 0.043. The standard deviation of an observation of unit weight was 1.14. The data were examined for secondary extinction effects with negative results.

The observed data and the final differences are given in Table 1. The final coordinates and thermal parameters of the atoms are given in Table 2.

Table 1. (3 pages)

Observed structure factors, standard deviations and final differences ($\times 10.0$). Reflections marked with an * were given zero weight in least-squares.

Table 2.(2 pages)

Final coordinates and thermal parameters. $\text{Ca}(\text{DPP})_2(\text{C}_2\text{H}_5\text{OH})_{\frac{1}{2}}$.^{a,b}

(a) Standard deviations of the least significant digit(s) is given in parentheses.

(b) The form of the temperature factor (B in units of $\text{\AA}^2$) is

$$T = \exp(-.25(B_{11}h^2a^{*2} + B_{22}k^2b^{*2} + B_{33}l^2c^{*2} + 2B_{12}hka^*b^* + 2B_{13}hla^*c^* + 2B_{23}klb^*c^*)) \text{ for anisotropic and}$$

$$T = \exp(-B \sin^2\theta/\lambda^2) \text{ for isotropic thermal parameters.}$$

ATOM	X	Y	Z	811	822	833	812	813	823
CA121	-15589(4)	62742(4)	44489(4)	2.25(3)	2.55(3)	2.90(3)	1.78(2)	1.77(2)	-448(2)
CA122	-16097(4)	54531(5)	30824(4)	2.62(3)	3.09(3)	3.09(3)	1.23(2)	-1.00(2)	-66(2)
0111	-10458(1)	2998(1)	6414(1)	3.1(1)	2.49(1)	3.8(1)	1.22(1)	1.04(1)	-95(1)
0121	-1235(1)	4498(1)	5458(1)	3.0(1)	2.57(1)	3.10(1)	1.30(1)	1.30(1)	29(1)
C111	-1316(1)	2897(1)	7420(1)	3.9(1)	2.9(1)	4.6(1)	1.9(1)	1.9(1)	-1.0(1)
C121	-1185(1)	-1016(1)	8148(1)	4.2(1)	4.0(1)	5.1(1)	1.9(1)	1.6(1)	-4(1)
C131	-1745(1)	-1058(1)	7934(1)	4.2(1)	3.1(1)	6.0(1)	1.4(1)	1.0(1)	-1(1)
C141	-1157(1)	-1078(1)	7186(1)	5.3(1)	2.7(1)	7.1(1)	1.4(1)	1.5(1)	-1.7(1)
C151	-16620(1)	-10195(3)	6661(1)	4.7(1)	3.3(1)	4.9(1)	1.1(1)	1.1(1)	-1.1(1)
C161	-1070(1)	-1091(1)	6875(1)	2.4(1)	2.8(1)	3.6(1)	1.0(1)	1.1(1)	-1.1(1)
C171	-10137(2)	2274(2)	6352(2)	3.0(2)	2.9(2)	3.3(1)	1.1(1)	-2(1)	-8(1)
C181	10704(2)	2544(3)	5863(2)	3.4(2)	2.5(2)	4.0(2)	1.5(1)	1.5(1)	-7(1)
C191	1323(2)	3614(3)	5407(2)	3.0(2)	3.3(2)	2.9(1)	1.4(1)	-2(1)	-7(1)
C1101	2177(2)	2957(2)	4651(2)	3.3(2)	3.1(1)	3.0(1)	1.8(1)	-1(1)	-3(1)
C1111	2181(1)	2987(1)	4591(2)	4.2(2)	3.7(2)	4.1(2)	1.5(2)	1.7(2)	-1.0(1)
C1121	2296(1)	3152(4)	2985(3)	5.5(2)	5.7(2)	5.8(2)	2.9(2)	1.7(2)	-1.4(2)
C1131	3778(1)	4085(4)	3897(4)	4.7(3)	7.7(3)	6.8(3)	3.0(2)	2.0(2)	-1.3(2)
C1141	13801(3)	4858(4)	4325(3)	3.1(2)	6.5(3)	9.1(3)	1.9(2)	1.2(2)	-2.5(2)
C1151	3000(3)	4707(3)	4622(3)	3.3(2)	4.7(2)	5.7(2)	1.4(2)	1.3(2)	-1.8(2)
0131	-10862(1)	2583(2)	3235(1)	2.7(1)	3.02(1)	3.43(1)	1.20(1)	1.02(1)	-1.04(1)
0141	-10801(1)	3648(1)	4218(1)	3.1(1)	2.76(1)	3.18(1)	1.26(1)	1.12(1)	-9.1(1)
C1161	2612(1)	2730(4)	2199(3)	3.8(2)	3.3(2)	6.1(2)	1.2(1)	1.0(1)	-3.9(1)
C1171	3148(1)	5991(1)	1498(4)	4.6(3)	12.4(4)	9.2(3)	-1.0(3)	2.4(2)	-6.2(3)
C1181	3145(4)	5289(1)	1093(4)	5.1(3)	11.4(4)	6.7(3)	3.2(3)	2.5(3)	-1.8(3)
C1191	22357(1)	4327(5)	11005(3)	10.1(4)	6.8(3)	5.8(3)	4.8(3)	3.3(2)	-5.5(2)
C1211	11607(1)	1819(1)	3161(1)	7.3(1)	4.8(1)	3.3(1)	3.3(1)	2.8(1)	-1.3(1)
C1211	11632(1)	4757(3)	2191(2)	3.4(2)	3.9(2)	3.1(2)	1.8(1)	1.1(1)	-8(1)
C1221	10841(2)	4495(2)	2882(2)	3.2(2)	3.2(2)	2.8(1)	1.7(1)	-2(1)	-1.0(1)
C1231	3427(1)	3427(1)	3121(2)	3.4(2)	2.8(2)	4.4(2)	1.7(1)	1.7(1)	-1.3(1)
C1241	-10623(2)	3024(2)	3798(2)	3.1(2)	2.7(1)	2.9(1)	1.4(1)	-8(1)	-7(1)
C1251	-1248(2)	1819(2)	4006(2)	3.0(2)	3.0(1)	3.4(1)	1.4(1)	-3(1)	-9(1)
C1261	-1228(1)	1104(3)	3492(4)	7.1(3)	5.9(2)	9.5(3)	1.5(2)	4.0(2)	-2.3(2)
C1271	-11845(1)	-10002(3)	3707(1)	8.4(3)	5.5(2)	10.7(4)	1.4(2)	3.3(3)	-3.0(2)
C1281	-12510(3)	-10421(3)	6423(3)	5.1(2)	2.8(2)	6.4(2)	1.7(2)	-1.5(2)	-4(1)
C1291	-12572(3)	-10271(3)	4929(3)	5.6(2)	6.0(2)	6.1(2)	-2(2)	1.7(2)	-1.5(2)
C1301	-1196(3)	1370(3)	4771(3)	5.3(2)	3.4(2)	6.1(2)	-10(2)	2.0(2)	-1.9(2)
0151	7142(2)	2380(2)	6385(1)	2.7(1)	4.4(1)	3.9(1)	1.49(1)	1.24(1)	-6.8(1)
0161	7790(1)	3340(1)	4360(1)	2.1(1)	3.9(1)	3.9(1)	1.9(1)	1.2(1)	-1.00(1)
C1311	16125(3)	10883(3)	7993(3)	3.9(2)	4.5(2)	4.2(2)	1.3(2)	1.0(2)	-1.0(2)
C1321	5619(3)	10091(3)	8905(3)	5.5(3)	5.6(2)	4.4(2)	2.3(2)	1.1(2)	1(1)
C1331	4679(3)	-10577(3)	8720(4)	5.0(3)	4.9(2)	6.8(3)	1.7(2)	2.9(2)	1.0(2)
C1341	4230(3)	-10444(3)	7895(4)	3.8(2)	4.7(2)	7.5(3)	1.5(2)	1.5(2)	1(1)
C1351	4720(3)	10301(3)	7022(3)	3.3(2)	4.5(2)	5.6(2)	1.0(2)	1.0(2)	-1.0(2)
C1361	5683(2)	10136(2)	7089(2)	3.0(2)	3.0(1)	4.6(2)	1.9(1)	1.2(1)	-4(1)
C1371	6261(2)	11862(2)	6220(2)	2.7(2)	3.2(1)	4.3(2)	1.9(1)	1.7(1)	-1.0(1)
C1381	5833(3)	12040(3)	5298(3)	2.1(2)	4.4(2)	4.4(2)	1.1(1)	1.1(1)	-9(1)
C1391	6530(3)	12758(3)	4428(2)	2.9(2)	3.5(2)	3.9(2)	1.3(1)	1.3(1)	-1.3(1)
C1401	5716(2)	2897(3)	3616(2)	2.4(2)	4.1(2)	4.1(2)	1.0(1)	1.1(1)	-1.6(1)
C1411	4925(3)	2027(4)	3303(3)	3.4(2)	4.9(2)	5.9(2)	1.1(2)	-2.3(2)	-1.3(2)
C1421	4417(1)	2191(5)	2444(4)	3.8(2)	8.0(3)	6.9(3)	1.5(2)	-1.0(2)	-4.1(3)
C1431	4688(4)	3206(5)	1810(4)	5.9(3)	10.3(4)	5.0(3)	4.6(3)	-2.4(2)	-3.1(3)
C1441	5468(3)	4009(5)	7005(3)	5.7(3)	7.8(3)	4.5(2)	3.3(2)	-1.0(2)	-1.4(2)
C1451	5594(3)	3923(3)	2853(3)	3.4(2)	5.6(2)	4.2(2)	1.5(2)	-1.0(2)	-1.7(2)
0171	-11821(1)	45180(1)	2453(1)	3.4(1)	4.0(1)	3.6(1)	1.84(1)	1.24(1)	-1.08(1)
0181	-10146(1)	6543(2)	1518(2)	4.0(1)	4.7(1)	3.8(1)	2.6(1)	1.59(1)	1.09(1)
C1461	-1311(1)	7178(1)	2481(1)	4.0(2)	4.3(2)	4.9(2)	2.4(2)	1.3(2)	-8(2)
C1471	-13883(1)	7475(1)	7466(1)	4.4(2)	6.0(2)	7.6(3)	2.6(2)	1.0(2)	-1.2(2)
C1481	-14278(1)	7786(3)	1578(5)	3.2(2)	5.2(2)	11.3(4)	2.6(2)	-1.0(2)	-2.1(2)
C1491	-13934(3)	7776(3)	10726(4)	4.9(2)	6.0(2)	7.6(3)	3.3(2)	-1.1(2)	-1.3(2)
C1501	-13151(1)	7510(1)	10755(3)	4.4(2)	5.2(2)	4.8(2)	2.7(2)	-0.8(2)	-0.9(2)
C1511	-12726(2)	7209(2)	13630(2)	2.9(2)	3.1(1)	4.0(2)	1.2(1)	1.1(1)	-4(1)
C1521	-11865(2)	6913(3)	1701(2)	2.8(2)	3.4(2)	3.6(2)	1.2(1)	1.1(1)	-9(1)
C1531	-11169(2)	7487(3)	10968(3)	3.4(2)	4.0(2)	1.9(2)	1.4(2)	1.4(2)	1(1)
C1541	-10326(1)	7326(3)	10930(2)	3.1(2)	3.7(2)	3.2(2)	1.2(1)	1.1(1)	-9(1)
C1551	-10445(2)	8127(3)	10303(2)	3.4(2)	3.9(2)	3.5(2)	1.2(1)	1.2(1)	-1.3(1)
C1561	-1037(3)	9011(3)	-10494(3)	4.3(2)	4.7(2)	4.7(2)	1.4(2)	1.0(2)	-1.1(2)
C1571	-1131(4)	9747(4)	-11141(3)	6.1(3)	5.0(2)	5.9(2)	1.1(2)	1.7(2)	1(1)
C1581	-19602(4)	9602(4)	-10324(4)	5.4(3)	5.8(3)	7.1(3)	-1.1(2)	3.0(2)	-1.1(2)
C1591	-20845(3)	8728(4)	-10884(4)	4.1(2)	6.9(3)	7.1(3)	1.3(2)	1.9(2)	-1.2(2)
C1601	-1305(3)	7990(4)	10261(3)	5.2(3)	5.6(2)	5.3(2)	2.5(2)	1.1(2)	-0.4(2)
0191	8123(2)	3902(2)	2456(2)	4.5(1)	4.7(1)	4.6(1)	1.4(1)	-1.0(1)	-1.7(1)
C1611	7826(4)	3822(5)	1504(4)	6.2(3)	12.2(4)	5.9(3)	2.2(3)	-1.7(2)	-3.8(3)
C1621	8634(5)	4059(5)	1883(4)	13.7(5)	12.3(4)	5.9(3)	7.3(4)	-1.4(3)	-0.0(3)

H(01)	-.134(2)	.150(3)	.777(2)	3.8(7)
H(02)	-.222(2)	-.028(2)	.064(2)	4.4(8)
H(03)	-.211(2)	-.102(3)	.827(2)	5.8(9)
H(04)	-.108(2)	-.145(3)	.702(2)	5.4(9)
H(05)	-.028(2)	.024(3)	.611(2)	5.3(9)
H(6)	.093(2)	.199(2)	.589(2)	3.1(7)
H(07)	.159(2)	.231(3)	.439(2)	3.4(7)
H(09)	.293(2)	.255(3)	.356(2)	5.9(9)
H(09)	.428(3)	.425(3)	.347(3)	7.4(11)
H(10)	.432(3)	.546(3)	.434(2)	5.5(10)
H(11)	.303(2)	.524(3)	.516(2)	4.9(8)
H(12)	.262(3)	.620(3)	.253(3)	7.0(10)
H(13)	.365(4)	.067(5)	.150(4)	12.3(18)
H(14)	.362(3)	.540(4)	.056(3)	9.0(13)
H(15)	.226(3)	.388(3)	.057(3)	6.5(11)
H(16)	.104(3)	.336(4)	.162(3)	9.2(13)
H(17)	.020(2)	.290(3)	.283(2)	4.4(8)
H(18)	-.078(3)	.135(4)	.257(3)	9.4(14)
H(19)	-.105(3)	-.043(4)	.335(3)	9.3(14)
H(20)	-.292(2)	-.116(3)	.455(2)	4.0(7)
H(21)	-.296(3)	.003(3)	.554(3)	7.6(11)
H(22)	-.206(3)	.179(4)	.506(3)	8.0(13)
H(23)	.679(3)	.137(3)	.809(2)	5.1(9)
H(24)	.596(2)	.000(3)	.939(3)	5.0(9)
H(25)	.429(3)	-.116(3)	.925(3)	5.6(9)
H(26)	.363(3)	-.094(3)	.774(3)	6.4(11)
H(27)	.440(2)	.037(3)	.643(3)	5.3(10)
H(28)	.518(2)	.167(2)	.520(2)	3.1(7)
H(29)	.473(2)	.130(3)	.371(2)	3.9(8)
H(30)	.386(3)	.151(3)	.236(2)	6.1(9)
H(31)	.440(3)	.339(3)	.126(3)	5.9(10)
H(32)	.569(3)	.477(3)	.156(3)	7.8(13)
H(33)	.653(2)	.454(2)	.301(2)	3.2(7)
H(34)	-.288(2)	.703(2)	.799(2)	1.9(7)
H(35)	-.416(3)	.737(3)	.107(3)	7.5(12)
H(36)	-.475(3)	.796(3)	.157(3)	6.3(11)
H(37)	-.419(3)	.796(3)	.007(3)	7.4(11)
H(38)	-.291(2)	.751(2)	.016(2)	3.8(8)
H(39)	-.127(2)	.798(2)	.050(2)	2.6(7)
H(40)	-.023(2)	.715(3)	.078(2)	4.9(8)
H(41)	.102(3)	1.039(3)	-.159(3)	7.4(11)
H(42)	.249(3)	1.004(3)	-.142(3)	7.5(12)
H(43)	.262(3)	.862(3)	-.023(3)	7.6(13)
H(44)	.136(2)	.743(2)	.070(2)	3.1(8)
H(45)	.779(5)	.375(6)	.281(5)	15.6(28)
H(46)	.717(5)	.312(4)	.136(3)	10.6(16)
H(47)	.747	.434	.1232	12.000
H(48)	.8860	.3438	.113	14.000
H(49)	.930(4)	.469(5)	.089(5)	15.000
H(50)	.841(4)	.396(4)	.015(4)	12.8(16)

C. Results and Discussion

The complex forms into clusters of composition $(\text{Ca}_2(\text{DPP})_4(\text{C}_2\text{H}_5\text{OH}))_2$ around the center of symmetry at $0, \frac{1}{2}, \frac{1}{2}$. There is only one such cluster per unit cell, and its neighbors are related to it by unit cell translations. There are no contact distances of less than 3.50 Å between non-hydrogen atoms in different clusters.

The two independent calcium atoms Ca(1) and Ca(2) are respectively six-fold and ^{seven}-fold coordinated by the oxygens of the DPP ligands and the ethanol (Figure 1). Each calcium shares two oxygens with each of two neighboring calcium atoms of the other kind so that four of them are linked around the center of symmetry. The bridging oxygens, O(1) and O(2) of DPP(1), O(3) and O(4) of DPP(2), form a slightly distorted parallelopiped, with Ca atoms capping each of the four rectangular faces and the DPP ligands projecting into the area above the non-rectangular face (Figure 2). The edge lengths of the parallelopiped are somewhat distorted (Table 3). The angles between the faces of the parallelopiped from the uncapped plane (core plane 1) to the two planes capped by Ca(1) and Ca(2) (core planes 2 and 3 respectively) are 89.1° and 89.4°. The angle between the two capped planes is 69.1°.

The non-bridging oxygens O(5) and O(6) of DPP(3), O(7) and O(8) of DPP(4) and O(9) of the ethanol and the calcium atoms lie on the outward sides of the planes in question. Ca(1) is 1.03 Å above core plane 2, and the two oxygens of DPP(3) are 3.09 Å and 2.55 Å away from that plane. Ca(2) is 1.39 Å from core plane 3. The other three oxygens coordinated to Ca(2) are

Figure 1.

Stereoscopic view of the coordination around the Ca atoms.

The center of symmetry at $0, \frac{1}{2}, \frac{1}{2}$ is in the center of the cluster. Thermal ellipsoids have been scaled to include 50% probability.

Figure 2.

Stereoscopic view of the complex cluster. Hydrogen atoms and phenyl carbons except for those attached directly to the diketone have been omitted for clarity. View direction and scaling of the thermal ellipsoids are the same as in Figure 1.

XBL 724-663

Table 3.

Bonded and non-bonded distances between calcium and oxygen in the coordination cluster (Å).^a (See Figure 1.)

Bonded			Non-bonded		
Atom(1)	Atom(2)	Distance	Atom(1)	Atom(2)	Distance
Ca(1)	O(1)	2.353(2)	O(1)	O(2)	2.772(3) ^{b, c}
Ca(1)	O(2)	2.333(2)	O(1)	O(3)	2.807(3) ^c
Ca(1)	O(3)	2.358(2)	O(1)	O(4)	3.037(3) ^c
Ca(1)	O(4)	2.353(2)	O(2)	O(3)	3.102(3) ^c
Ca(1)	O(5)	2.256(2)	O(2)	O(4)	2.995(3) ^c
Ca(1)	O(6)	2.395(2)	O(3)	O(4)	2.804(3) ^{b, c}
Ca(2)	O(1)	2.416(2)	O(1)	O(5)	3.410 ^d
Ca(2)	O(2)	2.411(2)	O(1)	O(8)	3.111
Ca(2)	O(3)	2.436(2)	O(2)	O(4)	3.333(3) ^e
Ca(2)	O(4)	2.508(2)	O(2)	O(6)	3.010
Ca(2)	O(7)	2.308(2)	O(2)	O(7)	2.999
Ca(2)	O(8)	2.266(2)	O(3)	O(5)	3.526
Ca(2)	O(9)	2.408(2)	O(3)	O(8)	3.205
			O(4)	O(6)	2.943
			O(4)	O(9)	2.851
			O(5)	O(6)	2.801(3) ^b
			O(6)	O(9)	2.941(4)
			O(7)	O(8)	2.803(3) ^b
			O(7)	O(9)	3.054(3)
			O(8)	O(9)	3.491(4)

(a) Standard deviations given in parentheses.

(b) "Bite" of the ligand.

(c) Edges of central parallelepiped.

(d) Atoms were refined in two different blocks of least-squares.

Standard deviations are probably the same as others, ± 0.003 Å.

(e) Diagonal across the face of core plane 1.

at distances from core plane 3 of 2.90 Å, 2.80 Å and 2.82 Å to O(7), O(8) and O(9) respectively, forming an almost parallel plane. The distances between core planes 1, 2 and 3 and the respective planes related by the center of symmetry are 2.90 Å for core plane 1, 2.60 Å for core plane 2 and 2.87 Å for core plane 3.

The Ca-O distances average to 2.367 Å with a standard deviation of ± 0.026 Å calculated from the scatter of the measurements. The average compares well to the sum of the crystal radii (Pauling, 1960) of 2.39 Å. The range of the Ca-O distances is from 2.256(2) Å to 2.508(2) Å. The Ca(1)-O distances are in general shorter than the Ca(2)-O distances, and the distances from Ca to the unbridged oxygens are generally shorter than those to the bridging oxygens. The Ca(1)-O(6) distance of 2.395 Å is an exception due to steric factors mentioned below. These variances from the averages are in keeping with the coordinations of the atoms involved, with longer bond distances to atoms with higher coordination.

The distances of the Ca atoms to the planes of the diketone portions of the inner DPP ligands are on the same order as the distances from the Ca atoms to the core planes. For the non-bridging DPP ligands the Ca atoms are much closer to the planes of the diketones. For DPP(3), the distance from the plane to Ca(1) is only 0.19 Å, while in the more crowded environment around Ca(2) the distance from Ca(2) to the diketone plane of DPP(4) is 0.24 Å. The planes of the diketone residues of DPP(1)

2. Construction of apparatus.

The apparatus designed on the above principles and used in this study is shown in Fig. VI. It is capable of making measurements on samples ranging from fractions of a microgram to tens of milligrams, the upper limit being primarily a function of available space in the sample tube and the lower limit being determined by the finite mass of a containment system for any sample. In general, the sample container is chosen to match not only the physical size of the sample, but also the expected magnetic susceptibility. Thus, a sample of low susceptibility would require a lighter container than one of high susceptibility in order that reasonable deflections (d) might be observed. For example, the protactinium metal samples used in this study were expected to be of low susceptibility, and thus containers of minimum weight were desired. Knowing that the minimum deflection which can be measured with this apparatus is approximately 2.5μ , the minimum detectable force on the sample can be calculated. For example, the second metal sample has a mass of 0.479 mg and its container 1.008 mg . From eq. (2)

$$\begin{aligned} F_1 L \cos \theta &= F_2 + F_3 \frac{d}{2} = (m_h + m_{sc} + m_s)gd + m_f \frac{d}{2} \\ F_{1 \min} &= \frac{(m_h + m_{sc} + m_s + m_f/2)gd_{\min}}{L \cos \theta} \\ &= \frac{((0.0644 + 1.008 + 0.479 + 0.080) \times 10^{-3})gd_{\min}}{40} \\ &= 1.01 \times 10^{-5} \text{ dyne.} \end{aligned}$$

This minimum detectable force is approximately ten times larger than that

Figure 3.

Generalized numbering system for DPP ligands. Carbons are large single circles, oxygens are large double circles and hydrogens are small single circles. Numbering shown is for DPP(1), for successive DPP molecules add 2 to the oxygen numbers, 15 to the carbon numbers and 11 to the hydrogen numbers.

Table 4

Rotation of phenyl ring planes with respect to diketone planes and each other (deg).

DPP	Phenyl 1 to diketone.	Phenyl 2 to diketone.	Phenyl 1 to Phenyl 2.
1	21.1	31.0	33.8
2	13.4	7.8	16.1
3	5.6	34.0	28.6
4	36.6	12.9	47.6

This rotation is due to packing forces between the clusters and between ligands within the clusters. The angles of rotation do not cause any significant variation in the phenyl-diketone bond length or the stressed interior angle adjacent to the diketone in the phenyl groups.

Interatomic distances in the DPP ligands and the ethanol are shown in Table 5. The average phenyl C-C distance is $1.374 \text{ \AA}$, in reasonable agreement with literature values, and the C-C distance in the diketone averages to $1.396 \text{ \AA}$, again in good agreement. The average C-C distance between the phenyl groups and the diketone residue of $1.502 \text{ \AA}$ agrees well with other values for $\text{sp}^2\text{-sp}^2$ single bonds, and the C-O average distance of $1.273 \text{ \AA}$ is in reasonable agreement for a C-O bond order of 1.5 (Pauling, 1960). The standard deviations of these averages as calculated from the scatter of the various independent distances are all on the order of $\pm 0.003 \text{ \AA}$. There is a statistically, but probably not chemically significant difference in the C-O bond lengths between those involving bridging oxygens and those involving non-bridging oxygens. The average C-O distance for the former is slightly longer at $1.281(2) \text{ \AA}$ than the average C-O distance for the latter at $1.264(3) \text{ \AA}$. The phenyl C-C bonds also show a statistically significant variation. The bonds to the para carbon average to a low $1.363(2) \text{ \AA}$ compared to the averages of the ortho-meta bonds ($1.382(2) \text{ \AA}$) and the bonds to the carbon adjacent to the diketone ($1.379(3) \text{ \AA}$). The average of all C-H bonds in the DPP molecules is $0.94 \text{ \AA}$ with a spread of $\pm 0.09 \text{ \AA}$, in agreement with values for many x-ray studies.

Table 5.

Intra-molecular bond distances (Å): $\text{Ca}(\text{DPP})_2(\text{C H OH})_{\frac{1}{2}}$.

DPP(1)			DPP(2)		
Atom(1)	Atom(2)	Distance	Atom(1)	Atom(2)	Distance
C(1)	C(2)	1.379(5)	C(16)	C(17)	1.385(6)
C(2)	C(3)	1.376(5)	C(17)	C(18)	1.350(7)
C(3)	C(4)	1.367(5)	C(18)	C(19)	1.367(7)
C(4)	C(5)	1.391(5)	C(19)	C(20)	1.378(6)
C(5)	C(6)	1.388(4)	C(20)	C(21)	1.379(5)
C(6)	C(1)	1.381(4)	C(21)	C(16)	1.364(5)
C(6)	C(7)	1.496(4)	C(21)	C(22)	1.500(4)
C(7)	O(1)	1.285(3)	C(22)	O(3)	1.280(3)
C(7)	C(8)	1.387(4)	C(22)	C(23)	1.391(4)
C(8)	C(9)	1.397(4)	C(23)	C(24)	1.403(4)
C(9)	O(2)	1.283(3)	C(24)	O(4)	1.277(3)
C(9)	C(10)	1.496(4)	C(24)	C(25)	1.502(4)
C(10)	C(11)	1.377(4)	C(25)	C(26)	1.379(5)
C(11)	C(12)	1.368(5)	C(26)	C(27)	1.384(6)
C(12)	C(13)	1.365(6)	C(27)	C(28)	1.349(6)
C(13)	C(14)	1.368(6)	C(28)	C(29)	1.348(5)
C(14)	C(15)	1.387(5)	C(29)	C(30)	1.373(5)
C(15)	C(10)	1.386(4)	C(30)	C(25)	1.368(5)
C(1)	H(1)	0.92(3)	C(16)	H(12)	0.97(4)
C(2)	H(2)	0.95(3)	C(17)	H(13)	0.92(5)
C(3)	H(3)	0.97(3)	C(18)	H(14)	0.90(5)
C(4)	H(4)	0.92(3)	C(19)	H(15)	0.93(4)
C(5)	H(5)	0.95(3)	C(20)	H(16)	1.00(4)
C(8)	H(6)	0.93(3)	C(23)	H(17)	0.95(3)
C(11)	H(7)	1.00(3)	C(26)	H(18)	0.95(4)
C(12)	H(8)	1.01(4)	C(27)	H(19)	0.86(4)
C(13)	H(9)	0.91(4)	C(28)	H(20)	0.94(3)
C(14)	H(10)	0.88(3)	C(29)	H(21)	1.00(4)
C(15)	H(11)	0.94(3)	C(30)	H(22)	0.90(4)

Table 5 (continued)

DPP(3)			LPP(4)		
Atom(1)	Atom(2)	Distance	Atom(1)	Atom(2)	Distance
C(31)	C(32)	1.389(5)	C(46)	C(47)	1.384(5)
C(32)	C(33)	1.367(6)	C(47)	C(48)	1.373(5)
C(33)	C(34)	1.364(6)	C(48)	C(49)	1.365(6)
C(34)	C(35)	1.378(5)	C(49)	C(50)	1.373(5)
C(35)	C(36)	1.393(5)	C(50)	C(51)	1.373(4)
C(36)	C(31)	1.381(5)	C(51)	C(46)	1.373(5)
C(36)	C(37)	1.506(4)	C(51)	C(52)	1.507(4)
C(37)	O(5)	1.272(3)	C(52)	O(7)	1.263(3)
C(37)	C(38)	1.399(4)	C(52)	C(53)	1.393(4)
C(38)	C(39)	1.399(4)	C(53)	C(54)	1.395(4)
C(39)	O(6)	1.262(3)	C(54)	O(8)	1.261(3)
C(39)	C(40)	1.502(4)	C(54)	C(55)	1.506(4)
C(40)	C(41)	1.381(4)	C(55)	C(56)	1.372(5)
C(41)	C(42)	1.386(6)	C(56)	C(57)	1.383(5)
C(42)	C(43)	1.354(6)	C(57)	C(58)	1.353(6)
C(43)	C(44)	1.374(6)	C(58)	C(59)	1.365(6)
C(44)	C(45)	1.388(5)	C(59)	C(60)	1.380(5)
C(45)	C(40)	1.382(5)	C(60)	C(55)	1.383(5)
C(31)	H(23)	0.98(3)	C(46)	H(34)	0.79(3)
C(32)	H(24)	0.93(3)	C(47)	H(35)	0.94(4)
C(33)	H(25)	0.97(4)	C(48)	H(36)	0.84(4)
C(34)	H(26)	0.92(4)	C(49)	H(37)	0.98(4)
C(35)	H(27)	0.92(3)	C(50)	H(38)	0.94(3)
C(38)	H(28)	0.93(3)	C(53)	H(39)	0.85(3)
C(41)	H(29)	0.94(3)	C(56)	H(40)	1.01(3)
C(42)	H(30)	0.99(4)	C(57)	H(41)	1.01(4)
C(43)	H(31)	0.88(4)	C(58)	H(42)	0.92(4)
C(44)	H(32)	0.92(4)	C(59)	H(43)	0.92(4)
C(45)	H(33)	0.96(3)	C(60)	H(44)	0.87(3)

Table 5 (continued)

Ethanol		
Atom(1)	Atom(2)	Distance
O(9)	C(61)	1.390(5)
C(61)	C(62)	1.464(7)
O(9)	H(45)	0.68(6)
C(61)	H(46)	1.12(5)
C(61)	H(47) ^b	1.00
C(62)	H(48) ^b	1.00
C(62)	H(49)	1.04(6)
C(62)	H(50)	1.09(6)

(a) Standard deviations given in parentheses.

(b) Position not refined in least-squares.

The inter-atomic angles in the ligands are normal for complexed DPP molecules (Section IV, Section VI, Zalkin, et al., 1969). The diketone ring is spread, with O-C-C and C-C-C bond angles $\sim 125^\circ$, to give the ligand a bigger "bite", and the interior angle in the phenyl ring at the juncture to the diketone is reduced from 120° to $\sim 118^\circ$ to reduce the steric interaction between the phenyl and the diketone.

The C-O and C-C bond distances in the ethanol are greatly shortened, probably due to the very large amounts of anisotropic thermal motion (Figure 2, Table 2).

Comparison of this structure to other polymerized diketone chelates will be given in Section VII.

D. References

- Cromer, D.T. and Liberman, D. (1970). J. Chem. Phys., 53(5), 1891.
- Doyle, P.A. and Turner, P.S. (1968). Acta Cryst., A24, 390.
- Johnson, C.K. (1965). Oak Ridge National Report-3794, Revised, Oak Ridge, Tenn.
- Long, R.E. (1965). Ph.D. Thesis, University of California, Los Angeles, CA .
- Pauling, L. (1960). "The Nature of the Chemical Bond", Third Ed., Cornell University Press, Ithaca, N.Y.
- Stewart, R.F., Davidson, E.R. and Simpson, W.T. (1965). J. Chem. Phys., 42, 3175.
- Wilson, A.J.C. (1942). Nature (London), 150, 152.
- Zalkin, A., Templeton, D.T. and Karraker, D.G. (1969). Inorg. Chem., 8(12), 2680.

VI. Crystal and Molecular Structure of bis(1,3-diphenyl-1,3-propanedionato)strontium hemiacetate. $\text{Sr}(\text{DPP})_2((\text{CH}_3)_2\text{CO})_{\frac{1}{2}}$.

A. Experimental Procedure

The complex was synthesized using a modification of the method of Hammond, et al. (1963). Strontium nitrate was dissolved in a small amount of water, and ethanol added until precipitation was just observed. This solution was added to an ethanolic solution of 1,3-diphenyl-1,3-propanedione (HDPP), and a strong aqueous $\text{NH}_3/\text{NH}_4\text{Cl}$, pH 10, buffer was added to the mixture. A whitish precipitate (probably either $\text{Sr}(\text{NO}_3)_2$ or HDPP) immediately formed, and the mixture was heated to reflux for an hour and stirred continuously. Heating caused the precipitate to dissolve, and cooling yielded a very pale yellow crystalline precipitate which was filtered and dried in air. The formal reaction is

$$\text{Sr}(\text{NO}_3)_2 + 2 \text{HDPP} \xrightarrow{\text{pH } 10} \text{Sr}(\text{DPP})_2 + 2 \text{HNO}_3.$$

Attempts to recrystallize the compound proceeded with difficulty, generally yielding a viscous oil, but finally a few crystals were obtained by very slow evaporation of an ethanol solution with a very small amount of acetone added to it. Of these few, only one was good enough for crystallographic study.

This crystal, a thin plate of approximate dimensions 0.03 mm x 0.17 mm x 0.32 mm was affixed to a glass fiber and precession photographs taken which indicated a triclinic unit cell. It was then mounted on a Picker/Nuclear four-circle diffractometer automated by a DEC PDP 8I computer and disc and equipped with a molybdenum x-ray tube.

Twelve reflections were carefully centered in χ , ω and 2θ using Mo K α radiation ($\lambda = 0.7107 \text{ \AA}$), and used to refine the cell dimensions and orientation matrix using the FACS-1 least-squares program on the PDP 8I. The cell dimensions were determined again by least squares on the positions of the same twelve reflections centered after the data collection and were found to have changed by as much as five standard deviations. The second set of cell parameters, $a = 13.759(10) \text{ \AA}$, $b = 14.182(9) \text{ \AA}$, $c = 16.031(9) \text{ \AA}$, $\alpha = 107.57(3)^\circ$, $\beta = 90.36(3)^\circ$, $\gamma = 113.07(3)^\circ$, were used throughout the structure analysis. Inspection of the intensities before data collection revealed a rapid fall-off of intensity with increasing 2θ , there being very few appreciable intensities beyond $2\theta = 25^\circ$ and none at all beyond $2\theta = 30^\circ$.

Data were collected using a θ - 2θ scan technique and graphite monochromatized ($2\theta_m = 11.80^\circ$) Mo K α radiation. Peaks were scanned 0.7° below the predicted K α_1 position to 0.7° above the predicted K α_2 position at a scan speed of $1^\circ/\text{min}$. Backgrounds were counted for 10 seconds at positions offset 0.3° from each end of the scan (all angles 2θ). Copper foil attenuators were automatically inserted in the diffracted beam any time the count rate exceeded 7000 cps and the peak was rescanned and the backgrounds measured with the attenuators in place. All reflections in the hemisphere of reciprocal space $+h, +k, +l$ were collected for $2\theta = 1^\circ$ - 30° ($\sin\theta/\lambda < 0.370$). Two reflections, the (3 0 0) and the (1 $\bar{3}$ 0), were measured after every 100 data to monitor crystal decay and alignment. The (1 $\bar{3}$ 0) showed a steady linear decline to 84% of

its original intensity over the data collection period, while the (3 0 0) showed large ($\pm 10\%$) random variations around a similar drop. 2371 reflections were measured, not including multiple measurements of the standards. Of these, 260 were measured with zero intensity, and in all 684 had $I < \sigma(I)$. Intensities and their standard deviations were calculated according to:

$$I = C - \frac{t_c}{2t_b} (B_1 + B_2) \quad \sigma(I) = C + \frac{t_c^2}{4t_b^2} (B_1 + B_2)$$

where C is the total counts over scan time t_c and B_1 and B_2 are the background counts, each measured for time t_b .

An absorption correction was applied to the intensities assuming that the crystal composition was $\text{Sr}(\text{DPP})_2(\text{C}_2\text{H}_5\text{OH})_{\frac{1}{2}}$ with $Z = 4$, in compositional isomorphism with the related Ca compound (Section V). This yielded $d_c = 1.36 \text{ gm/cm}^3$ and $\mu = 21.3 \text{ cm}^{-1}$. (The actual composition differs from this by one carbon atom, and does not affect the results to the accuracy given.) The correction factors ranged from 1.07 to 1.41. The data were then corrected for the linear crystal decay, Lorentz and polarization factors applied, and those reflections taken more than once averaged to give 2204 unique reflections of which 239 were measured as zero and another 303 had $F^2 < \sigma(F^2)$.

The atomic scattering factors of Doyle and Turner (1968) were used for neutral strontium, carbon and oxygen, together with the real and imaginary dispersion corrections of Cromer and Liberman (1970). Our least-squares program minimizes the function $\Sigma w(\Delta F)^2 / \Sigma w F_o^2$. The weighting scheme used throughout gave zero weight to those reflections with $F^2 < \sigma(F^2)$ and $w = 1/\sigma^2(F)$ for

all others. $\sigma(F)$ was calculated by finite differences from $\sigma(F^2)$ with $\sigma^2(F^2) = \sigma_o^2(F^2) + (pF^2)^2$, $\sigma_o(F^2) = (Lp)^{-1}\sigma(I)$ where p ($=0.05$ in this study) is a factor used to reduce the weights given to high intensity reflections, which are more subject to systematic errors.

Programs for the CDC 6600 and CDC 7600 used in the solution and analysis of this structure were: MAGPIK, INCOR and ORDER, general data reduction programs; ABSOR, our modified version of Hamilton's GON09 absorption correction program; FORDAP, A. Zalkin's Fourier analysis program; LSLONG and LESQ, our unpublished modifications of the Ganzel-Sparks-Trueblood least-squares program; DISMAT, a crystallographic distances and angles program using the correlation matrix from least-squares to calculate standard deviations; ORTEP, C. Johnson's (1965) thermal ellipsoid crystallographic plotting program; LSPLAN, our modification of the least-squares planes program from the University of Pittsburgh; and LIST1 and LISTAP, data presentation programs.

B. Determination and Refinement

A three-dimensional Patterson map was calculated, and the top four peaks other than the origin were interpreted as the interactions of the two independent Sr atoms. Other peaks were interpreted to give the positions of six oxygen atoms in an arrangement very similar to that encountered for $\text{Ca}(\text{DPP})_2(\text{C}_2\text{H}_5\text{OH})_{1/2}$. The coordinates and isotropic thermal parameters of these eight atoms and an overall scale factor were refined for three cycles of full-matrix least-squares to an R value ($R_1 = \Sigma |\Delta F| / \Sigma |F_o|$) of 0.34. A Fourier map revealed an additional 21 peaks interpretable as portions of DPP ligands. Three more cycles of least-squares on 29 atoms brought R_1 to 0.29. Another Fourier showed many peaks in unreasonable positions, including two peaks very close to the origin. It was decided that the original choice of oxygen atom positions might have biased the Fourier calculations, and a Fourier map was phased on the results from least-squares on the Sr atoms alone ($R_1 = 0.36$). This map was similar to the first, but allowed the positioning of 40 atoms, including the Sr atoms. These refined to an R_1 of 0.27 and a Fourier revealed the positions of all of the rest of the DPP atoms and a peak near Sr(1) which was assumed to be the oxygen of an ethanol, in keeping with the similarity between this structure and the one for the Ca complex. Three cycles of least-squares on these atoms, all refined with isotropic thermal parameters, brought R_1 to 0.14. A difference Fourier showed two peaks close to each of the Sr atoms, representing anisotropic thermal motion, and peaks near the atom labeled O(9) which suggested that O(9)

might have migrated to the center of gravity of what was still considered as an ethanol. (Due to the low resolution of the data ($d_{\min} = 1.64 \text{ \AA}$ for $2\theta = 25^\circ$, the effective limit of the data), the refinement did not distinguish well between atoms and groups of atoms, a factor which made interpretation of the Fourier maps tricky, and which accounts for poor agreement in bond distances, etc.) The O(9) atom was relabeled as C(61) and the two other peaks assigned labels of O(9) (close to Sr(1)) and C(62) (the methyl carbon). When these atoms were refined in least-squares for four cycles, R_1 reduced to 0.115. The coordinates of O(9), C(61) and C(62) refined so that the atoms formed an approximately equilateral triangle with edge 1.9 Å. A difference Fourier showed the anisotropy peaks noted above and one peak in the center of the triangle. Refinement of the structure with the Sr atoms anisotropic and the "ethanol" disordered gave an R_1 of 0.094 with poor refinement for the "ethanol" atoms. A difference Fourier phased after removal of the "ethanol" atoms ($R_1 = 0.13$) showed a large triangular area of electron density with a peak at each corner and high density in the center. On the assumption of similarity to the Ca structure, a disordered ethanol model was hand fitted to the peaks and the positions of the six atoms with occupancy factor of 0.5 were fixed in least-squares. Refinement of the rest of the structure and of four thermal parameters for the ethanol gave an R_1 of 0.094 and R_2 ($\sum w(\Delta F)^2 / \sum w F_o^2$) of 0.080. However, the thermal parameters of the methyl carbons of the two half-ethanols refined to the lowest values of any in the structure. It was then pointed out

that the triangular region of electron density could also be fitted by an acetone molecule and this possibility was tried. The positions of the disordered ethanol were relabeled and given full occupancy, and the positions allowed to refine in addition to the thermal parameters. Four cycles of least-squares brought R_1 to 0.088, R_2 to 0.072, and all shifts to less than 10% of their standard deviations. The thermal parameters of all of the acetone atoms were in keeping with those of the rest of the structure, and the refinement was considered complete at that point. A difference Fourier showed no peaks greater than $0.6 \text{ e}^-/\text{\AA}^3$. No attempt was made to locate the hydrogen atoms.

The observed F's and their standard deviations are compared to the final differences in Table 1. The final positional and thermal parameters of the atoms are given in Table 2.

Table 1.

Observed structure factors, standard deviations and final differences. Reflections marked with an * were given zero weight in least-squares.

Table 2.

Final Coordinates and Thermal Parameters. $\text{Sr}(\text{DPF})_2((\text{CH}_3)_2\text{CO})_{1/2}^{\text{a,b}}$

(a) Standard deviations of the least significant digit(s) are given in parentheses.

(b) The form of the temperature factor (B in units of $\text{\AA}^2$) is

$$T = \exp(-.25(B_{11}h^2a^{*2} + B_{22}k^2b^{*2} + B_{33}l^2c^{*2} + 2B_{12}hka^*b^* + 2B_{13}hla^*c^* + 2B_{23}k\ell b^*c^*)) \text{ for anisotropic and}$$

$$T = \exp(-B \sin^2\theta/\lambda^2) \text{ for isotropic thermal parameters.}$$

ATOM	X	Y	Z	811	822	833	812	813	823
S(1)	.2002(3)	-.0410(2)	.0449(2)	7.4(3)	4.5(2)	4.4(2)	3.1(2)	.4(2)	.5(2)
S(2)	.0158(3)	.1118(2)	.1623(2)	7.5(3)	4.2(2)	2.6(2)	2.5(2)	.5(2)	.1(1)
O(1)	.178(1)	.016(2)	-.088(1)	5.0(5)					
O(2)	.183(1)	.139(2)	.089(1)	6.3(5)					
C(101)	.241(3)	.003(3)	-.246(2)	8.6(10)					
C(102)	.263(3)	.005(3)	-.334(3)	12.3(12)					
C(103)	.284(3)	.101(4)	-.355(3)	15.8(13)					
C(104)	.326(3)	.205(4)	-.293(3)	13.3(13)					
C(105)	.298(3)	.203(3)	-.203(2)	10.2(11)					
C(106)	.265(3)	.110(3)	-.185(2)	8.4(10)					
C(107)	.235(2)	.113(3)	-.094(2)	7.2(9)					
C(108)	.274(2)	.211(2)	-.023(2)	5.0(7)					
C(109)	.240(2)	.218(2)	-.062(2)	5.4(8)					
C(110)	.275(2)	.351(2)	.126(2)	5.0(8)					
C(111)	.293(2)	.416(2)	.094(2)	5.3(8)					
C(112)	.334(2)	.529(2)	.163(2)	6.7(8)					
C(113)	.342(2)	.537(3)	.250(2)	7.8(9)					
C(114)	.322(2)	.449(3)	.275(2)	7.0(9)					
C(115)	.289(2)	.341(2)	.215(2)	6.5(8)					
O(3)	-.035(1)	.066(1)	-.125(1)	5.0(5)					
O(4)	-.027(1)	.189(2)	.059(1)	6.4(5)					
C(116)	-.026(3)	.068(3)	-.293(2)	8.3(9)					
C(117)	-.013(3)	.070(3)	-.383(3)	11.6(12)					
C(118)	.043(3)	.185(4)	-.399(3)	12.6(13)					
C(119)	.098(3)	.262(3)	-.326(3)	11.9(12)					
C(120)	.082(3)	.256(3)	-.237(2)	9.7(11)					
C(121)	.020(3)	.163(3)	-.226(2)	7.2(9)					
C(122)	.002(2)	.159(3)	-.138(2)	6.7(9)					
C(123)	.016(2)	.256(2)	-.066(2)	4.3(7)					
C(124)	.004(2)	.265(2)	.022(2)	5.0(8)					
C(125)	.024(2)	.379(2)	.089(2)	5.4(8)					
C(126)	.078(2)	.472(3)	.001(2)	6.9(8)					
C(127)	.104(2)	.576(2)	.128(2)	7.3(9)					
C(128)	.070(3)	.583(3)	.212(2)	8.4(10)					
C(129)	.020(3)	.406(3)	.234(2)	9.1(10)					
C(130)	-.003(2)	.381(3)	.168(2)	7.4(9)					
O(5)	.260(1)	-.078(1)	.171(1)	6.5(5)					
O(6)	.259(1)	-.187(1)	-.010(1)	6.7(5)					
C(32)	.257(3)	-.113(3)	.324(2)	9.7(10)					
C(32)	.270(3)	-.112(3)	.420(3)	12.5(12)					
C(33)	.360(3)	-.126(3)	.443(2)	10.6(11)					
C(34)	.435(3)	-.139(3)	.392(3)	13.5(14)					
C(35)	.428(3)	-.135(3)	.300(2)	10.6(11)					
C(36)	.336(3)	-.125(2)	.281(2)	6.6(8)					
C(37)	.311(2)	-.129(2)	.185(2)	6.0(8)					
C(38)	.344(2)	-.199(2)	.118(2)	6.2(8)					
C(39)	.315(2)	-.221(2)	.026(2)	5.7(8)					
C(40)	.337(3)	-.319(2)	-.039(2)	6.1(8)					
C(41)	.425(3)	-.356(3)	-.015(2)	8.5(10)					
C(42)	.440(3)	-.423(3)	-.080(3)	10.7(11)					
C(43)	.376(3)	-.478(3)	-.157(2)	9.1(10)					
C(44)	.298(3)	-.461(3)	-.186(2)	8.0(10)					
C(45)	.271(3)	-.444(3)	-.118(3)	10.1(10)					
O(7)	.129(1)	.177(1)	.302(1)	5.6(5)					
O(8)	-.048(1)	.219(1)	.283(1)	5.0(5)					
C(46)	.301(3)	.159(3)	.410(2)	8.8(10)					
C(47)	.399(3)	.232(3)	.463(2)	9.6(10)					
C(48)	.442(3)	.329(3)	.536(2)	10.1(11)					
C(49)	.387(3)	.390(3)	.555(2)	10.7(11)					
C(50)	.285(3)	.369(3)	.511(2)	9.0(10)					
C(51)	.252(3)	.268(3)	.434(2)	5.7(8)					
C(52)	.149(3)	.251(2)	.387(2)	6.8(9)					
C(53)	.073(3)	.286(2)	.414(2)	5.7(8)					
C(54)	-.016(3)	.271(2)	.370(2)	6.0(8)					
C(55)	-.105(3)	.309(2)	.404(2)	7.9(10)					
C(56)	-.087(3)	.351(2)	.502(2)	9.7(10)					
C(57)	-.188(4)	.391(3)	.537(2)	11.0(12)					
C(58)	-.248(3)	.387(2)	.488(2)	8.2(9)					
C(59)	-.258(3)	.347(3)	.395(3)	12.2(12)					
C(60)	-.177(3)	.311(2)	.357(2)	8.2(10)					
O(9)	.404(2)	.083(2)	.075(2)	13.3(8)					
C(61)	.479(4)	.104(4)	.134(3)	13.6(15)					
C(62)	.565(4)	.075(4)	.141(3)	16.1(15)					
C(63)	.474(3)	.202(3)	.214(3)	14.6(13)					

C. Results and Discussion

The complex consists of clusters of composition $(\text{Ca}_2(\text{DPP})_4(\text{CH}_3)_2\text{CO})_2$ around the center of symmetry at $\underline{0,0,0}$. There are two different Sr environments. The clusters are related only by unit cell translations and there is only one inter-cluster contact of less than $3.50 \text{ \AA}$, between C(26) and C(26) across the center of symmetry at $\underline{0, \frac{1}{2}, 0}$ ($3.42 \text{ \AA}$).

The Sr(1) and Sr(2) atoms are respectively seven and six-coordinated by the oxygens of the DPP ligands and the acetone (Figure 1). Each Sr atom shares two oxygen atoms with each of two neighboring Sr atoms, forming a cluster around the center of symmetry. The bridging oxygens, O(1) and O(2) of DPP(1) and O(3) and O(4) of DPP(2), occupy the corners of a slightly distorted parallelepiped, with the two rectangular faces capped by Sr atoms and the non-rectangular face occupied by the atoms of the DPP ligands (Figure 2). The uncapped face, core plane 1 (CP 1), forms almost right angles with the planes capped by Sr(1) and Sr(2) (CP 2 and CP 3 respectively). The acute angle between CP 2 and CP 3 is 83° , significantly different from right angles. The core planes are planar to less than a standard deviation of the coordinates of the four atoms that make each of them up. The distances between symmetry-related planes is $3.04 \text{ \AA}$ for CP 1, $3.30 \text{ \AA}$ for CP 2 and $2.83 \text{ \AA}$ for CP 3. The mean plane of the Sr atoms lies half way between CP 1 and CP 1'. Sr(1) is $1.50 \text{ \AA}$ away from CP 2 and the three other oxygens coordinated to Sr(1) have distances of $3.25 \text{ \AA}$, $3.17 \text{ \AA}$ and $3.21 \text{ \AA}$ for O(5), O(6) and O(9) respectively, forming an almost parallel plane. Sr(2) is

Figure 1.

Stereoscopic view of the coordination around the Sr atoms.

The center of symmetry at 0,0,0 is indicated. Thermal ellipsoids have been scaled to include 50% probability.

XBL 724-793

Figure 2.

Stereoscopic view of the complex cluster. Phenyl carbons except for those attached directly to the diketone have been omitted and the acetone atoms have been given artificial thermal parameters for clarity. View direction and scaling of the thermal ellipsoids are the same as in Figure 1.

XBL 724-792

1.14 Å from CP 3 with O(7) and O(8) of DPP(4) at distances of 3.08 Å and 3.24 Å respectively, indicating steric interaction between DPP(4) and DPP(3), forcing O(8) away from CP 3 and O(7) towards it.

The distances from Sr to the bridging oxygens average to 2.55 Å with no real difference between the two differently coordinated Sr atoms. The closer distance from CP 3 to Sr(2) is due to the fact that the oxygen atoms making up CP 3 are further apart on the average than those making up CP 2 (Table 3). The distances from the Sr atoms to the unshared oxygens average to 2.44 Å, indicating the expected tighter bonding to the unshared ligands, and the Sr(1) to O(9) distance is 2.60 Å, in keeping with the fact that the acetone is uncharged and thus not as tightly bound as the negative DPP ligands.

The angles of the diketone planes of the bridging DPP molecules to CP 1 are 89° for DPP(1) and 108° for DPP(2) with an angle between the two of 17°. The larger angle for DPP(2) reflects the less crowded environment it encounters (Figure 2), whereas DPP(1) is forced to near perpendicularity with CP 1 by interaction with the acetone molecule. DPP(3) forms an angle of 71° with CP 2, and the acetone plane forms a right angle with the same plane, again reflecting the steric interaction noted before. The diketone of DPP(4) forms an angle of 86° with CP 3, reflecting its nearly symmetrical environment with respect to CP 1 and CP 1'. The tilt of DPP(4) within that plane has already been noted.

Table 3.

Bonded and non-bonded distances (Å) between strontium and oxygen in the coordination cluster.^a (See Figure 1)

Bonded			Non-bonded		
Atom(1)	Atom(2)	Distance	Atom(1)	Atom(2)	Distance
Sr(1)	O(1)	2.55(2)	O(1)	O(2)	2.83(2) ^{b,c}
Sr(1)	O(2)	2.54(2)	O(1)	O(3)	3.37(2) ^c
Sr(1)	O(3)	2.57(2)	O(1)	O(4)	3.04(3) ^c
Sr(1)	O(4)	2.58(2)	O(2)	O(3)	3.04(3) ^c
Sr(1)	O(5)	2.46(2)	O(2)	O(4)	3.30(3) ^c
Sr(1)	O(6)	2.43(2)	O(3)	O(4)	2.86(2) ^{b,c}
Sr(1)	O(9)	2.60(3)	O(1)	O(8)	3.61(3)
Sr(2)	O(1)	2.59(2)	O(1)	O(9)	3.67(3)
Sr(2)	O(2)	2.53(2)	O(2)	O(7)	3.42(3)
Sr(2)	O(3)	2.53(2)	O(2)	O(9)	3.43(3)
Sr(2)	O(4)	2.47(2)	O(3)	O(5)	3.27(3)
Sr(2)	O(7)	2.42(2)	O(3)	O(7)	3.48(2)
Sr(2)	O(8)	2.47(2)	O(4)	O(6)	3.26(3)
			O(4)	O(8)	3.58(3)
			O(5)	O(6)	2.85(2) ^b
			O(5)	O(9)	3.18(3)
			O(6)	O(9)	3.36(3)
			O(7)	O(8)	2.76(2) ^b

(a) Standard deviations given in parentheses.

(b) Bite of the ligand.

(c) Edge of the central oxygen parallelopiped.

The DPP ligands can be described in terms of three planes each, the two phenyl rings (phenyl(1), 1-6; phenyl(2), 10-15, Figure 3) and the diketone residue (7-9, oxygen 1 and 2, Figure 3). All such planes in this structure are planar to within two standard deviations of the coordinates of the atoms involved. The plane of the acetone molecule is also good to two standard deviations. The phenyl planes are twisted along the 6-7 and 9-10 axes with respect to the diketone residue planes by angles ranging from 10° to 45° (Table 4).

Due to the low resolution of the data noted above, the distances within the DPP ligands (Table 5) are not well defined. The average C-C distance in the phenyl groups of 1.41 Å is close to other values where the hydrogens have been omitted, and the other average distances, C-O = 1.28 Å, C-C(diketone) = 1.40 Å, C(phenyl)-C(diketone) = 1.53 Å, are in reasonable agreement with other values for this ligand (Sections III-V). The trends visible in the acetone distances are also in chemical accord.

Further discussion of this structure is given in Section VII.

Figure 3.

Generalized labeling system for DPP ligands. Carbon atoms are single circles, oxygen atoms are double circles. Numbering shown is for DPP(1), for successive DPP molecules add 2 to the oxygen numbers and 15 to the carbon numbers.

Table 4.

Rotation of phenyl ring planes with respect to diketone planes and each other (deg).

DPP	Phenyl 1 to diketone.	Phenyl 2 to diketone.	Phenyl 1 to Phenyl 2.
1	16.0	27.2	14.6
2	19.6	10.4	18.4
3	32.8	25.3	45.0
4	22.8	14.8	37.1

Table 5.

Intra-molecular bond distances (Å). $\text{Sr}(\text{DPP})_2((\text{CH}_3)_2\text{CO})_{1/2}^{\text{a}}$

DPP(1)			DPP(2)		
Atom(1)	Atom(2)	Distance	Atom(1)	Atom(2)	Distance
C(01)	C(02)	1.45(4)	C(16)	C(17)	1.47(4)
C(02)	C(03)	1.41(4)	C(17)	C(18)	1.37(4)
C(03)	C(04)	1.39(5)	C(18)	C(19)	1.42(4)
C(04)	C(05)	1.51(4)	C(19)	C(20)	1.45(4)
C(05)	C(06)	1.34(4)	C(20)	C(21)	1.34(4)
C(06)	C(01)	1.44(4)	C(21)	C(16)	1.34(3)
C(06)	C(07)	1.51(3)	C(21)	C(22)	1.45(3)
C(07)	O(1)	1.32(3)	C(22)	O(3)	1.30(3)
C(07)	C(08)	1.41(3)	C(22)	C(23)	1.44(3)
C(08)	C(09)	1.43(3)	C(23)	C(24)	1.40(3)
C(09)	O(2)	1.31(3)	C(24)	O(4)	1.28(3)
C(09)	C(10)	1.50(3)	C(24)	C(25)	1.56(3)
C(10)	C(11)	1.38(3)	C(25)	C(26)	1.45(4)
C(11)	C(12)	1.52(3)	C(26)	C(27)	1.44(3)
C(12)	C(13)	1.36(3)	C(27)	C(28)	1.41(4)
C(13)	C(14)	1.35(3)	C(28)	C(29)	1.46(4)
C(14)	C(15)	1.43(3)	C(29)	C(30)	1.46(4)
C(15)	C(10)	1.40(3)	C(30)	C(25)	1.32(3)
DPP(3)			DPP(4)		
C(31)	C(32)	1.54(4)	C(46)	C(47)	1.43(4)
C(32)	C(33)	1.40(4)	C(47)	C(48)	1.42(4)
C(33)	C(34)	1.34(4)	C(48)	C(49)	1.33(4)
C(34)	C(35)	1.51(4)	C(49)	C(50)	1.44(4)
C(35)	C(36)	1.36(4)	C(50)	C(51)	1.48(3)
C(36)	C(31)	1.34(4)	C(51)	C(46)	1.36(3)
C(36)	C(37)	1.56(3)	C(51)	C(52)	1.51(3)
C(37)	O(5)	1.25(3)	C(52)	O(7)	1.39(3)
C(37)	C(38)	1.44(3)	C(52)	C(53)	1.34(3)
C(38)	C(39)	1.42(3)	C(53)	C(54)	1.32(4)

Table 5 (continued)

Atom(1)	Atom(2)	Distance	Atom(1)	Atom(2)	Distance
C(39)	O(6)	1.27(3)	C(54)	O(8)	1.33(3)
C(39)	C(40)	1.61(3)	C(54)	C(55)	1.57(4)
C(40)	C(41)	1.40(4)	C(55)	C(56)	1.47(4)
C(41)	C(42)	1.44(4)	C(56)	C(57)	1.48(4)
C(42)	C(43)	1.33(4)	C(57)	C(58)	1.32(4)
C(43)	C(44)	1.39(4)	C(58)	C(59)	1.41(4)
C(44)	C(45)	1.59(4)	C(59)	C(60)	1.46(4)
C(45)	C(40)	1.37(3)	C(60)	C(55)	1.26(4)
Acetone					
O(9)	C(61)	1.28(4)			
C(61)	C(62)	1.41(5)			
C(61)	C(63)	1.60(5)			

(a) Standard deviations in parentheses.

D. References

- Cromer, D.T. and Liberman, D. (1970). J. Chem. Phys., 53(5), 1891.
- Doyle, P.A. and Turner, P.S. (1968). Acta Cryst., A24, 390.
- Hammond, G.S., Nonhebel, D.C. and Wu, C-H.S. (1963). Inorg. Chem., 2(1), 73.
- Johnson, C.K. (1965). Oak Ridge National Report-3794, Revised, Oak Ridge, Tenn.

VII. Summary and Comparison

A. Introduction

Various aspects of the structures of 1,3-diphenyl-1,3-propanedione (HDPP), $\text{Mg}(\text{DPP})_2(\text{DMF})_2$, $\text{Ca}(\text{DPP})_2(\text{C}_2\text{H}_5\text{OH})_{\frac{1}{2}}$ and $\text{Sr}(\text{DPP})_2((\text{CH}_3)_2\text{CO})_{\frac{1}{2}}$, which were covered in Sections III - VI, will be compared. Comparisons of these complexes with various β -diketone complexes of divalent metals in the literature will be made, as will certain correlations on the structure of the DPP^- ligand.

B. $\text{Mg}(\text{DPP})_2(\text{DMF})_2$ and Other Octahedra.

The octahedral coordination of Mg in $\text{Mg}(\text{DPP})_2(\text{DMF})_2$ (Section IV) is similar to that exhibited by other β -diketone complexes of divalent metals, e.g. diaquobis(acetylacetonato)-magnesium, $\text{Mg}(\text{AA})_2(\text{H}_2\text{O})_2$ (Morosin, 1967), $\text{Co}(\text{AA})_2(\text{H}_2\text{O})_2$ (Bullen, 1959) and $\text{Ni}(\text{AA})_2(\text{H}_2\text{O})_2$ (Montgomery and Lingafelter, 1964). In each case, the divalent metal cation is octahedrally coordinated by two β -diketone ligands and two solvent molecules in a monomeric unit. In all of these cases, the oxygen atoms in the solvent molecules are significantly further away from the metal ion than are the ligand oxygen atoms. For the $\text{Mg}(\text{DPP})_2$ complex, the distortion (2.06, 2.06, 2.10 Å) is similar to the distortion for the acetylacetonato complexes of Mg (2.03, 2.04, 2.15 Å), Co (2.05, 2.06, 2.23 Å) and Ni (2.02, 2.01, 2.14 Å). This effect is also noted in a dimeric situation with octahedral coordination in $\text{Co}(\text{AA})_2(\text{H}_2\text{O})$ (Cotton and Elder, 1966), and in cases where the coordination is not octahedral, as in the Ca and Sr DPP complexes (Section V, VI) and the seven-coordinate $\text{Ho}(\text{DPP})_3(\text{H}_2\text{O})$ (Zalkin, *et al.*, 1969).

The octahedral complex $\text{Ni}(\text{HAA})_2(\text{H}_2\text{O})_2(\text{ClO}_4)_2$ (Anzenhofer and Hewitt, 1971), in which all the ligands on the Ni^{++} are neutral, does not show this distortion. The distances in Anzenhofer's structure are 2.07, 2.02, 2.04 Å and 2.04, 2.03, 2.05 Å for the three independent distances around the two different Ni atoms, the distance to the water given last in each trio.

The $\text{Mg}(\text{DPP})_2$ complex differs from the other octahedral

complexes, however, in that the solvent molecules are cis to one another on the coordination octahedron (Figure, p.89). The solvent molecules are all trans to one another in the other monomeric complexes. The cis arrangement is not required by the space group symmetry, which could just as easily accomodate the trans configuration, but, while all the other distorted octahedra could be described as tetragonally distorted, the octahedron in $\text{Mg}(\text{DPP})_2(\text{DMF})_2$ cannot.

Bullen explains the distortion of the $\text{Co}(\text{AA})_2(\text{H}_2\text{O})_2$ octahedron in terms of electronic combinations of the available d-orbitals of the Co. The same explanation could hold for the Ni complex as well, but not for the magnesium complexes, since the Mg has no available d-orbitals. Morosin concludes in his paper that the effect is due to packing forces rather than electronic ones since the diaquo Mg complex shows the same distortion as the Co and Ni complexes. The appearance of the same effect in $\text{Mg}(\text{DPP})_2(\text{DMF})_2$ and in other structures noted above, where the packing and coordination environments are radically different, as well as the absence of any effect in the octahedral nickel acetylacetonate perchlorate complex, leads me to suspect that the effect is primarily electrostatic in origin, with secondary contributions from d-orbital hybridization and packing effects.

C. $\text{Ca}(\text{DPP})_2(\text{C}_2\text{H}_5\text{OH})_{\frac{1}{2}}$, $\text{Sr}(\text{DPP})_2((\text{CH}_3)_2\text{CO})_{\frac{1}{2}}$ and Other Polymers.

1. Comparison

The two DPP complexes of Ca and Sr are not isomorphous in a compositional sense; one contains an ethanol and the other an acetone as a solvation molecule. However, the environments of the Ca and the Sr atoms are relatively similar and the basic shape of the polymeric complex is roughly the same. Despite this, the unit cells of the two structures are not of similar shape. The primary differences of molecular geometry lie in the degree of distortion of the central parallelepiped of O atoms (Figures, pp. 117, 145).

The central parallelepiped of oxygen atoms is strongly distorted from rectangularity in the Ca complex, while in the Sr complex the acute angle between the rectangular faces is much closer to 90° (69° for the Ca and 83° for the Sr complex). The planes that make up the central parallelepiped are closer together in the Ca compound than they are in the Sr complex (2.90, 2.87 and 2.60 Å vs. 3.04, 3.30 and 2.83 Å). The wider range in the distances between planes in the Sr complex is also reflected in the wider range of distances on the edges of the parallelepiped (2.83 - 3.37 Å for Sr vs. 2.77 - 3.10 Å for Ca).

All of these effects can be explained as due to the fact that the bonds between the Ca and the oxygens are shorter than those between the Sr and the oxygens. The differences in the Ca-O and Sr-O distances are compatible with the difference in ionic radius between Ca and Sr ($\text{Ca-O}_{\text{ave}} = 2.37$ Å, $\text{Sr-O}_{\text{ave}} = 2.55$ Å; $r_{\text{Ca}} = 0.99$ Å, $r_{\text{Sr}} = 1.13$ Å) (Pauling, 1960), and amount to almost

+0.2 Å.

Given the coordination geometry, shorter bonds between the metal and the bridging oxygens mean that the DPP ligands doing the bridging must move closer together (Figures, pp. 119, 147). In the Ca complex they are forced close enough together that they must slide parallel to one another to achieve a staggered configuration and increase the distances between the atoms, thus decreasing the acute angle of the parallelogram of core plane 1 (pp. 115, 143). In the case of the Sr complex, the distances from the Sr to the oxygen allow for enough room between DPP ligands so that they are not forced into close proximity and can assume a more eclipsed configuration with respect to one another.

The reason that $\text{Mg}(\text{DPP})_2$ does not polymerize in the same manner as the Ca and Sr complexes is thus apparent, since the Mg-O distances of around 2.05 Å would have required too much forcing of the DPP ligands towards one another even with full staggering of the diketone residues.

2. Contrast

The form of the novel polymerization of the Ca and Sr DPP complexes has been thoroughly described in Sections V and VI. Previous studies of polymeric β -diketone complexes of divalent metals have generally been conducted using acetylacetonate anion (AA^-) as the ligand, and the structures are relatively straightforward.

The general rationale of the acetylacetonate polymers seems to be the achievement of octahedral coordination for the metal

ions. $\text{Ni}(\text{AA})_2$ forms trimers, each Ni atom being surrounded octahedrally, and the octahedra sharing faces. The Ni atoms lie in a straight line (Bullen, *et al.*, 1961). $\text{Co}(\text{AA})_2$ forms a tetrameric linear chain with octahedra sharing faces between the two Co atoms at each end and sharing an edge between the two central Co atoms (Cotton and Elder, 1965). $\text{Co}(\text{AA})_2(\text{H}_2\text{O})$ dimerizes so that each cobalt can be octahedrally coordinated, and the octahedra share an edge (Cotton and Elder, 1966).

Gerlach and Holm (1969) note that β -diketones with bulky side groups act to suppress polymerization and force the complexes to become monomeric even though octahedral coordination may be lost. Thus the structures of bis(2,2,6,6-tetramethyl-3,5-heptanedionato)zinc and nickel ($\text{Zn}(\text{DTM})_2$ and $\text{Ni}(\text{DTM})_2$) have been determined by crystallographic means to be monomeric and tetrahedral and planar respectively (Cotton and Wood, 1964; Cotton and Wise, 1966). Other indications show that the structures of the DTM complexes of other divalent transition metals are monomeric and either tetrahedral or planar (Gerlach and Holm, 1969).

The results of the studies in Sections V and VI certainly indicate that bulky side groups cannot be counted on to reduce polymerization, even though the standard advantages are not present (i.e. none of the ions manage to achieve octahedral or other symmetrical coordination). The existence of the dimer in the study of $\text{Pr}(\text{DTM})_3$ (Erasmus and Boeyens, 1970), which involves sharing DTM ligands to achieve seven-fold coordination, indicates that this ability to polymerize is not specific to DPP. Further,

a recent study of the DPM complexes of the alkaline-earth metals by means of vapor-phase diffusion, mass spectroscopy and molecular weight measurements indicate that the Ca, Sr and Ba complexes are polymeric in solution and in the vapor phase, but that the Mg complex is monomeric (Schwarberg, et al., 1970).

What these results suggest is that a combination of the bulk of the ligand and the length of the M-O bond is the criterion for whether or not polymerization is favorable. With a short M-O bond length, as with Mg or the divalent first-row transition metals, ligands with bulky side groups such as t-butyl (DPM) or phenyl (DPP) are forced too close to one another in the configurations necessary for polymerization. When the M-O distance increases, as for Ca, Sr, or Pr(III), then neighboring ligands need not be so close together, and the advantages of sharing ligands so as to get greater coordination from the negative ions can be realized.

D. A Note on the DPP^- Ligand.

Three structures involving the DPP^- ligand were performed which were accurate enough to permit comparison of bond lengths. These were the structures of HDPP (Section III (Williams, 1966)), $\text{Mg}(\text{DPP})_2(\text{DMF})_2$ (Section IV) and $\text{Ca}(\text{DPP})_2(\text{C}_2\text{H}_5\text{OH})_2$. The structure of $\text{Sr}(\text{DPP})_2((\text{CH}_3)_2\text{CO})_{1/2}$ was not well enough determined to give the sort of comparison needed. There have been no published structures involving DPP^- with high enough accuracy either.

In all three of these structures it was noted that the bonds in the phenyl rings varies systematically depending on whether they were near to or far from the juncture to the β -diketone. Specifically the bonds to the para carbon in the phenyl group were shorter than the other C-C bonds in the ring (Table 1).

Table 1.

C-C distances in phenyl groups (Å).^a

	HDPP	$\text{Mg}(\text{DPP})_2$	$\text{Ca}(\text{DPP})_2$
<u>m-p</u>	1.37 - 1.37	1.36 - 1.37	1.363(2)
<u>o-m</u>	1.38 - 1.38	1.37 - 1.38	1.382(2)
<u>l-o</u>	1.38 - 1.40	1.38 - 1.39	1.379(3)

(a) Standard deviations of individual measurements are ± 0.003 Å for HDPP, ± 0.006 Å for Mg and ± 0.005 Å for Ca; the ranges are given for HDPP and Mg, and the average over 16 independent measurements given for Ca.

It is possible that this effect could be due to thermal motion, but the differences in the thermal motions of the l carbon and the p carbon is not that great in the three structures (Tables, pp. 60, 86, 113). It is more likely to be a steric effect due to the same interaction which causes the interior

angle on the 1 carbon of the phenyl groups to be significantly less than 120° (118° average) in all three of the above structures.

The bite of the ligand, the distance between the two oxygen atoms on the same molecule, is constant over the complex series from Mg to Sr and is the same as the bite determined for other β -diketon complexes. The average bite is 2.80 Å. This value is equal to the sum of the crystal radii for the two oxygens (Pauling, 1960). The O-C-C and C-C-C angles in the diketone open up from the expected 120° to, on the average, 125° to allow this O-O distance.

In HDPP, the value of the bite is much closer to the "ideal" value, based on 120° angles, of 2.41 Å, due to a strong internal hydrogen bond. The hydrogen bond reduces the bite from 2.80 Å to 2.46 Å, overcoming the steric interaction of the oxygens.

E. References

- Anzenhofer, K. and Hewitt, T.G. (1971). Zeit. für Krist., 134, 54.
- Bullen, G.J. (1959). Acta Cryst., 12, 703.
- Bullen, G.J., Mason, R. and Pauling, P. (1961). Nature, 189, 291.
- Cotton, F.A. and Elder, R.C. (1965). Inorg. Chem., 4(8), 1145.
- Cotton, F.A. and Elder, R.C. (1966). Inorg. Chem., 5(3), 423.
- Cotton, F.A. and Wise, J.J. (1966). Inorg. Chem., 5(7), 1201.
- Cotton, F.A. and Wood, J.S. (1964). Inorg. Chem., 3(2), 245.
- Erasmus, C.S. and Boeyens, J.C.A. (1970). Acta Cryst., B26, 1843.
- Gerlach, D.H. and Hom, R.H. (1969). Inorg. Chem., 8(11), 2292.
- Montgomery, H. and Lingafelter, E.C. (1963). Acta Cryst., 17, 1481.
- Morosin, B. (1967). Acta Cryst., 22, 316.
- Pauling, L. (1960). "The Nature of the Chemical Bond", Third Ed., Cornell University Press, Ithaca, N.Y.
- Schwarberg, J.E., Sievers, R.E. and Moshier, R.W. (1970). Anal. Chem., 42(14), 1828.
- Williams, D.E. (1966). Acta Cryst., 21, 340.
- Zalkin, A., Templeton, D.H. and Karraker, D.G. (1969). Inorg. Chem., 8(12), 2680.