12, 384

What Can We Measure in $\rho - \omega$ Interference Experiments?*

A. Rabl and N. W. Reay

Department of Physics
The Ohio State University
Columbus, Ohio 43210

ABSTRACT

We show that $\rho - \omega$ interference, if studied in pionic production of both 2π and 3π final states, may allow an experimental separation of the phase of the mixing parameter from the phase difference of the ρ and ω production amplitudes. We point out that the mixing parameter measured in 2π final states differs from the one measured in 3π final states, unless certain isospin breaking amplitudes are sufficiently small.

MASTER

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

* Research supported in part by the U.S. Atomic Energy Commission under Contract AT(11-1)-1545

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

In recent years, $\rho - \omega$ interference has been studied both in electromagnetic and in pionic production of ρ -mesons.¹⁻⁴ Pionic production experiments have so far yielded little detailed information because of uncertainties both in production phases and in the sharing of interference effects between participating helicity amplitudes. In this note, we discuss what can be learned by combining measurements of two and three pion final states in pionic experiments, and we pay special attention to the possible contributions of isospin violating amplitudes. These isospin violations are customarily assumed to be of electromagnetic origin and hence of order α ; in fact, until now discussions of $\rho - \omega$ interference⁵⁻⁹ have relied heavily on this assumption. We suggest that this point can be tested experimentally. Specifically, if isospin breaking terms in the $\rho - \omega$ system are unexpectedly large (say more than about $1/30$ the size of the isospin conserving terms) then significant discrepancies can be found between the values for the mixing parameter as determined from 2π and from 3π final states.

Let us start by defining a complex mixing parameter ϵ to relate the pure, unmixed isospin states ρ_0 and ω_0 to the physical vector mesons ρ and ω

$$\begin{aligned} |\rho\rangle &= |\rho_0\rangle - \epsilon |\omega_0\rangle \\ |\omega\rangle &= \epsilon |\rho_0\rangle + |\omega_0\rangle \end{aligned} \quad (1)$$

(N.B. Throughout this paper, ρ and ρ_0 refer only to the neutral member of the ρ triplet.) The data summary in Ref.8 gives a value $|\epsilon| \approx \frac{1}{30}$.

To lowest order in ϵ , the amplitude for producing a final state f from an initial state i near the $\rho - \omega$ resonance region can be written as

$$T_{fi} = f_{\rho_0} f_{\rho_0} P_{\rho_0} A_{\rho_0 i} + f_{\omega_0} f_{\omega_0} P_{\omega_0} A_{\omega_0 i} + \epsilon (P_{\omega_0} - P_{\rho_0}) [f_{\rho_0} f_{\omega_0} A_{\omega_0 i} + f_{\omega_0} f_{\rho_0} A_{\rho_0 i}] \quad (2)$$

where A_{Vi} is the production amplitude for the vector meson V , f_{Vf} is the decay amplitude and $P_V(m) = [m_V^2 - im_V \Gamma_V - m^2]^{-1}$ is the propagator¹⁰ (V stands for the pure isospin states ρ_0 and ω_0). In most of the following formulas, we shall not indicate the initial state i explicitly.

This expression can be simplified with some reasonable approximations. First of all, the smallness of the width ratio $\Gamma_{\omega_0} / \Gamma_{\rho_0} \approx \Gamma_{\omega} / \Gamma_{\rho} \approx 1/11$ allows us to drop ϵP_{ρ_0} terms relative to ϵP_{ω_0} to get

$$T(\pi^+ \pi^-) = f_{\rho_0} 2\pi P_{\rho_0} A_{\rho_0} + (f_{\omega_0} 2\pi + \epsilon f_{\rho_0} 2\pi) P_{\omega_0} A_{\omega_0} + \epsilon f_{\omega_0} 2\pi P_{\omega_0} A_{\rho_0} \quad (3)$$

and

$$T(\pi^+ \pi^0 \pi^-) = f_{\omega_0} 3\pi P_{\omega_0} (A_{\omega_0} + \epsilon A_{\rho_0}) + f_{\rho_0} 3\pi P_{\rho_0} A_{\rho_0} \quad (4)$$

For the specific cases of 2π and 3π final states, we can make further approximations based on the smallness of the isospin-breaking amplitudes $f_{\rho_0} 3\pi$ and $f_{\omega_0} 2\pi$. They are, presumably, due to electromagnetism and hence of order α relative to $f_{\rho_0} 2\pi$ and $f_{\omega_0} 3\pi$. But since we want to test this assumption, we have to proceed more cautiously. What if for some reason isospin violations are anomalously large in the ρ - ω system, possibly due to some non-electromagnetic isospin-breaking interaction?¹¹ (Also, how big can "order α " be: 2α , 4α? And 4α is already as big as $|\epsilon|$). On the other hand, $|f_{\rho_0} 3\pi / f_{\omega_0} 3\pi|$ and $|f_{\omega_0} 2\pi / f_{\rho_0} 2\pi|$ could really not be very much larger¹² than $|\epsilon|$; $|f_{\omega_0} 2\pi / f_{\rho_0} 2\pi|$, for example, is obviously bounded

by $(\Gamma_\omega/\Gamma_\rho)^{\frac{1}{2}} \approx 1/3$. Therefore, we keep the isospin-breaking amplitudes explicitly except where they are multiplied by ϵ or by P_{ρ_0} ; in other words, we drop the last term in each of Eqs.(3) and (4).

As a result, we obtain the differential production cross-sections for the 2π final state

$$\frac{d\sigma}{dt}(2\pi) \propto \left| P_{\rho_0}^{(\pi)} + P_{\omega_0}^{(\pi)} \frac{|A_{\omega_0}|}{|A_{\rho_0}|} \left| \epsilon_{2\pi} \right| \exp(i\psi_{2\pi}) \right|^2 \quad (5)$$

with

$$\epsilon_{2\pi} \equiv \epsilon + \frac{f_{\omega_0 2\pi}}{f_{\rho_0 2\pi}} \quad (6)$$

and

$$\psi_{2\pi} \equiv \psi_{\epsilon 2\pi} + \psi_{\omega_0} - \psi_{\rho_0} \quad (7)$$

$\psi_{2\pi}$ is the experimentally measured 2π state phase, $\psi_{\epsilon 2\pi}$ is the phase of $\epsilon_{2\pi}$, and ψ_{ω_0} and ψ_{ρ_0} are the production phases, i.e. the phases of A_{ω_0} and A_{ρ_0} . In the 3π case, both the leading term A_{ω_0} and the ϵ correction come with the same propagator P_{ω_0} ; hence we lose no information if we integrate over the ω mass region.

$$\frac{d\sigma}{dt}(3\pi) \propto \int d\omega |T(\pi^+\pi^0\pi^-)|^2 \quad (8)$$

We find

$$\frac{d\sigma}{dt}(3\pi) \propto \left| A_{\omega_0} + \epsilon A_{\rho_0} \right|^2 \quad (9)$$

The interference effect in the 3π cross-section becomes measurable if a pair of charge symmetric reactions, e.g.

$$\pi^+ n \rightarrow (\omega_0, \rho_0) p \quad \xrightarrow{\quad} 3\pi \quad (10+)$$

and

$$\pi^- p \rightarrow (\omega_0, \rho_0) n \xrightarrow{\quad} 3\pi \quad (10-)$$

are compared. These reactions are related by a rotation $R = e^{i\pi I_2}$ in isospin space, and hence their respective cross-sections would be proportional to $\left| A_{\omega_0} \pm \epsilon A_{\rho_0} \right|^2$ if the production amplitudes conserved isospin.

But again, just as with the decay process, we want to keep an open mind about the possibility of an unexpectedly large isospin violating production amplitude. So we only assume that the transition operator T in the production amplitude splits into parts which are even and odd under the isospin rotation R

$$T = T_0 + T_1$$

with

$$R T_0 R^{-1} = T_0 \quad \text{and} \quad R T_1 R^{-1} = -T_1. \quad (11)$$

This is satisfied by any theoretical scheme (we obviously exclude weak interactions), including the u_3 term of Oakes and Sokolow.¹¹ We shall neglect ϵT_1 terms, which is reasonable, but not essential for deriving our results.

Splitting the production amplitudes accordingly, e.g.

$$A_{\omega_0+} = \langle \omega_0 p | T | \pi^+ n \rangle = \langle \omega_0 p | T_0 | \pi^+ n \rangle + \langle \omega_0 p | T_1 | \pi^+ n \rangle$$

$$A_{\omega_0-} = \langle \omega_0 n | T | \pi^- p \rangle = \langle \omega_0 p | T_0 | \pi^+ n \rangle - \langle \omega_0 p | T_1 | \pi^+ n \rangle$$

and analogously for $A_{\rho_0\pm}$, we find that the cross-sections for the reactions (10 $\pm$) are given by

$$\frac{d\sigma}{dt} (3\pi|\pm) \propto \left| A_{\omega_0+} \pm \epsilon_{3\pi} A_{\rho_0+} \right|^2 \quad (12)$$

with

$$\epsilon_{3\pi} \equiv \epsilon + \frac{\langle \omega_0 p | T_1 | \pi^+ n \rangle}{\langle \rho_0 p | T_0 | \pi^+ n \rangle} . \quad (13)$$

Therefore, the ratio

$$\frac{\frac{d\sigma}{dt} (3\pi|+)}{\frac{d\sigma}{dt} (3\pi|-)} = \frac{1 + 2 \frac{|A_{\rho_0}|}{|A_{\omega_0}|} |\epsilon_{3\pi}| \cos \psi_{3\pi}}{1 - 2 \frac{|A_{\rho_0}|}{|A_{\omega_0}|} |\epsilon_{3\pi}| \cos \psi_{3\pi}} \quad (14)$$

allows us to determine $|\epsilon_{3\pi}| \cos \psi_{3\pi}$ where $\psi_{3\pi}$ is the 3π final state phase

$$\psi_{3\pi} \equiv \psi_{\epsilon 3\pi} + \psi_{\rho_0} - \psi_{\omega_0} . \quad (15)$$

If one incorporates helicity frame density matrix elements to derive the ω transverse spin projection $\rho_{00} \frac{d\sigma}{dt}$, the ratio (14) may be as large as three in the small momentum transfer region $t \leq 0.1 \text{ GeV}^2$ where

$$\left| A_{\rho_0} / A_{\omega_0} \right| \gtrsim 3.$$

There is an additional advantage in using this projection. $\rho_{00} \frac{d\sigma}{dt}$ may be written in terms of s-channel helicity amplitudes as

$$\rho_{00} \frac{d\sigma}{dt} = \left| M_{00\frac{1}{2}-\frac{1}{2}} + M_{00\frac{1}{2}\frac{1}{2}} \right|^2 .$$

There are both theoretical and experimental reasons to expect the suppression of the nucleon non-flip helicity amplitude $M_{00\frac{1}{2}\frac{1}{2}}$. On the theoretical side, only π exchange contributes to the projection for ρ production, and only B exchange for ω production, both coupling only to nucleon helicity flip. Experimentally, the turnover at small momentum

transfers for both ρ and ω production indicates suppression of the nucleon non-flip amplitude.^{13,14} If the turnover for ω production continues to hold in a forthcoming high statistics ZGS experiment,¹⁵ it would indicate that the projection selects only a single nucleon helicity flip amplitude. In this case, the coherence factor for participating helicity amplitudes would trivially reduce to unity.

Let us consider what information we can extract from ρ - ω interference experiments. Knowing $\left| \frac{A_{\omega_0}}{A_{\rho_0}} \right|$, we get from the 2π cross-section Eq.(5) both the absolute value $\left| \epsilon_{2\pi} \right|$ and $\psi_{2\pi}$, while from the 3π cross-section ratio (14), we get only the combination $\left| \epsilon_{3\pi} \right| \cos \psi_{3\pi}$ where the relevant quantities are defined by Eqs.(6), (7), (13) and (15).

Clearly, if the isospin violating amplitudes are small enough, we can set

$$\epsilon_{2\pi} \approx \epsilon \approx \epsilon_{3\pi} \quad (16a)$$

and extract both the mixing phase ψ_ϵ and the phase difference of the production amplitudes as

$$\psi_\epsilon = \frac{1}{2}(\psi_{2\pi} + \psi_{3\pi}) \quad (16b)$$

and

$$\psi_{\omega_0} - \psi_{\rho_0} = \frac{1}{2}(\psi_{2\pi} - \psi_{3\pi}), \quad (16c)$$

respectively.

We should like to emphasize, however, that no experiment can measure the mixing parameter ϵ itself; only $\epsilon_{2\pi}$ and $\epsilon_{3\pi}$ are measured directly. To test for a possible difference between $\epsilon_{2\pi}$ and $\epsilon_{3\pi}$, let us think for a moment about photoproduction and e^+e^- annihilation experiments; since here the initial states do not possess definite G-parity, ρ - ω interference can only be observed in the 2π spectrum. Again, only the quantity $\epsilon_{2\pi}$ can be measured, rather than ϵ itself. Using now the value for $\psi_{\epsilon_{2\pi}}$ obtained from

electromagnetic production,¹⁻³ together with the phase $\psi_{2\pi}$, Eq.(7) determined from pionic production of ρ -mesons,⁴ one can estimate the strong production phase difference $\psi_{\rho_0} - \psi_{\omega_0}$. With the present experimental values and uncertainties, we have $\psi_{\epsilon_{2\pi}} = 95^\circ \pm 15^\circ$ ⁸ and $\psi_{2\pi}$ between 0 and 45° ⁴ which puts $\cos(\psi_{\epsilon} + \psi_{\rho_0} - \psi_{\omega_0})$ between -0.8 and -1. This, together with ω production data for $|\epsilon_{3\pi}| \cos \psi_{3\pi}$, Eq.(14), can be used to extract $|\epsilon_{3\pi}|$ for comparison with $|\epsilon_{2\pi}|$. A discrepancy between these two values would be significant¹⁶ if it were larger than roughly $|\epsilon_{2\pi}/2|$.

We note that, even with infinite accuracy, experiments cannot decide how much of the discrepancy between $\epsilon_{2\pi}$ and $\epsilon_{3\pi}$ is due to the production and how much due to the decay. Therefore, such an effect, if observed, could not be confronted with theory unless all isospin breaking terms can be calculated with sufficient accuracy.

In summary, combining data from measurements of ρ - ω interference in pionic production of 2π and 3π final states with previous electromagnetic production data yields a measurement or upper bound on the isospin violating effects in production and decay of the pure isospin state ω_0 . If these are small enough, one can use the strong interaction experiments alone to separate experimentally the mixing phase from the difference of production phases. This would provide an independent check on the mixing phase, and a test of exchange degeneracy of the π and B mesons.¹⁷

We would like to thank Robert Oakes, William F. Palmer, Veronika Rabl and Ronald Torgerson for helpful discussions.

REFERENCES

1. See Ref.8 for a list of electro- and photoproduction experiments.
2. D. Rhines, et al., ANL HEP Report 7146 (1971).
3. B. N. Ratcliff, et al., Phys. Lett. 38B, 345 (1972).
4. R. Diebold, et al., Proceedings of the Tallahassee Conference on $\pi\pi$ Interactions, to be published. The data are preliminary. When the final analysis is complete, the error for $\psi_{2\pi}$ should be reduced considerably.
5. A. S. Goldhaber, et al., Phys. Lett. 30B, 249 (1969).
6. M. Gourdin, et al., Phys. Lett. 30B, 247 (1969).
7. D. Horn, Phys. Rev. D 1, 1421 (1970).
8. J. L. Lemke and R. G. Sachs, Phys. Rev. D 5, 590 (1972).
9. N. N. Achasov and G. N. Shestakov, Nuc. Phys. B45, 93 (1972).
10. We use a spin zero instead of a spin 1 propagator. This simplification does not affect our results.
11. Such interactions have been proposed at various times; see e.g., R. J. Oakes, Phys. Lett. 29B, 683 (1969) and 30B, 262 (1969); R. Sokolow, Phys. Rev. 137, B1221 (1965); G. Cigogna, F. Strocchi and R. Vergara-Caffarelli, Phys. Rev. Lett. 29, 1702 (1972).
12. Present experimental evidence (see Ref.13) sets only a rather weak upper limit $\left| f_{\rho_0 3\pi} / f_{\omega_0 3\pi} \right| \lesssim \frac{1}{3}$. However, in the 3π amplitude the $f_{\rho_0 2\pi}$ term is multiplied by the broad ρ propagator, and it is suppressed by an order of magnitude if the integral in Eq.(8) is restricted to a narrow band around the ω mass. (A forthcoming ZGS experiment, see Ref.15, will test for $\Gamma_{\rho 3\pi}$ at the level of 0.1 MeV or below.)
13. J. A. J. Matthews, et al., Phys. Rev. Lett. 26, 400 (1971).

14. F. Bulos, et al., Phys. Rev. Lett. 26, 1453 (1971).
15. The evidence for turnover of $\rho_{00} \frac{d\sigma}{dt}$ in ω -production is weak; an experiment is being constructed to provide more information. See N. R. Stanton et al., ZGS Experimental Proposal E-337.
16. If, for example, the entire violation were in the decay process, such a difference would correspond to a decay width $\Gamma_{\omega_0 2\pi} \approx 0.08$ MeV. (This estimate is made with the assumption that $(f_{\omega_0 2\pi}/f_{\rho_0 2\pi})$ and ϵ are relatively real.)
17. π -B degeneracy would imply the unusually flat Regge trajectory slope of 0.66 GeV^{-2} .