NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

RECEIVED BY TIC MAY 12 1972

AEC-COO-1515-41
April 26, 1972

TECHNICAL PROGRESS REPORT

Contract AT(11-1)-1515 between U.S. ATOMIC ENERGY COMMISSION and THE UNIVERSITY OF WISCONSIN, Principal Investigator, Marion L. Jackson and Co-Principal Investigator, J. Keith Syers, Department of Soil Science, University of Wisconsin, Madison, Wisconsin 53706
Phone: 608+262-0562 or 608+262-2633.

This progress report is submitted pursuant to Article B-XXI of Contract No. AT(11-1)-1515, which has been in operation since June 1, 1965. Under this contract, 20 scientific papers have been published or accepted for publication, five additional papers have been submitted for publication, and seven others are in various stages of development for publication (all listed near the end of this report).

Synopsis

Scientific knowledge has been advanced in the following subject matter areas:

(i). Blister-like areas were induced according to degree of hydration of cations through osmotic swelling in low charge areas of micaceous vermiculite basal cleavages. This phenomenon occurred after washing out the excess salts of the saturating cations in the lyotropic series:

No other cation induced this type of swelling.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

(ii). Specific sorption of $^{90}\text{Sr}^{2+}$ and $^{45}\text{Ca}^{2+}$ in the presence of 1N NaNO_3 was traced to proton replacement in solid surface functional groups of hydrous metal oxides of Fe and Al in gel form. The sorption was greatly influenced by the pH of the solutions and sorption influenced the charge of the gel. Amorphous coatings of such materials were previously found on the surfaces of micaceous vermiculites.

(iii). Perforations such as holes and cracks in micaceous vermiculites caused by fission of ^{238}U in the crystal layers were found by scanning electron microscopy. Coatings and crusts of Fe and Al hydrous oxides may have been formed by diffusion through these crystal defects. Ultramicrotomy and high resolution electron microscopy permitted direct photography of layers and interlayers of phyllosilicates. The defects are expected to affect the weathering release of K^+ and fixation of exchangeable cation species such as $^{137}\text{Cs}^+$.

(iv). The aerosolic (tropospheric) dusts, found in this project by the use of the oxygen isotope composition of the mineral quartz to have accreted in soils and sediments in a wide range of locations throughout the world, have been shown to contain micaceous vermiculite which is responsible for the sorption and strong fixation of $^{137}\text{Cs}^+$. The dusts, which vary in their $^{137}\text{Cs}^+$ content depending on locality and

time of deposition, all sorbed significant amounts of added $^{137}\text{Cs}^+$ and strongly fixed it against replacement by Ca^{2+} in 0.01N CaCl_2 solution used to simulate natural soil conditions. Eolian influenced soils developed in basalt in Australia were much less effective in fixing $^{137}\text{Cs}^+$ and $^{90}\text{Sr}^{2+}$ against this extractant.

The research conducted during the past year under this contract is outlined in greater detail, together with documentation to project papers, in the following pages.

Documentation

I. Osmotic swelling on micaceous vermiculite basal cleavages as revealed by electron microscopy: Osmotic swelling was found to blister-like swelling on phlogopitic, saponitic, vermiculite on Li^+ saturation and washing out the excess LiCl (Sridhar, Jackson and Syers, 1972 -- COO-1515-37)*. Blisters had earlier been noted in the project (Roth, Jackson, and Syers, 1969 -- COO-1515-23; Sridhar, Jackson and Syers, 1970). During the current year, research on ^{effects of} layer charge and hydration character of the cation has been carried out and prepared for publication. This blister-like

* (Authors, date) refers to publications or manuscripts accepted for publication given in the first and second list, respectively, of references appended. (Authors, in preparation) refers to manuscripts in the third list of references appended.

configuration creates wedge zones at the edges, which are expected to be important in cation exchange selectivity as reported earlier (Dolcater, Lotse, Syers and Jackson, 1968 -- COO-1515-15; Jackson, 1968 -- COO-1515-14). Project studies (Dolcater, Jackson, and Syers, 1972 -- COO-1515-24) revealed that layer charge density in addition to cation charge and hydration control cation exchange selectivity (CES) properties of micaceous vermiculite. Completion of the study (Sridhar, Jackson and Syers, 1971) on reversal of the swelling by the fixing cations K^+ , NH_4^+ , Rb^+ , and Cs^+ is in process (Sridhar and Jackson, in preparation). Literature on weathering of micas has been extensively reviewed (Mokma, Jackson, and Syers, 1972 -- COO-1515-31).

Although layer charge blocked by (Fe,Al) coatings was released by dithionite (CBD) treatment, additional structural iron was reduced and oxidized without being reflected in a corresponding layer charge change (Roth, Jackson, and Syers, 1969 -- COO-1515-23). This anomaly is being investigated (Veith and Jackson, in preparation; Sridhar and Jackson, in preparation).

II. Specific adsorption of cations such as strontium-90 by hydrous metal oxides: Sorption of $^{90}Sr^{2+}$ and $^{45}Ca^{2+}$ was carried out (Kinniburgh, Syers, and Jackson, 1971) in 10^{-6} molar concentrations in 1M $NaNO_3$ solution to measure specific sorption sites on hydrous gels of iron and aluminum.

The divalent cations were sorbed even in the presence of a million times more concentrated Na^+ solution. One mole of M^{2+} was specifically sorbed for each mole of H^+ released, so that the surface charge became more positive as a result of the sorption. The mass action law was obeyed and the divalent cations were completely isotopically exchangeable (Kinniburgh, Syers, and Jackson, in preparation).

Coatings of (Fe,Al) on micaceous vermiculite were shown (Roth, Jackson, de Villiers, and Volk, 1967 -- C00-1515-3) in the project to have pH dependent cation exchange charge characteristics (de Villiers and Jackson, 1967 -- C00-1515-5). The coatings blocked some of the layer charge (Roth, Jackson, Lotse, and Syers, 1968 -- C00-1515-13) which was released by CBD treatment to remove the hydrous oxides. Mainly ferruginous sesquioxide coatings on micaceous vermiculite surfaces appeared in the early stages of soil development (less than 250 years) and increased in thickness and amount in the older soils of a chronosequence of soils developed in greywacke and mica-schist materials of New Zealand (Mokma, Jackson, Syers, and Stevens, 1972' -- C00-1515-30). Redistribution of iron oxides was indicated by the increase in the quantity of discrete goethite particles in the older soils. The amount of $^{90}\text{Sr}^{2+}$ sorbed varied within a relatively narrow range but that fixed against replacement by Ca^{2+} in 0.01N CaCl_2 varied over a

much wider range. The tendency for a decrease in the amount of $^{90}\text{Sr}^{2+}$ sorbed and fixed with increasing age in the older soils indicates that the free iron oxides which sorb and fix Sr^{2+} increase in crystallinity and become less active.

Removal of ferruginous sesquioxides tended to decrease the extent of sorption and fixation of $^{90}\text{Sr}^{2+}$ but increased the extent of sorption and fixation of $^{137}\text{Cs}^+$, owing to factors of increased layer charge density and also of accessibility to the crystal interiors (discussed below).

III. Evaluate by electron microscopy and radiochemical analysis the origin of micaceous vermiculite configurations, such as fission particle tracks and sesquioxide coatings, and their effect on cation exchange properties: The structural configurations of micaceous vermiculites were investigated further by scanning electron microscopy (Jackson, Lee, Brown, Sachs, and Syers, 1972 -- C00-1515-34). The coatings of (Fe,Al) oxides were shown to occur as a spatter-pattern over intermittent cleavages while other cleavages were clean. The oxide distribution may build up over ^{238}U fission particle tracks, the latter being found in both primary micas and in vermiculite. Diffusion of octahedral ions and K^+ through the tracks may occur during weathering, since K^+ diffusion occurred through tracks induced by UO_2^{2+} saturation of mica surfaces and neutron activation of the samples, a subject being investigated further (Lee, Jackson, and Syers, in preparation).

Earlier the cation exchange capacity of muscovite macroflakes, determined radiochemically (Mokma, Syers, and Jackson, 1970 -- C00-1515-17), was found to be between two and three times higher than the theoretical value, a finding related to cation exchange in areas made accessible to solutions by cleavages and weathering along the (hk0) direction in the crystals, possibly along fission particle tracks, and along growth planes.

Similar morphological features on the cleavages of micaceous vermiculite were obtained by replica electron microscopic observation conducted under this project (Roth et al., 1967); the SEM pictures confirm the replica work (REM) and are more informative as to morphology of the system. Numerous microscopic cracks and pores occurred normal to the (001) cleavage surfaces on micaceous vermiculite macroflakes. Treatment of micaceous vermiculites with citrate-bicarbonate-dithionite produced smooth surfaces and released minute fragments by fracture along crystal growth planes which were previously held together by sesquioxidic crusts. Splitting along (hk0) planes, approximately normal to (001) planes, indicated weakened bonds within the octahedral layer caused by the protonation of structural oxygens as octahedral cations moved out.

The presence of crystallographic flaws, such as cracks and holes, appear to increase the accessibility of the interior

parts of micaceous vermiculite to fixation of $^{137}\text{Cs}^+$ and other cations. For example, $^{137}\text{Cs}^+$ was strongly fixed in micaceous New Zealand soils (Mokma, Jackson, Syers, and Stevens, 1972 -- C00-1515-30), increasing with removal of the free ferruginous oxides. The micaceous vermiculite content of a soil horizon largely determined its sorption and fixation of $^{137}\text{Cs}^+$. Removal of free iron oxides increased the amount of $^{137}\text{Cs}^+$ sorbed and fixed by organic-matter free samples, presumably because additional exchange charge density was made active by removal of the positively charged coatings. The strong fixation of $^{137}\text{Cs}^+$ by the micaceous vermiculite component of the soils suggests that the fixed $^{137}\text{Cs}^+$ may not be released short of the decomposition of the micaceous vermiculite by weathering.

The crystallographic layers and oxide interlayers of chlorite were directly observed by high resolution electron microscopy after ultramicrotome cutting of sections perpendicular to the layers (Brown and Jackson, 1972 -- C00-1515-35). In a study of hydroxy interlayer-layer relations in phyllosilicates, micrographs made before and after an appreciable irradiation from the electron beam revealed images of the 7 Å structural plane spacings of chloritized materials, representing the interlayer hydroxyl to layer hydroxyl (002) electron density nodes. Loss of the images at 7Å intervals and reenforcement of those at 14Å intervals

was caused by heating in the electron beam. Work is in progress (Lee, Jackson and Brown, in preparation) showing micaceous vermiculite in type specimens and those after a high degree of natural weathering. Establishment by these studies of the presence of micaceous vermiculite in clays of highly weathered kaolinitic soils (under study) should help explain properties important to fixations of radioactive wastes such as $^{137}\text{Cs}^+$.

IV. Aerosolic (tropospheric) dusts containing micaceous vermiculite react with radiocesium and radiostrontium:

Through the use of the oxygen isotope ratios of the 'marker mineral' quartz isolated by H_2SiF_6 solution (Syers, Chapman, Jackson, Rex, and Clayton, 1968 -- COO-1515-16), the addition has been shown of aerosolic dusts to soils and sediments in a wide range of locations throughout the world (Rex, Syers, Jackson, and Clayton, 1969 -- COO-1515-21; Syers, Jackson, Berkheiser, Clayton, and Rex, 1969 -- COO-1515-22). In the North Pacific area, the $\delta^{18}\text{O}$ values of quartz isolated from a wide range of soils developed over quartz-free basalt in several Hawaiian Islands (Jackson, Levelt, Syers, Rex, Clayton, Sherman, and Uehara, 1972 -- COO-1515-27) and from deep-sea pelagic sediments (Clayton, Rex, Syers, and Jackson, 1972 -- COO-1515-29) were remarkably similar (16 to 18 o/oo), suggesting a common eolian origin. The average $\delta^{18}\text{O}$ values of quartz from Hawaiian soils (17.6 o/oo) and pelagic

sediments (17.8 o/oo) were virtually identical and varied similarly for any given particle size range, indicating a common eolian origin for the quartz and associated micaceous minerals.

In contrast, the $\delta^{18}\text{O}$ values of quartz isolated from soils and sediments collected from several areas in the South Pacific area (Mokma, Syers, Jackson, Clayton, and Rex, 1972 -- C00-1515-32) were appreciably lower (12 to 15 o/oo) than the corresponding samples from the North Pacific area. Most particles in the pelagic sediments from the South Pacific fell within the aerosolic dust range. The $\delta^{18}\text{O}$ values of quartz isolated from a range of pelagic sediments between latitudes 35 and 45°S were similar to those (12.9 to 15.4 o/oo) of quartz from a wide range of New Zealand and Australian soils located between these latitudes. The dust samples collected from the several eolian-influenced soils in the South Pacific area contain, along with quartz, an appreciable amount of micaceous vermiculite minerals. An eolian transport of quartz (and associated micaceous vermiculite) from land to sea appears to explain this relationship and also that of decreasing oxygen delta values of quartz from sediments and soils with increasing latitude in the South Pacific area.

Since the soils and sediments derived from aerosolic dust have micaceous vermiculite as a characteristic feature

of their mineralogy, the $^{137}\text{Cs}^+$ fixing capacity was measured for a number of dusts. The specific activity of $^{137}\text{Cs}^+$ on eight aerosolic dusts collected from snow and ice, from mud-rains, and from the atmosphere at several localities in the world ranged from 0 to 264 pc/g (Syers, Mokma, Jackson, Dolcater, and Rex, 1972 -- C00-1515-28). Zero values were obtained for dust samples collected from a mud-rain in England (deposited in 1968) and from an ice-core in Greenland (deposited pre-1940). A sample separated from melted ice collected on the Franz Josef glacier in New Zealand contained 264 pc/g of $^{137}\text{Cs}^+$; a relatively high proportion of the $^{137}\text{Cs}^+$ on the latter sample could not be replaced by five consecutive washings with 0.01N CaCl_2 , indicating a strong fixation by the micaceous vermiculite component of the dust. The fact that all of the dust samples sorbed significant amounts (30 to 50%) of added carrier-free $^{137}\text{Cs}^+$, suggests that the absence or presence of only small amounts of $^{137}\text{Cs}^+$ on dusts is related to the lack of nuclear testing at the time when the dust was transported. A pelagic sediment from the north central Pacific and a soil from the Hawaiian Islands, both of which contain weathered micaceous minerals of eolian origin, also sorbed appreciable amounts of added $^{137}\text{Cs}^+$. This implies that aerosolic dusts following deposition as sediments or soils have the ability to sorb $^{137}\text{Cs}^+$ incident in rainwater. Most of the added $^{137}\text{Cs}^+$

sorbed by the dusts was fixed against replacement by Ca^{2+} in 0.01N CaCl_2 .

Eolian influenced soils of Victoria, Australia fixed $^{137}\text{Cs}^+$ and $^{90}\text{Sr}^{2+}$ much less strongly, reflecting a predominance of basalt-derived clays (Mokma, Jackson, Syers, and Gibbons, 1972 -- COO-1515-33). The oxygen isotope abundance . (13.6 to 15.4 o/oo) in quartz from soils developed in this chronosequence of quartz-free basalt and from associated soils in sedimentary rocks and the particle size distribution of the quartz (flöttsand, coarser than aerosolic silt) indicate that the quartz originated by eolian transport from stranded beach ridges and dunes occurring windward (Jackson, Syers, Gibbons, and Mokma, 1972 -- COO-1515-43).

In summary, aerosolic dusts appear to constitute a major vehicle for the global transport and fixation of $^{137}\text{Cs}^+$. The widespread addition of aerosolic dusts to soils and sediments, established in this study, also insures that the $^{137}\text{Cs}^+$ in rainfall is sorbed and strongly fixed by the micaceous vermiculite component and thus more easily washed from surface tissues on which it might be incident.

PUBLICATIONS

and Manuscripts Accepted for Publication

Under AEC contract AT(11-1)-1515, initiated June 1, 1965

1967

1. de Villiers, J. M., and M. L. Jackson
1967. ALUMINOUS CHLORITE ORIGIN OF pH-DEPENDENT CATION EXCHANGE CAPACITY VARIATIONS.
Soil Sci. Soc. Amer. Proc. 31:614-619.
(Paper COO-1515-5)
2. Roth, C. B., M. L. Jackson, J. M. de Villiers, and V. V. Volk
1967. SURFACE COLLOIDS ON MICACEOUS VERMICULITE.
In Soil Chemistry and Soil Fertility, G. V. Jacks, Editor, Trans. Commissions II and IV, Intl. Soc. Soil Sci., (Aberdeen), pp. 217-221.
(Paper COO-1515-3)

1968

3. Dolcater, D. L., E. G. Lotse, J. K. Syers, and M. L. Jackson
1968. CATION EXCHANGE SELECTIVITY OF SOME CLAY-SIZED MINERALS AND SOIL MATERIALS.
Soil Sci. Soc. Amer. Proc. 32:795-798.
(Paper COO-1515-15)
4. Jackson, M. L.
1968. WEATHERING OF PRIMARY AND SECONDARY MINERALS IN SOILS.
Trans. Intl. Soc. Soil Sci., 9th Congr. (Adelaide) 4:281-292.
(Paper COO-1515-14)
5. Roth, C. B., M. L. Jackson, E. G. Lotse, and J. K. Syers
1968. FERROUS-FERRIC RATIO AND CEC CHANGES ON DEFERRATION OF WEATHERED MICACEOUS VERMICULITE.
Israel J. Chem. 6:261-273.
(Paper COO-1515-13)
6. Syers, J. K., S. L. Chapman, M. L. Jackson, R. W. Rex, and R. N. Clayton
1968. QUARTZ ISOLATION FROM ROCKS, SEDIMENTS AND SOILS FOR DETERMINATION OF OXYGEN ISOTOPIC COMPOSITION.
Geochim. Cosmochim. Acta 32:1022-1024.
(Paper COO-1515-16)

1969

7. Rex, R. W., J. K. Syers, M. L. Jackson, and R. N. Clayton
1969. EOLIAN ORIGIN OF QUARTZ IN SOILS OF HAWAIIAN ISLANDS AND IN PACIFIC PELAGIC SEDIMENTS.
Science 163:277-279.
(Paper COO-1515-21)
8. Roth, C. B., M. L. Jackson, and J. K. Syers
1969. DEFERRATION EFFECT ON STRUCTURAL FERROUS-FERRIC IRON RATIO AND CEC OF VERMICULITES AND SOILS.
Clays Clay Minerals 17:253-264.
(Paper COO-1515-23)
9. Syers, J. K., M. L. Jackson, V. E. Berkheiser, R. N. Clayton, and R. W. Rex
1969. EOLIAN SEDIMENT INFLUENCE ON PEDOGENESIS DURING THE QUATERNARY.
Soil Sci. 107:421-427.
(Paper COO-1515-22)

1970

10. Dolcater, D. L., J. K. Syers, and M. L. Jackson
1970. TITANIUM AS FREE OXIDE AND SUBSTITUTED FORMS IN KAOLINITES AND OTHER SOIL MINERALS.
Clays Clay Minerals 18:71-79.
(Paper COO-1515-25)
11. Mokma, D. L., J. K. Syers, and M. L. Jackson
1970. CATION EXCHANGE CAPACITY AND WEATHERING OF MUSCOVITE MACROFLAKES.
Soil Sci. Soc. Amer. Proc. 34:146-151.
(Paper COO-1515-17)
12. Sridhar, K., M. L. Jackson, and J. K. Syers
1970. MICACEOUS VERMICULITE BASAL CLEAVAGES WITH BLISTER-LIKE LIFTING OF LAYERS AS A FUNCTION OF EXCHANGEABLE CATION CHARGE AND HYDRATION.
Abstracts, Geol. Soc. Amer. meeting, Milwaukee, WI, pp. 692-693.

1971

13. Jackson, M. L., T. W. M. Levelt, J. K. Syers, R. W. Rex, R. N. Clayton, G. D. Sherman, and G. Uehara
1971. GEOMORPHOLOGICAL RELATIONSHIPS OF TROPICOSPHERICALLY-DERIVED QUARTZ IN SOILS OF THE HAWAIIAN ISLANDS.
Soil Sci. Soc. Amer. Proc. 35:515-525.
(Paper COO-1515-27)

14. Kinniburgh, D. G., J. K. Syers, and M. L. Jackson
1971. SELECTIVE SORPTION AND EXCHANGE OF Ca^{2+} AND Sr^{2+} BY OXIDES AND HYDROUS OXIDES OF IRON AND ALUMINUM.
Agronomy Abstracts, New York meeting, NY (1971), pp. 123-124.
15. Sridhar, K., M. L. Jackson, and J. K. Syers
1971. INTERACTION OF FIXING CATIONS WITH BLISTERS ON MICACEOUS VERMICULITE CLEAVAGES PRODUCED BY OSMOTIC SWELLING.
Abstracts 20th Clay Min. Conf., Rapid City, SD (1971), pp. 32.

1972

16. Clayton, R. N., R. W. Rex, J. K. Syers, and M. L. Jackson
1972. OXYGEN ISOTOPE ABUNDANCE IN QUARTZ FROM PACIFIC PELAGIC SEDIMENTS.
J. Geophys. Res.
(Accepted for publication) (COO-1515-29).
17. Syers, J. K., D. L. Mokma, M. L. Jackson, D. L. Dolcater, and R. W. Rex
1972. MINERALOGICAL COMPOSITION AND CESIUM-137 RETENTION PROPERTIES OF CONTINENTAL AEROSOLIC DUSTS.
Soil Sci. 113:116-123.
(COO-1515-28).
18. Mokma, D. L., J. K. Syers, M. L. Jackson, R. N. Clayton, and R. W. Rex
1972a. AEOLIAN ADDITIONS TO SOILS AND SEDIMENTS IN THE SOUTH PACIFIC AREA.
J. Soil Sci. 23:
(Accepted for publication) (COO-1515-32).
19. Mokma, D. L., M. L. Jackson, J. K. Syers, and P. R. Stevens
1972b. MINERALOGY OF A CHRONOSEQUENCE OF SOILS FROM GREYWACKE AND MICA SCHIST ALLUVIUM, WESTLAND, NEW ZEALAND.
N. Z. J. Sci.
(Accepted for publication) (COO-1515-30).
20. Mokma, D. L., M. L. Jackson, and J. K. Syers
1972c. WEATHERING OF MICAS IN SOILS.
Encyclopedia of Soil Science, Springer, Berlin-Gottingen-Heidelberg
(Accepted for publication) (Paper COO-1515-31)

MANUSCRIPTS

Submitted for Publication

1. Brown, J. L., and M. L. Jackson
1972. CHLORITE EXAMINATION BY ULTRAMICROTOMY AND
HIGH RESOLUTION ELECTRON MICROSCOPY.
Clays Clay Miner. 20:
(Submitted for publication) (COO-1515-35).
2. Dolcater, D. L., M. L. Jackson, and J. K. Syers
1972. CATION EXCHANGE SELECTIVITY IN MICA AND
VERMICULITE.
Amer. Mineral.
(Submitted for publication) (COO-1515-24).
3. Jackson, M. L., S. Y. Lee, J. L. Brown, I. B. Sachs,
and J. K. Syers
1972. SCANNING ELECTRON MICROSCOPY OF HYDROUS METAL
OXIDE CRUSTS INTERCALATED IN NATURALLY
WEATHERED MICACEOUS VERMICULITE.
Soil Sci. Soc. Amer. Proc. 36:
(Submitted for publication) (COO-1515-34).
4. Sridhar, K., M. L. Jackson, and J. K. Syers
1972. MICACEOUS VERMICULITE BASAL CLEAVAGES WITH
BLISTER-LIKE INTERLAYER SWELLING AS A FUNCTION
OF EXCHANGEABLE CATION AND LAYER CHARGE.
Amer. Mineral.
(Submitted for publication) (COO-1515-37).
5. Mokma, D. L., M. L. Jackson, J. K. Syers, and F. R.
Gibbons
1972. MINERALOGY AND RADIOISOTOPE RETENTION
PROPERTIES OF A CHRONOSEQUENCE OF SOILS
DEVELOPED IN BASALT OF VICTORIA, AUSTRALIA.
J. Soil Sci.
(Submitted for publication) (COO-1515-33).

Manuscripts in Preparation

1. Jackson, M. L., F. R. Gibbons, J. K. Syers, and D. L. Mokma
1972. EOLIAN INFLUENCE ON SOILS DEVELOPED IN A CHRONOSEQUENCE OF BASALTS OF VICTORIA, AUSTRALIA.
Soil Sci.
(COO-1515-43) (To be submitted for publication)
2. Kinniburgh, D. G., J. K. Syers, and M. L. Jackson
1972. SELECTIVE SORPTION AND EXCHANGE OF CALCIUM AND STRONTIUM BY OXIDES AND HYDROUS METAL OXIDES OF IRON AND ALUMINUM.
Soil Sci. Soc. Amer. Proc. 36:
(COO-1515-38) (To be submitted for publication)
3. Sridhar, K., and M. L. Jackson
1972. INTERACTION OF FIXING CATIONS WITH BLISTERS ON MICACEOUS VERMICULITE CLEAVAGES PRODUCED BY OSMOTIC SWELLING.
Clays Clay Miner. 20:
(COO-1515-42) (To be submitted for publication)
(Presented before the Clay Conference, Rapid City, SD, Aug. 1971).
4. Lee, S. Y., M. L. Jackson, and J. L. Brown
1972. MICACEOUS VERMICULITE DOMAINS IN KAOLINITE WEATHERING PRODUCT OBSERVED BY ULTRAMICROTOMY AND HIGH RESOLUTION ELECTRON MICROSCOPY.
Clays Clay Miner. 20:
(COO-1515-44) (To be proposed for presentation at the 21st Annual Clay Mineral Conference, Woods Hole, MA, Sept. 1972)
5. Sridhar, K., and M. L. Jackson
1972. LAYER CHARGE DECREASE MECHANISMS DURING THE TRANSFORMATION OF PHLOGOPITE THROUGH VERMICULITE TO SAPONITE.
Soil Sci. Soc. Amer. Proc. 36:
(COO-1515-46) (To be proposed for presentation at Miami Beach, FL, meeting Oct., 1972).
6. Lee, S. Y., M. L. Jackson, and J. K. Syers
1972. FISSION PARTICLE TRACKS IN MICAS AND MICACEOUS VERMICULITES AS RELATED TO CHEMICAL WEATHERING AND CATION EXCHANGE PROPERTIES.
Soil Sci. Soc. Amer. Proc. 36:
(COO-1515-45) (To be proposed for presentation at the Miami Beach, FL meeting, Oct., 1972).
7. Veith, J. A., and M. L. Jackson
1972. MECHANISM OF LAYER CHARGE LOWERING DURING WEATHERING OF MICACEOUS MINERALS.
(Paper being developed currently).

Key to C00-1515 Report Series

Contract AT(11-1)-1515 between
U.S. ATOMIC ENERGY COMMISSION and THE UNIVERSITY OF WISCONSIN

Principal Investigator: Marion L. Jackson
Co-Principal Investigator: J. Keith Syers

Department of Soil Science, University of Wisconsin, Madison, WI 53706
Phone: 608+262-0562 or 608+262-2633

C00- 1515	Sent		Author, Journal, Content	Manuscript date
	Manu.	Reprint		
*				
1	x	na	Science Information Exchange form	15 July 65
2	x	na	Science Information Exchange form	1 Mar 66
3	x	x	Roth <u>et al.</u> , ISSS, Aberdeen (1967) Ver sesquioxide coatings	1 Oct 66
(4)	x	na	Jackson, Trip Report, Aberdeen	1 Oct 66
5	x	x	de Villiers and Jackson, SSSAP (1967) Al/(OH) ₂ pH/CEC	9 Jan 67
6	x	na	Technical Progress Report (1967)	1 Feb 67
(7)	x	na	Renewal Proposal	14 Feb 67
(8)	x	na	Single Column Budget	14 Feb 67
(9)	x	na	Statement of costs	14 Feb 67
(10)	x	na	Equipment report	14 Feb 68
11	x	na	Science Information Exchange form	14 Feb 67
12	x	na	Science Information Exchange form	24 Feb 68
13	x	x	Roth <u>et al.</u> , <u>Israel J. Chem.</u> (1968) Fe/CEC	30 June 67
14	x	x	Jackson, <u>ISSS 9th Congr.</u> (Adelaide) (1968) pH/CEC	1 July 67
15	x	x	Dolcater <u>et al.</u> , <u>SSSA Proc.</u> (1968) CES/mica-verm layer charge	15 Dec 67
16	x	x	Syers <u>et al.</u> , <u>Geochim. Cos. Acta</u> (1968) Quartz Isolation by H ₂ SiF ₆	14 Feb 68
17	x	x	Mokma <u>et al.</u> , <u>SSSA Proc.</u> (1970) Muscovite flakes/CEC, coatings	27 Feb 70
18	x	na	Technical Progress Report (1968)	24 Feb 68
19	x	na	Science Information Exchange form	24 Feb 69
20	x	na	Technical Progress Report (1969)	24 Feb 69

* Numbers refer to published papers and Technical Progress Reports, including C00-1515-6 (1967), C00-1515-18 (1968), and C00-1515-20 (1969), with abstracts of papers published under the contract. Numbers in () incorrectly used for other than Technical Progress Reports. na = Not applicable.

Key to C00-1515 Report Series (Cont.)

Sent		Author, Journal, Content	Manuscript date
C00-1515	Re-Ms. print		
21	x	x Rex <u>et al.</u> , <u>Science</u> (1969) Aerosolic dust	23 Sept 68
22	x	x Syers <u>et al.</u> , <u>S. Sci.</u> (1969) Dusts/loess	20 Feb 69
23	x	x Roth <u>et al.</u> , <u>CCMin</u> (1969) Fe,Al coat./CEC	6 Jan 69
24	x	Dolcater <u>et al.</u> , <u>Am. Min.</u> (1972) CES	26 Feb 71
25	x	x Dolcater <u>et al.</u> , <u>CCMin</u> (1970) CEC/Ti Subst.	27 Feb 70
26	x	na Technical Progress Report (1969-70)	28 Feb 70
27	x	xx Jackson <u>et al.</u> , <u>SSSAProc</u> (1971) dusts/geomor.	26 Feb 71
28	x	xx Syers <u>et al.</u> , <u>S. Sci.</u> (1972) dusts/ ¹³⁷ Cs	26 Feb 71
29	xx	Clayton <u>et al.</u> , <u>JGeophysRe</u> (1972) dusts/pelag	26 Apr 72
30	x	Mokma <u>et al.</u> , <u>NZJSci.</u> (1972) Chrono./ ¹³⁷ Cs	26 Feb 71
31	x	Mokma <u>et al.</u> , <u>Hbook S. Sci.</u> (1972) Mica Wea.	26 Feb 71
32	x	xx Mokma <u>et al.</u> , <u>J. S. Sci.</u> (1972) Eol./S.Pac.	26 Feb 71
33	x	Mokma <u>et al.</u> , <u>J. S. Sci.</u> (1972) ASL/Isotopes	26 Feb 71
34	xx	Jackson <u>et al.</u> , <u>SSSAProc.</u> SEM/Verm	26 Apr 72
35	xx	Brown, Jackson, <u>CCMin</u> Ultmicrot./layers	26 Apr 72
36	x	na Technical Progress Report (1971)	26 Feb 71
37	xx	Sridhar <u>et al.</u> , <u>Am. Min.</u> Blisters/cations	26 Apr 72
38	xx	Kinniburgh <u>et al.</u> , <u>SSSAP</u> Sr, Ca sorp/Sesq.	26 Apr 72
39	x	na Science Information Exchange Form	6 Feb 71
40	xx	na Science Information Exchange Form	6 Apr 72
41	xx	na Technical Progress Report (1972)	26 Apr 72
42		Sridhar <u>et al.</u> , <u>CCM</u> Blisters/fixation In preparation	"
43		Jackson <u>et al.</u> , <u>Soil Sci.</u> (1972) ASL/Isotopes	"
44		Lee <u>et al.</u> , <u>CCM</u> Mica domains	"
45		Lee <u>et al.</u> , <u>SSSAP</u> Fission trac	"

xx = Sent with this Technical Progress Report (C00-1515-41).

Effort levels:

Principal Investigator, Marion L. Jackson,

0.3 man year equivalent

(50% in two summer months; 20% during academic
year to present, and to end of current term
of contract)

Co-Principal Investigator, J. Keith Syers,

0.2 man-year equivalent

(to present and to end of current term of contract)