MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

KN Kinematics

P(LAB)---T(LAB)---INV MASS---P(CMS)---
(MEV/C) (MEV) (MEV) (MEV/C)

0	0	1432	0
20	0	1432	13
40	2	1433	26
60	4	1434	39
80	6	1436	52
100	10	1439	65
120	14	1441	78
140	19	1445	91
160	25	1449	104
180	32	1453	116
200	39	1457	129
220	47	1462	141
240	55	1468	153
260	64	1474	166
280	74	1480	178
300	84	1486	189
320	95	1493	201
340	106	1500	213
360	117	1507	224
380	129	1514	235
400	142	1522	247
420	154	1530	258
440	168	1538	268
460	181	1546	279
480	195	1554	290
500	209	1563	300
520	223	1572	310
540	238	1580	321
560	253	1589	331
580	268	1598	341
600	283	1607	350
620	299	1616	360
640	315	1625	370
660	330	1634	379
680	347	1644	388
700	363	1653	397
720	379	1662	406
740	396	1671	415
760	413	1681	424
780	429	1690	433
800	446	1699	442
820	463	1709	450
840	481	1718	459
860	498	1728	467
880	515	1737	475
900	533	1747	483
920	550	1756	492
940	568	1765	500
960	586	1775	508
980	604	1784	515
1000	621	1794	523
1020	639	1803	531
1040	657	1812	538
1060	676	1822	546
1080	694	1831	553
1100	712	1840	561
1120	730	1850	568
1140	749	1859	575
1160	767	1868	583
1180	785	1877	590
1200	804	1887	597
1220	822	1896	604
1240	841	1905	611
1260	860	1914	618
1280	878	1923	624
1300	897	1932	631
1320	916	1941	638
1340	934	1950	645
1360	953	1959	651

P(LAB)---T(LAB)---INV MASS---P(CMS)---
(MEV/C) (MEV) (MEV) (MEV/C)

1380	972	1968	658
1400	991	1977	664
1420	1010	1986	671
1440	1029	1995	677
1460	1047	2004	684
1480	1066	2013	690
1500	1085	2022	696
1520	1104	2031	702
1540	1123	2039	709
1560	1143	2048	715
1580	1162	2057	721
1600	1181	2065	727
1620	1200	2074	733
1640	1219	2083	739
1660	1238	2091	745
1680	1257	2100	751
1700	1276	2109	756
1720	1296	2117	762
1740	1315	2126	768
1760	1334	2134	774
1780	1353	2143	779
1800	1373	2151	785
1820	1392	2159	791
1840	1411	2168	796
1860	1431	2176	802
1880	1450	2184	808
1900	1469	2193	813
1920	1489	2201	818
1940	1508	2209	824
1960	1527	2217	829
1980	1547	2226	835
2000	1566	2234	840
2020	1586	2242	845
2040	1605	2250	851
2060	1625	2258	856
2080	1644	2266	861
2100	1663	2274	866
2120	1683	2282	872
2140	1702	2290	877
2160	1722	2298	882
2180	1741	2306	887
2200	1761	2314	892
2220	1780	2322	897
2240	1800	2330	902
2260	1820	2338	907
2280	1839	2346	912
2300	1859	2353	917
2320	1878	2361	922
2340	1898	2369	927
2360	1917	2377	932
2380	1937	2384	937
2400	1956	2392	941
2420	1976	2400	946
2440	1996	2407	951
2460	2015	2415	956
2480	2035	2423	960
2500	2055	2430	965
2520	2074	2438	970
2540	2094	2445	975
2560	2113	2453	979
2580	2133	2460	984
2600	2153	2468	988
2620	2172	2475	993
2640	2192	2483	998
2660	2212	2490	1002
2680	2231	2498	1007
2700	2251	2505	1011
2720	2271	2512	1016
2740	2290	2520	1020
2760	2310	2527	1025
2780	2330	2534	1029

P(LAB)---T(LAB)---INV MASS---P(CMS)---
(GEV/C) (GEV) (GEV) (GEV/C)

3.0	2.5	2.61	1.08
3.2	2.7	2.63	1.12
3.4	2.9	2.75	1.16
3.6	3.1	2.82	1.20
3.8	3.3	2.88	1.24
4.0	3.5	2.95	1.27
4.2	3.7	3.01	1.31
4.4	3.9	3.07	1.34
4.6	4.1	3.13	1.38
4.8	4.3	3.19	1.41
5.0	4.5	3.25	1.44
5.2	4.7	3.31	1.48
5.4	4.9	3.36	1.51
5.6	5.1	3.42	1.54
5.8	5.3	3.47	1.57
6.0	5.5	3.52	1.60
6.2	5.7	3.58	1.63
6.4	5.9	3.63	1.65
6.6	6.1	3.68	1.68
6.8	6.3	3.73	1.71
7.0	6.5	3.78	1.74
7.2	6.7	3.83	1.76
7.4	6.9	3.88	1.79
7.6	7.1	3.93	1.82
7.8	7.3	3.97	1.84
8.0	7.5	4.02	1.87
8.2	7.7	4.07	1.89
8.4	7.9	4.11	1.92
8.6	8.1	4.16	1.94
8.8	8.3	4.20	1.96
9.0	8.5	4.25	1.99
9.2	8.7	4.29	2.01
9.4	8.9	4.33	2.03
9.6	9.1	4.38	2.06
9.8	9.3	4.42	2.08
10.0	9.5	4.46	2.10
10.5	10.0	4.57	2.16
11.0	10.5	4.67	2.21
11.5	11.0	4.77	2.26
12.0	11.5	4.86	2.31
12.5	12.0	4.96	2.36
13.0	12.5	5.05	2.41
13.5	13.0	5.15	2.46
14.0	13.5	5.24	2.51
14.5	14.0	5.32	2.56
15.0	14.5	5.41	2.60
16.0	15.5	5.58	2.69
17.0	16.5	5.75	2.77
18.0	17.5	5.91	2.86
19.0	18.5	6.07	2.94
20.0	19.5	6.22	3.02
22.0	21.5	6.51	3.17
24.0	23.5	6.79	3.31
26.0	25.5	7.07	3.45
28.0	27.5	7.33	3.59
30.0	29.5	7.58	3.71
32.0	31.5	7.82	3.84
34.0	33.5	8.06	3.96
36.0	35.5	8.29	4.08
38.0	37.5	8.51	4.19
40.0	39.5	8.73	4.30
42.0	41.5	8.94	4.41
44.0	43.5	9.15	4.51
46.0	45.5	9.35	4.62
48.0	47.5	9.55	4.72
50.0	49.5	9.74	4.81
55.0	54.5	10.21	5.05
60.0	59.5	10.66	5.28
65.0	64.5	11.10	5.50
70.0	69.5	11.51	5.71

PAGES i to ii
WERE INTENTIONALLY
LEFT BLANK

PREFACE TO THE SERIES

This is the first in a new series of reports produced by the Particle Data Group. In this series we will collect and display $\sqrt{\text{total}}$ and differential cross sections, polarizations, mass spectra, and other similar data. Each report will cover one input channel. This one is K^+N (K^+p and K^+d). In the next few months we hope to bring out NN , π^+N , and $\bar{N}N$. Following later will be π^-N , K^-N , etc. All reports will be complete from January 1968, and will also contain selected results before that date. The reports will be updated periodically, as necessary.

At present there are seventeen physicists in the Particle Data Group who are working on one or more phases of these reports. They are:

System Development (LRL)

Art Rosenfeld
LeRoy Price
Odette Benary
Naomi Schmidt[†]

Encoding Data (LRL)

Odette Benary
LeRoy Price
Naomi Schmidt[†]

 K^+N Interactions (LRL)

Odette Benary
Roger Bland[†]
LeRoy Price
Naomi Schmidt[†]
* Charles Wohl[§]
V. Henri (CERN) starting Sept. '69

 K^-N Interactions (CERN)

* Claude Bricman (below 2.5 GeV/c)
Enzo Flaminio (above 2.5 GeV/c)

 π^+N Interactions (MIT)

* Henry Lubatti^{||}
Terry Watts[#]
Fred Winkelmann

 π^-N Interactions

* Tom Ferbel (BNL)
David Miller (MIT)

 NN Interactions

Gideon Alexander (Tel-Aviv)
Odette Benary (LRL)

 $\bar{N}N$ Interactions

Yoshio Sumi (Osaka)
* Alan Thorndike (BNL)
Frank Turkot (BNL)
Toshihiro Yoshida (Kyoto)

If you have any suggestions for improving these reports, please let us know. Our address is:

Particle Data Center
Lawrence Radiation Laboratory
Berkeley, California 94720
(415) 843-2740, Ext. 6301;
nights, weekends, and holidays
call 642-0466

* "Chairman."

[†] Now at Brandeis Univ., Waltham, Massachusetts.

[‡] Now at Ecole Polytechnique, Paris.

[§] Now at Oxford University, Oxford

^{||} Now at University of Washington, Seattle, Washington.

[#] Now at SLAC, Stanford, California.

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

TABLE OF CONTENTS

Section I. General Procedures

A. Introduction	5
B. Scope of the Compilation	5
C. Data Handling	5
D. Collaboration with Other Groups	8
E. Other Cross-Section Compilations	8
F. Acknowledgments	8

Section II. K⁺N Data

A. Text

1. Renormalization of Bubble Chamber Cross Sections	11
2. Correcting for Unmeasured Decay Modes	11
3. K ⁺ p Interactions	11
Cross Sections	11
Elastic Angular Distributions	12
Polarization in Elastic Scattering	12
Elastic Phase Shift Analyses	13
Quasi-Two-Body Angular Distributions	13
4. K ⁺ d and K ⁺ n Interactions	14
Special Problems with Deuterium	14
K ⁺ d Cross Sections	14
K ⁺ n Cross Sections	14
K ⁺ n Charge-Exchange Angular Distributions.	15

B. Figures and Tables	16
1. Data	
$K^+p \rightarrow$ total	19-25
elastic	19-23
$KN\pi$	19, 22-23, 26-27
$KN\pi\pi$	19, 22-23, 28-29
$KN\pi\pi\pi$	31
K^*N	32-34
$K\Delta$	32-34
$K^*\Delta$	32-34
"rare reactions"	37
$K^+d \rightarrow$ total	88-89
KNN	88-89
$Kd\pi$	91
$KNN\pi$	90
$K^+n \rightarrow$ total	94-95
$KN\pi$	96
$KN(I=0) \rightarrow$ total	19, 98-99
elastic	19, 98-99
$KN\pi$	19, 98-99
2. Fits to the Data	
Forward K^+p Elastic Scattering	55
Backward K^+p Elastic Scattering	65
Section III. <u>Data Listings</u>	107
Section IV. <u>Indices</u>	227
Reference List	234

A COMPILATION OF K^+N REACTIONS

Particle Data Group

LeRoy R. Price, Naomi Barash-Schmidt, * Odette Benary,
Roger W. Bland, † Arthur H. Rosenfeld, Charles G. Wohl‡

Lawrence Radiation Laboratory
University of California
Berkeley, California 94720

ABSTRACT—We compile 90 papers reporting K^+N and K^+d interactions. We display cross sections, angular distributions, and polarizations. Included are indices to the papers, as well as a complete listing of the selected K^+N data, and results of our fits to some of the data. The cutoff date for this report was 15 July 1969.

* Present address: Brandeis University, Waltham, Massachusetts.

† Present address: Ecole Polytechnique, Paris.

‡ Present address: Oxford University, Oxford

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

Section I.

GENERAL PROCEDURES

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

Introduction

It has been known for some time that a comprehensive, evaluated compilation of experimental high energy physics results is essential. In the past there have been numerous collections of data, generally covering rather narrow fields. But because they were not computerized, the authors became "exhausted" on the first edition, and updated versions were never published.

Encouraged by the success of the Particle Data Group's computerized "Particle Properties Tables,"* we have started also to compile cross-sectional data. Our system is completely computer based, so that we will be able to answer specific user requests, in addition to periodically publishing the collected data.

During the last two and one-half years, we have had two full-time physicists here at LRL developing and coding the system programs.

During the last few months we have been feeding data into the system—and this report on K^+N is our first result. (We consider this report and the next, on NN , to be more or less "debug" versions—we are still trying to figure out the best ways to organize the reports and present the data.)

We plan to continue our program development and would appreciate any comments you may have on the way the data are displayed, the types of data collected, etc. In the next few months we hope to go to a form of photo-composition. This means essentially that we will have an unlimited character set and can print Greek letters instead of having to spell them out, for example.

*Particle Data Group, Rev. Mod. Phys. **41**, 109 (1969).

Scope of the Compilation

1. We will collect all experimental high energy physics results that can be represented by simple tables or graphs, i. e., σ , $d\sigma/d\omega$, polarizations, angular distributions, density matrices, etc.

We leave it to Data Summary Tape Libraries to store Dalitz plots or other ≥ 2 -dimensional displays (although the presence of such data is indicated in our KEY WORDS). In any case our printed compilations should serve as a necessary "table of contents" to a DST library.

2. The data come primarily from published journals, e. g., Physical Review, Physical Review Letters, Nuclear Physics, Physics Letters, Nuovo Cimento, etc.

We do also compile unpublished theses and conference reports—if the reports give enough information to permit a valid evaluation of the experiment and analysis.

We do not record data that appear in abstract form only, nor do we generally accept preprints unless the article has already been accepted for publication.

3. The compilation is to be complete from January 1968. Before that time we will enter data that are particularly important. But the bulk of the pre-1968 papers will not be put into our system.

4. To reduce the number of errors to the very minimum, all punched information is checked by the physicist who read the article and by another physicist as well.

Data Handling

In order to make this compilation as accurate and complete as possible, many physicists are involved. These physicists fall into two general categories:

a) Those who read and evaluate the data. These physicists (referred to as "readers") are generally experimentalists chosen for their "expertise" in a particular field. In

general they are not from LRL. They are organized into small groups, each group being responsible for a different input channel.

b) Those physicists who encode the data, run the programs, write system programs, etc. These are all at LRL.

The list below indicates the most important steps that every article must go through in order to have its information entered onto the DATA TAPE (the magnetic tape that contains all of our data). This list is summarized in the "Flow Diagram" in Fig. A.

a) The "reader" (physicist) finds a relevant article, reads it, marks it, fills out a form, and mails a copy of the article plus the form to us.

b) Our secretary assigns it a number and it is logged in.

c) The LRL physicist responsible for this initial state quickly scans it, writes out the standard KEY WORDS, etc., to help in the next step.

d) Our secretary transcribes bibliographic information, putting abstract, citations, comments, beam information, and KEY WORDS, onto coding sheets.

e) The LRL physicist transcribes the data that the reader has selected onto coding sheets (this is much more laborious than you might suspect).

f) Key-punch operators punch the data.

g) The LRL physicist puts the cards for a particular article into the correct order.

h) Cards are put onto the DATA TAPE with the DATAPE program.

i) If any cards are out of order, essential information missing, etc., the article is rejected by DATAPE. The physicist repairs the deck, and it is again put through DATAPE.

j) The output DATA TAPE is read by the SKELM program, which makes a listing of all the data stored for each article.

k) The SKELM is looked over by the LRL physicist for obvious errors.

l) SKELM output is mailed to the original physicist "reader," who checks all entries carefully and returns SKELM plus corrections (if any) to us.

m) If any errors are found, steps

e) through l) are repeated as many times as necessary.

n) When the reader has no more corrections or changes to make, the LRL physicist gives the article its final verification (i. e., he rechecks all data with the original article). The name of this physicist is put on the tape, and the article is then ready to be used by any one of a number of programs.

Steps b) through n) take, on the average, about 1-3/4 hours per article (1-1/4 hour physicist + 1/2 hour secretary).

Even after being verified, an article can have its contents slightly increased, e. g., if renormalized data are added. In this case only steps e) through k) are repeated.

Again, all the above is just to get the data onto the DATA TAPE. When preparing a report such as this, many additional tasks are involved. A few typical ones are:

a) Collecting all the data on a particular set of reactions—plotting them, looking at systematic errors, removing obviously bad data from the graphs (but leaving it in the tables).

b) Ironing out normalization differences between experiments.

c) Worrying about the various ways in which different authors make resonance cuts and subtractions.

d) Deciding what types of curves (if any) should be fit to certain classes of data.

Fig. A.

Collaboration with Other Groups

Some physicists in Europe have formed a group called HERA (High Energy Reactions Analysis)* to also compile cross-section data. We are trying to keep in close contact with one another in order to minimize duplication of effort both in programming and data collection. Their first report[†] was published earlier this year and more reports are expected soon.

We also cooperate with HERA on report distribution: LRL prints and distributes both HERA and our reports for the Western Hemisphere and Japan, and CERN does the same for the rest of the world.

We originally planned to collaborate closely with John Hornbostel of BNL. He had been compiling cross-section information for a number of years and was going to bring out a series of reports covering the data appearing before January 1968. Sadly, however, he died early this year, with the project incomplete. We wish to thank BNL for sending all of his files to us.

We also thank the Michigan Cross Section Group for sending us all of their files, also. Their report is referred to in the next section under Williams et al.

Other Cross-Section Compilations

We present below (in chronological order) all of the previous cross-section compilations that we know of. In addition to just listing data, some of them have nice reviews, perform various fits to the data, etc.

- V. S. Barashenkov and V. M. Maltsev, Cross Sections for Elementary Particle Interactions, Fortsch. Physik 9, 549 (1961).

*See B. Sadoulet, "An Example of an Organization of Compilation of Data," Preprint CERN/D. Ph. II/PHYSICS 68-21.

[†]B. Sadoulet, "Data Compilation of Antiproton-Proton Reactions into Antihyperon-Hyperon," CERN-HERA 69-2 (1969).

- V. S. Barashenkov and J. Patera, Cross Sections for Antinucleon Production, Fortsch. Physik 11, 469 (1963).
- V. S. Barashenkov and J. Patera, Strange Particle Production, Fortsch. Physik 11, 479 (1963).
- M. N. Focacci and G. Giacomelli, Pion-Proton Elastic Scattering, CERN 66-18 (1966).
- J. T. Beale, S. D. Ecklund, and R. L. Walker, Pion Photoproduction Data Below 1.5 GeV, CALT-68-108 (1966).
- H. Yukawa, ed., Experimental Data on Hadron Interactions in GeV Region, Supplement of the Progress of Theoretical Physics (Kyoto), Extra Number (1967).
- P. K. Williams, D. M. Levine, J. A. Koschik, References and Some Two-Body Data for High Energy Reactions, University of Michigan, 1967 (unpublished).
- G. Alexander, O. Benary, and U. Maor, Data Compilation of Proton-Proton Interactions Between 1 and 32 GeV/c, Nucl. Phys. B5, 1 (1968).
- G. Alexander, O. Benary, and U. Maor, Data Compilation of Baryon-Baryon Interactions. (II) Proton-Neutron Collisions Between 1 and 27 GeV/c, Nucl. Phys. B7, 281 (1968).
- G. Alexander, O. Benary, U. Karshon, and U. Maor, Data Compilation of Baryon-Baryon Interactions. (III) Hyperon-Proton Collisions, Nucl. Phys. B10, 554 (1969).

Acknowledgments

We wish to thank Arlene Wells for her general help in handling the data. We also thank David Herndon for preparing the plots in Fig. 15b and Marjorie Hutchinson for her assistance with some of the programming.

Section II.

K^+N DATA

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

Renormalization of Bubble Chamber Cross Sections

The total K^+p and K^+d cross sections have been measured quite accurately between 0.6 and 3.3 GeV/c by the recent counter experiments of Bugg et al. (Ref. 30 at back of this report), Cool et al. (Ref. 49), and Abrams et al. (Ref. 40). (For the latter two experiments we use revised data received by private communication.) Most partial cross sections have been measured in bubble chambers and normalized to the number of observed beam-track decays. These values can be corrected and their errors slightly reduced by renormalizing to the more-precisely-measured counter total cross sections.

The method we have used is to least-square the (interpolated) counter measurement of the total cross section with the value obtained in the bubble chamber experiment. Having this number, one can then correct the partial cross sections (and their errors).

The corresponding angular distributions were also rescaled, except that we did not change the percentage errors in these.

Correcting for Unmeasured Decay Modes

We have made use of isotopic spin-invariance of strong interactions to simplify some results. Cross sections for resonance production are corrected for unmeasured decay modes by isotopic spin invariance. Various cross sections proceeding from a pure $I = 0$ state are also quoted; isotopic-spin relations among the inelastic final states are discussed in detail by Hirata et al. (Ref. 34).

K^+p Interactions

As an introduction to K^+N interactions we summarize the available cross-section data in Fig. 1. The $I = 1$ curves are adapted from Bland et al. (Ref. 72), and the $I = 0$ from Hirata et al. (Ref. 34). The πN $I = 1/2$ and $I = 3/2$ total cross sections are also shown for comparison. The KN total cross section is distinguished from the πN and $\bar{K}N$ cross sections by its relative lack of structure. This is also true for the angular distributions. The other striking feature of KN is the similarity between the $I = 0$ and $I = 1$ cross sections.

Cross Sections

The K^+p total cross section (Figs. 2-4) is now known quite precisely from about 1 to 20 GeV/c. Below 1 GeV/c the present data are considerably less precise and somewhat confused. Further counter measurements in this region are necessary. Above 1 GeV/c there is strikingly little structure, as compared with πN or K^-N total cross sections (see insert to Fig. 1). There is a peak at 1.25 GeV/c and small ripples at 1.8 and 2.6 GeV/c, but no resonant behavior has been seen at these momenta in the elastic or inelastic channels. The total cross section from 3 to 20 GeV/c is consistent with a constant value of about 17.2 mb.

The single-pion-production cross section (see Figs. 3 and 5) rises abruptly at $K\Delta$ threshold, 0.8 GeV/c (presumably causing the steep rise in the total cross section at the same momentum). It peaks near 1.5 GeV/c and falls off slowly to 3.5 GeV/c. Above that momentum only the charge mode $p K_1^0 \pi^+$ (i. e.,

with a K_1 decay) has been measured, because it is hard to identify the missing neutrals in the modes $p K^+ \pi^0$ or $n K^+ \pi^+$. The $p K^0 \pi^+$ cross-section drops smoothly up to 12.3 GeV/c.

Double pion production occurs in five charge states:

$$K^+ p \rightarrow K^+ p \pi^+ \pi^- \quad (1)$$

$$\rightarrow K^0 p \pi^+ \pi^0 \quad (2)$$

$$\rightarrow K^0 n \pi^+ \pi^+ \quad (3)$$

$$\rightarrow K^+ p \pi^0 \pi^0 \quad (\text{underconstrained in HBC}) \quad (4)$$

$$\rightarrow K^+ n \pi^0 \pi^0 \quad (\text{underconstrained in HBC}) \quad (5)$$

Reactions 4 and 5 have two missing neutrals and cannot be positively identified in the bubble chamber. Their cross sections can be estimated, however, under the assumption of $K^* \Delta$ dominance, introduced in Bland et al. (Ref. 72). $K^* \Delta$ production occurs in reactions 1 to 5 in the ratios 18:13:2:2:1, and so for the $K N \pi \pi$ cross section we take the sum of reactions 1 and 2 multiplied by 36/33. [The presence of other quasi-two-body final states does not drastically affect this procedure; for $K^* N^*$ (where N^* represents any $I = 1/2 \pi p$ state) production the multiplicative factor becomes 3/2; for $K^+ p \rightarrow K^* p$, $K^* \rightarrow K^* \pi$, 8/9; and for $K^+ p \rightarrow K^* p$, $K^* \rightarrow K p$, 1.0.] The double-pion production cross sections are given in Fig. 6 and the three-pion production cross sections in Fig. 7.

In Fig. 8 we show production cross sections for the three main quasi-two-body final states: $K \Delta$ (1236), $K^* (890) p$, and $K^* (890) \Delta$ (1236). Hyperon production of strange particles by K^+ mesons can occur only in three-body final states. It has not been observed below 1.96 GeV/c and is small at all momenta where measured. Cross sections for associated production and other rare reactions are given in Table 1.

Elastic Angular Distributions

The differential cross section for $K^+ p$ elastic scattering (Figs. 9, 10, and 12) is fairly well known from threshold to 2 GeV/c,

and at intervals of 1 to 2 GeV/c up to 14 GeV/c. The angular distributions are relatively uncomplicated (again compared with πN or $K^+ N$), and a forward diffraction peak dominates from 1.2 GeV/c up.)

We have fitted all data in the forward direction - $t < 0.6$ (GeV/c)² to the form $d\sigma/dt = A e^{Bt}$. The coefficients A and B are shown in Fig. 11. The high-energy forward differential cross section is larger than the minimum value calculated from the optical theorem, implying a substantial real part of the forward scattering amplitude. For a total cross section of 17.2 mb, the optical theorem predicts $d\sigma/dt|_{t=0} = 15.2 \text{ mb}/(\text{GeV}/c)^2$, but our exponential fits to the data for - $t < 0.6$ (see Fig. 11) give values of about 20 mb/(GeV/c)². This result is, however, sensitive to normalization errors (estimated at 7% for the highest energy data but often not well known) as well as a possible turndown of the differential cross section in the unmeasured region of t , - $t < 0.05$ (GeV/c)². Accurate measurement of small-angle elastic scattering would be very interesting in this respect.

$K^+ p$ backward scattering (Fig. 12) has been measured by several experiments (counter and bubble chamber) measuring only scatters near 180 deg and, with less accuracy, by the bubble chamber experiments measuring the entire angular region. There is substantial disagreement between the precise counter data of Carroll et al. [Ref. 61] and other measurements near 1 GeV/c, leaving the correct experimental facts in that region somewhat in doubt. A backward peak is present at all momenta above 0.8 GeV/c. We have fitted data for $\cos \theta < -0.6$ to the form $d\sigma/du = A e^{B(u-u_0)}$; the coefficients are displayed in Fig. 13.

Polarization in Elastic Scattering

Until very recently data on proton polarization in $K^+ p$ elastic scattering were almost nonexistent. Accurate data now exist over a wide range of momenta, as seen in Fig. 14.

Elastic Phase Shift Analyses

Until very recently the absence of polarization data precluded a model-independent phase shift analysis above inelastic threshold. Some preliminary results of analyses using the polarization data are now available. In Fig. 15 we show some results presented at the 1969 Lund Conference. The S and P Argand plots are summaries given by rapporteur R. Levi-Setti; the D plots are from Andersson et al. (Ref. 54). While there is evidently still disagreement among the various analyses, it appears that the 1250-MeV/c peak in the total cross section (Fig. 15c) is not due to an s-channel resonance. The possibility of resonant states at higher momenta is still open; the P_{13} wave is now the best resonance candidate. There are (at least) two possible interpretations of the P_{13} trajectories as shown in Fig. 15a: (1) it resonates, or (2) above about 900 MeV/c it just rapidly becomes inelastic, with little phase variation. More sophisticated analyses such as have been done for πN elastic scattering and higher-energy data are probably called for to resolve this question. The only amplitude at all suggestive of resonant behavior is P_{13} .

The P_{13} Argand plots in Fig. 15b show two sets of possible solutions of the CERN-Holland data (Ref. 54) and two of the Argonne-Maryland data (Ref. 90). One solution from each of these groups was presented by Levi-Setti in his rapporteur talk at Lund and also appears in Figs. 15a and 15c.

Next to each Argand plot we show the corresponding "speed," i. e., the distance between two points on the Argand plot divided by the corresponding difference in energy.

The speed plots are interesting because if a resonance is present, the speed should be a maximum at resonance (assuming the background in that partial wave does not have too rapid an energy variation).

This can easily be seen by first considering an elastic resonance with no background. Then

$$T = \frac{1}{\epsilon - i}.$$

Then the complex derivative of T is

$$\frac{dT}{d\epsilon} = -\frac{1}{(\epsilon - i)^2}$$

and the "speed," defined as $\left| \frac{dT}{d\epsilon} \right|$, is

$$\frac{1}{|(\epsilon - i)^2|} = \frac{1}{\sqrt{(\epsilon - i)^2 \cdot (\epsilon + i)^2}} = \frac{1}{\epsilon^2 + 1} = \text{Im}(T).$$

If the resonance is inelastic, the circle $(\epsilon - i)^{-1}$ is merely reduced by a constant elasticity factor, so the argument does not change. If there is a constant background it makes no difference to the velocity, but of course we have to measure the "imaginary" part of T from the base of the resonance circle rather than from $T = 0$.

When one considers the Argand plot as well as the speed diagram for the CERN-HOLL II solution in Fig. 15b, it is not clear whether the large peak in speed around 1.9 GeV is really due to a resonance or just to noise.

Figure 15c (as presented by Levi-Setti in his Lund rapporteur talk) shows the K^+p total cross section together with the contribution to the total cross section from different partial waves. These partial cross sections are calculated from the expression

$$\sigma_{el}^{\ell} = 4\pi \lambda^2 \left(J + \frac{1}{2}\right) \text{Im}[T^{\ell}(0)].$$

Quasi-Two-Body Angular Distributions

We have data on the differential cross section for the $K\Delta$ final state from 860 MeV/c to 12.7 GeV/c (see Fig. 16). The lower limit is below the 870-MeV/c threshold for Δ production at the nominal Δ mass of 1236 MeV, but Δ production through the low-mass tail still dominates single pion production.

All of the distributions include background, about 50% at low energies and decreasing to

$\approx 10\%$ above 1500 MeV/c.

The differential cross section for the K^+p final state (Fig. 17) has been measured from 1200 MeV/c to 12.7 GeV/c. At low momenta (below 1.5 GeV/c) this reaction is not well separated from Δ production; the effect of this bias is difficult to estimate.

The $K^+\Delta$ differential cross section is given in Fig. 18.

K^+d and K^+n Interactions

Special Problems with Deuterium

Use of deuterium as a target involves certain problems not encountered with hydrogen.

1) The spectator proton is invisible in the bubble chamber when its momentum is below about 90 MeV/c (range of 2 mm).

In the present generation of fitting programs (SIOUX, GRIND) an unseen proton is approximated by a "spectator proton" with momentum components constrained to roughly 0 ± 40 MeV/c. The unseen proton thus becomes a (poorly) measured track. This uncertainty in spectator momentum then has a smearing effect on mass distributions and makes resonance-background separation much more difficult.

2) There are frequently neutrons in the final states. This means that processes like π^0 production are often underconstrained.

K^+d Cross Sections

The K^+d total cross section is shown in Fig. 19, along with cross sections for the quasi-elastic reactions $K^+d \rightarrow K^+pn$, and $K^+d \rightarrow K^0pp$. The K^+pn cross section is not measured directly but is obtained by subtracting the K^0pp cross section from the total. Below 600 MeV/c the K^+d total cross section is essentially unknown, and little is known about $I = 0$ KN elastic scattering. There is reliable low-energy charge-exchange data, but more precise data on $K^+d \rightarrow K^+pn$ and $K^+d \rightarrow K^+d$ are needed to study $I = 0$ reactions

in this energy region.

It is interesting to note that K^+d elastic scattering has never been directly measured at any momentum.

The K^+d single-pion-production cross sections are given in Fig. 20. The total $KN\pi$ cross section was calculated by Hirata et al. (Ref. 34), using isospin conservation. The only reaction in which the deuteron has been positively identified is $K^+d \rightarrow K^0\pi^+d$, the cross section for which is shown in Fig. 21. It remains remarkably constant, over a momentum region in which the total single-pion production varies by almost a factor of 10.

K^+n Cross Sections

Deducing the properties of K^+n interactions from K^+d data is nontrivial. There are several effects involved:

a) Screening (Glauber effect) at the target nucleon by the "other" nucleon makes measured cross sections generally lower ($\approx 10\%$) than free-nucleon cross sections.

b) Double scattering; some, but not all, of the events are rejected by requiring at least one slow nucleon.

c) Pauli-principle suppression of forward scattering; two protons with no relative momentum may not be in the same spin state. The spin-nonflip amplitude is thereby suppressed more than the spin-flip, so correcting for this effect requires detailed knowledge of the free-nucleon amplitude involved, as well as the deuteron wave function. The correction is negligible for (roughly) $-t > 0.1$ (BeV/c)².

d) "Smearing"; the Fermi motion of the target nucleon gives a spread δm of the c.m. energy. Thus at a beam momentum of 1.5 GeV/c, $\delta m = \pm 35$ MeV. If there were no Fermi momentum, such a ± 35 -MeV δm would correspond to a spread in beam momentum of $\delta p = (\sqrt{s}/M_p) \delta m = (2000/938) \delta m \approx \pm 70$ MeV/c. Due to these effects, extracting or "unfolding" free-nucleon cross sections from K^+d data is

a somewhat controversial procedure at present. In most cases the K^+n data is uncorrected for at least some of the effects above.

The K^+n total cross section is given in Fig. 22 (Glauber corrections have been made, but in general other deuteron corrections have not been included), and the $I = 0$ total cross section, derived from those for K^+p and K^+n , in Fig. 24. (Glauber corrections are included in all but Hirata et al.) Both are subject to the theoretical problems of unfolding the K^+d total cross-section data. The difference between Carter's (Ref. 37) and Cool's (Ref. 49) $I = 0$ total cross section in Fig. 24 illustrates the uncertainties in this process.

There are few available data on K^+n single-pion-production cross sections. However, ignoring deuterium corrections of the order of 5 to 15%, one can assume that $\sigma(K^+n \rightarrow K^+\pi^-p) \approx \sigma(K^+d \rightarrow K^+\pi^-pp)$ and

$\sigma(K^+n \rightarrow K^0\pi^0p) \approx \sigma(K^+d \rightarrow K^0\pi^0pp)$. The reaction $K^+d \rightarrow K^0\pi^+pn$ can be divided into interactions on the neutron and on the proton by studying the nucleon lab momentum distributions. This has been done by Hirata et al. (Ref. 34), giving the cross sections for $K^+n \rightarrow K^0\pi^+n$ shown in Fig. 23. These cross sections are not corrected for shielding effects and are probably slightly low. In Fig. 24, $I = 0$ elastic and single-pion-production cross sections are given.

K^+n Charge-Exchange Angular Distributions

Although we plot only $K^+d \rightarrow K^0pp$, information on K^+n charge exchange on free neutrons is obtained from the reaction $K^+d \rightarrow K^0pp$, assuming that the slower nucleon is the spectator. Usually the spectator momentum is required to be below some cut-off value, 150 to 300 MeV/c. The scattering angle is then computed in the c.m. of the outgoing kaon and struck proton.

Figures and Tables

In the following pages are the figures and tables representing the data compilation. For every figure there is a corresponding table on the facing (or same) page. All tables and figures are photographed computer output.

Occasionally there are pieces of data that are tabulated but not plotted; generally these are obsolete data. Such data have parentheses around them in the tables.

Data in square brackets are numbers that have been calculated by us, e. g., by summing various partial modes given by different papers but at the same energy, or by applying isospin corrections.

K^+_p Cross Sections

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

Figure 1. KN cross sections in the $l = 0$ and $l = 1$ states. The curves are adapted from Bland [ref. 72] (for $l = 1$) and Hirata [ref. 34] (for $l = 0$) and are only approximate representations of the data. The $KN\pi\pi$ cross section assumes $K^+\Delta$ dominance, as discussed in our text. In the inset we show the pion-nucleon $l = 1/2$ and $l = 3/2$ total cross sections (A. N. Diddens et al., Phys. Rev. Letters **10**, 262 (1963) for comparison. On the abscissa we plot, instead of the pion beam momentum, the "equivalent" K^+ beam momentum, giving the same momentum in the center of mass.

$K^+ p$ interactions

P_{beam} (GeV/c)	σ_{total} (mb)	$\sigma_{elastic}$ (mb)	References	P_{beam} (GeV/c)	σ_{total} (mb)	$\sigma_{elastic}$ (mb)	References
.140	9.200 ± 2.100		GOLDHABER 62	2.147	(15.400 ± 1.200)		BURROWS 59 *
.140		(9.200 ± 2.100)	GOLDHABER 62	2.150	17.570	.189	ABRAMS 67
.175	12.500	2.200	GOLDHABER 62	2.196	17.540	.080	COOL 66
.175		(12.500 ± 2.200)	GOLDHABER 62	2.200	17.600	.189	ABRAMS 67
.205	11.500	1.700	GOLDHABER 62	2.260	17.100	.583	COOK 61
.205		(11.500 ± 1.700)	GOLDHABER 62	2.300	17.440	.184	ABRAMS 67
.235	11.200	1.600	GOLDHABER 62	2.350	17.520	.180	ABRAMS 67
.235		(11.200 ± 1.600)	GOLDHABER 62	2.387	(13.000 ± 1.100)		BURROWS 59 *
.265	10.000	1.600	GOLDHABER 62	2.396	17.550	.080	COOL 66
.265		(10.000 ± 1.600)	GOLDHABER 62	2.400	17.560	.186	ABRAMS 67
.355	11.700	1.200	GOLDHABER 62	2.450	17.480	.179	ABRAMS 67
.355		(11.700 ± 1.200)	GOLDHABER 62	2.473	17.250	.120	BUGG 68
.451	16.300	1.700	KYCIA 60	2.500	17.490	.182	ABRAMS 67
.520	12.200	1.300	GOLDHABER 62	2.550	17.100	.671	COOK 61
.520		(12.200 ± 1.300)	GOLDHABER 62	2.550	17.440	.179	ABRAMS 67
.522	15.200	1.300	KYCIA 60	2.600	17.500	.180	ABRAMS 67
.589	16.300	1.700	KYCIA 60	2.650	17.470	.179	ABRAMS 67
.592	14.360	1.240	BUGG 68	2.700	17.410	.179	ABRAMS 67
.620	12.910	.490	BUGG 68	2.750	17.410	.179	ABRAMS 67
.642	12.400	.900	GOLDHABER 62	2.800	17.400	.179	ABRAMS 67
.642		$(12.400 \pm .900)$	GOLDHABER 62	2.830	16.700	.583	COOK 61
.644	12.880	.480	BUGG 68	2.850	17.300	.178	ABRAMS 67
.713	11.140	.350	BUGG 68	2.900	17.340	.176	ABRAMS 67
.768	11.650	.330	BUGG 68	2.950	17.300	.178	ABRAMS 67
.770	13.000	.949	COOK 61	2.970			DEBAISIEUX 66
.778		$(12.600 \pm .400)$	FOCARDI 67	2.972	(20.000 ± 2.000)		SALLSTROM 67
.810		$(13.000 \pm .700)$	STUBBS 61	3.000	(24.500 ± 2.500)		VON DARDEL 60 *
.823	12.970	.250	BUGG 68	3.000	17.190	.176	ABRAMS 67
.864	13.210	.240	BUGG 68	3.050	17.140	.176	ABRAMS 67
.864	14.070	.550	BLAND 68	3.100	17.080	.178	ABRAMS 67
.864		$(11.980 \pm .270)$	BLAND 68	3.150	17.150	.176	ABRAMS 67
.891	14.390	.380	COOL 66	3.200	17.130	.174	ABRAMS 67
.904	14.230	.190	BUGG 68	3.250	17.500	1.200	BAKER 63
.938	14.590	.210	BUGG 68	3.250	17.130	.174	ABRAMS 67
.942	15.570	.300	COOL 66	3.300	17.140	.174	ABRAMS 67
.962	(17.500 ± 1.500)		HUMMUES 59 *	3.460			DEBAISIEUX 66
.969	15.630	.270	BUGG 68	3.500	(23.500 ± 2.500)		VON DARDEL 60 *
.969	$(15.280 \pm .900)$	11.680	BLAND 68	4.000	17.600	1.200	BAKER 63
.970	15.400	.671	COOK 61	4.970			DEBAISIEUX 66
.970		(10.200 ± 1.500)	COOK 63	5.000	(20.000 ± 1.500)		VON DARDEL 60 *
.992	15.970	.270	COOL 66	5.500	17.900	.800	BAKER 63
1.029	15.690	.170	BUGG 68	6.000	17.000	.100	GALBRAITH 65
1.043	17.390	.250	COOL 66	6.800			FOLEY 63
1.063	(19.600 ± 1.100)		BURROWS 59 *	7.000	18.400	.600	BAKER 63
1.084	17.040	.170	BUGG 68	7.300			CHIEN 69
1.094	17.120	.250	COOL 66	8.000	(19.500 ± 1.000)		VON DARDEL 60 *
1.140	18.020	.170	BUGG 68	8.000	17.300	.100	GALBRAITH 65
1.144	18.090	.150	COOL 66	8.500	18.700	.600	BAKER 63
1.170	18.100	.671	COOK 61	9.800			FOLEY 63
1.170		(10.700 ± 1.100)	COOK 63	10.000	18.800	.700	BAKER 63
1.189	18.060	.170	BUGG 68	10.000	17.300	.100	GALBRAITH 65
1.194	18.470	.170	COOL 66	10.900	18.100	.790	BAKER 63
1.207	(19.850 ± 1.280)	10.090	BLAND 68	11.500	19.000	.600	BAKER 63
1.238	18.110	.170	BUGG 68	12.000	17.300	.100	GALBRAITH 65
1.245	18.540	.120	COOL 66	12.500	18.300	.800	BAKER 63
1.293	18.440	.150	BUGG 68	12.800			FOLEY 63
1.295	18.610	.110	COOL 66	13.400	17.500	.600	BAKER 63
1.300	17.900	.949	COOK 61	14.000	17.400	.100	BAKER 63
1.345	18.440	.100	COOL 66	14.800			GALBRAITH 65
1.347	18.270	.160	BUGG 68	15.000	18.500	.700	FOLEY 63
1.370	$(18.800 \pm .800)$		BURROWS 59 *	15.000	17.000	.100	BAKER 63
1.395	18.270	.100	COOL 66	16.000	18.800	.600	GALBRAITH 65
1.408	17.970	.140	BUGG 68	16.000	17.100	.100	BAKER 63
1.440	18.100	.762	COOK 61	19.000	17.300	.900	GALBRAITH 65
1.445	18.040	.100	COOL 66	20.000	17.500	.100	BAKER 63
1.455	$(18.200 \pm .800)$	9.600	BETTINI 65				GALBRAITH 65
1.460	17.940	.160	BUGG 68				
1.495	17.930	.090	COOL 66				
1.512	$(16.300 \pm .800)$		BURROWS 59 *				
1.550	17.700	.208	ABRAMS 67				
1.563	17.660	.140	BUGG 68				
1.596	17.750	.090	COOL 66				
1.600	17.710	.203	ABRAMS 67				
1.646	17.860	.060	COOL 66				
1.654	$(18.300 \pm .900)$		BURROWS 59 *				
1.690	17.500	.671	COOK 61				
1.696	17.850	.060	COOL 66				
1.700	17.730	.195	ABRAMS 67				
1.746	17.800	.060	COOL 66				
1.750	17.830	.192	ABRAMS 67				
1.796	17.800	.060	COOL 66				
1.800	17.980	.206	ABRAMS 67				
1.850	17.770	.191	ABRAMS 67				
1.877	$(15.500 \pm .800)$		BURROWS 59 *				
1.896	17.810	.060	COOL 66				
1.900	17.790	.191	ABRAMS 67				
1.945	17.410	.160	BUGG 68				
1.950	17.750	.191	ABRAMS 67				
1.960	(19.400 ± 2.000)	6.800	CHINOWSKY 65				
1.970	16.900	.500	COOK 61				
1.970		$(5.600 \pm .400)$	COOL 66				
1.996	17.600	.080	COOK 63				
2.000	17.630	.186	COOL 66				
2.050	17.720	.191	ABRAMS 67				
2.076	17.510	.080	ABRAMS 67				
2.100	17.560	.189	COOL 66				
			ABRAMS 67				

* DATA WAS READ FROM A GRAPH

REFERENCES		REFERENCES	
1	BURROWS 59.....PRL	2	117
2	KYCIA 60.....PR	118	553
3	VON DARDEL 60.....PRL	5	333
4	COOK 61.....PRL	7	182
5	STUBBS 61.....PRL	7	188
6	GOLDHABER 62.....PRL	9	135
7	BAKER 63.....PR	129	2285
8	COOK 63.....PR	129	2743
9	FOLEY 63.....PRL	11	503
10	BETTINI 65.....PL	16	83
11	CHINOWSKY 65.....PR	13981411	
12	GALBRAITH 65.....PR	138	8913
13	COOL 66.....PRL	17	102
14	DEBAISIEUX 66.....NC	45A	885
15	DEBAISIEUX 66.....NC	43A	142
16	ABRAMS 67.....PRL	19	259
17	FOCARDI 67.....PL	248	314
18	SALLSTROM 67.....NC	49A	348
19	BLAND 68.....UCRL	18758	
20	BUGG 68.....PR	168	1466
21	CHIEN 69.....PL	288	615

Data in parentheses have not been included in the figure. ()

Figure 2. K⁺p total and elastic cross sections, over the full momentum range of existing measurements.

K⁺p interactions

P _{beam} (GeV/c)	σ_{total} (mb)	σ_{elastic} (mb)	$\sigma_{\text{KN}\pi}$ (mb)	$\sigma_{\text{KN}\pi\pi}$ (mb)	References	P _{beam} (GeV/c)	σ_{total} (mb)	σ_{elastic} (mb)	$\sigma_{\text{KN}\pi}$ (mb)	$\sigma_{\text{KN}\pi\pi}$ (mb)	References
1.140	9.200 ± 2.100	(9.200 ± 2.100)			GOLDBERGER 62	1.800	17.980 ± .206				ABRAMS 67
1.140					GOLDBERGER 62	1.850	17.770 ± .191				ABRAMS 67
1.175	12.500 ± 2.200	(12.500 ± 2.200)			GOLDBERGER 62	1.877	(15.500 ± .800)				BURROWES 59 *
1.175					GOLDBERGER 62	1.896	17.810 ± .060				COOL 66
1.205	11.500 ± 1.700	(11.500 ± 1.700)			GOLDBERGER 62	1.900	17.790 ± .191				ABRAMS 67
1.205					GOLDBERGER 62	1.945	17.410 ± .160				BUGG 68
1.235	11.200 ± 1.600	(11.200 ± 1.600)			GOLDBERGER 62	1.950	17.750 ± .191				ABRAMS 67
1.235					GOLDBERGER 62	1.960	(19.400 ± 2.000)	8.800 ± .530	7.400 ± .550	3.500 ± .300	CHINDENSKY 65
1.265	10.000 ± 1.600	(10.000 ± 1.600)			GOLDBERGER 62	1.980					BLAND 68
1.265					GOLDBERGER 62	1.970	16.900 ± .500				COOK 61
1.355	11.700 ± 1.200	(11.700 ± 1.200)			GOLDBERGER 62	1.970		2.600 ± .430			COOK 63
1.355					GOLDBERGER 62	1.986	17.600 ± .080				COOL 66
1.451	16.300 ± 1.700	(16.300 ± 1.700)			GOLDBERGER 62	2.000	17.630 ± .186				ABRAMS 67
1.520	12.200 ± 1.300	(12.200 ± 1.300)			GOLDBERGER 62	2.050	17.720 ± .191				ABRAMS 67
1.520					GOLDBERGER 62	2.096	17.510 ± .080				COOL 66
1.522	15.200 ± 1.300	(15.200 ± 1.300)			KYCIA 60	2.100	17.560 ± .189				ABRAMS 67
1.589	16.300 ± 1.700	(16.300 ± 1.700)			KYCIA 60	2.147	(15.400 ± 1.200)				BURROWES 59 *
1.592	14.360 ± 1.240	(14.360 ± 1.240)			BUGG 68	2.150	17.570 ± .189				ABRAMS 67
1.620	12.910 ± .490	(12.910 ± .490)			BUGG 68	2.196	17.540 ± .080				COOL 66
1.642					FISK 62	2.200	17.600 ± .189				ABRAMS 67
1.642	12.400 ± .900	(12.400 ± .900)	.060 ± .030		GOLDBERGER 62	2.260	17.100 ± .583				COOK 61
1.642					GOLDBERGER 62	2.300	17.440 ± .184				ABRAMS 67
1.644	12.880 ± .480	(12.880 ± .480)			BUGG 68	2.350	17.520 ± .180				ABRAMS 67
1.713	11.140 ± .350	(11.140 ± .350)	.180 ± .010		BUGG 68	2.387	(13.000 ± 1.100)				BURROWES 59 *
1.735					FILIPPAS 67	2.396	17.550 ± .080				COOL 66
1.768	11.650 ± .330	(11.650 ± .330)			BUGG 68	2.400	17.560 ± .186				ABRAMS 67
1.770	13.000 ± .949	(13.000 ± .949)			COOK 61	2.450	17.480 ± .179				ABRAMS 67
1.778					FOCARDI 67	2.473	17.250 ± .120				BUGG 68
1.785			.920 ± .033		FILIPPAS 67	2.500	17.490 ± .182				ABRAMS 67
1.810			.950 ± .163		FISK 62	2.500	17.100 ± .671				COOL 61
1.810					STUBBS 61	2.550	17.440 ± .179				ABRAMS 67
1.823	12.970 ± .250	(12.970 ± .250)			BUGG 68	2.600	17.500 ± .180				ABRAMS 67
1.864	13.210 ± .240	(13.210 ± .240)			BUGG 68	2.650	17.470 ± .179				ABRAMS 67
1.864	14.070 ± .550	(14.070 ± .550)	1.640 ± .073		BLAND 68	2.650				5.500 ± .400	BLAND 68
1.891	14.390 ± .380	(14.390 ± .380)			COOL 66	2.700	17.410 ± .179				ABRAMS 67
1.904	14.230 ± .190	(14.230 ± .190)			BUGG 68	2.750	17.410 ± .179				ABRAMS 67
1.938	14.590 ± .210	(14.590 ± .210)			BUGG 68	2.800	17.400 ± .179				ABRAMS 67
1.942	15.570 ± .300	(15.570 ± .300)			COOL 66	2.830	16.700 ± .583				COOK 61
1.962	(17.500 ± 1.500)	(17.500 ± 1.500)			BURROWES 59 *	2.850	17.300 ± .178				ABRAMS 67
1.969	15.630 ± .220	(15.630 ± .220)			BUGG 68	2.900	17.340 ± .176				ABRAMS 67
1.969	15.280 ± .900	(15.280 ± .900)	11.680 ± .210	3.670 ± .143	BLAND 68	2.950	17.300 ± .178				ABRAMS 67
1.970	15.400 ± .671	(15.400 ± .671)			COOK 61	2.970		8.800 ± .400			DEBAISIEUX 66
1.992	15.970 ± .270	(15.970 ± .270)			COOL 66	2.970				5.500 ± .700	BLAND 68
1.992	15.690 ± .170	(15.690 ± .170)			BUGG 68	2.972	(20.000 ± 2.000)				SALLSTROM 67
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66	3.000	(24.500 ± 2.500)				VON DARDEL 60 *
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66	3.000	17.190 ± .176				ABRAMS 67
1.992	15.690 ± .170	(15.690 ± .170)			BURROWES 59 *	3.050	17.140 ± .176				ABRAMS 67
1.992	15.690 ± .170	(15.690 ± .170)			BUGG 68	3.100	17.080 ± .178				ABRAMS 67
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66	3.150	17.150 ± .176				ABRAMS 67
1.992	15.690 ± .170	(15.690 ± .170)			BUGG 68	3.200	17.130 ± .174				ABRAMS 67
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66	3.250	17.500 ± 1.200				BAKER 63
1.992	15.690 ± .170	(15.690 ± .170)			COOK 61	3.250	17.130 ± .174				ABRAMS 67
1.992	15.690 ± .170	(15.690 ± .170)			COOK 61	3.300	17.140 ± .174				ABRAMS 67
1.992	15.690 ± .170	(15.690 ± .170)			BUGG 68	3.460		8.360 ± .360			DEBAERE 66
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66	3.460				5.300 ± .500	BLAND 68
1.992	15.690 ± .170	(15.690 ± .170)			BLAND 68	3.500	(23.500 ± 2.500)				VON DARDEL 60 *
1.992	15.690 ± .170	(15.690 ± .170)			BUGG 68	3.500	17.600 ± 1.200			4.000 ± .200	BASSOMPIER 68
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66	4.000		8.820 ± .410			BAKER 63
1.992	15.690 ± .170	(15.690 ± .170)			BUGG 68	4.970					DEBAERE 66
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66	5.000	(20.000 ± 1.500)				VON DARDEL 60 *
1.992	15.690 ± .170	(15.690 ± .170)			COOK 61	5.000				[8.000 ± .500]	
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66						
1.992	15.690 ± .170	(15.690 ± .170)			BUGG 68						
1.992	15.690 ± .170	(15.690 ± .170)			BLAND 68						
1.992	15.690 ± .170	(15.690 ± .170)			BURROWES 59 *						
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66						
1.992	15.690 ± .170	(15.690 ± .170)			BUGG 68						
1.992	15.690 ± .170	(15.690 ± .170)			COOK 61						
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66						
1.992	15.690 ± .170	(15.690 ± .170)			BLAND 68						
1.992	15.690 ± .170	(15.690 ± .170)			BUGG 68						
1.992	15.690 ± .170	(15.690 ± .170)			BETTINI 65						
1.992	15.690 ± .170	(15.690 ± .170)			BUGG 68						
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66						
1.992	15.690 ± .170	(15.690 ± .170)			BLAND 68						
1.992	15.690 ± .170	(15.690 ± .170)			BURROWES 59 *						
1.992	15.690 ± .170	(15.690 ± .170)			ABRAMS 67						
1.992	15.690 ± .170	(15.690 ± .170)			BUGG 68						
1.992	15.690 ± .170	(15.690 ± .170)			BLAND 68						
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66						
1.992	15.690 ± .170	(15.690 ± .170)			ABRAMS 67						
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66						
1.992	15.690 ± .170	(15.690 ± .170)			BURROWES 59 *						
1.992	15.690 ± .170	(15.690 ± .170)			COOK 61						
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66						
1.992	15.690 ± .170	(15.690 ± .170)			BLAND 68						
1.992	15.690 ± .170	(15.690 ± .170)			BUGG 68						
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66						
1.992	15.690 ± .170	(15.690 ± .170)			ABRAMS 67						
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66						
1.992	15.690 ± .170	(15.690 ± .170)			BURROWES 59 *						
1.992	15.690 ± .170	(15.690 ± .170)			COOK 61						
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66						
1.992	15.690 ± .170	(15.690 ± .170)			BLAND 68						
1.992	15.690 ± .170	(15.690 ± .170)			BUGG 68						
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66						
1.992	15.690 ± .170	(15.690 ± .170)			ABRAMS 67						
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66						
1.992	15.690 ± .170	(15.690 ± .170)			BURROWES 59 *						
1.992	15.690 ± .170	(15.690 ± .170)			COOK 61						
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66						
1.992	15.690 ± .170	(15.690 ± .170)			BLAND 68						
1.992	15.690 ± .170	(15.690 ± .170)			BUGG 68						
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66						
1.992	15.690 ± .170	(15.690 ± .170)			ABRAMS 67						
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66						
1.992	15.690 ± .170	(15.690 ± .170)			BURROWES 59 *						
1.992	15.690 ± .170	(15.690 ± .170)			COOK 61						
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66						
1.992	15.690 ± .170	(15.690 ± .170)			BLAND 68						
1.992	15.690 ± .170	(15.690 ± .170)			BUGG 68						
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66						
1.992	15.690 ± .170	(15.690 ± .170)			ABRAMS 67						
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66						
1.992	15.690 ± .170	(15.690 ± .170)			BURROWES 59 *						
1.992	15.690 ± .170	(15.690 ± .170)			COOK 61						
1.992	15.690 ± .170	(15.690 ± .170)			COOL 66						
1.992	15.690 ± .170	(15.690 ± .170)			BLAND 68						
1.992	15.690 ± .170	(15.690 ± .170)			BUGG 68

Figure 3. K⁺p total, elastic, total-single-pion production, and total-double-pion production cross sections. The K N $\pi\pi$ cross section assumes K^{*} Δ dominance, as discussed in the text.

K^+p total cross section

P_{beam} (GeV/c) $E_{\text{c.m.}}$ σ_{total} (mb) References

1.029	.001	1.807	15.690	.170	BUGG	68
1.043	.010	1.814	17.390	.250	COOL	66
1.084	.001	1.833	17.040	.170	BUGG	68
1.094	.011	1.838	17.120	.250	COOL	66
1.140	.001	1.859	18.020	.170	BUGG	68
1.144	.011	1.861	18.090	.150	COOL	66
1.189	.001	1.882	18.060	.170	BUGG	68
1.194	.012	1.884	18.470	.120	COOL	66
1.238	.001	1.904	18.110	.170	BUGG	68
1.245	.012	1.907	18.540	.120	COOL	66
1.293	.001	1.929	18.440	.150	BUGG	68
1.295	.013	1.930	18.610	.110	COOL	66
1.345	.013	1.953	18.440	.100	COOL	66
1.347	.001	1.954	18.270	.160	BUGG	68
1.395	.014	1.975	18.270	.100	COOL	66
1.408	.001	1.981	17.970	.140	BUGG	68
1.445	.014	1.997	18.040	.100	COOL	66
1.468	.001	2.008	17.940	.160	BUGG	68
1.495	.015	2.020	17.930	.090	COOL	66
1.550	.012	2.044	17.700	.110	ABRAMS	67
1.563	.001	2.049	17.660	.140	BUGG	68
1.596	.016	2.064	17.750	.090	COOL	66
1.600	.012	2.066	17.710	.100	ABRAMS	67
1.646	.016	2.085	17.860	.060	COOL	66
1.696	.017	2.107	17.850	.060	COOL	66
1.700	.013	2.109	17.730	.080	ABRAMS	67
1.746	.017	2.128	17.800	.060	COOL	66
1.750	.013	2.130	17.830	.070	ABRAMS	67
1.796	.018	2.149	17.800	.060	COOL	66
1.800	.013	2.151	17.980	.100	ABRAMS	67
1.850	.014	2.172	17.770	.070	ABRAMS	67
1.896	.019	2.191	17.810	.060	COOL	66
1.900	.014	2.193	17.790	.070	ABRAMS	67
1.945	.001	2.211	17.410	.160	BUGG	68
1.950	.015	2.213	17.750	.070	ABRAMS	67
1.996	.020	2.232	17.600	.080	COOL	66
2.000	.015	2.234	17.630	.060	ABRAMS	67
2.050	.015	2.254	17.720	.070	ABRAMS	67
2.096	.021	2.273	17.510	.080	COOL	66
2.100	.016	2.274	17.560	.070	ABRAMS	67
2.150	.016	2.294	17.570	.070	ABRAMS	67
2.196	.022	2.313	17.540	.080	COOL	66
2.200	.016	2.314	17.600	.070	ABRAMS	67
2.300	.017	2.353	17.440	.060	ABRAMS	67
2.350	.018	2.373	17.520	.040	ABRAMS	67
2.396	.024	2.391	17.550	.080	COOL	66
2.400	.018	2.392	17.560	.060	ABRAMS	67
2.450	.018	2.411	17.480	.040	ABRAMS	67
2.473	.001	2.420	17.250	.120	BUGG	68
2.500	.019	2.430	17.490	.050	ABRAMS	67
2.550	.019	2.449	17.440	.040	ABRAMS	67
2.600	.019	2.468	17.500	.040	ABRAMS	67
2.650	.020	2.486	17.470	.040	ABRAMS	67
2.700	.020	2.505	17.410	.040	ABRAMS	67
2.750	.021	2.523	17.410	.040	ABRAMS	67
2.800	.021	2.542	17.400	.040	ABRAMS	67
2.850	.021	2.560	17.300	.040	ABRAMS	67
2.900	.022	2.578	17.340	.030	ABRAMS	67
2.950	.022	2.596	17.300	.040	ABRAMS	67
3.000	.022	2.613	17.190	.040	ABRAMS	67
3.050	.023	2.631	17.140	.040	ABRAMS	67
3.100	.023	2.649	17.080	.050	ABRAMS	67
3.150	.024	2.666	17.150	.040	ABRAMS	67
3.200	.024	2.683	17.130	.030	ABRAMS	67
3.250	.024	2.701	17.130	.030	ABRAMS	67
3.300	.025	2.718	17.140	.030	ABRAMS	67

NO SYSTEMATIC ERRORS HAVE BEEN INCLUDED EVEN THOUGH SOME AUTHORS MAY HAVE GIVEN THEM

REFERENCES			
1	COOL	66.....PRL	17 102
2	ABRAMS	67.....PRL	19 259
3	BUGG	68.....PR	168 1466

CNTR
CNTR
CNTR

Figure 4. The $K^+ p$ total cross section from 1 to 3.5 GeV/c, as measured by three recent counter experiments. Systematic errors have not been included in this figure in order to see fine structure more easily.

K⁺ p interactions

P_{beam} (GeV/c)	$\sigma_{\text{KN}\pi}$ (mb)	$\sigma_{\text{K}^+\text{n}\pi^+}$ (mb)	$\sigma_{\text{K}^+\text{p}\pi^0}$ (mb)	$\sigma_{\text{K}^0\text{p}\pi^+}$ (mb)	References
.642	.060 $\pm$.030				FI SK 62
.735	.180 .010	.020 $\pm$.010	.030 $\pm$.010	.130 $\pm$.010	FILIPPAS 67
.755			.110 .019		FILIPPAS 67
.785	.520 .030	.070 .010		.340 .030	FILIPPAS 67
.810	.950 .160	.110 .040	.220 .050	.640 .080	FI SK 62
.864	1.640 .070	.131 .017	.360 .030	1.150 .060	BL AND 68
.910				2.100 .200	KEHOE 63
.969	3.670 .140	.300 .040	.800 .070	2.570 .120	BL AND 68
1.140				4.600 .300	ROLDT 64
1.207	7.400 .170	.560 .060	1.800 .100	5.040 .150	BL AND 68
1.367				5.360 .330	BL AND 68
1.455	8.160 .310	1.350 .100	1.900 .200	4.900 .300	BETTINI 65
1.585				5.000 .400	BL AND 68
1.960	7.400 .550	1.500 .300	1.900 .300	4.100 .500	CHINOWSKY 65
2.260				2.620 .300	BOHSE 67
2.650				2.700 .300	NEWMAN 67
2.965				2.100 .300	FERRO-LUZZ 65
2.972	4.400 .400	1.000 .200	1.100 .200	2.300 .300	SALLSTROM 67
3.500	4.000 .200	.800 .100	1.320 .070	1.850 .100	BASSOMPTIER 68
3.535				1.388 .029	BI SHOP 67
4.600				1.200 .120	FU 68 *
5.000				.900 .100	BASSOMPTIER 68

* DATA WAS READ FROM A GRAPH

REFERENCES				
1	FISK	62.....CERN CONF	358	HBC
2	KEHOE	63.....PRL	11 93	HLBC
3	BOLDT	64.....PR	133 8220	HBC
4	RAFTINI	65.....PL	16 83	HBC
5	CHINOWSKY	65.....PR	15981411	HBC
6	FERRER-LUZ	65.....NC	36 1101	HBC
7	BISHOP	67.....THESIS		HBC
8	BOMSE	67.....PR	158 1298	HBC
9	FILIPPAS	67.....NC	51A 1053	HBC
10	NEWMAN	67.....PR	158 1310	HBC
11	SALLSTROM	67.....NC	49A 348	HBC
12	BASSOMPIER	68.....PL	278 468	HBC
13	BLAND	68.....UCRL	18758	HBC
14	FU	68.....ICARL	18417	HBC

Figure 5. $K^+ p$ single-pion production cross sections from 0 to 5 GeV/c. The dashed line joins the $K\pi N$ cross section points. $K\pi N = nK^+\pi^+ + pK^+\pi^0 + pK^0\pi^+$.

$K^+ p$ interactions

P_{beam} (GeV/c)	$\sigma_{KN\pi\pi}$ (mb)	$\sigma_{K^0 n \pi^+ \pi^+}$ (mb)	$\sigma_{K^+ p \pi^+ \pi^-}$ (mb)	$\sigma_{K^0 p \pi^+ \pi^0}$ (mb)	References
1.267	$.029 \pm .011$		$.026 \pm .010$		BLAND 68
1.367	$.147 \pm .025$	$.009 \pm .012$ - .006	$.065 \pm .012$	$.060 \pm .018$	BLAND 68
1.450	$.280 \pm .070$				BLAND 68
1.455		$.020 \pm .020$	$.150 \pm .040$	$.080 \pm .050$	RETTINI 65
1.585	$.860 \pm .100$	$.080 \pm .030$	$.380 \pm .050$	$.320 \pm .070$	BLAND 68
1.960		$.310 \pm .100$	$1.640 \pm .200$	$1.240 \pm .200$	CHINOWSKY 65
1.960	$3.500 \pm .300$				BLAND 68
2.260			$1.700 \pm .200$		BOMSE 67
2.650	$5.500 \pm .400$				BLAND 68
2.650		$.640 \pm .120$	$2.500 \pm .200$	$1.900 \pm .200$	NEWMAN 67
2.970		$.600 \pm .200$	$2.300 \pm .300$	$2.100 \pm .300$	FERRO-LUZZ 65
2.970	$5.500 \pm .700$				BLAND 68
3.460	$5.300 \pm .500$				BLAND 68
3.500		$.600 \pm .100$	$2.100 \pm .300$	$2.500 \pm .300$	BASSOMPIER 68
3.535		$.506 \pm .017$		$1.831 \pm .034$	BISHOP 67
5.000	$[4.000 \pm .300]$	$.400 \pm .050$	$1.800 \pm .150$	$1.500 \pm .200$	BASSOMPIER 68
5.438			$1.790 \pm .100$	$1.450 \pm .150$	BOMSE 68
12.700		$.150 \pm .030$			BERLINGHIE 68
12.730	$[1.800 \pm .200]$		$.950 \pm .100$	$.565 \pm .070$	BERLINGHIE 68
REFERENCES					
1	BETTINI 65.....PL	16 83	HBC		
2	CHINOWSKY 65.....PR	13981411	HBC		
3	FERRO-LUZZ 65.....NC	39 417	HBC		
4	BISHOP 67.....THESIS		HBC		
5	BOMSE 67.....PR	158 1281	HBC		
6	NEWMAN 67.....PR	158 1310	HBC		
7	BASSOMPIER 68.....PL	278 468	HBC		
8	BERLINGHIE 68.....PL	278 665	HBC		
9	BERLINGHIE 68.....CERN 68-7	172	HBC		
10	BLAND 68.....UCRL	18758	HBC		
11	BOMSE 68.....PRL	20 1519	HBC		

Data in square brackets have been calculated by us.

Figure 6. Cross sections for the three double-pion-production charge states measured in the bubble chamber, and the $KN\pi\pi$ cross section, computed assuming $K^* \Delta$ dominance as discussed in the text.

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

Figure 7. Cross sections for triple pion production.

P_{beam} (GeV/c)	$\sigma_{K^+p\pi^+\pi^-\pi^0}$ (mb)	$\sigma_{K^+n\pi^+\pi^-\pi^-}$ (mb)	$\sigma_{K^0p\pi^+\pi^-\pi^-}$ (mb)	References
1.585	$.009 \pm .010$ - .005		$.006 \pm .015$ - .004	BLAND 68
1.960	.050 .020	$.010 \pm .006$	.020 .010	CHINOWSKY 65
2.650	.430 .060	.140 .030	.190 .050	NEWMAN 67
3.000	.620 .050	.150 .030	.350 .040	BASSOMPIER 68
3.500	.710 .040	.190 .020	.440 .040	BASSOMPIER 68
5.000	1.100 .100	.310 .030	.620 .050	BASSOMPIER 68
REFERENCES				
1	CHINOWSKY 65.....PR	13981411	HBC	
2	NEWMAN 67.....PR	158 1310	HBC	
3	BASSOMPIER 68.....PL	278 468	HBC	
4	BLAND 68.....UCRL	18758	HBC	

K⁺ p interactions

P _{beam}	$\sigma_{K^{*+}(890)p}$ ↳ K ⁰ π ⁺	$\sigma_{K^{*+}(890)p}$ ↳ K ⁺ π ⁰	$\sigma_{K^{*+}(890)N}$	$\sigma_{K\Delta^{++}(1238)}$ ↳ pπ ⁺	$\sigma_{K\Delta(1238)}$	$\sigma_{K^{*+}(890)\Delta^{++}(1238)}$ ↳ K ⁺ π ⁻ ↳ pπ ⁺	$\sigma_{K^{*+}(890)\Delta(1238)}$	References
(GeV/c)	(mb)	(mb)	(mb)	(mb)	(mb)	(mb)	(mb)	
.735				.080 ± .060	.110 ± .080			BLAND 68
.785				.190 .040	.250 .050			BLAND 68
.864				.860 .120	1.130 .150			BLAND 68
.969				2.420 .190	3.090 .230			BLAND 68
1.140	.900 ± .300		[1.343 ± .448]	3.600 .500	[4.824 .670]			BOLOT 64
1.207	1.060 .200	.550 ± .110	1.610 .228	3.430 .320	4.560 .350			BLAND 68
1.367	1.720 .240		[2.567 .358]	3.050 .330	4.100 .400			BLAND 68
1.455				2.400 .500	[3.216 .670]			BLAND 68
1.585	2.150 .350		[3.209 .522]	2.050 .340	2.700 .400	.250 ± .070	[.500 ± .140]	BETTINI 65
1.960	1.200 .200		[1.751 .299]	2.100 .300	[2.814 .402]			BLAND 68
2.260						.800 .100	[1.600 .200]	GOLDHABER 66
2.260	1.100 .200		[1.642 .299]	1.100 .200	[1.474 .268]			BOMSE 67
2.650	1.500 .300		[2.239 .448]	.900 .200	[1.206 .268]	1.330 .150	[2.660 .300]	90MSE 67
2.965	.800 .100		[1.194 .149]	.800 .100	[1.072 .134]			NEWMAN 67
2.970						1.200 .200	[2.400 .400]	FERRO-LUZZ 65
2.972	.950 .130	.440 .090	1.390 .158	.680 .110	.860 .130			FERRO-LUZZ 65
3.460						.920 .100	[1.840 .200]	SALLSTROM 68
3.535	.548 .023		[.813 .034]	.354 .015	[.474 .020]			GEORGE 67
4.600	.360 .040		[.537 .060]					BISHOP 67
4.970						.720 .060	[1.440 .120]	FU 68
12.700	.051 .012		[.076 .018]	.044 .007	[.059 .009]	.100 .030	[.200 .060]	GEORGE 67
								BERLINGHIE 68

↑ plotted	↑ plotted	↑ plotted
1 BOLDT 64.....PR	133 8220	HBC
2 BETTINI 65.....PL	16 83	HBC
3 FERRO-LUZZ 65.....NC	36 1101	HBC
4 FERRO-LUZZ 65.....NC	39 417	HBC
5 GOLDHABER 66.....PR	142 913	HBC
6 BISHOP 67.....THESIS		HBC
7 BOMSE 57.....PR	158 1298	HBC
8 BOMSE 57.....PR	158 1281	HBC
9 GEORGE 57.....NC	49A 9	HBC
10 NEWMAN 67.....PR	158 1310	HBC
11 BERLINGHIE 68.....NP	88 333	HBC
12 BLAND 68.....UCRL	18758	HBC
13 FU 68.....UCRL	18417	HBC
14 SALLSTROM 68.....AF	37 468	HBC

Data in square brackets have been calculated by us.

Figure 8a. K⁺p cross sections for resonance production via the K^{*}p, KΔ, and K^{*}Δ quasi-two-body final states over the full momentum range of existing measurements. Unmeasured decay modes have been corrected for using isotopic-spin branching ratios.

K⁺p interactions

Figure 8b. K⁺p cross sections for resonance production via the K^{*}p, KΔ, and K^{*}Δ, quasi-two-body final states for the momentum range from 0 to 5 GeV/c. Unmeasured decay modes have been corrected for using isotopic-spin branching ratios.

"Rare" K^+p Interactions

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

$K^+ p$ interactions

P_{beam} (GeV/c)	$\sigma_{K^+ p \pi^+ \pi^+ \pi^-} \text{ (mb)}$	$\sigma_{K^+ p \pi^+ \pi^+ \pi^-} \text{ (mb)}$	References
2.650	$.010 \pm .010$	$.040 \pm .040$	NEWMAN 67
1	REFERENCES NEWMAN 67.....PR	158 1310	HBC

P_{beam} (GeV/c)	$\sigma_{K^+ K^+ \Lambda} \text{ (mb)}$	$\sigma_{K^+ K^+ \Sigma^0} \text{ (mb)}$	$\sigma_{K^+ K^0 \Sigma^+} \text{ (mb)}$	$\sigma_{K^+ K^+ \Lambda \pi^0} \text{ (mb)}$	$\sigma_{K^+ K^0 \Lambda \pi^+} \text{ (mb)}$	References
2.650	$.030 \pm .020$	.010	$.020 \pm .020$	$.010 \pm .010$	$.030 \pm .030$	NEWMAN 67
2.970	.023 .004	$.007 \pm .002$	.015 .005	.007 .002	.012 .003	FERRO-LUZZ 65
3.000	.022 .003	.010 .002	.019 .005	.005 .001	.009 .002	JOLDERSMA 66
3.460	.022 .004	.007 .002	.008 .004	.009 .003		FERRO-LUZZ 65
3.535	.025 .003	.010 .002		.013 .002		BISHOP 67
4.970	.018 .006	.006 .003	.023 .009	.030 .008	.036 .009	FERRO-LUZZ 65
12.700				.014 .005	.026 .007	BERLINGHIE 68
12.730	.012 .003					BERLINGHIE 68
1	REFERENCES FERRO-LUZZ 65.....PL	17 155		HBC		
2	JOLDERSMA 66.....PRL	17 716		HBC		
3	BISHOP 67.....THESIS			HBC		
4	NEWMAN 67.....PR	158 1310		HBC		
5	BERLINGHIE 68.....PL	278 665		HBC		
6	BERLINGHIE 68.....CERN 68-7	172		HBC		

P_{beam} (GeV/c)	$\sigma_{K^0 p \omega \pi^+} \text{ (mb)}$	$\sigma_{K^0 p \eta \pi^+} \text{ (mb)}$	References
3.535	$.064 \pm .035$	$.049 \pm .007$	BISHOP 67
1	REFERENCES BISHOP 67.....THESIS		HBC

P_{beam} (GeV/c)	$\sigma_{pp \bar{\Lambda}} \text{ (mb)}$	$\sigma_{pp \bar{\Sigma}^0} \text{ (mb)}$	$\sigma_{pp \bar{\Lambda} \pi^0} \text{ (mb)}$	$\sigma_{pn \bar{\Lambda} \pi^+} \text{ (mb)}$	$\sigma_{pp \bar{\Lambda} \pi^+ \pi^-} \text{ (mb)}$	References
4.970	$.003 \pm .001$	$.001 \pm .001$				BASSOMPIER 67
9.000			$.009 \pm .002$	$.035 \pm .004$	$.003 \pm .001$	ALEXANDER 68
12.730	.006 .002					BERLINGHIE 68
1	REFERENCES BASSOMPIER 67.....NC	48 589		HBC		
2	ALEXANDER 68.....PRL	20 755		HBC		
3	BERLINGHIE 68.....CERN 68-7	172		HBC		

Table 1. Cross sections for associated production of strange particles and other rare reactions (not plotted).

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

K^+p Elastic Differential Cross Sections

All data below 2 GeV/c are plotted as functions of $\cos \theta_{c.m.}$. Above 1 GeV/c data are plotted as functions of t , except that experiments which report numbers of events (and do not give a factor which would enable us to convert to σ) have not been included.

For experiments above 0.75 GeV/c and which report data close enough to the backward direction, we plot $d\sigma/du$.

We have transformed from $t \leftrightarrow \cos \theta \leftrightarrow u$ as necessary.

Since for elastic scattering the expression for t in the c.m. is

$$t = -2q^2(1 - \cos \theta_{KK'}) , \quad (1)$$

we give in the tables $2q^2$ in $(\text{GeV}/c)^2$ and call it $2Q$ SQUARE.

$$\text{Also } u = \frac{(m_p^2 - m_K^2)^2}{s} - 2q^2(1 + \cos \theta_{KK'}) \quad (2)$$

so for backward scattering, u attains its maximum value u_0 ,

$$u_0 = \frac{(m_p^2 - m_K^2)^2}{s} \quad (3)$$

which we call UZERO.

We fit only data with $\cos \theta < -0.6$; u , given by (2), is then printed as CUT ON U in the appropriate table. If no CUT ON U is specified on a $d\sigma/du$ table, it means that all of the data in that distribution are already within our $\cos \theta < -0.6$ limit.

In addition we have fit both the forward and backward angular distributions to exponentials. The fitted parameters have been plotted in Figs. 11 and 13.

K^+p elastic differential cross section

DISTRIBUTION IN COS THETA OF THE SCATT BEAM WITH RESPECT TO THE BEAM DIRECTION IN THE C.M. SYSTEM

BEAM MOMENTUM= .140 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)	
-.900	-.600	.900	.500
-.600	-.200	1.100	.500
-.200	.200	.900	.300
.200	.600	1.400	.500
.600	.800	1.500	.700
.800	.950	11.000	2.500

2Q SQUARE= .017

GOLDHABER PRL 9 135(1962) HBC

BEAM MOMENTUM= .175 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)	
-1.000	-.600	1.100	.300
-.600	-.200	1.300	.400
-.200	.200	.800	.400
.200	.600	1.600	.400
.600	.800	2.300	.700
.800	.950	5.000	1.200

2Q SQUARE= .026

GOLDHABER PRL 9 135(1962) HBC

BEAM MOMENTUM= .205 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)	
-1.000	-.600	1.000	.200
-.600	-.200	1.000	.200
-.200	.200	1.300	.400
.200	.600	1.000	.200
.600	.800	1.500	.400
.800	.950	1.600	.400

2Q SQUARE= .035

GOLDHABER PRL 9 135(1962) HBC

BEAM MOMENTUM= .235 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)	
-1.000	-.600	1.100	.200
-.600	-.200	1.200	.200
-.200	.200	.500	.200
.200	.600	1.200	.200
.600	.800	1.000	.200
.800	.950	2.400	.500

2Q SQUARE= .045

GOLDHABER PRL 9 135(1962) HBC

BEAM MOMENTUM= .265 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)	
-1.000	-.600	.900	.200
-.600	-.200	.900	.200
-.200	.200	1.000	.200
.200	.600	.800	.200
.600	.800	1.300	.400
.800	.950	3.600	1.200

2Q SQUARE= .057

GOLDHABER PRL 9 135(1962) HBC

BEAM MOMENTUM= .355 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)	
-1.000	-.600	.900	.100
-.600	-.200	.800	.100
-.200	.200	1.150	.150
.200	.600	1.000	.150
.600	.950	1.250	.200

2Q SQUARE= .098

GOLDHABER PRL 9 135(1962) HBC

K^+p elastic differential cross section

Figure 9. Differential cross section $d\sigma/d\omega$ for K^+p elastic scattering up to 2 GeV/c. So as not to mask possible structure, systematic errors have not been included even though they may have been given by the authors. Some distributions given originally as $d\sigma/dt$ vs. t were transformed to $d\sigma/d\omega$ vs. $\cos \theta_{c.m.}$

K^+p elastic differential cross section

DISTRIBUTION IN COS THETA OF THE SCATT BEAM WITH RESPECT TO THE BEAM DIRECTION IN THE C.M. SYSTEM

BEAM MOMENTUM= .520 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)
-1.000	-.600	.900 .150
-.600	-.200	1.000 .150
-.200	.200	1.100 .150
.200	.600	1.100 .150
.600	.950	1.000 .150

2Q SQUARE= .193

GOLDHABER PRL 9 135(1962) HBC

BEAM MOMENTUM= .642 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)
-1.000	-.600	1.100 .120
-.600	-.200	.900 .120
-.200	.200	.950 .120
.200	.600	.950 .120
.600	.950	1.050 .120

2Q SQUARE= .274

GOLDHABER PRL 9 135(1962) HBC

BEAM MOMENTUM= .778 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)
-1.000	-.900	1.137 .064
-.900	-.800	1.073 .061
-.800	-.700	1.055 .060
-.700	-.600	.953 .057
-.600	-.500	1.137 .064
-.500	-.400	.984 .058
-.400	-.300	.885 .054
-.300	-.200	.929 .056
-.200	-.100	.929 .056
-.100	0.	.912 .055
0.	.100	.893 .055
.100	.200	.887 .055
.200	.300	.848 .053
.300	.400	.770 .052
.400	.500	.876 .055
.500	.600	1.064 .062
.600	.700	1.006 .060
.700	.800	1.059 .063
.800	.900	1.134 .065
.900	.950	1.375 .096

2Q SQUARE= .373

FOCARDI PL 248 314(1967) HBC

BEAM MOMENTUM= .810 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)
-1.000	-.720	1.130 + .090
-.720	-.440	1.100 + .080
-.440	-.160	.910 + .070
-.160	.120	.900 + .070
.120	.400	.890 + .060
.400	.680	.970 + .080
.680	.960	1.100 + .090
		- .070

2Q SQUARE= .398

STUBBS PRL 7 188(1961) HBC

BEAM MOMENTUM= .864 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)
-1.000	-1.000	1.600 .120
-.900	-.800	1.300 .060
-.800	-.700	1.100 .060
-.700	-.600	.990 .050
-.600	-.500	.900 .050
-.500	-.400	.870 .050
-.400	-.300	.910 .050
-.300	-.200	.770 .050
-.200	-.100	.800 .050
-.100	0.	.850 .050
0.	.100	.810 .050
.100	.200	.920 .050
.200	.300	.870 .050
.300	.400	.860 .050
.400	.500	.940 .050
.500	.600	1.040 .050
.600	.700	1.060 .060
.700	.800	1.060 .060
.800	.900	1.210 .060
.900	1.000	1.340 .130

2Q SQUARE= .439

BLAND UCRL 18889 (1969) HBC

BEAM MOMENTUM= .865 GEV/C

COS THETA	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)
-.910	.800 .100
-.720	1.000 .120
-.640	.860 .130
-.570	.850 .120
-.500	.850 .120
-.430	.800 .130
-.360	.790 .110
-.290	.890 .130
-.210	.900 .110
-.110	.860 .120
-.030	.870 .130
.060	.710 .100
.180	.780 .080
.310	1.050 .240
.370	.930 .230
.420	1.010 .310
.480	.880 .200
.530	.840 .200
.590	1.050 .220
.640	.880 .230
.700	1.310 .190
.790	1.040 .190

2Q SQUARE= .440

PLUS POSSIBLE SYSTEMATIC ERROR OF +- 15PER CENT

ANDERSSON CERN (1969) CNTR

K^+p elastic differential cross section

Figure 9 (Continued)

K^+p elastic differential cross section

DISTRIBUTION IN COS THETA OF THE SCATT BEAM WITH RESPECT TO THE BEAM DIRECTION IN THE C.M. SYSTEM

BEAM MOMENTUM= .910 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (NUMBER OF EVENTS)	HLBC
-1.000	-.800	116.000	10.770
-.800	-.600	106.000	10.296
-.600	-.400	112.000	10.583
-.400	-.200	100.000	10.000
-.200	0.	109.000	10.440
0.	.200	127.000	11.269
.200	.400	134.000	11.576
.400	.600	138.000	11.747
.600	.800	129.000	11.358
.800	.900	153.000	12.369

2Q SQUARE= .475

HIRSCH PR 135 8191(1964) HLBC 32

BEAM MOMENTUM= .969 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)
-1.000	-1.000	1.370
-.900	-.800	1.100
-.800	-.700	.950
-.700	-.600	.810
-.600	-.500	.750
-.500	-.400	.830
-.400	-.300	.740
-.300	-.200	.740
-.200	-.100	.810
-.100	0.	.820
0.	.100	.790
.100	.200	.780
.200	.300	.760
.300	.400	.960
.400	.500	1.050
.500	.600	1.170
.600	.700	.930
.700	.800	1.280
.800	.900	1.350
1.000	1.000	1.620

2Q SQUARE= .522

BLAND UCRL 18889 (1969) HBC

BEAM MOMENTUM= .970 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)	SPRK
-.960	-.800	.610	.110
-.800	-.600	.670	.090
-.600	-.400	.760	.100
-.400	-.200	.670	.080
-.200	0.	.570	.060
0.	.300	.750	.078
.300	.500	1.150	.110
.500	.700	.840	.090
.700	.820	1.070	.180

2Q SQUARE= .523

COOK PR 129 2743(1963) SPRK

BEAM MOMENTUM= .971 GEV/C

COS THETA	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)
-.840	.900
-.780	.870
-.720	.830
-.660	.870
-.600	.850
-.530	.740
-.460	.840
-.300	.720
-.310	.820
-.230	.820
-.150	.800
-.080	.850
-.020	.880
.030	.790
.170	.890
.380	.910
.440	1.100
.500	.990
.550	1.110
.600	1.220
.650	1.240
.740	1.150
.830	.960

2Q SQUARE= .524

PLUS POSSIBLE SYSTEMATIC ERROR OF +- 15PER CENT

ANDERSSON CERN (1969) CNTR

BEAM MOMENTUM= 1.087 GEV/C

COS THETA	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)
-.810	.510
-.720	.510
-.640	.520
-.580	.500
-.510	.410
-.430	.500
-.350	.460
-.230	.490
-.160	.450
-.110	.550
-.040	.490
.060	.500
.140	.600
.230	.830
.290	.920
.350	.800
.410	.920
.460	.870
.540	1.040
.630	.980
.710	1.390
.780	1.390
.850	1.240

2Q SQUARE= .618

PLUS POSSIBLE SYSTEMATIC ERROR OF +- 15PER CENT

ANDERSSON CERN (1969) CNTR

BEAM MOMENTUM= 1.170 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)
-.960	-.900	.730
-.900	-.800	.430
-.800	-.700	.360
-.700	-.600	.410
-.600	-.500	.240
-.500	-.400	.430
-.400	-.300	.360
-.300	-.200	.330
-.200	-.100	.350
-.100	0.	.420
0.	.300	.610
.300	.400	1.040
.400	.500	.950
.500	.600	1.190
.600	.700	1.260
.700	.860	1.890

2Q SQUARE= .687

COOK PR 129 2743(1963) SPRK

K^+p elastic differential cross section

Figure 9 (Continued)

K^+p elastic differential cross section

DISTRIBUTION IN COS THETA OF THE SCATT BEAM WITH RESPECT TO THE BEAM DIRECTION IN THE C.M. SYSTEM

BEAM MOMENTUM= 1.207 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)	
-1.000	-1.000	.570	.070
-1.000	-.900	.570	.070
-.900	-.800	.490	.050
-.800	-.700	.380	.050
-.700	-.600	.410	.050
-.600	-.500	.350	.050
-.500	-.400	.370	.050
-.400	-.300	.430	.050
-.300	-.200	.450	.050
-.200	-.100	.460	.050
-.100	0.	.510	.060
0.	.100	.540	.060
.100	.200	.660	.060
.200	.300	.710	.060
.300	.400	.880	.060
.400	.500	1.120	.070
.500	.600	1.270	.080
.600	.700	1.650	.080
.700	.800	1.630	.080
.800	.900	2.070	.100
1.000	1.000	2.540	.160

2Q SQUARE= .718

BLAND UCRL 18889 (1969) HBC

BEAM MOMENTUM= 1.215 GEV/C

COS THETA	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)	
-.820	.350	.120
-.740	.250	.060
-.660	.400	.090
-.550	.420	.070
-.460	.300	.070
-.350	.400	.100
-.260	.380	.070
-.160	.570	.100
-.070	.370	.070
-.010	.530	.100
.050	.500	.090
.120	.450	.070
.180	.770	.120
.250	.780	.120
.320	.730	.120
.390	.950	.100
.500	1.280	.150
.600	1.380	.130
.700	1.780	.180
.770	1.660	.190
.830	1.420	.250

2Q SQUARE= .725

PLUS POSSIBLE SYSTEMATIC ERROR OF +- 15PER CENT

ANDERSSON CERN (1969) CNTR

BEAM MOMENTUM= 1.372 GEV/C

COS THETA	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)	
-.860	.220	.090
-.790	.220	.040
-.710	.240	.050
-.650	.250	.050
-.590	.220	.050
-.520	.270	.050
-.440	.220	.050
-.360	.280	.050
-.290	.320	.050
-.240	.310	.100
-.180	.360	.070
-.140	.470	.070
-.450	.950	.130
-.520	1.000	.170
-.570	1.160	.150
-.630	1.360	.190
-.680	1.510	.200
-.730	1.650	.220
-.800	1.770	.160
-.870	1.740	.170

2Q SQUARE= .067

PLUS POSSIBLE SYSTEMATIC ERROR OF +- 15PER CENT

ANDERSSON CERN (1969) CNTR

BEAM MOMENTUM= 1.455 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)	
-1.000	-.950	.119	.038
-.950	-.900	.261	.056
-.900	-.850	.154	.043
-.850	-.800	.249	.054
-.800	-.750	.107	.036
-.750	-.700	.154	.043
-.700	-.650	.059	.027
-.650	-.600	.143	.041
-.600	-.550	.154	.043
-.550	-.500	.131	.039
-.500	-.450	.154	.043
-.450	-.400	.147	.041
-.400	-.350	.226	.052
-.350	-.300	.166	.044
-.300	-.250	.119	.038
-.250	-.200	.321	.062
-.200	-.150	.345	.064
-.150	-.100	.261	.056
-.100	-.050	.297	.059
-.050	0.	.309	.061
0.	.050	.475	.075
.050	.100	.439	.080
.100	.150	.475	.075
.150	.200	.737	.094
.200	.250	.511	.078
.250	.300	.594	.084
.300	.350	.772	.096
.350	.400	.701	.091
.400	.450	1.164	.118
.450	.500	.927	.105
.500	.550	1.010	.110
.550	.600	1.224	.121
.600	.650	1.414	.130
.650	.700	1.841	.148
.700	.750	2.138	.159
.750	.800	2.293	.165
.800	.850	2.412	.169
.850	.900	3.113	.192
.900	.950	3.778	.212

2Q SQUARE= .930

THE DATA HAVE BEEN RENORMALIZED(MULT.BY .99)

BETTINI PL 16 83(1965) HBC

BEAM MOMENTUM= 1.453 GEV/C

COS THETA	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)	
-.830	.210	.040
-.760	.200	.060
-.700	.140	.040
-.640	.150	.040
-.570	.160	.040
-.500	.200	.040
-.420	.210	.050
-.350	.300	.050
-.270	.300	.060
-.210	.320	.070
-.150	.410	.080
-.140	.480	.070
-.310	.670	.150
-.380	.720	.090
-.440	1.070	.090
-.550	1.170	.120
-.610	1.700	.180
-.660	1.550	.200
-.700	1.680	.220
-.770	1.790	.170
-.860	1.980	.190

2Q SQUARE= .928

PLUS POSSIBLE SYSTEMATIC ERROR OF +- 15PER CENT

ANDERSSON CERN (1969) CNTR

BEAM MOMENTUM= 1.970 GEV/C

COS THETA	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)	
-.940	.043	.021
-.850	.081	.019
-.750	.062	.017
-.650	.064	.017
-.550	.054	.016
-.450	.048	.016
-.350	.029	.014
-.250	.056	.021
-.150	.083	.028
-.050	.061	.023
.100	.120	.060
.250	.270	.040
.350	.320	.050
.450	.440	.060
.550	.490	.060
.650	1.040	.110
.750	1.470	.140
.850	1.840	.150
.888	2.750	.720
.895	2.410	.430
.903	2.080	.410
.920	1.940	.480
.935	2.080	.470
.949	2.620	.450
.962	2.620	.560
.973	3.940	.800
.983	4.360	1.120
.990	5.210	1.290

2Q SQUARE= 1.384

CULM PR 129 2743(1963) SPRK

K^+p elastic differential cross section

Figure 9 (Continued)

K⁺p elastic differential cross section

MOMENTUM TRANSFER BETWEEN BEAM AND SCATT BEAM

BEAM MOMENTUM= 1.087 GEV/C

MOM TRANSF	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
.093	12.601
.136	14.125
.179	14.125
.229	9.959
.284	10.569
.334	8.841
.385	9.349
.402	8.130
.439	9.349
.476	8.434
.532	6.097
.581	5.081
.643	4.979
.686	5.599
.717	4.573
.761	4.979
.835	4.675
.884	5.081
.934	4.166
.977	5.081
1.014	5.284
1.063	5.183
1.119	5.183

ZQ SQUARE= .618
PLUS POSSIBLE SYSTEMATIC ERROR OF +- 15PER CENT
ANDERSSON CERN (1969) CNTR

BEAM MOMENTUM= 1.215 GEV/C

MOM TRANSF	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
.123	12.307
.167	14.388
.217	15.428
.290	11.961
.362	11.094
.442	8.234
.493	6.327
.544	6.760
.594	6.674
.638	3.900
.689	4.334
.732	4.594
.776	3.207
.841	4.940
.913	3.294
.979	3.467
1.058	2.600
1.124	3.640
1.203	3.467
1.261	2.167
1.219	3.034

ZQ SQUARE= .725
PLUS POSSIBLE SYSTEMATIC ERROR OF +- 15PER CENT
ANDERSSON CERN (1969) CNTR

BEAM MOMENTUM= 1.170 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
.096	.206	17.281
.206	.275	11.521
.275	.344	10.881
.344	.412	8.686
.412	.481	9.509
.481	.687	5.577
.687	.756	3.860
.756	.825	3.200
.825	.893	3.017
.893	.962	3.292
.962	1.031	3.932
1.031	1.100	2.194
1.100	1.168	3.749
1.168	1.237	3.292
1.237	1.306	2.932
1.306	1.374	4.678

ZQ SQUARE= .687
COOK PR 129 2743(1963) SPRK

BEAM MOMENTUM= 1.372 GEV/C

MOM TRANSF	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
.112	12.881
.172	12.954
.232	12.076
.275	11.051
.318	9.954
.369	8.490
.412	7.977
.472	6.953
.738	3.586
1.013	2.635
1.065	2.269
1.107	2.342
1.168	2.049
1.236	1.610
1.305	1.976
1.365	1.610
1.417	1.820
1.468	1.767
1.537	1.610
1.597	1.610

ZQ SQUARE= .859
PLUS POSSIBLE SYSTEMATIC ERROR OF +- 15PER CENT
ANDERSSON CERN (1969) CNTR

BEAM MOMENTUM= 1.207 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
0.	0.	22.221
.072	.144	18.109
.144	.215	14.260
.215	.287	14.435
.287	.359	11.110
.359	.431	9.798
.431	.503	7.699
.503	.575	6.211
.575	.646	5.774
.646	.718	4.724
.718	.790	4.462
.790	.862	4.024
.862	.934	3.937
.934	1.005	3.762
1.005	1.077	3.237
1.077	1.149	3.062
1.149	1.221	3.587
1.221	1.293	3.324
1.293	1.365	4.287
1.365	1.436	4.987
1.436	1.436	4.987

ZQ SQUARE= .718
BLAND UCRL 18889 (1969) HBC

BEAM MOMENTUM= 1.453 GEV/C

MOM TRANSF	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
.130	13.401
.214	12.115
.279	11.370
.316	10.490
.362	8.798
.418	7.919
.501	6.903
.576	4.873
.641	4.535
.798	3.249
1.068	2.775
1.123	2.166
1.179	2.030
1.253	2.030
1.318	1.421
1.393	1.354
1.458	1.083
1.523	1.015
1.578	.948
1.634	1.354
1.699	1.421

ZQ SQUARE= .928
PLUS POSSIBLE SYSTEMATIC ERROR OF +- 15PER CENT
ANDERSSON CERN (1969) CNTR

K^+p elastic differential cross section

Figure 10. $\ln d\sigma/dt$ for K^+p elastic scattering above 1 GeV/c. The lines are the result of our least-squares fit to $d\sigma/dt = Ae^{Bt}$, using data in the interval $0 \leq -t \leq .6$ (GeV/c)². The fitted parameters are plotted in Fig. 11. So as not to mask possible structure, systematic errors have not been included even though they may have been given by the authors.

Some distributions given originally as $d\sigma/d\omega$ vs $\theta_{c.m.}$ were transformed to $d\sigma/dt$ vs t . Experiments which report numbers of events, rather than $d\sigma/dt$ (and do not give a factor which would enable us to convert to σ) have not been included.

K^+p elastic differential cross section

MOMENTUM TRANSFER BETWEEN BEAM AND SCATT BEAM

BEAM MOMENTUM= 1.455 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
0.	.047	25.521 1.431
.047	.093	21.027 1.299
.093	.140	16.292 1.143
.140	.186	15.489 1.115
.186	.233	14.446 1.077
.233	.279	12.439 .999
.279	.326	9.550 .875
.326	.372	8.266 .814
.372	.419	6.822 .740
.419	.465	6.260 .709
.465	.512	7.865 .794
.512	.558	4.735 .616
.558	.605	5.217 .647
.605	.651	4.013 .567
.651	.698	3.451 .526
.698	.744	4.976 .632
.744	.791	3.210 .508
.791	.837	3.611 .538
.837	.884	3.210 .508
.884	.930	2.087 .409
.930	.977	2.006 .401
.977	1.023	2.006 .401
1.023	1.070	1.766 .376
1.070	1.116	2.327 .432
1.116	1.163	2.167 .417
1.163	1.209	.803 .254
1.209	1.256	1.124 .300
1.256	1.302	1.525 .350
1.302	1.349	.963 .278
1.349	1.395	1.043 .289
1.395	1.442	.003 .266
1.442	1.488	1.043 .289
1.488	1.535	.963 .278
1.535	1.581	.401 .179
1.581	1.628	1.043 .289
1.628	1.674	.722 .241
1.674	1.721	1.685 .368
1.721	1.767	1.043 .289
1.767	1.814	1.766 .376
1.814	1.860	.803 .254

2Q SQUARE= .930

THE DATA HAVE BEEN RENORMALIZED(MULT.BY .99)

DETTINI PL 16 83(1965) HBC

BEAM MOMENTUM= 1.960 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
.010	.200	17.100 .071
.200	.400	7.482 .831
.400	.600	4.572 .623
.600	.800	2.910 .499
.800	1.000	1.143 .249
1.000	1.200	.727 .125
1.200	1.400	.357 .125
1.400	1.600	.470 .125
1.600	1.800	.208 .125
1.800	2.000	.166 .083
2.000	2.600	.104 .083
2.600	2.750	.146 .083

2Q SQUARE= 1.376

THE DATA HAVE BEEN RENORMALIZED(MULT.BY .91)

CHINOWSKY PR 1398141(1965) HBC

BEAM MOMENTUM= 1.970 GEV/C

MUM TRANSF	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
.014	23.644 5.854
.024	19.787 5.083
.037	17.881 3.631
.053	11.890 2.541
.071	11.890 2.042
.090	9.440 2.133
.111	8.804 2.178
.134	9.440 1.861
.159	10.937 1.951
.183	12.480 3.260
.208	8.350 .681
.346	6.671 .635
.485	4.720 .499
.623	2.224 .272
.761	1.997 .272
.900	1.452 .227
1.038	1.225 .182
1.246	.545 .272
1.454	.277 .104
1.592	.377 .127
1.731	.254 .095
1.869	.132 .064
2.008	.218 .073
2.146	.245 .073
2.284	.290 .077
2.423	.281 .077
2.561	.368 .086
2.686	.195 .095

2Q SQUARE= 1.384

COOK PR 129 2743(1963) SPRK

BEAM MOMENTUM= 2.970 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
.046	.115	14.650 1.510
.115	.229	8.980 .910
.229	.344	6.000 .650
.344	.459	3.550 .430
.459	.573	2.320 .310
.573	.688	1.450 .230
.688	.802	1.290 .210
.802	.917	.620 .130
.917	1.032	.480 .120
1.032	1.376	.190 .040
1.376	2.063	.040 .010
2.063	2.293	0.
2.293	4.471	.010 .010

2Q SQUARE= 2.293

DEBAISIEUX NC 43A 142(1966) HBC

BEAM MOMENTUM= 3.460 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
.027	.055	15.750 1.660
.055	.110	12.500 1.170
.110	.165	9.590 .940
.165	.219	7.620 .770
.219	.274	5.520 .590
.274	.329	5.380 .550
.329	.384	4.250 .490
.384	.439	2.990 .370
.439	.494	3.370 .410
.494	.549	2.530 .330
.549	.606	1.580 .180
.606	.623	1.120 .140
.623	.660	.460 .080
.660	1.097	.350 .070
1.097	2.058	.056 .010
2.058	5.487	.007 .002

2Q SQUARE= 2.744

DEBAERE NC 45A 885(1966) HBC

BEAM MOMENTUM= 3.550 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
.160	.200	9.200 .930
.200	.260	8.100 .280
.260	.320	6.460 .260
.320	.380	5.300 .240
.380	.440	4.100 .220
.440	.500	3.200 .200
.500	.560	2.320 .170
.560	.620	2.010 .170
.620	.680	1.690 .160
.680	.740	1.370 .150
.740	.800	1.030 .100
.800	.940	.630 .090
.940	1.040	.580 .100
1.040	1.140	.430 .090
1.140	1.340	.170 .040
1.340	1.740	.050 .020

2Q SQUARE= 2.827

PLUS POSSIBLE SYSTEMATIC ERROR OF +- 15PER CENT

BANAIGS NP 89 640(1969) SPRK

K^+p elastic differential cross section

Figure 10 (Continued)

K^+p elastic differential cross section

MOMENTUM TRANSFER BETWEEN BEAM AND SCATT BEAM

BEAM MOMENTUM= 4.970 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)	
.041	.083	14.230	1.780
.083	.166	9.640	1.140
.166	.249	6.840	.850
.249	.331	4.420	.590
.331	.414	2.720	.400
.414	.497	2.350	.390
.497	.580	1.250	.230
.580	.663	.880	.190
.663	.746	.520	.140
.746	.829	.610	.150
.829	1.036	.268	.064
1.036	8.287	.005	.001

2Q SQUARE= 4.143

DEBAERE NC 45A 885(1966) HBC

BEAM MOMENTUM= 6.800 GEV/C

MOM TRANSF	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)	
.251	4.810	.340
.321	2.930	.240
.399	2.110	.180
.481	1.460	.150
.569	.944	.117
.663	.671	.100
.762	.331	.072
.867	.266	.071
.977	.219	.057
1.091	.117	.049
1.193	.058	.042

2Q SQUARE= 5.849
PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7PER CENT

FOLEY PRL 11 503(1963) CNTR

BEAM MOMENTUM= 7.300 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)	
.100	.150	10.370	.540
.150	.200	8.110	.480
.200	.250	6.530	.430
.250	.300	4.970	.370
.300	.350	3.450	.310
.350	.400	3.080	.300
.400	.450	1.750	.220
.450	.500	1.750	.220
.500	.550	1.380	.200
.550	.600	.730	.140
.600	.650	.730	.140
.650	.700	.730	.140
.700	.750	.340	.100
.750	.800	.450	.110
.800	.850	.400	.110
.850	.900	.113	.057
.900	.950	.141	.063
.950	1.000	.085	.049
1.000	1.050	.170	.069
1.050	1.100	.086	.049
1.100	1.200	.028	.020
1.200	1.300	.056	.028
1.300	1.400	.028	.020
1.400	1.500	.042	.024

2Q SQUARE= 6.316
PLUS POSSIBLE SYSTEMATIC ERROR OF +- 9PER CENT

CHEN PL 288 615(1969) HBC

BEAM MOMENTUM= 9.800 GEV/C

MOM TRANSF	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)	
.049	14.740	1.540
.071	13.200	1.380
.098	10.150	1.080
.133	8.370	.970
.173	7.870	.940
.212	4.360	.780
.266	4.120	.290
.340	2.640	.210
.423	1.580	.140
.510	1.090	.120
.605	.716	.094
.707	.300	.062
.816	.270	.055
.931	.162	.048
1.051	.043	.027

2Q SQUARE= 8.455
PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7PER CENT

FOLEY PRL 11 503(1963) CNTR

BEAM MOMENTUM= 12.800 GEV/C

MOM TRANSF	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)	
.050	16.430	1.310
.083	12.760	.990
.120	8.470	.870
.166	7.210	.670
.224	5.050	.530
.277	3.800	.240
.291	3.620	.460
.354	2.250	.170
.441	1.430	.120
.531	.652	.085
.630	.521	.071
.736	.235	.046
.849	.112	.035
.971	.060	.025
1.099	.054	.019

2Q SQUARE= 11.465
PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7PER CENT

FOLEY PRL 11 503(1963) CNTR

BEAM MOMENTUM= 14.800 GEV/C

MOM TRANSF	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)	
.066	14.370	1.060
.111	9.610	.840
.161	7.810	.770
.222	5.030	.570
.290	3.230	.230
.300	3.630	.410
.371	2.000	.160
.389	1.750	.300
.462	1.230	.110
.558	.805	.086
.661	.451	.067
.774	.181	.045
.896	.153	.040
1.024	.115	.035
1.160	.069	.028

2Q SQUARE= 13.339
PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7PER CENT

FOLEY PRL 11 503(1963) CNTR

K^+p elastic differential cross section

Figure 10 (Continued)

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

K⁺p elastic differential cross section

Fitted parameters

P_{beam} (GeV/c)	$d\sigma/dt [t=0]$ mb/(GeV/c) ²	SLOPE (GeV/c) ⁻²	χ^2/N
1.087	17.11 ± 3.16	1.71 ± .39	.22
1.170	22.46 ± 2.70	2.31 ± .29	.64
1.207	22.96 ± .90	2.26 ± .14	.56
1.215	21.16 ± 3.89	2.05 ± .29	.72
1.372	17.04 ± 3.09	1.80 ± .34	.80
1.453	18.87 ± 3.39	2.10 ± .27	.98
1.455	25.58 ± .97	2.96 ± .14	.35
1.960	19.10 ± 1.58	2.94 ± .35	.80
1.970	15.29 ± 1.22	2.50 ± .29	1.47
2.970	19.62 ± 1.99	4.20 ± .34	.43
3.460	16.44 ± 1.09	3.76 ± .23	.99
3.550	19.09 ± 3.01	3.80 ± .14	.26
4.970	18.49 ± 1.89	4.89 ± .35	.47
6.800	16.23 ± 2.59	5.06 ± .38	.35
7.300	20.86 ± 2.26	5.37 ± .22	1.61
9.800	18.89 ± 1.78	5.74 ± .22	1.37
12.800	21.21 ± 1.89	6.29 ± .20	1.11
14.800	19.76 ± 1.83	6.00 ± .19	.95

Fig. 11. Coefficients from our least squares fit of the K⁺p elastic scattering differential cross section to the form

$$d\sigma/dt = \left. \frac{d\sigma}{dt} \right|_{t=0} e^{\text{"slope"} \cdot t}$$

over the interval $0 \leq -t \leq .6$ (GeV/c)². Systematic scaling errors in the data have been folded into the errors of $d\sigma/dt|_{t=0}$ given by the fit. The fitted values of the slopes and $d\sigma/dt$ at $t=0$ whenever given by the authors, are presented in the listing of the Data Tape. In the table, χ^2/N is the chi-square of the fit divided by the number of degrees of freedom.

K^+p elastic differential cross section

U BETWEEN BEAM AND SCATT TARG

BEAM MOMENTUM= .778 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
-.586	-.568	23.132 1.598
-.568	-.530	19.078 1.094
-.530	-.493	17.816 1.060
-.493	-.456	16.924 1.009
-.456	-.418	17.900 1.043
-.418	-.381	14.737 .925
-.381	-.344	16.318 .976
-.344	-.306	14.266 .892
-.306	-.269	14.922 .925
-.269	-.232	15.023 .925
-.232	-.194	15.343 .925
-.194	-.157	15.629 .942
-.157	-.119	15.629 .942
-.119	-.082	14.889 .908
-.082	-.045	16.554 .976
-.045	-.007	19.128 1.077
-.007	.030	16.033 .959
.030	.067	17.748 1.009
.067	.105	18.051 1.026
.105	.142	19.128 1.077

UZERO= .142

FOCARDI PL 248 314(1967) HBC
CUT ON U = -.007

BEAM MOMENTUM= .864 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
-.743	-.743	19.164 1.859
-.699	-.655	17.305 .858
-.655	-.611	15.160 .858
-.611	-.568	15.160 .858
-.568	-.524	14.874 .715
-.524	-.480	13.444 .715
-.480	-.436	12.299 .715
-.436	-.392	12.443 .715
-.392	-.348	13.158 .715
-.348	-.304	11.584 .715
-.304	-.260	12.156 .715
-.260	-.216	11.441 .715
-.216	-.172	11.012 .715
-.172	-.128	13.015 .715
-.128	-.084	12.443 .715
-.084	-.040	12.872 .715
-.040	.004	14.159 .715
.004	.048	15.732 .858
.048	.091	18.592 .858
.135	.135	22.883 1.716

UZERO= .135

BLAND UCRL 18889 (1969) HBC
CUT ON U = -.040

BEAM MOMENTUM= .969 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
-.917	-.917	19.487 2.045
-.864	-.812	16.240 .962
-.812	-.760	15.397 .962
-.760	-.708	11.187 .842
-.708	-.656	14.074 .962
-.656	-.603	12.631 .842
-.603	-.551	11.548 .842
-.551	-.499	9.142 .722
-.499	-.447	9.383 .722
-.447	-.394	9.503 .722
-.394	-.342	9.864 .842
-.342	-.290	9.744 .722
-.290	-.238	8.902 .722
-.238	-.185	8.902 .722
-.185	-.133	9.984 .842
-.133	-.081	7.022 .722
-.081	-.029	9.714 .842
-.029	.024	11.428 .842
.024	.076	13.232 .962
.128	.128	16.480 1.925

UZERO= .128

BLAND UCRL 18889 (1969) HBC
CUT ON U = -.081

BEAM MOMENTUM= .990 GEV/C

U	SIGMA +- DSIGMA (MICROBARNS/(GEV/C)**2)
-.031	4994.000 2233.000
-.020	7473.000 2491.000
-.009	5295.000 2001.000
.002	5726.000 2024.000
.013	4122.000 1683.000
.023	3465.000 1550.000
.034	5076.000 1919.000
.045	4350.000 1777.000
.056	6103.000 2034.000
.067	3658.000 1493.000
.078	7010.000 2024.000
.088	6497.000 1802.000
.099	7453.000 1757.000
.110	8290.000 1658.000
.121	5687.000 1376.000
.126	6210.000 1600.000

UZERO= .127

PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7PER CENT

CARROLL PRL 21 1282(1968) SPRK

BEAM MOMENTUM= 1.040 GEV/C

U	SIGMA +- DSIGMA (MICROBARNS/(GEV/C)**2)
-.069	4059.000 1224.000
-.057	5773.000 1260.000
-.046	4357.000 974.000
-.034	6154.000 1055.000
-.022	6281.000 1019.000
-.011	5780.000 950.000
.001	5261.000 889.000
.013	6386.000 985.000
.024	4588.000 852.000
.036	5713.000 952.000
.047	4283.000 795.000
.059	4551.000 936.000
.071	5603.000 845.000
.082	6325.000 831.000
.091	5651.000 999.000
.097	3704.000 726.000
.103	5424.000 880.000
.109	5987.000 855.000
.114	5567.000 647.000
.120	5257.000 628.000
.123	5570.000 580.000

UZERO= .123

PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7PER CENT

CARROLL PRL 21 1282(1968) SPRK

BEAM MOMENTUM= 1.110 GEV/C

U	SIGMA +- DSIGMA (MICROBARNS/(GEV/C)**2)
-.078	3967.000 887.000
-.065	5292.000 908.000
-.052	4631.000 761.000
-.040	3584.000 624.000
-.027	2519.000 776.000
-.014	3918.000 605.000
.001	3595.000 583.000
.011	3324.000 562.000
.024	4987.000 749.000
.037	3347.000 550.000
.049	4464.000 616.000
.062	4022.000 563.000
.075	4985.000 592.000
.084	5746.000 876.000
.091	4335.000 694.000
.097	4469.000 638.000
.103	4340.000 560.000
.110	4081.000 478.000
.116	4028.000 462.000
.119	4000.000 695.000

UZERO= .119

PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7PER CENT

CARROLL PRL 21 1282(1968) SPRK

K^+p elastic differential cross section

Figure 12. $\ln d\sigma/du$ for K^+p elastic scattering above 750 MeV/c. The straight lines are the results of our fit to $d\sigma/du = Ae^{B(u - u_0)}$. (u_0 corresponds to K^+ going backward ($\cos \theta = -1$) in the c.m.s. after scattering).

We have transformed data given as $d\sigma/d\omega$ to $d\sigma/du$ only when the experiments reported data close enough to $\cos \theta_{c.m.} = -1$. In the fit

we have used data in the region $-1 \leq \cos \theta_{c.m.} \leq -.6$; at higher momenta, where the range of measured data is much smaller, the fitted curve is drawn only as far as the measurements extend. The fitted parameters are plotted in Fig. 13.

The square symbols on the graphs represent the values of $d\sigma/du$ at $u = u_0$ obtained from our fits.

K^+ p elastic differential cross section

U BETWEEN BEAM AND SCATT TARG

BEAM MOMENTUM= 1.170 GEV/C

U	SIGMA +- DSIGMA (MICROBARN/(GEV/C)**2)
-.112	3666.000
-.098	5390.000
-.084	3766.000
-.071	2991.000
-.057	3010.000
-.043	3609.000
-.029	3229.000
-.016	3248.000
-.002	3327.000
.012	3832.000
.026	4457.000
.040	5029.000
.053	3723.000
.067	4143.000
.078	5204.000
.084	3630.000
.091	4049.000
.098	3537.000
.105	4650.000
.112	4464.000
.115	4360.000

UZERO= .115
PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7PER CENT

CARROLL PRL 21 1282(1968) SPRK

BEAM MOMENTUM= 1.170 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARN/(GEV/C)**2)
-1.163	-1.053	17.281
-1.053	-.984	11.521
-.984	-.915	10.881
-.915	-.847	8.686
-.847	-.778	9.509
-.778	-.572	5.577
-.572	-.503	3.840
-.503	-.434	3.200
-.434	-.366	3.017
-.366	-.297	3.292
-.297	-.228	3.432
-.228	-.159	2.194
-.159	-.091	3.749
-.091	-.022	3.292
-.022	.047	3.932
.047	.088	6.675

UZERO= .115

COOK PR 129 2743(1963) SHK
CUT ON U = -.159

BEAM MOMENTUM= 1.200 GEV/C

U	SIGMA +- DSIGMA (MICROBARN/(GEV/C)**2)
-.122	1074.000
-.108	2285.000
-.093	1497.000
-.079	1713.000
-.065	2411.000
-.050	2061.000
-.036	1958.000
-.022	2213.000
-.008	1924.000
.007	2865.000
.021	3165.000
.035	2046.000
.049	2833.000
.064	3008.000
.074	2047.000
.082	2883.000
.089	3027.000
.096	3333.000
.103	3652.000
.110	2976.000
.114	3560.000

UZERO= .114
PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7PER CENT

CARROLL PRL 21 1282(1968) SPRK

BEAM MOMENTUM= 1.207 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARN/(GEV/C)**2)
-1.323	-1.323	22.221
-1.251	-1.179	18.109
-1.179	-1.108	14.260
-1.108	-1.036	14.435
-1.036	-.964	11.110
-.964	-.892	9.798
-.892	-.820	7.699
-.820	-.748	6.211
-.748	-.677	5.774
-.677	-.605	4.724
-.605	-.533	4.462
-.533	-.461	4.024
-.461	-.389	3.937
-.389	-.318	3.762
-.318	-.246	3.237
-.246	-.174	3.062
-.174	-.102	3.587
-.102	-.030	3.324
-.030	.042	4.287
.042	.113	4.987
.113	.113	4.987

UZERO= .113

BLAND UCRL 18889 (1969) HBC
CUT ON U = -.174

BEAM MOMENTUM= 1.300 GEV/C

U	SIGMA +- DSIGMA (MICROBARN/(GEV/C)**2)
-.138	2105.000
-.122	1347.000
-.106	991.000
-.090	1671.000
-.074	1449.000
-.058	1028.000
-.042	1498.000
-.027	1276.000
-.011	1474.000
.005	1575.000
.021	1722.000
.037	1781.000
.053	1765.000
.065	1052.000
.073	1679.000
.081	1539.000
.089	1887.000
.097	1814.000
.105	1781.000
.109	1950.000

UZERO= .108
PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7PER CENT

CARROLL PRL 21 1282(1968) SPRK

BEAM MOMENTUM= 1.400 GEV/C

U	SIGMA +- DSIGMA (MICROBARN/(GEV/C)**2)
-.169	728.000
-.151	900.000
-.134	935.000
-.116	1654.000
-.098	1124.000
-.081	983.000
-.063	1013.000
-.046	870.000
-.028	1418.000
-.011	1423.000
.007	1199.000
.025	1406.000
.042	1357.000
.055	1502.000
.064	1487.000
.073	1899.000
.082	1547.000
.091	1678.000
.099	1591.000
.104	1650.000

UZERO= .104

PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7PER CENT

CARROLL PRL 21 1282(1968) SPRK

K^+p elastic differential cross section

Figure 12 (Continued)

K⁺p elastic differential cross section

U BETWEEN BEAM AND SCATT TARG

BEAM MOMENTUM= 1.455 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MICROBARN/(GEV/C)**2)	
-1.759	-1.713	25.521	1.431
-1.713	-1.666	21.027	1.299
-1.666	-1.620	16.292	1.143
-1.620	-1.573	15.489	1.115
-1.573	-1.527	14.446	1.077
-1.527	-1.480	12.439	.999
-1.480	-1.434	9.550	.875
-1.434	-1.387	8.266	.814
-1.387	-1.341	6.822	.740
-1.341	-1.294	6.260	.709
-1.294	-1.248	7.865	.794
-1.248	-1.201	4.735	.616
-1.201	-1.155	5.217	.647
-1.155	-1.108	4.013	.567
-1.108	-1.062	3.451	.526
-1.062	-1.015	4.976	.632
-1.015	-.969	3.210	.508
-.969	-.922	3.611	.538
-.922	-.876	3.210	.498
-.876	-.829	2.087	.409
-.829	-.783	2.006	.401
-.783	-.736	2.006	.401
-.736	-.689	1.766	.376
-.689	-.643	2.327	.432
-.643	-.596	2.167	.417
-.596	-.550	.803	.254
-.550	-.503	1.174	.300
-.503	-.457	1.525	.350
-.457	-.410	.963	.278
-.410	-.364	1.043	.289
-.364	-.317	.883	.266
-.317	-.271	1.043	.289
-.271	-.224	.963	.278
-.224	-.178	.401	.179
-.178	-.131	1.043	.289
-.131	-.085	.722	.241
-.085	-.038	1.685	.368
-.038	.008	1.043	.289
.008	.055	1.766	.376
.055	.101	.803	.254

UZERO= .101
THE DATA HAVE BEEN RENORMALIZED(MULT.BY .99)

BETTINI PL 16 83(1965) HBC
CUT ON U = -.271

BEAM MOMENTUM= 1.610 GEV/C

U	SIGMA +- DSIGMA (MICROBARN/(GEV/C)**2)	
-.214	536.000	103.000
-.193	493.000	82.000
-.172	442.000	77.000
-.151	418.000	73.000
-.129	516.000	59.000
-.108	598.000	61.000
-.087	516.000	62.000
-.065	546.000	58.000
-.044	539.000	60.000
-.023	522.000	53.000
-.001	716.000	59.000
.020	704.000	55.000
.036	513.000	59.000
.047	690.000	48.000
.057	658.000	65.000
.068	657.000	56.000
.079	852.000	57.000
.089	752.000	48.000
.095	829.000	50.000

UZERO= .095
PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7PER CENT

CARROLL PRL 21 1282(1968) SPRK

BEAM MOMENTUM= 1.710 GEV/C

U	SIGMA +- DSIGMA (MICROBARN/(GEV/C)**2)	
-.245	163.000	94.000
-.222	70.000	49.000
-.199	252.000	84.000
-.176	277.000	77.000
-.152	161.000	54.000
-.129	406.000	85.000
-.106	320.000	83.000
-.083	358.000	75.000
-.060	389.000	75.000
-.037	510.000	84.000
-.014	353.000	65.000
.010	558.000	77.000
.027	430.000	92.000
.038	521.000	97.000
.050	449.000	85.000
.062	422.000	75.000
.073	450.000	66.000
.085	517.000	67.000
.091	491.000	60.000

UZERO= .091
PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7PER CENT

CARROLL PRL 21 1282(1968) SPRK

BEAM MOMENTUM= 1.490 GEV/C

U	SIGMA +- DSIGMA (MICROBARN/(GEV/C)**2)	
-.179	755.000	308.000
-.160	1218.000	338.000
-.141	731.000	231.000
-.121	737.000	204.000
-.102	929.000	213.000
-.083	903.000	207.000
-.064	560.000	169.000
-.044	720.000	186.000
-.025	1185.000	228.000
-.006	911.000	194.000
.013	962.000	189.000
.032	963.000	176.000
.047	1312.000	257.000
.056	1217.000	234.000
.066	1182.000	219.000
.075	1380.000	215.000
.085	1530.000	194.000
.095	1357.000	172.000
.099	1460.000	164.000

UZERO= .100
PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7PER CENT
CARROLL PRL 21 1202(1960) SPRK

BEAM MOMENTUM= 1.670 GEV/C

U	SIGMA +- DSIGMA (MICROBARN/(GEV/C)**2)	
-.207	486.000	95.000
-.185	567.000	61.000
-.163	248.000	51.000
-.141	325.000	62.000
-.119	578.000	71.000
-.096	531.000	66.000
-.074	486.000	61.000
-.052	420.000	57.000
-.030	431.000	63.000
-.008	435.000	51.000
.015	542.000	53.000
.031	549.000	69.000
.048	709.000	70.000
.054	644.000	70.000
.065	558.000	60.000
.076	595.000	52.000
.087	625.000	49.000
.092	608.000	45.000

UZERO= .092
PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7PER CENT
CARROLL PRL 21 1282(1968) SPRK

BEAM MOMENTUM= 1.790 GEV/C

U	SIGMA +- DSIGMA (MICROBARN/(GEV/C)**2)	
-.241	225.000	44.000
-.216	299.000	51.000
-.192	218.000	36.000
-.168	284.000	34.000
-.143	347.000	44.000
-.119	288.000	42.000
-.095	354.000	33.000
-.070	295.000	37.000
-.046	451.000	37.000
-.021	439.000	32.000
.003	458.000	31.000
.021	482.000	43.000
.033	460.000	40.000
.046	528.000	41.000
.058	547.000	38.000
.070	511.000	31.000
.082	585.000	32.000
.088	580.000	32.000

UZERO= .088
PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7PER CENT
CARROLL PRL 21 1282(1968) SPRK

K^+p elastic differential cross section

Figure 12 (Continued)

K^+p elastic differential cross section

U BETWEEN BEAM AND SCATT TARG

BEAM MOMENTUM= 1.880 GEV/C

U	SIGMA +- DSIGMA (MICROBARN/(GEV/C)**2)
-.267	117.000 48.000
-.241	204.000 53.000
-.215	194.000 44.000
-.189	219.000 53.000
-.163	176.000 34.000
-.137	239.000 39.000
-.111	341.000 59.000
-.085	309.000 52.000
-.059	271.000 37.000
-.032	343.000 40.000
-.006	348.000 37.000
.013	375.000 46.000
.026	351.000 43.000
.039	358.000 40.000
.052	475.000 44.000
.065	455.000 38.000
.078	475.000 36.000
.091	490.000 36.000

UZERO= .085
PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7PER CENT
CARROLL PRL 21 1282(1968) SPRK

BEAM MOMENTUM= 2.000 GEV/C

U	SIGMA +- DSIGMA (MICROBARN/(GEV/C)**2)
-.288	52.000 17.000
-.258	118.000 21.000
-.229	101.000 18.000
-.200	107.000 16.000
-.170	142.000 18.000
-.141	157.000 18.000
-.112	167.000 22.000
-.082	226.000 19.000
-.053	237.000 19.000
-.023	216.000 17.000
-.002	236.000 23.000
.013	317.000 27.000
.028	354.000 26.000
.043	342.000 23.000
.057	389.000 22.000
.072	366.000 21.000
.081	405.000 21.000

UZERO= .081
PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7PER CENT
CARROLL PRL 21 1282(1968) SPRK

BEAM MOMENTUM= 2.200 GEV/C

U	SIGMA +- DSIGMA (MICROBARN/(GEV/C)**2)
-.290	32.000 12.000
-.258	63.000 16.000
-.226	95.000 20.000
-.194	83.000 21.000
-.163	91.000 26.000
-.131	113.000 29.000
-.099	126.000 15.000
-.067	196.000 22.000
-.036	205.000 17.000
-.012	194.000 23.000
.004	241.000 25.000
.020	266.000 25.000
.036	290.000 24.000
.052	304.000 21.000
.068	302.000 23.000
.076	373.000 23.000

UZERO= .076
PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7PER CENT
CARROLL PRL 21 1282(1968) SPRK

BEAM MOMENTUM= 2.330 GEV/C

U	SIGMA +- DSIGMA (MICROBARN/(GEV/C)**2)
-.318	40.000 12.000
-.284	36.000 10.000
-.250	69.000 13.000
-.216	91.000 16.000
-.182	86.000 14.000
-.148	124.000 18.000
-.114	122.000 15.000
-.080	136.000 18.000
-.046	174.000 15.000
-.021	196.000 18.000
-.004	194.000 16.000
.013	257.000 18.000
.030	271.000 17.000
.047	302.000 16.000
.064	331.000 16.000
.072	353.000 16.000

UZERO= .072
PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7PER CENT
CARROLL PRL 21 1282(1968) SPRK

BEAM MOMENTUM= 2.450 GEV/C

U	SIGMA +- DSIGMA (MICROBARN/(GEV/C)**2)
-.349	44.000 20.000
-.312	27.000 13.000
-.276	68.000 19.000
-.239	27.000 10.000
-.203	76.000 20.000
-.167	68.000 19.000
-.130	100.000 23.000
-.094	102.000 18.000
-.058	88.000 18.000
-.021	145.000 19.000
.015	231.000 24.000
.052	251.000 22.000
.070	305.000 33.000

UZERO= .070
PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7PER CENT
CARROLL PRL 21 1282(1968) SPRK

BEAM MOMENTUM= 2.760 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MICROBARN/(GEV/C)**2)
-.230	-.272	35.000 18.000
-.188	-.230	52.000 22.000
-.146	-.188	60.000 26.000
-.104	-.146	86.000 30.000
-.062	-.104	95.000 31.000
-.020	-.062	155.000 39.000
.023	-.020	146.000 40.000
.065	.023	241.000 53.000
.065	.065	261.000 48.000

UZERO= .063
ABRAMS PRL 21 1407(1968) HBC

K^+p elastic differential cross section

Figure 12 (Continued)

K⁺p elastic differential cross section

Figure 12 (Continued)

U BETWEEN BEAM AND SCATT TARG

BEAM MOMENTUM= 3.550 GEV/C

BEAM MOMENTUM= 5.200 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MICROBARN/(GEV/C)**2)
-1.790	-1.190	7.400 3.500
-1.180	-.780	9.600 4.200
-.780	-.480	8.000 3.500
-.480	-.280	10.200 4.200
-.280	-.180	19.300 6.900
-.180	-.080	20.200 7.700
-.080	-.023	33.100 12.900
-.023	.047	47.400 13.190

UZERO= .052
PLUS POSSIBLE SYSTEMATIC ERROR OF +- 15PER CENT

BAKAFS NP 89 640(1969) SPRK
CUT ON U = -1.019

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MICROBARN/(GEV/C)**2)
-.850	-.750	0. 4.700
-.750	-.650	0. 1.300
-.650	-.550	1.600 1.400
-.550	-.450	3.900 1.800
-.450	-.350	4.800 1.600
-.350	-.250	1.100 1.200
-.250	-.150	3.400 1.100
-.150	-.100	12.100 2.800
-.100	-.060	9.500 3.000
-.060	-.020	14.000 4.000
-.020	.020	14.600 5.700

UZERO= .037
PLUS POSSIBLE SYSTEMATIC ERROR OF +- 15PER CENT

BAKER PL 288 291(1968) SPRK+CNTR

BEAM MOMENTUM= 6.900 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MICROBARN/(GEV/C)**2)
-1.100	-.900	0. .400
-.900	-.700	.300 .500
-.700	-.500	.700 .500
-.500	-.300	1.400 .600
-.300	-.100	2.100 .700
-.100	0.	2.500 1.100

UZERO= .029
PLUS POSSIBLE SYSTEMATIC ERROR OF +- 15PER CENT

BAKER PL 288 291(1968) SPRK+CNTR

K⁺p elastic differential cross section

Fitted parameters

P_{beam} (GeV/c)	$d\sigma/du [u = u_0]$ $\mu b/(\text{GeV}/c)^2$	SLOPE $(\text{GeV}/c)^{-2}$	χ^2/N
.778	19735.63 ± 1149.91	1.45 ± .69	1.32
.864	22785.97 1303.04	3.17 .54	.07
.969	16556.53 1496.37	2.87 .73	.90
.990	6883.64 958.24	2.01 1.64	.12
1.040	5535.09 478.58	-.05 .60	.00
1.110	4337.31 377.97	.34 .58	.45
1.170	5586.95 1243.84	2.23 1.34	.49
1.170	4364.68 362.13	1.06 .45	.74
1.200	3381.74 283.85	2.99 .50	1.02
1.207	5058.89 437.85	1.63 .60	1.15
1.300	1804.56 149.59	1.24 .43	1.03
1.400	1652.73 131.74	1.96 .36	1.19
1.455	1523.22 297.05	1.64 1.00	1.45
1.490	1396.74 127.45	2.52 .57	.71
1.610	768.90 58.63	1.76 .26	1.38
1.670	624.12 49.64	1.75 .33	1.79
1.710	520.90 49.05	2.17 .55	.93
1.790	577.04 43.31	2.74 .24	.81
1.880	480.81 38.23	3.16 .35	.64
2.000	404.25 30.43	4.30 .24	1.32
2.200	363.54 28.25	5.41 .35	.65
2.330	342.90 25.56	5.42 .27	.75
2.450	280.05 25.78	5.55 .47	1.43
2.760	258.40 34.19	6.14 1.08	2.10
3.550	35.82 9.01	1.78 .46	1.69
5.200	15.68 4.00	3.32 .80	.13
6.900	3.32 1.50	2.21 1.38	.19

Figure 13. Coefficients from our least squares fit of the backward K⁺p elastic differential cross section to the form

$$d\sigma/du = \left. \frac{d\sigma}{du} \right|_{u=u_0} \cdot e^{\text{"slope"} \cdot (u - u_0)}$$

In the fit we have used only data corresponding to $-1 \leq \cos \theta_{\text{c.m.}} \leq -.6$. Systematic scaling errors in the data have been folded into the errors of

$$\left. \frac{d\sigma}{du} \right|_{u=u_0}$$

given by the fit.

In the table, χ^2/N is the chi square of the fit divided by the number of degrees of freedom.

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

K^+_p Elastic Polarization

The sign of the published polarization data has been reversed where necessary to conform to the Basel convention.*

*Proceedings of the International Symposium on Polarization Phenomena of Nucleons, Basel, Switzerland, 1960, Helv. Phys. Acta, Suppl. No. 6, 436 (1961).

Polarization in K^+p elastic scattering

DISTRIBUTION IN COS THETA OF THE SCATT BEAM WITH RESPECT TO THE BEAM DIRECTION IN THE C.M. SYSTEM

BEAM MOMENTUM= .865 GEV/C

COS THETA	POLARIZ+-DPOL	
-.810	.340	.245
-.720	.150	.152
-.640	.320	.186
-.570	.470	.193
-.500	.450	.183
-.430	.530	.206
-.360	.340	.168
-.290	.790	.216
-.210	.100	.141
-.110	.700	.191
.030	.450	.183
.060	.770	.214
.140	.850	.317
.200	.500	.261
.240	.490	.308
.330	.940	.213
.420	.490	.185
.480	.520	.281
.530	.850	.212
.590	.490	.251
.640	.230	.282
.700	.700	.159
.790	.610	.257

ANDERSSON 69.....CERN

BEAM MOMENTUM= .971 GEV/C

COS THETA	POLARIZ+-DPOL	
-.840	.090	.250
-.780	-.130	.161
-.720	.170	.172
-.660	.370	.160
-.600	.310	.157
-.530	.350	.159
-.450	.230	.134
-.380	.360	.169
-.310	.490	.149
-.230	.430	.163
-.150	.340	.140
-.050	.620	.168
.030	.930	.269
.120	.640	.195
.200	.630	.177
.240	.850	.182
.400	.630	.145
.500	.770	.159
.650	.760	.152
.740	.710	.139
.830	.820	.261

ANDERSSON 69.....CERN

BEAM MOMENTUM= 1.215 GEV/C

COS THETA	POLARIZ+-DPOL	
-.820	.040	.190
-.740	.010	.130
-.660	.120	.141
-.550	.230	.106
-.460	.090	.141
-.350	.350	.140
-.260	.480	.131
-.160	.480	.115
-.070	.750	.164
-.010	.720	.147
.050	.630	.130
.120	.820	.159
.180	.540	.114
.250	.750	.138
.320	.680	.136
.390	.730	.125
.500	.600	.108
.600	.710	.114
.700	.650	.110
.770	.600	.108
.830	.540	.121

ANDERSSON 69.....CERN

BEAM MOMENTUM= .910 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA	
-.940	-.766	-.700	.930
-.766	-.174	-.550	.930
-.174	.342	.740	.450
.342	.766	.800	.800
HIRSCH	64.....PR	135 8191	

BEAM MOMENTUM= 1.087 GEV/C

COS THETA	POLARIZ+-DPOL	
-.810	-.130	.161
-.720	.280	.146
-.640	.640	.195
-.580	.130	.191
-.510	.070	.220
-.430	.410	.162
-.350	.420	.172
-.290	.670	.242
-.190	.790	.191
-.110	.850	.221
-.040	.740	.195
.060	1.210	.589
.140	.820	.380
.230	.620	.276
.290	.900	.275
.350	.900	.168
.410	.690	.158
.460	.900	.233
.540	.680	.136
.630	.550	.122
.710	.600	.127
.780	.710	.146
.850	.750	.196

ANDERSSON 69.....CERN

BEAM MOMENTUM= 1.370 GEV/C

COS THETA	POLARIZ+-DPOL	
-.895	-.279	.268
-.820	.464	.346
-.760	.210	.274
-.690	.040	.300
-.600	.805	.280
-.485	.724	.173
-.415	.790	.250
-.360	-.007	.304
-.305	.792	.151
-.225	.669	.140
-.125	.627	.102
-.060	.768	.197
.020	.776	.089
.115	.763	.090
.230	.663	.090
.350	.727	.059
.430	.731	.088
.490	.765	.075
.580	.717	.075
.650	.590	.100
.695	.672	.070
.745	.592	.095
.790	.673	.060
.830	.698	.168
.870	.548	.090

ASBURY 69.....PRL 23 194

Polarization in K^+p elastic scattering

Figure 14. Proton polarization in K^+p elastic scattering. The sign of published polarization data has been reversed where necessary so as to conform to the Basel convention; the positive polarization direction is along the normal

$$\vec{n} = \vec{p}_{K-in.} \times \vec{p}_{K-out.}$$

Data have been transformed to $\cos \theta_{c.m.}$ when given in another variable.

Polarization in K^+p elastic scattering

DISTRIBUTION IN COS THETA OF THE SCATT BEAM WITH RESPECT TO THE BEAM DIRECTION IN THE C.M. SYSTEM

BEAM MOMENTUM= 1.372 GEV/C

COS THETA	POLARIZ+-DPOL
-.860	-.450 .475
-.790	-.170 .231
-.710	-.060 .130
-.650	-.040 .240
-.590	.160 .251
-.520	.390 .208
-.440	.230 .233
-.360	.440 .211
-.290	.620 .248
-.210	.370 .208
-.130	.920 .303
-.060	.710 .263
.200	.720 .610
.280	.620 .160
.370	.470 .135
.450	.700 .159
.520	.480 .123
.600	.680 .130
.680	.550 .146
.730	.480 .108
.800	.560 .109
.870	.420 .127

ANDERSSON 69.....CERN

BEAM MOMENTUM= 1.450 GEV/C

COS THETA	POLARIZ+-DPOL
-.895	.192 .256
-.815	.319 .365
-.710	.392 .166
-.590	.429 .203
-.510	.542 .201
-.330	.750 .161
-.247	.649 .145
-.170	.570 .170
-.080	.565 .134
.018	.598 .121
.105	.574 .085
.197	.660 .074
.326	.694 .071
.420	.710 .080
.475	.660 .100
.575	.752 .093
.674	.628 .062
.735	.536 .071
.777	.679 .050
.815	.552 .102
.860	.549 .099
.880	.172 .164

ASBURY 69.....PRL 23 194

BEAM MOMENTUM= 1.453 GEV/C

COS THETA	POLARIZ+-DPOL
-.830	.140 .441
-.760	.340 .373
-.700	-.240 .322
-.640	-.030 .280
-.570	.840 .325
-.500	.460 .250
-.420	.100 .900
-.350	.400 .199
-.270	.690 .208
-.180	.600 .166
-.100	.600 .219
-.020	.500 .654
.070	.340 .334
.140	1.070 .468
.190	.530 .536
.250	.960 .388
.310	.820 .302
.300	.700 .1F2
.460	.790 .143
.550	.480 .115
.610	.600 .120
.700	.550 .102
.770	.580 .148
.820	.590 .107
.890	.450 .129

ANDERSSON 69.....CERN

BEAM MOMENTUM= 1.710 GEV/C

COS THETA	POLARIZ+-DPOL
-.900	-.164 .800
-.830	-.151 .179
-.690	.362 .190
-.560	.589 .270
-.380	.764 .206
-.240	.570 .115
-.140	.495 .095
-.040	.626 .080
.050	.596 .080
.070	.658 .080
.150	.641 .118
.100	.440 .111
.320	.564 .120
.390	.625 .059
.510	.686 .081
.550	.674 .045
.615	.686 .073
.660	.652 .036
.720	.645 .058
.760	.660 .034
.800	.684 .057
.840	.576 .030
.880	.570 .070
.910	.555 .112

ASBURY 69.....PRL 23 194

BEAM MOMENTUM= 1.890 GEV/C

COS THETA	POLARIZ+-DPOL
-.915	-.080 .650
-.779	-.040 .400
-.730	-.360 .385
-.675	.560 .500
-.560	.450 .520
-.400	.350 .400
-.305	.910 .500
-.180	.425 .186
-.085	.326 .135
.015	.384 .092
.125	.472 .099
.220	.396 .076
.340	.387 .087
.400	.512 .086
.520	.552 .044
.600	.585 .045
.637	.593 .033
.695	.520 .066
.742	.563 .033
.790	.561 .044
.830	.570 .031
.875	.505 .056
.900	.410 .120

ASBURY 69.....PRL 23 194

BEAM MOMENTUM= 2.480 GEV/C

COS THETA	POLARIZ+-DPOL
.310	.430 .251
.370	.280 .181
.440	.280 .191
.500	.670 .125
.570	.480 .113
.610	.400 .132
.650	.490 .084
.710	.450 .083
.760	.500 .065
.810	.640 .068
.870	.510 .056
.900	.530 .066
.930	.490 .132

ANDERSSON 69.....PL 288, 611

Polarization in K^+p elastic scattering

Figure 14 (Continued)

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

K_p^+ Elastic Phase Shift Analyses

K⁺P PARTIAL WAVE AMPLITUDES

LEVI-SETTI

74

ANDERSSON, ET AL

- SOLUTION I
- SOLUTION II
- SOLUTION III
- DCS DATA
- DCS+POL DATA

Figure 15a. Argand plots for K⁺p elastic scattering phase-shift results presented to the XVI International Conference on High Energy Physics at Lund, Sweden, June 1969. The S₁₁, P₁₁, and P₁₃ diagrams were presented by R. Levi-Setti as Rapporteur; the labels refer to the following reports to the conference: CERN-Holland—S. Andersson et al.; ANL-MD (Argonne and Univ. of Maryland)—

F. Asbury et al; LMO—Lea, Martin and Odes; BGRT—Bologna, Glasgow, Roma, and Trieste collaboration. The D₁₃ and D₁₅ diagrams are from Andersson et al.; Solution II corresponds to the CERN-Holland solutions on the other three diagrams.

K^+p P_{13} PARTIAL WAVE

CERN - HOLL I

CERN - HOLL II

ANL - MD I

ANL - MD II

Figure 15b. K^+p elastic P_{13} partial wave Argand and "speed" plots. The top two sets correspond to CERN-HOLLAND solutions I and II (ref. 54); the bottom two sets to ANL-MD solutions I and II (ref. 90). The speed is the distance between two points on the Argand plot divided by the corresponding difference in the energies. For the significance of the speed plots, see the text.

Figure 15c. The K^+p total cross section together with the contributions to the total cross section from different partial waves (from Levi-Setti's rapporteur talk at Lund, 1969).

K^+p Inelastic Differential Cross Sections

We have plotted here only those distributions for which there were more than about two experiments.

Also each plot is only for one particular decay mode of a resonance. Background problems make the combining of more than one decay mode on a plot rather risky.

Differential cross section for $K^+p \rightarrow K^0 \Delta^{++}$

MOMENTUM TRANSFER BETWEEN BEAM AND K^0

BEAM MOMENTUM= .860 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (NUMBER OF EVENTS)	
.104	.123	36.000	6.000
.123	.141	52.000	7.211
.141	.160	76.000	8.718
.160	.178	99.000	9.950
.178	.197	108.000	10.392
.197	.216	113.000	10.630
.216	.234	125.000	11.180
.234	.253	118.000	10.863
.253	.271	101.000	10.050
.271	.290	72.000	8.485

BLAND 68.....UCRL 18131

BEAM MOMENTUM= .960 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (NUMBER OF EVENTS)	
.059	.102	39.000	6.245
.102	.144	85.000	9.220
.144	.187	119.000	10.909
.187	.230	145.000	12.042
.230	.273	171.000	13.077
.273	.315	181.000	13.454
.315	.358	158.000	12.570
.358	.401	178.000	13.342
.401	.443	151.000	12.288
.443	.486	93.000	9.644

BLAND 68.....UCRL 18131

BEAM MOMENTUM= 1.140 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)	
.033	.109	1.120	.659
.109	.185	6.986	2.307
.185	.262	9.556	2.307
.262	.338	3.954	1.318
.338	.414	5.272	1.648
.414	.490	4.745	1.648
.490	.567	3.361	1.318
.567	.643	4.943	1.648
.643	.719	2.373	.989
.719	.795	1.318	.659

BOLDT 64.....PR 133 8220

BEAM MOMENTUM= 1.360 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (NUMBER OF EVENTS)	
.020	.135	5.000	2.236
.135	.250	17.000	4.123
.250	.365	19.000	4.359
.365	.480	37.000	6.083
.480	.595	28.000	5.292
.595	.709	45.000	6.708
.709	.824	51.000	7.141
.824	.939	50.000	7.071
.939	1.054	66.000	8.124
1.054	1.169	44.000	6.633

BLAND 68.....UCRL 18131

BEAM MOMENTUM= 1.200 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (NUMBER OF EVENTS)	
.028	.115	20.000	4.472
.115	.202	43.000	6.557
.202	.289	76.000	8.718
.289	.376	74.000	8.602
.376	.463	83.000	9.110
.463	.550	103.000	10.149
.550	.636	120.000	10.954
.636	.723	106.000	10.296
.723	.810	130.000	11.402
.810	.897	73.000	8.544

BLAND 68.....UCRL 18131

BEAM MOMENTUM= 1.455 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)	
.017	.148	2.619	.259
.148	.280	3.338	.293
.280	.411	3.158	.285
.411	.543	2.182	.237
.543	.674	1.874	.219
.674	.806	1.232	.178
.806	.937	1.309	.183
.937	1.069	.822	.145
1.069	1.200	.770	.141
1.200	1.332	.385	.099

BETTINI 65.....PL 16 83

Differential cross section for $K^+p \rightarrow K^0\Delta^{++}$

Figure 16. The differential cross section for the reaction $K^+p \rightarrow K^0\Delta^{++}(1238)$. The $d\sigma/dt$ scale is in arbitrary units because experiments which report number of events rather than $d\sigma/dt$ (and

do not give a factor which would enable us to convert to σ) have also been included. The distributions given originally as $d\sigma/d\omega$ vs $\cos \theta_{c.m.}$ were transformed to $d\sigma/dt$ vs t [taking $m(\Delta) = 1.22$ GeV].

Differential cross section for $K^+p \rightarrow K^0\Delta^{++}$

MOMENTUM TRANSFER BETWEEN BEAM AND K0

BEAM MOMENTUM= 1.960 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
.008	.119	2.425 .369
.119	.229	2.583 .369
.229	.339	2.108 .369
.339	.449	3.268 .369
.449	.560	1.950 .369
.560	.780	.791 .105
.780	1.111	.474 .079
1.111	1.662	.158 .047
1.662	2.213	.158 .047

GOLDHABER 66.....PR 142 913

BEAM MOMENTUM= 2.972 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
.003	.104	1.347 .389
.104	.205	1.908 .463
.205	.306	1.122 .355
.306	.406	1.122 .355
.406	.507	.673 .275
.507	.608	.561 .251
.608	.709	.112 .112
.709	.810	.337 .194
.810	.910	.112 .112
.910	1.011	.112 .112
1.011	1.112	0. .112
1.112	1.213	.112 .112
1.213	1.314	.112 .112
1.314	1.414	0. .112
1.414	1.515	.112 .112
1.515	1.918	0. .112
1.918	2.019	.112 .112
2.019	3.128	0. .112
3.128	3.228	.112 .112
3.228	3.732	0. .112
3.732	3.833	.112 .112
3.833	4.035	0. .112

SALLSTROM 67.....NC 49A 348

BEAM MOMENTUM= 4.600 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (NUMBER OF EVENTS)
0.	.200	50.000 7.000
.200	.400	52.000 7.000
.400	.600	17.000 4.000
.600	1.000	5.500 1.500

FU 68.....UCRL 18417

BEAM MOMENTUM= 2.965 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
.003	.104	1.032 .235
.104	.204	1.505 .296
.204	.305	1.505 .296
.305	.405	1.290 .269
.405	.506	.602 .172
.506	.606	.473 .150
.606	.707	.215 .099
.707	.807	.215 .099
.807	.908	.086 .061
.908	1.008	.172 .088
1.008	1.108	.086 .061
1.108	1.209	0. .043
1.209	1.309	0. .043
1.309	1.410	0. .043
1.410	1.510	0. .043
1.510	1.611	0. .043
1.611	1.711	0. .043
1.711	1.812	.043 .043
1.812	1.912	0. .043
1.912	2.013	.043 .043
2.013	2.113	0. .043
2.113	2.214	.043 .043
2.214	2.314	.086 .061
2.314	2.415	.043 .043
2.415	2.515	0. .043
2.515	2.615	0. .043
2.615	2.716	.043 .043
2.716	2.816	0. .043
2.816	2.917	0. .043
2.917	3.017	0. .043
3.017	3.118	.043 .043
3.118	3.218	.043 .043
3.218	3.319	0. .043
3.319	3.419	.043 .043
3.419	3.520	0. .043
3.520	3.620	.043 .043
3.620	3.721	0. .043
3.721	3.821	0. .086
3.821	3.922	.086 .061
3.922	4.022	0. .061

FERRO-LUZZ 65.....NC 36 1101

BEAM MOMENTUM= 12.700 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (NUMBER OF EVENTS)
.015	.115	60.000 12.000
.115	.215	23.000 9.000
.215	.415	11.000 2.500
.415	.715	4.000 2.000

BERLINGHIE 68.....NP 88 333

Differential cross section for $K^+p \rightarrow K^0\Delta^{++}$

Figure 16 (Continued)

Differential cross section for $K^+ p \rightarrow p K^* (890)^+ \rightarrow K^0 \pi^+$

MOMENTUM TRANSFER BETWEEN BEAM AND $K^*(890)^+$

BEAM MOMENTUM= 1.200 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (NUMBER OF EVENTS)
.114	.169	56.000 7.483
.169	.223	42.000 6.481
.223	.278	57.000 7.550
.278	.333	49.000 7.000
.333	.387	41.000 6.403
.387	.442	36.000 6.000
.442	.497	43.000 6.557
.497	.552	31.000 5.568
.552	.606	27.000 5.196
.606	.661	12.000 3.464

BLAND 68.....UCRL 18131

BEAM MOMENTUM= 1.360 GEV/C

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (NUMBER OF EVENTS)
.071	.162	52.000 7.211
.162	.253	43.000 6.557
.253	.345	37.000 6.083
.345	.436	28.000 5.292
.436	.527	31.000 5.568
.527	.618	23.000 4.796
.618	.709	33.000 5.745
.709	.800	10.000 3.162
.800	.892	10.000 3.162
.892	.983	8.000 2.828

BLAND 68.....UCRL 18131

BEAM MOMENTUM= 1.960 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
.026	.130	3.433 .554
.130	.234	2.769 .443
.234	.337	1.163 .277
.337	.441	1.052 .277
.441	.649	.720 .166
.649	1.064	.343 .166
1.064	1.583	.127 .111
1.583	2.103	.188 .055

GOLDHABER 66.....PR 142 913

BEAM MOMENTUM= 2.650 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
.013	.181	3.285 .720
.181	.349	1.753 .526
.349	.518	1.990 .561
.518	.686	.363 .239
.686	.854	.379 .245
.854	1.023	0. .458
1.023	1.191	0. .269
1.191	2.369	0. .158
2.369	2.537	0. .158
2.537	3.042	0. .316
3.042	3.211	.316 .223
3.211	3.379	.316 .223

NEWMAN 67.....PR 158 1310

BEAM MOMENTUM= 2.965 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
.010	.109	1.442 .290
.109	.208	1.574 .306
.208	.307	1.455 .279
.307	.405	.874 .214
.405	.504	.568 .168
.504	.603	.393 .137
.603	.702	.262 .110
.702	.801	.175 .089
.801	.899	.087 .062
.899	.998	.131 .077
.998	1.097	.044 .044
1.097	1.196	0. .062
1.196	1.295	.087 .062
1.295	1.394	.131 .077
1.394	1.492	.087 .062
1.492	1.591	.044 .044
1.591	1.690	.087 .062
1.690	1.789	0. .062
1.789	1.888	0. .044
1.888	1.987	.087 .062
1.987	2.085	.044 .044
2.085	2.184	.044 .044
2.184	2.283	0. .044
2.283	2.382	.044 .044
2.382	2.481	0. .044
2.481	2.579	0. .044
2.579	2.678	.044 .044
2.678	2.777	0. .062
2.777	2.876	0. .044
2.876	2.975	0. .044
2.975	3.074	.044 .044
3.074	3.172	0. .062
3.172	3.271	.087 .062
3.271	3.370	.044 .044
3.370	3.469	.044 .044
3.469	3.568	0. .062
3.568	3.667	.087 .062
3.667	3.765	.044 .044
3.765	3.864	.044 .044
3.864	3.963	0. .044

FERRO-LUZZ 65.....NC 36 1101

BEAM MOMENTUM= 2.972 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
.020	.120	5.000 1.000
.120	.220	2.200 .500
.220	.320	1.500 .500
.320	.420	1.200 .300
.420	.520	.750 .200
.520	.600	.400 .150
.600	.700	.150 .090
.700	.800	.340 .190
.800	.900	.150 .090
.900	1.000	.080 .060

SALLSTROM 68.....AF 37. 468

BEAM MOMENTUM= 4.600 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (NUMBER OF EVENTS)
0.	.100	20.000 5.000
.100	.300	28.000 2.000
.300	.500	15.000 3.000
.500	.800	5.600 2.000
.800	1.000	2.000 1.000

FU 68.....UCRL 18417

BEAM MOMENTUM= 12.700 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (NUMBER OF EVENTS)
.070	.170	12.000 4.000
.170	.270	17.000 3.000
.270	.370	9.000 3.000
.370	.470	8.000 3.000
.470	.870	2.500 .700

BERLINGHIE 68.....NP 88 333

Figure 17. Differential cross section for the reaction

The $d\sigma/dt$ scale is in arbitrary units because experiments which re-

port number of events rather than $d\sigma/dt$ (and do not give a factor which would enable us to convert to σ) have also been included.

The distributions given originally as $d\sigma/d\omega$ vs $\cos \theta_{c.m.}$ were transformed to $d\sigma/dt$ vs t [taking $m(K^*) = .89$ GeV].

Differential cross section for $K^+ p \rightarrow K^*(890)^0 \Delta^{++}$

$\searrow K^+ \pi^- \quad \swarrow p \pi^+$

MOMENTUM TRANSFER BETWEEN BEAM AND $K^*(890)^0$

BEAM MOMENTUM= 2.650 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
.131	.259	4.523 .574
.259	.386	2.626 .438
.386	.514	1.240 .301
.514	.641	.511 .193
.641	.768	.365 .163
.768	.896	.365 .163
.896	1.023	.073 .073
1.023	1.151	.219 .126
1.151	1.278	.219 .126
1.278	1.406	.146 .103
1.406	1.533	.073 .073
1.533	1.661	0.
1.661	1.788	0.
1.788	1.916	0.
1.916	2.043	.073 .073
2.043	2.171	.073 .073
2.171	2.298	.219 .126
2.298	2.426	0.
2.426	2.553	.073 .073
2.553	2.680	.146 .103

NEWMAN 67.....PR 158 1310

BEAM MOMENTUM= 3.460 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
.135	.084	5.726 .716
.187	.135	3.757 .580
.239	.187	3.131 .529
.290	.239	1.789 .400
.342	.290	.716 .253
.393	.342	.537 .219
.445	.393	.537 .219
.497	.445	.179 .127
.548	.497	.447 .200
.600	.548	0.
.651	.600	.358 .179
.703	.651	.089 .089
.755	.703	.089 .089
.806	.755	.179 .127
.858	.806	.268 .155
.910	.858	.268 .155
.961	.910	.089 .089
1.013	.961	.089 .089

GEORGE 67.....NC 49A 9

BEAM MOMENTUM= 12.700 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (NUMBER OF EVENTS)
.060	.080	30.000 5.000
.080	.100	15.000 4.000
.100	.140	6.000 2.000
.140	.240	3.200 1.000
.240	.360	1.600 .500
.560	1.060	.110 .090

BERLINGHIE 68.....NP 88 333

BEAM MOMENTUM= 2.970 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
.107	.187	5.935 .626
.187	.266	3.363 .471
.266	.346	1.846 .349
.346	.425	.989 .255
.425	.505	.725 .219
.505	.584	.462 .174
.584	.664	.396 .162
.664	.743	.198 .114
.743	.823	.528 .187
.823	.903	.132 .093
.903	.982	.264 .132
.982	1.062	.132 .093
1.062	1.141	.132 .093
1.141	1.221	0.
1.221	1.300	.132 .093
1.300	1.380	.132 .093
1.380	1.459	0.
1.459	1.539	.264 .132
1.539	1.619	.198 .114
1.619	1.698	.066 .066
1.698	1.778	0.
1.778	1.857	0.
1.857	1.937	0.
1.937	2.016	.066 .066
2.016	2.096	0.
2.096	2.176	0.
2.176	2.255	.132 .093
2.255	2.335	.066 .066
2.335	2.414	.066 .066
2.414	2.494	.066 .066
2.494	2.573	0.
2.573	2.653	.066 .066
2.653	2.732	.066 .066
2.732	2.812	.066 .066
2.812	2.892	0.
2.892	2.971	.066 .066
2.971	3.051	0.
3.051	3.130	0.
3.130	3.210	.066 .066
3.210	3.289	.066 .066

FERRO-LUZZ 65.....NC 39 417

BEAM MOMENTUM= 4.970 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
.138	.050	4.026 .442
.225	.138	1.601 .279
.312	.225	.679 .182
.400	.312	.485 .153
.487	.400	.679 .182
.575	.487	.194 .097
.662	.575	.097 .069
.750	.662	.097 .069
.837	.750	.049 .049
.924	.837	.049 .049
1.012	.924	.049 .049
1.099	1.012	.097 .069
1.187	1.099	.049 .049
1.274	1.187	.049 .049
1.536	1.449	.097 .069

GEORGE 67.....NC 49A 9

Differential cross section for $K^+p \rightarrow K^{*0}(890)\Delta^{++}$
 $\downarrow_{K^+\pi^-} \downarrow_{p\pi^+}$

Figure 18. Differential cross section for the reaction
 $K^+p \rightarrow K^{*0}(890)\Delta^{++}(1238)$

$\downarrow_{K^+\pi^-}$

The $d\sigma/dt$ scale is in arbitrary units because experiments which report number of events rather than $d\sigma/dt$ (and do

not give a factor which would enable us to convert to σ) have also been included. The distributions given originally as $d\sigma/d\omega$ vs $\cos\theta_{c.m.}$, were transformed to $d\sigma/dt$ vs t [taking $m(K^*) = .89$ GeV and $m(\Delta) = 1.22$ GeV].

Differential cross section for $K^+ p \rightarrow K^* \Delta$

(Not plotted)

DIFFERENTIAL CROSS SECTION FOR K+ PROTON INTO DELTA+ K**

BEAM MOMENTUM= 2.970 GEV/C

THIS DATA WAS READ FROM A GRAPH

K*(890)+ DECAYS TO K0 PI+

DELTA+ DECAYS TO PROTON P10

MOMENTUM TRANSFER BETWEEN BEAM AND K*(890)+

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)	
.107	.187	1.224	.220
.187	.266	.395	.125
.266	.346	.197	.088
.346	.425	.118	.068
.425	.505	.079	.056
.505	.584	.237	.097
.584	.664	.079	.056
.664	.743	0.	
.743	.823	0.	
.823	.903	0.	
.903	.982	0.	
.982	1.062	0.	
1.062	1.141	.039	.039
1.141	1.221	0.	
1.221	1.300	0.	
1.300	1.380	0.	
1.380	1.459	.039	.039
1.459	1.539	.039	.039
1.539	1.619	0.	
1.619	1.698	0.	
1.698	1.778	0.	
1.778	1.857	0.	
1.857	1.937	0.	
1.937	2.016	.079	.056
2.016	2.096	0.	
2.096	2.176	0.	
2.176	2.255	0.	
2.255	2.335	0.	
2.335	2.414	0.	
2.414	2.494	0.	
2.494	2.573	0.	
2.573	2.653	.039	.039
2.653	2.732	.039	.039
2.732	2.812	0.	
2.812	2.892	0.	
2.892	2.971	0.	
2.971	3.051	0.	
3.051	3.130	0.	
3.130	3.210	0.	
3.210	3.289	0.	

FERRO-LUZZ 65.....NC

39 417

DIFFERENTIAL CROSS SECTION FOR K+ PROTON INTO DELTA++ K*0

BEAM MOMENTUM= 2.970 GEV/C

THIS DATA WAS READ FROM A GRAPH

DELTA++ DECAYS TO PROTON P1+

K*(890)0 DECAYS TO K0 P10

MOMENTUM TRANSFER BETWEEN BEAM AND K*(890)0

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)	
.107	.187	1.912	.355
.187	.266	1.187	.280
.266	.346	.396	.162
.346	.425	.528	.187
.425	.505	.462	.174
.505	.584	.132	.093
.584	.664	.396	.162
.664	.743	.132	.093
.743	.823	.198	.114
.823	.903	.132	.093
.903	.982	.132	.093
.982	1.062	.132	.093
1.062	1.141	0.	
1.141	1.221	.132	.093
1.221	1.300	0.	
1.300	1.380	.066	.066
1.380	1.459	.066	.066
1.459	1.539	0.	
1.539	1.619	0.	
1.619	1.698	.132	.093
1.698	1.778	0.	
1.778	1.857	.066	.066
1.857	1.937	.066	.066
1.937	2.016	0.	
2.016	2.096	0.	
2.096	2.176	.066	.066
2.176	2.255	0.	
2.255	2.335	0.	
2.335	2.414	0.	
2.414	2.494	0.	
2.494	2.573	.066	.066
2.573	2.653	0.	
2.653	2.732	0.	
2.732	2.812	.066	.066
2.812	2.892	.132	.093
2.892	2.971	0.	
2.971	3.051	0.	
3.051	3.130	0.	
3.130	3.210	.066	.066
3.210	3.289	.066	.066

FERRO-LUZZ 65.....NC

39 417

HBC

K^+ d Interactions

K⁺d interactions

P _{beam} (GeV/c)	σ_{total} (mb)	σ_{K^+pn} (mb)	σ_{K^+pp} (mb)	References
.233			1.000 + .400	SLATER 61
.330			- .300	
.376			2.700 .400	SLATER 61
.529			3.100 .400	SLATER 61
.591	26.710 ± 1.239		6.500 .600	SLATER 61
.619	28.610 .466			BUGG 68
.641			6.700 .600	BUGG 68
.643	28.130 .466			SLATER 61
.712	27.700 .363			BUGG 68
.767	28.980 .337			BUGG 68
.770	27.300 .671			BUGG 68
.811			6.600 .700	COOK 61
.822	29.560 .268			SLATER 61
.863	30.330 .260			BUGG 68
.863		20.870 ± .560	6.720 .400	BUGG 68
.888	30.100 .360			HIRATA 68
.903	31.270 .253			COOL 66
.937	32.140 .238			BUGG 68
.939	31.980 .340			BUGG 68
.968	32.720 .246			COOL 66
.970		20.910 .500	6.260 .290	BUGG 68
.989	32.400 .583			HIRATA 68
1.028	33.200 .240			COOK 61
1.040	34.290 .224			COOL 66
1.040	34.770 .240			BUGG 68
1.083	35.900 .218			COOL 66
1.091	36.040 .240			BUGG 68
1.139	37.340 .219			BUGG 68
1.141	37.220 .150			COOL 66
1.170	35.400 .583			COOK 61
1.188	37.360 .719			BUGG 68
1.191	37.690 .120			COOL 66
1.211		17.740 .780	4.990 .240	HIRATA 68
1.237	37.460 .280			BUGG 68
1.242	37.570 .110			COOL 66
1.292	37.430 .206			BUGG 68
1.292	37.430 .100			COOL 66
1.300	35.600 .671			COOK 61
1.342	36.990 .120			COOL 66
1.346	37.410 .219			BUGG 68
1.364		17.280 1.150	3.690 .290	HIRATA 68
1.392	36.780 .120			COOL 66
1.407	36.920 .206			BUGG 68
1.440	35.400 .583			COOK 61
1.442	36.430 .100			COOL 66
1.467	36.530 .219			BUGG 68
1.492	36.270 .100			COOL 66
1.550	36.030 .377			ABRAMS 67
1.562	36.290 .199			BUGG 68
1.593	35.890 .080			COOL 66
1.600	35.940 .376			ABRAMS 67
1.643	35.780 .080			COOL 66
1.670	35.100 .583			COOK 61
1.693	35.760 .060			COOL 66
1.700	35.780 .367			ABRAMS 67
1.743	35.890 .060			COOL 66
1.750	35.900 .370			ABRAMS 67
1.793	35.700 .060			COOL 66
1.800	35.680 .371			ABRAMS 67
1.850	35.630 .363			ABRAMS 67
1.893	35.530 .060			COOL 66
1.900	35.700 .364			ABRAMS 67
1.944	35.190 .205			BUGG 68
1.950	35.400 .367			ABRAMS 67
1.970	35.600 .363			COOK 61
1.993	35.410 .080			COOL 66
2.000	35.680 .360			ABRAMS 67
2.050	35.550 .362			ABRAMS 67
2.093	35.470 .080			COOL 66
2.100	35.340 .362			ABRAMS 67
2.150	35.280 .360			ABRAMS 67
2.193	35.220 .100			COOL 66
2.200	35.280 .360			ABRAMS 67
2.260	33.900 .671			COOK 61
2.300	34.980 .357			ABRAMS 67
2.300			1.500 .150	BUTTERWORT 65
2.350	35.050 .356			ABRAMS 67
2.393	35.450 .080			COOL 66
2.400	34.920 .354			ABRAMS 67
2.450	35.000 .354			ABRAMS 67
2.472	35.780 .192			BUGG 68
2.500	34.870 .354			ABRAMS 67
2.550	33.400 .583			COOK 61
2.550	34.870 .351			ABRAMS 67
2.600	34.850 .352			ABRAMS 67
2.650	34.750 .351			ABRAMS 67
2.700	34.770 .350			ABRAMS 67
2.750	34.720 .351			ABRAMS 67
2.800	34.670 .350			ABRAMS 67
2.850	33.400 .762			COOL 66
2.850	34.520 .349			ABRAMS 67
2.900	34.480 .347			ABRAMS 67
2.950	34.420 .347			ABRAMS 67
2.970			.750 .080	GOLDSCHMID 68
3.000	34.350 .346			ABRAMS 67
3.050	34.340 .346			ABRAMS 67
3.100	34.300 .345			ABRAMS 67
3.150	34.140 .344			ABRAMS 67
3.200	34.120 .344			ABRAMS 67
3.250	34.200 .344			ABRAMS 67
3.300	34.130 .344			ABRAMS 67
6.000	33.400 .300			GALBRAITH 65
8.000	33.900 .300			GALBRAITH 65
10.000	33.800 .300			GALBRAITH 65
12.000	33.900 .300			GALBRAITH 65
14.000	33.800 .300			GALBRAITH 65
16.000	33.400 .300			GALBRAITH 65
18.000	33.700 .300			GALBRAITH 65
20.000	34.200 .300			GALBRAITH 65

REFERENCES

1	COOK	61.....PRL	7 182
2	SLATER	61.....PRL	7 378
3	BUTTERWORT	65.....PRL	15 734
4	GALBRAITH	65.....PR	138 8913
5	COOL	66.....PRL	17 102
6	ABRAMS	67.....PRL	19 259
7	BUGG	68.....PR	168 1466
8	GOLDSCHMID	68.....PL	278 602
9	HIRATA	68.....PRL	21 1485

CNTR
HBC
DSC
CNTR
CNTR
CNTR
DSC
HBC

Figure 19. K⁺d total cross section, partial cross sections for K⁺d $\rightarrow$ K⁺pn (including K⁺d $\rightarrow$ K⁺d), and K⁺d $\rightarrow$ K⁰pp.

K^+d interactions

Figure 20. Cross sections for K^+d single pion production.

P_{beam} (GeV/c)	$\sigma_{K^0 pn \pi^+}$ (mb)	$\sigma_{K^+ pp \pi^-}$ (mb)	$\sigma_{K^0 pp \pi^0}$ (mb)	$\sigma_{K NN \pi}$ (mb)	References
.863	$1.240 \pm .140$	$.460 \pm .050$	$.480 \pm .080$	$2.740 \pm .240$	HIRATA 68
.968	$2.470 \pm .160$	$.910 \pm .050$	$.820 \pm .080$	$5.540 \pm .290$	HIRATA 68
1.211	$6.210 \pm .290$	$2.890 \pm .120$	$2.070 \pm .130$	$14.510 \pm .600$	HIRATA 68
1.364	$6.080 \pm .420$	$3.210 \pm .200$	$2.060 \pm .190$	$15.540 \pm .950$	HIRATA 68
REFERENCES					
1	HIRATA 68.....PRL	21 1485	HBC

Figure 21. Cross section for $K^+d \rightarrow K^0\pi^+d$.

P_{beam} (GeV/c)	$E_{c.m.}$ (GeV)	$\sigma_{K^0d\pi^+}$ (mb)	References	
.863	2.737	$.130 \pm .040$	HIRATA	68
.968	2.799	$.130 \pm .030$	HIRATA	68
1.211	2.944	$.140 \pm .030$	HIRATA	68
1.364	3.033	$.180 \pm .050$	HIRATA	68
2.300	3.547	$.150 \pm .035$	BUTTERWORT	65
2.970	3.880	$.228 \pm .025$	BUCHNER	69
REFERENCES				
1	BUTTERWORT 65.....PRL	15	500	HBC
2	HIRATA 68.....PRL	21	1485	HBC
3	BUCHNER 69.....NP	89	286	HBC

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

K^+n Interactions

For a brief discussion regarding problems in "unfolding" neutron information from deuterium data, see the text.

$K^+ n$ total cross section

P_{beam} (GeV/c)	$E_{\text{c.m.}}$ (GeV)	σ_{total} (mb)	References
.770		1.687 15.500 ± 1.140	COOK 61
.888 ± .009	1.742	16.850 .520	COOL 66
.939 .009	1.766	17.600 .450	COOL 66
.970	1.781	17.800 .854	COOK 61
.989 .010	1.790	18.530 .360	COOL 66
1.040 .010	1.814	18.910 .350	COOL 66
1.091 .011	1.838	20.610 .350	COOL 66
1.141 .011	1.861	21.250 .210	COOL 66
1.170	1.874	18.200 .854	COOK 61
1.191 .012	1.884	20.870 .170	COOL 66
1.242 .012	1.907	20.260 .160	COOL 66
1.292 .013	1.930	19.680 .150	COOL 66
1.300	1.934	18.500 1.140	COOK 61
1.342 .013	1.953	19.320 .160	COOL 66
1.392 .014	1.975	19.220 .160	COOL 66
1.440	1.997	18.100 .949	COOK 61
1.442 .014	1.998	19.070 .140	COOL 66
1.492 .015	2.020	18.950 .130	COOL 66
1.550 .012	2.045	18.910 .240	ABRAMS 67
1.593 .016	2.064	18.790 .120	COOL 66
1.600 .012	2.067	18.890 .241	ABRAMS 67
1.643 .016	2.086	18.670 .100	COOL 66
1.690	2.106	18.500 .854	COOK 61
1.693 .017	2.107	18.690 .080	COOL 66
1.700 .013	2.110	18.830 .218	ABRAMS 67
1.743 .017	2.129	18.880 .080	COOL 66
1.750 .013	2.132	18.860 .218	ABRAMS 67
1.793 .018	2.150	18.730 .080	COOL 66
1.800 .013	2.153	18.530 .232	ABRAMS 67
1.850 .014	2.174	18.660 .212	ABRAMS 67
1.893 .019	2.192	18.500 .080	COOL 66
1.900 .014	2.194	18.690 .212	ABRAMS 67
1.950 .015	2.215	18.700 .208	ABRAMS 67
1.970	2.223	18.600 .671	COOK 61
1.993 .020	2.233	18.550 .110	COOL 66
2.000 .015	2.236	18.790 .204	ABRAMS 67
2.050 .015	2.256	18.540 .211	ABRAMS 67
2.093 .021	2.273	18.670 .110	COOL 66
2.100 .016	2.276	18.490 .215	ABRAMS 67
2.150 .016	2.296	18.430 .210	ABRAMS 67
2.193 .022	2.313	18.400 .130	COOL 66
2.200 .016	2.316	18.400 .209	ABRAMS 67
2.260	2.340	17.700 .854	COOK 61
2.300 .017	2.355	18.270 .204	ABRAMS 67
2.350 .018	2.375	18.260 .196	ABRAMS 67
2.393 .024	2.391	18.630 .110	COOL 66
2.400 .018	2.394	18.090 .198	ABRAMS 67
2.450 .018	2.413	18.250 .192	ABRAMS 67
2.500 .019	2.432	18.110 .198	ABRAMS 67
2.550	2.451	17.100 .854	COOK 61
2.550 .019	2.451	18.170 .191	ABRAMS 67
2.600 .019	2.470	18.090 .191	ABRAMS 67
2.650 .020	2.488	18.020 .190	ABRAMS 67
2.700 .020	2.507	18.110 .191	ABRAMS 67
2.750 .021	2.525	18.060 .190	ABRAMS 67
2.800 .021	2.543	18.010 .190	ABRAMS 67
2.830	2.554	17.500 .949	COOK 61
2.850 .021	2.562	17.950 .189	ABRAMS 67
2.900 .022	2.580	17.850 .185	ABRAMS 67
2.950 .022	2.598	17.810 .188	ABRAMS 67
3.000 .022	2.615	17.810 .188	ABRAMS 67
3.050 .023	2.633	17.830 .188	ABRAMS 67
3.100 .023	2.651	17.850 .100	ABRAMS 67
3.150 .024	2.668	17.610 .186	ABRAMS 67
3.200 .024	2.685	17.610 .183	ABRAMS 67
3.250 .024	2.703	17.660 .184	ABRAMS 67
3.300 .025	2.720	17.550 .182	ABRAMS 67
6.000 .105	3.527	17.500 .400	GALBRAITH 65
8.000 .140	4.023	17.600 .400	GALBRAITH 65
10.000 .175	4.465	17.500 .400	GALBRAITH 65
12.000 .210	4.868	17.600 .400	GALBRAITH 65
14.000 .245	5.239	17.500 .400	GALBRAITH 65
16.000 .280	5.586	17.400 .400	GALBRAITH 65
18.000 .315	5.913	17.600 .400	GALBRAITH 65
20.000 .350	6.223	17.700 .400	GALBRAITH 65

REFERENCES

1	COOK	61.....PRL
2	GALBRAITH	65.....PR
3	COOL	66.....PRL
4	ABRAMS	67.....PRL

7	182
138	8913
17	102
19	259

CNTR
CNTR
CNTR
CNTR

Figure 22. K^+n total cross section, as deduced from the directly-measured K^+p and K^+d total cross sections. Glauber corrections have been made, but in general other deuteron corrections have not been included. Galbraith [ref. 39] assumes that the real part of the forward amplitudes are purely imaginary.

Figure 23. Cross section for $K^+ n \rightarrow n K^0 \pi^+$. No deuteron correction made.

P_{beam} (GeV/c)	$E_{c.m.}$ (GeV)	$\sigma_{K^0 n \pi^+}$ (mb)	References	
.863	1.731	$.130 \pm .040$	HIRATA	68
.968	1.780	.350 .050	HIRATA	68
1.211	1.893	1.310 .120	HIRATA	68
1.364	1.963	1.120 .150	HIRATA	68
REFERENCES				
1	HIRATA	68.....PRL	21	1485

KN Interactions ($I = 0$)

For a brief discussion regarding problems of "unfolding" $I = 0$ information from deuterium data, see the text.

KN interactions ($I=0$)

P_{beam} (GeV/c)	σ_{total} (mb)	σ_{elastic} (mb)	$\sigma_{\text{KN}\pi}$ (mb)	References
.775	21.100			CARTER 67
.825	22.100			CARTER 67
.863		21.500	$1.000 \pm .400$	HIRATA 68
.875	22.600			CARTER 67
.888	19.310 ± 1.350			COOL 66
.925	23.100			CARTER 67
.939	19.620	21.500	$2.500 \pm .600$	COOL 66
.968				HIRATA 68
.975	24.200			CARTER 67
.989	21.090	.940		COOL 66
1.025	25.300			CARTER 67
1.040	20.420	.890		COOL 66
1.075	25.600			CARTER 67
1.091	24.090	.890		COOL 66
1.125	25.200			CARTER 67
1.141	24.400	.540		COOL 66
1.175	24.500			CARTER 67
1.191	23.270	.430		COOL 66
1.211		16.100	8.000 ± 1.000	HIRATA 68
1.225	23.900			CARTER 67
1.242	21.970	.420		COOL 66
1.275	23.200			CARTER 67
1.292	20.740	.390		COOL 66
1.325	22.400			CARTER 67
1.342	20.200	.380		COOL 66
1.364		13.000	8.900 ± 1.800	HIRATA 68
1.375	21.700			CARTER 67
1.392	20.170	.380		COOL 66
1.425	21.400			CARTER 67
1.442	20.090	.360		COOL 66
1.475	21.400			CARTER 67
1.492	19.980	.340		COOL 66
1.525	21.200			CARTER 67
1.550	20.110	.448		ABRAMS 67
1.575	21.000			CARTER 67
1.593	19.830	.310		COOL 66
1.600	20.070	.421		ABRAMS 67
1.625	20.800			CARTER 67
1.643	19.480	.240		COOL 66
1.675	20.700			CARTER 67
1.693	19.530	.220		COOL 66
1.700	19.930	.352		ABRAMS 67
1.725	20.600			CARTER 67
1.743	19.960	.220		COOL 66
1.750	19.890	.343		ABRAMS 67
1.775	20.600			CARTER 67
1.793	19.650	.220		COOL 66
1.800	19.080	.407		ABRAMS 67
1.825	20.500			CARTER 67
1.850	19.540	.317		ABRAMS 67
1.875	20.400			CARTER 67
1.893	19.180	.220		COOL 66
1.900	19.590	.318		ABRAMS 67
1.925	20.300			CARTER 67
1.950	19.650	.310		ABRAMS 67
1.975	20.300			CARTER 67
1.993	19.500	.290		COOL 66
2.000	19.950	.290		ABRAMS 67
2.025	20.200			CARTER 67
2.050	19.370	.316		ABRAMS 67
2.075	20.200			CARTER 67
2.093	19.820	.290		COOL 66
2.100	19.420	.325		ABRAMS 67
2.125	20.200			CARTER 67
2.130	19.280	.316		ABRAMS 67
2.175	20.000			CARTER 67
2.193	19.260	.310		COOL 66
2.200	19.210	.315		ABRAMS 67
2.225	19.900			CARTER 67
2.275	19.800			CARTER 67
2.300	19.100	.299		ABRAMS 67
2.325	19.800			CARTER 67
2.350	19.000	.255		ABRAMS 67
2.375	19.700			LAKIER 67
2.393	19.710	.290		COOL 66
2.400	18.620	.288		ABRAMS 67
2.425	19.400			CARTER 67
2.450	19.020	.249		ABRAMS 67
2.475	19.500			CARTER 67
2.500	18.740	.267		ABRAMS 67
2.525	19.400			CARTER 67
2.550	18.900	.235		ABRAMS 67
2.575	19.400			CARTER 67
2.600	18.680	.246		ABRAMS 67
2.625	19.300			CARTER 67
2.650	18.570	.245		ABRAMS 67
2.675	19.300			CARTER 67
2.700	18.900	.234		ABRAMS 67
2.725	19.200			LAKIER 67
2.750	18.710	.246		ABRAMS 67
2.775	19.200			CARTER 67
2.800	18.620	.246		ABRAMS 67
2.825	19.200			CARTER 67
2.850	18.600	.245		ABRAMS 67
2.875	19.200			CARTER 67
2.900	18.360	.219		ABRAMS 67
2.925	19.200			CARTER 67
2.950	18.310	.230		ABRAMS 67
2.975	19.200			CARTER 67
3.000	18.420	.231		ABRAMS 67
3.025	19.200			CARTER 67
3.050	18.520	.232		ABRAMS 67
3.075	19.200			CARTER 67
3.100	18.610	.252		ABRAMS 67
3.125	19.100			CARTER 67
3.150	18.070	.229		ABRAMS 67
3.175	19.100			CARTER 67
3.200	18.090	.217		ABRAMS 67
3.225	19.100			CARTER 67
3.250	18.190	.218		ABRAMS 67

REFERENCES

1	COOL	66.....PRL	17 102
2	ABRAMS	67.....PRL	19 259
3	CARTER	67.....PRL	18 801
4	HIRATA	68.....PRL	21 1485

CNTR
CNTR

HBC

KN interactions ($I=0$)

Figure 24. Isospin zero KN total, elastic and single-pion production cross sections. The smoothed $I = 0$ total cross section calculated by Carter [Reference 37] using dispersion relations, is given as points without error bars. Glauber corrections are included in all but Hirata et al. data.

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

K^+ d Partial Differential Cross Sections

Differential cross section for $K^+ d \rightarrow ppK^0$

DISTRIBUTION IN COS THETA OF THE K^0 WITH RESPECT TO THE BEAM DIRECTION IN THE C.M. SYSTEM

BEAM MOMENTUM= .330 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)	
-1.000	-.600	.110	.040
-.600	-.200	.250	.050
-.200	.200	.400	.070
.200	.600	.290	.060
.600	1.000	.160	.040
SLATER	61.....PRL	7	378

BEAM MOMENTUM= .529 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)	
-1.000	-.700	.150	.070
-.700	-.400	.440	.100
-.400	-.100	.600	.120
-.100	.200	.730	.130
.200	.500	.040	.140
.500	.800	.580	.120
.800	1.000	.200	.080
SLATER	61.....PRL	7	378

BEAM MOMENTUM= .600 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)	
-1.000	-.800	.220	.020
-.800	-.600	.340	.020
-.600	-.400	.470	.020
-.400	-.200	.550	.020
-.200	0.	.590	.040
0.	.200	.680	.030
.200	.400	.670	.020
.400	.600	.690	.020
.600	.800	.600	.020
.800	1.000	.380	.020
RAY	69.....BNL	13616	

BEAM MOMENTUM= .641 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)	
-1.000	-.700	.320	.080
-.700	-.400	.440	.080
-.400	-.100	.500	.100
-.100	.200	.840	.120
.200	.500	.640	.120
.500	.800	.600	.100
.800	1.000	.420	.100
SLATER	61.....PRL	7	378

BEAM MOMENTUM= .811 GEV/C

THIS DATA WAS READ FROM A GRAPH

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)	
-1.000	-.700	.330	.090
-.700	-.400	.440	.090
-.400	-.100	.500	.100
-.100	.200	.600	.120
.200	.500	.720	.120
.500	.800	.610	.130
.800	1.000	.500	.120
SLATER	61.....PRL	7	378

Differential cross section for $K^+ d \rightarrow pp K^0$

Figure 25. Differential cross section for the charge exchange reaction $K^+ d \rightarrow K^0 pp$. The slower proton in the lab system is taken to be the spectator, and only events with lab spectator momenta below a cutoff value of 150 to 300 MeV/c (depending on the experiment) are included. The angle θ is usually taken to be that between the incident K^+ and the outgoing K^0 in the center of mass of the outgoing K^0 and the spectator proton. Some experimenters calculate θ in slightly different centers of mass, but the difference is claimed to be negligible. From these distributions one can presumably calculate the distributions for $K^+ n \rightarrow K^0 p$ by making the appropriate "deuterium corrections."

Differential cross section for $K^+ d \rightarrow ppK^0$

BEAM MOMENTUM= 2.300 GEV/C

THIS DATA WAS READ FROM A GRAPH

MOMENTUM TRANSFER BETWEEN BEAM AND K0

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
0.	.080	1.600 .400
.080	.140	2.000 .400
.140	.200	2.200 .400
.200	.260	2.000 .400
.260	.360	1.500 .300
.360	.460	1.400 .300
.460	.560	1.300 .300
.560	.650	1.400 .300
.650	.850	.600 .200
.850	1.050	.300 .100
1.100	1.800	.090 .020
1.800	3.400	.080 .020

BUTTERWORTH 65.....PRL

15 734

 $P_{beam}^{lab} = 2.300 \text{ GeV/c}$

BEAM MOMENTUM= 2.970 GEV/C

MOMENTUM TRANSFER BETWEEN BEAM AND K0

MDM TRANSF	SIGMA +- DSIGMA (MILLIBARNS/(GEV/C)**2)
.025	.870 .243
.075	1.320 .305
.125	1.130 .284
.175	1.570 .336
.225	1.020 .253
.275	1.320 .305
.325	1.310 .305
.375	1.170 .284
.425	.980 .223
.500	.510 .132
.610	.460 .112
.770	.170 .050
.990	.067 .030
1.410	.026 .013

GOLOSCHMID 68.....PL

278 602

 $P_{beam}^{lab} = 2.970 \text{ GeV/c}$

BEAM MOMENTUM= 2.300 GEV/C

THIS DATA WAS READ FROM A GRAPH

DISTRIBUTION IN COS THETA OF THE K0 WITH RESPECT TO THE BEAM

MINIMUM	MAXIMUM	SIGMA +- DSIGMA (MILLIBARNS/STERADIAN)
.610	.810	.030 + .020
		- .010
.810	.910	.200 + .100
		- .050
.915	.945	.800 + .300
		- .200
.940	.960	2.000 1.000
.965	.975	2.500 1.000
.975	.985	4.000 1.000
.985	.995	3.000 1.000
.990	1.000	3.200 1.000

BUTTERWORTH 65.....PRL

15 734

 $P_{beam}^{lab} = 2.300 \text{ GeV/c}$

Figure 25 (Continued)

Section III.

DATA LISTINGS

In this section we present a listing of all the K^+ articles on our DATA TAPE. These are the actual data used in forming the graphs and tables in Section II. The information is presented article-by-article, just as we store it.

We debated for some time whether or not we should give these listings because they are rather lengthy. As an experiment, we decided to put them in this edition. If you find them useful, let us know.

Actually, these listings do contain a large amount of information not included in the tables and graphs already presented in Section II.

- In particular there are a number of articles giving mass spectra etc., for which we have punched no data. Nevertheless, the person interested in K^+ interactions may find many of these papers quite useful. (These papers are not referred to in any way in Section II.)

- In addition we have also punched the title and abstract for every article, to assist you in your selection of articles for further reading.

- Also in this section you will find comments on many pieces of data—it is in general not yet practical to present these comments in Section II.

- Many articles give data that we feel we cannot meaningfully compile at present (only partially corrected, integrated only over a certain interval, etc.). These data have in many cases been punched and will be found in this section.

- You will also find in this section, data reported as upper and lower limits, approximate values, etc.

- We also indicate here how background-resonance separations were made and the values of the parameters used (or fitted).

- Occasionally we do not use the data

as originally given in the article. This section tells exactly where our data came from (private communications, unpublished companion report, etc.).

- The size of an experiment is frequently indicated by the total number of pictures taken, or by the number of events in various distributions.

- To give you an idea of the scope of a particular article, KEYWORDS are included for each article. These words can also be used to form classified indices (see Section IV).

- Some papers give fitted values for various parameters in which we are interested. Recently we have started encoding this information. In the future we plan to include more of these fitted quantities.

To repeat, the above items are some of the things you will find in this section that are not presented in Section II.

We have also found that theses are frequently hard to come by. Thus we feel that our listing of theses may help give their data greater distribution than they might otherwise have. We would like to make the general appeal that a copy of all particle physics theses be sent to us.

Finally this section may serve the useful function of permitting the reader to easily check on the accuracy of our input data. The data is arranged article-by-article, and in most cases we have indicated [in square brackets] the exact location of the data in the article (i.e., the figure, table, or page number). If you find any errors or misinterpretations, please let us know as soon as possible.

As for the organization of the information in this section, we should first mention that the order of the articles is "random," and has no physical meaning. The order is, however, the same as given in the Reference list,

and as given in the Indices in Section IV.

Above the double line in each article you will find the title, authors and institutions, abstract (if present), footnote references, KEY WORDS, comments, beam information, etc.

Many articles have references listed under two categories:

"CROSS REFERENCES" mean that these papers refer to the same data or same experiment as the present paper;

"ADDITIONAL FOOTNOTE REFERENCES" (in future we'll call it ADDITIONAL CITATIONS) are all other references found in the paper. They may refer to theory, experimental apparatus, similar experiments done by other groups, etc.

Below the double line in each article appear the data. We enter the data in exactly the same units as given by the authors. (This is done primarily to facilitate the verification of the data.) If we do alter the data in any way, we indicate this fact by an appropriate comment.

Occasionally authors give the same data in two different forms. We punch both, if we feel that both forms are useful, and display them side-by-side in the listings that follow.

We have tried to be particularly careful about including systematic errors, whenever given by the authors. We have also tried to indicate exactly how resonance and background separations have been made. In some cases it is quite unclear from the original article and we have had to contact the authors directly.

Another reason for contacting authors has been to get tables of data that correspond to the published graphs. If we are unable to get tables from an author, or if the article is more than a couple of years old, we read the data of the published graph, and then include the warning that "This data was read from a graph." (In some cases the tables we re-

ceived have been more up to date than the published graphs.)

Because of the limitations imposed by not having a printer with Greek letters, we have had to spell out many symbols. One exception is, however, the abbreviation for microbarns, μb : we use "UB." We hope that we will be able, in the near future, to use a more complete set of characters, so that our output will be easier to read.

1 SEARCH FOR NEW MESON RESONANCES IN K+P INTERACTIONS AT 12.7 GEV/C. (PHYS. LETTERS 278, 655 (1968))
J.C.BERLINGHIERI, N.S.FARRER, T.FERBEL, B.E.NORMAN, A.C.MELISSINOS, P.F.SLATTERY, T.YAMANOUCHI, H.YUTA (UNIV. OF ROCHESTER, ROCHESTER, N. Y., USA)

ABSTRACT IN A STUDY OF K+P INTERACTIONS AT 12.7 GEV/C WE HAVE CARRIED OUT A SEARCH FOR MESON RESONANCES HAVING ANY ONE OF THE FOLLOWING PROPERTIES (A) A STRANGENESS OF S=2, (B) S=1 AND ISOTOPIC SPIN OF I=1/2, OR (C) S=1, I=1/2 AND DECAYING INTO AN ANTIHYPHERON ANTI-Y AND A NUCLEON N.

CROSS REFERENCES
SEE ALSO PHYS. LETTERS 278, 177 (1968).
THIS ARTICLE SUPERSEDES PART OF CERN HADRON CONFERENCE 2 172 (1968).

ADDITIONAL FOOTNOTE REFERENCES
PHYS. REV. LETTERS 17, 1200 (1966), PHYS. REV. 135, 8779 (1964), AND PHYS. REV. LETTERS 23, 755 (1968).

ARTICLE READ BY ODETTE BENARY IN 12/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K+ ON PROTON AT 12.7 GEV/C.

THIS EXPERIMENT USES THE B.N.L. 80 IN. (H) BUBBLE CHAMBER.

KEY WORDS = CROSS SECTION MASS SPECTRUM

CROSS SECTION FOR K+ PROTON → K+ K+ P10 LAMBDA. (TABLE 2)

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
12.7	14. ± 5.	40

CROSS SECTION FOR K+ PROTON → K+ K0 P1+ LAMBDA. (TABLE 2)

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
12.7	26. ± 7.	68

CROSS SECTION FOR K+ PROTON → K+ K0 P1+ P10 LAMBDA. (TABLE 2)

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
12.7	30. ± 15.	16

CROSS SECTION FOR K+ PROTON → K0 K0 P1+ P1+ LAMBDA. (TABLE 2)

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
12.7	20. ± 10.	11

CROSS SECTION FOR K+ PROTON → K+ K0 P1+ P1+ P1- LAMBDA. (TABLE 2)

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
12.7	39. ± 13.	17

CROSS SECTION FOR K+ PROTON → NEUTRON K0 P1+ P1+. (PAGE 667)

(THIS DATA REPLACES THE VALUE GIVEN EARLIER IN BERLINGHIERI, ET. AL., CERN HADRON CONFERENCE 172 (1968))

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
12.7	150. ± 30.	243

CROSS SECTION FOR K+ PROTON → K+ K+ LAMBDA + K+ K+ SIGMA0. (TABLE 2)

* * THIS DATA SHOULD NOT BE USED * * (IN CERN HADRON CONF.(VOL.2) P.171(1968) THEY GIVE A VALUE FOR K+P → K+K+LAMBDA ALONE.)

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
12.7	14. ± 3.	36

- 2 EVIDENCE FOR KAPPA(725) IN K⁺P INTERACTIONS AT 3 GEV/C. [PHYS. LETTERS 12, 255 (1964)]
M. FERRO-LUZZI, R. GEORGE, Y. GOLOSCHMIDT-CLERMONT, V. P. HENRI, B. JONGEJANS, D. W. G. LEITH, G. R. LYNCH, F. MULLER, J. M. PERREAU (EUROPEAN
ORG. FOR NUC. RES., GENEVA, SWITZERLAND)
- CROSS REFERENCES
PART OF THIS ARTICLE SUPERSEDED BY PHYS. LETTERS 278, 468 (1968).
- ADDITIONAL FOOTNOTE REFERENCES
SIENNA CONFERENCE 1 189 (1963), PHYS. LETTERS 9, 359 (1964), PHYS. LETTERS 5, 279, PHYS. LETTERS 5, 283 (1963), AND
SIENNA CONFERENCE 1 125 (1963).
- ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 3/68, AND VERIFIED BY ROGER BLAND.
- BEAM IS K⁺ ON PROTON AT 3 GEV/C.
- THIS EXPERIMENT USES THE SACLAY 81 CM (H) BUBBLE CHAMBER.
- GENERAL COMMENTS ON THIS ARTICLE
1 EVIDENCE FOR THE KAPPA(725) NOT CODED, CROSS SECTIONS ONLY APPROXIMATE
- KEY WORDS = KAPPA MESON K*(1890) DELTA(1230) CROSS SECTION MASS SPECTRUM OMEGA MESON
-
- CROSS SECTION FOR K⁺ PROTON = K⁺ PROTON PI⁺ PI⁻ P10.
[THIS DATA SHOULD NOT BE USED - A MORE RECENT VALUE MAY BE FOUND IN BASSOMPIERRE, ET.AL., PHYS. LETTERS 278, 468 (1968)]
- LABORATORY
BEAM MOMENTUM
GEV/C (1) MILLI-BARNS (1)
3. .8
- (1) VALUE IS APPROXIMATE ONLY.
-
- CROSS SECTION FOR K⁺ PROTON = K⁰ PROTON PI⁺ PI⁻ PI⁻.
[THIS DATA SHOULD NOT BE USED - A MORE RECENT VALUE MAY BE FOUND IN BASSOMPIERRE, ET.AL., PHYS. LETTERS 278, 468 (1968)]
- LABORATORY
BEAM MOMENTUM
GEV/C (1) MILLI-BARNS (1)
3. .4
- (1) VALUE IS APPROXIMATE ONLY.
-
- CROSS SECTION FOR K⁺ PROTON = K⁺ NEUTRON PI⁺ PI⁻ PI⁻.
[THIS DATA SHOULD NOT BE USED - A MORE RECENT VALUE MAY BE FOUND IN BASSOMPIERRE, ET.AL., PHYS. LETTERS 278, 468 (1968)]
- LABORATORY
BEAM MOMENTUM
GEV/C (1) MILLI-BARNS (1)
3. .2
- (1) VALUE IS APPROXIMATE ONLY.
-
- CROSS SECTION FOR K⁺ PROTON = OMEGA MESON K⁺ PROTON.
OMEGA MESON = PI⁺ PI⁻ P10 (1)
- LABORATORY
BEAM MOMENTUM
GEV/C (2) MILLI-BARNS (2)
3. .1
- (1) USED SIMPLE MASS CUT.
(2) VALUE IS APPROXIMATE ONLY.
- 3 EVIDENCE FOR K*(1250) RESONANCE PRODUCTION IN K⁺P INTERACTIONS AT 9 GEV/C. [PHYS. REV. LETTERS 19, 972 (1967)]
G. GOLDBABER, A. FIRESTONE, B. C. SHEN [U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA, AND UNIV. OF CALIFORNIA, BERKELEY, CALIF.,
USA]
- ABSTRACT WE PRESENT EVIDENCE FOR K*(1250) RESONANCE PRODUCTION IN K⁺P INTERACTIONS AT 9 GEV/C. WE OBSERVE THAT THE
LARGE K PI PI MASS ENHANCEMENT IN THE REGION 1.1 TO 1.5 BEV PROBABLY CONSISTS OF THREE RESONANCES AT THE OBSERVED
MASSES 1.25, 1.36, AND 1.42 BEV ABOVE A BROAD BACKGROUND DUE PRIMARILY TO DIFFRACTION DISSOCIATION.
- ADDITIONAL FOOTNOTE REFERENCES
BERKELEY CONFERENCE 103 (1967), PHYS. REV. LETTERS 10, 1087 (1967), BULL. AM. PHYS. SOC. 12, 506 (1967), BULL. AM.
PHYS. SOC. 12, 540 (1967), BULL. AM. PHYS. SOC. 12, 557 (1967), PHYS. REV. LETTERS 19, 44 (1967), PHYS. REV. LETTERS 17, 726 (1966), PHYS. LETTERS 9, 207 (1964), DUBNA CONFERENCE 987 (1964),
PHYS. REV. LETTERS 19, 976 (1967), LRL INTERDEPARTMENTAL REPORT P-117 (1965), LRL INTERDEPARTMENTAL REPORT P-126, AND
PHYS. LETTERS 22, 357 (1966).
- ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 4/68, AND VERIFIED BY LEROY PRICE.
- BEAM IS K⁺ ON PROTON AT 9 GEV/C.
- THIS EXPERIMENT USES THE B.N.L. 80 IN. (H) BUBBLE CHAMBER.
- GENERAL COMMENTS ON THIS ARTICLE
1 NO CROSS SECTION DATA GIVEN.
- KEY WORDS = MASS SPECTRUM K*(1320) K*(1420) C MESON ANGULAR DISTRIBUTION
-
- *****
* NO DATA PUNCHED FOR THIS ARTICLE *

4 TOTAL CROSS SECTIONS FOR P, P-BAR, CHARGED K, AND CHARGED PI ON HYDROGEN BETWEEN 3 AND 10 GEV/C. [PHYS. REV. LETTERS 5, 333 (1960)]

G.VONDARDEL, O.H. FRISCH, R. MERMOD, R.H. MILBURN, P.A. PIRQUE, M. VIVARGENT, G. WEBER, K. WINTER (EUROPEAN ORG. FOR NUC. RES., GENEVA, SWITZERLAND)

ADDITIONAL FOOTNOTE REFERENCES

JETP 34 499 (1958), PHYS. REV. LETTERS 4, 89 (1960), REVIEW OF SCIENTIFIC INSTRUMENTS 25, 1070 (1954), PHYS. REV. LETTERS 3, 568 (1959), JETP 37 872 (1960), PHYS. REV. 119, 2068 (1953), PHYS. REV. LETTERS 2, 117 (1959), PHYS. REV. 118, 553 (1960), UCRL 3639 (1957), AND UCRL 8965.

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 6/68, AND VERIFIED BY LEROY PRICE.

BEAM NO. 1 IS K+ ON PROTON FROM 3 TO 10 GEV/C.
 NO. 2 IS PROTON ON PROTON FROM 3 TO 10 GEV/C.
 NO. 3 IS ANTI-PROTON ON PROTON FROM 3 TO 10 GEV/C.
 NO. 4 IS K- ON PROTON FROM 3 TO 10 GEV/C.
 NO. 5 IS PI+ ON PROTON FROM 3 TO 10 GEV/C.
 NO. 6 IS PI- ON PROTON FROM 3 TO 10 GEV/C.

THIS EXPERIMENT USES COUNTERS.

KEY WORDS = CROSS SECTION

K+ PROTON TOTAL CROSS SECTION.

.....
 . THIS DATA WAS READ FROM A GRAPH .

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
3.0	24.5 ± 2.5
3.5	23.5 ± 2.5
5.0	20.0 ± 1.5
8.0	19.5 ± 1.0

K- PROTON TOTAL CROSS SECTION.

(- NO DATA PUNCHED FOR THIS PARAMETER.)

PI+ PROTON TOTAL CROSS SECTION.

(- NO DATA PUNCHED FOR THIS PARAMETER.)

PI- PROTON TOTAL CROSS SECTION.

(- NO DATA PUNCHED FOR THIS PARAMETER.)

PROTON PROTON TOTAL CROSS SECTION.

(- NO DATA PUNCHED FOR THIS PARAMETER.)

PBAR PROTON TOTAL CROSS SECTION.

(- NO DATA PUNCHED FOR THIS PARAMETER.)

5 EVIDENCE FOR A K*(1320) RESONANCE. [PHYS. REV. LETTERS 17, 726 (1966)]

B.C. SHEN, I. BUTTERWORTH, C. FU, G. GOLDBERGER, S. GOLDBERGER, G.H. TRILLING (U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA, AND UNIV. OF CALIFORNIA, BERKELEY, CALIF., USA)

ABSTRACT IT IS SUGGESTED THAT THE BROAD K*(1890)PI ENHANCEMENT FROM 1.15 TO 1.55 BEV IS A COMPLEX PHENOMENON CONSISTING OF AT LEAST THREE PARTS -- KINEMATIC ENHANCEMENT VIA THE DECK MECHANISM, A POSSIBLE NEW K* RESONANCE AT 1320 MEV, AND AN ALTERNATIVE DECAY MODE OF THE K*(1430). IT IS FURTHER SPECULATED THAT THE A1 AND K*(1320) COULD BE MEMBERS OF AN OCTET.

ADDITIONAL FOOTNOTE REFERENCES

PHYS. LETTERS 16, 184 (1965), UCRL 16332 (1965), UCRL 16744 (1966), PHYS. REV. LETTERS 13, 159 (1964), PHYS. REV. LETTERS 15, 329 (1965), PHYS. LETTERS 15, 281 (1965), CAL TECH CALT 58-85, PHYS. REV. LETTERS 15, 731 (1965), PHYS. LETTERS 22, 112 (1966), PHYS. REV. LETTERS 14, 1074 (1965), AND PHYS. LETTERS 17, 347 (1965).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 4/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K+ ON PROTON AT 4.6 GEV/C.

THIS EXPERIMENT USES THE B.N.L. 80 IN. (H) RHAIFF CHAMBER.

KEY WORDS = K*(1320) K*(1420) MASS SPECTRUM MOMENTUM TRANSFER

.....
 * NO DATA PUNCHED FOR THIS ARTICLE *

6 PION PRODUCTION BY K⁺ MESONS ON HYDROGEN AND DEUTERIUM (CERN CONFERENCE 358 (1962))

J. FISK, H. K. TICHON, D. H. STORK (UNIV. OF CALIF., LOS ANGELES, CALIF., USA)
 W. CHINDENSKY, G. GOLDBERGER, S. GOLDBERGER, T. F. STUBBS (U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA)

CROSS REFERENCES
 THIS ARTICLE SUPERSEDES PART OF PHYS. REV. LETTERS 7, 188 (1961).

ADDITIONAL FOOTNOTE REFERENCES
 PHYS. REV. LETTERS 8, 447 (1962), AND PHYS. REV. LETTERS 7, 378 (1961).

ARTICLE READ BY LEROY PRICE IN 1/68, AND VERIFIED BY ROGER BLAND.

BEAM NO. 1 IS K⁺ ON PROTON FROM 642 TO 810 GEV/C.
 NO. 2 IS K⁺ ON DEUTERON FROM 642 TO 810 GEV/C.

THIS EXPERIMENT USES THE L.R.L. 15 IN. (H) BUBBLE CHAMBER.

KEY WORDS - CROSS SECTION MASS SPECTRUM MESON WITH $\gamma=1$ AT 720 MEV BARYON RESONANCE WITH $\gamma=2$ AT 1480 MEV
 ANGULAR DISTRIBUTION

CROSS SECTION FOR K⁺ PROTON - K MESON NUCLEON PION. (PAGE 358)

(AT THESE ENERGIES THIS IS THE ONLY INELASTIC CHANNEL OPEN.)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.642	.06 ± .03
.810	.95 ± .16 (1)

(1) THIS DATA REPLACES THE VALUE GIVEN EARLIER IN STUBBS, ET. AL., PHYS. REV. LETTERS 7, 188 (1961).

CROSS SECTION FOR K⁺ PROTON - K⁰ PI⁺ PROTON. (TABLE 1)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS
.81	.64 ± .08	63

CROSS SECTION FOR K⁺ PROTON - K⁺ PI⁰ PROTON. (TABLE 1)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS
.81	.22 ± .05	22

CROSS SECTION FOR K⁺ PROTON - K⁺ PI⁺ NEUTRON. (TABLE 1)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS
.81	.18 ± .04	11

7 K⁺P INTERACTION AT 455 MEV (PHYS. REV. LETTERS 7, 188 (1961))

T.F. STUBBS, H. BRADNER, M. CHINOWSKY, G. COLOMABER, S. GOLDMABER (U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA)
W.E. SLATER, D.H. STORK, H.K. TICHN (UNIV. OF CALIF., LOS ANGELES, CALIF., USA)

CROSS REFERENCES

PART OF THIS ARTICLE SUPERSEDED BY CERN CONFERENCE 358 (1962).

ADDITIONAL FOOTNOTE REFERENCES

ROCHESTER CONFERENCE 451 (1960), PHYS. REV. 109, 2121 (1958), NUOVO CIMENTO 12, 241 (1959), PHYS. REV. 107, 279 (1957),
PHYS. REV. 118, 553 (1960), PHYS. REV. LETTERS 2, 117 (1959), PHYS. REV. LETTERS 5, 333 (1960), ANNUAL REV. OF
NUCLEAR SCIENCE 9, 61 (1959), PHYS. REV. 94, 1799 (1954), PHYS. REV. 109, 2133 (1958), PHYS. REV. 113, 1303 (1959),
AND PHYS. REV. 120, 1898 (1960).

ARTICLE READ BY NAOMI BARASH-SCHMINT IN 1/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K⁺ ON PROTON AT .81 GEV/C.

THIS EXPERIMENT USES THE L.R.L. 15 IN. (H) BUBBLE CHAMBER.

GENERAL COMMENTS ON THIS ARTICLE

1 ASSUMES K⁺ INTO TAU = .061 ± .002

KEY WORDS = CROSS SECTION ANGULAR DISTRIBUTION PHASE SHIFT

K⁺ PROTON ELASTIC CROSS SECTION.

(CONTAINS SOME COULOMB EFFECTS. ANGULAR DISTRIBUTION IS EXTRAPOLATED FROM COS(THETA) = .96 TO 1.0.)

LABORATORY BEAM ENERGY GEV	MILLI-BARNS	NO. EVENTS
.455 ± .005	13.0 ± .7	1330

K⁺ PROTON TOTAL INELASTIC CROSS SECTION.

(THIS DATA SHOULD NOT BE USED - A MORE RECENT VALUE MAY BE FOUND IN FISK, ET. AL., CERN CONFERENCE 358 (1961))

LABORATORY BEAM ENERGY GEV	MILLI-BARNS	NO. EVENTS
.455 ± .005	1.0 ± .2	102

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM ENERGY = .455 ± .005 GEV.

(CONTAINS SOME COULOMB EFFECTS)

THIS DATA WAS READ FROM A GRAPH

COS(THETA)		D-SIGMA/D-OMEGA MB/SR
MIN	MAX	
-1.00	-.72	1.13 ± .09
		-.07
-.72	-.44	1.10 ± .08
		-.07
-.44	-.16	.91 ± .07
		-.08
-.16	.12	.90 ± .07
		-.07
.12	.40	.89 ± .06
		-.08
.40	.68	.97 ± .08
		-.08
.68	.96	1.10 ± .09
		-.07

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.

8 K⁺P INTERACTION FROM 140 TO 642 MEV/C. [PHYS. REV. LETTERS 9, 135 (1962)]

S. GOLDHABER, W. CHINOWSKY, G. GOLDHABER, W. LEE, T. A. O'MALLORAN, T. F. STUBBS (U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA, AND UNIV. OF CALIFORNIA, BERKELEY, CALIF., USA)

CROSS REFERENCES
SOME OF THIS DATA RENORMALIZED IN UCRL 18758 (1968).

ADDITIONAL FOOTNOTE REFERENCES
PHYS. REV. 118, 553 (1960), PHYS. REV. LETTERS 7, 188 (1961), AND PHYS. REV. LETTERS 7, 182 (1961).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 1/68, AND VERIFIED BY ROGER BLAND.

BEAM IS K⁺ ON PROTON FROM .140 TO .642 GEV/C.

THIS EXPERIMENT USES THE L.R.L. 15 IN. (H) BUBBLE CHAMBER.

GENERAL COMMENTS ON THIS ARTICLE

1 COULOMB CONTRIBUTIONS ARE REMOVED FROM TOTAL CROSS SECTIONS, BUT NOT FROM ANGULAR DISTRIBUTIONS.

KEY WORDS - CROSS SECTION PHASE SHIFT ANGULAR DISTRIBUTION MODELS

K⁺ PROTON ELASTIC CROSS SECTION.

DATA IS INTEGRATED OVER COS(THETA) FROM -1.00 TO .85. THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.

(INCLUDES COULOMB CONTRIBUTIONS)

LABORATORY BEAM MOMENTUM GEV/C		MILLI-BARNS	
.140 +- .020		14.9 +- 2.5	
.175 .015		16.0	2.4
.205 .015		13.7	1.8
.235 .015		12.7	1.6
.265 .015		11.0	1.6
.355 .025		11.9	1.2
.520 .015		11.9	1.2
.642 .007		11.7	.8

K⁺ PROTON ELASTIC CROSS SECTION.

(NO COS(THETA) CUT.)
COULOMB CORRECTIONS APPLIED

LABORATORY BEAM MOMENTUM GEV/C		MILLI-BARNS	
.140 +- .020		9.2 +- 2.1	
.175 .015		12.5	2.2
.205 .015		11.5	1.7
.235 .015		11.2	1.6
.265 .015		10.0	1.6
.355 .025		11.7	1.2
.520 .015		12.2	1.3
.642 .007		12.4	.9

K⁺ PROTON TOTAL CROSS SECTION.

(NO COS(THETA) CUT. TOTAL AND ELASTIC CROSS SECTIONS ARE THE SAME AT THESE ENERGIES.)
COULOMB CORRECTIONS APPLIED

LABORATORY BEAM MOMENTUM GEV/C		MILLI-BARNS	
.140 +- .020		9.2 +- 2.1	
.175 .015		12.5	2.2
.205 .015		11.5	1.7
.235 .015		11.2	1.6
.265 .015		10.0	1.6
.355 .025		11.7	1.2
.520 .015		12.2	1.3
.642 .007		12.4	.9

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = .14 +- .02 GEV/C.

.....
THIS DATA WAS READ FROM A GRAPH.
.....

COS(THETA)		D-SIGMA/D-OMEGA MB/SR	
MIN	MAX		
-.90	-.60	.9 +- .5	
-.60	-.20	1.1	.5
-.20	.20	.9	.3
.20	.60	1.4	.5
.60	.80	1.5	.7
.80	.95	11.0	2.5

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = .175 ± .015 GEV/C.

THIS DATA WAS READ FROM A GRAPH.

COS(THETA)		D-SIGMA/D-OMEGA MB/SR
MIN	MAX	
-1.00	-.60	1.1 ± .3
-.60	-.20	1.3 .4
-.20	.20	.8 .4
.20	.60	1.6 .4
.60	.80	2.3 .7
.80	.95	5.0 1.2

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = .205 ± .015 GEV/C.

THIS DATA WAS READ FROM A GRAPH.

COS(THETA)		D-SIGMA/D-OMEGA MB/SR
MIN	MAX	
-1.00	-.60	1.0 ± .2
-.60	-.20	1.0 .2
-.20	.20	1.3 .4
.20	.60	1.0 .2
.60	.80	1.5 .4
.80	.95	1.6 .4

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = .235 ± .015 GEV/C.

THIS DATA WAS READ FROM A GRAPH.

COS(THETA)		D-SIGMA/D-OMEGA MB/SR
MIN	MAX	
-1.00	-.60	1.1 ± .2
-.60	-.20	1.2 .2
-.20	.20	.5 .2
.20	.60	1.2 .2
.60	.80	1.0 .2
.80	.95	2.4 .5

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = .265 ± .015 GEV/C.

THIS DATA WAS READ FROM A GRAPH.

COS(THETA)		D-SIGMA/D-OMEGA MB/SR
MIN	MAX	
-1.00	-.60	.9 ± .2
-.60	-.20	.9 .2
-.20	.20	1.0 .2
.20	.60	.8 .2
.60	.80	1.3 .4
.80	.95	3.6 1.2

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = .355 ± .025 GEV/C.

THIS DATA WAS READ FROM A GRAPH.

COS(THETA)		D-SIGMA/D-OMEGA MB/SR
MIN	MAX	
-1.00	-.60	.90 ± .10
-.60	-.20	.80 .10
-.20	.20	1.15 .15
.20	.60	1.00 .15
.60	.95	1.25 .20

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.LABORATORY BEAM MOMENTUM = .520 $\pm$.015 GEV/C.

 . THIS DATA WAS READ FROM A GRAPH .

COS(THETA)		D-SIGMA/D-OMEGA MB/SR	
MIN	MAX		
-1.00	-.60	.90 $\pm$.15	
-.60	-.20	1.00	.15
-.20	.20	1.10	.15
.20	.60	1.10	.15
.60	.95	1.00	.15

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.LABORATORY BEAM MOMENTUM = .642 $\pm$.007 GEV/C.

 . THIS DATA WAS READ FROM A GRAPH .

COS(THETA)		D-SIGMA/D-OMEGA MB/SR	
MIN	MAX		
-1.00	-.60	1.10 $\pm$.12	
-.60	-.20	.90	.12
-.20	.20	.95	.12
.20	.60	.95	.12
.60	.95	1.05	.12

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.

9 EVIDENCE AGAINST THE EXISTENCE OF A STRANGENESS +2 MESON RESONANCE AT A MASS OF 1280 MEV. (PHYS. REV. LETTERS 17, 716 (1966))

T. JOLDERSMA, R. B. PALMER, N. P. SAMIOS (BROOKHAVEN NAT. LAB., UPTON, L.I., N. Y., USA)

ADDITIONAL FOOTNOTE REFERENCES

PHYS. LETTERS 17, 155 (1965), BNL H-10 (1962), PHYS. REV. 134, B383 (1964), AND PHYS. REV. LETTERS 16, 1063 (1966).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 4/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K⁺ ON PROTON AT 3 GEV/C.

THIS EXPERIMENT USES THE B.N.L. 80 IN. (H) BUBBLE CHAMBER. A TOTAL OF 12000 PICTURES ARE REPORTED ON.

GENERAL COMMENTS ON THIS ARTICLE

I LOOKS FOR BUT SEES NO K⁺K⁺ RESONANCE

KEY WORDS = CROSS SECTION DALITZ PLOT MASS SPECTRUM

CROSS SECTION FOR K⁺ PROTON $\rightarrow$ K⁺ K⁺ LAMBDA.

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
3.	22.3 $\pm$ 3.3	71

CROSS SECTION FOR K⁺ PROTON $\rightarrow$ K⁺ K⁺ SIGMA0.

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
3.	9.8 $\pm$ 2.0	31

CROSS SECTION FOR K⁺ PROTON $\rightarrow$ K⁺ K⁰ SIGMA+.

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
3.	19.4 $\pm$ 4.7	51

CROSS SECTION FOR K⁺ PROTON $\rightarrow$ K⁺ K⁺ LAMBDA PI0.

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
3.	4.7 $\pm$ 1.3	15

CROSS SECTION FOR K⁺ PROTON $\rightarrow$ K⁺ K⁰ LAMBDA PI+.

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
3.	9.1 $\pm$ 1.9	29

10 INTERACTIONS OF 900 MEV/C K⁺ ON NUCLEI LEADING TO THE PRODUCTION OF ONE PION. (NUOVO CIMENTO 35, 319 (1965))

E. FIORINI, A. PULLIA (UNIV. DI MILANO, MILANO, ITALY)
A. BEZAGUET, P. MUSSET (ECOLE POLYTECHNIQUE, PARIS, FRANCE)

ADDITIONAL FOOTNOTE REFERENCES

CERN TC-62-26 (1962), REVIEW OF SCIENTIFIC INSTRUMENTS 32, 1302 (1961), PHYS. LETTERS 5, 279 (1963), PHYS. REV. 135, 8484 (1964), PHYS. REV. 135, 8495 (1964), SIENNA CONFERENCE 1 125 (1963), DUBNA CONFERENCE PAPTR. TALK (1964), AND PHYS. REV. LETTERS 11, 93 (1963).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 3/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K⁺ ON HEAVY LIQUID FROM .77 TO .95 GEV/C.

THIS EXPERIMENT USES A HEAVY LIQUID BUBBLE CHAMBER. A TOTAL OF 70000 PICTURES ARE REPORTED ON.

GENERAL COMMENTS ON THIS ARTICLE

- 1 NO DATA PUNCHED BECAUSE OF LOW STATISTICS AND 200 MEV/C BEAM SPREAD
- 2 THEY QUOTE UPPER LIMIT FOR KAPPA CROSS SECTION.

KEY WORDS = ANGULAR DISTRIBUTION KAPPA MESON CROSS SECTION

* NO DATA PUNCHED FOR THIS ARTICLE *

11 EVIDENCE FOR A MASS ENHANCEMENT IN THE ANTI-LAMBDA N SYSTEM. (PHYS. REV. LETTERS 20, 755 (1968))

G. ALEXANDER, A. FIRESTONE, G. GOLDBERGER, B. C. SHEN (U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA)

CROSS REFERENCES

SEE ALSO NUOVO CIMENTO 48A, 589 (1967), PHYS. LETTERS 25B, 246 (1967), AND UNIV. OF ROCHESTER 875-218 (1967).

ADDITIONAL FOOTNOTE REFERENCES

PHYS. REV. LETTERS 17, 890 (1966), PHYS. REV. LETTERS 18, 1209 (1967), HEIDELBERG CONFERENCE PAPTR. TALK (1967), NUOVO CIMENTO 34, 1644 (1964), AND PHYS. REV. LETTERS 19, 972 (1967).

ARTICLE READ BY ODETTE BENARY IN 12/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K⁺ ON PROTON AT 9 GEV/C.

THIS EXPERIMENT USES THE B.N.L. 80 IN. (H) BUBBLE CHAMBER. A TOTAL OF 90000 PICTURES ARE REPORTED ON.

GENERAL COMMENTS ON THIS ARTICLE

- 1 REPORTS A LAMBDA-BAR NUCLEON ENHANCEMENT AT 2240 MEV WITH A WIDTH OF 70 MEV.

KEY WORDS = CROSS SECTION ANGULAR DISTRIBUTION MASS SPECTRUM ANTI-LAMBDA ANTI-SIGMA0

CROSS SECTION FOR K⁺ PROTON = ANTI-LAMBDA PROTON PROTON + ANTI-SIGMA0 PROTON PROTON. (PAGE 755)

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
9.	11.4 ± 2.3	25

CROSS SECTION FOR K⁺ PROTON = ANTI-LAMBDA PROTON NEUTRON PI⁺. (PAGE 755)

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
9.	35.2 ± 4.0	76

CROSS-SECTION FOR K⁺ PROTON = ANTI-LAMBDA PROTON PROTON PI⁰. (PAGE 755)

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
9.	9. ± 2.	21

CROSS SECTION FOR K⁺ PROTON = ANTI-LAMBDA PROTON PROTON PI⁺ PI⁻. (PAGE 755)

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
9.	2.7 ± 1.1	6

12 OBSERVATION OF CORRELATIONS BETWEEN VECTOR-MESON AND BARYON-RESONANCE DECAYS. (PHYS. LETTERS 18, 76 (1965))

G.GOLDHABER, J.L.BROWN, I.BUTTERWORTH, S.GOLDHABER, A.A.HIRATA, J.A.KADYK, B.C.SHEN, G.H.TRILLING (U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA, AND UNIV. OF CALIFORNIA, BERKELEY, CALIF., USA)

ADDITIONAL FOOTNOTE REFERENCES

PHYS. REV. 138, 8997 (1965), PHYS. LETTERS 6, 52 (1963), ATHENS CONFERENCE 130 (1963), NUOVO CIMENTO 24, 453 (1952), NUOVO CIMENTO 34, 735 (1964), PHYS. REV. LETTERS 12, 399 (1964), PHYS. REV. LETTERS 13, 456 (1964), NUCLEAR INSTRUMENTS AND METHODS 20, 37 (1963), PHYS. REV. LETTERS 12, 336 (1964), PHYS. REV. LETTERS 9, 330 (1962), PHYS. REV. 135, 8508 (1964), AND CORAL GABLES CONFERENCE 34 (1965).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 3/68, AND VERIFIED BY LEROY PRICE.

BEAM NO. 1 IS K⁺ ON PROTON AT 1.96 GEV/C.
NO. 2 IS K⁺ ON DEUTERON AT 2.3 GEV/C.
NO. 3 IS PI⁺ ON PROTON AT 3.65 GEV/C.

THIS EXPERIMENT USES THE B.N.L. 20 IN. (H) BUBBLE CHAMBER.

KEY WORDS - ANGULAR DISTRIBUTION K*(890) RHO MESON DELTA(1238) MASS SPECTRUM

* NO DATA PUNCHED FOR THIS ARTICLE *

13 K⁺P PHASE-SHIFT ANALYSIS BELOW 1500 MEV/C. (PHYS. REV. 165, 1770 (1968))

A.T.LEA (RUTHERFORD HIGH EN. LAB., CHILTON, DIO, BERK. ENGLAND)
B.R.MARTIN (BROOKHAVEN NAT. LAB., UPTON, L.I., N. Y., USA)
G.C.DADES (EUROPEAN ORG. FOR NUC. RES., GENEVA, SWITZERLAND)

ABSTRACT AN EXTENSIVE SYSTEMATIC ENERGY-DEPENDENT PHASE-SHIFT ANALYSIS IN THE KAON LABORATORY MOMENTUM RANGE 140-1435 MEV/C HAS BEEN MADE OF K⁺P SCATTERING DATA, SUPPLEMENTED BY INFORMATION FROM K⁺P FORWARD DISPERSION RELATIONS. S, P, AND D WAVES WERE INCLUDED AND VARIOUS ASSUMPTIONS WERE MADE CONCERNING THE SOURCE OF THE INELASTICITY. RESULTS ARE PRESENTED FOR SOLUTIONS ALLOWING ALL POSSIBLE COMBINATIONS OF UP TO THREE PARTIAL WAVES TO BE INELASTIC. FOUR BASIC TYPES OF SOLUTION ARE FOUND, AND EXPERIMENTS ARE SUGGESTED WHICH WILL BEST HELP TO DECIDE AMONG THEM. TWO OF THESE GROUPS OF SOLUTIONS SHOW SOME EVIDENCE FOR A RESONANCE IN THE P 1/2 PARTIAL WAVE, ALTHOUGH ITS PRECISE CHARACTERISTICS CANNOT BE DETERMINED WITH PRESENT DATA.

CROSS REFERENCES

SEE ALSO PHYS. REV. LETTERS 21, 1286 (1968).
THIS ARTICLE SUPERSEDES PART OF PHYS. LETTERS 23, 280 (1966).
ANALYSIS OF DATA FROM PHYS. REV. LETTERS 17, 102 (1966), PHYS. REV. 118, 553 (1960), PHYS. REV. LETTERS 9, 135 (1962), PHYS. REV. LETTERS 7, 188 (1961), PHYS. REV. LETTERS 7, 182 (1961), PHYS. REV. 129, 2743 (1963), PHYS. LETTERS 16, 83 (1965), PHYS. REV. 139, 8141 (1965), PHYS. LETTERS 24, 8314 (1967), PHYS. REV. 135, 8191 (1964), AND UCRL 17388 (1967).

ADDITIONAL FOOTNOTE REFERENCES

PHYS. REV. 139, 8156 (1965), PHYS. LETTERS 19, 420 (1965), PHYS. LETTERS 12, 76 (1964), PHYS. LETTERS 20, 306 (1966), PHYS. REV. LETTERS 16, 714 (1966), REV. MOD. PHYS. 35, 737 (1963), PHYS. REV. 129, 2311 (1963), NUOVO CIMENTO 12, 241 (1959), PHYS. REV. 139, 8141 (1965), PHYS. REV. LETTERS 17, 939 (1966), PHYS. REV. 155, 1744 (1967), PHYS. REV. 138, 8913 (1965), PHYS. REV. 129, 2285 (1963), PHYS. REV. 132, 2721 (1963), PHYS. REV. 131, 2248 (1963), PHYS. REV. 125, 1696 (1962), PHYS. REV. LETTERS 14, 29 (1965), PHYS. REV. 139, 8719 (1965), PHYS. LETTERS 21, 229 (1966), PHYS. REV. LETTERS 7, 204 (1961), PHYS. REV. 113, 338 (1959), PHYS. REV. LETTERS 19, 259 (1967), PHYS. REV. LETTERS 19, 255 (1967), PHYS. REV. 123, 2168 (1961), PHYS. REV. 139, 81336 (1965), ANNALS OF PHYSICS 10, 307 (1960), PHYS. REV. 138, 8947 (1965), NUOVO CIMENTO 50, A493 (1967), PROGRESS OF THEORETICAL PHYSICS 37, 398 (1967), AND PROGRESS OF THEORETICAL PHYSICS 11, 213 (1954).

ARTICLE READ BY ODETTE BEVARY IN 12/68, AND VERIFIED BY LEROY PRICE.

KEY WORDS - PHASE SHIFT

* NO DATA PUNCHED FOR THIS ARTICLE *

14 K⁺P SCATTERING AND A POSSIBLE Z⁺ RESONANCE. (PHYS. REV. LETTERS 21, 1286 (1968))

B.R.MARTIN (BROOKHAVEN NAT. LAB., UPTON, L.I., N. Y., USA)

ABSTRACT AN ENERGY-DEPENDENT PHASE-SHIFT ANALYSIS IN THE KAON LABORATORY-MOMENTUM RANGE BELOW 2.0 BEV/C HAS BEEN MADE OF K⁺P SCATTERING DATA, SUPPLEMENTED BY INFORMATION FROM K⁺P FORWARD DISPERSION RELATIONS. EVIDENCE FOR A POSSIBLE RESONANCE IN THE P 1/2 STATE IS PRESENTED.

CROSS REFERENCES

SEE ALSO PHYS. LETTERS 23, 380 (1966), AND PHYS. REV. 165, 1770 (1968).
ANALYSIS OF DATA FROM PHYS. REV. LETTERS 17, 102 (1966), PHYS. REV. 118, 553 (1960), PHYS. REV. LETTERS 9, 135 (1962), PHYS. REV. LETTERS 7, 188 (1961), PHYS. REV. LETTERS 7, 182 (1961), PHYS. REV. 129, 2743 (1963), PHYS. LETTERS 16, 83 (1965), PHYS. REV. 139, 8141 (1965), PHYS. LETTERS 24, 8314 (1967), PHYS. REV. 135, 8191 (1964), UCRL 17388, PHYS. REV. LETTERS 19, 259 (1967), PHYS. REV. 168, 1466 (1968), AND PHYS. REV. LETTERS 21, 1202 (1968).

ADDITIONAL FOOTNOTE REFERENCES

PHYS. REV. LETTERS 20, 689 (1968), PHYS. LETTERS 278, 22 (1968), AND PHYS. REV. 139, 8156 (1965).

ARTICLE READ BY ODETTE BEVARY IN 12/68, AND VERIFIED BY LEROY PRICE.

KEY WORDS - PHASE SHIFT

* NO DATA PUNCHED FOR THIS ARTICLE *

15 EVIDENCE FOR A MESON RESONANCE WITH STRANGENESS +2. (PHYS. LETTERS 17, 155 (1965))

M. FERRO-LUZZI, R. GEORGE, Y. GOLDSCHMIDT-CLERMONT, V. P. HENRI, B. JONGEJANS, D. W. G. LEITH, G. R. LYNCH, F. MULLER, J. M. PERREAU (EUROPEAN ORG. FOR NUC. RES., GENEVA, SWITZERLAND)

ADDITIONAL FOOTNOTE REFERENCES
SIENNA CONFERENCE 182 (1963), AND CERN TC/65-10 (1965).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 5/68, AND VERIFIED BY ROGER BLAND.

BEAM IS K+ ON PROTON FROM 2.97 TO 4.97 GEV/C.

THIS EXPERIMENT USES THE SACLAY 81 CM (H) BUBBLE CHAMBER. A TOTAL OF 400000 PICTURES ARE REPORTED ON.

GENERAL COMMENTS ON THIS ARTICLE
1 SEE EVIDENCE FOR K+K+ RESONANCES AT 1090 AND 1280 MEV/C

KEY WORDS = CROSS SECTION MASS SPECTRUM

CROSS SECTION FOR K+ PROTON = K+ K+ LAMBDA.

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
2.97	23.0 +- 4.5	44
3.46	22.0 4.5	47
4.97	18.0 6.0	14

CROSS SECTION FOR K+ PROTON = K+ K+ SIGMA0.

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
2.97	7.5 +- 2.5	13
3.46	7.0 2.5	9
4.97	6.0 3.0	4

CROSS SECTION FOR K+ PROTON = K+ K0 SIGMA+.

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
2.97	15. +- 5.	11
3.46	9. 4.	12
4.97	23. 9.	7

CROSS SECTION FOR K+ PROTON = K+ K+ LAMBDA PI0.

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
2.97	7.5 +- 2.5	11
3.46	9.5 3.0	20
4.97	30.0 8.0	26

CROSS SECTION FOR K+ PROTON = K+ K0 LAMBDA PI+.

(2.46 GEV/C VALUE NOT USED SINCE THERE APPEARS TO BE A TYPOGRAPHICAL ERROR IN THE ORIGINAL TABLE.)

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
2.97	12.5 +- 3.5	23
4.97	36.0 9.0	36

16 K+P PHASE SHIFT ANALYSIS. (PHYS. LETTERS 23, 380 (1968))

A. T. LEA, B. R. MARTIN (NIELS BOHR INSTITUTE, COPENHAGEN, DENMARK)
G. C. DADES (NORDISK INS. FOR TEOR. ATOMFYS., COPENHAGEN, DENMARK)

ABSTRACT IN A PRELIMINARY PHASE SHIFT ANALYSIS OF K+P DATA WE HAVE FOUND POSSIBLE EVIDENCE FOR A STRANGENESS +1 BARYON RESONANCE.

CROSS REFERENCES

PART OF THIS ARTICLE SUPERSEDED BY PHYS. REV. 165, 1770 (1968).
ANALYSIS OF DATA FROM PHYS. REV. LETTERS 4, 155 (1962), PHYS. REV. LETTERS 7, 182 (1961), PHYS. REV. 129, 2743 (1963),
PHYS. REV. LETTERS 7, 188 (1961), PHYS. REV. 135B, 191 (1964), PHYS. REV. LETTERS 2, 117 (1959), PHYS. LETTERS 15, 83
(1965), AND PHYS. REV. 118, 553 (1960).

ADDITIONAL FOOTNOTE REFERENCES

ANNALS OF PHYSICS 10, 307 (1960), PHYS. REV. LETTERS 14, 29 (1965), PHYS. REV. 139, 719 (1965), PHYS. LETTERS 21, 229
(1966), AND PHYS. REV. LETTERS 17, 102 (1966).

ARTICLE READ BY DOETTE BENARY IN 12/68, AND VERIFIED BY LEROY PRICE.

KEY WORDS = PHASE SHIFT

* NO DATA PINCHED FOR THIS ARTICLE *

- 17 RESONANT STATES IN THE $K^+PI^+PI^-P$ SYSTEM. [PHYS. LETTERS 16, 184 (1965)]
 S.P. ALMEIDA, M.W. ATHERTON, T.A. BYER, P.J. DORMAN, A.G. FORSON, J.M. SCHARENGUIVEL, D.M. SENDALL, B.A. WESTWOOD (CAVENDISH LAB., CAMB. UNIV., CAMBRIDGE, ENGLAND)
 ADDITIONAL FOOTNOTE REFERENCES
 CERN TC/02/1963, PHYS. REV. LETTERS 9, 330 (1962), SIENNA CONFERENCE I 189, NUOVO CIMENTO 33, 309 (1964), PHYS. LETTERS 9, 71 (1964), AND PHYS. LETTERS 9, 207 (1964).
 ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 3/68, AND VERIFIED BY LEROY PRICE.
 BEAM IS K^+ ON PROTON AT 5 GEV/C.
 THIS EXPERIMENT USES THE SACLAY 81 CM (H) BUBBLE CHAMBER.
 GENERAL COMMENTS ON THIS ARTICLE
 1 NO CROSS SECTIONS GIVEN. STUDIES $K^*(1320)$ $N^*(1238)$ PRODUCTION IN $K^+PI^+PI^-$ PROTON FINAL STATE.
 KEY WORDS = MASS SPECTRUM ANGULAR DISTRIBUTION DALITZ PLOT ONE PARTICLE EXCHANGE DELTA(1238)
 $K^*(890)$ $K^*(1320)$

 * * * * *
 * NO DATA PUNCHED FOR THIS ARTICLE *
 * * * * *
- 18 ANGULAR CORRELATIONS IN THE DECAY OF THE K^* AND N^* STATES. [PHYS. LETTERS 6, 309 (1963)]
 G.B. CHADWICK, D.J. CRENNELL, M.T. DAVIES, M. DERRICK, J.H. MULVEY, P.B. JONES, D. RADOJICIC, C.A. WILKINSUN (OXFORD UNIV., OXFORD, ENGLAND)
 A. BETTINI, M. CRESTI, S. LIMENTANI, L. PERUZZO, R. SANTANGELO (UNIV. OF PADOVA, PADOVA, ITALY)
 CROSS REFERENCES
 PART OF THIS ARTICLE SUPERSEDED BY PHYS. LETTERS 16, 83 (1965).
 ADDITIONAL FOOTNOTE REFERENCES
 PHYS. REV. LETTERS 10, 368 (1963), PHYS. REV. LETTERS 10, 371 (1963), NUOVO CIMENTO 27, 1450 (1963), PHYS. REV. LETTERS 11, 90 (1963), NATIONAL INST. FOR NUCLEAR RESEARCH NTL/M/38 (1962), PHYS. LETTERS 3, 283 (1963), PHYS. REV. LETTERS 6, 300 (1961), CERN CONFERENCE 295 (1962), PHYS. REV. LETTERS 9, 330 (1962), PHYS. REV. LETTERS 10, 138 (1963), PHYS. REV. 129, 1379 (1963), PHYS. REV. LETTERS 8, 140 (1962), AND PHYS. REV. LETTERS 11, 93 (1963).
 ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 5/68, AND VERIFIED BY LEROY PRICE.
 BEAM IS K^+ ON PROTON AT 1.455 GEV/C.
 THIS EXPERIMENT USES THE CERN 32 CM (H) BUBBLE CHAMBER.
 KEY WORDS = ANGULAR DISTRIBUTION ONE PARTICLE EXCHANGE

 * * * * *
 * NO DATA PUNCHED FOR THIS ARTICLE *
 * * * * *
- 19 INELASTIC PROCESSES NEAR THE $T=1$ K^+P PEAK AT 1250 MEV/C. [PHYS. REV. LETTERS 18, 1077 (1967)]
 R.W. BLAND, M.G. BOWLER, J.L. BROWN, G. GOLDBABER, S. GOLDBABER, V.H. SEEGER, G.M. TRILLING (U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA, AND UNIV. OF CALIFORNIA, BERKELEY, CALIF., USA)
 ABSTRACT K^+P INELASTIC FINAL STATES IN THE REGION OF THE 1250-MEV/C $NUCL$ PEAK ARE EXAMINED FOR EVIDENCE OF RESONANCE BEHAVIOR. THE DOMINANT KN^* FINAL STATE IS FOUND TO BE ABOUT EQUALLY DIVIDED BETWEEN $P 1/2$ AND $P 3/2$ FINAL STATES, WITH NO RAPID PHASE VARIATION BETWEEN 960 AND 1370 MEV/C.
 CROSS REFERENCES
 DATA SUPERSEDED BY NUCRL 10131 (1967).
 ADDITIONAL FOOTNOTE REFERENCES
 PHYS. REV. LETTERS 17, 102 (1966), PHYS. REV. LETTERS 17, 920 (1966), PHYS. REV. LETTERS 9, 135 (1962), PHYS. REV. LETTERS 7, 188 (1951), PHYS. REV. 133, 822 (1964), PHYS. LETTERS 16, 83 (1965), PHYS. REV. 142, 913 (1966), PHYS. REV. 139B, 1411 (1965), PHYS. REV. LETTERS 11, 90 (1963), NUOVO CIMENTO 35, 1101 (1965), NUOVO CIMENTO 33, 906 (1964), AND PHYS. REV. LETTERS 18, 1081 (1967).
 ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 4/68, AND VERIFIED BY LEROY PRICE.
 BEAM IS K^+ ON PROTON FROM .86 TO 1.58 GEV/C.
 THIS EXPERIMENT USES THE L.R.L. 25 IN. (H) BUBBLE CHAMBER.
 KEY WORDS = $K^*(890)$ DELTA(1238) ANGULAR DISTRIBUTION

 * * * * *
 * NO DATA PUNCHED FOR THIS ARTICLE *
 * * * * *
- 20 THE PRODUCTION MECHANISM FOR THE REACTION $K^+P \rightarrow K^+P$ AT 3 GEV/C. [PHYS. LETTERS 9, 359 (1964)]
 G.R. LYNCH, M. FERRO-LUZZI, R. GEORGE, V. GOLDSCHMIDT-CLERMONT, V.P. HENRI, B. JONGEJANS, D.W.G. LEITH, F. MULLER, J.M. PERREAU (EUROPEAN ORG. FOR NUC. RES., GENEVA, SWITZERLAND)
 CROSS REFERENCES
 DATA SUPERSEDED BY NUOVO CIMENTO 36, 1101 (1965).
 ADDITIONAL FOOTNOTE REFERENCES
 CERN 63-12, PHYS. REV. LETTERS 10, 138 (1963), PHYS. LETTERS 8, 144 (1964), PHYS. REV. LETTERS 8, 140 (1962), PHYS. LETTERS 6, 62 (1963), AND PHYS. LETTERS 6, 309 (1963).
 ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 3/68, AND VERIFIED BY LEROY PRICE.
 BEAM IS K^+ ON PROTON AT 2.965 GEV/C.
 THIS EXPERIMENT USES THE SACLAY 81 CM (H) BUBBLE CHAMBER.
 KEY WORDS = $K^*(890)$ DELTA(1238) DENSITY MATRIX ANGULAR DISTRIBUTION ONE PARTICLE EXCHANGE

 * * * * *
 * NO DATA PUNCHED FOR THIS ARTICLE *
 * * * * *

21 ISOBAR PRODUCTION BY 910-MEV/C K^+ MESONS. [PHYS. REV. LETTERS 11, 93 (1963)]

B. KEHO (UNIV. OF MARYLAND, COLLEGE PARK, MD., USA)

ADDITIONAL FOOTNOTE REFERENCES

PHYS. REV. LETTERS 11, 93 (1963), NUOVO CIMENTO 16, 1073 (1960), AND PHYS. REV. LETTERS 8, 447 (1962).

ARTICLE READ BY NAOMI BARASH-SCHMIOT IN 5/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K^+ ON HEAVY LIQUID AT .91 GEV/C.

THIS EXPERIMENT USES THE L.R.L. 30 IN. (HL) BUBBLE CHAMBER. A TOTAL OF 40000 PICTURES ARE REPORTED ON.

KEY WORDS = CROSS SECTION ONE PARTICLE EXCHANGE ANGULAR DISTRIBUTION MODELS

CROSS SECTION FOR K^+ PROTON $\rightarrow$ K^0 PROTON π^+ .

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.91 $\pm$.04	2.1 $\pm$.2

DIFFERENTIAL CROSS SECTION FOR K^+ PROTON.

(PROD. ANG DIST FOR 3-BODY STATE - NO DATA PUNCHED FOR THIS PARAMETER.)

22 EVIDENCE FOR THREE $K \pi \pi$ RESONANT STATES BELOW 1.4 GEV IN K^+P REACTIONS AT 5 GEV/C. [PHYS. LETTERS 268, 30 (1967)]

G. BASSOMPIERRE, Y. GOLDSCHMIDT-CLERMONT, A. GRANT, V. P. HENRI, I. HUGHES, B. JONGEJANS, R. L. LANDER, D. LINGLIN, F. MULLER, J. M. PERREAU, I. SAITOV, R. L. SEKU, L. V. G. WOLF (EUROPEAN ORG. FOR NUC. RES., GENEVA, SWITZERLAND)
W. DEBAERE, J. DEBAISIEUX, P. DUFOR, F. GRARD, J. HEUGHEBAERT, L. PAPE, P. PEETERS, F. VERREURE, R. WINDMOLDERS (INST. INTERUNIV. DES SCI. NUC., BRUXELLES, BELGIUM)
M. JOES, W. WATT (BIRMINGHAM UNIV., BIRMINGHAM, ENGLAND)

ABSTRACT THE STUDY OF K^+P REACTIONS AT AN INCIDENT MOMENTUM OF 4.97 GEV/C SUGGESTS THAT THE WIDE ENHANCEMENT IN THE ($K \pi \pi$) $^+$ MASS SPECTRUM MAY BE SPLIT INTO SEVERAL RESONANT STATES. IN PARTICULAR, EXAMINATION OF THE $K^*(1892)\pi^+$ DECAY MODE INDICATES THE EXISTENCE OF THE STATES $K^*(1230)$, $K^*(1280)$ AND $K^*(1320)$, IN ADDITION TO THE WELL KNOWN $K^*(1420)$. THE $K^*(1230)$ AND $K^*(1320)$ HAVE THE PROPERTIES EXPECTED FROM 1^+ STATES PRODUCED BY 0^+ EXCHANGE.

CROSS REFERENCES

PART OF THIS ARTICLE SUPERSEDED BY PHYS. LETTERS 278, 468 (1968).
CONTINUATION OF PREVIOUS EXPERIMENT IN NUOVO CIMENTO 49A, 9 (1967).

ADDITIONAL FOOTNOTE REFERENCES

NUOVO CIMENTO 49, 9 (1967), NUOVO CIMENTO 49, 373 (1967), PHYS. REV. LETTERS 17, 726 (1966), PHYS. REV. LETTERS 18, 1087 (1967), HEIDELBERG CONFERENCE RAPTR. TALK (1967), PHYS. REV. LETTERS 19, 546 (1967), PHYS. REV. LETTERS 19, 271 (1967), PHYS. REV. LETTERS 19, 972 (1967), PHYS. LETTERS 9, 207 (1964), AND PHYS. REV. LETTERS 19, 44 (1967).

ARTICLE READ BY ODETTE BENARY IN 12/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K^+ ON PROTON AT 4.97 GEV/C.

THIS EXPERIMENT USES THE CERN 2M (H) BUBBLE CHAMBER.

KEY WORDS = CROSS SECTION MASS SPECTRUM ANGULAR DISTRIBUTION ϕ PEAK

CROSS SECTION FOR K^+ PROTON $\rightarrow$ K^+ PROTON π^+ π^- . [PAGE 30]

(THIS DATA SHOULD NOT BE USED - A MORE RECENT VALUE MAY BE FOUND IN BASSOMPIERRE, ET. AL., PHYS. LETTERS 278, 468 (1968))

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS
4.97	1.80 $\pm$.15	6347

CROSS SECTION FOR K^+ PROTON $\rightarrow$ K^0 PROTON π^+ π^0 . [PAGE 30]

(THIS DATA SHOULD NOT BE USED - A MORE RECENT VALUE MAY BE FOUND IN BASSOMPIERRE, ET. AL., PHYS. LETTERS 278, 468 (1968))

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS
4.97	2.3 $\pm$.3	3075

CROSS SECTION FOR K^+ PROTON $\rightarrow$ K^0 NEUTRON π^+ π^+ . [PAGE 30]

(THIS DATA SHOULD NOT BE USED - A MORE RECENT VALUE MAY BE FOUND IN BASSOMPIERRE, ET. AL., PHYS. LETTERS 278, 468 (1968))

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS
4.97	.6 $\pm$.1	662

23 OBSERVATION OF THE REACTION $K^+ p \rightarrow p p \bar{\Lambda}$ (NUOVO CIMENTO 48, 589 (1967))

G. BASSOMPIERRE, Y. GOLOSCHMIDT-CLERMONT, A. GRANT, V. P. HENRI, B. JONCE JANS, R. L. LANDER, D. LINGLIN, F. MÜLLER, J. M. PERREAU, A. PROKES, R. L. SEKUL IN, J. K. TOUMINIEMI (EUROPEAN ORG. FOR NUC. RES., GENEVA, SWITZERLAND)
W. DEBAERE, J. DEBAISIEUX, P. DUFOUR, F. GRARD, J. HEUGHEBAERT, L. PAPE, P. PEETERS, F. VERBEURE, R. WINDMOLERS (INST. INTERUNIV. DES SCI. NUC., BRUXELLES, BELGIUM)

ADDITIONAL FOOTNOTE REFERENCES
NUOVO CIMENTO 132, 1273 (1963), NUOVO CIMENTO 10, 635 (1958), JETP 11 726 (1960), JETP 11 976 (1960), PHYS. LETTERS 21, 471 (1966), AND CERN TC/02-63-3.

ARTICLE READ BY LEROY PRICE IN 10/68, AND VERIFIED BY ROGER BLAND.

BEAM IS K^+ ON PROTON AT 4.97 GEV/C.

THIS EXPERIMENT USES A HYDROGEN BUBBLE CHAMBER. A TOTAL OF 20000 PICTURES ARE REPORTED ON.

GENERAL COMMENTS ON THIS ARTICLE
1 NORMALIZED TO TAU DECAYS. BRANCHING RATIO OF TAU/ALL = .055.
2 K^+ LIFETIME = 1.23×10^{-8} SEC.

KEY WORDS = DALITZ PLOT CROSS SECTION

CROSS SECTION FOR $K^+ p \rightarrow p p \bar{\Lambda}$. (PAGE 590)

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
4.97	3.2 ± 1.0	14

CROSS SECTION FOR $K^+ p \rightarrow p p \bar{\Lambda}$. (PAGE 590)

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
4.97	$.6 \pm .6$	2

DIFFERENTIAL CROSS SECTION FOR $K^+ p \rightarrow p p \bar{\Lambda}$. (FIGURE 2)

LABORATORY BEAM MOMENTUM = 4.97 GEV/C.

COS(THETA)	NO. EVENTS	
MIN	MAX	
-1.0	-.8	2
-.8	-.6	2
-.6	-.4	2
-.4	-.2	1
-.2	.0	2
.0	.2	0
.2	.4	2
.4	.6	1
.6	.8	1
.8	1.0	0

THETA IS THE ANGLE THAT THE ANTI-LAMBDA MAKES WITH THE BEAM IN THE GRAND C.M.

24 K-MESON-NUCLEON TOTAL CROSS SECTIONS FROM 0.6 TO 2.0 BEV. (PHYS. REV. LETTERS 2, 117 (1959))

H. C. BURROWES, D. O. CALDWELL, D. H. FRISCH, D. A. HILL, D. M. RITSON, R. A. SCHLUTER (MASSACHUSETTS INST. OF TECH., CAMBRIDGE, MASS., USA)

ADDITIONAL FOOTNOTE REFERENCES
PHYS. REV. 109, 2121 (1958), PHYS. REV. 107, 248 (1957), PHYS. REV. 100, 242 (1955), PHYS. REV. 110, 569 (1958),
PHYS. REV. 110, 572 (1958), PROGRESS IN THEORETICAL PHYSICS, JAPAN 19, 238 (1958), AND UCRL 8461 (1958).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 6/68, AND VERIFIED BY LEROY PRICE.

BEAM NO. 1 IS K^+ ON PROTON FROM .98 TO 2.45 GEV/C.
NO. 2 IS K^+ ON DEUTERON FROM .98 TO 2.45 GEV/C.

THIS EXPERIMENT USES COUNTERS.

KEY WORDS = CROSS SECTION

K^+ PROTON TOTAL CROSS SECTION.

.....
THIS DATA WAS READ FROM A GRAPH.
.....

C.M. ENERGY GEV	MILLI-BARNS
1.776	17.5 ± 1.5
1.823	19.6
1.964	18.8
2.027	16.3
2.089	18.3
2.183	15.5
2.293	15.4
2.387	13.0

K^+ NEUTRON TOTAL CROSS SECTION.

(ERRORS MUCH LARGER THAN RECENT MEASUREMENTS - NO DATA PUNCHED FOR THIS PARAMETER.)

25 ANALYSIS OF THE K11301 SYSTEM PRODUCED IN 5.5-GEV/C K⁺P INTERACTIONS. [PHYS. REV. LETTERS 20, 1519 (1968)]

F. BOMSE, S. BORENSTEIN, A. CALLAHAN, J. COLE, B. COX, D. ELLIS, L. ETTINGER, D. GILLESPIE, G. LUSTE, R. MERCER, E. MOSES, A. PEVSNER, R. ZDANIS
(JOHNS HOPKINS UNIV., BALTIMORE, MD., USA)

ADDITIONAL FOOTNOTE REFERENCES

PHYS. REV. LETTERS 18, 1087 (1967), PHYS. REV. LETTERS 19, 972 (1967), PHYS. REV. LETTERS 16, 1069 (1966), PHYS. REV. LETTERS 17, 726 (1966), PHYS. LETTERS 15, 184 (1965), NUOVO CIMENTO 49A, 373 (1957), PHYS. LETTERS 26B, 30 (1967), PHYS. REV. LETTERS 19, 271 (1967), PHYS. REV. LETTERS 20, 171 (1968), PHYS. REV. LETTERS 19, 44 (1967), PHYS. LETTERS 22, 357 (1966), PHYS. REV. LETTERS 19, 546 (1967), PHYS. REV. 136, 1207 (1964), PHYS. REV. 129, 1057 (1962), BERKELEY CONFERENCE 215 (1966), AND SLAC 43 (1965).

ARTICLE READ BY ODETTE BENARY IN 12/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K⁺ ON PROTON AT 5.438 GEV/C.

THIS EXPERIMENT USES THE 8.N.L. 80 IN. (H) BUBBLE CHAMBER.

KEY WORDS = CROSS SECTION MASS SPECTRUM MODELS DENSITY MATRIX ANGULAR DISTRIBUTION Q PEAK
SPIN AND PARITY

CROSS SECTION FOR K⁺ PROTON = K⁺ PI⁻ PI⁺ PROTON. [PAGE 1519]

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS
5.438 ± .025	1.79 ± .10	6500

CROSS SECTION FOR K⁺ PROTON = K⁰ PI⁺ PI⁰ PROTON. [PAGE 1519]

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS
5.438 ± .025	1.45 ± .15	900

26 K⁺P INTERACTIONS AT 13 GEV/C. [CERN HADRON CONFERENCE 172 (1968)]

J.C. BERLINCHERI, M.S. FARBER, T. FERBEL, B.E. FORMAN, A.C. MELISSINOS, P.F. SLATTERY, T. YAMAMUCHI, H. YUTA (UNIV. OF ROCHESTER, ROCHESTER, N. Y., USA)

CROSS REFERENCES

PART OF THIS ARTICLE SUPERSEDED BY PHYS. LETTERS 27B, 665 (1968), AND NUC. PHYS. 88, 333 (1968).
CONTINUATION OF PREVIOUS EXPERIMENT IN PHYS. REV. LETTERS 18, 1087 (1967), AND UNIV. OF ROCHESTER UR-875-218.

ADDITIONAL FOOTNOTE REFERENCES

REVIEW OF SCIENTIFIC INSTRUMENTS 38, 879 (1967), CERN TC/66-17, PHYS. REV. LETTERS 16, 485 (1966), CERN 65-24VOL.1, PHYS. LETTERS 25B, 426 (1967), NUOVO CIMENTO 48A, 589 (1967), UCRL 15744, NUOVO CIMENTO 50, 377 (1967), PHYS. REV. LETTERS 16, 855 (1966), PHYS. REV. LETTERS 17, 884 (1966), PHYS. REV. LETTERS 19, 546 (1967), AND PHYS. REV. 120, 1857 (1960).

ARTICLE READ BY ODETTE BENARY IN 12/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K⁺ ON PROTON AT 12.7 GEV/C.

THIS EXPERIMENT USES THE 8.N.L. 80 IN. (H) BUBBLE CHAMBER.

KEY WORDS = CROSS SECTION DALITZ PLOT MODELS ANGULAR DISTRIBUTION Q PEAK DELTA++ K*(890)
K*(1420) DENSITY MATRIX MASS SPECTRUM

CROSS SECTION FOR K⁺ PROTON = K⁺ PI⁺ PI⁻ NEUTRON. [PAGE 175]

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS
12.73	950. ± 100.

CROSS SECTION FOR K⁺ PROTON = K⁰ PI⁺ PI⁰ PROTON. [PAGE 175]

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS
12.73	565. ± 70.

CROSS SECTION FOR K⁺ PROTON = K⁰ NEUTRON PI⁺ PI⁺. [PAGE 175]

(THIS DATA SHOULD NOT BE USED - A MORE RECENT VALUE MAY BE FOUND IN BERLINCHERI, ET. AL., PHYS. LETTERS 27B, 665 (1968))

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS
12.73	130. ± 40.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ K^+ \text{ LAMBDA.}$ [TABLE 1]

LABORATORY BEAM MOMENTUM GEV/C 12.73	MICRO-BARNS 12. $\pm$ 3.
---	-----------------------------

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow \text{PROTON PROTON ANTI-LAMBDA.}$ [TABLE 1]

LABORATORY BEAM MOMENTUM GEV/C 12.73	MICRO-BARNS 6. $\pm$ 2.
---	----------------------------

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \text{ DELTA}^{++}.$ [TABLE 1]
 $\text{DELTA}^{++} \rightarrow \text{PROTON PI}^+ \text{ [1]}$

(THIS DATA SHOULD NOT BE USED - A MORE RECENT VALUE MAY BE FOUND IN BERLINGHIERI-ET. AL., NUC. PHYS. 88, 333 (1968))

LABORATORY BEAM MOMENTUM GEV/C 12.73	MICRO-BARNS 50. $\pm$ 9.
---	-----------------------------

[1] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow \text{NEUTRON PI}^+ K^0(890)^+.$ [TABLE 1]
 $K^0(890)^+ \rightarrow K^0 \text{ PI}^+ \text{ [1]}$

LABORATORY BEAM MOMENTUM GEV/C 12.73	MICRO-BARNS 75. $\pm$ 25.
---	------------------------------

[1] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow \text{PROTON } K^0(890)^+.$ [TABLE 1]
 $K^0(890)^+ \rightarrow K^0 \text{ PI}^+ \text{ [1]}$

(THIS DATA SHOULD NOT BE USED - A MORE RECENT VALUE MAY BE FOUND IN BERLINGHIERI-ET. AL., NUC. PHYS. 88, 333 (1968))

LABORATORY BEAM MOMENTUM GEV/C 12.73	MICRO-BARNS 57. $\pm$ 9.
---	-----------------------------

[1] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.

27 INTERACTION OF K^+ MESONS WITH PROTONS. [PHYS. REV. 118, 593 (1960)]

T.F.KYCIA, L.T.KERTH, R.G.BAENDER (U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA, AND UNIV. OF CALIFORNIA, BERKELEY, CALIF., USA)

ABSTRACT THE TOTAL K^+P CROSS SECTION WAS MEASURED AT THE THREE K^+ MESON ENERGIES 175 $\pm$ 25, 225 $\pm$ 25, AND 275 $\pm$ 5 MEV, AND THE DIFFERENTIAL SCATTERING CROSS SECTION WAS MEASURED AT 225 MEV. THE K^+P NUCLEAR FORCE WAS SHOWN TO BE REPULSIVE, FROM THE OBSERVED CONSTRUCTIVE INTERFERENCE WITH COULOMB SCATTERING. THE DIFFERENTIAL CROSS SECTION WAS OTHERWISE ISOTROPIC AND COULD ARISE FROM EITHER PURE S-WAVE OR PURE P-WAVE SCATTERING. SUBTRACTED DISPERSION RELATIONS WERE APPLIED TO THESE DATA AND THE REST OF THE AVAILABLE K^+ -PROTON SCATTERING DATA. THE STATISTICAL ERRORS IN THE DATA WERE FOUND TO BE TOO LARGE TO DETERMINE THE K^+ -HYPERON RELATIVE PHASE. ON THE ASSUMPTION THAT THE K^+ LAMBDA AND K^+ LAMBDA $\bar{K}$ RELATIVE PHASES ARE THE SAME, THE $\tan^2 \delta$ FOR SCALAR COUPLING, $(G^2/4\pi)$ WOULD BE LESS THAN 0.6; FOR PSEUDOSCALAR COUPLING, IT WOULD BE LESS THAN 10.

ADDITIONAL FOOTNOTE REFERENCES

PHYS. REV. 112, 1793 (1958), PHYS. REV. 88, 1358 (1952), PHYS. REV. LETTERS 2, 117 (1959), NUOVO CIMENTO 7, 190 (1958), PROGRESS OF THEORETICAL PHYSICS 3, 238 (1958), PHYS. REV. 110, 572 (1958), PHYS. REV. 113, 565 (1958), PHYS. REV. 110, 569 (1958), PROGRESS OF THEORETICAL PHYSICS 4, 403 (1958), PHYS. REV. 106, 167 (1957), BULL. AM. PHYS. SOC. 4, 288 (1959), PHYS. REV. LETTERS 2, 312 (1959), BULL. AM. PHYS. SOC. 3, 363 (1958), AND PHYS. REV. LETTERS 2, 510 (1959).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 6/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K^+ ON PROTON FROM .45 TO .60 GEV/C.

THIS EXPERIMENT USES COUNTERS.

KEY WORDS - CROSS SECTION PHASE SHIFT ANGULAR DISTRIBUTION

K^+ PROTON TOTAL CROSS SECTION.

LABORATORY BEAM ENERGY GEV	MILLI-BARNS
.175 $\pm$.025	16.3 $\pm$ 1.7
.225 $\pm$.025	15.2 $\pm$ 1.3
.275 $\pm$.025	16.3 $\pm$ 1.7

28 REACTION $K^+ \rightarrow K^+ p \pi^+ \pi^-$ AT 2.26 GEV/C. [PHYS. REV. 158, 1281 (1967)]

F. BOMSE, S. ROSENSTEIN, J. COLE, D. GILLESPIE, R. KRAEMER, G. LUSTE, I. MILLER, E. MOSES, A. PEVSNER, R. SINGH, R. ZDANIS [JOHNS HOPKINS UNIV., BALTIMORE, MD., USA]

ABSTRACT THE REACTION $K^+ \rightarrow K^+ p \pi^+ \pi^-$ AT 2.26 GEV/C IS INVESTIGATED. IT IS FOUND TO BE DOMINATED BY THE SIMULTANEOUS PRODUCTION OF THE $K^*(895 \text{ MEV})$ AND $N^*(1238 \text{ MEV})$ RESONANCES. THE PRODUCTION AND DECAY ANGULAR DISTRIBUTIONS OF THESE RESONANCES IN THE DOUBLE-RESONANCE CHANNEL ARE ANALYZED AND THE RESULTS COMPARED WITH THE PREDICTIONS OF A SINGLE-PARTICLE-EXCHANGE MODEL. THESE DISTRIBUTIONS ARE CONSISTENT WITH THE ASSUMPTION THAT THE PRIMARY PRODUCTION MECHANISM RESPONSIBLE FOR THE REACTION $K^+ \rightarrow K^+ p \pi^+ \pi^-$ AT 2.26 GEV/C IS THE EXCHANGE OF A SINGLE π MESON. A SEARCH IS CONDUCTED IN THIS CHANNEL FOR OTHER POSSIBLE RESONANT STATES. A MONTE CARLO ANALYSIS IS UTILIZED FOR THIS PURPOSE. THE DATA ARE NOT FOUND TO BE CONSISTENT WITH THE PRODUCTION OF ANY OTHER RESONANCES EXCEPT THE AFOREMENTIONED K^* (895 MEV) AND N^* (1238 MEV).

ADDITIONAL FOOTNOTE REFERENCES

NUOVO CIMENTO 34, 1644 (1964), NUOVO CIMENTO 33, 309 (1964), NUOVO CIMENTO 34, 735 (1964), REV. MOD. PHYS. 37, 484 (1965), PHYS. REV. 139, 8428 (1965), NUOVO CIMENTO 39, 417 (1965), PHYS. LETTERS 9, 71 (1964), PHYS. LETTERS 15, 74 (1965), PHYS. REV. LETTERS 16, 1069 (1966), PHYS. LETTERS 9, 207 (1964), PHYS. LETTERS 15, 104 (1965), AND PHYS. LETTERS 14, 338 (1965).

ARTICLE READ BY NAOMI BARASH-SCHMIOT IN 5/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K^+ ON PROTON AT 2.26 GEV/C.

THIS EXPERIMENT USES THE R.N.L. 20 IN. (H) BUBBLE CHAMBER. A TOTAL OF 100000 PICTURES ARE REPORTED ON.

KEY WORDS $K^*(895)$ DELTA++ ANGULAR DISTRIBUTION ONE PARTICLE EXCHANGE DENSITY MATRIX
MASS SPECTRUM CROSS SECTION

CROSS SECTION FOR K^+ PROTON $\rightarrow K^+$ PROTON $\pi^+ \pi^-$.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS
2.260 $\pm$.023	1.7 $\pm$.2	1296

CROSS SECTION FOR K^+ PROTON $\rightarrow K^*(890) \text{ DELTA++}$.
 $K^*(890) \rightarrow K^+ \pi^-$ [1]
 $\text{DELTA++} \rightarrow \text{PROTON } \pi^+$ [2]

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.260 $\pm$.023	.8 $\pm$.1

[1] COUNTED ONLY EVENTS ABOVE BACKGROUND [MASS = .895 GEV; WIDTH = .050 GEV].
[2] COUNTED ONLY EVENTS ABOVE BACKGROUND [MASS = 1.210 GEV; WIDTH = .120 GEV].

CROSS SECTION FOR K^+ PROTON $\rightarrow K^+$ PROTON $\pi^+ \pi^-$.
CROSS SECTION IS ONLY FOR THE NON-RESONANT PRODUCTION OF THESE PARTICLES.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.260 $\pm$.023	.5 $\pm$.1

CROSS SECTION FOR K^+ PROTON $\rightarrow K^*(890) \text{ PROTON } \pi^+$.
 $K^*(890) \rightarrow K^+ \pi^-$ [1]

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.260 $\pm$.023	.2 $\pm$.1

[1] OVERLAP REGION EXCLUDED.

CROSS SECTION FOR K^+ PROTON $\rightarrow K^+ \pi^- \text{ DELTA++}$.
 $\text{DELTA++} \rightarrow \text{PROTON } \pi^+$ [1]

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.260 $\pm$.023	.3 $\pm$.1

[1] OVERLAP REGION EXCLUDED.

29 K⁺P ELASTIC SCATTERING AT 3.0 GEV/C. [NUOVO CIMENTO 43A, 142 (1966)]

J. DEBAISIEUX, F. GRARD, J. HEUCHEBAERT, L. PAPE, R. WINDMOLDS (LAB. DES HAUTES ENERGIES, BRUXELLES, BELGIUM)
R. GEORGE, Y. GOLDSCHMIDT-CLERMONT, V. P. HENRI, D. A. G. LEITH, G. R. LYNCH, F. MULLER, J. L. PERREAU (EUROPEAN ORG. FOR NUC. RES., GENEVA, SWITZERLAND)

ABSTRACT IN THE COURSE OF A SYSTEMATIC STUDY OF K⁺P INTERACTIONS AT 3.0 GEV/C, THE ELASTIC-SCATTERING REACTION HAS BEEN INVESTIGATED. A TOTAL OF 1720 EVENTS WERE IDENTIFIED AS ELASTIC SCATTERS, GIVING A CROSS-SECTION OF (4.8 ± 0.4) MB. THE ANGULAR DISTRIBUTION SHOWS CHARACTERISTIC DIFFRACTION PEAKING AND WAS FITTED USING $D\text{-SIGMA}/D\text{-T} = 10\text{-SIGMA}/D\text{-T}$ (AT T=0) $\exp(\alpha \text{ALPHA-T} + \beta \text{BETA-T SQUARED})$, IN THE MOMENTUM-TRANSFER REGION $(0.05 / 1.14) \text{ (GEV/C)}^2$. THE BEST FIT GAVE $\text{ALPHA} = (4.55 \pm 0.39) \text{ (GEV/C)}^{**2}$ AND $\text{BETA} = (0.64 \pm 0.42) \text{ (GEV/C)}^{**4}$. THE EXTRAPOLATED EXPERIMENTAL CROSS-SECTION AT 0 DEG., $[D\text{-SIGMA}/D\text{-T (AT T=0)]$, IS FOUND TO BE $(19.5 \pm 2.3) \text{ MB/(GEV/C)}^2$ -SQUARED, AND EXCEEDS THE OPTICAL-THEOREM PREDICTION BY $(3.8 \pm 2.3) \text{ MB/(GEV/C)}^2$ -SQUARED, IMPLYING THAT THERE IS A CONTRIBUTION FROM THE REAL PART OF THE K⁺P SCATTERING AMPLITUDE AT 3.0 GEV/C.

ADDITIONAL FOOTNOTE REFERENCES

PHYS. REV. 129, 2743 (1963), PHYS. REV. 139, 81411 (1965), PHYS. REV. LETTERS 11, 503 (1963), CERN 63-12 (1963), UCRL 8033 REV. (1965), PHYS. REV. LETTERS 7, 188 (1961), PHYS. REV. LETTERS 9, 135 (1962), PHYS. REV. LETTERS 7, 182 (1961), PHYS. REV. 129, 2285 (1963), PHYS. REV. 138, 8913 (1965), NUCLEAR INSTRUMENTS AND METHODS 34, 242 (1965), SIENNA CONFERENCE 634 (1963), PHYS. REV. LETTERS 12, 229 (1964), PHYS. REV. 132, 1252 (1963), PHYS. REV. LETTERS 11, 287 (1963), PHYS. REV. LETTERS 12, 429 (1964), AND PHYS. REV. 135, 1263 (1964).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 1/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K⁺ ON PROTON AT 2.97 GEV/C.

THIS EXPERIMENT USES THE SACLAY 81 CM (H) BUBBLE CHAMBER. A TOTAL OF 40000 PICTURES ARE REPORTED ON.

GENERAL COMMENTS ON THIS ARTICLE

- 1 K⁺ LIFETIME TAKEN AS 1.229 ± 0.008 ($\times 10^{**8}$ SEC)
- 2 K⁺ TO TAU BRANCHING RATIO TAKEN AS 0.055 ± 0.001 .

KEY WORDS = CROSS SECTION ANGULAR DISTRIBUTION

K⁺ PROTON ELASTIC CROSS SECTION.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS
2.970 $\pm$.015	4.0 $\pm$.4	1720

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = 2.970 $\pm$.015 GEV/C.

COS(THETA)		D-SIGMA/D-T MB/(GEV/C)**2		NO. EVENTS
MIN	MAX			
-.95	.00	.01 $\pm$.01		12
.00	.10	.00	.00	0
.10	.40	.04	.01	15
.40	.55	.19	.04	24
.55	.60	.48	.12	21
.60	.65	.62	.13	27
.65	.70	1.29	.21	36
.70	.75	1.45	.23	63
.75	.80	2.32	.31	101
.80	.85	3.55	.43	155
.85	.90	6.00	.65	261
.90	.95	8.98	.91	391
.95	.98	14.65	1.51	383

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.

30 KAON-NUCLEON TOTAL CROSS SECTIONS FROM 0.6 TO 2.65 GEV/C. [PHYS. REV. 168, 1466 (1968)]

D. V. BUGG, R. S. GILMORE, K. M. KNIGHT, D. C. SALTER, G. H. STAFFORD, E. J. WILSON (RUTHERFORD HIGH EN. LAB., CHILTON, DID. BERK. ENGLAND)
J. D. DAVIES, J. D. DOWELL, P. M. HATTERSLEY, R. J. HOMER, A. W. ODELL (BIRMINGHAM UNIV., BIRMINGHAM, ENGLAND)
A. A. CARTER, R. J. TAPPER, K. F. RILEY (CAVENDISH LAB., CAMB. UNIV., CAMBRIDGE, ENGLAND)

ABSTRACT TOTAL CROSS SECTIONS OF K⁺ AND K⁻ MESONS ON PROTONS AND DEUTERONS HAVE BEEN MEASURED IN A TRANSMISSION EXPERIMENT OVER THE RANGE OF LABORATORY MOMENTUM 0.6 TO 2.65 GEV/C. MEASUREMENTS HAVE BEEN MADE ON K⁻ AT 50 MOMENTA AT INTERVALS OF 25 TO 50 MEV/C; THE EXPERIMENTAL ACCURACY IS BETTER THAN 1 PER CENT ABOVE 700 MEV/C, AND THE MOMENTUM RESOLUTION OF THE BEAM IS ± 0.6 PER CENT. STRUCTURE IS OBSERVED IN THE TOTAL CROSS SECTIONS SUGGESTING OR CONFIRMING $\gamma(11)$ RESONANCES AT MASSES OF 1665, 1768, 1905, 2020, 2250, AND 2455 MEV/C-SQUARED AND $\gamma(13)$ RESONANCES AT MASSES OF 1695, 1819, 1870, 2130, AND 2340 MEV/C-SQUARED. THE K⁺ MEASUREMENTS ARE LESS EXTENSIVE, AND ARE CONCENTRATED IN THE MOMENTUM RANGE BELOW 1.5 GEV/C; THE EXPERIMENTAL ERRORS ARE TYPICALLY ± 0.2 MB. STRUCTURE PREVIOUSLY REPORTED IN THE K⁺P AND K⁺D TOTAL CROSS SECTIONS NEAR A LABORATORY MOMENTUM OF 1.2 GEV/C IS CONFIRMED. TOTAL CROSS SECTIONS OF K⁺ AND K⁻ ON CARBON HAVE BEEN MEASURED AT A NUMBER OF MOMENTA WITH AN ACCURACY OF ABOUT ± 2 PER CENT.

ADDITIONAL FOOTNOTE REFERENCES

PHYS. REV. 123, 320 (1967), PHYS. REV. 125, 1696 (1962), PHYS. REV. LETTERS 10, 188 (1963), PHYS. LETTERS 6, 296 (1963), PHYS. REV. LETTERS 16, 1228 (1966), PHYS. REV. LETTERS 18, 62 (1967), PHYS. REV. LETTERS 17, 102 (1966), NUCLEAR INSTRUMENTS AND METHODS 17, 20 (1962), PHYS. REV. LETTERS 17, 561 (1966), RUTHERFORD HIGH ENERGY LAB. RPP/H/28, PHYS. REV. 99, 530 (1955), PHYS. REV. 100, 242 (1953), CERN PHYS. 67-17 (1967), PHYS. REV. LETTERS 19, 678 (1967), PHYS. REV. 118, 553 (1960), PHYS. REV. LETTERS 7, 187 (1961), PHYS. REV. 127, 2745 (1953), PHYS. REV. LETTERS 4, 135 (1962), DUBNA D-721 (1961), FORTSCHRITTE DER PHYSIK 9, 549 (1961), AND PHYS. REV. 126, 360 (1962).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 2/68, AND VERIFIED BY LEROY PRICE.

BEAM NO. 1 IS K⁺ ON PROTON FROM .592 TO 2.473 GEV/C.
NO. 2 IS K⁺ ON DEUTERON FROM .591 TO 2.472 GEV/C.
NO. 3 IS K⁻ ON PROTON FROM .59 TO 2.66 GEV/C.
NO. 4 IS K⁻ ON DEUTERON FROM .59 TO 2.66 GEV/C.

THIS EXPERIMENT USES COUNTERS.

GENERAL COMMENTS ON THIS ARTICLE

- 1 FINDS THAT GLAUBER THEORY IS NOT RELIABLE FOR EXTRACTING NEUTRON CROSS SECTIONS

KEY WORDS = CROSS SECTION

K+ PROTON TOTAL CROSS SECTION.

LABORATORY BEAM MOMENTUM GEV/C (1)	MILLI-BARNS
.592 +- .001	14.36 +- 1.24
.620 .001	12.91 .49
.644 .001	12.89 .48
.713 .001	11.14 .35
.768 .001	11.65 .33
.823 .001	12.97 .25
.864 .001	13.21 .24
.904 .001	14.23 .19
.938 .001	14.59 .21
.969 .001	15.63 .22
1.029 .001	15.69 .17
1.084 .001	17.04 .17
1.140 .001	18.02 .17
1.189 .001	18.06 .17
1.238 .001	18.11 .17
1.293 .001	18.44 .15
1.347 .001	18.27 .16
1.408 .001	17.97 .14
1.468 .001	17.94 .16
1.563 .001	17.66 .14
1.945 .001	17.41 .16
2.473 .001	17.25 .12

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF +- .5 PER CENT.

K+ NEUTRON TOTAL CROSS SECTION.

LABORATORY BEAM MOMENTUM GEV/C (1)	MILLI-BARNS (2)
.591 +- .001	26.71 +- 1.23
.619 .001	28.61 .44
.643 .001	28.13 .44
.712 .001	27.70 .33
.767 .001	28.90 .30
.822 .001	29.56 .22
.863 .001	30.33 .21
.903 .001	31.27 .20
.937 .001	32.14 .18
.968 .001	32.72 .19
1.028 .001	34.29 .16
1.083 .001	35.90 .15
1.139 .001	37.34 .15
1.188 .001	37.36 .15
1.237 .001	37.46 .23
1.292 .001	37.43 .13
1.346 .001	37.41 .15
1.407 .001	36.92 .13
1.467 .001	36.53 .15
1.562 .001	36.29 .12
1.944 .001	35.19 .13
2.472 .001	34.78 .11

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF +- .5 PER CENT.

(2) PLUS POSSIBLE SYSTEMATIC ERRORS OF +- .1414 MILLI-BARNS AND +- .2 PER CENT.

31 BACKWARD KP ELASTIC SCATTERING AT 3.55 GEV/C. [PHYS. LETTERS 248, 317 (1967)]

J. BANAIGS, J. BERGER, C. BONNEL, J. DUFLO, L. GOLOZAH, F. PLOUIN (CNTR. D'ETUDES NUC. SACLAY, GIF-SUR-YVETTE, FRANCE)
 W. F. BAKER, P. J. CARLSON, V. CHABAUD, A. LUNDBY (EUROPEAN ORG. FOR NUC. RES., GENEVA, SWITZERLAND)

ABSTRACT BACKWARD ELASTIC K+P AND K-P SCATTERING HAS BEEN MEASURED IN THE ANGULAR INTERVAL THETA(C.M.) FROM 168 DEGREES TO 177 DEGREES. WE FIND A K+ DIFFERENTIAL CROSS SECTION OF 17 +- 4 MICROBARNS/SR AND A K- DIFFERENTIAL CROSS SECTION OF LESS THAN 0.6 MICROBARNS/SR. K+ P ELASTIC SCATTERING EXHIBITS A BACKWARD PEAK.

ADDITIONAL FOOTNOTE REFERENCES

PHYS. LETTERS 23, 505 (1966), NUOVO CIMENTO 45A, 885 (1966), PHYS. LETTERS 21, 117 (1966), PHYS. LETTERS 19, 441 (1965),
 NUOVO CIMENTO 43, 142 (1966), PHYS. LETTERS 23, 171 (1966), PHYS. REV. LETTERS 15, 913 (1966), JINR P1-3008 (1966),
 PHYS. REV. LETTERS 16, 1228 (1966), UCRL 8030 JAN. (1967), AND PHYS. REV. LETTERS 17, 102 (1966).

ARTICLE READ BY NAOMI BARASH-SCHMIOT IN 1/68, AND VERIFIED BY LEROY PRICE.

BEAM NO. 1 IS K+ ON PROTON AT 3.55 GEV/C.
 NO. 2 IS K- ON PROTON AT 3.55 GEV/C.

THIS EXPERIMENT USES SPARK CHAMBERS. A TOTAL OF 108000 PICTURES ARE REPORTED ON.

KEY WORDS - CROSS SECTION ANGULAR DISTRIBUTION

ELASTIC DIFFERENTIAL CROSS SECTION FOR K+ PROTON.

LABORATORY BEAM MOMENTUM = 3.55 +- .05 GEV/C.

(THIS DATA SHOULD NOT BE USED - MORE RECENT VALUES MAY BE FOUND IN BAKER, ET. AL., NUC. PHYS. BY, 249 (1969))

COS(THETA)	D-SIGMA/D-OMEGA UB/SR (1)	NO. EVENTS
-1.000 (2)	26.1 +- 12.0	
-.983 .008	14.9 5.6	7
-.994 .004	21.3 5.5	15

THETA IS THE ANGLE THAT THE K+ MAKES WITH THE BEAM IN THE GRAND C.M.

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF +- 25 PER CENT.

(2) EXTRAPOLATED POINT.

32 TOTAL CROSS-SECTION MEASUREMENTS OF K⁺ P AND K⁺ N INTERACTIONS [PHYS. REV. LETTERS 7, 182 (1961)]

V. COOK, D. KEEFE, L. T. KERTH, P. G. MURPHY, W. A. WENZEL, T. F. ZIPF (U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA, AND UNIV. OF CALIFORNIA, BERKELEY, CALIF., USA)

ADDITIONAL FOOTNOTE REFERENCES

NUOVO CIMENTO 12, 241 (1959), PHYS. REV. 118, 553 (1960), ROCHESTER CONFERENCE 451 (1960), PHYS. REV. LETTERS 2, 117 (1959), PHYS. REV. LETTERS 5, 333 (1960), ROCHESTER CONFERENCE 444 (1960), DUBNA D721 (1961), PHYS. REV. 123, 320 (1961), PHYS. REV. 100, 242 (1955), AND PHYS. REV. LETTERS 7, 188 (1961).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 1/68, AND VERIFIED BY LEROY PRICE.

BEAM NO. 1 IS K⁺ ON PROTON FROM .77 TO 2.83 GEV/C.

NO. 2 IS K⁺ ON DEUTERON FROM .77 TO 2.83 GEV/C.

THIS EXPERIMENT USES COUNTERS.

GENERAL COMMENTS ON THIS ARTICLE

1 NEUTRON CROSS SECTIONS HAVE NOT BEEN CORRECTED FOR PAULI PRINCIPLE EFFECTS

KEY WORDS = CROSS SECTION

K⁺ PROTON TOTAL CROSS SECTION.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS [1]
.77	12.0 ± .9
.97	15.4 ± .6
1.17	10.1 ± .6
1.30	17.9 ± .9
1.44	18.1 ± .7
1.69	17.5 ± .6
1.97	16.9 ± .6
2.26	17.1 ± .5
2.55	17.1 ± .6
2.83	16.7 ± .5

[1] PLUS POSSIBLE SYSTEMATIC ERROR OF ± .3 MILLI-BARNS.

K⁺ DEUTERON TOTAL CROSS SECTION.
COULOMB CORRECTIONS APPLIED

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS [1]
.77	27.3 ± .6
.97	32.4 ± .5
1.17	35.4 ± .5
1.30	35.6 ± .6
1.44	35.4 ± .5
1.69	35.1 ± .5
1.97	34.6 ± .5
2.26	33.9 ± .6
2.55	33.4 ± .5
2.83	33.4 ± .7

[1] PLUS POSSIBLE SYSTEMATIC ERROR OF ± .3 MILLI-BARNS.

K⁺ NEUTRON TOTAL CROSS SECTION.
GLAUBER CORRECTION APPLIED

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS [1]
.77	15.5 ± 1.1
.97	17.8 ± .8
1.17	10.2 ± .8
1.30	18.5 ± 1.1
1.44	18.1 ± .9
1.69	18.5 ± .8
1.97	18.6 ± .6
2.26	17.7 ± .8
2.55	17.1 ± .8
2.83	17.5 ± .9

[1] PLUS POSSIBLE SYSTEMATIC ERROR OF ± .3 MILLI-BARNS.

33 K⁺ P AND K⁻ P TOTAL CROSS SECTIONS IN THE MOMENTUM RANGE 3-19 BEV/C. [PHYS. REV. 129, 2295 (1963)]

W.F. BAKER, R.L. COOL, E.W. JENKINS, T.F. KYCIA, R.H. PHILLIPS, A.L. READ (BROOKHAVEN NAT. LAB., UPTON, L.I., N. Y., USA)

ABSTRACT THE TOTAL CROSS SECTIONS OF K⁺ AND K⁻ MESONS ON PROTONS HAVE BEEN MEASURED BETWEEN 3 AND 19 BEV/C AT THE BROOKHAVEN ALTERNATING GRADIENT SYNCHROTRON. THE DATA WERE OBTAINED FROM MEASUREMENTS OF ATTENUATION IN A LIQUID HYDROGEN TARGET. THE K MESONS WERE DEFINED IN MOMENTUM BY MAGNETIC DEFLECTION AND A SCINTILLATION COUNTER TELESCOPE. THEY WERE IDENTIFIED BY MEASURING THE VELOCITY WITH A HIGH-PRESSURE GAS-FILLED CERENKOV COUNTER. VALUES OF THE CROSS SECTIONS WERE OBTAINED AT INTERVALS OF APPROXIMATELY 1 BEV/C WITH AN ERROR FOR EACH VALUE OF APPROXIMATELY 5 PER CENT. THE K⁺P TOTAL CROSS SECTION IS, WITHIN ERRORS, CONSTANT OVER THE ENTIRE MOMENTUM RANGE WITH A VALUE OF 18.4 MICROBARN. THE K⁻P TOTAL CROSS SECTION DECREASES GRADUALLY FROM A VALUE OF 28 MICROBARN AT 6 BEV/C TO 21.6 MICROBARN AT 19 BEV/C. THE CROSS SECTIONS ARE SIGNIFICANTLY DIFFERENT UP TO THE HIGHEST MOMENTA REACHED IN THIS EXPERIMENT.

ADDITIONAL FOOTNOTE REFERENCES

FORTSCHRITTE DER PHYSIK 9, 549 (1961), PHYS. REV. LETTERS 7, 188 (1961), PHYS. REV. 125, 1696 (1962), PHYS. REV. LETTERS 9, 135 (1962), PHYS. REV. LETTERS 2, 117 (1959), CERN CONFERENCE 385 (1962), JETP 19 29 (1962), PHYS. REV. LETTERS 7, 182 (1961), PHYS. REV. LETTERS 123, 320 (1961), PHYS. REV. LETTERS 5, 333 (1960), REV. MOD. PHYS. 33, 382 (1961), JETP 7 499 (1958), PHYS. REV. LETTERS 7, 394 (1961), PHYS. REV. LETTERS 8, 41 (1952), NUOVO CIMENTO 14, 951 (1959), NUOVO CIMENTO 18, 947 (1960), PHYS. REV. A, 142 (1962), BNL EDC-36, BNL WFB-1, NUCLEAR ELECTRONICS 1, 63 (1962), BNL 5390, BNL GTD-2, REVIEW OF SCIENTIFIC INSTRUMENTS 25, 1070 (1954), PHYS. REV. LETTERS 7, 185 (1961), PHYS. REV. LETTERS 7, 352 (1961), PHYS. REV. LETTERS 8, 173 (1962), PHYS. REV. LETTERS 7, 35 (1961), PHYS. REV. LETTERS 9, 76 (1962), AND PHYS. REV. LETTERS 9, 183 (1962).

ARTICLE READ BY ODETTE BENARY IN 12/68, AND VERIFIED BY LEROY PRICE.

BEAM NO. 1 IS K⁺ ON PROTON FROM 3.25 TO 19.00 GEV/C.
NO. 2 IS K⁻ ON PROTON FROM 4.0 TO 19.1 GEV/C.

THIS EXPERIMENT USES COUNTERS.

KEY WORDS = CROSS SECTION

K⁺ PROTON TOTAL CROSS SECTION. [TABLE 1]

(MOMENTUM SPREAD OF THE BEAM IS GIVEN ONLY AT 3 AND 19 GEV/C. WE HAVE INTERPOLATED FOR INTERMEDIATE VALUES.)

LABORATORY BEAM MOMENTUM GEV/C	PER CENT	MILLI-BARN
3.25	1.2	17.5 ± 1.2
4.00	1.2	17.6 ± 1.2
5.50	1.3	17.9 ± .8
7.00	1.4	18.4 ± .6
8.50	1.5	18.7 ± .6
10.00	1.5	18.8 ± .7
10.90	1.6	18.1 ± .7
11.50	1.6	19.0 ± .6
12.50	1.7	18.3 ± .8
13.40	1.7	17.5 ± .6
15.00	1.8	18.5 ± .7
16.90	1.9	18.8 ± .6
19.00	2.0	17.3 ± .9

34 K⁺D PARTIAL CROSS SECTIONS AROUND 1 BEV/C. [PHYS. REV. LETTERS 21, 1485 (1968)]

A.A. MIRATA, C.G. WOHL, G. GLONHABER, G.H. TRILLING (U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA, AND UNIV. OF CALIFORNIA, BERKELEY, CALIF., USA)

CROSS REFERENCES
SEE ALSO UCRL 18322 (1968).

ADDITIONAL FOOTNOTE REFERENCES

PHYS. REV. LETTERS 17, 102 (1966), PHYS. REV. 168, 1466 (1968), PHYS. REV. LETTERS 18, 1377 (1967), UCRL 18131 (1968), PHYS. REV. 140, 8104 (1965), PHYS. REV. LETTERS 7, 378 (1961), PHYS. REV. LETTERS 19, 734 (1965), PHYS. REV. 134, 8111 (1964), CAVENDISH LAB. REP 69-10 (1961), AND PHYS. REV. LETTERS 9, 135 (1962).

ARTICLE READ BY LEROY PRICE IN 9/68, AND VERIFIED BY ROGER BLAND.

BEAM IS K⁺ ON DEUTERON FROM .863 TO 1.364 GEV/C.

THIS EXPERIMENT USES THE L.R.L. 25 IN. (H) BUBBLE CHAMBER. A TOTAL OF 100000 PICTURES ARE REPORTED ON.

KEY WORDS = CROSS SECTION

CROSS SECTION FOR K⁺ DEUTERON = K⁰ PROTON PROTON.

(FROM TABLE 1, UCRL-18322.)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARN
.863	6.72 ± .40
.968	6.26 ± .29
1.211	4.99 ± .24
1.364	3.69 ± .29

CROSS SECTION FOR K⁺ DEUTERON = K⁰ PROTON PROTON π^0 .

(FROM TABLE 1, UCRL-18322.)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARN
.863	.48 ± .08
.968	.82 ± .08
1.211	2.09 ± .13
1.364	2.06 ± .19

CROSS SECTION FOR $K^+ \text{ DEUTERON} \rightarrow K^0 \text{ PROTON NEUTRON } \pi^+$.

(FROM TABLE I, UCRL-18322.)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.863	1.24 $\pm$.14
.968	2.47 $\pm$.16
1.211	6.21 $\pm$.29
1.364	6.08 $\pm$.42

CROSS SECTION FOR $K^+ \text{ DEUTERON} \rightarrow K^+ \text{ PROTON PROTON } \pi^-$.

(FROM TABLE I, UCRL-18322.)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.863	.46 $\pm$.05
.968	.91 $\pm$.05
1.211	2.89 $\pm$.12
1.364	3.21 $\pm$.20

CROSS SECTION FOR $K^+ \text{ DEUTERON} \rightarrow K^0 \pi^+ \text{ DEUTERON}$.(FROM TABLE I, UCRL-18322. THIS IS INCLUDED IN $K^0 \text{ P N } \pi^+$ ABOVE. DEUTERON DOES NOT BREAK UP.)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.863	.13 $\pm$.04
.968	.13 $\pm$.03
1.211	.14 $\pm$.03
1.364	.18 $\pm$.05

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \pi^+ \text{ PROTON}$.* * THIS DATA SHOULD NOT BE USED * * (DEUTERON CORRECTIONS NOT APPLIED. THIS IS INCLUDED IN $K^0 \text{ P N } \pi^+$ ABOVE. FROM TABLE I, UCRL-18322.)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.863	.98 $\pm$.13
.968	1.99 $\pm$.15
1.211	4.76 $\pm$.26
1.364	4.78 $\pm$.38

CROSS SECTION FOR $K^+ \text{ NEUTRON} \rightarrow K^0 \pi^+ \text{ NEUTRON}$.(FROM TABLE I, UCRL-18322. THIS IS INCLUDED IN $K^0 \text{ P N } \pi^+$ ABOVE. DEUTERON CORRECTIONS NOT APPLIED.)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.863	.13 $\pm$.04
.968	.35 $\pm$.05
1.211	1.31 $\pm$.12
1.364	1.12 $\pm$.15

CROSS SECTION FOR $K^+ \text{ DEUTERON} \rightarrow K^+ \text{ DEUTERON} + K^+ \text{ PROTON (NEUTRON SPECTATOR)} + K^+ \text{ NEUTRON (PROTON SPECTATOR)}$.

(FROM TABLE I, UCRL-18322. THESE WERE DERIVED FROM DATA BY ASSUMING 1-SPIN, ETC.)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.863	27.59 $\pm$.32
.968	27.17 $\pm$.35
1.211	22.73 $\pm$.63
1.364	20.97 $\pm$.99

CROSS SECTION FOR $K^+ \text{ DEUTERON} \rightarrow K^+ \text{ PROTON NEUTRON}$.

(FROM TABLE I, UCRL-18322. THESE WERE DERIVED FROM DATA BY ASSUMING 1-SPIN, ETC.)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.863	20.87 $\pm$.56
.968	20.91 $\pm$.50
1.211	17.74 $\pm$.78
1.364	17.28 $\pm$ 1.15

CROSS SECTION FOR $K^+ \text{ DEUTERON} \rightarrow K \text{ MESON NUCLEON NUCLEON PION.}$

(FROM TABLE I, UCRL-18322. THESE WERE DERIVED FROM DATA BY ASSUMING I-SPIN, ETC.)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.863	2.74 $\pm$.24
.968	5.54 .29
1.211	14.51 .60
1.364	15.54 .95

CROSS SECTION FOR $K^+ \text{ DEUTERON} \rightarrow K \text{ MESON NUCLEON NUCLEON PION PION.}$

(FROM TABLE I, UCRL-18322. THESE WERE DERIVED FROM DATA BY ASSUMING I-SPIN, ETC.)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.968	.010 $\pm$.005
1.211	.170 .050
1.364	.760 .180

$I = 0$ K NUCLEON ELASTIC CROSS SECTION.

(FROM TABLE II, UCRL-18322.)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS (1)
.863	21.5
.968	21.5
1.211	16.1
1.364	13.0

(1) VALUES ARE APPROXIMATE ONLY.

$I = 0$ CROSS SECTION FOR $K \text{ NUCLEON} \rightarrow K \text{ MESON NUCLEON PION.}$

(FROM TABLE II, UCRL-18322.)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.863	1.0 $\pm$.4
.968	2.5 .6
1.211	8.0 1.0
1.364	8.9 1.8

35 INTERFERENCE EFFECTS IN THE REACTION $K^+ p \rightarrow \text{KAON NUCLEON PION}$ AT 1.2 BEV/C. [PHYS. REV. LETTERS 17, 939 (1966)]

R.W. BLAND, M.G. BOWLER, J.L. BROWN, G. GOLDBABER, S. GOLDBABER, J.A. KADYK, G.H. TRILLING (U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA, AND UNIV. OF CALIFORNIA, BERKELEY, CALIF., USA)

ABSTRACT SUBSTANTIAL INTERFERENCE HAS BEEN OBSERVED BETWEEN N^* AND K^* PRODUCTION IN 1.2-BEV/C $K^+ p$ INTERACTIONS.

CROSS REFERENCES
DATA SUPERSEDED BY UCRL 18758 (1969).

ADDITIONAL FOOTNOTE REFERENCES
PHYS. REV. LETTERS 10, 562 (1961), PHYS. REV. LETTERS 15, 737 (1965), PHYS. LETTERS 21, 462 (1966), UCRL 11527 (1964),
PHYS. REV. LETTERS 10, 351 (1963), PHYS. REV. LETTERS 11, 90 (1963), UCRL 16370 (1965), PHYS. REV. LETTERS 14, 118
(1965), NUDVO CIMENTO 33, 906 (1964), AND PHYS. REV. LETTERS 17, 102 (1966).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 4/68, AND VERIFIED BY LERDY PRICE.

REMARK IS K^+ ON PROTON AT 1.2 GEV/C.

THIS EXPERIMENT USES THE L.R.L. 25 IN. (H) BUBBLE CHAMBER.

KEY WORDS = $K^*(890)$ DELTA(1238) DALITZ PLOT CROSS SECTION

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \text{ PROTON PI}^+.$

(THIS DATA SHOULD NOT BE USED - MORE RECENT VALUES MAY BE FOUND IN BLAND, ET AL., UCRL 18758 (1969))

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
1.2	5.1 $\pm$.3

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \text{ PROTON } \pi^0$.

[THIS DATA SHOULD NOT BE USED - A MORE RECENT VALUE MAY BE FOUND IN BLAND, ET AL., UCRL 18758 (1969)]

LABORATORY
BEAM MOMENTUM
GEV/C
1.2

MILLI-BARNS
2.0 $\pm$.2

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \text{ NEUTRON } \pi^+$.

[THIS DATA SHOULD NOT BE USED - A MORE RECENT VALUE MAY BE FOUND IN BLAND, ET AL., UCRL 18758 (1969)]

LABORATORY
BEAM MOMENTUM
GEV/C
1.2

MILLI-BARNS
.56 $\pm$.10

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow \text{DELTA}^{++} K^0$.

[THIS DATA SHOULD NOT BE USED - A MORE RECENT VALUE MAY BE FOUND IN BLAND, ET AL., UCRL 18758 (1969)]

LABORATORY
BEAM MOMENTUM
GEV/C
1.2

MILLI-BARNS
2.9 $\pm$.3

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)^+ \text{ PRUTON.}$

$K^*(890)^+ \rightarrow K^0 \pi^+$ (1)

[THIS DATA SHOULD NOT BE USED - A MORE RECENT VALUE MAY BE FOUND IN BLAND, ET AL., UCRL 18758 (1969)]

LABORATORY
BEAM MOMENTUM
GEV/C
1.2

MILLI-BARNS
.87 $\pm$.14

(1) COUNTED ONLY EVENTS ABOVE BACKGROUND.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow \text{DELTA}^+ K^+ \text{ PROTON } \pi^0$ (1)

[THIS DATA SHOULD NOT BE USED - A MORE RECENT VALUE MAY BE FOUND IN BLAND, ET AL., UCRL 18758 (1969)]

LABORATORY
BEAM MOMENTUM
GEV/C
1.2

MILLI-BARNS
.67 $\pm$.15

(1) COUNTED ONLY EVENTS ABOVE BACKGROUND.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)^+ \text{ PROTON.}$

$K^*(890)^+ \rightarrow K^+ \pi^0$ (1)

[THIS DATA SHOULD NOT BE USED - A MORE RECENT VALUE MAY BE FOUND IN BLAND, ET AL., UCRL 18758 (1969)]

LABORATORY
BEAM MOMENTUM
GEV/C
1.2

MILLI-BARNS
.66 $\pm$.12

(1) COUNTED ONLY EVENTS ABOVE BACKGROUND.

36 ANTIPROTON AND KADN ELASTIC SCATTERING AT HIGH ENERGIES. [PHYS. REV. LETTERS 11, 503 (1963)]

K.J.FOLEY, S.J.LINDENBAUM, W.A.LOVE, S.OZAKI, J.J.RUSSELL, L.C.L.YUAN [BROOKHAVEN NAT. LAB., UPTON, L.I., N. Y., USA]

ADDITIONAL FOOTNOTE REFERENCES

PHYS. REV. LETTERS 10, 376 (1963), PHYS. REV. LETTERS 10, 543 (1963), PHYS. REV. LETTERS 11, 425 (1963), PHYS. REV. 129, 2285 (1963), AND PHYS. REV. LETTERS 7, 185 (1961).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 1/68, AND VERIFIED BY LEROY PRICE.

BEAM NO. 1 IS K^+ ON PROTON FROM 6.8 TO 14.8 GEV/C.
NO. 2 IS K^- ON PROTON FROM 7.2 TO 9.0 GEV/C.
NO. 3 IS ANTI-PROTON ON PROTON FROM 7.2 TO 12.0 GEV/C.

THIS EXPERIMENT USES COUNTERS.

GENERAL COMMENTS ON THIS ARTICLE
1 K^- AND ANTIPROTON DATA NOT PUNCHED FOR NOW.

KEY WORDS = CROSS SECTION MODELS

K⁺ PROTON ELASTIC CROSS SECTION.

LABORATORY BEAM MOMENTUM		MILLI-BARNS (1)	
GEV/C			
6.8		3.48	± .43
9.8		3.34	.17
12.8		3.34	.15
14.8		3.41	.17

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF ± 7 PER CENT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = 6.8 GEV/C.

-T	D-SIGMA/D-T	
(GEV/C)**2 (1)	MB/(GEV/C)**2 (2)	
.251	4.810	± .340
.321	2.930	.240
.399	2.110	.180
.441	1.460	.150
.569	.944	.117
.663	.671	.100
.762	.331	.072
.867	.266	.071
.977	.219	.057
1.091	.117	.049
1.193	.058	.042

T IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K⁺) AND THE (OUTGOING K⁺).

(1) VALUES ARE APPROXIMATE ONLY.

(2) PLUS POSSIBLE SYSTEMATIC ERROR OF ± 7 PER CENT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = 9.8 GEV/C.

-T	D-SIGMA/D-T	
(GEV/C)**2 (1)	MB/(GEV/C)**2 (2)	
.049	14.740	± 1.560
.071	13.200	1.380
.098	10.150	1.080
.133	8.370	.970
.173	7.870	.940
.212	4.360	.780
.266	4.120	.290
.340	2.690	.210
.423	1.580	.140
.510	1.090	.120
.605	.716	.094
.707	.300	.062
.816	.270	.055
.931	.162	.048
1.051	.043	.027

T IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K⁺) AND THE (OUTGOING K⁺).

(1) VALUES ARE APPROXIMATE ONLY.

(2) PLUS POSSIBLE SYSTEMATIC ERROR OF ± 7 PER CENT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = 12.8 GEV/C.

-T	D-SIGMA/D-T	
(GEV/C)**2 (1)	MB/(GEV/C)**2 (2)	
.050	16.430	± 1.310
.083	12.760	.990
.120	8.470	.870
.166	7.210	.670
.224	5.050	.530
.277	3.800	.240
.291	3.620	.460
.354	2.250	.170
.441	1.430	.120
.531	.652	.085
.630	.521	.071
.736	.235	.046
.849	.112	.035
.971	.060	.025
1.099	.054	.019

T IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K⁺) AND THE (OUTGOING K⁺).

(1) VALUES ARE APPROXIMATE ONLY.

(2) PLUS POSSIBLE SYSTEMATIC ERROR OF ± 7 PER CENT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K+ PROTON.

LABORATORY BEAM MOMENTUM = 14.8 GEV/C.

-T (GEV/C)**2 [1]	D-SIGMA/D-T MB/(GEV/C)**2 [2]
.066	14.370 +- 1.060
.111	9.610 .840
.161	7.810 .770
.222	5.030 .570
.290	3.230 .230
.300	3.630 .410
.371	2.000 .160
.389	1.750 .300
.462	1.230 .110
.558	.805 .086
.661	.451 .067
.774	.181 .045
.896	.153 .040
1.024	.115 .035
1.160	.069 .028

T IS THE MOMENTUM TRANSFER BETWEEN THE [INCOMING K+] AND THE [OUTGOING K+].

[1] VALUES ARE APPROXIMATE ONLY.

[2] PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7 PER CENT.

37 EVIDENCE FOR AN S=+1, I=0 RESONANCE IN THE K+ NUCLEON SYSTEM. (PHYS. REV. LETTERS 18, 801 (1967))

A.A.CARTER (CAVENDISH LAB., CAMB. UNIV., CAMBRIDGE, ENGLAND)

ABSTRACT WE EVALUATE THE PHASE OF THE K+ NUCLEON FORWARD-SCATTERING AMPLITUDE FROM DISPERSION RELATIONS AND FIND A QUALITATIVE DIFFERENCE BETWEEN THE I=0 AND I=1 STATE. WITH ASSUMPTIONS FOR THE BACKGROUND CONTRIBUTIONS TO THE AMPLITUDE, WE PRESENT EVIDENCE FOR AN S=+1, I=0 RESONANCE.

CROSS REFERENCES

ANALYSIS OF DATA FROM PHYS. REV. LETTERS 17, 102 (1966), AND PHYS. REV. 168, 1466 (1968).

ADDITIONAL FOOTNOTE REFERENCES

PHYS. LETTERS 21, 229 (1966), NUOVO CIMENTO 45A, 792 (1966), AND PHYS. LETTERS 23, 380 (1966).

ARTICLE READ BY ODETTE BENARY IN 5/69, AND VERIFIED BY ROGER BLAND.

KEY WORDS - PHASE SHIFT I*

I = 0 K NUCLEON TOTAL CROSS SECTION.

(CALCULATED USING DATA FROM A.A.CARTER - CAVENDISH LABORATORY INTERVAL REPORT HEP 68-10(1968))

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS [1]	LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS [1]
.525	14.6	3.025	19.2
.575	16.0	3.075	19.2
.625	17.3	3.125	19.1
.675	18.6	3.175	19.1
.725	19.9	3.225	19.1
.775	21.1	3.275	19.1
.825	22.6	3.325	19.0
.875	22.6	3.375	19.0
.925	24.2	3.425	18.9
.975	25.3	3.475	18.9
1.025	25.6	3.525	18.8
1.075	25.6	3.575	18.8
1.125	25.2	3.625	18.8
1.175	24.5	3.675	18.8
1.225	23.9	3.725	18.7
1.275	23.2	3.775	18.7
1.325	22.4	3.825	18.7
1.375	21.7	3.875	18.7
1.425	21.4	3.925	18.7
1.475	21.4	3.975	18.6
1.525	21.2	4.025	18.6
1.575	21.0	4.075	18.6
1.625	20.8	4.125	18.5
1.675	20.7	4.175	18.4
1.725	20.6	4.225	18.3
1.775	20.6	4.275	18.3
1.825	20.5	4.325	18.3
1.875	20.4	4.375	18.3
1.925	20.1	4.425	18.2
1.975	20.3	4.475	18.2
2.025	20.2	4.525	18.2
2.075	20.2	4.575	18.1
2.125	20.2	4.625	18.1
2.175	20.0	4.675	18.1
2.225	19.9	4.725	18.1
2.275	19.8	4.775	18.1
2.325	19.8	4.825	18.0
2.375	19.7	4.875	18.0
2.425	19.6	4.925	18.0
2.475	19.5	4.975	18.0
2.525	19.4	5.025	18.0
2.575	19.4	5.075	18.0
2.625	19.3	5.125	18.0
2.675	19.3	5.175	18.0
2.725	19.2	5.225	17.9
2.775	19.2	5.275	17.9
2.825	19.2	5.325	17.9
2.875	19.2	5.375	17.9
2.925	19.2	5.425	17.9
2.975	19.2	5.475	17.9

[1] VALUES ARE APPROXIMATE ONLY.

38 DOUBLE REGGE DESCRIPTION OF OMEGA PRODUCTION IN $K^+ p$ INTERACTIONS. [PHYS. REV. 177, 2092 (1969)]

G. ALEXANDER, A. FIRESTONE, C. FU, G. GOLDBABER, A. PIGNOTTI (U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA, AND UNIV. OF CALIFORNIA, BERKELEY, CALIF., USA)

ABSTRACT WE PRESENT RESULTS ON THE OMEGA AND PHI PRODUCTION IN $K^+ p \rightarrow K^+ \Omega$ AND $K^+ p \rightarrow K^+ \phi$ INTERACTIONS AT INCIDENT MOMENTA OF 4.6 AND 9.0 GEV/C. THERE IS NO STATISTICALLY SIGNIFICANT EVIDENCE FOR A OMEGA OR K PHI RESONANCE PRODUCTION. THE 437 $K^+ \Omega$ AND 437 $K^+ \phi$ EVENTS AT 9 GEV/C ARE ANALYZED IN TERMS OF A MULTIPERIPHERAL REGGE MODEL ON THE ENTIRE DALITZ PLOT. THE EVENTS HAVE BEEN ASSIGNED TO THE VARIOUS PERIPHERAL DIAGRAMS ON THE BASIS OF CRITERIA IN THE FOUR-MOMENTUM TRANSFERS. A SATISFACTORY DESCRIPTION OF THE DATA HAS BEEN OBTAINED ON THE BASIS OF DIAGRAMS INVOLVING POMERANCHUK AND MESON EXCHANGES.

ADDITIONAL FOOTNOTE REFERENCES

PHYS. REV. 131, 2282 (1963), JETP 17 233 (1963), NUOVO CIMENTO 49A, 157 (1967), NUOVO CIMENTO 51A, 696 (1967), PHYS. REV. LETTERS 21, 701 (1968), CERN HADRON CONFERENCE 1 611 (1968), PHYS. REV. 166, 1525 (1968), PHYS. REV. 166, 1768 (1968), PHYS. REV. LETTERS 20, 1078 (1968), PHYS. REV. LETTERS 15, 118 (1965), PHYS. REV. LETTERS 12, 206 (1964), PHYS. REV. LETTERS 20, 518 (1968), JETP 19 1148 (1964), PHYS. REV. 168, 1597 (1968), PHYS. REV. 160, 1322 (1967), AND PHYS. REV. 165, 1699 (1968).

ARTICLE READ BY ODETTE REYARY IN 6/69, AND VERIFIED BY LEROY PRICE.

BEAM IS K^+ ON PROTON FROM 4.6 TO 9.0 GEV/C.

THIS EXPERIMENT USES THE B.N.L. 80 IN. (H) BUBBLE CHAMBER. A TOTAL OF 140000 PICTURES ARE REPORTED ON.

KEY WORDS = MASS SPECTRUM DALITZ PLOT MODELS ANGULAR DISTRIBUTION

* NO DATA PUNCHED FOR THIS ARTICLE *

39 TOTAL CROSS SECTIONS OF PROTONS, ANTIPROTONS, AND π AND K MESONS ON HYDROGEN AND DEUTERIUM IN THE MOMENTUM RANGE 6-22 GEV/C. [PHYS. REV. 138, 8913 (1965)]

W. GALBRAITH, E. W. JENKINS, T. F. KYCIA, B. A. LEONTIC, R. H. PHILLIPS, A. L. READ (BROOKHAVEN NAT. LAB., UPTON, L.I., N. Y., USA)
R. RUBENSTEIN (CORNELL UNIV., ITHACA, N. Y., USA)

ABSTRACT THE TOTAL CROSS SECTIONS σ_{TOT} OF p , $\bar{p}$, CHARGED π AND CHARGED K ON HYDROGEN AND DEUTERIUM HAVE BEEN MEASURED BETWEEN 6 AND 22 GEV/C AT INTERVALS OF 2 GEV/C TO AN ACCURACY GREATER THAN PREVIOUSLY REPORTED. THE METHOD UTILIZED WAS A CONVENTIONAL GOOD-GEOMETRY TRANSMISSION EXPERIMENT WITH SCINTILLATION COUNTERS SUBTENDING VARIOUS SOLID ANGLES OF LIQUID H_2 AND D_2 . WITH THE INCREASE IN STATISTICAL ACCURACY OF THE DATA, IT WAS FOUND THAT A PREVIOUSLY ADOPTED PROCEDURE OF LINEARLY EXTRAPOLATING TO ZERO SOLID ANGLE THE PARTIAL CROSS SECTIONS MEASURED AT FINITE SOLID ANGLES WAS NOT A SUFFICIENTLY ACCURATE PROCEDURE FROM WHICH TO DEDUCE σ_{TOT} . THE PARTICLE-NEUTRON CROSS SECTIONS ARE DERIVED BY APPLYING THE GLAUBER SCREENING CORRECTION TO THE DIFFERENCE BETWEEN THE PARTICLE-DEUTERON AND PARTICLE-PROTON CROSS SECTIONS. THE TOTAL CROSS SECTIONS OF π^+D AND π^-D ARE EQUAL AT ALL MEASURED MOMENTA, WHICH CONFIRMS THE VALIDITY OF CHARGE SYMMETRY UP TO 20 GEV/C. RESULTS ARE PRESENTED SHOWING THE VARIATION OF CROSS SECTIONS WITH MOMENTUM; EVIDENCE IS PRESENTED FOR A SMALL BUT SIGNIFICANT DECREASE IN pp TOTAL CROSS SECTIONS AND $p\bar{n}$ TOTAL CROSS SECTIONS IN THE MOMENTUM REGION ABOVE 12 GEV/C.

ADDITIONAL FOOTNOTE REFERENCES

FORTSCHRITTE DER PHYSIK 9, 549 (1961), PHYS. REV. LETTERS 10, 262 (1963), PHYS. REV. LETTERS 13, 205 (1964), PHYS. REV. LETTERS 7, 185 (1961), PHYS. REV. LETTERS 7, 352 (1961), PHYS. REV. LETTERS 9, 576 (1960), PHYS. REV. LETTERS 7, 127 (1961), PHYS. REV. LETTERS 8, 173 (1962), PHYS. REV. LETTERS 9, 32 (1962), NUOVO CIMENTO 34, 825 (1964), PHYS. REV. 129, 2285 (1963), PHYS. REV. 123, 1850 (1961), INT'L CONGRESS ON NUCLEAR PHYSICS, PARIS, FRANCE 162 (1964), JETP 15 272 (1962), JETP 18 1239 (1963), PHYS. REV. LETTERS 3, 285 (1959), PHYS. REV. 128, 869 (1962), PHYS. REV. 100, 242 (1955), PHYS. REV. 123, 320 (1961), PHYS. REV. LETTERS 7, 182 (1961), JETP 34 499 (1958), NUOVO CIMENTO 14, 951 (1959), NUOVO CIMENTO 18, 947 (1960), PHYS. REV. LETTERS 9, 108 (1962), PHYS. REV. LETTERS 10, 376 (1963), PHYS. REV. LETTERS 12, 543 (1963), PHYS. REV. LETTERS 11, 425 (1963), PHYS. REV. LETTERS 11, 503 (1963), UCRL 3421 (1956), NUCLEAR ELECTRONICS 1, 63 (1962), REVIEW OF SCIENTIFIC INSTRUMENTS 25, 1070 (1954), PHYS. REV. 135, 8358 (1964), PHYS. LETTERS 14, 164 (1965), PHYS. REV. 134, 8633 (1964), AND PHYS. REV. 134, 8630 (1964).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 3/68, AND VERIFIED BY LEROY PRICE.

BEAM NO. 1 IS π^+ ON PROTON FROM 6 TO 20 GEV/C.
NO. 2 IS π^- ON PROTON FROM 6 TO 20 GEV/C.
NO. 3 IS π^+ ON DEUTERON FROM 6 TO 20 GEV/C.
NO. 4 IS π^- ON DEUTERON FROM 6 TO 20 GEV/C.
NO. 5 IS K^+ ON DEUTERON FROM 6 TO 20 GEV/C.
NO. 6 IS K^+ ON PROTON FROM 6 TO 20 GEV/C.
NO. 7 IS K^- ON DEUTERON FROM 6 TO 18 GEV/C.
NO. 8 IS K^- ON PROTON FROM 6 TO 20 GEV/C.
NO. 9 IS PROTON ON DEUTERON FROM 6 TO 22 GEV/C.
NO. 10 IS PROTON ON PROTON FROM 6 TO 22 GEV/C.
NO. 11 IS ANTI-PROTON ON DEUTERON FROM 6 TO 18 GEV/C.
NO. 12 IS ANTI-PROTON ON PROTON FROM 6 TO 18 GEV/C.

THIS EXPERIMENT USES COUNTERS.

KEY WORDS = CROSS SECTION

K^+ DEUTERON TOTAL CROSS SECTION.

LABORATORY BEAM MOMENTUM GEV/C	PER CENT	MILLI-BARNS
6. +- 1.75	33.4 +- .3	
8. 1.75	33.9 .3	
10. 1.75	33.8 .3	
12. 1.75	33.9 .3	
14. 1.75	33.8 .3	
16. 1.75	33.4 .3	
18. 1.75	33.7 .3	
20. 1.75	34.2 .3	

K+ PROTON TOTAL CROSS SECTION.

LABORATORY BEAM MOMENTUM GEV/C		MILLI-BARNS	
	PER CENT		
6.	1.75	17.0	±.1
8.	1.75	17.3	.1
10.	1.75	17.3	.1
12.	1.75	17.3	.1
14.	1.75	17.4	.1
16.	1.75	17.0	.1
18.	1.75	17.1	.1
20.	1.75	17.5	.1

K+ NEUTRON TOTAL CROSS SECTION.

(UNFOLDED FROM DEUTERIUM DATA)

Glauber correction applied. Assume $\text{Re } \{f(0)\} = 0$.

LABORATORY BEAM MOMENTUM GEV/C		MILLI-BARNS	
	PER CENT		
6.	1.75	17.5	±.4
8.	1.75	17.6	.4
10.	1.75	17.5	.4
12.	1.75	17.6	.4
14.	1.75	17.5	.4
16.	1.75	17.4	.4
18.	1.75	17.6	.4
20.	1.75	17.7	.4

K- DEUTERON TOTAL CROSS SECTION.

LABORATORY BEAM MOMENTUM GEV/C		MILLI-BARNS	
	PER CENT		
6.	1.75	44.1	±.3
8.	1.75	41.7	.3
10.	1.75	41.5	.3
12.	1.75	40.3	.3
14.	1.75	40.1	.3
16.	1.75	40.1	.4
18.	1.75	39.9	.7

K- PROTON TOTAL CROSS SECTION.

LABORATORY BEAM MOMENTUM GEV/C		MILLI-BARNS	
	PER CENT		
6.	1.75	24.0	±.3
8.	1.75	23.6	.2
10.	1.75	22.5	.2
12.	1.75	21.6	.2
14.	1.75	21.5	.2
16.	1.75	21.2	.4
18.	1.75	21.0	.8
20.	1.75	22.4	4.6

K- NEUTRON TOTAL CROSS SECTION.

(UNFOLDED FROM DEUTERIUM DATA)

Glauber correction applied.

LABORATORY BEAM MOMENTUM GEV/C		MILLI-BARNS	
	PER CENT		
6.	1.75	21.9	±.4
8.	1.75	19.7	.4
10.	1.75	20.6	.4
12.	1.75	20.2	.4
14.	1.75	20.1	.4
16.	1.75	20.3	.6
18.	1.75	20.3	1.1

PI+ PROTON TOTAL CROSS SECTION.

LABORATORY BEAM MOMENTUM GEV/C		MILLI-BARNS	
	PER CENT		
6.	1.75	26.2	±.2
8.	1.75	25.1	.2
10.	1.75	24.8	.2
12.	1.75	24.2	.2
14.	1.75	23.9	.2
16.	1.75	23.4	.2
18.	1.75	23.5	.2
20.	1.75	23.4	.2

PI- PROTON TOTAL CROSS SECTION.

LABORATORY BEAM MOMENTUM GEV/C		MILLI-BARNS
PER CENT		
6. +- 1.75	28.5 +- .3	
8. 1.75	27.5 .3	
10. 1.75	26.5 .3	
12. 1.75	25.9 .3	
14. 1.75	25.4 .3	
16. 1.75	25.1 .3	
18. 1.75	25.0 .3	
20. 1.75	24.8 .3	

PI+ DEUTERON TOTAL CROSS SECTION.

LABORATORY BEAM MOMENTUM GEV/C		MILLI-BARNS
PER CENT		
6. +- 1.75	52.8 +- .5	
8. 1.75	50.5 .5	
10. 1.75	49.3 .5	
12. 1.75	48.2 .5	
14. 1.75	46.9 .5	
16. 1.75	46.6 .5	
18. 1.75	46.3 .5	
20. 1.75	45.9 .5	

PI- DEUTERON TOTAL CROSS SECTION.

LABORATORY BEAM MOMENTUM GEV/C		MILLI-BARNS
PER CENT		
6. +- 1.75	52.7 +- .5	
8. 1.75	51.0 .5	
10. 1.75	49.3 .5	
12. 1.75	47.9 .5	
14. 1.75	47.1 .5	
16. 1.75	46.4 .5	
18. 1.75	46.4 .5	
20. 1.75	45.8 .5	

PROTON DEUTERON TOTAL CROSS SECTION.

LABORATORY BEAM MOMENTUM GEV/C		MILLI-BARNS
PER CENT		
6. +- 1.75	77.4 +- 1.3	
8. 1.75	76.2 1.3	
10. 1.75	75.8 1.3	
12. 1.75	74.4 1.3	
14. 1.75	74.0 1.3	
16. 1.75	73.7 1.3	
18. 1.75	72.8 1.3	
20. 1.75	72.1 1.3	
22. 1.75	71.6 1.3	

PROTON PROTON TOTAL CROSS SECTION.

LABORATORY BEAM MOMENTUM GEV/C		MILLI-BARNS
PER CENT		
6. +- 1.75	40.6 +- .6	
8. 1.75	40.0 .6	
10. 1.75	39.9 .6	
12. 1.75	39.4 .6	
14. 1.75	39.1 .6	
16. 1.75	38.7 .6	
18. 1.75	38.7 .6	
20. 1.75	38.4 .6	
22. 1.75	38.3 .6	

NEUTRON PROTON TOTAL CROSS SECTION.

(UNFOLDED FROM DEUTERIUM DATA)

LABORATORY BEAM MOMENTUM GEV/C		MILLI-BARNS
PER CENT		
6. +- 1.75	42.6 +- 1.7	
8. 1.75	41.8 1.7	
10. 1.75	41.5 1.7	
12. 1.75	40.4 1.7	
14. 1.75	40.2 1.7	
16. 1.75	40.2 1.7	
18. 1.75	39.2 1.7	
20. 1.75	38.7 1.7	
22. 1.75	38.2 1.7	

PBAR DEUTERON TOTAL CROSS SECTION.

LABORATORY BEAM MOMENTUM GEV/C	PER CENT	MILLI-BARNS
6. $\pm$ 1.75		106.9 $\pm$ 1.3
8. 1.75		102.7 1.3
12. 1.75		96.1 1.3
14. 1.75		95.0 1.4
16. 1.75		93.2 1.6
18. 1.75		87.2 6.1

PBAR PROTON TOTAL CROSS SECTION.

LABORATORY BEAM MOMENTUM GEV/C	PER CENT	MILLI-BARNS
6. $\pm$ 1.75		59.3 $\pm$ 1.1
8. 1.75		56.4 .8
16. 1.75		49.2 .8
18. 1.75		50.3 3.6
12. 1.75		51.7 .8
14. 1.75		50.7 .9

NEUTRON PBAR TOTAL CROSS SECTION.

(UNFOLDED FROM DEUTERIUM DATA)

Glauber correction applied.

LABORATORY BEAM MOMENTUM GEV/C	PER CENT	MILLI-BARNS
6. $\pm$ 1.75		59.5 $\pm$ 4.0
8. 1.75		57.3 3.9
17. 1.75		43.8 3.7
14. 1.75		53.4 3.7
16. 1.75		52.7 3.7
18. 1.75		44.4 9.0

40 NEW STRUCTURES IN THE K⁺ P AND K⁺ D TOTAL CROSS SECTIONS BETWEEN 1.55 AND 3.30 GEV/C. (PHYS. REV. LETTERS 19, 259 (1967))

R.J.ABRAMS, R.L.COOL, G.GIACOMELLI, T.F.KYCIA, B.A.LEONTIC, K.K.LI, D.N.MICHAEL (BROOKHAVEN NAT. LAB., UPTON, L.I., N. Y., USA)

ABSTRACT TWO NEW I=1 STRUCTURES ABOUT 0.2 MB IN HEIGHT ARE OBSERVED IN K⁺ P AND K⁺ D TOTAL CROSS SECTION MEASUREMENTS AT C.M. ENERGIES OF 2190 $\pm$ 10 MEV.

ADDITIONAL FOOTNOTE REFERENCES

PHYS. REV. LETTERS 18, 1209 (1967), PHYS. REV. LETTERS 17, 102 (1966), PHYS. REV. LETTERS 2, 117 (1959), PHYS. REV. LETTERS 7, 182 (1961), RHEI. AM. PHYS. SOC. 17, A6 (1967), AND PHYS. REV. LETTERS 19, 259 (1967).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 4/68, AND VERIFIED BY LEROY PRICE.

BEAM NO. 1 IS K⁺ ON PROTON FROM 1.55 TO 3.30 GEV/C.
NO. 2 IS K⁺ ON DEUTERON FROM 1.55 TO 3.30 GEV/C.

THIS EXPERIMENT USES COUNTERS.

KEY WORDS * CROSS SECTION

K⁺ PROTON TOTAL CROSS SECTION.

ORIGINAL DATA			CORRECTED VALUES [1]		
LABORATORY BEAM MOMENTUM GEV/C	PER CENT	MILLI-BARNS [2]	LABORATORY BEAM MOMENTUM GEV/C	PER CENT	MILLI-BARNS [3]
1.55 $\pm$.75		18.03 $\pm$.25	1.55 $\pm$.75		17.70 $\pm$.11
1.60 .75		17.95 .25	1.60 .75		17.71 .10
1.70 .75		17.95 .25	1.70 .75		17.73 .08
1.75 .75		18.04 .25	1.75 .75		17.83 .07
1.80 .75		18.20 .25	1.80 .75		17.98 .10
1.85 .75		18.08 .25	1.85 .75		17.77 .07
1.90 .75		18.08 .25	1.90 .75		17.79 .07
2.00 .75		17.79 .25	1.95 .75		17.75 .07
2.05 .75		17.89 .25	2.00 .75		17.83 .06
2.10 .75		17.71 .25	2.05 .75		17.72 .07
2.15 .75		17.71 .25	2.10 .75		17.56 .07
2.20 .75		17.75 .25	2.15 .75		17.57 .07
2.30 .75		17.65 .25	2.20 .75		17.60 .07
2.35 .75		17.65 .25	2.30 .75		17.44 .06
2.40 .75		17.68 .25	2.35 .75		17.52 .04
2.45 .75		17.59 .25	2.40 .75		17.56 .06
2.50 .75		17.59 .25	2.45 .75		17.48 .04
2.55 .75		17.54 .25	2.50 .75		17.49 .05
2.60 .75		17.59 .25	2.55 .75		17.44 .04
2.65 .75		17.57 .25	2.60 .75		17.50 .04
2.70 .75		17.50 .25	2.65 .75		17.47 .04
2.75 .75		17.49 .25	2.70 .75		17.41 .04
2.80 .75		17.48 .25	2.75 .75		17.41 .04
2.85 .75		17.38 .25	2.80 .75		17.40 .04
2.90 .75		17.40 .25	2.85 .75		17.30 .04
2.95 .75		17.27 .25	2.90 .75		17.34 .03
3.00 .75		17.26 .25	2.95 .75		17.30 .04
3.05 .75		17.22 .25	3.00 .75		17.19 .04
3.10 .75		17.08 .25	3.05 .75		17.14 .04
3.15 .75		17.19 .25	3.10 .75		17.08 .05
3.20 .75		17.16 .25	3.15 .75		17.19 .04
3.25 .75		17.16 .25	3.20 .75		17.13 .03
3.30 .75		17.16 .25	3.25 .75		17.13 .03
			3.30 .75		17.14 .03

[1] PRIVATE COMMUNICATION FROM T.KYCIA, APRIL, 1969.

[2] PLUS POSSIBLE SYSTEMATIC ERROR OF $\pm$ 1 PER CENT.[3] PLUS POSSIBLE SYSTEMATIC ERROR OF $\pm$ 1 PER CENT.

K+ DEUTERON TOTAL CROSS SECTION.

ORIGINAL DATA			CORRECTED VALUES [1]		
LABORATORY BEAM MOMENTUM GEV/C	PER CENT	MILLI-BARNS [2] PER CENT	LABORATORY BEAM MOMENTUM GEV/C	PER CENT	MILLI-BARNS [3]
1.55	±.75	36.30 ±.13	1.55	±.75	36.03 ±.11
1.60	.75	36.10 .13	1.60	.75	35.94 .11
1.70	.75	36.00 .13	1.70	.75	35.78 .08
1.75	.75	36.10 .13	1.75	.75	35.90 .09
1.80	.75	35.82 .13	1.80	.75	35.68 .10
1.85	.75	35.79 .13	1.85	.75	35.63 .07
1.90	.75	35.84 .13	1.90	.75	35.70 .07
1.95	.75	35.81 .13	1.95	.75	35.69 .06
2.00	.75	35.79 .13	2.00	.75	35.68 .05
2.10	.75	35.42 .13	2.05	.75	35.55 .07
2.15	.75	35.37 .13	2.10	.75	35.34 .08
2.20	.75	35.31 .13	2.15	.75	35.28 .07
2.30	.75	35.00 .13	2.20	.75	35.28 .07
2.35	.75	35.05 .13	2.30	.75	34.98 .07
2.40	.75	34.95 .13	2.35	.75	35.05 .06
2.45	.75	35.00 .13	2.40	.75	34.92 .06
2.50	.75	34.84 .13	2.45	.75	35.00 .05
2.55	.75	34.84 .13	2.50	.75	34.87 .06
2.60	.75	34.79 .13	2.55	.75	34.87 .04
2.65	.75	34.74 .13	2.60	.75	34.85 .05
2.70	.75	34.74 .13	2.65	.75	34.75 .05
2.75	.75	34.65 .13	2.70	.75	34.77 .04
2.80	.75	34.63 .13	2.75	.75	34.72 .05
2.85	.75	34.42 .13	2.80	.75	34.67 .05
2.90	.75	34.37 .13	2.85	.75	34.52 .05
2.95	.75	34.32 .13	2.90	.75	34.48 .04
3.00	.75	34.21 .13	2.95	.75	34.42 .04
3.05	.75	34.21 .13	3.00	.75	34.35 .04
3.10	.75	34.16 .13	3.05	.75	34.34 .04
3.15	.75	34.00 .13	3.10	.75	34.30 .04
3.20	.75	33.95 .13	3.15	.75	34.14 .04
3.25	.75	34.00 .13	3.20	.75	34.12 .04
3.30	.75	33.95 .13	3.25	.75	34.20 .04
			3.30	.75	34.13 .04

[1] PRIVATE COMMUNICATION FROM T.KYCIA, APRIL, 1969.

[2] PLUS POSSIBLE SYSTEMATIC ERROR OF ± 1 PER CENT.

[3] PLUS POSSIBLE SYSTEMATIC ERROR OF ± 1 PER CENT.

I = 0 K NUCLEON TOTAL CROSS SECTION.

(PRIVATE COMMUNICATION FROM T.KYCIA APRIL 1969)

LABORATORY BEAM MOMENTUM GEV/C	PER CENT	MILLI-BARNS [1]
1.55	±.75	20.11 ±.40
1.60	.75	20.07 .37
1.70	.75	19.93 .29
1.75	.75	19.89 .28
1.80	.75	19.08 .36
1.85	.75	19.54 .25
1.90	.75	19.59 .25
1.95	.75	19.65 .24
2.00	.75	19.95 .21
2.05	.75	19.37 .25
2.10	.75	19.42 .26
2.15	.75	19.28 .25
2.20	.75	19.21 .25
2.30	.75	19.10 .23
2.35	.75	19.00 .17
2.40	.75	18.62 .22
2.45	.75	19.02 .16
2.50	.75	18.74 .19
2.55	.75	18.90 .14
2.60	.75	18.68 .16
2.65	.75	18.57 .16
2.70	.75	18.80 .14
2.75	.75	18.71 .16
2.80	.75	18.62 .16
2.85	.75	18.60 .16
2.90	.75	18.36 .17
2.95	.75	18.31 .14
3.00	.75	18.42 .14
3.05	.75	18.52 .14
3.10	.75	18.61 .17
3.15	.75	18.07 .14
3.20	.75	18.09 .12
3.25	.75	18.19 .12
3.30	.75	17.95 .12

[1] PLUS POSSIBLE SYSTEMATIC ERROR OF ± 1 PER CENT.

K⁺ NEUTRON TOTAL CROSS SECTION.

(PRIVATE COMMUNICATION FROM KYCIA APRIL 1969)

LABORATORY BEAM MOMENTUM GEV/C	PER CENT	MILLI-BARNS (1)
1.55 +- .75		18.91 +- .16
1.60 -.75		18.89 .15
1.70 .75		18.83 .11
1.75 .75		18.86 .11
1.80 .75		18.53 .14
1.85 .75		18.66 .10
1.90 .75		18.69 .10
1.95 .75		18.70 .09
2.00 .75		18.79 .08
2.05 .75		18.54 .10
2.10 .75		18.49 .11
2.15 .75		18.43 .10
2.20 .75		18.40 .10
2.30 .75		18.27 .09
2.35 .75		18.26 .07
2.40 .75		18.09 .08
2.45 .75		18.25 .06
2.50 .75		18.11 .08
2.55 .75		18.17 .06
2.60 .75		18.09 .06
2.65 .75		18.02 .06
2.70 .75		18.11 .06
2.75 .75		18.06 .06
2.80 .75		18.01 .06
2.85 .75		17.95 .06
2.90 .75		17.85 .05
2.95 .75		17.81 .06
3.00 .75		17.81 .06
3.05 .75		17.83 .06
3.10 .75		17.85 .06
3.15 .75		17.61 .06
3.20 .75		17.61 .05
3.25 .75		17.66 .05
3.30 .75		17.55 .05

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF +- 1 PER CENT.

41 FURTHER EVIDENCE FOR BARYON EXCHANGE FROM K⁺P BACKWARD ELASTIC SCATTERING. (PHYS. REV. LETTERS 19, 675 (1967))

D. CLINE, C. MOORE, D. REEDER (UNIV. OF WISCONSIN, MADISON, WISC., USA)

ABSTRACT WE HAVE STUDIED K⁺P ELASTIC SCATTERING NEAR THE BACKWARD DIRECTION AT A BEAM MOMENTUM OF 3.5 GEV/C. IN ADDITION TO PROVIDING EVIDENCE FOR BARYON EXCHANGE, THE EXPERIMENTAL RESULTS SHOW A POSSIBLE TURNOVER OF THE ANGULAR DISTRIBUTION FOR POSITIVE VALUES OF θ WHICH IS INTERPRETED IN TERMS OF A REGGEIZED BARYON EXCHANGE MODE. WE HAVE MADE A COMPARISON OF K⁺P, P⁺P, AND P⁺P BACKWARD SCATTERING WITHIN THE FRAMEWORK OF SU(3) WHICH LEADS TO PREDICTIONS CONCERNING OTHER, AS YET UNMEASURED, K⁺N PROCESSES.

ADDITIONAL FOOTNOTE REFERENCES

PHYS. LETTERS 22, 528 (1966), PHYS. REV. LETTERS 16, 913 (1966), PHYS. LETTERS 248, 317 (1967), PHYS. LETTERS 21, 117 (1966), NUOVO CIMENTO 45A, 805 (1966), PHYS. REV. LETTERS 16, 1228 (1966), PHYS. REV. LETTERS 17, 102 (1966), NUOVO CIMENTO 23, 1061 (1962), NUOVO CIMENTO 41A, 481 (1966), PHYS. LETTERS 23, 605 (1966), PHYS. REV. LETTERS 16, 28 (1966), BULL. AM. PHYS. SOC. 12, 469 (1967), AND PHYS. REV. 153, 1575 (1967).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 4/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K⁺ ON PROTON AT 3.53 GEV/C.

THIS EXPERIMENT USES THE B.N.L. 80 IN. (H) BUBBLE CHAMBER.

KEY WORDS = CROSS SECTION ANGULAR DISTRIBUTION

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = 3.53 GEV/C.
NUMBER OF EVENTS = 45.

THIS DATA WAS READ FROM A GRAPH.

COS(THETA)	D-SIGMA/D-OMEGA IN/5R
-0.54 +- .14	1.67 +- .90
-0.74 .06	3.00 1.50
-0.85 .05	3.40 1.40
-0.92 .02	10.00 3.20
-0.96 .02	12.50 5.00

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.

42 PRODUCTION OF K PI PI RESONANCES AT HIGH ENERGY. (PHYS. REV. LETTERS 18, 1087 (1967))

J. C. BERLIN, H. EAT, M. J. FARBER, I. PERDEL, D. E. UHLMAN, A. L. MELISSINUS, T. YAMANOUCHI, H. YUTA (UNIV. OF ROCHESTER, ROCHESTER, N. Y., USA)

ABSTRACT THE PREVIOUSLY OBSERVED ENHANCEMENT BETWEEN 1100 AND 1400 MEV IN THE K PI PI MASS SPECTRUM IS INTERPRETED AS $T = 1/2$, $J^P = 1^+$ RESONANCE AT 1280 +- 20 MEV AND WIDTH OF 130 PLUS OR MINUS 15 MEV WHICH DECAYS INTO PI⁺ + K⁰ AND INTO K⁺ + RHO⁰. EVIDENCE IS ALSO PRESENTED SUPPORTING THE EXISTENCE OF THE NEWLY-REPORTED L MESON.

ADDITIONAL FOOTNOTE REFERENCES

UCRL 16462 (1965), LRL INTERDEPARTMENTAL REPORT UCID 2873 (1966), PHYS. REV. LETTERS 17, 726 (1966), PHYS. REV. LETTERS 16, 1069 (1966), PHYS. LETTERS 22, 357 (1966), CERN TC/65-19, CERN TC/65-17, BULL. AM. PHYS. SOC. 12, 44 (1967), UCRL 8030 (1967), UCRL 16744 (1966), PHYS. REV. LETTERS 13, 169 (1964), AND PHYS. LETTERS 9, 207 (1964).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 4/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K⁺ ON PROTON AT 12.7 GEV/C.

THIS EXPERIMENT USES THE B.N.L. 80 IN. (H) BUBBLE CHAMBER.

KEY WORDS = K*(1320) L MESON MASS SPECTRUM ANGULAR DISTRIBUTION O PEAK

NO DATA PUNCHED FOR THIS ARTICLE

43 COHERENT K^+ PRODUCTION IN THE K^+ D REACTION AT 2.3 GEV/C. [PHYS. REV. LETTERS 15, 500 (1955)]

I. BUTTERWORTH, J. L. BROWN, G. GOLDBABER, S. GOLDBABER, A. A. MIRATA, J. A. KADYK, G. H. TRILLING [U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA, AND UNIV. OF CALIFORNIA, BERKELEY, CALIF., USA]

ADDITIONAL FOOTNOTE REFERENCES

NUCLEAR INSTRUMENTS AND METHODS 30, 37 (1963), NUOVO CIMENTO 11, 670 (1959), NUOVO CIMENTO 36, 1101 (1965), UCRL 11465 (1964), ATHENS CONFERENCE 92 (1963), PHYS. LETTERS 6, 6 (1963), NUC. PHYS. 29, 680 (1952), PROC. OF THE ROYAL SOCIETY OF LONDON A244, 491 (1958), AND PHYS. REV. LETTERS 15, 125 (1965).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 4/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K^+ ON DEUTERON AT 2.3 GEV/C.

THIS EXPERIMENT USES THE B.N.L. 20 IN. (III) BUBBLE CHAMBER. A TOTAL OF 100000 PICTURES ARE REPORTED ON.

KEY WORDS = CROSS SECTION DALITZ PLOT ANGULAR DISTRIBUTION MASS SPECTRUM K^+ (890) $\Delta(1238)$

CROSS SECTION FOR K^+ DEUTERON $\rightarrow$ K^0 π^+ DEUTERON.

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
2.3	150. $\pm$ 35.	33

CROSS SECTION FOR K^+ DEUTERON $\rightarrow$ K^+ π^- π^+ DEUTERON.

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
2.3	110. $\pm$ 16.	81

DIFFERENTIAL CROSS SECTION FOR K^+ DEUTERON $\rightarrow$ K^0 π^+ DEUTERON.

LABORATORY BEAM MOMENTUM = 2.3 GEV/C.

.....
THIS DATA WAS READ FROM A GRAPH.
.....

$-T$ (GEV/C)**2	D-SIGMA/D-T UB/(GEV/C)**2 (1)	NO. EVENTS
MIN	MAX	
.02	.04	18.0 $\pm$ 9.0
.04	.06	31.5 11.9
.06	.08	9.0 6.4
.08	.10	4.5 4.5
.10	.12	4.5 4.5
.12	.14	13.5 7.0
.14	.16	9.0 6.4
.16	.18	4.5 4.5
.18	.20	4.5 4.5
.20	.22	4.5 4.5
.22	.32	.0 4.5
.32	.34	4.5 4.5

T IS THE MOMENTUM TRANSFER BETWEEN THE [INCOMING K^+] AND THE [K^0 AND π^+].

(1) COUNTS WERE MULTIPLIED BY 4.5 TO GET THESE. ERRORS ARE TAKEN AS PROPORTIONAL TO THE SQUARE-ROOT OF THE COUNTS.

44 K π 0-NUCLEON INTERACTION IN THE $I=0$ STATE AT LOW ENERGIES. [PHYS. REV. 134, B1111 (1964)]

V. J. STENGER, W. E. SLATER, D. H. STORK, H. K. TICHON [UNIV. OF CALIF., LOS ANGELES, CALIF., USA]
G. GOLDBABER, S. GOLDBABER [U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA, AND UNIV. OF CALIFORNIA, BERKELEY, CALIF., USA]

ABSTRACT THE STUDY OF THE $I=0$ $K\pi$ INTERACTION AT LOW ENERGIES USING K^+ DEUTERIUM SCATTERING HAS BEEN EXTENDED ON THE EXPERIMENTAL SIDE BY THE INCLUSION OF ELASTIC AND INELASTIC NONCHARGE-EXCHANGE SCATTERING, AND ON THE THEORETICAL SIDE BY A MORE DETAILED INVESTIGATION OF THE FINAL-STATE INTERACTION BETWEEN THE TWO NUCLEONS. THE NEW RESULT FOR THE $I=0$ PHASE SHIFTS EXHIBIT THE SAME TRENDS AS THOSE PUBLISHED BEFORE; HOWEVER, THE NEW S-WAVE DATA ARE MORE ACCURATE. IT IS SHOWN THAT AT VERY LOW ENERGIES, THE $I=0$ S-WAVE SCATTERING MAY BE INTERPRETED AS DUE TO A VERY SMALL SCATTERING LENGTH $A_0 = +0.04 \pm 0.04$ F. THIS SHOULD BE CONTRASTED WITH $I=1$ SCATTERING LENGTH $A_1 = -0.31 \pm 0.01$ F.

CROSS REFERENCES

CONTINUATION OF PREVIOUS EXPERIMENT IN PHYS. REV. LETTERS 7, 378 (1961).

ADDITIONAL FOOTNOTE REFERENCES

PHYS. REV. LETTERS 9, 135 (1962), NUOVO CIMENTO 12, 241 (1959), PHYS. REV. 118, 553 (1960), PHYS. REV. 109, 2133 (1958), PHYS. REV. 109, 2121 (1958), PHYS. REV. 113, 1303 (1959), PHYS. REV. 120, 1898 (1960), PHYS. REV. LETTERS 7, 188 (1961), UCRL 10950, PHYS. REV. 129, 2743 (1963), PHYS. REV. LETTERS 1, 112 (1958), PHYS. REV. LETTERS 5, 108 (1950), NUOVO CIMENTO 10, 205 (1958), NUOVO CIMENTO 10, 168 (1958), PHYS. REV. LETTERS 7, 378 (1961), UCRL 9691, PHYS. REV. 80, 196 (1950), PHYS. REV. 84, 1084 (1951), PHYS. REV. 105, 256 (1957), PHYS. REV. 115, 1727 (1959), NOTAS DE FISICA 8, 4 (1961), NUOVO CIMENTO 23, 1163 (1962), LRL INTERDEPARTMENTAL REPORT REV-403 (1960), PHYS. REV. LETTERS 7, 346 (1961), NUOVO CIMENTO 11, 670 (1959), NUOVO CIMENTO 14, 722 (1959), PHYS. REV. LETTERS 5, 266 (1960), UCRL 6082, AND PHYS. REV. 122, 1606 (1961).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 3/60, AND VERIFIED BY LEROY PRICE.

BEAM IS K^+ ON DEUTERON FROM .23 TO .53 GEV/C.

THIS EXPERIMENT USES THE L.R.L. 15 IN. (II) BUBBLE CHAMBER.

GENERAL COMMENTS ON THIS ARTICLE

1 LUMPS TOGETHER FINAL STATES K^0 WITH $K^+\pi^0$

KEY WORDS = ANGULAR DISTRIBUTION CROSS SECTION PHASE SHIFT

CROSS SECTION FOR $K^+ \text{ DEUTERON} = K^+ \text{ DEUTERON} + K^+ \text{ PROTON (NEUTRON SPECTATOR)} + K^+ \text{ NEUTRON (PROTON SPECTATOR)}$.
 DATA IS INTEGRATED OVER THETA FROM 5. TO 180. DEGREES. THETA IS THE ANGLE THAT THE K^+ MAKES WITH THE BEAM IN THE LAB.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.377 $\pm$.018	19.7 $\pm$ 1.9
.530 $\pm$.015	18.9 $\pm$ 1.8

ELASTIC DIFFERENTIAL CROSS SECTION FOR K^+ NEUTRON.

LABORATORY BEAM MOMENTUM = .377 $\pm$.018 GEV/C.

 * THIS DATA WAS READ FROM A GRAPH *

COS(THETA)		D-SIGMA/D-OMEGA MB/SR
MIN	MAX	
-1.000	-.600	1.9 $\pm$.3
-.600	-.200	1.4 $\pm$.3
-.200	.200	1.4 $\pm$.3
.200	.600	1.5 $\pm$.3
.600	.996	1.7 $\pm$.3

THETA IS THE ANGLE THAT THE K^+ MAKES WITH THE BEAM IN THE GRAND C.M.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K^+ NEUTRON.

LABORATORY BEAM MOMENTUM = .530 $\pm$.015 GEV/C.

 * THIS DATA WAS READ FROM A GRAPH *

COS(THETA)		D-SIGMA/D-OMEGA MB/SR
MIN	MAX	
-1.000	-.600	1.40 $\pm$.26
-.600	-.200	1.45 $\pm$.26
-.200	.200	1.16 $\pm$.25
.200	.600	1.16 $\pm$.25
.600	.996	2.23 $\pm$.38

THETA IS THE ANGLE THAT THE K^+ MAKES WITH THE BEAM IN THE GRAND C.M.

45 COMPARISON OF K^+ PRODUCTION IN THE REACTIONS $K^+ N \rightarrow K^+ \pi^- p$ AND $K^+ p \rightarrow K^0 \pi^+ p$. [PHYS. REV. LETTERS 15, 737 (1965)]
 S. GOLDBABER, J. L. BROWN, I. BUTTERWORTH, G. GOLDBABER, A. A. MIKATA, J. A. KADYK, G. H. TRILLING (U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA, AND UNIV. OF CALIFORNIA, BERKELEY, CALIF., USA)

ADDITIONAL FOOTNOTE REFERENCES
 PHYS. REV. LETTERS 12, 522 (1964), PHYS. LETTERS 18, 76 (1965), PHYS. REV. 139, 8428 (1965), AND NUOVO CIMENTO 36, 1101 (1965).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 4/68, AND VERIFIED BY ROGER BLAND.

BEAM IS K^+ ON DEUTERON AT 2.3 GEV/C.

THIS EXPERIMENT USES THE S.N.L. 20 IN. (H) BUBBLE CHAMBER. A TOTAL OF 10000 PICTURES ARE REPRODUCED ON:

GENERAL COMMENTS ON THIS ARTICLE
 1 A SPECTATOR PROTON CUT IS MADE AT 300 MEV/C.

KEY WORDS = CROSS SECTION DALITZ PLOT ANGULAR DISTRIBUTION DELTA(1238) N*(1512) N*(1688)
 $K^*(890)$ DENSITY MATRIX MODELS

CROSS SECTION FOR K^+ NEUTRON $\rightarrow K^+ \pi^-$ PROTON.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS
2.3	3.0 $\pm$.2	1576

CROSS SECTION FOR K^+ PROTON $\rightarrow K^0 \pi^+ \pi^-$ PROTON.

* * THIS DATA SHOULD NOT BE USED * * / NO DEUTERON CORRECTIONS MADE.)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS
2.3	3.0 $\pm$.3	602

CROSS SECTION FOR K^+ NEUTRON $\rightarrow K^*(890) \pi^-$ PROTON.
 $K^*(890) \rightarrow K^+ \pi^-$ [1]

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.3	1.4 $\pm$.2

[1] USED SIMPLE MASS CUT [MASS = .920 GEV; WIDTH = .100 GEV].

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)^+ \text{ PROTON}$
 $K^*(890)^+ \rightarrow K^0 \pi^+ \{1\}$
 * * THIS DATA SHOULD NOT BE USED * * (NO DEUTERON CORRECTIONS MADE.)

LABORATORY
 BEAM MOMENTUM
 GEV/C 2.3
 MILLI-BARNS 1.3 $\pm$.2

\{1\} USED SIMPLE MASS CUT (MASS = .920 GEV; WIDTH = .100 GEV).

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)^+ \text{ PROTON}$
 $K^*(890)^+ \rightarrow K^0 \pi^+ \{1\}$

LABORATORY BEAM MOMENTUM = 2.3 GEV/C.

* * THIS DATA SHOULD NOT BE USED * * (NO DEUTERON CORRECTIONS MADE.)

-T (GEV/C)**2		NO. EVENTS
MIN	MAX	
.00	.04	8
.04	.08	21
.08	.12	21
.12	.16	19
.16	.20	17
.20	.24	14
.24	.28	19
.28	.32	15
.32	.36	15
.36	.40	9
.40	.44	9
.44	.48	11
.48	.52	10
.52	.56	9
.56	.60	1
.60	.64	7
.64	.68	6
.68	.72	6
.72	.76	3
.76	.80	3
.80	.84	4
.84	.88	4
.88	.92	1
.92	.96	3
.96	1.00	3

T IS THE MOMENTUM TRANSFER BETWEEN THE [INCOMING K+] AND THE [K*(890)+].

\{1\} USED SIMPLE MASS CUT (MASS = .920 GEV; WIDTH = .100 GEV).

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ NEUTRON} \rightarrow K^*(890)^0 \text{ PROTON}$
 $K^*(890)^0 \rightarrow K^+ \pi^- \{1\}$

LABORATORY BEAM MOMENTUM = 2.3 GEV/C.

-T (GEV/C)**2		NO. EVENTS
MIN	MAX	
.00	.04	49
.04	.08	149
.08	.12	95
.12	.16	52
.16	.20	54
.20	.24	40
.24	.28	34
.28	.32	34
.32	.36	26
.36	.40	20
.40	.44	18
.44	.48	9
.48	.52	9
.52	.56	7
.56	.60	11
.60	.64	9
.64	.68	8
.68	.72	10
.72	.76	8
.76	.80	3
.80	.84	9
.84	.88	2
.88	.92	3
.92	.96	2
.96	1.00	7

T IS THE MOMENTUM TRANSFER BETWEEN THE [INCOMING K+] AND THE [K*(890)0].

\{1\} USED SIMPLE MASS CUT (MASS = .920 GEV; WIDTH = .100 GEV).

46 SEARCH FOR EVIDENCE OF A $K^+ K^+$ RESONANCE. (PHYS. REV. LETTERS 16, 1053 (1966))

A.P. ERWIN, W.D. WALKER, A. WEINBERG (UNIV. OF WISCONSIN, MADISON, WISC., USA)

ADDITIONAL FOOTNOTE REFERENCES

PHYS. LETTERS 17, 155 (1965), PHYS. REV. LETTERS 8, 140 (1962), NUOVO CIMENTO 33, 309 (1964), PHYS. REV. LETTERS 12, 522 (1964), NUOVO CIMENTO 33, 906 (1964), PHYS. REV. 113, 1640 (1959), PHYS. REV. LETTERS 139, 81355 (1965), PHYS. REV. LETTERS 15, 805 (1965), PHYS. REV. LETTERS 13, 291 (1964), PHYS. REV. LETTERS 12, 399 (1964), PHYS. REV. 139, 8956 (1965), PHYS. REV. 130, 2455 (1963), PHYS. REV. LETTERS 11, 90 (1963), AND PHYS. REV. 134, 81099 (1964).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 5/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K^+ ON PROTON AT 3.535 GEV/C.

THIS EXPERIMENT USES THE B.N.L. 80 IN. (H) BUBBLE CHAMBER.

GENERAL COMMENTS ON THIS ARTICLE

1 SEE NO DIRECT EVIDENCE FOR $K^+ K^+$ RESONANCE BELOW 1.7 GEV

KEY WORDS - CROSS SECTION MASS SPECTRUM

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ K^+ \text{ LAMBDA} + K^+ K^+ \text{ SIGMA}^0$.

LABORATORY
BEAM MOMENTUM
GEV/C
3.535

MICRO-BARNS
31. $\pm$ 4.

NO. EVENTS
105

47 $K^+ P$ AND $\pi^+ P$ BACKWARD ELASTIC SCATTERING AT 5.2 AND 6.9 GEV/C. (PHYS. LETTERS 288, 291 (1968))

M.F. BAKER, K. BERKELMAN, P.J. CARLSON, G.P. FISHER, P. FLEURY, D. MARTILL, R. KALBACH, A. LUNDBY, S. MUKHIN, R. NIERHAUS, R.P. PRETZL, J. WOULDs (EUROPEAN ORG. FOR NUC. RES., GENEVA, SWITZERLAND)

ABSTRACT THE ANGULAR DISTRIBUTIONS OF $K^+ P$ AND $\pi^+ P$ ELASTIC SCATTERING WERE STUDIED AT 6.9 GEV/C. BACKWARD PEAKS ARE OBSERVED IN $K^+ P$ SCATTERING WITH AN ENERGY DEPENDENCE OF THE FORM $S \exp -4$.

ADDITIONAL FOOTNOTE REFERENCES

PHYS. REV. LETTERS 21, 389 (1968), PHYS. LETTERS 248, 317 (1967), PHYS. REV. LETTERS 19, 675 (1967), PHYS. REV. LETTERS 20, 607 (1968), PHYS. REV. LETTERS 21, 580 (1968), PHYS. REV. LETTERS 21, 396 (1968), NUOVO CIMENTO 45, 88 (1966), AND PHYS. REV. LETTERS 20, 689 (1968).

ARTICLE READ BY ODETTE BENARY IN 12/68, AND VERIFIED BY ROGER BLAND.

BEAM NO. 1 IS K^+ ON PROTON FROM 5.2 TO 6.9 GEV/C.

NO. 2 IS π^+ ON PROTON FROM 5.2 TO 6.9 GEV/C.

NO. 3 IS π^- ON PROTON AT 6.9 GEV/C.

NO. 4 IS K^- ON PROTON AT 6.9 GEV/C.

THIS EXPERIMENT USES SPARK CHAMBERS AND COUNTERS.

KEY WORDS - DIFFERENTIAL CROSS SECTION MODELS

ELASTIC DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON}$. (TABLE 1)

LABORATORY BEAM MOMENTUM = 5.2 GEV/C.

U (GEV/C) $\pm$ 2	D-SIGMA/D-U UB/(GEV/C) $\pm$ 2 (1)
0.00 $\pm$ 0.02	14.6 $\pm$ 5.7
-0.04 $\pm$ 0.02	14.0 4.0
-0.08 $\pm$ 0.02	9.5 3.0
-0.12 $\pm$ 0.02	12.1 2.8
-0.20 $\pm$ 0.05	3.4 1.1
-0.30 $\pm$ 0.05	3.3 1.2
-0.40 $\pm$ 0.05	4.8 1.6
-0.50 $\pm$ 0.05	3.5 1.5
-0.60 $\pm$ 0.05	1.4 1.4
-0.70 $\pm$ 0.05	0 1.3
-0.80 $\pm$ 0.05	0 4.7

U IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K^+) AND THE (PROTON).

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF $\pm$ 15 PER CENT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON}$. (TABLE 1)

LABORATORY BEAM MOMENTUM = 6.9 GEV/C.

U (GEV/C) $\pm$ 2	D-SIGMA/D-U UB/(GEV/C) $\pm$ 2 (1)
-0.05 $\pm$ 0.05	2.5 $\pm$ 1.1
-0.20 $\pm$ 0.10	2.1 0.7
-0.40 $\pm$ 0.10	1.4 0.6
-0.60 $\pm$ 0.10	0.7 0.5
-0.80 $\pm$ 0.10	0.3 0.5
-1.00 $\pm$ 0.10	0 0.4

U IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K^+) AND THE (PROTON).

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF $\pm$ 15 PER CENT.

48 RESONANCE PRODUCTION IN K⁺P INTERACTIONS AT 3.5 BEV/C. (THESIS (1967))

J.M.BISHOP (UNIV. OF WISCONSIN, MADISON, WISC., USA)

ADDITIONAL FOOTNOTE REFERENCES

REV. MOD. PHYS. 39, 1 (1967), PHYS. REV. LETTERS 8, 447 (1962), PHYS. LETTERS 5, 279 (1963), PHYS. LETTERS 12, 255 (1964), PHYS. LETTERS 22, 347 (1966), PHYS. REV. LETTERS 9, 300 (1961), CERN CONFERENCE 295, PHYS. REV. 135B, 484 (1964), PHYS. LETTERS 9, 71 (1964), PHYS. LETTERS 15, 74 (1965), THESIS GOSHAW (1966), PHYS. REV. LETTERS 18, 1059 (1966), PHYS. LETTERS 9, 207 (1964), PHYS. LETTERS 12, 65 (1964), PHYS. LETTERS 16, 184 (1965), PHYS. REV. LETTERS 17, 726 (1966), BULL. AM. PHYS. SOC. 11, 76 (1966), UCRL 16744, PHYS. LETTERS 22, 357 (1966), PHYS. REV. LETTERS 18, 1087 (1967), PHYS. LETTERS 14, 338 (1965), PHYS. REV. LETTERS 14, 401 (1965), PHYS. LETTERS 15, 351 (1965), PHYS. LETTERS 19, 612 (1965), PHYS. REV. LETTERS 18, 680 (1967), NUCLEAR INSTRUMENTS AND METHODS 20, 100 (1963), APPLIED OPTICS 2, 1003 (1963), BNL E-41, BNL BC-01-2-C, BNL BC-05-2-G, PHYS. REV. 129, 2285 (1963), PHYS. REV. LETTERS 16, 1063 (1966), PHYS. REV. LETTERS 18, 801 (1967), NUOVO CIMENTO 34, 1644 (1964), NUOVO CIMENTO 33, 309 (1964), PHYS. REV. LETTERS 11, 90 (1963), NUOVO CIMENTO 27, 724 (1953), PHYS. LETTERS 8, 144 (1964), PHYS. REV. LETTERS 17, 1200 (1966), PHYS. REV. LETTERS 13, 149 (1964), DUBNA CONFERENCE 1 480 (1964), PHYS. REV. LETTERS 17, 102 (1966), PHYS. REV. 134, 81062 (1964), PHYS. REV. 134, 81079 (1964), PHYS. REV. 137, 81115 (1965), PHYS. REV. 98, 1416 (1955), PHYS. LETTERS 15, 281 (1965), IMPERIAL COLLEGE, LONDON ICTP/65/18, PHYS. LETTERS 18, 357 (1965), PHYS. REV. LETTERS 14, 806 (1965), PHYS. REV. 139, 81023 (1965), PHYS. LETTERS 8, 134 (1964), PHYS. LETTERS 18, 167 (1965), PHYS. REV. LETTERS 17, 884 (1966), PHYS. REV. LETTERS 13, 555 (1964), PHYS. REV. LETTERS 14, 1043 (1965), PHYS. LETTERS 18, 177 (1965), BULL. AM. PHYS. SOC. 10, 503 (1965), PHYS. REV. LETTERS 15, 329 (1965), PHYS. REV. LETTERS 17, 423 (1966), PHYS. LETTERS 17, 155 (1965), PHYS. REV. LETTERS 17, 716 (1966), NUOVO CIMENTO 36, 1101 (1965), PHYS. REV. 100, 1540 (1955), PHYS. REV. LETTERS 8, 140 (1962), AND NUOVO CIMENTO 33, 906 (1964).

ARTICLE READ BY ODETTE BENARY IN 12/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K⁺ ON PROTON AT 3.535 GEV/C.

THIS EXPERIMENT USES THE B.N.L. 80 IN. (H) BUBBLE CHAMBER.

GENERAL COMMENTS ON THIS ARTICLE

1 CROSS SECTIONS ARE NORMALIZED TO TAU DECAYS.

KEY WORDS = CROSS SECTION MASS SPECTRUM DENSITY MATRIX DALITZ PLOT ANGULAR DISTRIBUTION MODELS
K*(1420) BRANCHING RATIO K*(1320) STRANGE PARTICLES DELTA(1236) N*(1400) K*(890)
N*(1512) N*(1688) ETA MESON(548) OMEGA MESON Q PEAK

COMPOUND KEY WORDS = K*(1420) BRANCHING RATIO K*(1320) BRANCHING RATIO

CROSS SECTION FOR K⁺ PROTON → K⁰ PI⁺ PROTON. (TABLE 4)

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
3.535 ± .025	1388. ± 29.	2361

CROSS SECTION FOR K⁺ PROTON → K⁰ PI⁺ PROTON PI⁰. (TABLE 4)

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
3.535 ± .025	1831. ± 34.	3114

CROSS SECTION FOR K⁺ PROTON → K⁰ PI⁺ PI⁰ NEUTRON. (TABLE 4)

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
3.535 ± .025	506. ± 17.	861

CROSS SECTION FOR K⁺ PROTON → LAMBDA K⁺ K⁺. (TABLE 4)

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
3.535 ± .025	24.7 ± 2.7	84

CROSS SECTION FOR K⁺ PROTON → SIGMA⁰ K⁺ K⁺. (TABLE 4)

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
3.535 ± .025	9.7 ± 1.8	33

CROSS SECTION FOR K⁺ PROTON → LAMBDA K⁺ K⁺ PI⁰. (TABLE 4)

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
3.535 ± .025	73.5 ± 2.1	46

CROSS SECTION FOR K⁺ PROTON → K⁰ BAR K⁺ K⁺ NEUTRON. (TABLE 4)

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
3.535 ± .025	7.6 ± 2.3	13

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \pi^+ \text{ PROTON } M \geq 2\pi^0$. [TABLE 4]

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
3.535 $\pm$.025	238. $\pm$ 33.	157

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \pi^+ \text{ PROTON } \eta \text{ MESON}(548)$. [TABLE 4]

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
3.535 $\pm$.025	49.1 $\pm$ 6.7	61

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \pi^+ \text{ PROTON } \Omega \text{ MESON}$. [TABLE 4]

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
3.535 $\pm$.025	64. $\pm$ 35.	11

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \pi^+ \pi^+ \text{ } M \geq 2\pi^0$. [TABLE 4]

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
3.535 $\pm$.025	166. $\pm$ 56.	282

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \Delta^{++}$. [TABLE 7]
 $\Delta^{++} \rightarrow \text{PROTON } \pi^+ [1]$

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
3.535 $\pm$.025	351. $\pm$ 15.	603

[1] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)^+ \text{ PROTON}$. [TABLE 7]
 $K^*(890)^+ \rightarrow [\text{DETECTED MODE}] K^0 \pi^+ + K^+ \pi^0 [1]$

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS [2]	NO. EVENTS
3.535 $\pm$.025	548. $\pm$ 23.	620

[1] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.

[2] THE FOLLOWING CORRECTION FACTOR WAS USED TO GO FROM DETECTED TO REPORTED QUANTITIES
 $R = 2$ (A CLEBSCH-GORDON FACTOR), WHERE R IS THE RATIO FOR $[K^*(890)^+ \rightarrow K^0 \pi^+] \text{ DIVIDED BY } [K^*(890)^+ \rightarrow K^+ \pi^0]$.CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(1420)^+ \text{ PROTON}$. [TABLE 7]
 $K^*(1420)^+ \rightarrow [\text{DETECTED MODE}] K^0 \pi^+ + K^+ \pi^0 [1]$

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS [2]	NO. EVENTS
3.535 $\pm$.025	120. $\pm$ 13.	136

[1] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.

[2] THE FOLLOWING CORRECTION FACTOR WAS USED TO GO FROM DETECTED TO REPORTED QUANTITIES
 $R = 2$ (A CLEBSCH-GORDON FACTOR), WHERE R IS THE RATIO FOR $[K^*(1420)^+ \rightarrow K^0 \pi^+] \text{ DIVIDED BY } [K^*(1420)^+ \rightarrow K^+ \pi^0]$.CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \text{ PROTON } \pi^+$. [TABLE 7]
CROSS SECTION IS ONLY FOR THE NON-RESONANT PRODUCTION OF THESE PARTICLES.

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
3.535 $\pm$.025	588. $\pm$ 24.	1002

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow \Delta^{++} K^0 \pi^0$. [TABLE 11]
 $\Delta^{++} \rightarrow \text{PROTON } \pi^+ [1]$

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
3.535 $\pm$.025	407. $\pm$ 18.	691

[1] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow \Delta^{++} K^0 \pi^+ [1]$. [TABLE 11]
 $\Delta^{++} \rightarrow [\text{DETECTED MODE}] \text{ PROTON } \pi^0 + \text{ NEUTRON } \pi^+ [1]$

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS [2]	NO. EVENTS
3.535 $\pm$.025	195. $\pm$ 16.	220

[1] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.

[2] THE FOLLOWING CORRECTION FACTOR WAS USED TO GO FROM DETECTED TO REPORTED QUANTITIES
 $R = 2$ (A CLEBSCH-GORDON FACTOR), WHERE R IS THE RATE FOR $[\Delta^{++} \rightarrow \text{PROTON } \pi^0]$ DIVIDED BY $[\Delta^{++} \rightarrow \text{NEUTRON } \pi^+]$.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)^+ \text{ PROTON } \pi^0$. [TABLE 11]
 $K^*(890)^+ \rightarrow [\text{DETECTED MODE}] K^0 \pi^+ + K^+ \pi^0 [1]$

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS [2]	NO. EVENTS
3.535 $\pm$.025	457. $\pm$ 19.	516

[1] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.

[2] THE FOLLOWING CORRECTION FACTOR WAS USED TO GO FROM DETECTED TO REPORTED QUANTITIES
 $R = 2$ (A CLEBSCH-GORDON FACTOR), WHERE R IS THE RATE FOR $[K^*(890)^+ \rightarrow K^0 \pi^+]$ DIVIDED BY $[K^*(890)^+ \rightarrow K^+ \pi^0]$.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)^0 \text{ PROTON } \pi^+ [1]$. [TABLE 11]
 $K^*(890)^0 \rightarrow [\text{DETECTED MODE}] K^0 \pi^0 + K^+ \pi^- [1]$

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS [2]	NO. EVENTS
3.535 $\pm$.025	858. $\pm$ 38.	486

[1] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.

[2] THE FOLLOWING CORRECTION FACTOR WAS USED TO GO FROM DETECTED TO REPORTED QUANTITIES
 $R = 2$ (A CLEBSCH-GORDON FACTOR), WHERE R IS THE RATE FOR $[K^*(890)^0 \rightarrow K^+ \pi^-]$ DIVIDED BY $[K^*(890)^0 \rightarrow K^0 \pi^0]$.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)^0 \Delta^{++}$. [TABLE 11]
 $\Delta^{++} \rightarrow \text{PROTON } \pi^+ [1]$
 $K^*(890)^0 \rightarrow [\text{DETECTED MODE}] K^0 \pi^0 + K^+ \pi^- [2]$

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS [3]	NO. EVENTS
3.535 $\pm$.025	560. $\pm$ 33.	316

[1] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.

[2] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.

[3] THE FOLLOWING CORRECTION FACTOR WAS USED TO GO FROM DETECTED TO REPORTED QUANTITIES
 $R = 2$ (A CLEBSCH-GORDON FACTOR), WHERE R IS THE RATE FOR $[K^*(890)^0 \rightarrow K^+ \pi^-]$ DIVIDED BY $[K^*(890)^0 \rightarrow K^0 \pi^0]$.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)^+ \Delta^{++}$. [TABLE 11]
 $\Delta^{++} \rightarrow [\text{DETECTED MODE}] \text{ PROTON } \pi^0 + \text{ NEUTRON } \pi^+ [1]$
 $K^*(890)^+ \rightarrow [\text{DETECTED MODE}] K^0 \pi^+ + K^+ \pi^0 [2]$

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS [3]	NO. EVENTS
3.535 $\pm$.025	261. $\pm$ 21.	195

[1] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.

[2] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.

[3] THE FOLLOWING CORRECTION FACTORS WERE USED TO GO FROM DETECTED TO REPORTED QUANTITIES
 1. $R = 2$ (A CLEBSCH-GORDON FACTOR), WHERE R IS THE RATE FOR $[\Delta^{++} \rightarrow \text{PROTON } \pi^0]$ DIVIDED BY $[\Delta^{++} \rightarrow \text{NEUTRON } \pi^+]$
 2. $R = 2$ (A CLEBSCH-GORDON FACTOR), WHERE R IS THE RATE FOR $[K^*(890)^+ \rightarrow K^0 \pi^+]$ DIVIDED BY $[K^*(890)^+ \rightarrow K^+ \pi^0]$.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \pi^+ \pi^0 \text{ PROTON}$. [TABLE 11]
 CROSS SECTION IS ONLY FOR THE NON-RESONANT PRODUCTION OF THESE PARTICLES.

LABORATORY BEAM MOMENTUM GEV/C	MICRO-BARNS	NO. EVENTS
3.535 $\pm$.025	825. $\pm$ 82.	1400

[2] THE FOLLOWING CORRECTION FACTOR WAS USED TO GO FROM DETECTED TO REPERTED QUANTITIES

$$R = 2 \cdot \frac{1}{\lambda} \text{ (A CLEBSCH-GORDON FACTOR)}, \text{ WHERE } R \text{ IS THE RATE FOR } [K^*(890)] \cdot \lambda \cdot K \cdot P(+) \text{ DIVIDED BY } [K^*(890)] + K + P(+) .$$

CROSS SECTION FOR K+ PROTON - DELTA+ KO PI+. (TABLE 18)
DELTA+ - (DETECTED MODE) NEUTRON PI+ + PROTON PI0 (1)

[2] THE FOLLOWING CORRECTION FACTOR WAS USED TO GO FROM DETECTED TO REPORTED QUANTITIES
 $R = 2 \cdot (1 - \text{GLEBSCH-GORDON FACTOR})$, WHERE R IS THE RATE FOR $(\Delta^+ \rightarrow \pi^- \text{ PROTON } \pi^0)$ DIVIDED BY $(\Delta^+ \rightarrow \pi^- \text{ NEUTRON } \pi^0)$.
 $R \approx 1$.

CROSS SECTION FOR K+ PROTON = N*(1400)+ K0 PI+. [TABLE 18]
N*(1400)+ = [DETECTED MODE] NEUTRON PI+ + PROTON PI0 [1]

[1] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
[2] THE FOLLOWING CORRECTION FACTOR WAS USED TO GO FROM DETECTED TO REPORTED QUANTITIES:
R = 2 * (A CLEBSCH-GORDON FACTOR), WHERE R IS THE RATE FOR $1N \rightarrow (1400)^+$ - PROTON PION DIVIDED BY $[N \rightarrow (1400)^+$ -
NEUTRON PION]

CROSS SECTION FOR K+ PROTON = N*(1512)+ KO PI+. [TABLE 18]
N*(1512)+ = [DETECTED MODE] NEUTRON PI+ + PROTON PI0 (1)

(1) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND WITH FORMS ONLY ABOVE FITTED BACKGROUND.
 (2) THE FOLLOWING CORRECTION FOR WAS USED TO GO FROM DETECTED TO REPORTED QUANTITIES

$$N = A \text{ CLEO5000 DARGON FACTOR} \times \frac{1}{1 + \frac{1}{2} \left(\frac{1}{\text{PRTON P10}} \right)^2}$$
 WHERE B IS THE DATE FOR INY(1512) + NEUTRON P10.

CROSS SECTION FOR $n + \text{PROTON} \rightarrow n + \text{PI}^+ + \text{PI}^0$ (TABLE 18)
 $N(1688) + \text{DETECTED MODE} \text{ NEUTRON PI}^+ + \text{PROTON PI}^0$ (1)

[1] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
[2] THE FOLLOWING CORRECTION FACTOR WAS USED TO GO FROM DETECTED TO REPORTED QUANTITIES
 $R = 2 \left(\frac{1}{1 + \text{CLEBSCH-GORDON FACTOR}} \right)$, WHERE R IS THE RATE FOR $IN(11500)^+$ - PROTON P10 DIVIDED BY $IN(1600)^+$ - NEUTRON P10.

CROSS SECTION FOR K+ PROTON - DELTA+ K*(890)+. [TABLE 18]

[1] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 [2] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 [3] THE FOLLOWING CORRECTION FACTORS WERE USED TO GO FROM DETECTED TO REPORTED QUANTITIES
 1. $R = 2$ (A CLEBSCH-GORDON FACTOR), WHERE R IS THE RATE FOR $(\Delta E)^+ \rightarrow \text{PROTON } \pi^0$ DIVIDED BY $(\Delta E)^+ \rightarrow \text{NEUTRON } \pi^0$
 2. $R = 2$ (A CLEBSCH-GORDON FACTOR), WHERE R IS THE RATE FOR $(K^+ \pi^0)^+ \rightarrow K^0 \pi^+$ DIVIDED BY $(K^+ \pi^0)^+ \rightarrow K^+ \pi^0$

CROSS SECTION FOR $K^+ \text{ PROTON} = N^*(1400)^+ K^*(890)^+ \dots$ [TABLE 18]
 $N^*(1400)^+ = \{\text{DETECTED MODE}\} \text{ NEUTRON } \pi^+ + \text{ PROTON } \pi^0 \{1\}$
 $K^*(890)^+ = \{\text{DETECTED MODE}\} K^0 \pi^+ + K^+ \pi^0 \{2\}$

LABORATORY
 BEAM MOMENTUM
 GEV/C
 3.535 $\pm$.025
 MICRO-BARNS {3}
 26. $\pm$ 7.
 NO. EVENTS
 20

- (1) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 (2) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 (3) THE FOLLOWING CORRECTION FACTORS WERE USED TO GO FROM DETECTED TO REPORTED QUANTITIES
 1. $R = 2$ (A CLEBSCH-GORDON FACTOR), WHERE R IS THE RATE FOR $[N^*(1400)^+ + \text{PROTON } \pi^0] \text{ DIVIDED BY } [N^*(1400)^+ + \text{NEUTRON } \pi^+]$
 2. $R = 2$ (A CLEBSCH-GORDON FACTOR), WHERE R IS THE RATE FOR $[K^*(890)^+ + K^0 \pi^+] \text{ DIVIDED BY } [K^*(890)^+ + K^+ \pi^0]$.

CROSS SECTION FOR $K^+ \text{ PROTON} = N^*(1512)^+ K^*(890)^+ \dots$ [TABLE 18]
 $N^*(1512)^+ = \{\text{DETECTED MODE}\} \text{ NEUTRON } \pi^+ + \text{ PROTON } \pi^0 \{1\}$
 $K^*(890)^+ = \{\text{DETECTED MODE}\} K^0 \pi^+ + K^+ \pi^0 \{2\}$

LABORATORY
 BEAM MOMENTUM
 GEV/C
 3.535 $\pm$.025
 MICRO-BARNS {3}
 47. $\pm$ 8.
 NO. EVENTS
 35

- (1) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 (2) USED SIMPLE MASS CUT.
 (3) THE FOLLOWING CORRECTION FACTORS WERE USED TO GO FROM DETECTED TO REPORTED QUANTITIES
 1. $R = 2$ (A CLEBSCH-GORDON FACTOR), WHERE R IS THE RATE FOR $[N^*(1512)^+ + \text{PROTON } \pi^0] \text{ DIVIDED BY } [N^*(1512)^+ + \text{NEUTRON } \pi^+]$
 2. $R = 2$ (A CLEBSCH-GORDON FACTOR), WHERE R IS THE RATE FOR $[K^*(890)^+ + K^0 \pi^+] \text{ DIVIDED BY } [K^*(890)^+ + K^+ \pi^0]$.

CROSS SECTION FOR $K^+ \text{ PROTON} = N^*(1688)^+ K^*(890)^+ \dots$ [TABLE 18]
 $N^*(1688)^+ = \{\text{DETECTED MODE}\} \text{ NEUTRON } \pi^+ + \text{ PROTON } \pi^0 \{1\}$
 $K^*(890)^+ = \{\text{DETECTED MODE}\} K^0 \pi^+ + K^+ \pi^0 \{2\}$

LABORATORY
 BEAM MOMENTUM
 GEV/C
 3.535 $\pm$.025
 MICRO-BARNS {3}
 57. $\pm$ 19.
 NO. EVENTS
 43

- (1) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 (2) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 (3) THE FOLLOWING CORRECTION FACTORS WERE USED TO GO FROM DETECTED TO REPORTED QUANTITIES
 1. $R = 2$ (A CLEBSCH-GORDON FACTOR), WHERE R IS THE RATE FOR $[N^*(1688)^+ + \text{PROTON } \pi^0] \text{ DIVIDED BY } [N^*(1688)^+ + \text{NEUTRON } \pi^+]$
 2. $R = 2$ (A CLEBSCH-GORDON FACTOR), WHERE R IS THE RATE FOR $[K^*(890)^+ + K^0 \pi^+] \text{ DIVIDED BY } [K^*(890)^+ + K^+ \pi^0]$.

CROSS SECTION FOR $K^+ \text{ PROTON} = K^0 \pi^+ \pi^+ \text{ NEUTRON}$. [TABLE 18]
 CROSS SECTION IS ONLY FOR THE NON-RESONANT PRODUCTION OF THESE PARTICLES.

LABORATORY
 BEAM MOMENTUM
 GEV/C
 3.535 $\pm$.025
 MICRO-BARNS
 202. $\pm$ 75.
 NO. EVENTS
 348

49 NEW STRUCTURE IN THE $K^+ \text{ P}$ AND $K^+ \text{ D}$ TOTAL CROSS SECTIONS BETWEEN 0.9 AND 2.4 GEV/C. [PHYS. REV. LETTERS 17, 102 (1966)]

R.L.COOL, G.GIACOMELLI, T.F.KYCIA, B.A.LEONTIC, K.K.LI, A.LUNDBY, J.TEIGER (BROOKHAVEN NAT. LAB., UPTON, L.I., N. Y., USA)

ADDITIONAL FOOTNOTE REFERENCES

PHYS. REV. LETTERS 15, 1228 (1966); PHYS. REV. LETTERS 7, 188 (1961); PHYS. REV. LETTERS 7, 182 (1961); PHYS. REV. LETTERS 9, 135 (1962); PHYS. REV. 100, 242 (1955); UCRL 11446 (1964); PHYS. LETTERS 6, 329 (1963); PHYS. REV. LETTERS 11, 93 (1963); PHYS. REV. 139, 8220 (1964); NUOVO CIMENTO 36, 1101 (1955); PHYS. REV. 129, 2743 (1963); PHYS. REV. LETTERS 7, 378 (1961); PHYS. REV. 134, 8111 (1964); PHYS. REV. 138, 8947 (1965); AND CERN TH-628 (1965).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 4/68, AND VERIFIED BY LEROY PRICE.

BEAM NO. 1 IS K^+ ON PROTON FROM .9 TO 2.4 GEV/C.
 NO. 2 IS K^+ ON DEUTERON FROM .9 TO 2.4 GEV/C.

THIS EXPERIMENT USES COUNTERS.

KEY WORDS = CROSS SECTION

K+ PROTON TOTAL CROSS SECTION.

ORIGINAL DATA			CORRECTED VALUES (1)		
LABORATORY BEAM MOMENTUM GEV/C	PER CENT	MILLI-BARNS	LABORATORY BEAM MOMENTUM GEV/C	PER CENT	MILLI-BARNS
.90 +- 1		14.60 +- 2.1	.891 +- 1		14.39 +- .38
.95 1		15.63 1.6	.942 1		15.57 .30
1.00 1		15.97 1.5	.992 1		15.97 .27
1.05 1		17.35 1.3	1.043 1		17.39 .25
1.10 1		17.10 1.3	1.094 1		17.12 .25
1.15 1		17.98 .9	1.144 1		18.09 .15
1.20 1		18.34 .5	1.194 1		18.47 .12
1.25 1		18.42 .5	1.245 1		18.54 .12
1.30 1		18.42 .5	1.295 1		18.61 .11
1.35 1		18.30 .5	1.345 1		18.44 .10
1.40 1		18.14 .5	1.395 1		18.27 .10
1.45 1		17.90 .5	1.445 1		18.04 .10
1.50 1		17.74 .5	1.495 1		17.93 .09
1.60 1		17.54 .5	1.596 1		17.75 .09
1.65 1		17.64 .5	1.646 1		17.86 .06
1.70 1		17.64 .5	1.696 1		17.85 .06
1.75 1		17.64 .5	1.746 1		17.80 .06
1.80 1		17.54 .5	1.796 1		17.80 .06
1.90 1		17.54 .5	1.896 1		17.81 .06
2.00 1		17.34 .5	1.996 1		17.60 .08
2.10 1		17.25 .5	2.096 1		17.51 .08
2.20 1		17.25 .5	2.196 1		17.54 .08
2.30 1		17.25 .5	2.296 1		17.55 .08
2.40 1		17.25 .5			

(1) PRIVATE COMMUNICATION FROM T.KYCIA, APRIL, 1969.

K+ DEUTERON TOTAL CROSS SECTION.

ORIGINAL DATA			CORRECTED VALUES (1)		
LABORATORY BEAM MOMENTUM GEV/C	PER CENT	MILLI-BARNS	LABORATORY BEAM MOMENTUM GEV/C	PER CENT	MILLI-BARNS
.90 +- 1		30.60 +- 1.10	.888 +- 1		30.10 +- .36
.95 1		32.51 1.22	.939 1		31.98 .34
1.00 1		33.58 .60	.989 1		33.20 .24
1.05 1		35.06 .57	1.040 1		34.77 .24
1.10 1		36.24 .39	1.091 1		36.04 .24
1.15 1		37.41 .32	1.141 1		37.22 .15
1.20 1		37.90 .25	1.191 1		37.69 .12
1.25 1		37.80 .25	1.242 1		37.57 .11
1.30 1		37.61 .25	1.292 1		37.43 .10
1.35 1		37.16 .25	1.342 1		36.99 .12
1.40 1		36.86 .25	1.392 1		36.78 .12
1.45 1		36.52 .25	1.442 1		36.43 .10
1.50 1		36.33 .25	1.492 1		36.27 .10
1.60 1		35.94 .25	1.593 1		35.89 .08
1.65 1		35.84 .25	1.643 1		35.78 .08
1.70 1		35.74 .25	1.693 1		35.76 .09
1.75 1		35.84 .25	1.743 1		35.89 .06
1.80 1		35.84 .25	1.793 1		35.70 .06
1.90 1		35.35 .25	1.893 1		35.33 .06
2.00 1		35.25 .25	1.993 1		35.41 .08
2.10 1		35.25 .25	2.093 1		35.47 .08
2.25 1		35.00 .25	2.193 1		35.22 .10
2.45 1		35.06 .25	2.393 1		35.45 .08

(1) PRIVATE COMMUNICATION FROM T.KYCIA, APRIL, 1969.

I = 0 K NUCLEON TOTAL CROSS SECTION.

(PRIVATE COMMUNICATION FROM T.KYCIA APRIL 1969)

LABORATORY BEAM MOMENTUM GEV/C	PER CENT	MILLI-BARNS
.888 +- 1		19.31 +- 1.35
.939 1		19.62 1.13
.989 1		21.09 .94
1.040 1		20.42 .89
1.091 1		24.09 .89
1.141 1		24.40 .54
1.191 1		23.27 .43
1.242 1		21.97 .42
1.292 1		20.74 .39
1.342 1		20.20 .38
1.392 1		20.17 .38
1.442 1		20.09 .36
1.492 1		19.98 .34
1.593 1		19.83 .31
1.643 1		19.48 .24
1.693 1		19.53 .22
1.743 1		19.96 .22
1.793 1		19.65 .22
1.893 1		19.18 .22
1.993 1		19.50 .29
2.093 1		19.92 .29
2.193 1		19.26 .31
2.393 1		19.71 .29

K+ NEUTRON TOTAL CROSS SECTION.

(PRIVATE COMMUNICATION FROM T.KYCIA APRIL 1969)

LABORATORY BEAM MOMENTUM GEV/C	PER CENT	MILLI-BARNS
.888 +- 1		16.85 +- .52
.939 1		17.60 .45
.989 1		18.53 .36
1.040 1		18.91 .35
1.091 1		20.61 .35
1.141 1		21.25 .21
1.191 1		20.87 .17
1.242 1		20.26 .16
1.292 1		19.68 .15
1.342 1		19.32 .16
1.392 1		19.22 .16
1.442 1		19.07 .14
1.492 1		18.95 .13
1.593 1		18.79 .12
1.643 1		18.67 .10
1.693 1		18.69 .08
1.743 1		18.88 .08
1.793 1		18.73 .08
1.893 1		18.50 .08
1.993 1		18.55 .11
2.093 1		18.67 .11
2.193 1		18.40 .13
2.393 1		18.63 .11

50 STUDY OF THE REACTIONS $K^+P \rightarrow K^+N^*$ AT 3.5 AND 5 GEV/C. [NUOVO CIMENTO 49A, 9 (1967)]

R. GEORGE, Y. GOLDSCHMIDT-CLERMONT, V. P. HENRI, B. JONGEJANS, M. KRAMMER, F. MULLER, J. M. PERREAU (EUROPEAN ORG. FOR NUC. RES., GENEVA, SWITZERLAND)
 W. DEBAERE, J. DEBAISIEUX, P. DUFOUR, F. GRARD, J. HEUGHEBAERT, L. PAPE, P. PEETERS, F. VERBEURE, R. WINDHOLDERS (INST. INTERUNIV. DES SCI. NUC., BRUXELLES, BELGIUM)

ABSTRACT FROM A STUDY OF THE FOUR-BODY FINAL STATE $K^+P \rightarrow K^+N^*PI$ AT 3.5 AND 5 GEV/C IT IS FOUND THAT THE QUASI-TWO-BODY REACTION $K^+P \rightarrow K^*(892)N^*(1236)$ IS PRODUCED IN LARGE AMOUNT. THIS REACTION SHOWS THE SAME GENERAL BEHAVIOUR AS OBTAINED AT 3 GEV/C SHARPLY PEAKED PRODUCTION ANGULAR DISTRIBUTIONS SUGGESTING PERIPHERALISM AS THE MECHANISM OF PRODUCTION. THE EXPERIMENTAL ANGULAR DISTRIBUTIONS ARE COMPARED WITH THE PREDICTIONS OF THREE MODELS A) ONE-PION EXCHANGE WITH ARBITRARY FORM FACTORS, B) ABSORPTION WITH PION EXCHANGE, C) SIMPLE REGGE-POLE EXCHANGE. THE RESULTS SHOW THAT THE ABSORPTION MODEL WITH PI EXCHANGE PREDICTS THE SHAPE FOR THE PRODUCTION ANGULAR DISTRIBUTION REASONABLY WELL BUT LESS WELL THE ENERGY DEPENDENCE OF THE PRODUCTION CROSS-SECTION. IT PREDICTS RATHER WELL THE GENERAL TREND OF THE DENSITY MATRIX ELEMENTS CHARACTERIZING THE DECAY DISTRIBUTIONS. IT ALSO SUCCEEDS IN ACCOUNTING FOR THE EXPERIMENTAL ANGULAR CORRELATIONS BETWEEN THE K^+ AND N^* DECAY. THE SIMPLE REGGE-POLE EXCHANGE GIVES ACCEPTABLE RESULTS FOR BOTH THE SHAPE AND ENERGY VARIATION OF THE PRODUCTION ANGULAR DISTRIBUTIONS. THE OBSERVED $K^*(1420)N^*(1236)$ REACTION AT 5 GEV/C GAVE PRODUCTION AND DECAY DISTRIBUTIONS IN QUALITATIVE AGREEMENT WITH A $2^+K^*(1400)$ PRODUCED IN A PI EXCHANGE PROCESS.

CROSS REFERENCES

SEE ALSO PHYS. LETTERS 26B, 30 (1967).
 PART OF THIS ARTICLE SUPERSEDED BY PHYS. LETTERS 27B, 46B (1968).
 CONTINUATION OF PREVIOUS EXPERIMENT IN NUOVO CIMENTO 39, 417 (1965).

ADDITIONAL FOOTNOTE REFERENCES

DUBNA CONFERENCE 1 718 (1964), CERN 63-12, CERN TC/02 63-3, PHYS. LETTERS 6, 62 (1963), NUOVO CIMENTO 33, 906 (1954), NUOVO CIMENTO 31, 363 (1964), PHYS. REV. 139, 842B (1965), NUOVO CIMENTO 45A, 885 (1966), NUOVO CIMENTO 42, 323 (1966), PHYS. LETTERS 18, 76 (1965), NUOVO CIMENTO 38, 518 (1965), BULL. AM. PHYS. SOC. 11, 77 (1966), AND UCRL 16332.

ARTICLE READ BY ODETTE BENARY IN 12/68, AND VERIFIED BY ROGER BLAND.

BEAM IS K^+ ON PROTON FROM 3.46 TO 4.97 GEV/C.

THIS EXPERIMENT USES THE SACLAY 81 CM (H) BUBBLE CHAMBER. A TOTAL OF 110000 PICTURES ARE REPORTED ON.

GENERAL COMMENTS ON THIS ARTICLE

1 CROSS SECTIONS ARE NORMALIZED TO TAU DECAYS.

KEY WORDS = CROSS SECTION DIFFERENTIAL CROSS SECTION MASS SPECTRUM DELTA(123B) $K^*(890)$ $K^*(1420)$
 MODELS DENSITY MATRIX ANGULAR DISTRIBUTION

CROSS SECTION FOR K^+ PROTON $\rightarrow$ K^0 PROTON PI^+PI^0 . [TABLE 1]

(THIS DATA SHOULD NOT BE USED - MORE RECENT VALUES MAY BE FOUND IN BASSOMPIERRE, ET. AL., PHYS. LETTERS 27B, 46B (1968))

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS
3.460 +- .015	2.45 +- .30	558
4.970 .015	2.30 .30	569

CROSS SECTION FOR K^+ PROTON $\rightarrow$ K^+ PROTON PI^+PI^- . [TABLE 1]

(THIS DATA SHOULD NOT BE USED - MORE RECENT VALUES MAY BE FOUND IN BASSOMPIERRE, ET. AL., PHYS. LETTERS 27B, 46B (1968))

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS
3.460 +- .015	2.10 +- .20	1482
4.970 .015	1.80 .15	1222

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \text{ NEUTRON } \pi^+ \pi^+$ (TABLE 1)

(THIS DATA SHOULD NOT BE USED - MORE RECENT VALUES MAY BE FOUND IN BASSOMPIERRE, ET. AL., PHYS. LETTERS 27B, 468 (1968))

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS
3.460 $\pm$.015	.55 $\pm$.10	121
4.970 $\pm$.015	.60 $\pm$.10	145

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890) \Delta^{++}$ (TABLE 2)
 $\Delta^{++} \rightarrow \text{PROTON } \pi^+ \pi^+$ (1)
 $K^*(890) \rightarrow K^0 \pi^0$ (2)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
3.460 $\pm$.015	.56 $\pm$.10

- (1) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH (MASS = 1.234 GEV; WIDTH = .140 GEV) AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND. NON-INTERFERING AMPLITUDES ASSUMED.
 (2) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH (MASS = .891 GEV; WIDTH = .050 GEV) AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND. NON-INTERFERING AMPLITUDES ASSUMED.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890) \Delta^{++}$ (TABLE 2)
 $\Delta^{++} \rightarrow \text{PROTON } \pi^+ \pi^+$ (1)
 $K^*(890) \rightarrow K^+ \pi^-$ (2)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
4.970 $\pm$.015	.72 $\pm$.06

- (1) FITTED FOR MASS AND/OR WIDTH (MASS = 1.244 GEV; WIDTH = .125 GEV), AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND. NON-INTERFERING AMPLITUDES ASSUMED.
 (2) FITTED FOR MASS AND/OR WIDTH (MASS = .892 GEV; WIDTH = .050 GEV), AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND. NON-INTERFERING AMPLITUDES ASSUMED.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890) \Delta^{++}$ (TABLE 2)
 $\Delta^{++} \rightarrow \text{PROTON } \pi^+ \pi^+$ (1)
 $K^*(890) \rightarrow K^+ \pi^-$ (2)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
3.460 $\pm$.015	.92 $\pm$.10

- (1) FITTED FOR MASS AND/OR WIDTH (MASS = 1.228 GEV; WIDTH = .142 GEV), AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND. NON-INTERFERING AMPLITUDES ASSUMED.
 (2) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH (MASS = .890 GEV; WIDTH = .050 GEV) AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND. NON-INTERFERING AMPLITUDES ASSUMED.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890) \Delta^{++}$ (TABLE 2)
 $\Delta^{++} \rightarrow \text{PROTON } \pi^0 \pi^0$ (1)
 $K^*(890) \rightarrow K^0 \pi^+$ (2)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
3.460 $\pm$.015	.15 $\pm$.03

- (1) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH (MASS = 1.234 GEV; WIDTH = .140 GEV) AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND. NON-INTERFERING AMPLITUDES ASSUMED.
 (2) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH (MASS = .891 GEV; WIDTH = .050 GEV) AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND. NON-INTERFERING AMPLITUDES ASSUMED.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(1420) \Delta^{++}$ (TABLE 2)
 $\Delta^{++} \rightarrow \text{PROTON } \pi^+ \pi^+$ (1)
 $K^*(1420) \rightarrow K^+ \pi^-$ (2)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
4.970 $\pm$.015	.11 $\pm$.02

- (1) FITTED FOR MASS AND/OR WIDTH (MASS = 1.244 GEV; WIDTH = .125 GEV), AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND. NON-INTERFERING AMPLITUDES ASSUMED.
 (2) FITTED FOR MASS AND/OR WIDTH (MASS = 1.430 GEV; WIDTH = .110 GEV), AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND. NON-INTERFERING AMPLITUDES ASSUMED.

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890) \Delta^{++}$ (FIGURE 5)
 $\Delta^{++} \rightarrow \text{PROTON } \pi^+ \text{ (1)}$
 $K^*(890) \rightarrow K^+ \pi^- \text{ (2)}$

LABORATORY BEAM MOMENTUM = $3.460 \pm .015$ GEV/C.

.....
 . THIS DATA WAS READ FROM A GRAPH .

COS(THETA)		D-SIGMA/D-OMEGA UB/SR (3)		NO. EVENTS
MAX	MIN			
1.000	.975	1.88	±.24	64
.975	.950	1.23	±.19	42
.950	.925	1.03	±.17	35
.925	.900	.59	±.13	20
.900	.875	.24	±.08	8
.875	.850	.18	±.07	6
.850	.825	.18	±.07	6
.825	.800	.06	±.04	2
.800	.775	.15	±.07	5
.775	.750	.00	±.03	0
.750	.725	.12	±.06	4
.725	.700	.03	±.03	1
.700	.675	.03	±.03	1
.675	.650	.06	±.04	2
.650	.625	.09	±.05	3
.625	.600	.09	±.05	3
.600	.575	.03	±.03	1
.575	.550	.03	±.03	1

THETA IS THE ANGLE THAT THE $K^*(890)$ MAKES WITH THE BEAM IN THE GRAND C.M.

- (1) COUNTED ALL EVENTS IN MASS BAND (MASS CUT FROM 1.160 TO 1.290 GEV).
 (2) COUNTED ALL EVENTS IN MASS BAND (MASS CUT FROM .860 TO .940 GEV).
 (3) COUNTS WERE MULTIPLIED BY .0294 TO GET THESE. ERRORS ARE TAKEN AS PROPORTIONAL TO THE SQUARE-ROOT OF THE COUNTS.

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890) \Delta^{++}$ (FIGURE 5)
 $\Delta^{++} \rightarrow \text{PROTON } \pi^+ \text{ (1)}$
 $K^*(890) \rightarrow K^+ \pi^- \text{ (2)}$

LABORATORY BEAM MOMENTUM = $4.970 \pm .015$ GEV/C.

.....
 . THIS DATA WAS READ FROM A GRAPH .

COS(THETA)		D-SIGMA/D-OMEGA UB/SR (3)		NO. EVENTS
MAX	MIN			
1.000	.975	2.24	±.25	83
.975	.950	.89	±.16	33
.950	.925	.38	±.10	14
.925	.900	.27	±.09	10
.900	.875	.38	±.10	14
.875	.850	.11	±.05	4
.850	.825	.05	±.04	2
.825	.800	.05	±.04	2
.800	.775	.03	±.03	1
.775	.750	.03	±.03	1
.750	.725	.03	±.03	1
.725	.700	.05	±.04	2
.700	.675	.03	±.03	1
.675	.650	.03	±.03	1
.600	.575	.05	±.04	2

THETA IS THE ANGLE THAT THE $K^*(890)$ MAKES WITH THE BEAM IN THE GRAND C.M.

- (1) COUNTED ALL EVENTS IN MASS BAND (MASS CUT FROM 1.160 TO 1.290 GEV).
 (2) COUNTED ALL EVENTS IN MASS BAND (MASS CUT FROM .860 TO .940 GEV).
 (3) COUNTS WERE MULTIPLIED BY .027 TO GET THESE. ERRORS ARE TAKEN AS PROPORTIONAL TO THE SQUARE-ROOT OF THE COUNTS.

51 K⁺P ELASTIC SCATTERING AT 3.5 AND 5.0 GEV/C. (NUOVO CIMENTO 45A, 685 (1966))

W. DEBAERE, J. DEBAISIEUX, P. DUFOUR, F. GRARD, J. HEUGHEBAERT, L. PAPE, P. PEETERS, F. VERBEURE, R. WINDMOLDERS (INST. INTERUNIV. DES SCI. NUC., BRUXELLES, BELGIUM)
 R. GEORGE, V. GOLDSCHMIDT-CLERMONT, V. P. MENRI, B. JONGEJANS, D. W. G. LEITH, A. MOISSEEV, F. MULLER, J. M. PERREAU, V. YARBA (EUROPEAN ORG. FOR NUC. RES., GENEVA, SWITZERLAND)

ABSTRACT THE ELASTIC SCATTERING OF K^+ MESONS ON PROTONS IS STUDIED AT 3.5 AND 5 GEV/C. THE TOTAL ELASTIC CROSS SECTIONS ARE FOUND TO BE (4.36 ± 0.36) MB AND (3.82 ± 0.41) MB RESPECTIVELY. THE DIFFERENTIAL ELASTIC CROSS-SECTIONS, WHICH EXHIBIT CHARACTERISTIC DIFFRACTION PEAKS, ARE FITTED BY $D-SIGMA/D-T = (D-SIGMA/D-TAT T=0) \exp(-\alpha T^2)$, GIVING $\alpha = (3.85 \pm 0.21)$ AND (4.70 ± 0.21) (GEV/C)² FOR THE TWO MOMENTA RESPECTIVELY, WITH $T/T_0 < 0.65$ (GEV/C) SQUARED. THE RESULTS ARE COMPARED TO THOSE AT NEIGHBORING ENERGIES, GIVING SOME SUPPORT TO THE PRESENCE OF A REAL PART OF THE FORWARD SCATTERING AMPLITUDE. THE DIFFRACTION PEAK SHOWS DEFINITE SHRINKING WITH INCREASING MOMENTA. THE DATA ARE EXAMINED IN THE LIGHT OF MODELS FOR HIGH-ENERGY SCATTERING.

CROSS REFERENCES
 SEE ALSO NUOVO CIMENTO 43A, 142 (1966).
 SOME OF THIS DATA RENORMALIZED IN UCRL 18758 (1968).

ADDITIONAL FOOTNOTE REFERENCES
 NUOVO CIMENTO 43A, 142 (1966), NUCLEAR INSTRUMENTS AND METHODS 34, 242 (1965), REV. MOD. PHYS. 37, 633 (1964), PHYS. REV. LETTERS 7, 188 (1961), PHYS. REV. LETTERS 7, 182 (1961), PHYS. REV. 129, 2743 (1963), PHYS. REV. 129, 2285 (1963), SIENNA CONFERENCE 1 634 (1963), PHYS. REV. 138, 8913 (1965), UCRL 11446, PHYS. REV. 139, 8141 (1955), PHYS. REV. LETTERS 11, 607 (1963), CERN 45-22, AND PHYS. REV. 139, D1936 (1963).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 1/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K^+ ON PROTON FROM 3.46 TO 4.97 GEV/C.

THIS EXPERIMENT USES THE SACLAY 81 CM (H) BUBBLE CHAMBER. A TOTAL OF 120000 PICTURES ARE REPORTED ON.

GENERAL COMMENTS ON THIS ARTICLE
 1 K^+ LIFETIME TAKEN AS 1.229 ± 0.008 ($\times 10^{-8}$ SEC)
 2 K^+ TO τ BRANCHING RATIO TAKEN AS 0.055.

KEY WORDS = CROSS SECTION ANGULAR DISTRIBUTION MODELS

K+ PROTON ELASTIC CROSS SECTION.

LABORATORY BEAM MOMENTUM GEV/C	PER CENT	MILLI-BARNS	NO. EVENTS
3.46 +- .5		4.36 +- .36	2949
4.97 .5		3.82 .41	1436

ELASTIC DIFFERENTIAL CROSS SECTION FOR K+ PROTON.

LABORATORY BEAM MOMENTUM = 3.46 GEV/C +- .5 (PER CENT).

COS(THETA)	D-SIGMA/D-T MB/(GEV/C)**2	NO. EVENTS
MIN MAX		
-1.00 .25	.007 +- .002	17
.25 .60	.056 .010	37
.60 .65	.350 .070	33
.65 .70	.460 .080	43
.70 .75	1.120 .140	106
.75 .80	1.580 .180	147
.80 .82	2.530 .330	94
.82 .84	3.370 .410	125
.84 .86	2.990 .370	111
.86 .88	4.250 .490	158
.88 .90	5.380 .550	200
.90 .92	5.520 .590	205
.92 .94	7.620 .770	283
.94 .96	9.590 .940	356
.96 .98	12.500 1.170	464
.98 .99	15.750 1.660	220

THETA IS THE ANGLE THAT THE K+ MAKES WITH THE BEAM IN THE GRAND C.M.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K+ PROTON.

LABORATORY BEAM MOMENTUM = 4.97 GEV/C +- .5 (PER CENT).

COS(THETA)	D-SIGMA/D-T MB/(GEV/C)**2	NO. EVENTS
MIN MAX		
-1.00 .75	.005 +- .001	14
.75 .80	.228 .064	22
.80 .82	.610 .150	20
.82 .84	.520 .140	17
.84 .86	.880 .190	29
.86 .88	1.250 .230	41
.88 .90	2.350 .390	77
.90 .92	2.720 .400	89
.92 .94	4.420 .590	145
.94 .96	6.840 .850	224
.96 .98	9.640 1.140	313
.98 .99	14.230 1.780	204

THETA IS THE ANGLE THAT THE K+ MAKES WITH THE BEAM IN THE GRAND C.M.

52 K+-P ELASTIC SCATTERING AT 3.55 GEV/C. [NUC. PHYS. 89, 640 (1969)]

J. BANAIOS, J. BERGER, C. BONNEL, J. DUFLU, L. GOLDZAM, F. PLOUIN (CNTR. D'ETUDES NUC. SACLAY, Gif-sur-Yvette, France);
 W. F. BAKER, P. J. CARLSON, V. CHABAUD, A. LUNDQVIST (EUROPEAN ORG. FOR NUC. RES., GENEVA, SWITZERLAND)

ABSTRACT: ELASTIC SCATTERING OF K+ ON PROTONS AT 3.55 GEV/C IN THE C.M. ANGULAR RANGES FROM 10 DEGREES TO 70 DEGREES ($0.16 < -T < 2$ (GEV/C)²) AND FROM 100 DEGREES TO 170 DEGREES ($3.4 < -T < 5.5$ (GEV/C)²). THESE RESULTS COMPLEMENT PREVIOUSLY PUBLISHED DATA FROM THIS GROUP WHICH COVERED THE ANGULAR REGION NEAR 180 DEGREES. FORWARD K+-P ELASTIC SCATTERING HAS A STRUCTURE NEAR $T = -1$ (GEV/C)², WHILST THE FORWARD K+-P ANGULAR DISTRIBUTION IS SMOOTH. A BACKWARD PEAK IS OBSERVED IN THE K+-P ANGULAR DISTRIBUTION, WHILST THE K+-P DIFFERENTIAL CROSS SECTION DECREASES TOWARDS 180 DEGREES.

CROSS REFERENCES

CONTINUATION OF PREVIOUS EXPERIMENT IN PHYS. LETTERS 248, 317 (1967), AND NUC. PHYS. 89, 249 (1969).

ADDITIONAL FOOTNOTE REFERENCES

NUOVO CIMENTO 45A, 889 (1966), PHYS. LETTERS 21, 117 (1966), PHYS. REV. LETTERS 19, 675 (1967), NUC. PHYS. 88, 31 (1968), INT'L. CONGRESS ON NUCLEAR PHYSICS, PARIS, FRANCE 349 (1963), CERN 65-11 57 (1965), CONF. ON PROGRAM. FOR FLYING SPOT DEVICES, MUNICH 86 (1967), PHYS. REV. LETTERS 21, 1282 (1968), UNIV. OF ROCHESTER 875-254 (1968), PHYS. LETTERS 288, 292 (1968), PHYS. REV. LETTERS 21, 1407 (1968), PHYS. REV. LETTERS 15, 838 (1965), PHYS. REV. LETTERS 17, 458 (1966), VIENNA CONFERENCE 173 (1968), VIENNA CONFERENCE 391 (1968), CERN TH-914 (1968), PHYS. REV. 166, 1768 (1968), PHYS. REV. LETTERS 20, 689 (1968), CERN TH-960 (1968), PHYS. REV. 153, 1575 (1967), PHYS. REV. LETTERS 19, 1079 (1967), UCL 8030 AUG. (1968), PHYS. LETTERS 288, 61 (1968), CERN HADRON CONFERENCE 580, AND ANL HEP 6819 (1968).

ARTICLE READ BY ODETTE BENARY IN 5/69, AND VERIFIED BY LEROY PRICE.

BEAM NO. 1 IS K+ ON PROTON AT 3.55 GEV/C.
 NO. 2 IS K- ON PROTON AT 3.55 GEV/C.

THIS EXPERIMENT USES SPARK CHAMBERS. A TOTAL OF 127000 PICTURES ARE REPORTED ON.

KEY WORDS = DIFFERENTIAL CROSS SECTION FITS MODELS

COMPOUND KEY WORDS = FITS DIFFERENTIAL CROSS SECTION

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON. (TABLE 1)

LABORATORY BEAM MOMENTUM = 3.55 $\pm$.05 GEV/C.

-T (GEV/C)**2	D-SIGMA/D-T MB/(GEV/C)**2 (1)	NO. EVENTS	COS(THETA)	D-SIGMA/D-OMEGA MB/SR (2)
.18 $\pm$.02	9.20 $\pm$.35	697	.9353	4.14 $\pm$.16
.23 $\pm$.03	8.10 $\pm$.28	859	.9186	3.64 $\pm$.12
.29 $\pm$.03	6.46 $\pm$.26	626	.8974	2.91 $\pm$.12
.35 $\pm$.03	5.30 $\pm$.24	493	.8762	2.39 $\pm$.11
.41 $\pm$.03	4.10 $\pm$.22	361	.8550	1.84 $\pm$.10
.47 $\pm$.03	3.20 $\pm$.20	266	.8337	1.44 $\pm$.09
.53 $\pm$.03	2.32 $\pm$.17	178	.8125	1.05 $\pm$.08
.59 $\pm$.03	2.01 $\pm$.17	142	.7913	.90 $\pm$.08
.65 $\pm$.03	1.69 $\pm$.16	110	.7700	.76 $\pm$.07
.71 $\pm$.03	1.37 $\pm$.15	83	.7488	.62 $\pm$.07
.79 $\pm$.03	1.03 $\pm$.10	93	.7255	.47 $\pm$.05
.89 $\pm$.05	.63 $\pm$.09	52	.6851	.29 $\pm$.04
.99 $\pm$.05	.58 $\pm$.10	35	.6498	.26 $\pm$.04
1.09 $\pm$.05	.43 $\pm$.09	24	.6144	.19 $\pm$.04
1.24 $\pm$.10	.17 $\pm$.04	19	.5513	.07 $\pm$.02
1.54 $\pm$.20	.05 $\pm$.02	9	.4552	.02 $\pm$.01

T IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K⁺) AND THE (OUTGOING K⁺).
 THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF $\pm$ 15 PER CENT.
 (2) PLUS POSSIBLE SYSTEMATIC ERROR OF $\pm$ 15 PER CENT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON. (TABLE 3)

LABORATORY BEAM MOMENTUM = 3.55 $\pm$.05 GEV/C.

(TABLE 3 SAYS THAT BELOW ARE IN MB, BUT FIGURES 3 AND 4 CLEARLY SHOW THAT THEY ARE ACTUALLY IN MICROBARN.)

U (GEV/C)**2	D-SIGMA/D-U UB/(GEV/C)**2 (1)	NO. EVENTS	COS(THETA)	D-SIGMA/D-OMEGA UB/SR (2)
.035 $\pm$.012	47.4 $\pm$ 13.1	15	-.9943	21.3 $\pm$ 5.9
.005 $\pm$.018	33.1 $\pm$ 12.9	7	-.9835	14.9 $\pm$ 5.8
-.130 $\pm$.050	20.2 $\pm$ 7.7	7	-.9358	9.1 $\pm$ 3.5
-.230 $\pm$.050	19.3 $\pm$ 6.9	7	-.9004	8.7 $\pm$ 3.1
-.380 $\pm$.100	10.2 $\pm$ 4.2	6	-.8473	4.6 $\pm$ 1.9
-.630 $\pm$.150	8.0 $\pm$ 3.5	6	-.7589	3.6 $\pm$ 1.6
-.980 $\pm$.200	9.6 $\pm$ 4.2	6	-.6351	4.3 $\pm$ 1.9
-1.490 $\pm$.300	7.4 $\pm$ 3.5	5	-.4546	3.3 $\pm$ 1.6

U IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K⁺) AND THE (PROTON).
 THETA IS THE ANGLE THAT THE PROTON MAKES WITH THE BEAM IN THE GRAND C.M.

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF $\pm$ 15 PER CENT.
 (2) PLUS POSSIBLE SYSTEMATIC ERROR OF $\pm$ 15 PER CENT.

FIT TO ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON. (TABLE 4)

LABORATORY BEAM MOMENTUM = 3.55 $\pm$.05 GEV/C.

DATA IS FIT OVER -T FROM .16 TO 1.74 (GEV/C)**2. T IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K⁺) AND THE (OUTGOING K⁺).
 FITTED FORMULA IS D-SIGMA/D-T = A*EXP(B*T)

WHERE D-SIGMA/D-T IS IN MB/(GEV/C)**2 AND -T IS IN (GEV/C)**2.

FITTED VALUES

A = 18.5 $\pm$.6
 B = 3.69 $\pm$.08

53 K⁺P ELASTIC SCATTERING AT 864, 969, AND 1207 MEV/C. (UCRL 18889 (1969))

R.W.BLAND, C.GOLDHABER, G.H.TRILLING [U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA, AND UNIV. OF CALIFORNIA, BERKELEY, CALIF., USA]

ABSTRACT THE DIFFERENTIAL CROSS SECTION FOR K⁺P ELASTIC SCATTERING HAS BEEN MEASURED AT 864, 969, AND 1207 MEV/C. OUR DATA SHOW A SMOOTH TRANSITION FROM LOW-ENERGY S-WAVE SCATTERING TO HIGH-ENERGY DIFFRACTION, AND ARE SOMEWHAT IN DISAGREEMENT WITH A RECENT EXPERIMENT ON K⁺P BACKWARD SCATTERING.

ADDITIONAL FOOTNOTE REFERENCES

PHYS. REV. LETTERS 9, 135 (1962); PHYS. REV. LETTERS 7, 188 (1961); PHYS. REV. 135B, 191 (1964); PHYS. REV. 129, 2743 (1963); PHYS. LETTERS 16, 83 (1965); PHYS. REV. 139B, 1411 (1965); NJDND CONFINT 51A, 401 (1967); PHYS. REV. LETTERS 21, 1242 (1968); PHYS. REV. LETTERS 17, 102 (1966); PHYS. REV. 168, 1466 (1968); PHYS. LETTERS 24B, 314 (1967); AND CAVENDISH LAB. REP 68-10 (1968).

ARTICLE READ BY ODETTE BENARY IN 6/69, AND VERIFIED BY ROGER BLAND.

BEAM IS K⁺ ON PROTON FROM .864 TO 1.207 GEV/C.

THIS EXPERIMENT USES THE L.R.L. 25 IN. (H) BUBBLE CHAMBER.

GENERAL COMMENTS ON THIS ARTICLE

1 THE DATA HAVE BEEN NORMALIZED TO THE ACCURATE TOTAL CROSS SECTION MEASUREMENTS OF COUL ET AL. (PRL 17, 102 (66)) AND BUGG ET AL. (PRL 168, 1466(68)). THE RESULTING ERROR IN MOMENTUM IS 2 TO 3 PER CENT.
 2 LEGENDRE-POLYNOMIAL FITS TO ELASTIC SCATTERING DATA ARE USED TO CALCULATE THE DIFFERENTIAL CROSS SECTION IN THE FORWARD AND BACKWARD DIRECTIONS.

KEY WORDS - DIFFERENTIAL CROSS SECTION FITS

COMPOUND KEY WORDS - FITS DIFFERENTIAL CROSS SECTION

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON. (TABLE 1)

LABORATORY BEAM MOMENTUM = .864 GEV/C.

COS(THETA)		D-SIGMA/D-OMEGA MB/SR		NO. EVENTS
MIN	MAX			
-1.0 [1]		1.60 +- .12		
-.9	-.8	1.30	.06	535
-.8	-.7	1.10	.06	454
-.7	-.6	.99	.05	409
-.6	-.5	.90	.05	372
-.5	-.4	.87	.05	357
-.4	-.3	.91	.05	377
-.3	-.2	.77	.05	316
-.2	-.1	.80	.05	328
-.1	.0	.85	.05	350
.0	.1	.81	.05	335
.1	.2	.92	.05	378
.2	.3	.87	.05	360
.3	.4	.86	.05	357
.4	.5	.94	.05	389
.5	.6	1.04	.05	431
.6	.7	1.06	.06	436
.7	.8	1.06	.06	437
.8	.9	1.21	.06	499
1.0 [1]		1.34	.13	

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.

[1] EXTRAPOLATED POINT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON. (TABLE 1)

LABORATORY BEAM MOMENTUM = .969 GEV/C.

COS(THETA)		D-SIGMA/D-OMEGA MB/SR		NO. EVENTS
MIN	MAX			
-1.0 [1]		1.37 +- .16		
-.9	-.8	1.70	.08	256
-.8	-.7	.95	.07	221
-.7	-.6	.81	.07	189
-.6	-.5	.75	.06	176
-.5	-.4	.83	.07	193
-.4	-.3	.74	.06	172
-.3	-.2	.74	.06	172
-.2	-.1	.81	.06	188
-.1	.0	.82	.07	192
.0	.1	.79	.06	185
.1	.2	.78	.06	184
.2	.3	.76	.06	176
.3	.4	.96	.07	225
.4	.5	1.05	.07	244
.5	.6	1.17	.08	272
.6	.7	.93	.07	218
.7	.8	1.28	.08	299
.8	.9	1.35	.08	315
1.0 [1]		1.62	.17	

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.

[1] EXTRAPOLATED POINT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON. (TABLE 1)

LABORATORY BEAM MOMENTUM = 1.207 GEV/C.

COS(THETA)		D-SIGMA/D-OMEGA MB/SR		NO. EVENTS
MIN	MAX			
-1.0 [1]		.57 +- .07		
-1.0	-.9	.57	.07	165
-.9	-.8	.49	.05	141
-.8	-.7	.38	.05	110
-.7	-.6	.41	.05	118
-.6	-.5	.35	.05	100
-.5	-.4	.51	.05	109
-.4	-.3	.43	.05	125
-.3	-.2	.45	.05	150
-.2	-.1	.46	.05	134
-.1	.0	.51	.06	147
.0	.1	.54	.06	157
.1	.2	.64	.06	192
.2	.3	.71	.06	204
.3	.4	.88	.06	255
.4	.5	1.12	.07	325
.5	.6	1.27	.08	368
.6	.7	1.65	.08	476
.7	.8	1.65	.08	470
.8	.9	2.07	.10	598
1.0 [1]		2.84	.14	

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.

[1] EXTRAPOLATED POINT.

54 POLARIZATION IN ELASTIC K^+ P SCATTERING BETWEEN 0.86 GEV/C AND 1.45 GEV/C -- RESULTS AND PHASE-SHIFT ANALYSIS. (CERN (1969))

S. ANDERSSON, C. DAUM, F. C. ERNE, J. P. LAGNAUX, J. C. SENS, F. UDDO, F. WAGNER (EUROPEAN ORG. FOR NUC. RES., GENEVA, SWITZERLAND)

ABSTRACT POLARIZATION AND DIFFERENTIAL CROSS-SECTION DATA AT 0.86, 0.97, 1.09, 1.37, AND 1.45 GEV/C ARE PRESENTED. AN ENERGY-INDEPENDENT PHASE-SHIFT ANALYSIS FROM THRESHOLD UP TO 1.45 GEV/C USING RANDOM SEARCHES AT 19 MOMENTA AND THE SHORTEST PATH METHOD TO LINK SOLUTIONS AT DIFFERENT MOMENTA, YIELDS THREE SOLUTIONS. ONE OF THESE IS UNLIKELY; THE OTHER TWO COINCIDE UP TO 0.86 GEV/C, AND BOTH SHOW AN ANTICLOCKWISE HALF CIRCLE IN THE P_{13} WAVE.

ADDITIONAL FOOTNOTE REFERENCES

UCRL 18323, PHYS. REV. 165, 1770 (1968), PHYS. LETTERS 288, 611 (1969), VIENNA CONFERENCE 485, 254 (1968), PHYS. REV. LETTERS 9, 135 (1962), PHYS. REV. LETTERS 7, 188 (1961), UCRL 18131 (1968), PHYS. REV. 135B, 191 (1964), PHYS. REV. 129, 2743 (1963), PHYS. REV. LETTERS 21, 1282 (1968), PHYS. LETTERS 16, 83 (1965), PHYS. REV. LETTERS 17, 102 (1966), PHYS. REV. 168, 1466 (1968), RUTHERFORD HIGH ENERGY LAB. RHEL 68/10 (1968), AND UCRL 18031.

ARTICLE READ BY ODETTE BENARY IN 7/69, AND VERIFIED BY ROGER BLAND.

BEAM IS K^+ ON HYDROGEN COMPOUND FROM .86 TO 1.45 GEV/C. TARGET IS POLARIZED 35 PER CENT (NORMAL TO THE BEAM DIRECTION).

THIS EXPERIMENT USES COUNTERS.

GENERAL COMMENTS ON THIS ARTICLE

1 THE DIFFERENTIAL CROSS SECTION DATA HAVE BEEN NORMALIZED TO EXISTING DATA. THEIR STATISTICAL ERRORS HAVE BEEN DOUBLED TO ACCOUNT FOR SYSTEMATIC ERRORS.

KEY WORDS = POLARIZATION DIFFERENTIAL CROSS SECTION PHASE SHIFT

ELASTIC DIFFERENTIAL CROSS SECTION FOR K^+ PROTON. (TABLE 1)

LABORATORY BEAM MOMENTUM = .865 GEV/C.

COS(THETA)	D-SIGMA/D-OMEGA MB/SR (1)
.79	1.04 +- .19
.70	1.31 .19
.64	.88 .23
.59	1.05 .22
.53	.84 .20
.48	.88 .20
.42	1.01 .31
.37	.93 .23
.31	1.05 .24
.18	.78 .08
.06	.71 .10
-.03	.87 .13
-.11	.86 .12
-.21	.90 .11
-.29	.89 .13
-.36	.79 .11
-.43	.80 .13
-.50	.85 .12
-.57	.85 .12
-.64	.86 .13
-.72	1.00 .12
-.81	.88 .18

THETA IS THE ANGLE THAT THE K^+ MAKES WITH THE BEAM IN THE GRAND C.M.

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF +- 15 PER CENT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K^+ PROTON. (TABLE 2)

LABORATORY BEAM MOMENTUM = .971 GEV/C.

COS(THETA)	D-SIGMA/D-OMEGA MB/SR (1)
.83	.96 +- .17
.74	1.15 .18
.65	1.24 .15
.60	1.22 .21
.55	1.11 .20
.50	.99 .19
.44	1.10 .22
.38	.91 .18
.17	.89 .10
.03	.79 .22
-.02	.88 .24
-.08	.85 .17
-.15	.80 .10
-.23	.82 .12
-.31	.82 .09
-.38	.72 .10
-.45	.84 .10
-.53	.74 .10
-.60	.85 .11
-.66	.87 .11
-.72	.83 .12
-.78	.87 .13
-.84	.90 .21

THETA IS THE ANGLE THAT THE K^+ MAKES WITH THE BEAM IN THE GRAND C.M.

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF +- 15 PER CENT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

[TABLE 3]

LABORATORY BEAM MOMENTUM = 1.087 GEV/C.

COS(THETA)	D-SIGMA/D-OMEGA MB/SR (1)
.85	1.24 +- .17
.78	1.39 .16
.71	1.39 .14
.63	.98 .17
.54	1.04 .09
.46	.87 .14
.41	.92 .12
.35	.80 .14
.29	.92 .30
.23	.83 .20
.14	.60 .26
.06	.50 .25
-.04	.49 .08
-.11	.55 .09
-.16	.45 .11
-.23	.49 .06
-.35	.46 .06
-.43	.50 .06
-.51	.41 .06
-.58	.50 .07
-.64	.52 .07
-.72	.51 .06
-.81	.51 .07

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF +- 15 PER CENT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

[TABLE 4]

LABORATORY BEAM MOMENTUM = 1.215 GEV/C.

COS(THETA)	D-SIGMA/D-OMEGA MB/SR (1)
.83	1.42 +- .25
.77	1.66 .19
.70	1.78 .18
.60	1.38 .13
.50	1.28 .15
.39	.95 .10
.32	.73 .12
.25	.78 .12
.18	.77 .12
.12	.45 .07
.05	.50 .09
-.01	.53 .10
-.07	.37 .07
-.16	.57 .10
-.26	.38 .07
-.35	.40 .10
-.46	.30 .07
-.55	.42 .07
-.66	.40 .09
-.74	.25 .06
-.82	.35 .12

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF +- 15 PER CENT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

[TABLE 5]

LABORATORY BEAM MOMENTUM = 1.372 GEV/C.

COS(THETA)	D-SIGMA/D-OMEGA MB/SR (1)
.87	1.76 +- .17
.80	1.77 .16
.73	1.65 .22
.68	1.51 .20
.63	1.36 .19
.57	1.16 .15
.52	1.09 .12
.45	.73 .18
.14	.49 .07
-.18	.36 .07
-.24	.31 .10
-.29	.32 .05
-.36	.28 .05
-.44	.22 .05
-.52	.27 .05
-.59	.22 .05
-.65	.25 .05
-.71	.24 .05
-.79	.22 .04
-.86	.22 .09

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF +- 15 PER CENT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON. (TABLE 6)

LABORATORY BEAM MOMENTUM = 1.453 GEV/C.

COS(THETA)	D-SIGMA/D-OMEGA MB/SR (1)
.86	1.98 ± .19
.77	1.79 .17
.70	1.68 .22
.66	1.55 .20
.61	1.30 .18
.55	1.17 .12
.46	1.02 .09
.38	.72 .09
.31	.67 .14
.14	.48 .07
-.15	.41 .00
-.21	.32 .07
-.27	.30 .06
-.35	.30 .05
-.42	.21 .04
-.50	.20 .04
-.57	.16 .04
-.64	.15 .04
-.70	.14 .04
-.76	.20 .06
-.83	.21 .04

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF ± 15 PER CENT.

ELASTIC POLARIZATION FOR K⁺ PROTON. (TABLE 1)

LABORATORY BEAM MOMENTUM = .865 GEV/C.

COS(THETA)	POLARIZATION (1)
.79	.61 ± .24
.70	.70 .12
.64	.23 .28
.59	.49 .24
.53	.85 .17
.48	.52 .27
.42	.49 .17
.33	.94 .16
.25	.69 .29
.20	.50 .25
.14	.85 .29
.06	.77 .10
-.03	.45 .17
-.11	.70 .16
-.21	.10 .14
-.29	.79 .10
-.36	.34 .16
-.43	.53 .19
-.50	.45 .17
-.57	.47 .18
-.64	.32 .18
-.72	.15 .15
-.81	.34 .24

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.

THE POLARIZATION IS OF THE PROTON ALONG THE NORMAL TO THE PRODUCTION PLANE IN THE GRAND C.M.

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF ± 15 PER CENT.

ELASTIC POLARIZATION FOR K⁺ PROTON. (TABLE 2)

LABORATORY BEAM MOMENTUM = .971 GEV/C.

COS(THETA)	POLARIZATION (1)
.83	.82 ± .23
.74	.71 .09
.65	.76 .10
.50	.77 .11
.40	.63 .11
.29	.85 .13
.20	.63 .15
.12	.84 .17
.03	.93 .23
-.05	.62 .14
-.15	.34 .13
-.23	.43 .13
-.31	.49 .13
-.38	.36 .16
-.45	.23 .13
-.53	.35 .15
-.60	.31 .15
-.66	.37 .15
-.72	.17 .17
-.78	.13 .16
-.84	.09 .25

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.

THE POLARIZATION IS OF THE PROTON ALONG THE NORMAL TO THE PRODUCTION PLANE IN THE GRAND C.M.

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF ± 15 PER CENT.

ELASTIC POLARIZATION FOR K+ PROTON. [TABLE 3]

LABORATORY BEAM MOMENTUM = 1.087 GEV/C.

COS(THETA)	POLARIZATION (1)
.85	.75 $\pm$.16
.78	.71 .10
.71	.60 .09
.63	.55 .09
.54	.68 .09
.46	.90 .19
.41	.69 .12
.35	.90 .10
.29	.90 .24
.23	.62 .26
.14	.82 .36
.06	1.21 .56
-.04	.74 .16
-.11	.85 .18
-.19	.79 .15
-.29	.67 .22
-.35	.42 .16
-.43	.41 .15
-.51	.07 .22
-.58	.13 .19
-.64	.64 .17
-.72	.28 .14
-.81	-.13 .16

THETA IS THE ANGLE THAT THE K+ MAKES WITH THE BEAM IN THE GRAND C.M.
 THE POLARIZATION IS OF THE PROTON ALONG THE NORMAL TO THE PRODUCTION PLANE IN THE GRAND C.M.

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF $\pm$ 15 PER CENT.

ELASTIC POLARIZATION FOR K+ PROTON. [TABLE 4]

LABORATORY BEAM MOMENTUM = 1.215 GEV/C.

(THIS DATA REPLACES VALUES GIVEN EARLIER IN ANDERSSON, ET AL, PHYS. LETTERS 288, 611 (1969))

(THE DATA REPORTED PREVIOUSLY IN PHYS. LETT. 288, 611 (1969) HAVE BEEN MULTIPLIED BY .85 TO ACCOUNT FOR AN ERROR IN THE TARGET POLARIZATION)

COS(THETA)	POLARIZATION (1)
.83	.54 $\pm$.09
.77	.60 .06
.70	.65 .05
.60	.71 .04
.50	.60 .06
.39	.73 .06
.32	.68 .09
.25	.75 .08
.18	.54 .08
.12	.82 .10
.05	.63 .07
-.01	.72 .10
-.07	.75 .12
-.16	.48 .09
-.26	.48 .11
-.35	.35 .13
-.46	.09 .14
-.54	.23 .10
-.66	.12 .14
-.76	.01 .13
-.82	.04 .13

THETA IS THE ANGLE THAT THE K+ MAKES WITH THE BEAM IN THE GRAND C.M.
 THE POLARIZATION IS OF THE PROTON ALONG THE NORMAL TO THE PRODUCTION PLANE IN THE GRAND C.M.

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF $\pm$ 15 PER CENT.

ELASTIC POLARIZATION FOR K+ PROTON. [TABLE 5]

LABORATORY BEAM MOMENTUM = 1.372 GEV/C.

COS(THETA)	POLARIZATION (1)
.87	.42 $\pm$.11
.80	.56 .07
.73	.48 .08
.68	.88 .17
.60	.68 .08
.52	.48 .10
.45	.70 .12
.37	.67 .09
.28	.62 .13
.20	.72 .60
-.06	.71 .24
-.13	.92 .27
-.21	.37 .20
-.29	.62 .23
-.36	.44 .20
-.44	.23 .23
-.52	.39 .20
-.59	.16 .25
-.65	.04 .24
-.71	-.06 .13
-.79	.17 .23
-.86	.43 .47

THETA IS THE ANGLE THAT THE K+ MAKES WITH THE BEAM IN THE GRAND C.M.
 THE POLARIZATION IS OF THE PROTON ALONG THE NORMAL TO THE PRODUCTION PLANE IN THE GRAND C.M.

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF $\pm$ 15 PER CENT.

ELASTIC POLARIZATION FOR K⁺ PROTON. (TABLE 6)

LABORATORY BEAM MOMENTUM = 1.453 GEV/C.

COS(THETA)	POLARIZATION (1)
.89	.45 +- .11
.82	.59 .06
.77	.58 .12
.70	.55 .06
.61	.60 .08
.55	.48 .09
.46	.79 .08
.38	.70 .11
.31	.82 .16
.25	.96 .36
.19	.53 .53
.14	1.07 .44
.07	.34 .33
-.02	.50 .65
-.10	.60 .20
-.18	.60 .14
-.27	.69 .18
-.35	.40 .19
-.42	.10 .90
-.50	.46 .24
-.57	.84 .30
-.64	-.03 .28
-.70	-.24 .32
-.76	.34 .37
-.83	.14 .44

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.
THE POLARIZATION IS OF THE PROTON ALONG THE NORMAL TO THE PRODUCTION PLANE IN THE GRAND C.M.

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF +- 15 PER CENT.

55 K⁺N CHARGE EXCHANGE REACTION AT 3 GEV/C. (PHYS. LETTERS 278, 602 (1968))

Y. GOLDSCHMIDT-CLERMONT, Y. P. MENRI, B. JONGEJANS, U. KUNDT, F. MULLER, R. L. SEKULIN, N. SHAFI, G. WOLF [EUROPEAN ORG. FOR NUC. RES., GENEVA, SWITZERLAND]
J. M. CRISPELS, J. DEBAISTEUX, M. DELABAYE, P. DUFOUR, F. GRARD, J. HEUGHEBAERT, J. NAISSE, G. THILL, R. WINDMOLERS [INST. INTERUNIV. DES SCI. NUC., BRUXELLES, BELGIUM]
K. BUCHNER, G. DEMM, G. GOEBEL, H. HUPE, T. JOLDERSMA, I. S. MITTRA, W. WITTEK [MAX-PLANCK-INST. FUR PHYS.-ASTROPHYS., MUNICH, GERM.]

ABSTRACT ABOUT 100,000 PICTURES, WITH AN AVERAGE OF ABOUT 12 K⁺ PER PICTURE, TAKEN IN THE 81 CM SACLAY DEUTERIUM BUBBLE CHAMBER EXPOSED TO A SEPARATED 3 GEV/C K⁺ BEAM HAVE BEEN ANALYZED FOR THE REACTION K⁺ + KO PP IN THE 1-PRONG + VO TOPOLOGIES. 214 SUCH EVENTS HAVE BEEN FOUND ALLOWING A DETERMINATION OF THE DIFFERENTIAL CROSS-SECTION. A COMPARISON WITH PREDICTION OF RARITA AND SCHWARZSCHILD YIELDS REASONABLE AGREEMENT; IN PARTICULAR A LARGE REAL PART IS INFERRED FOR THE AMPLITUDE FOR THE REACTION K⁺N → KO P.

ADDITIONAL FOOTNOTE REFERENCES

PHYS. REV. 162, 1378 (1967), PHYS. REV. 115, 1020 (1959), PHYS. REV. LETTERS 19, 259 (1967), PHYS. REV. 156, 1685 (1967), PHYS. REV. LETTERS 19, 734 (1965), PHYS. REV. 139, 81336 (1955), AND PHYS. REV. 156, 1522 (1967).

ARTICLE READ BY ODETTE BENARY IN 12/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K⁺ ON DEUTERON AT 2.97 GEV/C.

THIS EXPERIMENT USES THE SACLAY 81 CM (H) BUBBLE CHAMBER.

GENERAL COMMENTS ON THIS ARTICLE

1 CROSS SECTIONS ARE NORMALIZED TO TAU DECAYS.

KEY WORDS = CROSS SECTION DIFFERENTIAL CROSS SECTION MODELS CHARGE EXCHANGE

CROSS SECTION FOR K⁺ DEUTERON → KO PROTON PROTON. (PAGE 602)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS
2.97	.75 +- .08	214

DIFFERENTIAL CROSS SECTION FOR K⁺ DEUTERON → KO PROTON PROTON. (TABLE 1)

LABORATORY BEAM MOMENTUM = 2.97 GEV/C.

-T (GEV/C)**2	D-SIGMA/D-T MB/(GEV/C)**2 (1)
.025	.870 +- .240
.075	1.320 .300
.125	1.130 .280
.175	1.570 .330
.225	1.020 .250
.275	1.320 .300
.325	1.310 .300
.375	1.170 .280
.425	.980 .220
.500	.510 .130
.610	.460 .110
.770	.170 .050
.990	.067 .030
1.410	.026 .013

T IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K⁺) AND THE (KO).

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF +- 4 PER CENT.

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ NEUTRON} \rightarrow K^0 \text{ PROTON}$. [TABLE 1]

LABORATORY BEAM MOMENTUM = 2.97 GEV/C.

(CORRECTED FOR DEUTERON EFFECT, IGNORING SPIN FLIP)

-T (GEV/C)**2	D-SIGMA/D-T MB/(GEV/C)**2 [1]
.000 [2]	1.400 +- .700
.025	1.400 -.390
.075	1.560 .360
.125	1.240 .310
.175	1.670 .350
.225	1.060 .260
.275	1.320 .300
.325	1.310 .300
.375	1.170 .280
.425	.980 .220
.500	.510 .130
.610	.460 .110
.770	.170 .050
.990	.067 .030
1.410	.026 .013

T IS THE MOMENTUM TRANSFER BETWEEN THE [INCOMING K^+] AND THE [K^0].

[1] PLUS POSSIBLE SYSTEMATIC ERROR OF +- 4 PER CENT.

[2] EXTRAPOLATED POINT.

56 THE REACTION $K^+ N \pi^+ \pi^-$ AT 3 GEV/C AND THE PRODUCTION MECHANISM OF $K^+ N^*$. [NUOVO CIMENTO 39, 417 (1965)]

M. FERRO-LUZZI, R. GEORGE, Y. GOLDSCHMIDT-CLERMONT, V. P. HENRI, B. JONGEJANS, D. M. G. LEITH, G. R. LYNCH, F. MULLER, J. M. PERRFAU (EUROPEAN ORG. FOR NUC. RES., GENEVA, SWITZERLAND)

ABSTRACT IN A STUDY OF THE FOUR-PARTICLE STATES $K^+ p \rightarrow K^+ n \pi^+ \pi^-$ AT 3 GEV/C, IT IS FOUND THAT THE QUASI-TWO-BODY PROCESS $K^+ p \rightarrow K^+ n^*$ OCCURS IN LARGE PROPORTION. THE CROSS-SECTIONS FOR THE VARIOUS CHARGE AND DECAY MODES OF $K^+ n^*$ ARE IN GOOD AGREEMENT WITH THE PREDICTIONS BASED ON ISOTOPIC-SPIN INVARIANCE. THE SHARPLY PEAKED C.M. PRODUCTION ANGULAR DISTRIBUTION SUGGESTS PERIPHERALISM. WITHIN THE FRAMEWORK OF A SIMPLE ONE-MESON-EXCHANGE MODEL THE DECAY ANGULAR DISTRIBUTIONS OF K^+ AND n^* ARE CONSISTENT WITH THE ASSUMPTION THAT THE REACTION PROCEEDS PREDOMINANTLY THROUGH THE EXCHANGE OF A SINGLE PION. SUCH A MODEL IS UNABLE TO EXPLAIN THE NONZERO VALUE OF THE 'INTERFERENCE' PARAMETER PE (RMD 1.0) ($\sim 0.13 \pm 0.02$). IN ADDITION, THE MODEL REQUIRES THE INCLUSION OF A DRASTIC AD-HOC FORM FACTOR, SIMILAR TO THE ONE USED AT 1.96 GEV/C, IN ORDER TO FIT THE OBSERVED PRODUCTION ANGULAR DISTRIBUTION. A MODIFIED ONE-MESON-EXCHANGE MODEL TAKING INTO ACCOUNT THE ABSORPTION DUE TO COMPETING PROCESSES AND DEVELOPED BY JACKSON, DONOHUE, GOTTFRIED, KEYSER AND SVENSSON, DOES NOT EXPLAIN THE ABSOLUTE VALUE OF THE CROSS-SECTION, BUT PREDICTS WITH A MINIMUM OF ASSUMPTIONS, A PRODUCTION ANGULAR DISTRIBUTION AND DECAY CORRELATIONS IN REASONABLE ACCORD WITH EXPERIMENT.

ADDITIONAL FOOTNOTE REFERENCES

SIENNA CONFERENCE 1 189 (1963), CERN 63-12 (1963), PHYS. LETTERS 9, 359 (1964), NUOVO CIMENTO 36, 1101 (1965), PHYS. LETTERS 12, 255 (1964), PHYS. LETTERS 8, 144 (1964), NUOVO CIMENTO 33, 309 (1964), PHYS. REV. LETTERS 8, 140 (1962), NUOVO CIMENTO 34, 1644 (1964), PHYS. REV. LETTERS 11, 90 (1963), PHYS. LETTERS 12, 352 (1964), PHYS. REV. LETTERS 12, 522 (1964), NUOVO CIMENTO 33, 906 (1964), AND NUOVO CIMENTO 34, 753 (1964).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 3/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K^+ ON PROTON AT 2.97 GEV/C.

THIS EXPERIMENT USES THE SACLAY 81 CM (H) BUBBLE CHAMBER. A TOTAL OF 100000 PICTURES ARE REPORTED ON.

KEY WORDS = CROSS SECTION MASS SPECTRUM ANGULAR DISTRIBUTION DENSITY MATRIX ONE-PARTICLE EXCHANGE
MODELS $K^+(890)$ $\Delta(1232)$

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \text{ PROTON } \pi^+ \pi^-$.

(TOTAL, INCLUDING RESONANCE PRODUCTION)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS
2.970 +- .015	2.3 +- .3	1009

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \text{ PROTON } \pi^+ \pi^0$.

(TOTAL, INCLUDING RESONANCE PRODUCTION)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS
2.970 +- .015	2.1 +- .3	739

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \text{ NEUTRON } \pi^+ \pi^+$.

(TOTAL, INCLUDING RESONANCE PRODUCTION)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS
2.970 +- .015	.6 +- .2	208

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+(890) \Delta(1232) \rightarrow K^+ \pi^- \pi^+$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.970 +- .015	1.2 +- .2

[1] COUNTED ONLY EVENTS ABOVE BACKGROUND.

[2] COUNTED ONLY EVENTS ABOVE BACKGROUND.

CROSS SECTION FOR $K^+ \text{ PROTON} = K^*(89010) \text{ DELTA}^{++}$
 $K^*(89010) = K^0 \text{ P}^0 \{1\}$
 $\text{DELTA}^{++} = \text{PROTON P}^0 \{2\}$

LABORATORY
 BEAM MOMENTUM
 GEV/C
 2.970 $\pm$.015
 MILLI-BARNS
 .46 $\pm$.09

\{1\} COUNTED ONLY EVENTS ABOVE BACKGROUND.
 \{2\} COUNTED ONLY EVENTS ABOVE BACKGROUND.

CROSS SECTION FOR $K^+ \text{ PROTON} = K^*(89010) \text{ DELTA}^+$
 $K^*(89010) = K^0 \text{ P}^0 \{1\}$
 $\text{DELTA}^+ = \text{PROTON P}^0 \{2\}$

LABORATORY
 BEAM MOMENTUM
 GEV/C
 2.970 $\pm$.015
 MILLI-BARNS
 .18 $\pm$.04

\{1\} COUNTED ONLY EVENTS ABOVE BACKGROUND.
 \{2\} COUNTED ONLY EVENTS ABOVE BACKGROUND.

CROSS SECTION FOR $K^+ \text{ PROTON} = K^*(89010) \text{ DELTA}^+$
 $K^*(89010) = K^0 \text{ P}^0 \{1\}$
 $\text{DELTA}^+ = \text{NEUTRON P}^0 \{2\}$

LABORATORY
 BEAM MOMENTUM
 GEV/C
 2.970 $\pm$.015
 MILLI-BARNS
 .11 $\pm$.05

\{1\} COUNTED ONLY EVENTS ABOVE BACKGROUND.
 \{2\} COUNTED ONLY EVENTS ABOVE BACKGROUND.

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} = K^*(89010) \text{ DELTA}^{++}$
 $K^*(89010) = K^+ \text{ P}^0 \{1\}$
 $\text{DELTA}^{++} = \text{PROTON P}^0 \{2\}$

LABORATORY BEAM MOMENTUM = 2.970 $\pm$.015 GEV/C.

.....
 THIS DATA WAS READ FROM A GRAPH

COS(THETA)		D-SIGMA/D-OMEGA UB/SR \{3\}		NO. EVENTS
MIN	MAX			
-1.00	-.95	.02 $\pm$.02		1
-.95	-.90	.02 .02		1
-.90	-.85	.00 .02		0
-.85	-.80	.00 .02		0
-.80	-.75	.02 .02		1
-.75	-.70	.00 .02		0
-.70	-.65	.02 .02		1
-.65	-.60	.02 .02		1
-.60	-.55	.02 .02		1
-.55	-.50	.00 .02		0
-.50	-.45	.02 .02		1
-.45	-.40	.02 .02		1
-.40	-.35	.02 .02		1
-.35	-.30	.03 .02		2
-.30	-.25	.00 .02		0
-.25	-.20	.00 .02		0
-.20	-.15	.02 .02		1
-.15	-.10	.00 .02		0
-.10	-.05	.00 .02		0
-.05	.00	.00 .02		0
.00	.05	.02 .02		1
.05	.10	.05 .03		3
.10	.15	.07 .03		4
.15	.20	.00 .02		0
.20	.25	.03 .02		2
.25	.30	.03 .02		2
.30	.35	.00 .02		0
.35	.40	.03 .02		2
.40	.45	.03 .02		2
.45	.50	.07 .03		4
.50	.55	.03 .02		2
.55	.60	.13 .05		8
.60	.65	.05 .03		3
.65	.70	.10 .04		6
.70	.75	.12 .04		7
.75	.80	.18 .06		11
.80	.85	.25 .06		15
.85	.90	.47 .09		28
.90	.95	.85 .12		51
.95	1.00	1.50 .16		90

THETA IS THE ANGLE THAT THE $K^*(89010)$ MAKES WITH THE BEAM IN THE GRAND C.M.

\{1\} COUNTED ALL EVENTS IN MASS BAND [MASS = .900 GEV; WIDTH = .080 GEV].
 \{2\} COUNTED ALL EVENTS IN MASS BAND [MASS = 1.225 GEV; WIDTH = .130 GEV].
 \{3\} COUNTS WERE MULTIPLIED BY .0167 TO GET THESE. ERRORS ARE TAKEN AS PROPORTIONAL TO THE SQUARE-ROOT OF THE COUNTS.

DIFFERENTIAL CROSS SECTION FOR K+ PROTON = K*(89010 DELTA+
 K*(89010 = K0 P10 [1]
 DELTA+ = PROTON P1+ [2]

LABORATORY BEAM MOMENTUM = 2.970 +- .015 GEV/C.

 . THIS DATA WAS READ FROM A GRAPH .

COS(THETA)		D-SIGMA/D-OMEGA UB/SR [3]		NO. EVENTS
MIN	MAX			
-1.00	-.95	.02 +- .02		1
-.95	-.90	.02	.02	1
-.90	-.85	.00	.02	0
-.85	-.80	.00	.02	0
-.80	-.75	.00	.02	0
-.75	-.70	.03	.02	2
-.70	-.65	.02	.02	1
-.65	-.60	.00	.02	0
-.60	-.55	.00	.02	0
-.55	-.50	.02	.02	1
-.50	-.45	.00	.02	0
-.45	-.40	.00	.02	0
-.40	-.35	.00	.02	0
-.35	-.30	.00	.02	0
-.30	-.25	.02	.02	1
-.25	-.20	.00	.02	0
-.20	-.15	.00	.02	0
-.15	-.10	.02	.02	1
-.10	-.05	.02	.02	1
-.05	.00	.00	.02	0
.00	.05	.03	.02	2
.05	.10	.00	.02	0
.10	.15	.00	.02	0
.15	.20	.02	.02	1
.20	.25	.02	.02	1
.25	.30	.00	.02	0
.30	.35	.03	.02	2
.35	.40	.00	.02	0
.40	.45	.03	.02	2
.45	.50	.03	.02	2
.50	.55	.03	.02	2
.55	.60	.05	.03	3
.60	.65	.03	.02	2
.65	.70	.10	.04	6
.70	.75	.03	.02	2
.75	.80	.12	.04	7
.80	.85	.13	.05	8
.85	.90	.10	.04	6
.90	.95	.30	.07	18
.95	1.00	.48	.09	29

THETA IS THE ANGLE THAT THE K*(89010 MAKES WITH THE BEAM IN THE GRAND C.M.

[1] COUNTED ALL EVENTS IN MASS BAND I MASS = .900 GEV; WIDTH = .080 GEV I.

[2] COUNTED ALL EVENTS IN MASS BAND I MASS = 1.225 GEV; WIDTH = .130 GEV I.

[3] COUNTS WERE MULTIPLIED BY .0167 TO GET THESE. ERRORS ARE TAKEN AS PROPORTIONAL TO THE SQUARE-ROOT OF THE COUNTS.

 DIFFERENTIAL CROSS SECTION FOR K+ PROTON = K*(89010 DELTA+
 K*(89010 = K0 P1+ [1]
 DELTA+ = PROTON P1+ [2]

LABORATORY BEAM MOMENTUM = 2.970 +- .015 GEV/C.

 . THIS DATA WAS READ FROM A GRAPH .

COS(THETA)		D-SIGMA/D-OMEGA UB/SR [3]		NO. EVENTS
MIN	MAX			
-1.00	-.95	.00 +- .01		0
-.95	-.90	.00	.01	0
-.90	-.85	.00	.01	0
-.85	-.80	.00	.01	0
-.80	-.75	.00	.01	0
-.75	-.70	.00	.01	0
-.70	-.65	.00	.01	0
-.65	-.60	.01	.01	1
-.60	-.55	.01	.01	1
-.55	-.50	.00	.01	0
-.50	-.45	.00	.01	0
-.45	-.40	.00	.01	0
-.40	-.35	.00	.01	0
-.35	-.30	.00	.01	0
-.30	-.25	.00	.01	0
-.25	-.20	.00	.01	0
-.20	-.15	.02	.01	2
-.15	-.10	.00	.01	0
-.10	-.05	.00	.01	0
-.05	.00	.00	.01	0
.00	.05	.00	.01	0
.05	.10	.00	.01	0
.10	.15	.01	.01	1
.15	.20	.01	.01	1
.20	.25	.00	.01	0
.25	.30	.00	.01	0
.30	.35	.00	.01	0
.35	.40	.01	.01	1
.40	.45	.00	.01	0
.45	.50	.00	.01	0
.50	.55	.00	.01	0
.55	.60	.00	.01	0
.60	.65	.00	.01	0
.65	.70	.02	.01	2
.70	.75	.06	.02	6
.75	.80	.02	.01	2
.80	.85	.03	.02	3
.85	.90	.05	.02	5
.90	.95	.10	.03	10
.95	1.00	.31	.06	31

THETA IS THE ANGLE THAT THE K*(89010 MAKES WITH THE BEAM IN THE GRAND C.M.

[1] COUNTED ALL EVENTS IN MASS BAND I MASS = .900 GEV; WIDTH = .080 GEV I.

[2] COUNTED ALL EVENTS IN MASS BAND I MASS = 1.225 GEV; WIDTH = .130 GEV I.

[3] COUNTS WERE MULTIPLIED BY .01 TO GET THESE. ERRORS ARE TAKEN AS PROPORTIONAL TO THE SQUARE-ROOT OF THE COUNTS.

57 K⁺P INTERACTIONS AT 2 BEV/C--ELASTIC SCATTERING AND TOTAL CROSS SECTIONS. [PHYS. REV. 139, B1411 (1965)]

W. CHINOWSKY, G. GOLDHABER, S. GOLDHABER, T. A. O'HALLORAN, B. M. SCHWARZSCHILD (U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA, AND UNIV. OF CALIFORNIA, BERKELEY, CALIF., USA)

ABSTRACT STUDY OF 1.96-BEV/C K⁺P INTERACTIONS IN THE BROOKHAVEN 20-IN. LIQUID-HYDROGEN BUBBLE CHAMBER HAS YIELDED A MEASUREMENT OF THE ELASTIC-SCATTERING CROSS SECTION $\sigma_{\text{EL}} = 7.5 \pm 0.7$ MB. A FIT TO THE DIFFERENTIAL CROSS SECTION VERSUS MOMENTUM TRANSFER OF THE FORM $(\sigma_{\text{EL}}(t=0)) \exp(-\alpha|t|)$ GIVES, FOR THE INTERVAL $0.01 (\text{BEV}/C)^2 < t < 0.60 (\text{BEV}/C)^2$, $\alpha = 3.1 \pm 0.3 (\text{BEV}/C)^{-2}$, AND $\sigma_{\text{EL}}(t=0) = 4.8 \pm 0.5$ MB/SR. THE TOTAL CROSS SECTION OBTAINED IS 19.4 ± 2.0 MB WITH SINGLE-PION PRODUCTION DOMINANT.

CROSS REFERENCES
SOME OF THIS DATA RENORMALIZED IN UCRL 18758 (1969).

ADDITIONAL FOOTNOTE REFERENCES
PHYS. REV. 73, 1532 (1949), NUOVO CIMENTO 21, 127 (1961), NUOVO CIMENTO 21, 823 (1961), PHYS. REV. LETTERS 10, 357 (1963), PHYS. REV. LETTERS 7, 394 (1961), PHYS. REV. 126, 2204 (1952), PHYS. REV. 126, 1232 (1962), PHYS. REV. 126, 766 (1962), PHYS. REV. LETTERS 8, 142 (1962), UCRL 11068 (1963), UCRL 9099 (1961), PHYS. LETTERS 6, 62 (1963), PHYS. REV. LETTERS 7, 346 (1961), UCRL 8030, I (1964), PHYS. REV. 124, 2743 (1953), AND PHYS. REV. 135, B1263 (1964).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 1/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K⁺ ON PROTON AT 1.96 GEV/C.

THIS EXPERIMENT USES THE B.N.L. 20 IN. (H) BUBBLE CHAMBER.

KEY WORDS = CROSS SECTION ANGULAR DISTRIBUTION

K⁺ PROTON TOTAL CROSS SECTION.

(RENORMALIZED VALUES BELOW TAKE THIS TO BE $17.65 \pm .08$ MB.)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
1.96 $\pm$.02	19.4 $\pm$ 2.0

K⁺ PROTON ELASTIC CROSS SECTION. [PAGE 1412]

ORIGINAL DATA		RENORMALIZED VALUE [1]	
LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS	MILLI-BARNS
1.96 $\pm$.02	7.5 $\pm$.7	636	6.8 $\pm$.5

[1] BLAND, ET. AL., UCRL 18758 (1969).

CROSS SECTION FOR K⁺ PROTON $\rightarrow$ K⁰ PI⁺ PROTON. [TABLE III]

ORIGINAL DATA		RENORMALIZED VALUE [1]	
LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS		MILLI-BARNS
1.96 $\pm$.02	4.6 $\pm$.6		4.1 $\pm$.5

[1] BLAND, ET. AL., UCRL 18758 (1969).

CROSS SECTION FOR K⁺ PROTON $\rightarrow$ K⁺ PI⁰ PROTON. [TABLE III]

ORIGINAL DATA		RENORMALIZED VALUE [1]	
LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS		MILLI-BARNS
1.96 $\pm$.02	2.0 $\pm$.3		1.9 $\pm$.3

[1] BLAND, ET. AL., UCRL 18758 (1969).

CROSS SECTION FOR K⁺ PROTON $\rightarrow$ K⁺ PI⁺ NEUTRON. [TABLE III]

ORIGINAL DATA		RENORMALIZED VALUE [1]	
LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS		MILLI-BARNS
1.96 $\pm$.02	1.6 $\pm$.3		1.5 $\pm$.3

[1] BLAND, ET. AL., UCRL 18758 (1969).

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K \text{ MESON NUCLEON PION.}$

ORIGINAL DATA		RENORMALIZED VALUE [1]
LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	MILLI-BARNS
1.96 $\pm$.02	8.20 $\pm$.74	7.40 $\pm$.55

[1] BLAND, ET AL., UCRL 18758 (1969).

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \text{ PI}^- \text{ PROTON PI}^+.$

[TABLE III]

ORIGINAL DATA		RENORMALIZED VALUE [1]
LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	MILLI-BARNS
1.96 $\pm$.02	1.7 $\pm$.2	1.64 $\pm$.20

[1] BLAND, ET. AL., UCRL 18758 (1969).

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \text{ PI}^0 \text{ PROTON PI}^+.$

[TABLE III]

ORIGINAL DATA		RENORMALIZED VALUE [1]
LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	MILLI-BARNS
1.96 $\pm$.02	1.3 $\pm$.2	1.24 $\pm$.20

[1] BLAND, ET. AL., UCRL 18758 (1969).

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \text{ PI}^+ \text{ NEUTRON PI}^+.$

[TABLE III]

ORIGINAL DATA		RENORMALIZED VALUE [1]
LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	MILLI-BARNS
1.96 $\pm$.02	.33 $\pm$.10	.31 $\pm$.13

[1] BLAND, ET. AL., UCRL 18758 (1969).

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \text{ PI}^0 \text{ NEUTRON PI}^+.$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS [1]
1.96 $\pm$.02	.3

[1] VALUE IS APPROXIMATE ONLY.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \text{ PI}^- \text{ PI}^+ \text{ PI}^0 \text{ PROTON.}$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
1.96 $\pm$.02	.05 $\pm$.02

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \text{ PI}^+ \text{ PI}^+ \text{ PI}^- \text{ PROTON.}$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
1.96 $\pm$.02	.02 $\pm$.01

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \text{ PI}^- \text{ PI}^+ \text{ PI}^+ \text{ NEUTRON.}$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
1.96 $\pm$.02	.010 $\pm$.006

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = 1.96 ± .02 GEV/C.
NUMBER OF EVENTS = 636.

THIS DATA IS EQUIVALENT TO OTHER DATA FROM THIS SAME ARTICLE. IT IS ALSO CALLED SET NO. 10

.....
THIS DATA WAS READ FROM A GRAPH.
.....

ORIGINAL DATA		RENORMALIZED VALUES (1)	
-T (GEV/C)**2	D-SIGMA/D-OMEGA MB/SR		MILLI-BARNS
MIN	MAX		
.01	.20	3.400 ± .200	3.094 ± .182
.20	.40	1.800 .200	1.638 .182
.40	.60	1.100 .150	1.001 .136
.60	.80	.700 .120	.537 .109
.80	1.00	.275 .060	.250 .055
1.00	1.20	.175 .030	.159 .027
1.20	1.40	.086 .030	.078 .027
1.40	1.60	.113 .030	.103 .027
1.60	1.80	.050 .030	.045 .027
1.80	2.00	.040 .020	.036 .018
2.00	2.60	.025 .020	.023 .018
2.60	2.75	.035 .020	.032 .018

T IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K⁺ AND THE (OUTGOING K⁺).

(1) ORIGINAL VALUES HAVE BEEN MULTIPLIED BY .91 - THE SAME FACTOR THAT THE INTEGRATED CROSS SECTION WAS MULTIPLIED BY TO GET THE RE-NORMALIZED VALUES OF BLAND, ET. AL., UCRL 18759 (1969).

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = 1.96 ± .02 GEV/C.
NUMBER OF EVENTS = 636.

THIS DATA IS EQUIVALENT TO OTHER DATA FROM THIS SAME ARTICLE. IT IS ALSO CALLED SET NO. 10

.....
THIS DATA WAS READ FROM A GRAPH.
.....

ORIGINAL DATA		RENORMALIZED VALUES (1)	
COS(THETA)	D-SIGMA/D-OMEGA MB/SR	NO. EVENTS	MILLI-BARNS
MIN	MAX		
-1.0	-.9	.03 ± .03	.03 ± .03
-.9	-.8	.03 .03	.03 .03
-.8	-.7	.02 .02	.02 .02
-.7	-.6	.03 .03	.03 .03
-.6	-.5	.02 .02	.02 .02
-.5	-.4	.00 .02	.00 .02
-.4	-.3	.06 .04	.05 .04
-.3	-.2	.05 .03	.05 .03
-.2	-.1	.06 .04	.05 .04
-.1	0	.17 .06	.15 .05
0	.1	.08 .05	.07 .05
.1	.2	.08 .05	.07 .05
.2	.3	.26 .06	.24 .05
.3	.4	.30 .06	.27 .05
.4	.5	.54 .09	.49 .08
.5	.6	.89 .11	.81 .10
.6	.7	1.11 .11	1.01 .10
.7	.8	1.65 .17	1.50 .15
.8	.9	2.47 .20	2.25 .18
.9	1.0	6.02 .25	3.66 .23

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.

(1) ORIGINAL VALUES HAVE BEEN MULTIPLIED BY .91 - THE SAME FACTOR THAT THE INTEGRATED CROSS SECTION WAS MULTIPLIED BY TO GET THE RE-NORMALIZED VALUES OF BLAND, ET. AL., UCRL 18758 (1969).

58 COHERENT K⁺ PRODUCTION IN K⁺D REACTIONS AT 3 GEV/C. [NUC. PHYS. 89, 286 (1969)]

K. BUCHNER, G. DEHM, G. GOEBEL, H. HUPE, T. JOLDERSMA, I. S. MITTRA, W. WITTEK [MAX-PLANCK-INST. FÜR PHYS.-ASTROPHYS., MUNICH, GERM.]
J. M. CRISPEL, J. DEBAIS, J. DELABAY, P. DUFOUR, F. GRARD, J. HEUGHERAERT, J. NATSSE, G. THILL [INST. INTERUNIV. DES SCI. NUC.,
BRUXELLES, BELGIUM]
A. GRANT, V. P. HENRI, B. JONGEJANS, U. KUNDT, F. MÜLLER, R. L. SEKULIN, G. WOLF [EUROPEAN ORG. FOR NUC. RES., GENEVA, SWITZERLAND]

ABSTRACT IN A STUDY OF K⁺D INTERACTIONS AT 3 GEV/C WE HAVE OBSERVED THE FOLLOWING REACTIONS IN WHICH THE DEUTERON DOES NOT BREAK UP K⁺D → K⁺ p (K⁺ SEEN), K⁺D → K⁺ p + p + D. AN ABUNDANT PRODUCTION OF K⁺(892) IS OBSERVED IN THESE REACTIONS. THE PRODUCTION AND DECAY ANGULAR DISTRIBUTIONS FOR K⁺(892) ARE PRESENTED.

ADDITIONAL FOOTNOTE REFERENCES
PHYS. REV. LETTERS 15, 500 (1965), NUCOVO CIMENTO 34, 1644 (1964), NUCOVO CIMENTO 11, 670 (1959), HEIDELBERG CONFERENCE (1967), AND PHYS. LETTERS 26B, 636 (1968).

ARTICLE READ BY ODETTE BENARY IN 3/69, AND VERIFIED BY LEROY PRICE.

BEAM IS K⁺ ON DEUTERON AT 2.97 GEV/C.

THIS EXPERIMENT USES THE SACLAY 81 CM (H) BUBBLE CHAMBER. A TOTAL OF 100000 PICTURES ARE REPRODUCED ON.

GENERAL COMMENTS ON THIS ARTICLE

1 THE CROSS SECTIONS ARE NORMALIZED TO TAU DECAYS.

KEY WORDS = CROSS SECTION DALITZ PLOT K⁺(890) DENSITY MATRIX ANGULAR DISTRIBUTION FITS

COMPOUND KEY WORDS = K⁺(890) DENSITY MATRIX FITS ANGULAR DISTRIBUTION

CROSS SECTION FOR $K^+ \text{ DEUTERON} \rightarrow K^0 \pi^+ \text{ DEUTERON}$. (PAGE 287)

LABORATORY
BEAM MOMENTUM
GEV/C
2.97
MICRO-BARNS
228. $\pm$ 25.

CROSS SECTION FOR $K^+ \text{ DEUTERON} \rightarrow K^+ \pi^- \pi^+ \text{ DEUTERON}$. (PAGE 289)

LABORATORY
BEAM MOMENTUM
GEV/C
2.97
MICRO-BARNS
210. $\pm$ 22.

FIT TO DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ DEUTERON} \rightarrow K^*(890)^+ \text{ DEUTERON}$. (PAGE 289)
 $K^*(890)^+ \rightarrow K^0 \pi^+ \{1\}$

LABORATORY BEAM MOMENTUM = 2.97 GEV/C.

-T (GEV/C)**2 < .250. T IS THE MOMENTUM TRANSFER BETWEEN THE [INCOMING K+] AND THE [K*(890)+].
FITTED FORMULA IS NO. EVENTS = EXP(A-B*T)

WHERE -T IS IN (GEV/C)**2.

FITTED VALUE

B = 11.9 $\pm$ 3.5

{1} COUNTED ALL EVENTS IN MASS BAND.

FIT TO DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ DEUTERON} \rightarrow K^*(890) \pi^+ \text{ DEUTERON}$. (PAGE 292)
 $K^*(890) \rightarrow K^+ \pi^- \{1\}$

LABORATORY BEAM MOMENTUM = 2.97 GEV/C.

(THE FIT WAS DONE TAKING ONLY EVENTS HAVING (T-TMIN) LESS THAN .2 GEV/C**2)
FITTED FORMULA IS NO. EVENTS = EXP(A-B*(T-TMIN))

WHERE -T IS IN (GEV/C)**2. T IS THE MOMENTUM TRANSFER BETWEEN THE [INCOMING K+] AND THE [K*(890)0].

FITTED VALUE

B = 11.9 $\pm$ 1.8

{1} COUNTED ALL EVENTS IN MASS BAND.

FIT TO DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ DEUTERON} \rightarrow \text{DEUTERON } K^+ \pi^- \pi^+$. (PAGE 292)

LABORATORY BEAM MOMENTUM = 2.97 GEV/C.

(THE FIT WAS DONE TAKING ONLY EVENTS HAVING (T-TMIN) LESS THAN .2 GEV/C**2)
FITTED FORMULA IS NO. EVENTS = EXP(A-B*(T-TMIN))

WHERE -T IS IN (GEV/C)**2. T IS THE MOMENTUM TRANSFER BETWEEN THE [TARGET] AND THE [DEUTERON].

FITTED VALUE

B = 21.3 $\pm$ 1.6

59 A SEARCH FOR THE KAPPA MESON. [PHYS. LETTERS 22, 347 (1966)]

A.T.GOSHAW, A.R.ERWIN, W.D.WALKER, A.WEINBERG (UNIV. OF WISCONSIN, MADISON, WISC., USA)

ABSTRACT AN EXPERIMENT HAS BEEN COMPLETED IN WHICH WE LOOKED AT THE REACTION $K^+ p \rightarrow K^0 \pi^+ \pi^- \pi^+ p$ USING 3.52 GEV/C K^+ MESONS IN THE NBL 80 IN. CHAMBER. THIS TYPE OF REACTION PRODUCED A HIGH YIELD OF K^+ 'S FOR AN INCOMING K^+ OF 3.0 GEV/C. NO EVIDENCE IS FOUND OF THE KAPPA AT THIS ENERGY. SOME EVIDENCE OF KAPPA PRODUCTION IS SEEN IN THE REACTION $K^+ p \rightarrow K^0 \Omega(10^1) \pi^+ p$. THE CONCLUSION IS THAT THE K IS NOT A RESONANT STATE.

ADDITIONAL FOOTNOTE REFERENCES

PHYS. LETTERS 5, 279 (1963), PHYS. LETTERS 12, 255 (1964), PHYS. LETTERS 5, 283 (1963), PHYS. REV. 139, 81093 (1965), OXFORD CONFERENCE RAPTR.TALK (1965), PHYS. LETTERS 19, 350 (1965), AND PHYS. REV. 143, 1034 (1966).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 4/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K^+ ON PROTON AT 3.535 GEV/C.

THIS EXPERIMENT USES THE R.N.L. 80 IN. (H) BUBBLE CHAMBER.

KEY WORDS = KAPPA MESON MASS SPECTRUM

* NO DATA PUNCHED FOR THIS ARTICLE *

60 STUDY OF THE REACTIONS $K^+p \rightarrow K^+p$ AT 735 AND 785 MEV/C. [NUOVO CIMENTO 51A, 1053 (1967)]

T.A. FILIPPAS, V.P. HENRI, B. JONGEJANS, M. KRAMMER, J.M. PERREAU (EUROPEAN ORG. FOR NUC. RES., GENEVA, SWITZERLAND)
S. FOCARDI, A. MINGUZZI-RANZI, L. MONARI, G. SALTINI, P. SERRA (UNIV. DI BOLOGNA, BOLOGNA, ITALY, AND ISTITUTO DI FISICA DELLA
UNIVERSITA, MODENA, ITALY)
E. BARRELET, E. HUFFER, F. MULLER (ECOLE POLYTECHNIQUE, PARIS, FRANCE)

ABSTRACT. INCOMING K^+ MOMENTA OF 735 AND 785 MEV/C WERE CHOSEN TO STUDY THE REACTIONS $K^+p \rightarrow K^+p$ BELOW THE $K^*(1236)$ AND $N^*(1236)$ PRODUCTION THRESHOLDS IN THE HOPE OF OBSERVING THE PRODUCTION OF THE KAPPA-MESON IF IT EXISTS. A BROAD ENHANCEMENT IS OBSERVED IN THE $(K^+p)^+$ MASS SPECTRUM, WHICH COULD BE ATTRIBUTED TO THE PRODUCTION OF A KAPPA-MESON OF MASS APPROXIMATELY 725 MEV, WIDTH APPROXIMATELY 50 MEV AND CROSS-SECTION < 200 MICROBARNS AT 785 MEV/C. A MORE LIKELY EXPLANATION IS OFFERED BY ASSUMING AN INTERFERENCE BETWEEN THE LOW-ENERGY TAIL OF THE $N^*(1236)$ AND AN S-WAVE $p \rightarrow p$ BACKGROUND WITHOUT INVOKING THE PRODUCTION OF KAPPA-MESONS. SIMPLE ASSUMPTIONS CONCERNING THE AMPLITUDES OF THE N^* TAIL AND OF THE $p \rightarrow p$ BACKGROUND LEAD TO PRODUCTION AND DECAY DISTRIBUTIONS IN GOOD AGREEMENT WITH THE EXPERIMENTAL SPECTRA.

CROSS REFERENCES
SEE ALSO PHYS. LETTERS 240, 314 (1967).

ADDITIONAL FOOTNOTE REFERENCES

CERN CONFERENCE 322 (1962), PHYS. LETTERS 5, 279 (1963), PHYS. LETTERS 5, 283 (1963), PHYS. LETTERS 12, 255 (1964),
PHYS. LETTERS 19, 350 (1965), SIENNA CONFERENCE 125 (1963), PHYS. REV. 143, 1034 (1966), DUBNA CONFERENCE 607 (1964),
PHYS. LETTERS 22, 347 (1966), PHYS. REV. LETTERS 17, 838 (1966), CERN 65-2 (1965), PHYS. LETTERS 24, 8314 (1967),
REV. MOD. PHYS. 39, 1 (1967), PHYS. REV. 133, 8220 (1964), UCRL 16370, ANNALS OF PHYSICS 29, 19 (1964), PHYS. REV.
LETTERS 11, 93 (1963), PHYS. LETTERS 16, 83 (1965), AND NUOVO CIMENTO 44A, 647 (1966).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 6/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K^+ ON PROTON FROM .735 TO .785 GEV/C.

THIS EXPERIMENT USES THE SACLAY 81 CM (H) BUBBLE CHAMBER. A TOTAL OF 175000 PICTURES ARE REPORTED ON.

GENERAL COMMENTS ON THIS ARTICLE

1 CROSS SECTIONS ARE NORMALIZED TO TAU DECAYS. BRANCHING RATIO OF TAU/ALL = .056. K^+ LIFETIME = 1.235×10^{-8} SEC.

KEY WORDS = CROSS SECTION DALITZ PLOT MASS SPECTRUM KAPPA MESON ANGULAR DISTRIBUTION DELTA(1236)

CROSS SECTION FOR $K^+ \text{ PROTON } \rightarrow K^0 \text{ PI}^+ \text{ PROTON.}$ [TABLE 1]

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.735 $\pm$.030	.13 $\pm$.01
.785 .030	.34 .03

CROSS SECTION FOR $K^+ \text{ PROTON } \rightarrow K^+ \text{ P}^0 \text{ PROTON.}$ [TABLE 1]

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.735 $\pm$.030	.03 $\pm$.01
.755 .815	.11 .01

CROSS SECTION FOR $K^+ \text{ PROTON } \rightarrow K^+ \text{ PI}^+ \text{ NEUTRON.}$ [TABLE 1]

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.735 $\pm$.030	.02 $\pm$.01
.785 .030	.07 .01

CROSS SECTION FOR $K^+ \text{ PROTON } \rightarrow K^+ \text{ MESON NUCLEON PION.}$ [TABLE 1]

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.735 $\pm$.030	.18 $\pm$.01
.785 .030	.52 .03

CROSS SECTION FOR $K^+ \text{ PROTON } \rightarrow K^0 \text{ DELTA}^{++}$. [TABLE 4]

(ASSUMES COHERENT BACKGROUND WITHOUT KAPPA.)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS (1)
.785 $\pm$.030	.14

(1) VALUE IS APPROXIMATE ONLY.

61 K⁺P BACKWARD SCATTERING IN THE REGION FROM 1.0 TO 2.5 GEV/C. [PHYS. REV. LETTERS 21, 1282 (1968)]

A.S. CARROLL, J. FISCHER, A. LUNDBY, R. H. PHILLIPS, C. L. WANG [BROOKHAVEN NAT. LAB., UPTON, L.I., N. Y., USA]
F. LOBSCOWICZ, A. C. MELISSINOS, Y. NAGASHIMA, C. A. SMITH, S. TENKSURRY [UNIV. OF ROCHESTER, ROCHESTER, N. Y., USA]

ABSTRACT ELASTIC SCATTERING OF K⁺ MESONS FROM PROTONS IN THE BACKWARD DIRECTION HAS BEEN MEASURED TO HIGH ACCURACY. THE DATA INDICATE THAT BARYON EXCHANGE IS EXTREMELY IMPORTANT FOR P-LAB(K) > 1.8 GEV/C. A PHASE SHIFT ANALYSIS OF THE DATA SUGGESTS THE EXISTENCE OF A P 1/2 RESONANCE IN THE K⁺P SYSTEM AT A MASS OF APPROXIMATELY 2 GEV/C-SQUARED.

ADDITIONAL FOOTNOTE REFERENCES

PHYS. REV. LETTERS 20, 607 (1968), DUBNA CONFERENCE 4 109 (1967), BNL 11331, PHYS. REV. LETTERS 19, 460 (1967), PHYS. REV. LETTERS 19, 1504 (1967), PHYS. REV. 153, 1575 (1967), PHYS. REV. 155, 1792 (1967), PHYS. REV. 165, 1770 (1968), PHYS. LETTERS 248, 317 (1967), PHYS. REV. LETTERS 17, 102 (1966), PHYS. REV. LETTERS 19, 259 (1967), PHYS. REV. 168, 1466 (1968), PHYS. REV. LETTERS 16, 709 (1966), PHYS. REV. LETTERS 18, 798 (1966), AND PHYS. REV. 155, 1792 (1967).

ARTICLE READ BY LEROY PRICE IN 10/68, AND VERIFIED BY ROGER BLAND.

BEAM IS K⁺ ON PROTON FROM 1.0 TO 2.5 GEV/C.

THIS EXPERIMENT USES SPARK CHAMBERS.

GENERAL COMMENTS ON THIS ARTICLE

1. THIS PAPER REPORTS ONLY ON BACKWARD K⁺P ELASTIC SCATTERING (COS(THETA) = -1 TO -.7).

KEY WORDS • CROSS SECTION BARYON RESONANCE WITH Y=2 AT 2000 MEV

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = .99 +- .02 GEV/C.

(PRIVATE COMMUNICATION FROM CARROLL, ET AL., DECEMBER, 1968.)

COS(THETA)	D-SIGMA/D-OMEGA UB/SR [1]	U (GEV/C)**2	D-SIGMA/D-U UB/(GEV/C)**2 [2]
-1.00 [3]	535. +- 138.	.1254 [3]	6210. +- 1500.
-.99	490. 119.	.1209	5687. 1376.
-.97	715. 143.	.1131	8290. 1658.
-.95	642. 151.	.0993	7453. 1757.
-.93	560. 155.	.0884	6497. 1802.
-.91	604. 174.	.0776	7010. 2024.
-.89	315. 129.	.0558	3658. 1493.
-.87	526. 175.	.0559	6103. 2034.
-.85	375. 153.	.0451	4352. 1777.
-.83	438. 165.	.0343	5076. 1919.
-.81	290. 134.	.0235	3465. 1550.
-.79	355. 145.	.0124	4122. 1683.
-.77	494. 174.	.0018	5726. 2024.
-.75	456. 173.	-.0090	5295. 2001.
-.73	644. 215.	-.0199	7473. 2491.
-.71	430. 193.	-.0307	4994. 2233.

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.
U IS THE MOMENTUM TRANSFER BETWEEN THE [INCOMING K⁺] AND THE [OUTGOING K⁺].

- [1] PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7 PER CENT.
- [2] PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7 PER CENT.
- [3] EXTRAPOLATED POINT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = 1.04 +- .02 GEV/C.

(PRIVATE COMMUNICATION FROM CARROLL, ET AL., DECEMBER, 1968.)

COS(THETA)	D-SIGMA/D-OMEGA UB/SR [1]	U (GEV/C)**2	D-SIGMA/D-U UB/(GEV/C)**2 [2]
-1.000 [3]	517. +- 54.	.1231 [3]	5910. +- 580.
-.995	487. 58.	.1144	5557. 647.
-.985	516. 60.	.1086	5987. 855.
-.975	555. 79.	.1027	5424. 880.
-.965	503. 82.	.0959	3704. 726.
-.955	343. 67.	.0911	5651. 999.
-.945	538. 93.	.0824	6325. 831.
-.930	586. 77.	.0737	5603. 845.
-.910	519. 78.	.0591	6551. 935.
-.890	607. 87.	.0474	4293. 795.
-.870	397. 74.	.0350	5713. 952.
-.850	529. 88.	.0241	4588. 852.
-.830	425. 79.	.0125	6386. 985.
-.810	592. 91.	.0009	5261. 889.
-.790	487. 82.	-.0108	5780. 950.
-.770	536. 88.	-.0224	6281. 1019.
-.750	582. 94.	-.0341	6154. 1055.
-.730	570. 98.	-.0457	4357. 974.
-.710	404. 90.	-.0574	5773. 1260.
-.690	535. 117.	-.0590	4059. 1224.
-.670	376. 113.		

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.
U IS THE MOMENTUM TRANSFER BETWEEN THE [INCOMING K⁺] AND THE [OUTGOING K⁺].

- [1] PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7 PER CENT.
- [2] PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7 PER CENT.
- [3] EXTRAPOLATED POINT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = 1.11 ± .02 GEV/C.

(PRIVATE COMMUNICATION FROM CARROL, ET AL., DECEMBER, 1968.)

COS(THETA)	D-SIGMA/D-OMEGA UB/SR (1)	U (GEV/C)**2	D-SIGMA/D-U UB/(GEV/C)**2 (2)
-1.000 (3)	404. ± 70.	.1192 (3)	4030. ± 695.
-.995	407. 47.	.1150	4028. 462.
-.985	412. 48.	.1096	4081. 478.
-.975	438. 57.	.1033	4340. 560.
-.965	452. 65.	.0959	4469. 638.
-.955	438. 70.	.0906	4335. 694.
-.945	580. 89.	.0863	5766. 876.
-.930	504. 60.	.0747	4985. 592.
-.910	406. 57.	.0620	4022. 563.
-.890	449. 62.	.0493	4444. 616.
-.870	338. 56.	.0366	3347. 550.
-.850	504. 76.	.0239	4987. 749.
-.830	336. 57.	.0113	3324. 562.
-.810	363. 59.	-.0014	3595. 583.
-.790	396. 61.	-.0141	3918. 605.
-.770	255. 78.	-.0268	2519. 776.
-.750	362. 63.	-.0395	3584. 824.
-.730	468. 77.	-.0522	4631. 761.
-.710	535. 92.	-.0649	5293. 708.
-.690	401. 40.	-.0776	3967. 887.

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.U IS THE MOMENTUM TRANSFER BETWEEN THE [INCOMING K⁺] AND THE [OUTGOING K⁺].

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF ± 7 PER CENT.

(2) PLUS POSSIBLE SYSTEMATIC ERROR OF ± 7 PER CENT.

(3) EXTRAPOLATED POINT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = 1.17 ± .02 GEV/C.

(PRIVATE COMMUNICATION FROM CARROL, ET AL., DECEMBER, 1968.)

COS(THETA)	D-SIGMA/D-OMEGA UB/SR (1)	U (GEV/C)**2	D-SIGMA/D-U UB/(GEV/C)**2 (2)
-1.000 (3)	477. ± 44.	.1154 (3)	4350. ± 402.
-.995	489. 50.	.1119	4454. 460.
-.985	510. 54.	.1050	4650. 490.
-.975	388. 52.	.0981	3537. 477.
-.965	444. 63.	.0913	4049. 578.
-.955	390. 66.	.0844	3610. 605.
-.945	571. 91.	.0775	5234. 833.
-.930	454. 51.	.0671	4143. 466.
-.910	408. 51.	.0534	3723. 465.
-.890	551. 62.	.0396	5029. 566.
-.870	489. 62.	.0258	4457. 566.
-.850	420. 58.	.0120	3832. 531.
-.830	365. 54.	-.0017	3327. 496.
-.810	356. 54.	-.0155	3248. 490.
-.790	354. 53.	-.0293	3229. 487.
-.770	396. 57.	-.0431	3639. 516.
-.750	330. 55.	-.0568	3010. 502.
-.730	328. 59.	-.0706	2991. 537.
-.710	413. 75.	-.0844	3766. 688.
-.690	591. 106.	-.0982	5390. 968.
-.670	402. 107.	-.1120	3656. 979.

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.U IS THE MOMENTUM TRANSFER BETWEEN THE [INCOMING K⁺] AND THE [OUTGOING K⁺].

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF ± 7 PER CENT.

(2) PLUS POSSIBLE SYSTEMATIC ERROR OF ± 7 PER CENT.

(3) EXTRAPOLATED POINT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = 1.20 ± .02 GEV/C.

(PRIVATE COMMUNICATION FROM CARROL, ET AL., DECEMBER, 1968.)

COS(THETA)	D-SIGMA/D-OMEGA UB/SR (1)	U (GEV/C)**2	D-SIGMA/D-U UB/(GEV/C)**2 (2)
-1.000 (3)	400. ± 38.	.1144 (3)	3550. ± 338.
-.995	334. 37.	.1101	2976. 330.
-.985	412. 41.	.1030	3652. 369.
-.975	375. 48.	.0958	3333. 429.
-.965	340. 50.	.0887	3027. 443.
-.955	324. 51.	.0816	2883. 453.
-.945	237. 46.	.0744	2097. 410.
-.930	340. 40.	.0637	3008. 351.
-.910	319. 40.	.0494	2833. 358.
-.890	230. 36.	.0351	2046. 318.
-.870	356. 47.	.0209	3165. 421.
-.850	323. 46.	.0056	2855. 406.
-.830	207. 37.	-.0077	1826. 324.
-.810	250. 40.	-.0220	2213. 357.
-.790	221. 38.	-.0353	1958. 334.
-.770	232. 39.	-.0505	2061. 344.
-.750	272. 45.	-.0648	2411. 396.
-.730	177. 40.	-.0791	1713. 358.
-.710	169. 43.	-.0934	1497. 381.
-.690	257. 63.	-.1077	2285. 555.
-.670	121. 51.	-.1219	1074. 450.

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.U IS THE MOMENTUM TRANSFER BETWEEN THE [INCOMING K⁺] AND THE [OUTGOING K⁺].

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF ± 7 PER CENT.

(2) PLUS POSSIBLE SYSTEMATIC ERROR OF ± 7 PER CENT.

(3) EXTRAPOLATED POINT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K+ PROTON.

LABORATORY BEAM MOMENTUM = 1.30 +- .02 GEV/C.

(PRIVATE COMMUNICATION FROM CARROL, ET AL., DECEMBER, 1968.)

COS(THETA)	D-SIGMA/D-OMEGA UB/SR (1)	U (GEV/C)**2	D-SIGMA/D-U UB/(GEV/C)**2 (2)
-1.000 (3)	247. +- 22.	-.1086 (3)	1950. +- 175.
-.995	225. 22.	-.1046	1781. 175.
-.985	230. 24.	-.0956	1814. 191.
-.975	239. 29.	-.0887	1887. 230.
-.965	195. 27.	-.0807	1539. 213.
-.955	212. 31.	-.0728	1679. 245.
-.945	133. 28.	-.0648	1052. 219.
-.930	223. 25.	-.0529	1755. 207.
-.910	225. 28.	-.0370	1781. 219.
-.890	218. 29.	-.0211	1722. 228.
-.870	199. 29.	-.0052	1575. 227.
-.850	186. 28.	-.0107	1474. 225.
-.830	162. 27.	-.0256	1276. 213.
-.810	190. 29.	-.0425	1498. 228.
-.790	130. 23.	-.0584	1028. 182.
-.770	183. 29.	-.0743	1449. 229.
-.750	211. 34.	-.0902	1671. 268.
-.730	125. 29.	-.1051	991. 227.
-.710	171. 38.	-.1220	1347. 301.
-.690	266. 53.	-.1379	2105. 421.

THETA IS THE ANGLE THAT THE K+ MAKES WITH THE BEAM IN THE GRAND C.M.

U IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K+) AND THE (OUTGOING K+).

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7 PER CENT.

(2) PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7 PER CENT.

(3) EXTRAPOLATED POINT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K+ PROTON.

LABORATORY BEAM MOMENTUM = 1.40 +- .02 GEV/C.

(PRIVATE COMMUNICATION FROM CARROL, ET AL., DECEMBER, 1968.)

COS(THETA)	D-SIGMA/D-OMEGA UB/SR (1)	U (GEV/C)**2	D-SIGMA/D-U UB/(GEV/C)**2 (2)
-1.000 (3)	221. +- 19.	-.1038 (3)	1650. +- 128.
-.995	232. 19.	-.0994	1591. 155.
-.985	235. 21.	-.0936	1678. 151.
-.975	216. 23.	-.0818	1547. 166.
-.965	266. 29.	-.0730	1899. 205.
-.955	208. 26.	-.0642	1487. 188.
-.945	210. 28.	-.0554	1502. 203.
-.930	190. 19.	-.0422	1357. 138.
-.910	197. 21.	-.0247	1406. 151.
-.890	168. 21.	-.0071	1199. 147.
-.870	199. 23.	-.0135	1423. 164.
-.850	198. 24.	-.0281	1418. 172.
-.830	122. 19.	-.0457	870. 136.
-.810	142. 20.	-.0633	1013. 145.
-.790	138. 20.	-.0808	983. 143.
-.770	157. 23.	-.0984	1124. 165.
-.750	231. 30.	-.1160	1654. 215.
-.730	171. 28.	-.1336	935. 181.
-.710	136. 30.	-.1512	708. 180.
-.690	102. 30.	-.1688	728. 216.

THETA IS THE ANGLE THAT THE K+ MAKES WITH THE BEAM IN THE GRAND C.M.

U IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K+) AND THE (OUTGOING K+).

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7 PER CENT.

(2) PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7 PER CENT.

(3) EXTRAPOLATED POINT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K+ PROTON.

LABORATORY BEAM MOMENTUM = 1.49 +- .02 GEV/C.

(PRIVATE COMMUNICATION FROM CARROL, ET AL., DECEMBER, 1968.)

COS(THETA)	D-SIGMA/D-OMEGA UB/SR (1)	U (GEV/C)**2	D-SIGMA/D-U UB/(GEV/C)**2 (2)
-1.000 (3)	223. +- 25.	-.0995 (3)	1460. +- 164.
-.995	207. 26.	-.0947	1357. 172.
-.985	234. 30.	-.0851	1530. 194.
-.975	211. 33.	-.0755	1380. 215.
-.965	181. 34.	-.0659	1182. 219.
-.955	186. 36.	-.0563	1217. 234.
-.945	201. 39.	-.0467	1312. 257.
-.930	147. 27.	-.0323	963. 176.
-.910	147. 27.	-.0131	962. 189.
-.890	139. 30.	-.0051	911. 194.
-.870	181. 35.	-.0295	1185. 228.
-.850	110. 28.	-.0445	720. 186.
-.830	86. 26.	-.0638	560. 189.
-.810	138. 32.	-.0830	903. 207.
-.790	142. 33.	-.1022	929. 213.
-.770	113. 31.	-.1214	737. 204.
-.750	112. 35.	-.1406	731. 231.
-.730	186. 42.	-.1598	1218. 338.
-.710	115. 47.	-.1790	755. 308.

THETA IS THE ANGLE THAT THE K+ MAKES WITH THE BEAM IN THE GRAND C.M.

U IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K+) AND THE (OUTGOING K+).

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7 PER CENT.

(2) PLUS POSSIBLE SYSTEMATIC ERROR OF +- 7 PER CENT.

(3) EXTRAPOLATED POINT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = 1.61 ± .02 GEV/C.

(PRIVATE COMMUNICATION FROM CARROL, ET AL., DECEMBER, 1968.)

COS(THETA)	D-SIGMA/D-OMEGA UB/SR [1]	U (GEV/C)**2	D-SIGMA/D-U UB/(GEV/C)**2 [2]
-1.000 [3]	141. ± 8.	.0945 [3]	829. ± 50.
-.995	128. 8.	.0892	752. 48.
-.985	144. 10.	.0786	852. 57.
-.975	111. 9.	.0679	657. 56.
-.965	112. 11.	.0573	658. 65.
-.955	117. 11.	.0456	690. 68.
-.945	87. 10.	.0359	911. 59.
-.930	119. 9.	.0270	704. 55.
-.910	121. 10.	-.0013	716. 59.
-.890	89. 9.	-.0226	522. 53.
-.870	91. 10.	-.0439	539. 60.
-.850	93. 10.	-.0652	546. 58.
-.830	87. 10.	-.0855	516. 62.
-.810	95. 10.	-.1079	558. 61.
-.790	88. 10.	-.1292	516. 59.
-.770	88. 12.	-.1505	518. 73.
-.750	94. 13.	-.1718	552. 77.
-.730	84. 14.	-.1931	493. 82.
-.710	91. 17.	-.2144	535. 103.

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.
U IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K⁺) AND THE (OUTGOING K⁺).

[1] PLUS POSSIBLE SYSTEMATIC ERROR OF ± 7 PER CENT.
[2] PLUS POSSIBLE SYSTEMATIC ERROR OF ± 7 PER CENT.
[3] EXTRAPOLATED POINT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = 1.67 ± .02 GEV/C.

(PRIVATE COMMUNICATION FROM CARROL, ET AL., DECEMBER, 1968.)

COS(THETA)	D-SIGMA/D-OMEGA UB/SR [1]	U (GEV/C)**2	D-SIGMA/D-U UB/(GEV/C)**2 [2]
-1.000 [3]	108. ± 8.	.0924 [3]	608. ± 45.
-.995	111. 9.	.0870	625. 49.
-.985	105. 9.	.0759	595. 52.
-.975	99. 11.	.0647	558. 60.
-.965	114. 12.	.0536	644. 70.
-.955	126. 14.	.0425	709. 78.
-.945	97. 12.	.0314	549. 69.
-.930	96. 9.	.0148	542. 53.
-.910	77. 9.	-.0075	435. 51.
-.890	76. 11.	-.0297	431. 63.
-.870	74. 10.	-.0519	420. 57.
-.850	86. 11.	-.0741	486. 61.
-.830	94. 12.	-.0953	531. 66.
-.810	102. 13.	-.1186	578. 71.
-.790	58. 11.	-.1408	325. 62.
-.770	44. 9.	-.1630	248. 51.
-.750	47. 11.	-.1852	267. 61.
-.730	86. 17.	-.2074	486. 95.

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.
U IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K⁺) AND THE (OUTGOING K⁺).

[1] PLUS POSSIBLE SYSTEMATIC ERROR OF ± 7 PER CENT.
[2] PLUS POSSIBLE SYSTEMATIC ERROR OF ± 7 PER CENT.
[3] EXTRAPOLATED POINT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = 1.71 ± .02 GEV/C.

(PRIVATE COMMUNICATION FROM CARROL, ET AL., DECEMBER, 1968.)

COS(THETA)	D-SIGMA/D-OMEGA UB/SR [1]	U (GEV/C)**2	D-SIGMA/D-U UB/(GEV/C)**2 [2]
-1.000 [3]	90. ± 11.	.0926 [3]	491. ± 60.
-.995	95. 12.	.0848	517. 67.
-.985	83. 12.	.0732	450. 65.
-.975	78. 14.	.0617	422. 75.
-.965	85. 15.	.0521	449. 85.
-.955	96. 18.	.0385	521. 97.
-.945	79. 17.	.0270	430. 92.
-.930	103. 14.	.0096	558. 77.
-.910	65. 12.	-.0135	353. 65.
-.890	94. 15.	-.0357	510. 84.
-.870	72. 14.	-.0598	389. 75.
-.850	66. 14.	-.0829	353. 75.
-.830	59. 15.	-.1051	320. 83.
-.810	75. 16.	-.1292	406. 85.
-.790	30. 10.	-.1523	161. 54.
-.770	51. 14.	-.1755	277. 77.
-.750	46. 15.	-.1986	252. 84.
-.730	15. 9.	-.2217	70. 49.
-.710	30. 17.	-.2449	163. 94.

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.
U IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K⁺) AND THE (OUTGOING K⁺).

[1] PLUS POSSIBLE SYSTEMATIC ERROR OF ± 7 PER CENT.
[2] PLUS POSSIBLE SYSTEMATIC ERROR OF ± 7 PER CENT.
[3] EXTRAPOLATED POINT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = 1.79 ± .02 GEV/C.

(PRIVATE COMMUNICATION FROM CARROL, ET AL., DECEMBER, 1968.)

COS(THETA)	D-SIGMA/D-OMEGA UB/SR [1]	U (GEV/C)**2	D-SIGMA/D-U UB/(GEV/C)**2 [2]
-1.000 [3]	113. ± 6.	.3879 [3]	580. ± 32.
-.995	114. 6.	.0820	585. 32.
-.985	99. 6.	.0699	511. 31.
-.975	106. 7.	.0577	547. 38.
-.965	103. 8.	.0455	528. 41.
-.955	89. 8.	.0333	460. 40.
-.945	94. 8.	.3211	482. 43.
-.930	89. 6.	.0029	458. 31.
-.910	85. 6.	-.0215	439. 32.
-.890	88. 7.	-.0458	451. 37.
-.870	57. 7.	-.0702	295. 37.
-.850	69. 6.	-.0945	254. 34.
-.830	56. 8.	-.1189	288. 42.
-.810	67. 8.	-.1433	347. 44.
-.790	55. 7.	-.1676	284. 34.
-.770	42. 7.	-.1920	218. 36.
-.750	58. 10.	-.2163	299. 51.
-.730	44. 9.	-.2407	225. 44.

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.
U IS THE MOMENTUM TRANSFER BETWEEN THE [INCOMING K⁺] AND THE [OUTGOING K⁺].

- [1] PLUS POSSIBLE SYSTEMATIC ERROR OF ± 7 PER CENT.
[2] PLUS POSSIBLE SYSTEMATIC ERROR OF ± 7 PER CENT.
[3] EXTRAPOLATED POINT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = 1.82 ± .02 GEV/C.

(PRIVATE COMMUNICATION FROM CARROL, ET AL., DECEMBER, 1968.)

COS(THETA)	D-SIGMA/D-OMEGA UB/SR [1]	U (GEV/C)**2	D-SIGMA/D-U UB/(GEV/C)**2 [2]
-1.000 [3]	84. ± 12.	.0858 [3]	420. ± 60.
-.995	100. 14.	.0805	500. 72.
-.985	96. 10.	.0690	281. 52.
-.975	59. 12.	.0554	295. 60.
-.965	95. 20.	.0429	478. 100.
-.955	71. 17.	.0304	356. 86.
-.945	72. 19.	.0178	361. 93.
-.930	58. 13.	-.0010	289. 63.
-.910	60. 14.	-.0250	302. 69.
-.890	68. 16.	-.0511	340. 78.
-.870	44. 13.	-.0752	222. 64.
-.850	23. 19.	-.1012	467. 97.
-.830	39. 13.	-.1263	195. 65.
-.810	61. 17.	-.1514	304. 84.
-.790	81. 23.	-.1764	406. 101.
-.770	11. 8.	-.2015	56. 39.
-.750	40. 18.	-.2266	202. 90.
-.730	46. 23.	-.2516	229. 114.

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.
U IS THE MOMENTUM TRANSFER BETWEEN THE [INCOMING K⁺] AND THE [OUTGOING K⁺].

- [1] PLUS POSSIBLE SYSTEMATIC ERROR OF ± 7 PER CENT.
[2] PLUS POSSIBLE SYSTEMATIC ERROR OF ± 7 PER CENT.
[3] EXTRAPOLATED POINT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = 1.88 ± .02 GEV/C.

(PRIVATE COMMUNICATION FROM CARROL, ET AL., DECEMBER, 1968.)

COS(THETA)	D-SIGMA/D-OMEGA UB/SR [1]	U (GEV/C)**2	D-SIGMA/D-U UB/(GEV/C)**2 [2]
-1.000 [3]	104. ± 7.	.0849 [3]	498. ± 36.
-.995	98. 7.	.0784	475. 36.
-.985	75. 9.	.0653	455. 38.
-.975	99. 9.	.0523	479. 41.
-.965	74. 8.	.0392	358. 40.
-.955	73. 9.	.0262	351. 43.
-.945	78. 10.	.0132	375. 46.
-.930	72. 8.	-.0004	348. 37.
-.910	71. 8.	-.0325	343. 40.
-.890	56. 8.	-.0586	271. 37.
-.870	64. 11.	-.0846	309. 52.
-.850	71. 12.	-.1107	341. 59.
-.830	50. 8.	-.1358	239. 39.
-.810	36. 7.	-.1629	176. 34.
-.790	45. 11.	-.1890	219. 53.
-.770	40. 9.	-.2151	194. 44.
-.750	42. 11.	-.2411	204. 93.
-.730	24. 10.	-.2672	117. 48.

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.
U IS THE MOMENTUM TRANSFER BETWEEN THE [INCOMING K⁺] AND THE [OUTGOING K⁺].

- [1] PLUS POSSIBLE SYSTEMATIC ERROR OF ± 7 PER CENT.
[2] PLUS POSSIBLE SYSTEMATIC ERROR OF ± 7 PER CENT.
[3] EXTRAPOLATED POINT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.LABORATORY BEAM MOMENTUM = 2.00 $\pm$.02 GEV/C.

(PRIVATE COMMUNICATION FROM CARROL, ET AL., DECEMBER, 1968.)

COS(THETA)	D-SIGMA/D-OMEGA UB/SR (1)	U (GEV/C)**2	D-SIGMA/D-U UB/(GEV/C)**2 (2)
-1.000 (3)	93. $\pm$ 5.	.0810 (3)	405. $\pm$ 21.
-.995	84. 5.	.0720	366. 21.
-.985	89. 5.	.0573	389. 22.
-.975	79. 5.	.0426	342. 23.
-.965	82. 6.	.0279	354. 26.
-.955	73. 6.	.0132	317. 27.
-.945	54. 5.	-.0015	236. 23.
-.930	50. 4.	-.0235	216. 17.
-.910	54. 4.	-.0329	237. 19.
-.890	52. 4.	-.0822	226. 19.
-.870	38. 5.	-.1116	157. 22.
-.850	36. 4.	-.1410	157. 18.
-.830	33. 4.	-.1704	142. 18.
-.810	25. 4.	-.1997	107. 16.
-.790	23. 4.	-.2291	101. 18.
-.770	27. 5.	-.2585	118. 21.
-.750	12. 4.	-.2879	52. 17.

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.
U IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K⁺) AND THE (OUTGOING K⁺).

- (1) PLUS POSSIBLE SYSTEMATIC ERROR OF $\pm$ 7 PER CENT.
(2) PLUS POSSIBLE SYSTEMATIC ERROR OF $\pm$ 7 PER CENT.
(3) EXTRAPOLATED POINT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.LABORATORY BEAM MOMENTUM = 2.20 $\pm$.02 GEV/C.

(PRIVATE COMMUNICATION FROM CARROL, ET AL., DECEMBER, 1968.)

COS(THETA)	D-SIGMA/D-OMEGA UB/SR (1)	U (GEV/C)**2	D-SIGMA/D-U UB/(GEV/C)**2 (2)
-1.000 (3)	95. $\pm$ 6.	.0757 (3)	373. $\pm$ 23.
-.995	92. 6.	.0677	362. 23.
-.985	77. 5.	.0518	304. 21.
-.975	75. 6.	.0359	290. 24.
-.965	67. 6.	.0201	266. 25.
-.955	61. 6.	.0042	241. 25.
-.945	49. 6.	-.0117	194. 23.
-.930	52. 4.	-.0336	205. 17.
-.910	50. 6.	-.0674	196. 22.
-.890	32. 4.	-.0992	126. 15.
-.870	29. 7.	-.1309	113. 29.
-.850	23. 7.	-.1627	91. 26.
-.830	21. 5.	-.1945	83. 21.
-.810	24. 5.	-.2263	95. 20.
-.790	16. 4.	-.2581	63. 16.
-.770	8. 3.	-.2899	32. 12.

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.
U IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K⁺) AND THE (OUTGOING K⁺).

- (1) PLUS POSSIBLE SYSTEMATIC ERROR OF $\pm$ 7 PER CENT.
(2) PLUS POSSIBLE SYSTEMATIC ERROR OF $\pm$ 7 PER CENT.
(3) EXTRAPOLATED POINT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.LABORATORY BEAM MOMENTUM = 2.33 $\pm$.02 GEV/C.

(PRIVATE COMMUNICATION FROM CARROL, ET AL., DECEMBER, 1968.)

COS(THETA)	D-SIGMA/D-OMEGA UB/SR (1)	U (GEV/C)**2	D-SIGMA/D-U UB/(GEV/C)**2 (2)
-1.000 (3)	96. $\pm$ 4.	.0724 (3)	353. $\pm$ 16.
-.995	90. 4.	.0642	331. 16.
-.985	82. 4.	.0472	302. 16.
-.975	73. 5.	.0302	271. 17.
-.965	70. 5.	.0133	257. 18.
-.955	53. 4.	-.0037	194. 16.
-.945	53. 5.	-.0227	196. 18.
-.930	47. 4.	-.0452	174. 15.
-.910	37. 5.	-.0801	136. 18.
-.890	33. 4.	-.1141	122. 15.
-.870	33. 5.	-.1480	124. 18.
-.850	23. 4.	-.1820	86. 14.
-.830	25. 4.	-.2159	91. 16.
-.810	19. 3.	-.2499	69. 13.
-.790	10. 3.	-.2839	36. 10.
-.770	11. 3.	-.3178	40. 12.

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.
U IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K⁺) AND THE (OUTGOING K⁺).

- (1) PLUS POSSIBLE SYSTEMATIC ERROR OF $\pm$ 7 PER CENT.
(2) PLUS POSSIBLE SYSTEMATIC ERROR OF $\pm$ 7 PER CENT.
(3) EXTRAPOLATED POINT.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = 2.45 ± .02 GEV/C.

(PRIVATE COMMUNICATION FROM CARROL, ET AL., DECEMBER, 1968.)

COS(THETA)	D-SIGMA/D-OMEGA UB/SR [1]	U (GEV/C)**2	D-SIGMA/D-U UB/(GEV/C)**2 [2]
-1.00 [3]	89. ± 10.	.0696 [3]	339. ± 35.
-.99	73. 5.	.0515	251. 22.
-.97	67. 7.	.0151	231. 24.
-.95	42. 5.	-.0213	145. 19.
-.93	26. 5.	-.0577	89. 18.
-.91	30. 5.	-.0940	102. 18.
-.89	29. 7.	-.1304	100. 23.
-.87	20. 8.	-.1668	68. 19.
-.85	22. 5.	-.2031	76. 20.
-.83	7. 2.	-.2395	27. 10.
-.81	20. 6.	-.2759	68. 19.
-.79	8. 3.	-.3123	27. 13.
-.77	12. 5.	-.3486	44. 20.

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.
U IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K⁺) AND THE (OUTGOING K⁺).

[1] PLUS POSSIBLE SYSTEMATIC ERROR OF ± 7 PER CENT.
[2] PLUS POSSIBLE SYSTEMATIC ERROR OF ± 7 PER CENT.
[3] EXTRAPOLATED POINT.

62 STUDY OF THE REACTION K⁺ P → K⁰ P⁺ P AT 1.14 GEV/C. [PHYS. REV. 133, B220 (1964)]

E. BOLDT, J. DUBOC, N. H. DUONG, P. EBERHARD, R. GEORGE, V. P. HENRI, F. LEVY, J. POYEN, M. PRIPSTEIN (COLLEGE DE FRANCE, PARIS, FRANCE)
J. FRIESSBACH, B. TRON TIENTS, N. P. THOMPSON, N. G. W. R. GIP-SIR-YVETTE, FRANCE

ABSTRACT WE REPORT HERE THE STUDY OF INELASTIC PROCESS K⁺ P → K⁰ P⁺ P AT 1.14 GEV/C (TOTAL CENTER OF MASS ENERGY = 1.859 GEV). THE STUDY WAS DONE USING THE SACLAY 35-CM HYDROGEN BUBBLE CHAMBER AND ANALYZING EVENTS HAVING A TWO-PRONG + V₀ TOPOLOGY IN THE CHAMBER. THE TOTAL CROSS SECTION FOR THE REACTION IS 4.6 ± 0.3 MB. THE REACTION IS DOMINATED BY THE PRODUCTION OF THE N*(13,3) AND K*(890) RESONANCES, THE NONRESONATING BACKGROUND, IF ANY, CONTRIBUTES LESS THAN 0.4 MB TO THE TOTAL CROSS SECTION. NO EVENTS WERE FOUND WITH TWO PIONS IN THE FINAL STATE. THE CROSS SECTION FOR N*(13,3) PRODUCTION IS 3.6 ± 0.5 MB. THE K⁺ PRODUCTION CROSS SECTION (APART FROM POSSIBLE INTERFERENCE CONTRIBUTIONS) IS 0.9 ± 0.3 MB. OUR MEASURED VALUE FOR THE MASS OF THE N(13,3) RESONANCE IS 1212 ± 8 MEV WITH A FULL WIDTH OF 72 ± 13 MEV. A SYSTEMATIC ANALYSIS OF THE PRODUCTION AND DECAY PROPERTIES OF THE N*(13,3) WAS MADE. THE N*(13,3) IS PRODUCED WITH A PRONOUNCED SPIN ALIGNMENT. OUR DATA ARE CONSISTENT WITH A COMPLETE ALIGNMENT OF ± 1/2 WITH RESPECT TO THE PRODUCTION PLANE NORMAL OR WITH A ALIGNMENT OF ± 3/2 WITH RESPECT TO SOME DIRECTION IN THE PRODUCTION PLANE, THE MOST PROBABLE DIRECTION BEING THE DIRECTION OF THE INCIDENT K⁺. NO EVIDENCE IS SEEN FOR THE EXISTENCE OF THE K* (730 MEV) RESONANCE.

CROSS REFERENCES
THIS ARTICLE SUPERSEDES PHYS. LETTERS 6, 233 (1963).

ADDITIONAL FOOTNOTE REFERENCES
PHYS. LETTERS 6, 233 (1963), UCL 8030 REV. (1961), COMPTES RENDUS. ACAD. SCI. 255, 890 (1962), PHYS. REV. 128, 1868 (1962), REV. MOD. PHYS. 25, 314 (1963), PHYS. REV. LETTERS 10, 351 (1963), JINR 584 (1963), CERN CONFERENCE 703, AND PHYS. LETTERS 6, 273 (1963).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 3/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K⁺ ON PROTON AT 1.14 GEV/C.

THIS EXPERIMENT USES THE SACLAY 35 CM BUBBLE CHAMBER.

KEY WORDS = CROSS SECTION DALITZ PLOT DELTA(1238) MASS LIFETIME ANGULAR DISTRIBUTION K*(890)
KAPPA MESON MASS SPECTRUM DENSITY MATRIX

COMPOUND KEY WORDS = DELTA(1238) MASS DELTA(1238) LIFETIME

CROSS SECTION FOR K⁺ PROTON → K⁰ P⁺ PROTON:

LABORATORY
BEAM MOMENTUM
GEV/C
1.140 ± .027
MILLI-BARNS
4.6 ± .3

CROSS SECTION FOR K⁺ PROTON → DELTA++ K⁰.
DELTA++ PROTON P⁺ [1]

LABORATORY
BEAM MOMENTUM
GEV/C
1.140 ± .027
MILLI-BARNS
3.6 ± .5

[1] FITTED FOR MASS AND/OR WIDTH (MASS = 1.212 GEV; WIDTH = .072 GEV), AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.

CROSS SECTION FOR K⁺ PROTON → K*(890)+ PROTON.
K*(890)+ → K⁰ P⁺ [1]

LABORATORY
BEAM MOMENTUM
GEV/C
1.140 ± .027
MILLI-BARNS
.9 ± .3

[1] COUNTED ONLY EVENTS ABOVE BACKGROUND (MASS = .888 GEV; WIDTH = .350 GEV).

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow \Delta^{++} K^0$
 $\Delta^{++} \rightarrow \text{PROTON } \pi^+$ [1]
 LABORATORY BEAM MOMENTUM = $1.140 \pm .027 \text{ GEV/C.}$

.....
 . THIS DATA WAS READ FROM A GRAPH .

COS(THETA)		D-SIGMA/D-OMEGA MB/SR	
MIN	MAX		
.8	1.0	.068 $\pm$.040	
.6	.8	.424 .140	
.4	.6	.580 .140	
.2	.4	.240 .080	
.0	.2	.320 .100	
-.2	.0	.288 .100	
-.4	-.2	.204 .080	
-.6	-.4	.300 .100	
-.8	-.6	.144 .060	
-1.0	-.8	.080 .040	

THETA IS THE ANGLE THAT THE K^0 MAKES WITH THE BEAM IN THE GRAND C.M.

[1] COUNTED ALL EVENTS IN MASS BAND [MASS = 1.215 GEV ; WIDTH = $.110 \text{ GEV}$].

63 THE $K^+ - \text{PROTON}$ INTERACTION AT 1.45 GEV/C. [PHYS. LETTERS 16, 83 (1965)]

A.BETTINI, M.CRESTI, S.LIMENTANI, L.PERRUZZO, R.SANTANGELO (UNIV. OF PADOVA, PADOVA, ITALY)
 D.LOCKE, O.J. CRENNELL, M.T.DAVIES, P.B.JONES (OXFORD UNIV., OXFORD, ENGLAND)

CROSS REFERENCES
 SOME OF THIS DATA RENORMALIZED IN UCRL 18758 (1968).

ADDITIONAL FOOTNOTE REFERENCES
 PHYS. LETTERS 9, 359 (1964), PHYS. LETTERS 6, 309 (1963), UCRL 8030 (1964), PHYS. REV. LETTERS 11, 90 (1963), PHYS. LETTERS 6, 273 (1963), PHYS. REV. LETTERS 11, 93 (1963), PHYS. REV. 133, 8220 (1964), PHYS. REV. 134, 81099 (1964), NUOVO CIMENTO 33, 905 (1964), NUOVO CIMENTO 33, 309 (1964), PHYS. REV. LETTERS 12, 627 (1964), AND NUOVO CIMENTO 34, 735 (1964).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 1/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K^+ ON PROTON AT 1.455 GEV/C.

THIS EXPERIMENT USES THE CERN 32 CM (H) BUBBLE CHAMBER.

GENERAL COMMENTS ON THIS ARTICLE
 1 K^+ LIFETIME TAKEN AS 1.23 ± 0.01 ($\times 10^{-8} \text{ SEC}$)

KEY WORDS = $K^+(890)$ CROSS SECTION $\Delta^{++}(1238)$ ONE PARTICLE EXCHANGE ANGULAR DISTRIBUTION

$K^+ \text{ PROTON ELASTIC CROSS SECTION.}$

ORIGINAL DATA			RENORMALIZED VALUE [1]
LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS	MILLI-BARNS
$1.455 \pm .025$	$9.7 \pm .5$	2559	$9.6 \pm .3$

[1] BLAND, ET. AL., UCRL 18758 (1969).

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \text{ PROTON } \pi^+$.

ORIGINAL DATA			RENORMALIZED VALUE [1]
LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS		MILLI-BARNS
$1.455 \pm .025$	$4.95 \pm .30$		$4.9 \pm .3$

[1] BLAND, ET. AL., UCRL 18758 (1969).

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \text{ PROTON } \pi^0$.

ORIGINAL DATA			RENORMALIZED VALUE [1]
LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS		MILLI-BARNS
$1.455 \pm .025$	$1.95 \pm .20$		$1.9 \pm .2$

[1] BLAND, ET. AL., UCRL 18758 (1969).

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \text{ NEUTRON } \pi^+$.

ORIGINAL DATA			RENORMALIZED VALUE [1]
LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS		MILLI-BARNS
$1.455 \pm .025$	$1.35 \pm .10$		$1.35 \pm .10$

[1] BLAND, ET. AL., UCRL 18758 (1969).

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K \text{ MESON NUCLEON PION.}$

ORIGINAL DATA		RENORMALIZED VALUE [1]
LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	MILLI-BARNS
1.455 $\pm$.025	8.25 $\pm$.37	8.16 $\pm$.31

[1] BLAND, ET AL., UCRL 18758 (1969).

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \text{ PROTON } \pi^+ \pi^-.$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
1.455 $\pm$.025	.15 $\pm$.04

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \text{ PROTON } \pi^+ \pi^0.$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
1.455 $\pm$.025	.08 $\pm$.03

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \text{ NEUTRON } \pi^+ \pi^+.$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
1.455 $\pm$.025	.02 $\pm$.02

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)^+ \text{ PROTON.}$

$K^*(890)^+ \rightarrow K^0 \pi^+ + K^+ \pi^0$ [1]

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
1.455 $\pm$.025	2.4 $\pm$.3

[1] USED SIMPLE MASS CUT [MASS = .900 GEV; WIDTH = .100 GEV].

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow \Delta^{++} K^0.$

$\Delta^{++} \rightarrow \pi^+ \text{ PROTON}$ [1]

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
1.455 $\pm$.025	2.4 $\pm$.5

[1] USED SIMPLE MASS CUT [MASS = 1.220 GEV; WIDTH = .100 GEV].

$K^+ \text{ PROTON TOTAL CROSS SECTION.}$

(IN RENORMALIZING VALUES FROM THIS EXPERIMENT, A VALUE OF 18.00 $\pm$.10 MB HAS BEEN USED FOR THIS.)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
1.455 $\pm$.025	18.2 $\pm$.8

ELASTIC DIFFERENTIAL CROSS SECTION FOR K+ PROTON.

LABORATORY BEAM MOMENTUM = 1.455 +- .025 GEV/C.

ORIGINAL DATA				RENORMALIZED VALUES (1)		
COS(THETA)		D-SIGMA/D-OMEGA UB/SR (2)		NO. EVENTS	D-SIGMA/D-OMEGA UB/SR	
MIN	MAX					
-1.00	-.95	.12	+.04	10	.12	+.04
-.95	-.90	.26	.06	22	.26	.05
-.90	-.85	.16	.04	13	.15	.04
-.85	-.80	.25	.05	21	.25	.05
-.80	-.75	.11	.04	9	.11	.04
-.75	-.70	.16	.04	13	.15	.04
-.70	-.65	.06	.03	5	.05	.03
-.65	-.60	.14	.04	12	.14	.04
-.60	-.55	.16	.04	13	.15	.04
-.55	-.50	.13	.04	11	.13	.04
-.50	-.45	.16	.04	13	.15	.04
-.45	-.40	.14	.04	12	.14	.04
-.40	-.35	.23	.05	19	.23	.05
-.35	-.30	.17	.04	14	.17	.04
-.30	-.25	.12	.04	10	.12	.04
-.25	-.20	.32	.06	27	.32	.05
-.20	-.15	.35	.06	29	.34	.06
-.15	-.10	.26	.06	22	.26	.05
-.10	-.05	.30	.06	25	.30	.05
-.05	.00	.30	.06	25	.30	.05
.00	.05	.31	.06	26	.31	.05
.05	.10	.48	.08	40	.48	.08
.10	.15	.54	.08	45	.53	.08
.15	.20	.48	.08	40	.48	.08
.20	.25	.74	.09	62	.74	.09
.25	.30	.52	.08	43	.51	.08
.30	.35	.60	.08	50	.59	.08
.35	.40	.78	.10	65	.77	.10
.40	.45	.71	.09	59	.70	.09
.45	.50	1.18	.12	98	1.16	.12
.50	.55	.94	.11	78	.93	.10
.55	.60	1.02	.11	85	1.01	.11
.60	.65	1.24	.12	103	1.22	.12
.65	.70	1.43	.13	119	1.41	.13
.70	.75	1.86	.15	155	1.84	.15
.75	.80	2.16	.16	180	2.14	.15
.80	.85	2.32	.17	193	2.29	.17
.85	.90	2.44	.17	203	2.41	.17
.90	.95	3.14	.19	262	3.11	.19
.95	1.00	3.82	.21	318	3.78	.21

THETA IS THE ANGLE THAT THE K+ MAKES WITH THE BEAM IN THE GRAND C.M.

- (1) ORIGINAL VALUES HAVE BEEN MULTIPLIED BY .99 - THE SAME FACTOR THAT THE INTEGRATED CROSS SECTION WAS MULTIPLIED BY TO GET THE RE-NORMALIZED VALUES OF BLAND, ET. AL., UCRL 18758 (1969).
 (2) COUNTS WERE MULTIPLIED BY .012 TO GET THESE. ERRORS ARE TAKEN AS PROPORTIONAL TO THE SQUARE-ROOT OF THE COUNTS.

DIFFERENTIAL CROSS SECTION FOR K+ PROTON - DELTA++ KO.
DELTA++ - PROTON PI+ (1)

LABORATORY BEAM MOMENTUM = 1.455 +- .025 GEV/C.

 . THIS DATA WAS READ FROM A GRAPH .

ORIGINAL DATA				RENORMALIZED VALUES (2)		
COS(THETA)		D-SIGMA/D-OMEGA UB/SR (2)		NO. EVENTS	D-SIGMA/D-OMEGA UB/SR	
MIN	MAX					
.8	1.0	.275	+- .027	102	.274	+- .027
.6	.8	.351	.031	130	.349	.031
.4	.6	.332	.030	123	.330	.030
.2	.4	.229	.025	85	.228	.025
.0	.2	.197	.023	73	.196	.023
-.2	.0	.130	.019	48	.129	.019
-.4	-.2	.138	.019	51	.137	.019
-.6	-.4	.086	.015	32	.086	.015
-.8	-.6	.081	.015	30	.081	.015
-1.0	-.8	.040	.010	15	.040	.010

THETA IS THE ANGLE THAT THE KO MAKES WITH THE BEAM IN THE GRAND C.M.

- (1) USED SIMPLE MASS CUT [MASS = 1.220 GEV; WIDTH = .100 GEV].
 (2) ORIGINAL VALUES HAVE BEEN MULTIPLIED BY .995 - THE SAME FACTOR THAT THE INTEGRATED CROSS SECTION WAS MULTIPLIED BY TO GET THE RE-NORMALIZED VALUES OF BLAND, ET AL., UCRL 18758 (1969).
 (3) COUNTS WERE MULTIPLIED BY .0027 TO GET THESE. ERRORS ARE TAKEN AS PROPORTIONAL TO THE SQUARE-ROOT OF THE COUNTS.

DIFFERENTIAL CROSS SECTION FOR K^+ PROTON $\rightarrow K^*(890)^+ \rightarrow$ PROTON.
 $K^*(890)^+ \rightarrow K^+ \pi^0 \rightarrow K^0 \pi^+ \pi^-$
 LABORATORY BEAM MOMENTUM = 1.455 $\pm$.025 GEV/C.

.....
 . THIS DATA WAS READ FROM A GRAPH .

ORIGINAL DATA			RENORMALIZED VALUES [2]		
COS(THETA)		D-SIGMA/D-OMEGA UB/SR [3]	NO. EVENTS		D-SIGMA/D-OMEGA UB/SR
MIN	MAX				
-1.0	-.8	.083 $\pm$.021	15	.	.082 $\pm$.021
-.8	-.6	.049 .016	9	.	.049 .016
-.6	-.4	.055 .017	10	.	.055 .017
-.4	-.2	.104 .024	19	.	.104 .024
-.2	.0	.181 .032	33	.	.181 .031
.0	.2	.181 .032	33	.	.181 .031
.2	.4	.192 .033	35	.	.192 .032
.4	.6	.264 .038	48	.	.263 .038
.6	.8	.330 .043	60	.	.328 .042
.8	1.0	.473 .051	86	.	.471 .051

THETA IS THE ANGLE THAT THE $K^*(890)^+$ MAKES WITH THE BEAM IN THE GRAND C.M.

[1] USED SIMPLE MASS CUT [MASS = .900 GEV; WIDTH = .100 GEV].
 [2] ORIGINAL VALUES HAVE BEEN MULTIPLIED BY .995 - THE SAME FACTOR THAT THE INTEGRATED CROSS SECTION WAS MULTIPLIED BY TO GET THE RE-NORMALIZED VALUES OF BLAND, ET AL., UCRL 18758 (1969).
 [3] COUNTS WERE MULTIPLIED BY .0055 TO GET THESE. ERRORS ARE TAKEN AS PROPORTIONAL TO THE SQUARE-ROOT OF THE COUNTS.

64 ELASTIC K^+ PROTON SCATTERING AT 970, 1170, AND 1970 MEV/C. [PHYS. REV. 129, 2743 (1963)]

..COOK,D.,KEEFE,L.T.,KERTH,P.G.,MURPHY,W.A.,MENZEL,T.F.,ZIFF (U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA, AND UNIV. OF CALIFORNIA, BERKELEY, CALIF., USA)

ABSTRACT: A LIQUID-HYDROGEN TARGET SURROUNDED BY SPARK CHAMBERS WAS USED TO STUDY K^+ PROTON ELASTIC SCATTERING AT 970, 1170, AND 1970 MEV/C INCIDENT K^+ MOMENTUM. IN CONTRAST WITH RESULTS AT LOW ENERGIES, THE ANGULAR DISTRIBUTIONS WERE NOT ISOTROPIC: THE ANISOTROPY INCREASES WITH ENERGY, WHILE THE ELASTIC CROSS SECTION DECREASES. A NUMBER OF PHASE-SHIFT SOLUTIONS USING (A) $S_{1/2}$, $P_{1/2}$, AND $P_{3/2}$ COMPLEX PHASE SHIFTS; (B) REAL $S_{1/2}$, $P_{1/2}$, AND $P_{3/2}$, AND COMPLEX $D_{3/2}$ PHASE SHIFTS; AND (C) COMPLEX $S_{1/2}$ AND $D_{3/2}$, AND REAL $P_{1/2}$ AND $P_{3/2}$ PHASE SHIFTS HAVE BEEN FOUND FOR THE 970- AND 1170-MEV/C DATA. THE 1970-MEV/C DISTRIBUTION HAS BEEN FITTED BY AN OPTICAL MODEL. THE DATA PRESENTED HERE HAVE BEEN INCLUDED WITH OTHER INFORMATION ON K^+ -NUCLEON SCATTERING IN A TEST OF FORWARD-SCATTERING DISPERSION RELATIONS. THE DATA ARE STILL INSUFFICIENT TO PROVIDE A DEFINITE TEST; HOWEVER, AN ACCEPTABLE FIT WAS FOUND FOR GAMMA (EFFECTIVE POLE RESIDUE) = -0.1 ± 0.3 .

ADDITIONAL FOOTNOTE REFERENCES

NUOVO CIMENTO 12, 241 (1959), PHYS. REV. 118, 553 (1960), PHYS. REV. 107, 279 (1957), PHYS. REV. LETTERS 7, 188 (1961), PHYS. REV. LETTERS 2, 117 (1959), PHYS. REV. LETTERS 5, 333 (1960), DUBNA 0 721 (1961), PHYS. REV. LETTERS 7, 182 (1961), BULL. AM. PHYS. SOC. 6, 433 (1961), PHYS. REV. 123, 695 (1961), BERKELEY CONFERENCE 84 (1960), PHYS. REV. 123, 320 (1961), PHYS. REV. 75, 1352 (1949), PHYS. REV. 110, 569 (1958), PHYS. REV. 110, 572 (1958), PROGRESS OF THEORETICAL PHYSICS 19, 238 (1958), PHYS. REV. LETTERS 2, 510 (1959), PHYS. REV. LETTERS 4, 89 (1960), PHYS. REV. 124, 2049 (1961), PHYS. REV. LETTERS 9, 135 (1962), PHYS. REV. 125, 1696 (1962), PHYS. REV. LETTERS 8, 28 (1962), ANNALS OF PHYSICS 10, 307 (1960), PHYS. REV. 127, 1305 (1962), AND PHYS. REV. LETTERS 8, 142 (1962).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 1/68, AND VERIFIED BY ROGER BLAND.

BEAM IS K^+ ON PROTON FROM .97 TO 1.97 GEV/C.

THIS EXPERIMENT USES SPARK CHAMBERS.

KEY WORDS: CROSS SECTION ANGULAR DISTRIBUTION MODELS PHASE SHIFT

ELASTIC DIFFERENTIAL CROSS SECTION FOR K^+ PROTON.

LABORATORY BEAM MOMENTUM = .97 GEV/C $\pm$.4 PER CENT.

COS(THETA)		D-SIGMA/D-OMEGA MB/SR
.75 $\pm$.06		1.07 $\pm$.18
.60 .10		.84 .09
.40 .10		1.15 .11
.15 .15		.75 .07
-.10 .10		.57 .06
-.30 .10		.67 .08
-.50 .10		.76 .10
-.70 .10		.87 .09
-.90 .08		.61 .11

THETA IS THE ANGLE THAT THE K^+ MAKES WITH THE BEAM IN THE GRAND C.M.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K^+ PROTON.

LABORATORY BEAM MOMENTUM = 1.17 GEV/C $\pm$.4 PER CENT.

COS(THETA)		D-SIGMA/D-OMEGA MB/SR
.78 $\pm$.08		1.89 $\pm$.24
.65 .05		1.26 .14
.55 .05		1.19 .12
.45 .05		.95 .10
.35 .05		1.04 .10
.15 .15		.61 .05
-.05 .05		.42 .05
-.15 .05		.35 .05
-.25 .05		.33 .05
-.35 .05		.26 .06
-.45 .05		.43 .07
-.55 .05		.24 .05
-.65 .05		.41 .07
-.75 .05		.36 .06
-.85 .05		.43 .07
-.93 .03		.73 .17

THETA IS THE ANGLE THAT THE K^+ MAKES WITH THE BEAM IN THE GRAND C.M.

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.LABORATORY BEAM MOMENTUM = 1.97 GEV/C $\pm$ 4(PER CENT).

COS(THETA)	D-SIGMA/D-OMEGA MB/SR
.92 $\pm$.02	1.520 $\pm$.210
.85 $\pm$.05	1.840 $\pm$.150
.75 $\pm$.05	1.470 $\pm$.140
.65 $\pm$.05	1.040 $\pm$.110
.55 $\pm$.05	.490 $\pm$.060
.45 $\pm$.05	.440 $\pm$.060
.35 $\pm$.05	.320 $\pm$.050
.25 $\pm$.05	.270 $\pm$.040
.10 $\pm$.10	.120 $\pm$.060
-.05 $\pm$.05	.061 $\pm$.023
-.15 $\pm$.05	.083 $\pm$.028
-.25 $\pm$.05	.056 $\pm$.021
-.35 $\pm$.05	.029 $\pm$.014
-.45 $\pm$.05	.048 $\pm$.016
-.55 $\pm$.05	.054 $\pm$.016
-.65 $\pm$.05	.064 $\pm$.017
-.75 $\pm$.05	.062 $\pm$.017
-.85 $\pm$.05	.081 $\pm$.019
-.94 $\pm$.04	.043 $\pm$.021

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.LABORATORY BEAM MOMENTUM = 1.97 GEV/C $\pm$ 4(PER CENT).

THETA DEGREES	D-SIGMA/D-OMEGA MB/SR	NO. EVENTS
8.1	5.21 $\pm$ 1.29	153
10.7	4.36 $\pm$ 1.12	117
13.3	3.94 $\pm$.80	119
15.8	2.62 $\pm$.56	80
18.3	2.62 $\pm$.45	81
20.7	2.08 $\pm$.47	62
23.1	1.94 $\pm$.48	57
25.4	2.08 $\pm$.41	48
27.7	2.41 $\pm$.43	39
29.8	2.75 $\pm$.72	23

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.LABORATORY BEAM MOMENTUM = 1.17 GEV/C $\pm$ 4(PER CENT).

THETA DEGREES	D-SIGMA/D-OMEGA MB/SR	NO. EVENTS
9. $\pm$ 2.	3.36 $\pm$.92	150
20. $\pm$ 5.	2.36 $\pm$.45	120

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PHOTON.LABORATORY BEAM MOMENTUM = .97 GEV/C $\pm$ 4(PER CENT).

THETA DEGREES	D-SIGMA/D-OMEGA MB/SR	NO. EVENTS
15. $\pm$ 8.	2.40 $\pm$.78	92

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.K⁺ PROTON ELASTIC CROSS SECTION.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS
.97 $\pm$ 4	10.2 $\pm$ 1.5	1261
1.17 $\pm$ 4	10.7 $\pm$ 1.1	1699
1.97 $\pm$ 4	5.6 $\pm$.4	1593

65 REACTION $K^+p \rightarrow K^0 p \pi^+$ AT 2.26 GEV/C. (PHYS. REV. 198, 1298 (1967))

F. BOMSE, S. BORNSTEIN, J. COLE, D. GILLESPIE, R. KRAEMER, G. LUSTE, I. MILLER, E. MOSES, A. PEVSNER, R. SINGH, R. ZDANIS (JOHNS HOPKINS UNIV., BALTIMORE, MD., USA)

ABSTRACT THE PROPERTIES OF THE $K^*(895 \text{ MEV})$ AND $N^*(1238 \text{ MEV})$ RESONANCES PRODUCED IN THE REACTION $K^+p \rightarrow K^0 p \pi^+$ AT 2.26 GEV/C ARE STUDIED. THE PRODUCTION OF ONE OR THE OTHER OF THESE RESONANCES IS FOUND TO OCCUR IN THIS CHANNEL 80 PER CENT OF THE TIME. THE DECAY CORRELATIONS OF THE K^* AND N^* INDICATE THAT THE DOMINANT MECHANISM INVOLVED IN THEIR PRODUCTION MAY BE SINGLE-VECTOR MESON EXCHANGE. IN PARTICULAR, THE DECAY ANGULAR DISTRIBUTIONS OF THE K^* ARE IN GOOD AGREEMENT WITH THE PREDICTIONS OF THE STODOLSKY-SAKURAI MODEL. THE EXPERIMENTAL DIFFERENTIAL CROSS SECTIONS FOR PRODUCTION OF EACH OF THESE RESONANCES ARE ALSO COMPARED WITH THEORETICAL PREDICTIONS BASED ON A SINGLE-MESON-EXCHANGE MODEL WHICH INCLUDES THE EFFECTS OF ABSORPTION DUE TO COMPETING CHANNELS. THE AGREEMENT OF THE DATA WITH THE THEORY IS FOUND TO BE ONLY MODERATE HERE.

ADDITIONAL FOOTNOTE REFERENCES

BNL F-18 (1960), BNL F-28 (1960), BNL F-55 (1961), BNL F-49 (1961), BNL F-62 (1962), BNL F-68 (1962), BNL F-67 (1960), BNL F-94 (1962), PHYS. REV. LETTERS 11, 93 (1963), PHYS. REV. 133, 8220 (1954), PHYS. LETTERS 6, 309 (1963), PHYS. REV. 142, 913 (1966), PHYS. REV. LETTERS 15, 737 (1965), NUOVO CIMENTO 35, 1105 (1965), NUOVO CIMENTO 34, 1644 (1964), NUOVO CIMENTO 33, 906 (1964), NUOVO CIMENTO 37, 714 (1965), NUOVO CIMENTO 33, 309 (1964), NUOVO CIMENTO 34, 735 (1964), REV. MOD. PHYS. 37, 484 (1965), PHYS. REV. 139, 8428 (1965), PHYS. REV. LETTERS 11, 90 (1963), PHYS. REV. LETTERS 10, 351 (1963), AND PHYS. REV. LETTERS 11, 167 (1963).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 5/68, AND VERIFIED BY LERUY PRICE.

BEAM IS K^+ ON PROTON AT 2.26 GEV/C.

THIS EXPERIMENT USES THE B.N.L. 20 IN. (H) BUBBLE CHAMBER. A TOTAL OF 100000 PICTURES ARE REPORTED ON.

KEY WORDS = $K^*(890)$ DELTA++ DALITZ PLOT ONE PARTICLE EXCHANGE ANGULAR DISTRIBUTION DENSITY MATRIX

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \text{ PROTON } \pi^+$.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS
2.260 $\pm$.023	2.62 $\pm$.30	707

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)^+ \text{ PROTON.}$
 $K^*(890)^+ \rightarrow K^0 \pi^+ [1]$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.260 $\pm$.023	1.1 $\pm$.2

[1] FITTED FOR MASS AND/OR WIDTH (MASS = .895 GEV; WIDTH = .050 GEV), AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND. NON-INTERFERING AMPLITUDES ASSUMED.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow \Delta^{++} K^0.$
 $\Delta^{++} \rightarrow \text{PROTON } \pi^+ [1]$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.260 $\pm$.023	1.1 $\pm$.2

[1] FITTED FOR MASS AND/OR WIDTH (MASS = 1.234 GEV; WIDTH = .120 GEV), AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND. NON-INTERFERING AMPLITUDES ASSUMED.

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)^+ \text{ PROTON.}$
 $K^*(890)^+ \rightarrow K^0 \pi^+ [1]$

LABORATORY BEAM MOMENTUM = 2.260 $\pm$.023 GEV/C.

***** THIS DATA WAS READ FROM A GRAPH. *****

COS(THETA)		D-SIGMA/D-OMEGA MB/SR [2]
MIN	MAX	
-1.0	-.9	.042
-.9	-.8	.011
-.8	-.7	.031
-.7	-.6	.031
-.6	-.5	.000
-.5	-.4	.020
-.4	-.3	.020
-.3	-.2	.000
-.2	-.1	.031
-.1	.0	.021
.0	.1	.000
.1	.2	.031
.2	.3	.051
.3	.4	.071
.4	.5	.109
.5	.6	.151
.6	.7	.261
.7	.8	.331
.8	.9	.302
.9	1.0	.400

THETA IS THE ANGLE THAT THE $K^*(890)^+$ MAKES WITH THE BEAM IN THE GRAND C.M.

[1] USED SIMPLE MASS CUT.

[2] VALUES ARE APPROXIMATE ONLY.

DIFFERENTIAL CROSS SECTION FOR K^+ PROTON $\rightarrow$ DELTA++ K_0 .
 DELTA++ $\rightarrow$ PROTON P_1 [1]
 LABORATORY BEAM MOMENTUM = 2.260 $\pm$.023 GEV/C.

.....
 . THIS DATA WAS READ FROM A GRAPH .

COS(THETA)		D-SIGMA/D-OMEGA MB/SR [2]
MIN	MAX	
.9	1.0	.418
.8	.9	.339
.7	.8	.369
.6	.7	.290
.5	.6	.081
.4	.5	.130
.3	.4	.070
.2	.3	.029
.1	.2	.029
.0	.1	.000
-.1	.0	.009
-.2	-.1	.000
-.3	-.2	.009
-.4	-.3	.000
-.5	-.4	.018
-.6	-.5	.009
-.7	-.6	.050
-.8	-.7	.009
-.9	-.8	.018
-1.0	-.9	.018

THETA IS THE ANGLE THAT THE K_0 MAKES WITH THE BEAM IN THE GRAND C.M.

[1] USED SIMPLE MASS CUT.
 [2] VALUES ARE APPROXIMATE ONLY.

66 K^+P ELASTIC SCATTERING IN THE BACKWARD DIRECTION AT 2.76 BEV/C. (PHYS. REV. LETTERS 21, 1407 (1968))

G.S. ABRAMS, L. EISENSTEIN, T.A. O'HALLORAN, W. SHUFELT, J. WHITMORE (UNIV. OF ILLINOIS, URBANA, ILL., USA)

ABSTRACT THE DIFFERENTIAL CROSS SECTION FOR BACKWARD ($\cos(\theta) \leq -0.8$) K^+P ELASTIC SCATTERING HAS BEEN MEASURED AT 2.76 BEV/C. NO EVIDENCE IS SEEN FOR THE Z(2505) OR FOR A DIP IN THE D-SIGMA/D-U AT 180 DEGREES. A GOOD FIT WITH AN EXPONENTIAL IN $\cos(\theta)$ IS OBTAINED, YIELDING A SLOPE 13.0 ± 2.4 AND AN INTERCEPT D-SIGMA/D-OMEGA(180 DEG.) = 87 ± 16 MICROBARN/SR.

ADDITIONAL FOOTNOTE REFERENCES
 ANL HEP 6802, PHYS. REV. LETTERS 19, 259 (1967), PHYS. REV. 165, 1770 (1968), PHYS. REV. LETTERS 19, 675 (1967),
 PHYS. LETTERS 248, 317 (1967), AND PHYS. REV. 155, 1792 (1967).

ARTICLE READ BY ODETTE BENARY IN 12/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K^+ ON PROTON AT 2.76 GEV/C.

THIS EXPERIMENT USES THE ANL-WISC. 30 IN. (H) BUBBLE CHAMBER.

GENERAL COMMENTS ON THIS ARTICLE
 1 CROSS SECTIONS ARE NORMALIZED TO τ DECAYS.

KEY WORDS = DIFFERENTIAL CROSS SECTION MODELS

ELASTIC DIFFERENTIAL CROSS SECTION FOR K^+ PROTON. (TABLE 1)

LABORATORY BEAM MOMENTUM = 2.76 GEV/C $\pm$ 1(PER CENT).
 NUMBER OF EVENTS = 90.

COS(THETA)		D-SIGMA/D-OMEGA UB/SR	U (GEV/C)**2		D-SIGMA/D-U UB/(GEV/C)**2
MIN	MAX		MAX	MIN	
-1.00 [1]		87. $\pm$ 16.	.065 [1]		251. $\pm$ 48.
-1.00	-.98	80. 18.	.065	-.023	241. 53.
-.98	-.96	49. 13.	.023	-.020	146. 40.
-.96	-.94	52. 13.	-.020	-.062	155. 39.
-.94	-.92	32. 10.	-.062	-.104	95. 31.
-.92	-.90	29. 10.	-.104	-.146	86. 30.
-.90	-.88	20. 9.	-.146	-.188	50. 26.
-.88	-.86	17. 7.	-.188	-.230	52. 22.
-.86	-.84	12. 5.	-.230	-.272	35. 18.

THETA IS THE ANGLE THAT THE K^+ MAKES WITH THE BEAM IN THE GRAND C.M.
 U IS THE MOMENTUM TRANSFER BETWEEN THE [INCOMING K^+] AND THE [OUTGOING K^+].

[1] EXTRAPOLATED POINT.

FIT TO ELASTIC DIFFERENTIAL CROSS SECTION FOR K^+ PROTON. (PAGE 1408)

LABORATORY BEAM MOMENTUM = 2.76 GEV/C $\pm$ 1(PER CENT).
 FITTED FORMULA IS D-SIGMA/D-OMEGA = [D-SIGMA/D-OMEGA(AT 180 DEGREES)] * EXP[-A(1+COS(THETA))]

WHERE D-SIGMA/D-OMEGA IS IN UB/SR. THETA IS THE ANGLE THAT THE K^+ MAKES WITH THE BEAM IN THE GRAND C.M.

FITTER VALUE

A = 13.0 $\pm$ 2.4

67 ELASTIC BACKWARD SCATTERING OF PIONS AND KAONS [NUC. PHYS. 89, 249 (1969)]

W.F. BAKER, P.J. CARLSON, V. CHABAUD, A. LUNDY (EUROPEAN ORG. FOR NUC. RES., GENEVA, SWITZERLAND)
J. BANAIGS, J. BERGER, C. BOYNEL, J. DUFLO, L. GOLDSZAL, F. PLOUIN (CNTR. D'ETUDES NUC. SACLAY, GIF-SUR-YVETTE, FRANCE)

ABSTRACT THE ELASTIC SCATTERING OF π^+ MESONS BY PROTONS NEAR 180 DEGREES HAS BEEN MEASURED AT 2.85, 3.30, AND 3.55 GEV/C. THAT OF π^- MESONS AT 3.30 AND 3.55 GEV/C, AND THAT OF K^+ AND K^- MESONS AT 3.55 GEV/C. STEEPLY RISING BACKWARD PEAKS ARE OBSERVED AT ALL ENERGIES FOR π^+ SCATTERING, WHEREAS THE π^+ -P ANGULAR DISTRIBUTIONS ARE NOT AS STEEP. K^+ -P ELASTIC SCATTERING ALSO SHOWS A BACKWARD PEAK, NO BACKWARD K^+ -P ELASTIC SCATTERING EVENT WAS FOUND. THE UPPER LIMIT (90 PER CENT CONFIDENCE) FOR THE K^+ -P DIFFERENTIAL CROSS SECTION IS TWELVE TIMES SMALLER THAN THE CORRESPONDING K^+ -P CROSS SECTION. THE BEHAVIOUR OF THE DIFFERENTIAL CROSS SECTION CLOSE TO 180 DEGREES CAN BE EXPLAINED USING DIRECT CHANNEL RESONANCES. THE LARGE RATIO OF THE K^+ TO K^+ -P CROSS SECTION CAN BE UNDERSTOOD IN TERMS OF BARYON EXCHANGE.

CROSS REFERENCES
THIS ARTICLE SUPERSEDES PHYS. LETTERS 248, 317 (1967).

ADDITIONAL FOOTNOTE REFERENCES
PHYS. REV. 159, 1169 (1967), PHYS. LETTERS 10, 248 (1964), NUOVO CIMENTO 38, 60 (1965), PHYS. REV. LETTERS 19, 476 (1967), PHYS. REV. 132, 1252 (1963), PHYS. REV. 138, 8707 (1965), PHYS. REV. LETTERS 12, 429 (1964), PHYS. LETTERS 23, 605 (1966), PHYS. LETTERS 248, 317 (1967), PHYS. LETTERS 258, 361 (1967), NUCLEAR INSTRUMENTS 20, 86 (1963), CERN NP/44-1 (1964), NUCLEAR INSTRUMENTS 26, 137 (1964), PHYS. REV. 144, 1101 (1966), PHYS. REV. LETTERS 19, 678 (1967), PHYS. REV. LETTERS 19, 460 (1967), NUOVO CIMENTO 45, 885 (1966), PHYS. REV. LETTERS 19, 575 (1967), PHYS. LETTERS 21, 117 (1966), CERN HADRON CONFERENCE, PHYS. REV. LETTERS 17, 102 (1966), PHYS. REV. LETTERS 18, 798 (1967), PHYS. REV. LETTERS 20, 607 (1968), INT. CONF. ON PARTICLES AND FIELDS, ROCHESTER, NY, 1967 57 (1967), AND UCRL 8030 AUG. (1968).

ARTICLE READ BY ODETTE BENVARY IN 3/69, AND VERIFIED BY LEROY PRICE.

BEAM NO. 1 IS π^+ ON PROTON FROM 2.85 TO 3.55 GEV/C.
NO. 2 IS π^- ON PROTON FROM 3.30 TO 3.55 GEV/C.
NO. 3 IS K^+ ON PROTON AT 3.55 GEV/C.
NO. 4 IS K^- ON PROTON AT 3.55 GEV/C.

THIS EXPERIMENT USES SPARK CHAMBERS. A TOTAL OF 54000 PICTURES ARE REPORTED ON.

KEY WORDS = DIFFERENTIAL CROSS SECTION FITS MODELS

COMPOUND KEY WORDS = FITS DIFFERENTIAL CROSS SECTION

ELASTIC DIFFERENTIAL CROSS SECTION FOR K^+ PROTON. [TABLE 7]

LABORATORY BEAM MOMENTUM = 3.55 GEV/C $\pm$ 1 (PER CENT).

[THIS DATA REPLACES THE VALUE GIVEN EARLIER IN BANAIGS, ET AL., PHYS. LETTERS 248, 317 (1967)]

COS(THETA)	D-SIGMA/D-UMEGA UB/SA	NO. EVENTS	U (GEV/C)**2	D-SIGMA/D-U UB/(GEV/C)**2	NO. EVENTS
.9940 $\pm$.0080	21.3 $\pm$ 5.9	15	.035	47.4 $\pm$ 13.1	15
.9835 .0130	14.9 5.8	7	.005	33.1 12.9	7

THETA IS THE ANGLE THAT THE K^+ MAKES WITH THE BEAM IN THE GRAND C.M.
U IS THE MOMENTUM TRANSFER BETWEEN THE [INCOMING K^+] AND THE [OUTGOING K^+].

68 SINGLE PION PRODUCTION IN THE K^+ CHANNEL FROM 860 TO 1360 TO 1360 MEV/C. [UCRL 18131 (1968)]

R.W. BLAND (U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA)

ABSTRACT WE HAVE STUDIED SINGLE PION PRODUCTION BY K^+ MESONS ON PROTONS FROM 860 TO 1360 MEV/C. THE INCIDENT BEAM MOMENTUM, WE FIND STRONG $\pi^+(1236)$ PRODUCTION IN THIS REGION, WITH ITS MAXIMUM CROSS SECTION OCCURRING SLIGHTLY BELOW THE 1250 MEV/C PEAK IN THE TOTAL CROSS SECTION REPORTED BY COOL ET AL. A PARTIAL WAVE ANALYSIS OF THE K^+ FINAL STATE INDICATES THAT THE π^+ PRODUCTION IS DOMINATED BY THE $P_{1/2}$ AND $P_{3/2}$ STATES IN THE REGION OF THE COOL PEAK, WITH NO EVIDENT PHASE VARIATION FROM 860 TO 1360 MEV/C. THE π^+ PRODUCTION IS QUALITATIVELY CONSISTENT WITH THE PREDICTIONS OF THE MAGNETIC DIPOLE (M_1) RHO-EXCHANGE MODEL; NEAR THRESHOLD, HOWEVER, THE EXPERIMENTAL CROSS SECTION IS TOO HIGH BY A FACTOR OF 5 AND THE DETAILED PARTIAL WAVE STRUCTURE DEVIATES CONSIDERABLY FROM THE M_1 PREDICTIONS. THE SMOOTH VARIATION WITH THE BEAM MOMENTUM OF ALL PROPERTIES OF THE K^+ SYSTEM SUGGESTS THAT THE K^+ CHANNEL HAS NO STRONG RESONANT ACTIVITY, AND SHOULD BE INTERPRETED IN TERMS OF T-CHANNEL PHENOMENA.

CROSS REFERENCES
THIS ARTICLE SUPERSEDES PART OF PHYS. REV. LETTERS 17, 939 (1966).
PART OF THIS ARTICLE SUPERSEDED BY UCRL 18758 (1968).

ADDITIONAL FOOTNOTE REFERENCES
PHYS. REV. LETTERS 9, 135 (1962), PHYS. REV. LETTERS 7, 188 (1961), PHYS. REV. LETTERS 7, 182 (1961), PHYS. REV. LETTERS 17, 102 (1966), UCRL 11527 (1964), LRL INTERDEPARTMENTAL REPORT TN-29 (1964), UCRL 8030 JAN. (1967), PHYS. REV. LETTERS 9, 135 (1962), CERN PHYS. 67-10 (1967), PHYS. LETTERS 248, 314 (1967), PHYS. REV. LETTERS 7, 189 (1961), CERN CONFERENCE 358 (1962), PHYS. REV. LETTERS 11, 93 (1963), PHYS. REV. 133, 8220 (1964), PHYS. LETTERS 16, 83 (1964), PHYS. LETTERS 6, 309 (1963), UCRL 16370 (1965), PHYS. REV. 139B, 1411 (1965), PHYS. REV. 142, 913 (1966), NUOVO CIMENTO 33, 90A (1964), NUOVO CIMENTO 34, 1841E (1964), PHYS. REV. 158, 1298 (1967), PHYS. REV. 158, 1310 (1967), NUOVO CIMENTO 36, 1101 (1965), PHYS. REV. LETTERS 11, 90 (1963), BULL. AM. PHYS. SOC. 8, 20 (1963), PHYS. LETTERS 6, 62 (1962), PHYS. REV. LETTERS 1F, 777 (1964), CERN CONFERENCE 721 (1967), LRL INTERDEPARTMENTAL REPORT TG-149 (1966), PHYS. REV. LETTERS 17, 939 (1966), BULL. AM. PHYS. SOC. 12, 506 (1967), AND PHYS. REV. 140, 897 (1965).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 5/68, AND VERIFIED BY ROGER BLAND.

BEAM IS K^+ ON PROTON FROM .865 TO 1.365 GEV/C.

THIS EXPERIMENT USES THE L.R.L. 25 IN. (H) BUBBLE CHAMBER. A TOTAL OF 198000 PICTURES ARE REPORTED ON.

GENERAL COMMENTS ON THIS ARTICLE

1 CONTAINS CORRECTIONS TO PREVIOUS GOLDBERGER GROUP K^+ DATA. (NOW NORMALIZED TO COOL RATHER THAN TAU DECAYS).
2 ALL DATA (EXCEPT AT 1.360 GEV/C AND THE TOTAL CROSS SECTIONS) ARE NORMALIZED TO COOL.

3 GIVES UPPER LIMITS FOR K^+ MESON.

4 MUCH OF THE DATA IN THIS THESIS IS REPEATED IN A LATER PUBLICATION (UCRL 18758), SO IT HAS NOT BEEN INCLUDED HERE.

KEY WORDS = CROSS SECTION ANGULAR DISTRIBUTION PHASE SHIFT $K^+(890)$ DELTA(1236) KAPPA MESON
MODELS DENSITY MATRIX

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \Delta^{++}$ (FIGURE 9)
 $\Delta^{++} \rightarrow \text{PROTON } \pi^+$ [1]

LABORATORY BEAM MOMENTUM = .85 GEV/C.

(CUT MADE ON DALITZ PLOT TO REMOVE INTERFERENCE REGION. VALUES GIVEN HERE ARE FROM R. BLAND - PRIVATE COMMUNICATION.)

COS(THETA)		NO. EVENTS
MIN	MAX	
.8	1.0	36
.6	.8	52
.4	.6	76
.2	.4	99
.0	.2	108
-.2	.0	114
-.4	-.2	125
-.6	-.4	118
-.8	-.6	101
-1.0	-.8	72

THETA IS THE ANGLE THAT THE K^0 MAKES WITH THE BEAM IN THE GRAND C.M.

[1] COUNTED ALL EVENTS IN MASS BAND (MASS = 1.220 GEV; WIDTH = .120 GEV). OVERLAP REGION EXCLUDED.

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \Delta^{++}$ (FIGURE 9)
 $\Delta^{++} \rightarrow \text{PROTON } \pi^+$ [1]

LABORATORY BEAM MOMENTUM = .96 GEV/C.

(CUT MADE ON DALITZ PLOT TO REMOVE INTERFERENCE REGION. VALUES GIVEN HERE ARE FROM R. BLAND - PRIVATE COMMUNICATION.)

COS(THETA)		NO. EVENTS
MIN	MAX	
.8	1.0	39
.6	.8	85
.4	.6	119
.2	.4	145
.0	.2	171
-.2	.0	181
-.4	-.2	158
-.6	-.4	178
-.8	-.6	151
-1.0	-.8	93

THETA IS THE ANGLE THAT THE K^0 MAKES WITH THE BEAM IN THE GRAND C.M.

[1] COUNTED ALL EVENTS IN MASS BAND (MASS = 1.220 GEV; WIDTH = .120 GEV). OVERLAP REGION EXCLUDED.

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \Delta^{++}$ (FIGURE 9)
 $\Delta^{++} \rightarrow \text{PROTON } \pi^+$ [1]

LABORATORY BEAM MOMENTUM = 1.2 GEV/C.

(CUT MADE ON DALITZ PLOT TO REMOVE INTERFERENCE REGION. VALUES GIVEN HERE ARE FROM R. BLAND - PRIVATE COMMUNICATION.)

COS(THETA)		NO. EVENTS
MIN	MAX	
.8	1.0	20
.6	.8	43
.4	.6	76
.2	.4	74
.0	.2	83
-.2	.0	103
-.4	-.2	120
-.6	-.4	106
-.8	-.6	130
-1.0	-.8	73

THETA IS THE ANGLE THAT THE K^0 MAKES WITH THE BEAM IN THE GRAND C.M.

[1] COUNTED ALL EVENTS IN MASS BAND (MASS = 1.220 GEV; WIDTH = .120 GEV). OVERLAP REGION EXCLUDED.

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \Delta^{++}$ (FIGURE 9)
 $\Delta^{++} \rightarrow \text{PROTON } \pi^+$ [1]

LABORATORY BEAM MOMENTUM = 1.36 GEV/C.

(CUT MADE ON DALITZ PLOT TO REMOVE INTERFERENCE REGION. VALUES GIVEN HERE ARE FROM R. BLAND - PRIVATE COMMUNICATION.)

COS(THETA)		NO. EVENTS
MIN	MAX	
.8	1.0	5
.6	.8	17
.4	.6	19
.2	.4	37
.0	.2	28
-.2	.0	45
-.4	-.2	41
-.6	-.4	50
-.8	-.6	66
-1.0	-.8	44

THETA IS THE ANGLE THAT THE K^0 MAKES WITH THE BEAM IN THE GRAND C.M.

[1] COUNTED ALL EVENTS IN MASS BAND (MASS = 1.220 GEV; WIDTH = .120 GEV). OVERLAP REGION EXCLUDED.

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} = \text{PROTON } K^*(890)^+$ (FIGURE 26)
 $K^*(890)^+ = K^0 \pi^+$ (1)

LABORATORY BEAM MOMENTUM = 1.2 GEV/C.

(CUT MADE ON DALITZ PLOT TO REMOVE INTERFERENCE REGION. VALUES GIVEN HERE ARE FROM R. BLAND - PRIVATE COMMUNICATION.)

COS(THETA)		NO. EVENTS
MIN	MAX	
-1.0	-0.8	12
-0.8	-0.6	27
-0.6	-0.4	31
-0.4	-0.2	43
-0.2	0.0	36
0.0	0.2	41
0.2	0.4	49
0.4	0.6	57
0.6	0.8	42
0.8	1.0	56

THETA IS THE ANGLE THAT THE $K^*(890)^+$ MAKES WITH THE BEAM IN THE GRAND C.M.

(1) COUNTED ALL EVENTS IN MASS BAND [MASS = .890 GEV; WIDTH = .133 GEV]. OVERLAP REGION EXCLUDED.

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} = \text{PROTON } K^*(890)^+$ (FIGURE 26)
 $K^*(890)^+ = K^0 \pi^+$ (1)

LABORATORY BEAM MOMENTUM = 1.36 GEV/C.

(CUT MADE ON DALITZ PLOT TO REMOVE INTERFERENCE REGION. VALUES GIVEN HERE ARE FROM R. BLAND - PRIVATE COMMUNICATION.)

COS(THETA)		NO. EVENTS
MIN	MAX	
-1.0	-.8	8
-.8	-.6	10
-.6	-.4	10
-.4	-.2	33
-.2	.0	23
.0	.2	31
.2	.4	28
.4	.6	37
.6	.8	43
.8	1.0	52

THETA IS THE ANGLE THAT THE $K^*(890)^+$ MAKES WITH THE BEAM IN THE GRAND C.M.

(1) COUNTED ALL EVENTS IN MASS BAND [MASS = .890 GEV; WIDTH = .100 GEV]. OVERLAP REGION EXCLUDED.

69 THE REACTION $K^+p \rightarrow K^+n \pi^+$ IN ALL THREE FINAL CHARGE STATES AT 3 GEV/C. (NUOVO CIMENTO 49A, 348 (1967))

P. SALLSTRÖM, G. OTTER, G. EKSPONG (STOCKHOLMS UNIV., STOCKHOLM, SWEDEN)

ABSTRACT THE REACTION $K^+p \rightarrow K^+n \pi^+$ HAS BEEN STUDIED IN ALL THREE FINAL CHARGE STATES AT A MOMENTUM OF THE INCIDENT K^+ -MESON OF 2.97 GEV/C. THE RESONANCE $K^*(890)$ IS PRODUCED WITH A CROSS-SECTION OF (1.56 ± 0.19) MB AND A DECAY BRANCHING RATIO $(K^+ \rightarrow K^0 \pi^+)/(K^+ \rightarrow K^0 \pi^+ + 0.43 \pm 0.11)$. THE RESONANCE $N^*(1236)$ IS PRODUCED AS $N^+ \pi^+$ WITH THE CROSS-SECTION (0.70 ± 0.13) MB AND AS $N^+ \pi^0$ WITH THE CROSS-SECTION (0.27 ± 0.08) MB. THE DECAY BRANCHING RATIO OF N^+ , $(N^+ \rightarrow K^0 \pi^+)/(N^+ \rightarrow K^0 \pi^0)$ IS 0.05 ± 0.20 . THE CONSEQUENCES OF AN INTERFERENCE BETWEEN STATES WITH DIFFERENT ISOTOPIC SPIN OF THE $(K^+ \pi^+)$ -SYSTEM ARE DISCUSSED. THE DECAY CORRELATIONS OF THE RESONANCE EVENTS ARE PRESENTED AND THE SPIN SPACE DENSITY MATRIX ELEMENTS FOR THE RESONANCES ARE CALCULATED. THE N^+ DECAY DISTRIBUTIONS AGREE WITH THE PREDICTIONS OF THE MODEL OF STODOLSKY AND SAKURAI. THE DECAY ANGULAR CORRELATIONS OF THE TWO DECAY MODES OF K^+ , $K^+ \rightarrow K^0 \pi^+$ AND $K^+ \rightarrow K^0 \pi^0$, ARE FOUND TO BE SIGNIFICANTLY DIFFERENT. THIS MAY BE EXPLAINED AS DUE TO AN INTERFERENCE BETWEEN STATES WITH DIFFERENT ISOTOPIC SPIN OF THE $(K^+ \pi^+)$ -SYSTEM. THE K^+ DECAY CORRELATIONS, WHEN INTERFERENCE EFFECTS ARE ELIMINATED, INDICATE THAT THE RESONANCE IS PRODUCED TO A GREAT EXTENT VIA THE EXCHANGE OF A VECTOR MESON (ONE-MESON EXCHANGE MODEL). THE PRODUCTION ANGULAR DISTRIBUTIONS OF BOTH RESONANCES ARE SHARPLY PEAKED. THE MODIFIED ONE-PARTICLE EXCHANGE MODEL OF JACKSON ET AL. IS SHOWN TO GIVE A REASONABLE ACCOUNT OF THESE DISTRIBUTIONS AS WELL AS OF THE DECAY CORRELATIONS AND THE VARIATION OF THE DENSITY MATRIX ELEMENTS WITH THE PRODUCTION ANGLE OF THE RESONANCE.

CROSS REFERENCES
 SAME UP THIS UNIT RENORMALIZED IN UCL 18753 (1969).

ADDITIONAL FOOTNOTE REFERENCES

NUOVO CIMENTO 49A, 348 (1967); UCL 11068 (1963); NUOVO CIMENTO 36, 1101 (1955); PHYS. LETTERS 16, 83 (1965); NUOVO CIMENTO 34, 1844 (1954); NUOVO CIMENTO 33, 309 (1964); PHYS. REV. 139, 8428 (1965); CERN 83-12 (1963); NUOVO CIMENTO 43A, 142 (1966); PHYS. REV. LETTERS 10, 351 (1963); AND PHYS. REV. LETTERS 11, 90 (1963).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 5/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K^+ ON PROTON AT 2.972 GEV/C.

THIS EXPERIMENT USES THE SACLAY 81 CM (H) BUBBLE CHAMBER. A TOTAL OF 20000 PICTURES ARE REPORTED ON.

KEY WORDS = $K^*(890)$ DELTA++ CROSS SECTION DALITZ PLOT MASS SPECTRUM ANGULAR DISTRIBUTION
 DENSITY MATRIX MUONS

K^+ PROTON TOTAL CROSS SECTION.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.972 $\pm$.015	20. $\pm$ 2.

CROSS SECTION FOR $K^+ \text{ PROTON} = K^+ \text{ PROTON } \pi^0$.

LABORATORY BEAM MOMENTUM GEV/C	ORIGINAL DATA	NO. EVENTS	RENORMALIZED VALUE (1).
2.972 $\pm$.015	MILLI-BARNS 1.3 $\pm$.2	165	MILLI-BARNS 1.1 $\pm$.2

(1) BLAND, ET. AL., UCL 18753 (1969).

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \text{ PROTON } \pi^+$.

ORIGINAL DATA		RENORMALIZED VALUE (1)	
LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS	MILLI-BARNS
2.972 $\pm$.015	2.7 $\pm$.3	221	2.3 $\pm$.3

(1) BLAND, ET. AL., UCRL 18758 (1969).

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \text{ NEUTRON } \pi^+$.

ORIGINAL DATA		RENORMALIZED VALUE (1)	
LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS	MILLI-BARNS
2.972 $\pm$.015	1.2 $\pm$.2	165	1.0 $\pm$.2

(1) BLAND, ET. AL., UCRL 18758 (1969).

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \text{ MESON NUCLEON } \pi^0$.

ORIGINAL DATA		RENORMALIZED VALUE (1)	
LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS	MILLI-BARNS
2.972 $\pm$.015	5.2 $\pm$.4	551	4.4 $\pm$.4

(1) BLAND, ET AL., UCRL 18758 (1969).

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)^+ \text{ PROTON.}$
 $K^*(890)^+ \rightarrow K^+ \pi^0$ (1)

(THIS DATA SHOULD NOT BE USED - A MORE RECENT VALUE MAY BE FOUND IN SALLSTROM, ET AL., ARKIV FOR FYSIK 37, 463 (1959))

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.972 $\pm$.015	.47 $\pm$.10

(1) FITTED FOR MASS AND/OR WIDTH (MASS = .883 GEV; WIDTH = .047 GEV), AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)^+ \text{ PROTON.}$
 $K^*(890)^+ \rightarrow K^0 \pi^+$ (1)

(THIS DATA SHOULD NOT BE USED - A MORE RECENT VALUE MAY BE FOUND IN SALLSTROM, ET AL., ARKIV FOR FYSIK 37, 463 (1959))

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.972 $\pm$.015	1.09 $\pm$.16

(1) FITTED FOR MASS AND/OR WIDTH (MASS = .898 GEV; WIDTH = .068 GEV), AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \Delta^{++}. \Delta^{++} \rightarrow \text{PROTON } \pi^+$ (1)

(THIS DATA SHOULD NOT BE USED - A MORE RECENT VALUE MAY BE FOUND IN SALLSTROM, ET AL., ARKIV FOR FYSIK 37, 463 (1959))

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.972 $\pm$.015	.70 $\pm$.13

(1) FITTED FOR MASS AND/OR WIDTH (MASS = 1.221 GEV; WIDTH = .115 GEV), AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \Delta^{++}. \Delta^{++} \rightarrow \text{NEUTRON } \pi^+$ (1)

(THIS DATA SHOULD NOT BE USED - A MORE RECENT VALUE MAY BE FOUND IN SALLSTROM, ET AL., ARKIV FOR FYSIK 37, 463 (1959))

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.972 $\pm$.015	.01 $\pm$.05

(1) COUNTED ONLY EVENTS ABOVE BACKGROUND.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \Delta^{++}$
 $\Delta^{++} \rightarrow \text{PROTON } \pi^0$ (1)

(THIS DATA SHOULD NOT BE USED - A MORE RECENT VALUE MAY BE FOUND IN SALLSTROM, ET AL., ARKIV FOR FYSIK 37, 463 (1969))

LABORATORY
 BEAM MOMENTUM
 GEV/C
 2.972 $\pm$.015
 MILLI-BARNS
 .26 $\pm$.08

(1) FITTED FOR MASS AND/OR WIDTH (MASS = 1.267 GEV; WIDTH = .272 GEV), AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)^+ \text{ PROTON}$
 $K^*(890)^+ \rightarrow \pi^0 \pi^+$ (1)

LABORATORY BEAM MOMENTUM = 2.972 $\pm$.015 GEV/C.

(THIS DATA SHOULD NOT BE USED - A MORE RECENT VALUE MAY BE FOUND IN SALLSTROM, ET AL., ARKIV FOR FYSIK 37, 463 (1969))

.....
 . THIS DATA WAS READ FROM A GRAPH .

COS(THETA)		D-SIGMA/D-OMEGA UR/SR (2)		NO. EVENTS
MIN	MAX			
-1.00	-.50	.00	$\pm$.05	0
-.50	-.45	.05	.05	1
-.45	-.40	.14	.08	3
-.40	-.35	.05	.05	1
-.35	.00	.00	.05	0
.00	.05	.05	.05	1
.05	.25	.00	.05	0
.25	.30	.09	.07	2
.30	.35	.00	.05	0
.35	.40	.05	.05	1
.40	.55	.00	.05	0
.55	.60	.09	.07	2
.60	.65	.05	.05	1
.65	.70	.00	.05	0
.70	.75	.19	.09	4
.75	.80	.14	.08	3
.80	.85	.30	.10	8
.85	.90	.42	.14	9
.90	.95	.52	.16	11
.95	1.00	1.32	.25	28

THETA IS THE ANGLE THAT THE $K^*(890)^+$ MAKES WITH THE BEAM IN THE GRAND C.M.

(1) COUNTED ALL EVENTS IN MASS BAND.

(2) COUNTS WERE MULTIPLIED BY .047 TO GET THESE. ERRORS ARE TAKEN AS PROPORTIONAL TO THE SQUARE-ROOT OF THE COUNTS.

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)^+ \text{ PROTON}$
 $K^*(890)^+ \rightarrow \pi^0 \pi^+$ (1)

LABORATORY BEAM MOMENTUM = 2.972 $\pm$.015 GEV/C.

.....
 . THIS DATA WAS READ FROM A GRAPH .

COS(THETA)		D-SIGMA/D-OMEGA UR/SR (2)		NO. EVENTS
MIN	MAX			
-1.00	-.75	.03	$\pm$.03	1
-.95	-.75	.00	.03	0
-.75	-.70	.03	.03	1
-.70	-.65	.00	.03	0
-.65	-.60	.03	.03	1
-.60	.65	.00	.03	0
.65	.70	.03	.03	1
.70	.75	.05	.04	2
.75	.80	.07	.04	3
.80	.85	.10	.05	4
.85	.90	.20	.07	8
.90	.95	.30	.09	12
.95	1.00	.72	.13	29

THETA IS THE ANGLE THAT THE $K^*(890)^+$ MAKES WITH THE BEAM IN THE GRAND C.M.

(1) COUNTED ALL EVENTS IN MASS BAND.

(2) COUNTS WERE MULTIPLIED BY .025 TO GET THESE. ERRORS ARE TAKEN AS PROPORTIONAL TO THE SQUARE-ROOT OF THE COUNTS.

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow \Delta^{++} K^+$.
LABORATORY BEAM MOMENTUM = $2.972 \pm .015 \text{ GEV/C}$.

THIS DATA WAS READ FROM A GRAPH.

COS(THETA)		D-SIGMA/D-OMEGA UB/SR [1]		NO. EVENTS
MIN	MAX			
.95	1.00	.29	± .09	10
.90	.95	.26	± .09	9
.85	.90	.14	± .06	5
.80	.85	.09	± .05	3
.75	.80	.03	± .03	1
.70	.75	.06	± .04	2
.65	.70	.06	± .04	2
.60	.65	.00	± .03	0
.45	.50	.06	± .04	2
.40	.45	.00	± .03	0
.35	.40	.03	± .03	1
.30	.35	.03	± .03	1
-.40	.30	.00	± .03	0
-.45	-.40	.03	± .03	1
-.50	-.45	.03	± .03	1
-.75	-.50	.00	± .03	0
-.80	-.75	.03	± .03	1
-1.00	-.80	.00	± .03	0

THETA IS THE ANGLE THAT THE K^+ MAKES WITH THE BEAM IN THE GRAND C.M.

[1] COUNTS WERE MULTIPLIED BY .029 TO GET THESE. ERRORS ARE TAKEN AS PROPORTIONAL TO THE SQUARE-ROOT OF THE COUNTS.

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow \Delta^{++} K^0$.
LABORATORY BEAM MOMENTUM = $2.972 \pm .015 \text{ GEV/C}$.

THIS DATA WAS READ FROM A GRAPH.

COS(THETA)		D-SIGMA/D-OMEGA UB/SR [1]		NO. EVENTS
MIN	MAX			
.95	1.00	.43	± .12	12
.90	.95	.61	± .15	17
.85	.90	.36	± .11	10
.80	.85	.36	± .11	10
.75	.80	.22	± .09	6
.70	.75	.18	± .08	5
.65	.70	.04	± .04	1
.60	.65	.11	± .06	3
.55	.60	.04	± .04	1
.50	.55	.04	± .04	1
.45	.50	.00	± .04	0
.40	.45	.04	± .04	1
.35	.40	.04	± .04	1
.30	.35	.00	± .04	0
.25	.30	.04	± .04	1
.05	.25	.00	± .04	0
.00	.05	.04	± .04	1
-.55	.00	.00	± .04	0
-.60	-.55	.04	± .04	1
-.85	-.60	.00	± .04	0
-.90	-.85	.04	± .04	1
-1.00	-.90	.00	± .04	0

THETA IS THE ANGLE THAT THE K^0 MAKES WITH THE BEAM IN THE GRAND C.M.

[1] COUNTS WERE MULTIPLIED BY .036 TO GET THESE. ERRORS ARE TAKEN AS PROPORTIONAL TO THE SQUARE-ROOT OF THE COUNTS.

70 THE REACTION $K^+ P \rightarrow K^0 P \pi^+$ AT 3 GEV/C. (NUOVO CIMENTO 36, 1101 (1955))

M.FERRO-LUZZI, R.GEORGE, Y.GOLDSCHMIDT-CLERMONT, V.P.HENRI, B.JONGEJANS, D.W.G.LEITH, G.R.LYNCH, F.MULLER, J.M.PERRAU (EUROPEAN ORG. FOR NUC. RES., GENÈVE, SWITZERLAND)

ABSTRACT THE REACTION $K^+ P \rightarrow K^0 P \pi^+$ HAS BEEN STUDIED AT 3.0 GEV/C. THE TWO DOMINANT REACTIONS IN THIS FINAL STATE ARE $K^+ P \rightarrow K^* P$, $K^* \rightarrow K^0 \pi^+$ AND $K^+ P \rightarrow \pi^+ K^* P$, ACCOUNTING EACH FOR (38 ± 3) PER CENT OF THE EVENTS AND CORRESPONDING TO A CROSS-SECTION OF (0.8 ± 0.1) MB. THE PRODUCTION ANGULAR DISTRIBUTIONS FOR BOTH K^* AND π^+ ARE SUGGESTIVE OF A PERIPHERAL MECHANISM OF PRODUCTION. WITHIN THE FRAMEWORK OF A SIMPLE ONE-MESON-EXCHANGE MODEL, THE DECAY ANGULAR DISTRIBUTIONS FOR K^* ARE CONSISTENT WITH THE ASSUMPTION THAT THESE EVENTS ARE PREDOMINANTLY PRODUCED BY THE EXCHANGE OF A VECTOR MESON WITH ONLY (7 ± 6) PER CENT OF THE EVENTS BEING PRODUCED BY THE EXCHANGE OF A PION. THE π^+ DECAY DISTRIBUTIONS AGREE WITH THE PREDICTION OF THE MODEL OF STODOLSKY AND SAKURAI. FOR BOTH K^* AND π^+ , A FORM FACTOR OF THE TYPE $\exp(-\Delta^2/\Lambda^2)$, WITH Λ EQUAL TO ABOUT 0.6 GEV/C, IS NEEDED TO FIT THE VERY SHARPLY PEAKED PRODUCTION DISTRIBUTIONS. A MODIFIED ONE-MESON-EXCHANGE MODEL, TAKING INTO ACCOUNT ABSORPTION EFFECTS DUE TO COMPETING PROCESSES, DERIVED BY GOTTFRIED, JACKSON AND SVENSSON, GIVES A REASONABLE ACCOUNT OF THE PRODUCTION ANGULAR DISTRIBUTION AND OF THE DECAY CORRELATIONS FOR BOTH K^* AND π^+ . $N^0 K^0$, $N^0 \pi^0$

CROSS REFERENCES
THIS ARTICLE SUPERSEDES PHYS. LETTERS 9, 359 (1964).

ADDITIONAL FOOTNOTE REFERENCES
CERN 63-12 (1963), PHYS. LETTERS 9, 359 (1964), NUOVO CIMENTO 36, 1644 (1964), PHYS. REV. 133, B220 (1964), PHYS. LETTERS 6, 309 (1953), PHYS. REV. LETTERS 10, 138 (1963), PHYS. LETTERS 8, 144 (1964), NUOVO CIMENTO 33, 309 (1954), PHYS. REV. LETTERS 8, 140 (1962), PHYS. LETTERS 12, 352 (1964), PHYS. REV. LETTERS 11, 93 (1963), NUOVO CIMENTO 33, 906 (1964), CERN TH-428, SIENNA CONFERENCE I 189 (1963), AND PHYS. LETTERS 6, 62 (1963).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 3/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K^+ ON PROTON AT 2.965 GEV/C.

THIS EXPERIMENT USES THE SACLAY 81 CM (H) BUBBLE CHAMBER. A TOTAL OF 100000 PICTURES ARE REPORTED ON.

KEY WORDS = CROSS SECTION DELTA(1238) MASS MODELS $K^*(893)$ ONE PARTICLE EXCHANGE DALITZ PLOT
ANGULAR DISTRIBUTION DENSITY MATRIX LIFETIME

COMPOUND KEY WORDS = DELTA(1238) MASS $K^*(890)$ MASS DELTA(1238) LIFETIME $K^*(890)$ LIFETIME

CROSS SECTION FOR $K^+ \text{ PROTON} = K^0 \text{ PROTON } \pi^+.$

(TOTAL, INCLUDING RESONANCE PRODUCTION)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS
2.965 $\pm$.015	2.1 $\pm$.3	747

CROSS SECTION FOR $K^+ \text{ PROTON} = K^*(890)^+ \text{ PROTON}.$
 $K^*(890)^+ = K^0 \pi^+ \{1\}$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.965 $\pm$.015	.8 $\pm$.1

\{1\} FITTED FOR MASS AND/OR WIDTH [MASS = .891 GEV; WIDTH = .047 GEV], AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.

CROSS SECTION FOR $K^+ \text{ PROTON} = K^0 \text{ DELTA}^{++}.$
 $\text{DELTA}^{++} = \pi^+ \text{ PROTON } \{1\}$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.965 $\pm$.015	.8 $\pm$.1

\{1\} FITTED FOR MASS AND/OR WIDTH [MASS = 1.232 GEV; WIDTH = .125 GEV], AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} = K^*(890)^+ \text{ PROTON}.$
 $K^*(890)^+ = K^0 \pi^+ \{1\}$

LABORATORY BEAM MOMENTUM = 2.965 $\pm$.015 GEV/C.

 . THIS DATA WAS READ FROM A GRAPH .

COS(THETA)		D-SIGMA/D-OMEGA UB/SR (2,3)		NO. EVENTS
MIN	MAX			
-1.00	-.95	.00	.01	0
-.95	-.90	.01	.01	1
-.90	-.85	.01	.01	1
-.85	-.80	.03	.02	2
-.80	-.75	.00	.01	0
-.75	-.70	.01	.01	1
-.70	-.65	.01	.01	1
-.65	-.60	.03	.02	2
-.60	-.55	.00	.01	0
-.55	-.50	.01	.01	1
-.50	-.45	.00	.01	0
-.45	-.40	.00	.01	0
-.40	-.35	.00	.01	0
-.35	-.30	.01	.01	1
-.30	-.25	.00	.01	0
-.25	-.20	.00	.01	0
-.20	-.15	.01	.01	1
-.15	-.10	.00	.01	0
-.10	-.05	.01	.01	1
-.05	.00	.01	.01	1
.00	.05	.03	.02	2
.05	.10	.00	.01	0
.10	.15	.00	.01	0
.15	.20	.03	.02	2
.20	.25	.01	.01	1
.25	.30	.03	.02	2
.30	.35	.04	.02	3
.35	.40	.03	.02	2
.40	.45	.00	.01	0
.45	.50	.01	.01	1
.50	.55	.04	.02	3
.55	.60	.03	.02	2
.60	.65	.05	.03	4
.65	.70	.08	.03	6
.70	.75	.12	.04	9
.75	.80	.18	.04	13
.80	.85	.27	.06	20
.85	.90	.43	.08	31
.90	.95	.49	.08	36
.95	1.00	.45	.08	33

THETA IS THE ANGLE THAT THE $K^*(890)^+$ MAKES WITH THE BEAM IN THE GRAND C.M.

\{1\} OVERLAP REGION EXCLUDED. COUNTED ONLY EVENTS ABOVE BACKGROUND [MASS = .900 GEV; WIDTH = .080 GEV].

\{2\} PLUS POSSIBLE SYSTEMATIC ERROR OF $\pm$ 10 PER CENT.

\{3\} COUNTS WERE MULTIPLIED BY .01375 TO GET THESE. ERRORS ARE TAKEN AS PROPORTIONAL TO THE SQUARE-ROOT OF THE COUNTS .

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \Delta^{++}$.
 $\Delta^{++} \rightarrow p \pi^+$ PROTON (1)
 LABORATORY BEAM MOMENTUM = 2.965 $\pm$.015 GEV/C.

 . THIS DATA WAS READ FROM A GRAPH .

COS(THETA)		D-SIGMA/D-OMEGA UB/SR [2,3]		NO. EVENTS
MIN	MAX			
.95	1.00	.33 $\pm$.07		24
.90	.95	.48 .08		35
.85	.90	.48 .08		35
.80	.85	.41 .08		30
.75	.80	.19 .05		14
.70	.75	.15 .05		11
.65	.70	.07 .03		5
.60	.65	.07 .03		5
.55	.60	.03 .02		2
.50	.55	.05 .03		4
.45	.50	.03 .02		2
.40	.45	.00 .01		0
.35	.40	.00 .01		0
.30	.35	.00 .01		0
.25	.30	.00 .01		0
.20	.25	.00 .01		0
.15	.20	.00 .01		0
.10	.15	.01 .01		1
.05	.10	.00 .01		0
.00	.05	.01 .01		1
-.05	.00	.00 .01		0
-.10	-.05	.01 .01		1
-.15	-.10	.03 .02		2
-.20	-.15	.01 .01		1
-.25	-.20	.00 .01		0
-.30	-.25	.00 .01		0
-.35	-.30	.01 .01		1
-.40	-.35	.00 .01		0
-.45	-.40	.00 .01		0
-.50	-.45	.00 .01		0
-.55	-.50	.01 .01		1
-.60	-.55	.01 .01		1
-.65	-.60	.00 .01		0
-.70	-.65	.01 .01		1
-.75	-.70	.00 .01		0
-.80	-.75	.01 .01		1
-.85	-.80	.00 .01		0
-.90	-.85	.00 .01		0
-.95	-.90	.03 .02		2
-1.00	-.95	.00 .01		0

THETA IS THE ANGLE THAT THE K^0 MAKES WITH THE BEAM IN THE GRAND C.M.

- (1) COUNTED ALL EVENTS IN MASS BAND [MASS = 1.225 GEV; WIDTH = .130 GEV]. OVERLAP REGION EXCLUDED.
 (2) PLUS POSSIBLE SYSTEMATIC ERROR OF $\pm$ 10 PER CENT.
 (3) COUNTS WERE MULTIPLIED BY .01375 TO GET THESE. ERRORS ARE TAKEN AS PROPORTIONAL TO THE SQUARE-ROOT OF THE COUNTS .

71 PROTON POLARIZATION IN $K^+ p$ ELASTIC SCATTERING AT 780 MEV/C. [NUOVO CIMENTO 50A, 372 (1967)]

S.FEMINO, S.JANNELLI, F.MEZANARES (INS. DI FISICA DELL UNIV., MESSINA, ITALY)
 L.MONARI, P.SERRA (UNIV. DI BOLOGNA, BOLOGNA, ITALY)

ADDITIONAL FOOTNOTE REFERENCES

PHYS. LETTERS 24, 8314 (1967), ANNUAL REV. OF NUCLEAR SCIENCE 10, 291 (1960), PHYS. REV. 106, 564 (1957), UCRL 10813 (1964), PHYS. REV. 135, 8191 (1964), AND NUOVO CIMENTO 44, 879 (1965).

ARTICLE READ BY ODETTE BENARY IN 5/69. AND VERIFIED BY LEROY PRICE.

BEAM IS K^+ ON PROTON AT .78 GEV/C.

THIS EXPERIMENT USES THE SACLAY 81 CM (H) BUBBLE CHAMBER. A TOTAL OF 120000 PICTURES ARE REPORTED ON.

KEY WORDS = POLARIZATION

ELASTIC POLARIZATION FOR $K^+ \text{ PROTON}$. [TABLE 1]

LABORATORY BEAM MOMENTUM = .78 GEV/C.
 NUMBER OF EVENTS = 225.

(THE SAMPLE CONSISTS ONLY FROM ELASTIC EVENTS IN WHICH THE PROTON IS GOING FORWARD IN THE $K^+ p$ C.M.S. WE HAVE CHANGED THE SIGN OF THE POLARIZATION TO CONFORM TO THE BASEL CONVENTION)

COS(THETA)		POLARIZATION		NO. EVENTS
MIN	MAX			
-1.00	-.40	.32 $\pm$.40		146
-.40	.25	.21 .63		109

THETA IS THE ANGLE THAT THE K^+ MAKES WITH THE BEAM IN THE GRAND C.M.
 THE POLARIZATION IS OF THE PROTON ALONG THE NORMAL TO THE PRODUCTION PLANE IN THE GRAND C.M.

72 THE K+ P INTERACTION FROM 964 TO 1585 MEV/C: CROSS SECTIONS AND MASS DISTRIBUTIONS. [UCRL 18798 (1968)]

R.W.BLAND, M.G.BOWLER, J.L.BROWN, J.A.KADYK, G.GOLDHABER, S.GOLDHABER, V.H.SEEGER, G.H.TRILLING (U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA)

ABSTRACT IN THIS PAPER WE PRESENT, FOR FIVE MOMENTA FROM 864 TO 1585 MEV/C, CROSS SECTIONS FOR STABLE FINAL STATES AND FOR RESONANCE PRODUCTION, AND AN ANALYSIS OF THE K⁺ N⁺ P⁺ DALITZ PLOTS INCLUDING THE EFFECTS OF K*(891)-DELTA(1238) INTERFERENCE. WE SEE STRONG INTERFERENCE AT ALL MOMENTA WHERE THE K⁺ IS PRESENT, WITH THE RELATIVE PHASE OF THE K⁺ N⁺ AND K-DELTA AMPLITUDES REMAINING CONSTANT AS A FUNCTION OF MOMENTUM. WE COMBINE OUR CROSS-SECTION RESULTS WITH THOSE FROM OTHER EXPERIMENTS TO STUDY THE MOMENTUM DEPENDENCE OF THE PARTIAL CROSS SECTIONS IN THE REGION OF THE 1250 MEV/C PEAK IN THE TOTAL CROSS SECTION; WE FIND THAT THE TOTAL CROSS-SECTION CURVE CAN BE REPRESENTED AS THE SUM OF PARTIAL CROSS SECTION CURVES, EACH ONE SMOOTH AND WITHOUT A PEAK CORRESPONDING TO THAT IN THE TOTAL CROSS SECTION. THUS THIS STRUCTURE IS NOT DUE TO STRUCTURE IN ANY SINGLE PARTIAL CROSS SECTION, BUT RATHER TO THE SHARP RISES OF THE SINGLE- AND DOUBLE-PION PRODUCTION CHANNELS AT WIDELY SEPARATED THRESHOLDS. THIS INTERPRETATION IS AT VARIANCE WITH A CONVENTIONAL RESONANCE INTERPRETATION OF THE OBSERVED STRUCTURE IN THE TOTAL CROSS SECTION.

CROSS REFERENCES

THIS ARTICLE SUPERSEDES PART OF UCRL 18131 (1958), DUBNA CONFERENCE 715 (1964), UCRL 16370 (1965), AND PHYS. REV. LETTERS 18, 1077 (1967).
CONTINUATION OF PREVIOUS EXPERIMENT IN PHYS. REV. LETTERS 17, 939 (1966).
ANALYSIS OF DATA FROM NUOVO CIMENTO 51A, 1053 (1967).
CONTAINS RENORMALIZATION OF DATA FROM NUOVO CIMENTO 49A, 348 (1967), PHYS. LETTERS 16, 83 (1965), AND PHYS. REV. LETTERS 9, 135 (1962).

ADDITIONAL FOOTNOTE REFERENCES

PHYS. REV. LETTERS 17, 102 (1966), PHYS. REV. LETTERS 19, 259 (1967), PHYS. REV. 168, 1456 (1968), UCRL 11527 (1964), UCRL 8030 JAN. (1968), PHYS. LETTERS 248, 314 (1967), PHYS. REV. LETTERS 7, 188 (1961), PHYS. REV. LETTERS 11, 93 (1963), PHYS. REV. 133, 2220 (1964), PHYS. LETTERS 16, 83 (1965), PHYS. LETTERS 6, 309 (1963), PHYS. REV. 142, 913 (1966), PHYS. LETTERS 6, 62 (1963), PHYS. REV. 158, 1281 (1967), PHYS. REV. 158, 1310 (1967), NUOVO CIMENTO 36, 1101 (1965), ARKIV FOR FYSIK 37, 463 (1968), NUOVO CIMENTO 39, 417 (1965), NUOVO CIMENTO 49A, 9 (1967), NUOVO CIMENTO 45A, 885 (1966), PHYS. REV. LETTERS 7, 182 (1961), NUCLEAR INSTRUMENTS AND METHODS 20, 393 (1963), PHYS. REV. LETTERS 15, 737 (1965), NUOVO CIMENTO 33, 926 (1964), NUOVO CIMENTO 43A, 142 (1966), AND NUOVO CIMENTO 51A, 401 (1967).

ARTICLE READ BY ODETTE BENARY AND LEROY PRICE IN 7/69, AND VERIFIED BY ROGER BLAND.

BEAM IS K⁺ ON PROTON FROM .864 TO 1.585 GEV/C.

THIS EXPERIMENT USES THE L.R.L. 25 IN. (H) RUBBLE CHAMBER.

GENERAL COMMENTS ON THIS ARTICLE

1 CROSS SECTIONS AT .864, .969, AND 1.207 GEV/C ARE NORMALIZED TO TOTAL CROSS SECTION DATA OF COOL (PHYS. REV. LETTERS 17, 102(1966)) AND BUGG (PHYS. REV. 168, 1466(1968)). CROSS SECTIONS AT 1.367 AND 1.585 GEV/C ARE NORMALIZED TO TAU DECAYS.

KEY WORDS = CROSS SECTION K*(890) DELTA(1238) MASS SPECTRUM FITS DALITZ PLOT

COMPOUND KEY WORDS = FITS DALITZ PLOT

K+ PROTON TOTAL CROSS SECTION.

(NORMALIZED TO TAU DECAYS.)

LABORATORY BEAM MOMENTUM

GEV/C	MILLI-BARNS
.864	14.07 +- .55
.969	15.28 .90
1.207	19.85 1.28

K+ PROTON ELASTIC CROSS SECTION. COULOMB CORRECTIONS APPLIED

LABORATORY BEAM MOMENTUM

GEV/C	MILLI-BARNS
.864	11.98 +- .27
.969	11.68 .21
1.207	10.89 .18

CROSS SECTION FOR K+ PROTON = K0 PROTON P1+.

LABORATORY BEAM MOMENTUM

GEV/C	MILLI-BARNS
.864	1.15 +- .06
.969	2.57 .12
1.207	5.04 .15
1.367	5.36 .33
1.585	5.00 .40

CROSS SECTION FOR K+ PROTON = K+ PROTON P10.

LABORATORY BEAM MOMENTUM

GEV/C	MILLI-BARNS
.864	.36 +- .03
.969	.80 .07
1.207	1.80 .10

CROSS SECTION FOR K+ PROTON = K+ NEUTRON P1+.

LABORATORY BEAM MOMENTUM

GEV/C	MILLI-BARNS
.864	.131 +- .017
.969	.300 .040
1.207	.560 .060

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K \text{ MESON NUCLEON PION.}$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.864	1.64 $\pm$.07
.969	3.67 .14
1.207	7.40 .17

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \text{ PROTON PI}^+ \text{ PI}^-.$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
1.207	.026 $\pm$.010
1.367	.065 .012
1.585	.380 .050

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \text{ PROTON PI}^+ \text{ PI}^0.$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
1.367	.060 $\pm$.018
1.585	.320 .070

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \text{ NEUTRON PI}^+ \text{ PI}^+.$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
1.367	.009 $\pm$.012
	- .006
1.585	.080 $\pm$.030
	- .030

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K \text{ MESON NUCLEON PION PION.}$ (ASSUMES K^*-N^* DOMINANCE.)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
1.207	.029 $\pm$.011
1.367	.147 .025
1.450	.280 .070 (1)
1.585	.860 .100
1.960	3.500 .300 (2)
2.650	5.500 .400 (3)
2.970	5.500 .700 (4)
3.460	5.300 .500 (5)

- (1) USING DATA FROM BETTINI, ET AL., PHYS. LETTERS 16, 83 (1965).
 (2) USING DATA FROM CHINOWSKY, ET AL., PHYS. REV. 139, B1411 (1965).
 (3) USING DATA FROM NEWMAN, ET AL., PHYS. REV. 158, 1310 (1967).
 (4) USING DATA FROM SALLSTROM, ET AL., NUOVO CIMENTO 49A, 348 (1967).
 (5) USING DATA FROM GEORGE, ET AL., NUOVO CIMENTO 49A, 9 (1967).

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \text{ PROTON PI}^+ \text{ PI}^- \text{ PI}^0.$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
1.585	.009 $\pm$.010
	- .005

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \text{ PROTON PI}^+ \text{ PI}^- \text{ PI}^0.$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
1.585	.004 $\pm$.015
	- .004

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K \text{ MESON } \Delta(1238),$
 $\Delta(1238) \rightarrow \text{NUCLEON PION (1)}$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.735	.11 $\pm$.08 (2)
.785	.25 $\pm$.05 (2)
.864	1.13 $\pm$.15
.969	3.09 $\pm$.23
1.207	4.56 $\pm$.35
1.367	4.10 $\pm$.40
1.585	2.70 $\pm$.40

(1) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND. NON-INTERFERING AMPLITUDES ASSUMED.

(2) USING DATA FROM FILIPPAS, ET AL., NUOVO CIMENTO 51A, 1053 (1967).

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890) \text{ NUCLEON},$
 $K^*(890) \rightarrow K \text{ MESON PION (1)}$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
1.207	1.74 $\pm$.23
1.367	2.60 $\pm$.30
1.585	3.20 $\pm$.40

(1) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND. NON-INTERFERING AMPLITUDES ASSUMED.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \Delta^{++},$
 $\Delta^{++} \rightarrow \text{PROTON PION (1)}$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.735	.08 $\pm$.06 (2)
.785	.19 $\pm$.04 (2)
.864	.86 $\pm$.12
.969	2.42 $\pm$.19
1.207	3.43 $\pm$.32
1.367	3.05 $\pm$.33
1.585	2.05 $\pm$.34

(1) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND. NON-INTERFERING AMPLITUDES ASSUMED.

(2) USING DATA FROM FILIPPAS, ET AL., NUOVO CIMENTO 51A, 1053 (1967).

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)^+ \text{ PROTON},$
 $K^*(890)^+ \rightarrow K^0 \text{ PION (1)}$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
1.207	1.06 $\pm$.20
1.367	1.72 $\pm$.24
1.585	2.13 $\pm$.35

(1) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND. NON-INTERFERING AMPLITUDES ASSUMED.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \text{ PION PROTON},$
 CROSS SECTION IS ONLY FOR THE NON-RESONANT PRODUCTION OF THESE PARTICLES.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.735	.05 $\pm$.06 (1)
.785	.15 $\pm$.04 (1)
.864	.29 $\pm$.12
.969	.15 $\pm$.15
1.207	.25 $\pm$.30

(1) USING DATA FROM FILIPPAS, ET AL., NUOVO CIMENTO 51A, 1053 (1967).

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \Delta^{++},$
 $\Delta^{++} \rightarrow \text{PROTON PION (1)}$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.864	.19 $\pm$.08
.969	.43 $\pm$.11
1.207	.81 $\pm$.14

(1) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND. NON-INTERFERING AMPLITUDES ASSUMED.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \Delta^{++}$
 $\Delta^{++} \rightarrow \pi^+ \text{ NEUTRON} [1]$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.864	.08 $\pm$.04
.969	.24 $\pm$.07
1.207	.32 $\pm$.06

[1] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND. NON-INTERFERING AMPLITUDES ASSUMED.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)^+ \text{ PROTON}$
 $K^*(890)^+ \rightarrow K^+ \pi^0 [1]$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
1.207	.55 $\pm$.11

[1] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND. NON-INTERFERING AMPLITUDES ASSUMED.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \pi^0 \text{ PROTON}$
 CROSS SECTION IS ONLY FOR THE NON-RESONANT PRODUCTION OF THESE PARTICLES.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.864	.17 $\pm$.08
.969	.37 $\pm$.11
1.207	.20 $\pm$.14

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \pi^+ \text{ NEUTRON}$
 CROSS SECTION IS ONLY FOR THE NON-RESONANT PRODUCTION OF THESE PARTICLES.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.864	.05 $\pm$.04
.969	.06 $\pm$.06
1.207	.24 $\pm$.06

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890) \Delta^{++}$
 $\Delta^{++} \rightarrow \pi^+ \text{ PROTON} [1]$
 $K^*(890) \rightarrow K^+ \pi^- [2]$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
1.585	.25 $\pm$.07

[1] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND. NON-INTERFERING AMPLITUDES ASSUMED.
 [2] USED SIMPLE MASS CUT.

73

$K^+ \text{ PROTON ELASTIC SCATTERING AT } 910 \text{ MEV/C: POLARIZATION OF THE RECOIL PROTONS. [PHYS. REV. 135, 8191 (1964)]$

W. HIRSCH, G. GIDAL (U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA, AND UNIV. OF CALIFORNIA, BERKELEY, CALIF., USA)

ABSTRACT THE BERKELEY 30-IN. PROPANE BUBBLE CHAMBER WAS USED TO STUDY THE ELASTIC $K^+ \text{ PROTON}$ INTERACTION AT 910 MEV/C. THIS IS THE REGION OF TRANSITION FROM THE ISOTROPY IN ANGULAR DISTRIBUTION FOUND BELOW 810 MEV/C TO THE RAPIDLY INCREASING ANISOTROPY ABOVE 1 BEV/C. RESULTS BASED ON 1154 EVENTS SHOW THAT THE SERIES $(1 + A \cos(\theta_{\text{CM}}))$ CAN FIT THE ANGULAR DISTRIBUTION WITH $A = 0.18 \pm 0.05$. POLARIZATION WAS MEASURED ON THE SECONDARY PROTON BY USING 53 PROTON-PROTON AND 41 PROTON-CARBON RECOILS IN THE LIQUID OF THE CHAMBER. THE RESULTS OF A PHASE-SHIFT ANALYSIS INCORPORATING THESE POLARIZATION DATA ARE PRESENTED.

ADDITIONAL FOOTNOTE REFERENCES

UCRL 10813 (1963), PHYS. REV. LETTERS 9, 135 (1962), PHYS. REV. 118, 553 (1960), PHYS. REV. LETTERS 7, 188 (1961), UCRL 10130 (1962), PHYS. REV. 129, 2743 (1963), REVIEW OF SCIENTIFIC INSTRUMENTS 29, 874 (1958), LRL INTERDEPARTMENTAL REPORT BEV-483 (1960), PHYS. REV. LETTERS 5, 254 (1960), UCRL 10622, PHYS. REV. LETTERS 10, 27 (1963), PHYS. REV. LETTERS 11, 94 (1963), BULL. AM. PHYS. SOC. 8, 21 (1963), UCRL 10813 (1963), UCRL 9191 (1960), AND PHYS. REV. LETTERS 11, 95 (1963).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 3/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K^+ ON HEAVY LIQUID AT .91 GEV/C.

THIS EXPERIMENT USES THE L.R.L. 30 IN. (HL) BUBBLE CHAMBER. A TOTAL OF 42500 PICTURES ARE REPORTED ON.

KEY WORDS = POLARIZATION ANGULAR DISTRIBUTION PHASE SHIFT

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = .91 ± .10 GEV/C.

.....
 . THIS DATA WAS READ FROM A GRAPH .

COS(THETA)		NO. EVENTS
MIN	MAX	
-1.0	-.8	116
-.8	-.6	106
-.6	-.4	112
-.4	-.2	100
-.2	.0	109
.0	.2	127
.2	.4	134
.4	.6	138
.6	.8	129
.8	.9	153

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.ELASTIC POLARIZATION FOR K⁺ PROTON.

LABORATORY BEAM MOMENTUM = .91 ± .10 GEV/C.

(WE HAVE CHANGED TO SIGN OF THE POLARIZATION TO CONFORM WITH THE BASEL CONVENTION.)

THETA DEGREES		POLARIZATION
MIN	MAX	
40.	70.	.80 ± .80
70.	100.	.74 .45
100.	140.	-.55 .93
140.	160.	-.70 .93

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.
 THE POLARIZATION IS OF THE PROTON ALONG THE NORMAL TO THE PRODUCTION PLANE IN THE GRAND C.M.

74 SEARCH FOR BARYONIC RESONANCES WITH S = +1. IN K⁺P PRODUCTION EXPERIMENTS. [PHYS. LETTERS 278, 468 (1968)]

G. BASSOMPIERRE, Y. GOLDSCHMIDT-CLERMONT, A. GRANT, V. P. HENRI, R. JENNINGS, B. JONGEJANS, D. LINGLIN, F. MULLER, J. M. PERREAU, R. L. SEKULIN
 (EUROPEAN ORG. FOR NUC. RES., GENEVA, SWITZERLAND)
 W. DEBAERE, J. DEBAISIEUX, P. DUFODUR, F. GRARD, J. HEUGHEBAERT, L. PAPE, P. PEETERS, F. VERBEURE, R. WINDMOLDERS (INST. INTERUNIV. DES SCI.
 NUC., BRUXELLES, BELGIUM)

ABSTRACT IN ABOUT 30000 IDENTIFIED EVENTS PRODUCED IN K⁺P REACTIONS AT INCIDENT MOMENTA 3.0, 3.5 AND 5.0 GEV/C WE HAVE SEARCHED FOR THE PRODUCTION OF Z⁰S, I.E. S = +1, T = 0 OR 1 BARYONIC RESONANCES. NO INDICATION HAS BEEN FOUND IN THE EXAMINATION OF THE VARIOUS AVAILABLE PRODUCTION REACTIONS OF Z⁰ DECAYING WITH VISIBLE MODES. UPPER LIMITS FOR THE CORRESPONDING CROSS-SECTIONS ARE GENERALLY LESS THAN 30 MICROBARN AT 3.0 AND 3.5 GEV/C, 10 MICROBARN AND 5.0 GEV/C. SUMMING FOR A GIVEN PRODUCTION REACTION OVER ALL POSSIBLE Z⁰ DECAY MODES, I.E. PERFORMING A MISSING MASS TYPE OF ANALYSIS, DOES NOT REVEAL ANY FURTHER EVIDENCE.

CROSS REFERENCES

THIS ARTICLE SUPERSEDES PART OF PHYS. LETTERS 12, 255 (1964), NUOVO CIMENTO 49A, 9 (1967), AND PHYS. LETTERS 26A, 30 (1967).
 CONTINUATION OF PREVIOUS EXPERIMENT IN NUOVO CIMENTO 39, 417 (1965), AND NUOVO CIMENTO 36, 1102 (1965).

ADDITIONAL FOOTNOTE REFERENCES

PHYS. REV. LETTERS 17, 102 (1966), HEIDELBERG CONFERENCE 123 (1968), PHYS. REV. LETTERS 19, 259 (1967), NUOVO CIMENTO 46, 539 (1966), NUOVO CIMENTO 48, 589 (1967), UCRL 17388, BULL. AM. PHYS. SOC. 12, 4A (1967), PHYS. REV. LETTERS 18, 1077 (1967), PHYS. REV. 163, 1770 (1968), PHYS. REV. LETTERS 19, 253 (1967), AND HEIDELBERG CONFERENCE 117 (1968).

ARTICLE READ BY DOETTE BENARY IN 12/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K⁺ ON PROTON FROM 3 TO 5 GEV/C.

THIS EXPERIMENT USES BUBBLE CHAMBERS.

GENERAL COMMENTS ON THIS ARTICLE

1 THIS ARTICLE REPEATS THE CROSS SECTIONS FOR K⁺P → KO P PI⁺, K⁺P PI⁺ PI⁻, KO N PI⁺ PI⁺, AND KO P PI⁺ PI⁰ AT 3 GEV/C.
 = THE SAME VALUES WERE PREVIOUSLY PUBLISHED IN PL 12, 255 (1964).

KEY WORDS = CROSS SECTION MASS SPECTRUM Z⁰CROSS SECTION FOR K⁺ PROTON → KO PROTON PI⁺. [TABLE 21]

LABORATORY BEAM MOMENTUM	
GEV/C	MILLI-BARNS
3.5	1.85 ± .10
5.0	.90 ± .10

CROSS SECTION FOR K⁺ PROTON → K⁺ PROTON PI⁰. [TABLE 21]

LABORATORY BEAM MOMENTUM	
GEV/C	MILLI-BARNS
3.5	1.32 ± .07

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \text{ NEUTRON } \pi^+ \pi^-$ [TABLE 2]

(COMMENTS IN DEBAERE, NC 51A, 401 (1967) SUGGEST THAT THIS ERROR SHOULD BE MORE NEARLY 0.4 MB.)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
3.5	.8 $\pm$.1

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \text{ MESON NUCLEON } \pi^0$.

(THIS IS THE SUM OF THE ABOVE THREE REACTIONS.)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
3.5	4.0 $\pm$.2

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \text{ PROTON } \pi^+ \pi^-$ [TABLE 2]

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
3.5	2.10 $\pm$.30
5.0	1.80 .15 [1]

[1] THIS DATA REPLACES THE VALUE GIVEN EARLIER IN BASSOMPIERRE, ET.AL., PHYS. LETTERS 268, 30 (1967).

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \text{ NEUTRON } \pi^+ \pi^-$ [TABLE 2]

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
3.5	.60 $\pm$.10
5.0	.40 .05 [1]

[1] THIS DATA REPLACES THE VALUE GIVEN EARLIER IN BASSOMPIERRE, ET.AL., PHYS. LETTERS 268, 30 (1967).

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \text{ PROTON } \pi^+ \pi^0$ [TABLE 2]

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
3.5	2.5 $\pm$.3
5.0	1.5 .2 [1]

[1] THIS DATA REPLACES THE VALUE GIVEN EARLIER IN BASSOMPIERRE, ET.AL., PHYS. LETTERS 268, 30 (1967).

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \text{ PROTON } \pi^+ \pi^- \pi^0$ [TABLE 2]

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
3.0	.35 $\pm$.04 [1]
3.5	.44 .04
5.0	.62 .08

[1] THIS DATA REPLACES THE VALUE GIVEN EARLIER IN FERRO-LUZZI, ET.AL., PHYS. LETTERS 12, 255 (1964).

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \text{ PROTON } \pi^+ \pi^- \pi^0$ [TABLE 2]

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
3.0	.62 $\pm$.05 [1]
3.5	.71 .04
5.0	1.10 .10

[1] THIS DATA REPLACES THE VALUE GIVEN EARLIER IN FERRO-LUZZI, ET.AL., PHYS. LETTERS 12, 255 (1964).

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \text{ NEUTRON } \pi^+ \pi^- \pi^0$ [TABLE 2]

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
3.0	.15 $\pm$.03 [1]
3.5	.19 .02
5.0	.31 .03

[1] THIS DATA REPLACES THE VALUE GIVEN EARLIER IN FERRO-LUZZI, ET.AL., PHYS. LETTERS 12, 255 (1964).

75 INELASTIC K⁺ PROTON INTERACTIONS AT 2.65 BEV/C. [PHYS. REV. 158, 1310 (1967)]

R. NEWMAN, W. CHINOWSKY, J. SCHULTZ (U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA, AND UNIV. OF CALIFORNIA, BERKELEY, CALIF., USA)
W. B. JOHNSON, R. R. LARSEN (STANFORD LINEAR ACCEL. CNTR., STANFORD, CALIF., USA)

ABSTRACT PION PRODUCTION BY 2.65 BEV/C K⁺ MESONS INTERACTING IN THE RNL 20-IN. HYDROGEN BUBBLE CHAMBER HAS BEEN INVESTIGATED. IT IS FOUND THAT THESE INELASTIC EVENTS ARE DOMINATED BY K*(891) AND N*(1236) PRODUCTION. BASED ON 537 THREE- AND FOUR-BODY EVENTS, THE CROSS SECTIONS FOR PRODUCTION OF K⁰ P PI⁺ AND K⁺ P PI⁺ PI⁻ ARE 2.7 ± 0.3 AND 2.5 ± 0.2 MB, RESPECTIVELY. THE THREE-BODY FINAL STATES CONTAIN 56 PER CENT K*P AND 32 PERCENT N*KO, WHILE THE FOUR-BODY STATES CONTAIN 9 PER CENT K*P PI⁺, 19 PER CENT N*K PI⁻, AND 53 PER CENT K* N*. RESONANCE PRODUCTION IS COMPARED WITH PREDICTIONS OF SEVERAL VERSIONS OF THE ONE-MESON-EXCHANGE MODEL. ALTHOUGH AGREEMENT WITH THE DATA IS NOT EXACT, THE MODELS TEND TO REPRODUCE A NUMBER OF QUALITATIVE FEATURES OF THE EXPERIMENTAL RESULTS.

ADDITIONAL FOOTNOTE REFERENCES

NUOVO CIMENTO 14, 951 (1959), PHYS. REV. LETTERS 7, 394 (1961), PHYS. REV. LETTERS 8, 41 (1962), NUCLEAR INSTRUMENTS AND METHODS 20, 37 (1963), NUCLEAR INSTRUMENTS AND METHODS 23, 42 (1963), ANNUAL REV. OF NUCLEAR SCIENCE 13, 103 (1965), UCRL 9099 (1961), UCRL 16473, PHYS. REV. 139, B1097 (1965), PHYS. REV. LETTERS 10, 351 (1963), NUOVO CIMENTO 33, 906 (1964), NUOVO CIMENTO 34, 1841E (1964), PHYS. REV. LETTERS 15, 737 (1965), PHYS. REV. LETTERS 12, 522 (1964), NUOVO CIMENTO 36, 1101 (1965), PHYS. REV. 142, 913 (1966), PHYS. LETTERS 6, 52 (1963), NUOVO CIMENTO 39, 417 (1965), UCRL 16370 (1965), PHYS. REV. 133, B220 (1964), PHYS. LETTERS 6, 309 (1963), PHYS. REV. LETTERS 11, 93 (1963), PHYS. REV. 139, B428 (1965), NUOVO CIMENTO 39, 941 (1965), PHYS. REV. 145, 1072 (1966), PHYS. REV. LETTERS 7, 182 (1961), PHYS. REV. 139, B1411 (1965), NUOVO CIMENTO 43, A142 (1966), NUOVO CIMENTO 42, A323 (1966), PHYS. REV. LETTERS 17, 102 (1966), PHYS. LETTERS 9, 71 (1964), AND PHYS. LETTERS 15, 74 (1965).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 5/68, AND VERIFIED BY ROGER BLAND.

BEAM IS K⁺ ON PROTON AT 2.65 GEV/C.

THIS EXPERIMENT USES THE B.N.L. 20 IN. (H) BUBBLE CHAMBER. A TOTAL OF 50000 PICTURES ARE REPORTED ON.

GENERAL COMMENTS ON THIS ARTICLE

1 ASSUMES K⁺ LIFETIME = (1.234 ± .004) X 10⁻⁸ SEC.

KEY WORDS * K*(891) DELIA(1236) CROSS SECTION DALITZ PLOT MASS SPECTRUM ANGULAR DISTRIBUTION
DENSITY MATRIX

CROSS SECTION FOR K⁺ PROTON → K⁰ PROTON PI⁺.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.65 ± .03	2.7 ± .3

CROSS SECTION FOR K⁺ PROTON → K⁰ PROTON PI⁺ PI⁰.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.65 ± .03	1.9 ± .2

CROSS SECTION FOR K⁺ PROTON → K⁰ PI⁺ PI⁺ NEUTRON.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.65 ± .03	.64 ± .12

CROSS SECTION FOR K⁺ PROTON → K⁺ PROTON PI⁺ PI⁻.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.65 ± .03	2.5 ± .2

CROSS SECTION FOR K⁺ PROTON → K⁺ PROTON PI⁺ PI⁻ PI⁰.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.65 ± .03	.43 ± .08

CROSS SECTION FOR K⁺ PROTON → K⁺ PI⁺ PI⁺ PI⁻ NEUTRON.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.65 ± .03	.14 ± .03

CROSS SECTION FOR K⁺ PROTON → PROTON PI⁺ PI⁺ PI⁻ K⁰.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.65 ± .03	.19 ± .05

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow \text{PROTON } \pi^+ \pi^+ \pi^- K^0 \pi^0$.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.65 $\pm$.03	.04 $\pm$.04

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \text{ PROTON } \pi^+ \pi^+ \pi^- \pi^-$.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.65 $\pm$.03	.01 $\pm$.01

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ K^0 \text{ SIGMA}^+$.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.65 $\pm$.03	.02 $\pm$.02

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ K^+ \text{ SIGMA}^0$.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.65 $\pm$.03	.01

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ K^+ \text{ LAMBDA}$.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.65 $\pm$.03	.03 $\pm$.02

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ K^+ \text{ LAMBDA } \pi^0$.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.65 $\pm$.03	.01 $\pm$.01

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 K^+ \pi^+ \text{ LAMBDA}$.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.65 $\pm$.03	.03 $\pm$.03

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)^+ \text{ PROTON}$,
 $K^*(890)^+ \rightarrow K^0 \pi^+ [1]$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.65 $\pm$.03	1.5 $\pm$.3

[1] COUNTED ONLY EVENTS ABOVE BACKGROUND [MASS = .891 GEV; WIDTH = .350 GEV].

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow \text{DELTA}^{++} K^0$,
 $\text{DELTA}^{++} \rightarrow \text{PROTON } \pi^+ [1]$

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
2.65 $\pm$.03	.9 $\pm$.2

[1] COUNTED ONLY EVENTS ABOVE BACKGROUND [MASS = 1.236 GEV; WIDTH = .120 GEV].

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)0 \text{ DELTA}^{++}$
 $K^*(890)0 \rightarrow K^+ \text{ PI}^-$ [1]
 $\text{DELTA}^{++} \rightarrow \text{PROTON PI}^+$ [2]

LABORATORY
 BEAM MOMENTUM
 GEV/C
 2.65 $\pm$.03
 MILLI-BARNS
 1.33 $\pm$.15

[1] COUNTED ONLY EVENTS ABOVE BACKGROUND [MASS = .891 GEV; WIDTH = .050 GEV].
 [2] COUNTED ONLY EVENTS ABOVE BACKGROUND [MASS = 1.236 GEV; WIDTH = .120 GEV].

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)0 \text{ PROTON PI}^+$
 $K^*(890)0 \rightarrow K^+ \text{ PI}^-$ [1]

LABORATORY
 BEAM MOMENTUM
 GEV/C
 2.65 $\pm$.03
 MILLI-BARNS
 .23 $\pm$.10

[1] COUNTED ONLY EVENTS ABOVE BACKGROUND [MASS = .891 GEV; WIDTH = .050 GEV].

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow \text{DELTA}^{++} K^+ \text{ PI}^-$
 $\text{DELTA}^{++} \rightarrow \text{PROTON PI}^+$ [1]

LABORATORY
 BEAM MOMENTUM
 GEV/C
 2.65 $\pm$.03
 MILLI-BARNS
 .48 $\pm$.11

[1] COUNTED ONLY EVENTS ABOVE BACKGROUND [MASS = 1.236 GEV; WIDTH = .120 GEV].

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \text{ PROTON PI}^+ \text{ PI}^-$
 CROSS SECTION IS ONLY FOR THE NON-RESONANT PRODUCTION OF THESE PARTICLES.

LABORATORY
 BEAM MOMENTUM
 GEV/C
 2.65 $\pm$.03
 MILLI-BARNS
 .48 $\pm$.11

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)0 \text{ DELTA}^{++}$
 $K^*(890)0 \rightarrow K^+ \text{ PI}^-$ [1]
 $\text{DELTA}^{++} \rightarrow \text{PROTON PI}^+$ [2]

LABORATORY BEAM MOMENTUM = 2.65 $\pm$.03 GEV/C.

 . THIS DATA WAS READ FROM A GRAPH .

COS(THETA)		D-SIGMA/D-OMEGA UB/SR [3]		NO. EVENTS
MIN	MAX			
-1.0	-.9	.03 $\pm$.02		2
-.9	-.8	.01 .01		1
-.8	-.7	.00 .01		0
-.7	-.6	.04 .03		3
-.6	-.5	.01 .01		1
-.5	-.4	.01 .01		1
-.4	-.3	.00 .01		0
-.3	-.2	.00 .01		0
-.2	-.1	.00 .01		0
-.1	.0	.01 .01		1
.0	.1	.03 .02		2
.1	.2	.04 .03		3
.2	.3	.04 .03		3
.3	.4	.01 .01		1
.4	.5	.07 .03		5
.5	.6	.07 .03		5
.6	.7	.10 .04		7
.7	.8	.25 .06		17
.8	.9	.53 .09		36
.9	1.0	.92 .12		62

THETA IS THE ANGLE THAT THE $K^*(890)0$ MAKES WITH THE BEAM IN THE GRAND C.M.

[1] COUNTED ALL EVENTS IN MASS BAND [MASS = .890 GEV; WIDTH = .040 GEV].
 [2] COUNTED ALL EVENTS IN MASS BAND [MASS = 1.220 GEV; WIDTH = .120 GEV].
 [3] COUNTS WERE MULTIPLIED BY .0148 TO GET THESE. ERRORS ARE TAKEN AS PROPORTIONAL TO THE SQUARE-ROOT OF THE COUNTS.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \text{ PROTON OMEGA MESON}$
 $\text{OMEGA MESON} \rightarrow \text{PI}^+ \text{ PI}^- \text{ PI}^0$ [1]

LABORATORY
 BEAM MOMENTUM
 GEV/C
 2.65 $\pm$.03
 MILLI-BARNS
 .07 $\pm$.03

[1] USED SIMPLE MASS CUT.

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} = K^*(890)^+ \text{ PROTON}$
 $K^*(890)^+ = K^0 \text{ P}^+ [1]$
 LABORATORY BEAM MOMENTUM = $2.65 \pm .03 \text{ GEV/C}$.

.....
 : THIS DATA WAS READ FROM A GRAPH :

COS(THETA)		D-SIGMA/D-OMEGA UB/SR (2)		NO. EVENTS
MIN	MAX			
-1.0	-.9	.08	±.06	2.0
-.9	-.8	.08	±.06	2.0
-.8	-.5	.00	.04	.0
-.5	-.4	.04	.04	1.0
-.4	.3	.00	.04	.0
.3	.4	.12	.07	2.9
.4	.5	.00	.04	.0
.5	.6	.10	.07	2.4
.6	.7	.10	.06	2.3
.7	.8	.53	.15	12.6
.8	.9	.47	.14	11.1
.9	1.0	.88	.19	20.8

THETA IS THE ANGLE THAT THE $K^*(890)^+$ MAKES WITH THE BEAM IN THE GRAND C.M.

[1] OVERLAP REGION EXCLUDED.

[2] COUNTS WERE MULTIPLIED BY .0423 TO GET THESE. ERRORS ARE TAKEN AS PROPORTIONAL TO THE SQUARE-ROOT OF THE COUNTS.

76 PHASE SHIFT ANALYSIS OF K^+P ELASTIC SCATTERING AT 780 MEV/C. (PHYS. LETTERS 248, 314 (1967))

S.FOCARDI, A.MINGUZZI-RANZI, L.MONARI, G.SALTINI, P.SERRA (UNIV. DI BOLOGNA, BOLOGNA, ITALY, AND ISTITUTO DI FISICA DELLA
 UNIVERSITA, MODENA, ITALY)
 T.A.FILIPPAS, V.P.MENRI (EUROPEAN ORG. FOR NUC. RES., GENEVA, SWITZERLAND)

ABSTRACT A PHASE SHIFT ANALYSIS OF THE K^+P ELASTIC SCATTERING AT 780 MEV/C HAS BEEN PERFORMED. THE EXPERIMENTAL
 DIFFERENTIAL CROSS SECTION IS BEST EXPLAINED BY A SOLUTION WITH DOMINANT S WAVE, NEGATIVE S WAVE PHASE SHIFT (-42.7
 ± 1 DEG.) AND SMALL CONTRIBUTIONS OF P AND D WAVES.

ADDITIONAL FOOTNOTE REFERENCES

PHYS. REV. LETTERS 9, 135 (1962); PHYS. REV. LETTERS 7, 188 (1961); PHYS. REV. 137, B1115 (1965); UCRL 8030 AUG. (1966),
 AND NUOVO CIMENTO 45, 792 (1966).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 1/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K^+ ON PROTON AT .778 GEV/C.

THIS EXPERIMENT USES THE SACLAY 81 CM (H) BUBBLE CHAMBER. A TOTAL OF 45000 PICTURES ARE REPORTED ON.

GENERAL COMMENTS ON THIS ARTICLE

- 1 K^+ LIFETIME TAKEN AS $1.235 \pm .003$ ($\times 10^{-8}$ SEC)
- 2 K^+ TO τ BRANCHING RATIO TAKEN AS 0.057 ± 0.003 .

KEY WORDS = CROSS SECTION PHASE SHIFT

K^+ PROTON ELASTIC CROSS SECTION.

(CONTAINS SOME COULOMB EFFECTS. EXTRAPOLATED FROM COS(THETA) = .95 TO 1.0.)

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS
.778 ± .015	12.6 ± .4	9640

K^+ PROTON TOTAL INELASTIC CROSS SECTION.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS
.778 ± .015	.52 ± .03

ELASTIC DIFFERENTIAL CROSS SECTION FOR K^+ PROTON.

LABORATORY BEAM MOMENTUM = .778 ± .015 GEV/C.

COS(THETA)		D-SIGMA/D-OMEGA MB/3A	
MIN	MAX		
-1.00	-.90	1.137	±.064
-.90	-.80	1.073	.061
-.80	-.70	1.055	.060
-.70	-.60	.993	.057
-.60	-.50	1.137	.064
-.50	-.40	.984	.058
-.40	-.30	.885	.054
-.30	-.20	.929	.056
-.20	-.10	.929	.056
-.10	.00	.912	.055
.00	.10	.893	.055
.10	.20	.887	.055
.20	.30	.848	.053
.30	.40	.970	.058
.40	.50	.876	.055
.50	.60	1.064	.062
.60	.70	1.006	.060
.70	.80	1.059	.063
.80	.90	1.134	.065
.90	.95	1.375	.095

THETA IS THE ANGLE THAT THE K^+ MAKES WITH THE BEAM IN THE GRAND C.M.

77 ELASTIC SCATTERING OF POSITIVE KAONS ON POLARIZED PROTONS AT 1.22 AND 2.48 GEV/C. [PHYS. LETTERS 288, 511 (1969)]

S. ANDERSSON, C. DAUM, F. C. ERNE, J. P. LAGNAUX, J. C. SENS, F. UDD [EUROPEAN ORG. FOR NUC. RES., GENEVA, SWITZERLAND]

ABSTRACT POLARIZATION ANGULAR DISTRIBUTIONS OF K⁺P ELASTIC SCATTERING HAVE BEEN MEASURED AT 1.22 AND 2.48 GEV/C. OVER THE MEASURED RANGE OF -T THE POLARIZATION IS GENERALLY LARGE AND POSITIVE. THE DATA AT 1.22 GEV/C RESOLVE AN AMBIGUITY IN A RECENTLY PUBLISHED PHASE-SHIFT ANALYSIS, FAVORING THE SOLUTION WHICH REQUIRES NO RESONANCE IN THE K⁺P SYSTEM.

ADDITIONAL FOOTNOTE REFERENCES

PHYS. REV. LETTERS 17, 102 (1966), PHYS. REV. 168, 1466 (1968), UCRL 18323, PHYS. LETTERS 278, 468 (1968), PHYS. REV. LETTERS 19, 255 (1967), PHYS. REV. 165, 1770 (1968), NUOVO CIMENTO 45, 885 (1966), PHYS. REV. 153, 1523 (1967), NUC. PHYS. B6, 273 (1968), VIENNA CONFERENCE (1968), AND ANL 6819.

ARTICLE READ BY ODETTE BENARY IN 3/69, AND VERIFIED BY ROGER BLAND.

BEAM IS K⁺ ON HYDROGEN COMPOUND FROM 1.22 TO 2.48 GEV/C. TARGET IS POLARIZED 63 PER CENT (NORMAL TO THE BEAM DIRECTION).

THIS EXPERIMENT USES COUNTERS.

GENERAL COMMENTS ON THIS ARTICLE

1 BASEL CONVENTION USED FOR THE SIGN OF THE POLARIZATION.

KEY WORDS = POLARIZATION MODELS

ELASTIC POLARIZATION FOR K⁺ PROTON. [TABLE 1]

LABORATORY BEAM MOMENTUM = 1.22 GEV/C.

(THIS DATA SHOULD NOT BE USED - MORE RECENT VALUES MAY BE FOUND IN ANDERSSON, ET AL, CERN (1969))

COS(THETA)	POLARIZATION (1)	
.83	.63	±.11
.77	.71	.07
.70	.76	.06
.60	.84	.05
.50	.71	.07
.39	.86	.07
.32	.80	.10
.25	.88	.09
.18	.64	.09
.12	.96	.12
.05	.74	.11
-.01	.85	.12
-.07	.88	.14
-.16	.57	.11
-.26	.57	.13
-.35	.41	.15
-.46	.11	.16
-.55	.27	.12
-.66	.14	.16
-.74	.01	.15
-.82	.05	.22

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.

THE POLARIZATION IS OF THE PROTON ALONG THE NORMAL TO THE PRODUCTION PLANE IN THE GRAND C.M.

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF ± 5 PER CENT.

ELASTIC POLARIZATION FOR K⁺ PROTON. [TABLE 1]

LABORATORY BEAM MOMENTUM = 2.48 GEV/C.

COS(THETA)	POLARIZATION (1)	
.93	.49	±.13
.90	.53	.06
.87	.51	.05
.81	.64	.06
.76	.50	.06
.71	.45	.08
.65	.49	.08
.61	.40	.13
.57	.48	.11
.50	.67	.12
.44	.28	.19
.37	.28	.18
.31	.43	.25

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.

THE POLARIZATION IS OF THE PROTON ALONG THE NORMAL TO THE PRODUCTION PLANE IN THE GRAND C.M.

(1) PLUS POSSIBLE SYSTEMATIC ERROR OF ± 5 PER CENT.

78 RESULTS FROM 3.0-GEV/C K⁺ INTERACTIONS IN HYDROGEN. [PHYS. REV. 177, 1991 (1969)]

W.P.DODD, T.JOLDERSMA, R.B.PALMER, N.P.SAMIOS (BROOKHAVEN NAT. LAB., UPTON, L.I., N. Y., USA)

ABSTRACT THE INTERACTION OF 3.0-GEV/C K⁺ MESONS ON PROTONS WAS STUDIED IN AN EXPOSURE OF THE BROOKHAVEN-NATIONAL LABORATORY (BNL) 80-IN. HYDROGEN BUBBLE CHAMBER AT THE ALTERNATING GRADIENT SYNCHROTRON (AGS). RESULTS ARE PRESENTED CONCERNING A SEARCH FOR RESONANCES REQUIRING HIGHER SU(3) REPRESENTATIONS THAN 8 OR 10, NAMELY, AN ISOTOPIC-SPIN-1 (K⁺K⁺) OR -3/2 (K⁺PI⁺) BOSON, AND AN ISOTOPIC-SPIN-0 OR -1 (K⁺N) BARYON. EVIDENCE IS ALSO GIVEN FOR THE EXISTENCE OF THE N*(1400) RESONANCE, AS WELL AS A NEW (K PI) RESONANCE WITH MASS 1260 ± 20 MEV AND WIDTH 70 MEV.

ADDITIONAL FOOTNOTE REFERENCES

BNL M-10 (1962), PHYS. LETTERS 17, 155 (1965), PHYS. REV. LETTERS 17, 155 (1965), PHYS. REV. LETTERS 17, 102 (1966), PHYS. REV. LETTERS 19, 259 (1967), PHYS. REV. 134, B383 (1964), PHYS. REV. LETTERS 16, 1363 (1966), PHYS. REV. LETTERS 19, 972 (1967), PHYS. REV. LETTERS 18, 1087 (1967), NUOVO CIMENTO 46, 539 (1966), PHYS. REV. LETTERS 16, 1059 (1966), NUOVO CIMENTO 51A, 401 (1967), NUOVO CIMENTO 36, 1101 (1965), PHYS. REV. LETTERS 17, 726 (1966), CERN HADRON CONFERENCE 127 (1958), PHYS. LETTERS 16, 351 (1965), RUTHERFORD HIGH ENERGY LAB. M-29, NUOVO CIMENTO 49, 373 (1957), BERKELEY CONFERENCE 215 (1967), PHYS. REV. LETTERS 19, 255 (1967), BERKELEY CONFERENCE 137 (1967), PHYS. LETTERS 12, 65 (1964), PHYS. REV. 138, B190 (1965), PHYS. LETTERS 18, 342 (1965), PHYS. REV. 139, B1566 (1965), PHYS. LETTERS 18, 167 (1965), PHYS. REV. LETTERS 16, 855 (1966), PHYS. REV. LETTERS 17, 789 (1966), PHYS. REV. LETTERS 17, 884 (1966), PHYS. REV. LETTERS 13, 169 (1964), PHYS. REV. LETTERS 19, 397 (1967), PHYS. REV. LETTERS 20, 164 (1968), AND PHYS. REV. LETTERS 11, 503 (1963).

ARTICLE READ BY ODETTE BENARY IN 5/69, AND VERIFIED BY LEROY PRICE.

BEAM IS K⁺ ON PROTON AT 3 GEV/C.

THIS EXPERIMENT USES THE B.N.L. 80 IN. (H) BUBBLE CHAMBER. A TOTAL OF 100000 PICTURES ARE REPORTED ON.

GENERAL COMMENTS ON THIS ARTICLE

1 CROSS SECTIONS ARE NORMALIZED TO TAU DECAYS.

2 CROSS SECTIONS IN TABLE 1 ARE IDENTICAL TO THOSE GIVEN EARLIER IN JOLDERSMA, ET AL. PRL 18,716 (1966).

KEY WORDS = CROSS SECTION N*(1400) MESON WITH Y=1 AT 1260 MEV MASS SPECTRUM D&LIT7 PLOT
ANGULAR DISTRIBUTION FITS

COMPOUND KEY WORDS = FITS ANGULAR DISTRIBUTION

FIT TO DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON = K⁺ N*(1512)⁺ [PAGE 2001]
N*(1512)⁺ = NEUTRON PI⁺ [1]

LABORATORY BEAM MOMENTUM = 3. GEV/C ± .75(1 PER CENT).

(USING A SAMPLE OF EVENTS WITH BACKWARD DECAY PION IN THE N* C.M.S)
FITTED FORMULA IS NO. EVENTS = A*EXP(B*T)

WHERE -T IS IN (GEV/C)**2. T IS THE MOMENTUM TRANSFER BETWEEN THE [INCOMING K⁺] AND THE [OUTGOING K⁺].

FITTED VALUE

B = 1.6

[1] COUNTED ALL EVENTS IN MASS BAND [MASS = 1.485 GEV; WIDTH = .080 GEV].

FIT TO DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON = K⁺ N*(1400)⁺ [PAGE 2001]
N*(1400)⁺ = NEUTRON PI⁺ [1]

LABORATORY BEAM MOMENTUM = 3. GEV/C ± .75(1 PER CENT).

(USING A SAMPLE OF EVENTS WITH BACKWARD DECAY PION IN THE N* C.M.S)
FITTED FORMULA IS NO. EVENTS = A*EXP(B*T)

WHERE -T IS IN (GEV/C)**2. T IS THE MOMENTUM TRANSFER BETWEEN THE [INCOMING K⁺] AND THE [OUTGOING K⁺].

FITTED VALUE

B = 4.3

[1] COUNTED ALL EVENTS IN MASS BAND [MASS = 1.385 GEV; WIDTH = .342 GEV].

FIT TO ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON. [PAGE 2001]

LABORATORY BEAM MOMENTUM = 3. GEV/C ± .75(1 PER CENT).
FITTED FORMULA IS NO. EVENTS = A*EXP(B*T)

WHERE -T IS IN (GEV/C)**2. T IS THE MOMENTUM TRANSFER BETWEEN THE [INCOMING K⁺] AND THE [OUTGOING K⁺].

FITTED VALUE

B = 6.6

79 K⁺P ELASTIC SCATTERING AT 7.3 GEV/C. [PHYS. LETTERS 288, 615 (1969)]

C-Y. CHEN, E. MALAMUD, D. J. MELLENA, P. E. SCHLEIN, W. E. SLATER, D. H. STORK, H. K. TICHIO (UNIV. OF CALIF., LOS ANGELES, CALIF., USA)

ABSTRACT A STUDY OF K⁺P ELASTIC SCATTERING AT 7.3 GEV/C HAS YIELDED A CROSS SECTION OF 3.86 ± 0.38 MB. FOR EVENTS PRODUCED IN THE FORWARD DIRECTION, A DIFFRACTIVE PEAK IS OBSERVED WHICH FITS THE DISTRIBUTION $(D-SIGMA/DT) = [D-SIGMA/DT(0)] \exp(\alpha T + \beta T^2)$ WITH $[D-SIGMA/DT(0)] = (20.6 \pm 2.6)$ MB/(GEV/C)-SQUARED, $\alpha = (5.46 \pm 0.29)$ (GEV/C)⁻² AND $\beta = (0.38 \pm 0.25)$ (GEV/C)⁻⁴. THE RATIO $\int RE F(0)/\int F(0)/\int F(0)$ FOR THE FORWARD SCATTERING AMPLITUDE IS 0.60 ± 0.14 . ONE ELASTIC SCATTER WAS FOUND IN THE BACKWARD DIRECTION.

ADDITIONAL FOOTNOTE REFERENCES

REVIEW OF SCIENTIFIC INSTRUMENTS 38, 879 (1967), LRL INTERDEPARTMENTAL REPORT UCID 2743, LRL INTERDEPARTMENTAL REPORT UCID 2746, PHYS. REV. 139B, 1411 (1965), NUOVO CIMENTO 43A, 142 (1965), NUOVO CIMENTO 45A, 885 (1966), PHYS. REV. 139B, 1336 (1965), PHYS. REV. LETTERS 21, 1895 (1968), UNIV. OF WISCONSIN 100-206 (1968), PHYS. REV. LETTERS 11, 503 (1963), PHYS. REV. 129, 2743 (1963), PHYS. LETTERS 16, 83 (1965), PHYS. REV. 138B, 913 (1965), NUOVO CIMENTO 45A, 792 (1966), NUC. PHYS. 84, 467 (1968), PHYS. REV. 166, 1768 (1968), PHYS. REV. LETTERS 20, 589 (1968), PHYS. LETTERS 248, 317 (1967), PHYS. REV. LETTERS 19, 675 (1967), AND PHYS. LETTERS 288, 291 (1968).

ARTICLE READ BY ODETTE BENARY IN 3/69, AND VERIFIED BY ROGER BLAND.

BEAM IS K⁺ ON PROTON AT 7.3 GEV/C.

THIS EXPERIMENT USES THE B.N.L. 80 IN. (H) BUBBLE CHAMBER. A TOTAL OF 57000 PICTURES ARE REPORTED ON.

GENERAL COMMENTS ON THIS ARTICLE

1 THE TOTAL CROSS SECTION WAS NORMALIZED TO TAU DECAYS

KEY WORDS = CROSS SECTION DIFFERENTIAL CROSS SECTION FITS MODELS

COMPOUND KEY WORDS = FITS DIFFERENTIAL CROSS SECTION

K⁺ PROTON ELASTIC CROSS SECTION. [PAGE 616]

(DETERMINED FROM THE CORRECTED NUMBER OF EVENTS FOUND BY EXTRAPOLATING THE FITTED DIFFERENTIAL CROSS SECTION TO T = 0.)

LABORATORY
BEAM MOMENTUM
GEV/C
7.3

MILLI-BARNS
3.86 ± .38

NO. EVENTS
2738

ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON. [FIGURE 1]

LABORATORY BEAM MOMENTUM = 7.3 GEV/C.

(PRIVATE COMMUNICATION FROM D. J. MELLENA, MARCH 1969. THE FIRST TWO POINTS HAVE NOT BEEN INCLUDED SINCE THEY CONTAIN UNCORRECTED SCANNING BIASES.)

-T (GEV/C)**2	D-SIGMA/D-T MB/(GEV/C)**2 (1)
MIN MAX	
.10 .15	10.370 ± .540
.15 .20	8.110 .480
.20 .25	6.530 .430
.25 .30	4.970 .370
.30 .35	3.450 .310
.35 .40	3.080 .300
.40 .45	1.770 .250
.45 .50	1.750 .220
.50 .55	1.380 .200
.55 .60	.710 .140
.60 .65	.730 .140
.65 .70	.730 .140
.70 .75	.340 .100
.75 .80	.450 .110
.80 .85	.400 .110
.85 .90	.113 .057
.90 .95	.161 .063
.95 1.00	.085 .049
1.00 1.05	.170 .069
1.05 1.10	.085 .049
1.10 1.20	.028 .020
1.20 1.30	.056 .028
1.30 1.40	.028 .020
1.40 1.50	.042 .024

1 IS THE MOMENTUM TRANSFER BETWEEN THE INCOMING K⁺ AND THE OUTGOING K⁺.

11 PLUS POSSIBLE SYSTEMATIC ERROR OF ± 8.9 PER CENT.

FIT TO ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON. [TABLE 1]

LABORATORY BEAM MOMENTUM = 7.3 GEV/C.

DATA IS FIT OVER -T FROM .1 TO 1.2 (GEV/C)**2. T IS THE MOMENTUM TRANSFER BETWEEN THE INCOMING K⁺ AND THE OUTGOING K⁺.FITTED FORMULA IS $D-SIGMA/D-T = A \exp(BT + CT^2)$

WHERE D-SIGMA/D-T IS IN MB/(GEV/C)**2 AND -T IS IN (GEV/C)**2.

FITTED VALUES

A = 20.6 ± 2.6
B = 5.16 ± .40
C = -.02 ± .43

FIT TO ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON. (TABLE 1)

LABORATORY BEAM MOMENTUM = 7.3 GEV/C.

DATA IS FIT OVER -T FROM .1 TO 1.5 (GEV/C)**2. T IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K⁺) AND THE (OUTGOING K⁺).FITTED FORMULA IS $D-SIGMA/D-T = A \cdot \exp(BT+CT)**2$

WHERE D-SIGMA/D-T IS IN MB/(GEV/C)**2 AND -T IS IN (GEV/C)**2.

FITTED VALUES

A = 20.6 +- 2.6
 B = 5.46 +- .54
 C = .39 +- .50

FIT TO ELASTIC DIFFERENTIAL CROSS SECTION FOR K⁺ PROTON. (TABLE 1)

LABORATORY BEAM MOMENTUM = 7.3 GEV/C.

DATA IS FIT OVER -T FROM .1 TO 2.0 (GEV/C)**2. T IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K⁺) AND THE (OUTGOING K⁺).FITTED FORMULA IS $D-SIGMA/D-T = A \cdot \exp(BT+CT)**2$

WHERE D-SIGMA/D-T IS IN MB/(GEV/C)**2 AND -T IS IN (GEV/C)**2.

FITTED VALUES

A = 20.6 +- 2.6
 B = 5.46 +- .29
 C = .38 +- .25

80 THE REACTION K⁺ P → K⁺ N PI AT 3 GEV/C. INCLUDING THE PRODUCTION AND DECAY OF RESONANCES. (ARKIV FYSIK 37, 468 (1968))

P. SALLSTRÖM, G. BLOMQUIST (STOCKHOLMS UNIV., STOCKHOLM, SWEDEN)

ABSTRACT THE REACTION K⁺ P → K⁺ N PI HAS BEEN STUDIED IN ALL THREE FINAL CHARGE STATES AT MOMENTUM OF THE INCIDENT K⁺-MESONS OF 2.97 GEV/C, I.E. A TOTAL CM ENERGY OF 2.60 GEV. THE PRODUCTION CROSS SECTION FOR K⁺ IS FOUND TO BE (1.4 +- 0.2) MB AND FOR N⁰ (0.9 +- 0.1) MB. THE DECAY ANGULAR CORRELATIONS FOR THE TWO DECAY MODES OF K⁺, K⁺ → K⁰ PI⁺ AND K⁺ → K⁺ PI⁰ ARE FOUND TO BE DIFFERENT. THIS CAN BE EXPLAINED AS DUE TO AN INTERFERENCE BETWEEN STATES OF DIFFERENT TOTAL ISOSPIN OF THE (K PI)-SYSTEM. THE K⁺ DECAY ANGULAR CORRELATIONS, WHEN INTERFERENCE EFFECTS ARE ELIMINATED, INDICATE THAT THE RESONANCE IS PRODUCED TO A GREAT EXTENT BY EXCHANGE OF A VECTOR MESON. THE SPIN DENSITY MATRIX ELEMENTS ARE -- RHO(0,0) = 0.25 +- 0.05; RHO(1,-1) = 0.18 +- 0.04; AND {RE RHO(1,0)} = -0.11 +- 0.03. THE SPIN DENSITY MATRIX ELEMENTS FOR N⁰ ARE -- RHO(3,3) = 0.33 +- 0.05; {RE RHO(3,-1)} = 0.32 +- 0.04; AND {RE RHO(3,1)} = 0.00 +- 0.02, IN AGREEMENT WITH THE PREDICTIONS OF THE MODEL OF STODOLSKY AND SAKURAI. THE MODIFIED ONE-PARTICLE EXCHANGE MODEL OF JACKSON ET AL. GIVES A REASONABLE ACCOUNT OF THE PRODUCTION DIFFERENTIAL CROSS SECTIONS AS WELL AS OF THE VARIATIONS OF THE MATRIX ELEMENTS WITH THE PRODUCTION ANGLE OF THE RESONANCE. THE SAME IS TRUE OF THE REGGE POLE MODEL. THE DOUBLE PERIPHERAL MODEL, APPLIED TO THE REACTION K⁺ P → K⁺ N PI⁺, GIVES A REASONABLE DESCRIPTION OF THE CROSS SECTIONS IF (PI⁺, K⁰), AND (RHO⁺, OMEGA)-EXCHANGES ARE ASSUMED TOGETHER WITH A NON-PERIPHERAL BACKGROUND.

CROSS REFERENCES

THIS ARTICLE SUPERSEDES PART OF NUOVO CIMENTO 49A, 348 (1967).

ADDITIONAL FOOTNOTE REFERENCES

PHYS. LETTERS 16, 83 (1965), PHYS. REV. 142, 913 (1966), PHYS. REV. 158, 1298 (1967), PHYS. REV. LETTERS 15, 737 (1965), NUOVO CIMENTO 36, 1101 (1965), NUOVO CIMENTO 51A, 401 (1967), PHYS. REV. LETTERS 11, 90 (1963), NUOVO CIMENTO 34, 735 (1964), PHYS. REV. 139, 8428 (1965), CERN PHYS. 67-33, NUOVO CIMENTO 43, 142 (1965), LAB. DES HAUTES EN. AT BRUXELLES REPORT 74 (1966), PHYS. LETTERS 17, 155 (1965), NUOVO CIMENTO 34, 1644 (1964), UCRL 8030 SEPT. (1967), PHYS. REV. LETTERS 16, 1228 (1966), PHYS. REV. LETTERS 17, 102 (1966), PHYS. REV. LETTERS 19, 259 (1967), PHYS. LETTERS 12, 235 (1966), PHYS. REV. LETTERS 16, 1089 (1966), PHYS. REV. LETTERS 17, 726 (1966), NUOVO CIMENTO 49A, 9 (1967), CERN PHYS. 67-26 (1967), PHYS. LETTERS 22, 347 (1966), PHYS. REV. LETTERS 17, 939 (1966), PHYS. LETTERS 21, 442 (1966), PHYS. REV. 10, 351 (1963), CERN PHYS. 67-19 (1967), PHYS. REV. 148, 1410 (1966), PHYS. REV. 161, 1581 (1967), PHYS. LETTERS 12, 352 (1964), PHYS. REV. LETTERS 16, 485 (1966), NUOVO CIMENTO 46, 539 (1966), PHYS. REV. 161, 1581 (1967), PHYS. LETTERS 12, 352 (1964), PHYS. REV. LETTERS 16, 485 (1966), NUOVO CIMENTO 33, 1407 (1964), AND NUOVO CIMENTO 33, 906 (1964).

ARTICLE READ BY ODETTE BENARY IN 7/69, AND VERIFIED BY ROGER BLAND.

BEAM IS K⁺ UN PHUION AT 2.972 GEV/C.

THIS EXPERIMENT USES THE SACLAY 81 CM (N) BUBBLE CHAMBER.

GENERAL COMMENTS ON THIS ARTICLE

1 ONLY TWO PRONG EVENTS WITH NO VISIBLE K⁰ DECAY WERE ANALYZED.

KEY WORDS → K*(890) DELTA(1238) DALITZ PLOT MASS SPECTRUM DENSITY MATRIX MODELS CROSS SECTION
 ANGULAR DISTRIBUTION

CROSS SECTION FOR K⁺ PROTON → K*(890)⁺ PROTON. (TABLE 2)K*(890)⁺ → K⁺ PI⁰ [1]

[THIS DATA REPLACES VALUES GIVEN EARLIER IN SALLSTRÖM, ET AL., NUOVO CIMENTO 49A, 348 (1967)]

LABORATORY

BEAM MOMENTUM

GEV/C

2.972 +- .015

MILLI-BARNS

.44 +- .09

[1] FITTED FOR MASS AND/OR WIDTH (MASS = .890 GEV; WIDTH = .057 GEV), AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.

CROSS SECTION FOR K⁺ PROTON → K*(890)⁺ PROTON. (TABLE 2)K*(890)⁺ → K⁰ PI⁺ [1]

[THIS DATA REPLACES THE VALUE GIVEN EARLIER IN SALLSTRÖM, ET AL., NUOVO CIMENTO 49A, 348 (1967)]

LABORATORY

BEAM MOMENTUM

GEV/C

2.972 +- .015

MILLI-BARNS

.95 +- .13

[1] FITTED FOR MASS AND/OR WIDTH (MASS = .894 GEV; WIDTH = .071 GEV), AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow \Delta^{++} K^0$. [TABLE 2]
 $\Delta^{++} \rightarrow \text{PROTON } \pi^+ [1]$

[THIS DATA REPLACES THE VALUE GIVEN EARLIER IN SALLSTROM, ET AL., NUOVO CIMENTO 49A, 348 (1967)]

LABORATORY
 BEAM MOMENTUM
 GEV/C
 2.972 $\pm$.015
 MILLI-BARNS
 .68 $\pm$.11

[1] FITTED FOR MASS AND/OR WIDTH [MASS = 1.216 GEV; WIDTH = .095 GEV], AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow \Delta^{++} K^+$. [TABLE 2]
 $\Delta^{++} \rightarrow \text{H MESON } \pi^0 [1]$

[THIS DATA REPLACES THE VALUE GIVEN EARLIER IN SALLSTROM, ET AL., NUOVO CIMENTO 49A, 348 (1967)]

LABORATORY
 BEAM MOMENTUM
 GEV/C
 2.972 $\pm$.015
 MILLI-BARNS
 .16 $\pm$.06

[1] FITTED FOR MASS AND/OR WIDTH, AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow \Delta^{++} K^+$. [TABLE 2]
 $\Delta^{++} \rightarrow \text{H MESON } \pi^+ [1]$

[THIS DATA REPLACES THE VALUE GIVEN EARLIER IN SALLSTROM, ET AL., NUOVO CIMENTO 49A, 348 (1967)]

LABORATORY
 BEAM MOMENTUM
 GEV/C
 2.972 $\pm$.015
 MILLI-BARNS
 .02 $\pm$.05

[1] FITTED FOR MASS AND/OR WIDTH, AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow \Delta^{++}(1238) K \text{ MESON}$. [TABLE 2]

LABORATORY
 BEAM MOMENTUM
 GEV/C
 2.972 $\pm$.015
 MILLI-BARNS
 .86 $\pm$.13

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)^+ \text{ PROTON}$. [FIGURE 10]
 $K^*(890)^+ \rightarrow K^0 \pi^+ [1]$

LABORATORY BEAM MOMENTUM = 2.972 $\pm$.015 GEV/C.

[THIS DATA REPLACES THE VALUE GIVEN EARLIER IN SALLSTROM, ET AL., NUOVO CIMENTO 49A, 348 (1967)]

.....
 . THIS DATA WAS READ FROM A GRAPH .

-T (GEV/C)**2		D-SIGMA/D-T MB/(GEV/C)**2	
MIN	MAX		
.02	.12	5.00	$\pm$ 1.00
.12	.22	2.20	.50
.22	.32	1.50	.50
.32	.42	1.20	.30
.42	.52	.75	.20
.52	.60	.40	.15
.60	.70	.15	.09
.70	.80	.34	.19
.80	.90	.15	.09
.90	1.00	.08	.06

T IS THE MOMENTUM TRANSFER BETWEEN THE [INCOMING K^+] AND THE [$K^*(890)^+$].

[1] COUNTED ALL EVENTS IN MASS BAND.

FIT TO DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)^+ \text{ PROTON}$. [PAGE 480]
 $K^*(890)^+ \rightarrow K^0 \pi^+ [1]$

LABORATORY BEAM MOMENTUM = 2.972 $\pm$.015 GEV/C.
 FITTED FORMULA IS $D-SIGMA/D-T = F \exp(-B \cdot T)$

WHERE $D-SIGMA/D-T$ IS IN MB/(GEV/C)**2 AND $-T$ IS IN (GEV/C)**2. T IS THE MOMENTUM TRANSFER BETWEEN THE [INCOMING K^+] AND THE [$K^*(890)^+$].

FITTED VALUE

B = 4.6 $\pm$.2

[1] USED SIMPLE MASS CUT.

81 STUDY OF THE REACTION $K^+ p \rightarrow K^0 \pi^+ p$ AT 4.6 GEV/C AND A REVIEW OF THE AVAILABLE DATA. [UCRL 18417 (1968)]

C.F. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA, AND UNIV. OF CALIFORNIA, BERKELEY, CALIF., USA

ABSTRACT THE PRODUCTION AND THE DECAY PROPERTIES OF THE DOMINANT CHANNELS FOR THE REACTION $K^+ p \rightarrow K^0 \pi^+ p$, $K^0 \Delta^{++}(1236)$, AND $K^*(890) p$, ARE STUDIED AT 4.6 GEV/C. AS A RESULT IN COMPARING WITH THE EXISTING DATA AT VARIOUS MOMENTA, IT IS FOUND THAT THE PRODUCTION MECHANISMS FOR THIS REACTION ARE APPROXIMATELY INDEPENDENT OF THE LABORATORY MOMENTUM FROM 2-3 GEV/C UP TO 13 GEV/C, THE HIGHEST LABORATORY MOMENTUM AVAILABLE. THIS PHENOMENON CAN BE UNDERSTOOD IN TERMS OF THE REGGE-POLE MODEL.

ADDITIONAL FOOTNOTE REFERENCES

PHYS. REV. LETTERS 17, 939 (1966), UCRL 16370 (1965), PHYS. REV. 142, 913 (1966), PHYS. REV. 158, 1298 (1967), PHYS. REV. 158, 1310 (1967), PHYS. LETTERS 9, 359 (1964), CERN 67-5 (1967), NUOVO CIMENTO 33, 309 (1954), PHYS. REV. 139, 8428 (1965), NUOVO CIMENTO 34, 1644 (1964), NUOVO CIMENTO 33, 906 (1964), NUOVO CIMENTO 34, 1842E (1964), PHYS. LETTERS 8, 144 (1964), PHYS. REV. 139, 8546 (1965), NUOVO CIMENTO 45A, 59 (1966), NUOVO CIMENTO 24, 454 (1962), PHYS. REV. 134, 81099 (1964), PHYS. REV. LETTERS 11, 90 (1963), BULL. AM. PHYS. SOC. 8, 20 (1963), UCRL 16332 (1966), PHYS. REV. LETTERS 10, 351 (1963), PHYS. REV. LETTERS 22, 528 (1966), NUOVO CIMENTO 52A, 308 (1957), PHYS. REV. LETTERS 19, 614 (1967), PHYS. REV. 163, 1572 (1967), NUOVO CIMENTO 49, 157 (1967), NUOVO CIMENTO 51, 696 (1967), PHYS. REV. 160, 1323 (1967), PHYS. REV. 160, 1326 (1967), PHYS. REV. 166, 1525 (1968), PHYS. REV. LETTERS 15, 737 (1965), PHYS. REV. 147, 1045 (1966), PHYS. LETTERS 20, 79 (1966), PHYS. LETTERS 19, 598 (1965), PHYS. REV. 166, 1768 (1968), UCRL 18124 (1968), AND UCRL 18275 (1968).

ARTICLE READ BY ODETTE BENARY IN 7/69, AND VERIFIED BY ROGER BLAND.

BEAM IS K^+ ON PROTON AT 4.6 GEV/C.

THIS EXPERIMENT USES THE B.N.L. 80 IN. (H) BUBBLE CHAMBER.

KEY WORDS = CROSS SECTION DALITZ PLOT MODELS DENSITY MATRIX DIFFERENTIAL CROSS SECTION FITS
MASS SPECTRUM

COMPOUND KEY WORDS = FITS DIFFERENTIAL CROSS SECTION

CROSS SECTION FOR $K^+ \text{ PROTON } \rightarrow K^0 \pi^+ \text{ PROTON. [FIGURE 4]}$

THIS DATA WAS READ FROM A GRAPH.

LABORATORY BEAM MOMENTUM GEV/C 4.6	MILLI-BARNS 1.20 +- .12
---	----------------------------

CROSS SECTION FOR $K^+ \text{ PROTON } \rightarrow K^0 \Delta^{++}. [TABLE 1]$

LABORATORY BEAM MOMENTUM GEV/C 4.6	MILLI-BARNS .36 +- .04	NO. EVENTS 144
---	---------------------------	-------------------

CROSS SECTION FOR $K^+ \text{ PROTON } \rightarrow K^*(890)^+ \text{ PROTON. [TABLE 1]}$
 $K^*(890)^+ \rightarrow K^0 \pi^+ [1]$

LABORATORY BEAM MOMENTUM GEV/C 4.6	MILLI-BARNS .36 +- .04	NO. EVENTS 140
---	---------------------------	-------------------

[1] USED SIMPLE MASS CUT.

CROSS SECTION FOR $K^+ \text{ PROTON } \rightarrow K^*(1420)^+ \text{ PROTON. [TABLE 1]}$
 $K^*(1420)^+ \rightarrow K^0 \pi^+ [1]$

LABORATORY BEAM MOMENTUM GEV/C 4.6	MILLI-BARNS .13 +- .02	NO. EVENTS 51
---	---------------------------	------------------

[1] USED SIMPLE MASS CUT.

CROSS SECTION FOR $K^+ \text{ PROTON } \rightarrow K^0 \pi^+ \text{ PROTON. [TABLE 1]}$
CROSS SECTION IS ONLY FOR THE NON-RESONANT PRODUCTION OF THESE PARTICLES.

LABORATORY BEAM MOMENTUM GEV/C 4.6	MILLI-BARNS .37 +- .04	NO. EVENTS 147
---	---------------------------	-------------------

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \Delta^{++}$. [FIGURE 9A]
LABORATORY BEAM MOMENTUM = 4.6 GEV/C.

THIS DATA WAS READ FROM A GRAPH.

-T (GEV/C)**2		NO. EVENTS
MIN	MAX	
.0	.2	50.0 +- 7.0
.2	.4	52.0 7.0
.4	.6	17.0 4.0
.6	1.0	5.5 1.5

T IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K^+) AND THE (K^0).

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)^+ \text{ PROTON}$. [FIGURE 9B]
 $K^*(890)^+ \rightarrow K^0 \pi^+ [1]$

LABORATORY BEAM MOMENTUM = 4.6 GEV/C.

THIS DATA WAS READ FROM A GRAPH.

-T (GEV/C)**2		NO. EVENTS
MIN	MAX	
.0	.1	20.0 +- 5.0
.1	.3	28.0 2.0
.3	.5	15.0 3.0
.5	.8	5.6 2.0
.8	1.0	2.0 1.0

T IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K^+) AND THE ($K^*(890)^+$).

[1] USED SIMPLE MASS CUT.

FIT TO DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow \Delta^{++} K^0$. [TABLE 2]

LABORATORY BEAM MOMENTUM = 4.6 GEV/C.

DATA IS FIT OVER -T FROM .3 TO 1.0 (GEV/C)**2. T IS THE MOMENTUM TRANSFER BETWEEN THE (TARGET) AND THE (Δ^{++}).
FITTED FORMULA IS $D\text{-SIGMA}/D\text{-T} = A \cdot \exp(B \cdot T)$

WHERE $D\text{-SIGMA}/D\text{-T}$ IS IN MB/(GEV/C)**2 AND -T IS IN (GEV/C)**2.

FITTED VALUE

B = 4.6 +- .5

FIT TO DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)^+ \text{ PROTON}$. [TABLE 2]
 $K^*(890)^+ \rightarrow K^0 \pi^+ [1]$

LABORATORY BEAM MOMENTUM = 4.6 GEV/C.

DATA IS FIT OVER -T FROM .2 TO 1.0 (GEV/C)**2. T IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K^+) AND THE ($K^*(890)^+$).
FITTED FORMULA IS $D\text{-SIGMA}/D\text{-T} = A \cdot \exp(B \cdot T)$

WHERE $D\text{-SIGMA}/D\text{-T}$ IS IN MB/(GEV/C)**2 AND -T IS IN (GEV/C)**2.

FITTED VALUE

B = 3.6 +- .3

[1] USED SIMPLE MASS CUT.

82 DOUBLE-RESONANCE PRODUCTION BY 5 GEV/C K^+ MESONS ON PROTONS. [NUOVO CIMENTO 61A, 397 (1959)]

M. DEBAERE, J. DEBAISIEUX, P. DUFOUR, F. GRARD, J. HEUGHEBAERT, L. PAPE, P. PEETERS, F. VERBEURE, R. WINDMOLDERS (INST. INTERUNIV. DES SCI. NUC., BRUXELLES, BELGIUM)
G. BASSOMPIERRE, Y. GOLOSCHNIGOT-CLERMONT, A. GRANT, V. P. MENRI, B. JONGEJANS, D. LINGLIN, W. MATT, F. MULLER, J. M. PERREAU, H. PIOTROWSKA, T. SATTOV, R. L. SEKULIN, G. WOLF (EUROPEAN ORG. FOR NUC. RES., GENEVA, SWITZERLAND)

ABSTRACT AMONG THE INTERACTIONS OF K^+P AT 5 GEV/C PRODUCED IN THE CERN 2 M AND SACLAY 81 CM HYDROGEN BUBBLE CHAMBERS WE HAVE STUDIED THE FOUR-BODY REACTIONS $K^+P \rightarrow K^+N\pi^+\pi^+$. THE CROSS-SECTIONS FOR THE VARIOUS FINAL STATES CONTRIBUTING TO THESE REACTIONS HAVE BEEN DETERMINED. A DETAILED ANALYSIS HAS BEEN PERFORMED ON THE DOUBLE-RESONANCE PRODUCTION $K^+P \rightarrow K^*(892)\pi^+\pi^+\pi^+$. DECAY ANGULAR DISTRIBUTIONS FOR BOTH RESONANCES HAVE BEEN STUDIED AND SINGLE AND JOINT SPIN-DENSITY MATRIX ELEMENTS CALCULATED AS A FUNCTION OF FOUR-MOMENTUM TRANSFER. THE RESULTS HAVE BEEN COMPARED TO THE ABSORPTION-MODEL PREDICTIONS ASSUMING PION EXCHANGE. VARIOUS QUARK-MODEL PREDICTIONS FOR THE DECAY ANGULAR DISTRIBUTIONS HAVE BEEN TESTED. THE QUARK-MODEL RELATIONS DERIVED ON THE BASIS OF THE ADDITIVITY ASSUMPTION ONLY, ARE EXPERIMENTALLY WELL SATISFIED. IF ADDITIONAL CONDITIONS ABOUT THE QUARK-QUARK SCATTERING AMPLITUDES ARE IMPOSED, ONLY SOME OF THE QUARK-MODEL PREDICTIONS AGREE WITH EXPERIMENT.

CROSS REFERENCES

SEE ALSO PHYS. LETTERS 27B, 468 (1968), AND NUOVO CIMENTO 49A, 9 (1957).

ADDITIONAL FOOTNOTE REFERENCES

PHYS. LETTERS 26B, 33 (1967), NUOVO CIMENTO 46A, 539 (1966), PHYS. REV. LETTERS 16, 672 (1966), PHYS. REV. 151, 1306 (1966), NUOVO CIMENTO 38, 518 (1965), PHYS. LETTERS 18, 1 (1965), PHYS. LETTERS 28B, 72 (1968), NUC. PHYS. 87, 681 (1968), NUOVO CIMENTO 32A, 1152 (1967), NUC. PHYS. 36, 449 (1968), NUC. PHYS. 84, 559 (1968), AND PHYS. LETTERS 25B, 554 (1967).

ARTICLE READ BY ODETTE BENARY IN 6/69, AND VERIFIED BY ROGER BLAND.

BEAM IS K^+ ON PROTON AT 4.97 GEV/C.

THIS EXPERIMENT USES BUBBLE CHAMBERS.

GENERAL COMMENTS ON THIS ARTICLE

1 THIS ARTICLE REPORTS CROSS SECTIONS FOR $K^+P \rightarrow K^+P\pi^+\pi^+$, $K^0P\pi^+\pi^0$, AND $K^0N\pi^+\pi^+$ WHICH HAVE BEEN PREVIOUSLY PUBLISHED IN BASSOMPIERRE PL 27B, 468 (1968).

KEY WORDS = CROSS SECTION DIFFERENTIAL CROSS SECTION FITS ANGULAR DISTRIBUTION DENSITY MATRIX
MODELS MASS SPECTRUM

COMPOUND KEY WORDS = FITS DIFFERENTIAL CROSS SECTION

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \text{ PI}^- \text{ DELTA}^{++}$ (TABLE 1A)
 $\text{DELTA}^{++} \rightarrow \text{PROTON PI}^+ \{1\}$

LABORATORY	FRACTION
BEAM MOMENTUM	
GEV/C	PER CENT {2}
5.000 $\pm$.015	14.4 $\pm$ 1.0

- {1} FITTED FOR MASS AND/OR WIDTH (MASS = 1.225 GEV; WIDTH = .110 GEV), AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.
 {2} THIS FRACTION SHOULD BE MULTIPLIED BY 1.80 $\pm$.15 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)0 \text{ DELTA}^{++}$ (TABLE 1A)
 $K^*(890)0 \rightarrow K^+ \text{ PI}^- \{1\}$

LABORATORY	FRACTION
BEAM MOMENTUM	
GEV/C	PER CENT {2}
5.000 $\pm$.015	34.4 $\pm$.9

- {1} FITTED FOR MASS AND/OR WIDTH (MASS = .890 GEV; WIDTH = .058 GEV), AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.
 {2} THIS FRACTION SHOULD BE MULTIPLIED BY 1.80 $\pm$.15 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(1420)0 \text{ DELTA}^{++}$ (TABLE 1A)
 $K^*(1420)0 \rightarrow K^+ \text{ PI}^- \{1\}$

LABORATORY	FRACTION
BEAM MOMENTUM	
GEV/C	PER CENT {2}
5.000 $\pm$.015	7.3 $\pm$.6

- {1} FITTED FOR MASS AND/OR WIDTH (MASS = 1.422 GEV; WIDTH = .110 GEV), AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.
 {2} THIS FRACTION SHOULD BE MULTIPLIED BY 1.80 $\pm$.15 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^+ \text{ PI}^+ \text{ DELTA}^{++}$ (TABLE 1A)
 $\text{DELTA}^{++} \rightarrow \text{PROTON PI}^+ \{1\}$

LABORATORY	FRACTION
BEAM MOMENTUM	
GEV/C	PER CENT {2}
5.000 $\pm$.015	6.2 $\pm$.7

- {1} FITTED FOR MASS AND/OR WIDTH (MASS = 1.243 GEV; WIDTH = .130 GEV), AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.
 {2} THIS FRACTION SHOULD BE MULTIPLIED BY 1.80 $\pm$.15 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(1420)^+ \text{ PROTON}$ (TABLE 1A)
 $K^*(1420)^+ \rightarrow K^*(890)0 \text{ PI}^+ \{1\}$
 $K^*(1070)0 \rightarrow K^+ \text{ PI}^- \{2\}$

LABORATORY	FRACTION
BEAM MOMENTUM	
GEV/C	PER CENT {3}
5.000 $\pm$.015	4.0 $\pm$.7

- {1} FITTED FOR MASS AND/OR WIDTH (MASS = 1.396 GEV; WIDTH = .100 GEV), AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.
 {2} USED SIMPLE MASS CUT.
 {3} THIS FRACTION SHOULD BE MULTIPLIED BY 1.80 $\pm$.15 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(1420)^+ \text{ PROTON}$ (TABLE 1A)
 $K^*(1420)^+ \rightarrow K^+ \text{ PI}^+ \text{ PI}^- \{1\}$

LABORATORY	FRACTION
BEAM MOMENTUM	
GEV/C	PER CENT {2}
5.000 $\pm$.015	1.0 $\pm$.6

- {1} FITTED FOR MASS AND/OR WIDTH (MASS = 1.396 GEV; WIDTH = .100 GEV), AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.
 {2} THIS FRACTION SHOULD BE MULTIPLIED BY 1.80 $\pm$.15 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(1225)^+ \text{ PROTON}$ (TABLE 1A)
 $K^*(1225)^+ \rightarrow K^*(890)0 \text{ PI}^+ \{1\}$
 $K^*(890)0 \rightarrow K^+ \text{ PI}^- \{2\}$

LABORATORY	FRACTION
BEAM MOMENTUM	
GEV/C	PER CENT {3}
5.000 $\pm$.015	17.7 $\pm$.9

- {1} FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH (MASS = 1.225 GEV; WIDTH = .200 GEV) AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 {2} USED SIMPLE MASS CUT.
 {3} THIS FRACTION SHOULD BE MULTIPLIED BY 1.80 $\pm$.15 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = K^*(1225)^+ \text{ PROTON.}$ [TABLE 1A]
 $K^*(1225)^+ = K^+ \text{ RHO0 [1]}$
 $\text{RHO0} = \text{PI}^+ \text{ PI}^- \text{ [2]}$

LABORATORY	FRACTION
BEAM MOMENTUM	
GEV/C	PER CENT [3]
5.000 $\pm$.015	5.5 $\pm$.9

- [1] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH [MASS = 1.225 GEV; WIDTH = .200 GEV] AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 [2] USED SIMPLE MASS CUT.
 [3] THIS FRACTION SHOULD BE MULTIPLIED BY 1.80 $\pm$.15 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = K^*(1225)^+ \text{ PROTON.}$ [TABLE 1A]
 $K^*(1225)^+ = K^+ \text{ PI}^+ \text{ [1]}$

LABORATORY	FRACTION
BEAM MOMENTUM	
GEV/C	PER CENT [2]
5.000 $\pm$.015	1.

- [1] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH [MASS = 1.225 GEV; WIDTH = .200 GEV] AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 [2] THIS FRACTION SHOULD BE MULTIPLIED BY 1.80 $\pm$.15 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = \text{RHO0 } K^+ \text{ PROTON.}$ [TABLE 1A]
 $\text{RHO0} = \text{PI}^+ \text{ PI}^- \text{ [1]}$

LABORATORY	FRACTION
BEAM MOMENTUM	
GEV/C	PER CENT [2]
5.000 $\pm$.015	2.7 $\pm$.8

- [1] FITTED FOR MASS AND/OR WIDTH [MASS = .742 GEV; WIDTH = .164 GEV]; AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.
 [2] THIS FRACTION SHOULD BE MULTIPLIED BY 1.80 $\pm$.15 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = K^*(890)0 \text{ PROTON PI}^+.$ [TABLE 1A]
 $K^*(890)0 = K^+ \text{ PI}^- \text{ [1]}$

LABORATORY	FRACTION
BEAM MOMENTUM	
GEV/C	PER CENT [2]
5.000 $\pm$.015	.7

- [1] FITTED FOR MASS AND/OR WIDTH, AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.
 [2] THIS FRACTION SHOULD BE MULTIPLIED BY 1.80 $\pm$.15 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = K^*(1420)0 \text{ PROTON PI}^+.$ [TABLE 1A]
 $K^*(1420)0 = K^+ \text{ PI}^- \text{ [1]}$

LABORATORY	FRACTION
BEAM MOMENTUM	
GEV/C	PER CENT [2]
5.000 $\pm$.015	.5

- [1] FITTED FOR MASS AND/OR WIDTH, AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.
 [2] THIS FRACTION SHOULD BE MULTIPLIED BY 1.80 $\pm$.15 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = K^*(1420)^+ \text{ PROTON.}$ [TABLE 1A]
 $K^*(1420)^+ = K^+ \text{ RHO0 [1]}$

LABORATORY	FRACTION
BEAM MOMENTUM	
GEV/C	PER CENT [2]
5.000 $\pm$.015	.5

- [1] USED SIMPLE MASS CUT.
 [2] THIS FRACTION SHOULD BE MULTIPLIED BY 1.80 $\pm$.15 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = K^*(1660)^+ \text{ PROTON.}$ [TABLE 1A]
 $K^*(1660)^+ = K^*(1420)0 \text{ PI}^+ \text{ [1]}$

LABORATORY	FRACTION
BEAM MOMENTUM	
GEV/C	PER CENT [2]
5.000 $\pm$.015	1.3 $\pm$.3

- [1] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH [MASS = 1.660 GEV] AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 [2] THIS FRACTION SHOULD BE MULTIPLIED BY 1.80 $\pm$.15 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = K^*(1660)^+ \text{ PROTON.}$ (TABLE 1A)
 $K^*(1560)^+ = K^*(1890) \text{ PI}^+ \{1\}$

LABORATORY BEAM MOMENTUM FRACTION
 GEV/C PER CENT {2}
 5.000 $\pm$.015 .2

- {1} FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH (MASS = 1.660 GEV) AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 {2} THIS FRACTION SHOULD BE MULTIPLIED BY 1.80 $\pm$.15 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = K^+ \text{ PROTON PI}^+ \text{ PI}^-.$ (TABLE 1A)
 CROSS SECTION IS ONLY FOR THE NON-RESONANT PRODUCTION OF THESE PARTICLES.

LABORATORY BEAM MOMENTUM FRACTION
 GEV/C PER CENT {1}
 5.000 $\pm$.015 5.5 $\pm$ 1.0

- {1} THIS FRACTION SHOULD BE MULTIPLIED BY 1.80 $\pm$.15 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = \text{DELTA}^{++} \text{ KO PI}^0.$ (TABLE 1B)
 $\text{DELTA}^{++} = \text{PROTON PI}^+ \{1\}$

LABORATORY BEAM MOMENTUM FRACTION
 GEV/C PER CENT {2}
 5.000 $\pm$.015 10.5 $\pm$ 1.5

- {1} FITTED FOR MASS AND/OR WIDTH (MASS = 1.227 GEV; WIDTH = .132 GEV), AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.
 {2} THIS FRACTION SHOULD BE MULTIPLIED BY 1.5 $\pm$.2 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = K^*(890)^+ \text{ PROTON PI}^0.$ (TABLE 1B)
 $K^*(890)^+ = \text{KO PI}^+ \{1\}$

LABORATORY BEAM MOMENTUM FRACTION
 GEV/C PER CENT {2}
 5.000 $\pm$.015 1.8 $\pm$ 1.0

- {1} FITTED FOR MASS AND/OR WIDTH (MASS = .897 GEV; WIDTH = .058 GEV), AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.
 {2} THIS FRACTION SHOULD BE MULTIPLIED BY 1.5 $\pm$.2 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = \text{DELTA}^+ \text{ KO PI}^+.$ (TABLE 1B)
 $\text{DELTA}^+ = \text{PROTON PI}^0 \{1\}$

LABORATORY BEAM MOMENTUM FRACTION
 GEV/C PER CENT {2}
 5.000 $\pm$.015 4.8 $\pm$ 1.1

- {1} FITTED FOR MASS AND/OR WIDTH (MASS = 1.224 GEV; WIDTH = .130 GEV), AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.
 {2} THIS FRACTION SHOULD BE MULTIPLIED BY 1.5 $\pm$.2 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = \text{RHO}^+ \text{ KO PROTON.}$ (TABLE 1B)
 $\text{RHO}^+ = \text{PI}^+ \text{ PI}^0 \{1\}$

LABORATORY BEAM MOMENTUM FRACTION
 GEV/C PER CENT {2}
 5.000 $\pm$.015 3.1 $\pm$ 1.2

- {1} FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH (MASS = .750 GEV; WIDTH = .124 GEV) AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 {2} THIS FRACTION SHOULD BE MULTIPLIED BY 1.5 $\pm$.2 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = \text{DELTA}^{++} K^*(890)^0.$ (TABLE 1B)
 $\text{DELTA}^{++} = \text{PROTON PI}^+ \{1\}$
 $K^*(890)^0 = \text{KO PI}^0 \{2\}$

LABORATORY BEAM MOMENTUM FRACTION
 GEV/C PER CENT {3}
 5.000 $\pm$.015 21.2 $\pm$ 1.2

- {1} FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 {2} FITTED FOR MASS AND/OR WIDTH (MASS = .894 GEV; WIDTH = .058 GEV), AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.
 {3} THIS FRACTION SHOULD BE MULTIPLIED BY 1.5 $\pm$.2 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = \Delta K^+ K^+ (890)^+ \dots$ [TABLE 18]
 $\Delta K^+ = \dots$ PROTON P_{10} [1]
 $K^+ (890)^+ = \dots$ $K^0 P_{10}$ [2]

LABORATORY	FRACTION
BEAM MOMENTUM	
GEV/C	PER CENT [3]
5.000 $\pm$.015	8.7 $\pm$.9

- (1) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 (2) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 (3) THIS FRACTION SHOULD BE MULTIPLIED BY 1.5 $\pm$.2 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = \Delta K^+ K^+ (1420)^+ \dots$ [TABLE 18]
 $\Delta K^+ = \dots$ PROTON P_{10} [1]
 $K^+ (1420)^+ = \dots$ $K^0 P_{10}$ [2]

LABORATORY	FRACTION
BEAM MOMENTUM	
GEV/C	PER CENT [3]
5.000 $\pm$.015	6.7 $\pm$.8

- (1) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 (2) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH [MASS = 1.415 GEV; WIDTH = .125 GEV] AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 (3) THIS FRACTION SHOULD BE MULTIPLIED BY 1.5 $\pm$.2 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = \Delta K^+ K^+ (1420)^+ \dots$ [TABLE 18]
 $\Delta K^+ = \dots$ PROTON P_{10} [1]
 $K^+ (1420)^+ = \dots$ $K^0 P_{10}$ [2]

LABORATORY	FRACTION
BEAM MOMENTUM	
GEV/C	PER CENT [3]
5.000 $\pm$.015	2.8 $\pm$.7

- (1) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 (2) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH [MASS = 1.415 GEV; WIDTH = .125 GEV] AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 (3) THIS FRACTION SHOULD BE MULTIPLIED BY 1.5 $\pm$.2 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = K^+ (1420)^+ \text{ PROTON} \dots$ [TABLE 18]
 $K^+ (1420)^+ = \dots$ $K^+ (890)^+ P_{10}$ [1]

LABORATORY	FRACTION
BEAM MOMENTUM	
GEV/C	PER CENT [2]
5.000 $\pm$.015	1.0 $\pm$.8

- (1) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH [MASS = 1.416 GEV; WIDTH = .125 GEV] AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 (2) THIS FRACTION SHOULD BE MULTIPLIED BY 1.5 $\pm$.2 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = K^+ (1420)^+ \text{ PROTON} \dots$ [TABLE 18]
 $K^+ (1420)^+ = \dots$ $K^+ (890)^+ P_{10}$ [1]

LABORATORY	FRACTION
BEAM MOMENTUM	
GEV/C	PER CENT [2]
5.000 $\pm$.015	1.0 $\pm$.8

- (1) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH [MASS = 1.416 GEV; WIDTH = .125 GEV] AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 (2) THIS FRACTION SHOULD BE MULTIPLIED BY 1.5 $\pm$.2 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = K^+ (1420)^+ \text{ PROTON} \dots$ [TABLE 18]
 $K^+ (1420)^+ = \dots$ $K^0 P_{10}$ P_{10} [1]

LABORATORY	FRACTION
BEAM MOMENTUM	
GEV/C	PER CENT [2]
5.000 $\pm$.015	2.0 $\pm$ 1.1

- (1) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH [MASS = 1.415 GEV; WIDTH = .125 GEV] AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 (2) THIS FRACTION SHOULD BE MULTIPLIED BY 1.5 $\pm$.2 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = K^+ (1225)^+ \text{ PROTON} \dots$ [TABLE 18]
 $K^+ (1225)^+ = \dots$ $K^+ (890)^+ P_{10}$ [1]

LABORATORY	FRACTION
BEAM MOMENTUM	
GEV/C	PER CENT [2]
5.000 $\pm$.015	7.1 $\pm$ 1.1

- (1) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH [MASS = 1.225 GEV; WIDTH = .220 GEV] AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 (2) THIS FRACTION SHOULD BE MULTIPLIED BY 1.5 $\pm$.2 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = K^*(1225)^+ \text{ PROTON.}$ [TABLE 18]
 $K^*(1225)^+ = K^*(890)^+ \text{ P}^0$ (1)

LABORATORY BEAM MOMENTUM GEV/C	FRACTION PER CENT (2)
5.000 $\pm$.015	10.5 $\pm$ 1.2

- (1) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH (MASS = 1.225 GEV; WIDTH = .220 GEV) AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 (2) THIS FRACTION SHOULD BE MULTIPLIED BY 1.5 $\pm$.2 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = K^*(1225)^+ \text{ PROTON.}$ [TABLE 18]
 $K^*(1225)^+ = K^0 \text{ RHO}^+ (1)$

LABORATORY BEAM MOMENTUM GEV/C	FRACTION PER CENT (2)
5.000 $\pm$.015	13.5 $\pm$ 1.5

- (1) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH (MASS = 1.225 GEV; WIDTH = .220 GEV) AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 (2) THIS FRACTION SHOULD BE MULTIPLIED BY 1.5 $\pm$.2 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = K^*(890)^0 \text{ PROTON P}^0.$ [TABLE 18]
 $K^*(890)^0 = K^0 \text{ P}^0$ (1)

LABORATORY BEAM MOMENTUM GEV/C	FRACTION PER CENT (2)
5.000 $\pm$.015	.9

- (1) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 (2) THIS FRACTION SHOULD BE MULTIPLIED BY 1.5 $\pm$.2 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = K^*(1420)^+ \text{ PROTON.}$ [TABLE 18]
 $K^*(1420)^+ = K^0 \text{ RHO}^+ (1)$

LABORATORY BEAM MOMENTUM GEV/C	FRACTION PER CENT (2)
5.000 $\pm$.015	1.

- (1) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 (2) THIS FRACTION SHOULD BE MULTIPLIED BY 1.5 $\pm$.2 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = \text{PROTON K}^0 \text{ P}^0 \text{ P}^0.$ [TABLE 18]
 CROSS SECTION IS ONLY FOR THE NON-RESONANT PRODUCTION OF THESE PARTICLES.

LABORATORY BEAM MOMENTUM GEV/C	FRACTION PER CENT (1)
5.000 $\pm$.015	7.3 $\pm$ 1.0

- (1) THIS FRACTION SHOULD BE MULTIPLIED BY 1.5 $\pm$.2 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = \text{DELTA}^+ \text{ K}^0 \text{ P}^0.$ [TABLE 1C]
 $\text{DELTA}^+ = \text{NEUTRON P}^0$ (1)

LABORATORY BEAM MOMENTUM GEV/C	FRACTION PER CENT (2)
5.000 $\pm$.015	4.

- (1) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 (2) THIS FRACTION SHOULD BE MULTIPLIED BY .40 $\pm$.05 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = \text{DELTA}^+ K^*(890)^+.$ [TABLE 1C]
 $\text{DELTA}^+ = \text{NEUTRON P}^0$ (1)
 $K^*(890)^+ = K^0 \text{ P}^0$ (2)

LABORATORY BEAM MOMENTUM GEV/C	FRACTION PER CENT (3)
5.000 $\pm$.015	20.7 $\pm$ 3.0

- (1) FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 (2) FITTED FOR MASS AND/OR WIDTH (MASS = .898 GEV; WIDTH = .053 GEV), AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.
 (3) THIS FRACTION SHOULD BE MULTIPLIED BY .40 $\pm$.05 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = \Delta K^*(1420)^+ +$ [TABLE 1C]
 $\Delta K^+ + \text{ NEUTRON } \pi^+ [1]$
 $K^*(1420)^+ = K^0 \pi^+ [2]$

LABORATORY FRACTION
 BEAM MOMENTUM PER CENT [3]
 GEV/C
 5.000 $\pm$.015 2.

- [1] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 [2] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH [MASS = 1.415 GEV; WIDTH = .125 GEV] AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 [3] THIS FRACTION SHOULD BE MULTIPLIED BY .40 $\pm$.05 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = K^*(890)^+ \text{ NEUTRON } \pi^+ +$ [TABLE 1C]
 $K^*(890)^+ = K^0 \pi^+ [1]$

LABORATORY FRACTION
 BEAM MOMENTUM PER CENT [2]
 GEV/C
 5.000 $\pm$.015 13.2 $\pm$ 3.5

- [1] FITTED FOR MASS AND/OR WIDTH [MASS = .898 GEV; WIDTH = .053 GEV], AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.
 [2] THIS FRACTION SHOULD BE MULTIPLIED BY .40 $\pm$.05 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = K^*(1420)^+ \text{ NEUTRON } \pi^+ +$ [TABLE 1C]
 $K^*(1420)^+ = K^0 \pi^+ [1]$

LABORATORY FRACTION
 BEAM MOMENTUM PER CENT [2]
 GEV/C
 5.000 $\pm$.015 8. $\pm$ 3.

- [1] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH [MASS = 1.415 GEV; WIDTH = .125 GEV] AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 [2] THIS FRACTION SHOULD BE MULTIPLIED BY .40 $\pm$.05 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = N^*(1590)^+ K^*(890)^+ +$ [TABLE 1C]
 $N^*(1590)^+ = \text{ NEUTRON } \pi^+ [1]$
 $K^*(890)^+ = K^0 \pi^+ [2]$

LABORATORY FRACTION
 BEAM MOMENTUM PER CENT [3]
 GEV/C
 5.000 $\pm$.015 17. $\pm$ 3.

- [1] FITTED FOR MASS AND/OR WIDTH, AND THEN TOOK ONLY EVENTS ABOVE (FITTED) BACKGROUND.
 [2] FITTED DISTRIBUTION WITH FIXED MASS AND WIDTH AND TOOK EVENTS ONLY ABOVE (FITTED) BACKGROUND.
 [3] THIS FRACTION SHOULD BE MULTIPLIED BY .40 $\pm$.05 TO GET CROSS SECTION IN MILLIBARNS.

CROSS SECTION FOR $K^+ \text{ PROTON} = \text{ NEUTRON } \pi^+ \pi^+ K^0$. [TABLE 1C]
 CROSS SECTION IS ONLY FOR THE NON-RESONANT PRODUCTION OF THESE PARTICLES.

LABORATORY FRACTION
 BEAM MOMENTUM PER CENT [1]
 GEV/C
 5.000 $\pm$.015 41.1 $\pm$ 5.0

- [1] THIS FRACTION SHOULD BE MULTIPLIED BY .40 $\pm$.05 TO GET CROSS SECTION IN MILLIBARNS.

83 INTERFERENCE ANALYSIS OF THE $K^*(1400)$. [PHYS. REV. LETTERS 21, 1842 (1968)]

P. ANTICH, A. CALLAHAN, R. CARSON, B. COX, D. DENEGR, L. ETTLINGER, D. GILLESPIE, G. LUSTE, R. MERCER, A. PEVSNER, R. ZDANIS (JOHNS HOPKINS UNIV., BALTIMORE, MD., USA)

ABSTRACT A STUDY OF THE $K^+ \pi^-$ DECAY MODE OF THE $K^*(1400)$ PRODUCED WITH AN N^{*++} HAS BEEN MADE USING THE REACTION $K^+ p \rightarrow K^+ \pi^- \pi^+ p$ AT 5.5 BEV/C. WE HAVE FITTED THE $K^*(1400)$ ANGULAR DISTRIBUTIONS USING A MODEL INVOLVING INTERFERENCE BETWEEN 1^- AND $2^+ K^+ \pi^-$ STATES. IN THIS MODEL A CONSIDERABLE AMOUNT OF 1^- IS NEEDED IN THE $K^*(1400)$ MASS REGION, SUGGESTING THAT THERE IS A $J^P = 1^-$ RESONANCE UNDER THE $K^*(1400)$. SUCH A $J^P = 1^-$ RESONANCE COULD BE THE FIRST DAUGHTER OF THE $K^*(1400)$.

ADDITIONAL FOOTNOTE REFERENCES

PHYS. LETTERS 18, 76 (1965), PHYS. REV. LETTERS 19, 1052 (1967), NUOVO CIMENTO 39, 417 (1965), PHYS. REV. 139, 8428 (1955), BULL. AM. PHYS. SOC. 13, 114 (1968), PHYS. REV. 157, 1371 (1967), NUOVO CIMENTO 33, 309 (1964), PHYS. REV. 165, 1317 (1968), REV. MOD. PHYS. 40, 77 (1968), PHYS. LETTERS 258, 44 (1967), AND PHYS. REV. LETTERS 20, 1318 (1968).

ARTICLE READ BY ODETTE BENARY IN 4/69, AND VERIFIED BY LEROY PRICE.

BEAM IS K^+ ON PROTON AT 5.5 GEV/C.

THIS EXPERIMENT USES THE B.N.L. 80 IN. (H) BUBBLE CHAMBER.

KEY WORDS = MASS SPECTRUM $K^*(1420)$ ANGULAR DISTRIBUTION MODELS DENSITY MATRIX

* * * * *
 * NO DATA PUNCHED FOR THIS ARTICLE *
 * * * * *

84 STUDY OF MULTI-BODY FINAL STATES IN K⁺P REACTIONS AT INCIDENT MOMENTUM OF 5 GEV/C. [NUC. PHYS. B9, 295 (1969)]

G. BASSOMPIERRE, Y. GOLDSCHMIDT-CLERMONT, A. GRANT, V. P. HENRI, B. JONGEJANS, D. LINGLI, N. F. MULLER, H. PIOTROWSKA, J. M. PERREAU, I. SAITOV, R. L. SEKULIN, G. W. DLF (EUROPEAN ORG. FOR NUC. RES., GENEVA, SWITZERLAND)
 W. DEBAERE, J. DEBAISIEUX, E. DEWOLF, P. DUFOUR, F. GRARD, J. HEUGHEBAERT, L. PAPE, P. PEETERS, F. VERBEURE, R. WINDMOLDERS (INST. INTERUNIV. DES SCI. NUC., BRUXELLES, BELGIUM)

ABSTRACT RESULTS ARE GOOD ON THE COMPARISON OF THE PREDICTIONS OF THE MULTI-PERIPHERAL MODEL OF CHAY, LOSKIEWICZ AND ALLISON WITH THE EXPERIMENTAL DATA OBTAINED FROM A STUDY OF THE INTERACTIONS OF 5 GEV/C K⁺ MESONS WITH PROTONS. COMPARISONS HAVE BEEN MADE IN THE THREE- FOUR- FIVE- AND SIX-BODY FINAL STATES CONTAINING A PROTON, KAON AND PIONS. WITH THE ADDITION OF RESONANCE PRODUCTION TO THE MODEL, ADEQUATE PREDICTIONS OF EXPERIMENTAL DISTRIBUTIONS ARE OBTAINED. AN ATTEMPT IS MADE TO DESCRIBE THE ENERGY VARIATION OF THE THREE- AND FOUR-BODY FINAL-STATE CROSS SECTIONS.

CROSS REFERENCES
 ANALYSIS OF DATA FROM NUOVO CIMENTO 57A, 93 (1968), NUOVO CIMENTO 48, 589 (1967), CERN W.505 (1968), PHYS. LETTERS 26B, 30 (1967), UCRL 11446 (1964), UCRL 16370 (1965), PHYS. REV. LETTERS 11, 93 (1963), PHYS. REV. 133, 8220 (1964), PHYS. REV. 142, 913 (1966), PHYS. REV. LETTERS 15, 737 (1965), NUOVO CIMENTO 36, 1101 (1965), NUOVO CIMENTO 51A, 401 (1967), CERN MADRUM CONFERENCE 2 172, AND BULL. AM. PHYS. SOC. 13, 114 (1968).

ARTICLE READ BY ODETTE BENARY IN 3/69, AND VERIFIED BY LEROY PRICE.

BEAM IS K⁺ ON PROTON AT 5 GEV/C.

THIS EXPERIMENT USES BUBBLE CHAMBERS.

KEY WORDS = MODELS

 * NO DATA PUNCHED FOR THIS ARTICLE *

85 K⁺D CHARGE EXCHANGE REACTION FROM 52 TO 456 MEV. [PHYS. REV. LETTERS 7, 378 (1961)]

W. E. SLATER, D. H. STORK, H. K. TICHON (UNIV. OF CALIF., LOS ANGELES, CALIF., USA)
 W. LEE, W. CHINOWSKY, G. GOLDBABER, S. GOLDBABER, T. A. O'HALLORAN (U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA, AND UNIV. OF CALIFORNIA, BERKELEY, CALIF., USA)

ADDITIONAL FOOTNOTE REFERENCES
 PHYS. REV. 112, 624 (1958), PHYS. REV. 113, 1303 (1959), PHYS. REV. 130, 1998 (1960), NUOVO CIMENTO 12, 241 (1959), PHYS. REV. 118, 300 (1960), NUOVO CIMENTO 19, 179 (1961), LRL INTERDEPARTMENTAL REPORT BEV 483 (1960), PHYS. REV. LETTERS 2, 266 (1959), ANNUAL REV. OF NUCLEAR SCIENCE 9, 61 (1959), ROCHESTER CONFERENCE 451 (1960), PHYS. REV. 84, 1084 (1951), PHYS. REV. 105, 256 (1957), PHYS. REV. 105, 247 (1957), PHYS. REV. 115, 1727 (1959), NUOVO CIMENTO 11, 670 (1959), NUOVO CIMENTO 14, 722 (1959), UCRL 9691 (1961), PHYS. REV. LETTERS 7, 188 (1961), PHYS. REV. 118, 553 (1960), AND PHYS. REV. LETTERS 7, 182 (1961).

ARTICLE READ BY ODETTE BENARY IN 6/69, AND VERIFIED BY ROGER BLAND.

BEAM IS K⁺ ON DEUTERON FROM .233 TO .811 GEV/C. (BEAM KINETIC ENERGY = .052 TO .456 GEV)

THIS EXPERIMENT USES THE L.R.L. 15 IN. (H) BUBBLE CHAMBER.

GENERAL COMMENTS ON THIS ARTICLE

1 THE K⁺ PATH LENGTH WAS OBTAINED BY TWO INDEPENDENT METHODS 1) BY DIRECT MEASUREMENTS OF THE K⁺ TRACKS IN EVERY TENTH PICTURE 2) BY INFERRING THE TOTAL PATH LENGTH FROM THE NUMBER OF TAU DECAYS.

KEY WORDS = CHARGE EXCHANGE CROSS SECTION DIFFERENTIAL CROSS SECTION PHASE SHIFT

COMPOUND KEY WORDS = CHARGE EXCHANGE CROSS SECTION

CROSS SECTION FOR K⁺ DEUTERON = K⁰ PROTON PROTON. [TABLE 1]

(IT IS NOT CLEAR WHETHER OR NOT A CUTOFF ON THE SPECTATOR PROTON WAS APPLIED)

LABORATORY BEAM ENERGY GEV		MILLI-BARNS	NO. EVENTS
.052 ± .017		1.0 ± .4	13
.100 ± .013		2.7 ± .4	46
.127 ± .011		3.1 ± .4	65
.230 ± .011		6.5 ± .6	161
.315 ± .006		6.7 ± .6	216
.456 ± .005		6.6 ± .7	196

DIFFERENTIAL CROSS SECTION FOR K⁺ DEUTERON = K⁰ PROTON PROTON. [FIGURE 2A]

LABORATORY BEAM ENERGY = .100 ± .013 GEV.

 * THIS DATA WAS READ FROM A GRAPH *

COS(THETA)		D-SIGMA/D-OMEGA MB/SR
MIN	MAX	
-.6	-.4	.05 ± .10
-.4	-.2	.08 ± .10
-.2	.0	.10 ± .10
.0	.2	.20 ± .15
.2	.4	.25 ± .15
.4	.6	.40 ± .15
.6	.8	.80 ± .20
.8	1.0	.40 ± .20

THETA IS THE ANGLE THAT THE K⁰ MAKES WITH THE BEAM IN THE LAB.

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ DEUTERON} \rightarrow K^0 \text{ PROTON PROTON.}$ (FIGURE 2B)
 LABORATORY BEAM ENERGY = .230 $\pm$.011 GEV.

 THIS DATA WAS READ FROM A GRAPH .

COS(THETA)		D-SIGMA/D-OMEGA MB/SR
MIN	MAX	
-1.0	-.8	.00 $\pm$.05
		-.05
-.8	-.6	.02 $\pm$.10
		-.10
-.6	-.4	.02 $\pm$.10
		-.10
-.4	-.2	.20 $\pm$.20
		-.10
-.2	.0	.30 $\pm$.20
		-.10
.0	.2	.46 $\pm$.20
		-.15
.2	.4	.50 $\pm$.20
		-.20
.4	.6	1.00 $\pm$.20
		-.20
.6	.8	1.85 $\pm$.30
		-.30
.8	1.0	.90 $\pm$.20
		-.20

THETA IS THE ANGLE THAT THE K^0 MAKES WITH THE BEAM IN THE LAB.

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ DEUTERON} \rightarrow K^0 \text{ PROTON PROTON.}$ (FIGURE 2C)
 LABORATORY BEAM ENERGY = .315 $\pm$.006 GEV.

 THIS DATA WAS READ FROM A GRAPH .

COS(THETA)		D-SIGMA/D-OMEGA MB/SR
MIN	MAX	
-1.0	-.8	.00 $\pm$.05
		-.10
-.8	-.6	.10 $\pm$.10
		-.15
-.6	-.4	.15 $\pm$.15
		-.15
-.4	-.2	.20 $\pm$.15
		-.15
-.2	.0	.30 $\pm$.20
		-.15
.0	.2	.28 $\pm$.15
		-.20
.2	.4	.43 $\pm$.20
		-.25
.4	.6	1.15 $\pm$.25
		-.20
.6	.8	1.40 $\pm$.20
		-.20
.8	1.0	1.35 $\pm$.20

THETA IS THE ANGLE THAT THE K^0 MAKES WITH THE BEAM IN THE LAB.

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ DEUTERON} \rightarrow K^0 \text{ PROTON PROTON.}$ (FIGURE 2D)
 LABORATORY BEAM ENERGY = .456 $\pm$.005 GEV.

 THIS DATA WAS READ FROM A GRAPH .

COS(THETA)		D-SIGMA/D-OMEGA MB/SR
MIN	MAX	
-1.0	-.8	.00 $\pm$.05
		-.08
-.8	-.6	.08 $\pm$.10
		-.10
-.6	-.4	.10 $\pm$.10
		-.15
-.4	-.2	.15 $\pm$.10
		-.15
-.2	.0	.30 $\pm$.15
		-.20
.0	.2	.52 $\pm$.20
		-.20
.2	.4	.40 $\pm$.20
		-.20
.4	.6	.65 $\pm$.20
		-.30
.6	.8	1.30 $\pm$.30
		-.30
.8	1.0	1.70 $\pm$.30

THETA IS THE ANGLE THAT THE K^0 MAKES WITH THE BEAM IN THE LAB.

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ DEUTERON} \rightarrow K^0 \text{ PROTON PROTON.}$ (FIGURE 2A*)
 LABORATORY BEAM ENERGY = .100 $\pm$.013 GEV.

 THIS DATA WAS READ FROM A GRAPH .

COS(THETA)		D-SIGMA/D-OMEGA MB/SR
MIN	MAX	
-1.0	-.6	.11 $\pm$.04
		-.05
-.6	-.2	.25 $\pm$.05
		-.07
-.2	.2	.40 $\pm$.07
		-.06
.2	.6	.29 $\pm$.06
		-.04
.6	1.0	.16 $\pm$.04

THETA IS THE ANGLE THAT THE K^0 MAKES WITH THE BEAM IN THE GRAND C.M.

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ DEUTERON} \rightarrow K^0 \text{ PROTON PROTON}$. (FIGURE 2B')
 LABORATORY BEAM ENERGY = $.230 \pm .011 \text{ GEV}$.

.....
 . THIS DATA WAS READ FROM A GRAPH .

COS(THETA)		D-SIGMA/D-OMEGA MB/SR
MIN	MAX	
-1.0	-.7	.15 $\pm$.07
-.7	-.4	.44 .10
-.4	-.1	.60 .12
-.1	.2	.73 .13
.2	.5	.84 .14
.5	.8	.58 .12
.8	1.0	.20 .08

THETA IS THE ANGLE THAT THE K^0 MAKES WITH THE BEAM IN THE GRAND C.M.

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ DEUTERON} \rightarrow K^0 \text{ PROTON PROTON}$. (FIGURE 2C')
 LABORATORY BEAM ENERGY = $.315 \pm .006 \text{ GEV}$.

.....
 . THIS DATA WAS READ FROM A GRAPH .

COS(THETA)		D-SIGMA/D-OMEGA MB/SR
MIN	MAX	
-1.0	-.7	.32 $\pm$.08
-.7	-.4	.44 .08
-.4	-.1	.50 .10
-.1	.2	.84 .12
.2	.5	.64 .12
.5	.8	.60 .10
.8	1.0	.42 .10

THETA IS THE ANGLE THAT THE K^0 MAKES WITH THE BEAM IN THE GRAND C.M.

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ DEUTERON} \rightarrow K^0 \text{ PROTON PROTON}$. (FIGURE 2D')
 LABORATORY BEAM ENERGY = $.456 \pm .005 \text{ GEV}$.

.....
 . THIS DATA WAS READ FROM A GRAPH .

COS(THETA)		D-SIGMA/D-OMEGA MB/SR
MIN	MAX	
-1.0	-.7	.33 $\pm$.09
-.7	-.4	.44 .09
-.4	-.1	.50 .10
-.1	.2	.60 .12
.2	.5	.72 .12
.5	.8	.61 .13
.8	1.0	.50 .12

THETA IS THE ANGLE THAT THE K^0 MAKES WITH THE BEAM IN THE GRAND C.M.

86 K^+N CHARGE-EXCHANGE SCATTERING AT 2.3 BEV/C [PHYS. REV. LETTERS 15, 734 (1965)]

I. BUTTERWORTH, J. L. BROWN, G. GOLDBERGER, S. GOLDBERGER, A. A. MIRATA, J. A. KADYK, B. M. SCHWARZSCHILD, G. M. TRILLING (U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA, AND UNIV. OF CALIFORNIA, BERKELEY, CALIF., USA)

CROSS REFERENCES
 SEE ALSO UCRL 17572.

ADDITIONAL FOOTNOTE REFERENCES
 NUCLEAR INSTRUMENTS AND METHODS 20, 37 (1963), UCRL 9691 (1961), PHYS. REV. LETTERS 7, 182 (1961), PHYS. LETTERS 16, 328 (1965), AND PHYS. REV. 139, 81336 (1965).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 4/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K^+ ON DEUTERON AT 2.3 GEV/C.

THIS EXPERIMENT USES THE B.N.L. 20 IN. (H) BUBBLE CHAMBER. A TOTAL OF 100000 PICTURES ARE REPORTED ON.

GENERAL COMMENTS ON THIS ARTICLE
 1 NORMALIZED TO TAU DECAYS.
 2 USED ONLY EVENTS WITH SPECTATOR MOMENTUM $> 300 \text{ MEV/C}$.

KEY WORDS - CROSS SECTION ANGULAR DISTRIBUTION

CROSS SECTION FOR $K^+ \text{ DEUTERON} \rightarrow K^0 \text{ PROTON PROTON}$.

LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	NO. EVENTS
2.3	$1.50 \pm .15$	257

DIFFERENTIAL CROSS SECTION FOR K^+ DEUTERON $\rightarrow$ K^0 PROTON PROTON. (FIGURE 1)

LABORATORY BEAM MOMENTUM = 2.3 GEV/C.

.....
 . THIS DATA WAS READ FROM A GRAPH .

COS(THETA)	D-SIGMA/D-OMEGA MB/SR
.995 $\pm$.005	3.20 $\pm$ 1.00
	- 1.00
.990 .005	3.00 $\pm$ 1.00
	- 1.00
.980 .005	4.00 $\pm$ 1.00
	- 1.00
.970 .005	2.50 $\pm$ 1.00
	- 1.00
.950 .010	2.00 $\pm$ 1.00
	- 1.00
.930 .015	.80 $\pm$.30
	- .20
.860 .050	.20 $\pm$.10
	- .05
.710 .100	.03 $\pm$.02
	- .01

THETA IS THE ANGLE THAT THE K^0 MAKES WITH THE BEAM IN THE LAB.

DIFFERENTIAL CROSS SECTION FOR K^+ DEUTERON $\rightarrow$ K^0 PROTON PROTON. (FIGURE 2)

LABORATORY BEAM MOMENTUM = 2.3 GEV/C.

.....
 . THIS DATA WAS READ FROM A GRAPH .

-T (GEV/C)**2	D-SIGMA/D-T MB/(GEV/C)**2
.04 $\pm$.04	1.60 $\pm$.40
.11 .03	2.00 .40
.17 .03	2.20 .40
.23 .03	2.00 .40
.31 .05	1.50 .30
.41 .05	1.40 .20
.51 .05	1.30 .30
.60 .05	1.40 .30
.75 .10	.60 .20
.95 .10	.30 .10
1.45 .35	.09 .02
2.60 .80	.08 .02

T IS THE MOMENTUM TRANSFER BETWEEN THE (INCOMING K^+) AND THE (K^0).

87 RESOLUTION OF THE $T = 0$ FERMI-YANG AMBIGUITY IN K NUCLEON SCATTERING. [BNL 13616 (1969)]

A.K.RAY, R.W.BURRIS, H.E.FISK, R.W.KRAEMER (CARNEGIE INST. OF TECH., PITTSBURGH, PA., USA)
 D.G.MILL, M.SAKITT (BROOKHAVEN NAT. LAB., UPTON, L.I., N. Y., USA)

ABSTRACT FROM AN ANALYSIS OF THE PROTON POLARIZATION AND DIFFERENTIAL CROSS SECTION IN $K^+ N$ CHARGE EXCHANGE SCATTERING AT 600 MEV/C A YANG TYPE PHASE SHIFT SOLUTION FOR $T = 0$ SCATTERING IS ESTABLISHED.

ADDITIONAL FOOTNOTE REFERENCES

PHYS. REV. LETTERS 9, 135 (1962), PHYS. REV. LETTERS 7, 378 (1961), PHYS. REV. 134, B111 (1964), PHYS. REV. 121, 1229 (1961), NUOVO CIMENTO 23, 1163 (1962), NUOVO CIMENTO 11, 670 (1959), PHYS. REV. LETTERS 20, 1272 (1968), PHYS. REV. LETTERS 7, 187 (1961), PHYS. REV. 135, B191 (1964), NUOVO CIMENTO 32A, 371 (1967), NUOVO CIMENTO 45A, 792 (1966), PHYS. REV. LETTERS 17, 599 (1966), AND PHYS. REV. LETTERS 19, 1094 (1967).

ARTICLE READ BY ODETTE BENARY IN 7/69, AND VERIFIED BY ROGER BLAND.

BEAM IS K^+ ON DEUTERON AT .6 GEV/C.

THIS EXPERIMENT USES THE B.N.L. 30 IN. (H) BUBBLE CHAMBER.

KEY WORDS = CHARGE EXCHANGE DIFFERENTIAL CROSS SECTION POLARIZATION FITS PHASE SHIFT

COMPOUND KEY WORDS = CHARGE EXCHANGE DIFFERENTIAL CROSS SECTION FITS DIFFERENTIAL CROSS SECTION
 FITS POLARIZATION

DIFFERENTIAL CROSS SECTION FOR K^+ DEUTERON $\rightarrow$ K^0 PROTON PROTON. (FIGURE 1A)

LABORATORY BEAM MOMENTUM = .6 GEV/C.

.....
 . THIS DATA WAS READ FROM A GRAPH .

COS(THETA)	D-SIGMA/D-OMEGA MB/SR
MIN MAX	
-1.0 -.8	.22 $\pm$.02
-.8 -.6	.34 .02
-.6 -.4	.47 .02
-.4 -.2	.55 .02
-.2 .0	.59 .04
.0 .2	.68 .03
.2 .4	.67 .02
.4 .6	.69 .02
.6 .8	.60 .02
.8 1.0	.38 .02

THETA IS THE ANGLE THAT THE K^0 MAKES WITH THE BEAM IN THE GRAND C.M.

POLARIZATION FOR K^+ DEUTERON $\rightarrow$ K^0 PROTON PROTON. (FIGURE 21.)

LABORATORY BEAM MOMENTUM = .6 GEV/C.

(THE POLARIZATION DATA ARE OBTAINED FROM 853 SELECTED EVENTS IN WHICH THE NON-SPECTATOR PROTON UNDERGOES A SECOND SCATTERING.)

THIS DATA WAS READ FROM A GRAPH

COS(THETA)	POLARIZATION	NO. EVENTS
-.00	-.92 $\pm$.10	80
-.40	-.50 $\pm$.38	153
-.05	-.75 $\pm$.35	153
.25	-.95 $\pm$.30	263
.55	.10 $\pm$.65	100

THETA IS THE ANGLE THAT THE K^0 MAKES WITH THE BEAM IN THE GRAND C.M.
THE POLARIZATION IS OF THE PROTON ALONG THE NORMAL TO THE PRODUCTION PLANE IN THE GRAND C.M.

88 QUASI-TWO-BODY REACTIONS IN 13 GEV/C K^+ P INTERACTIONS. (NUC. PHYS. 98, 333 (1968))

J.C. BERLINGHIERI, M.S. FARBER, T. FERBEL, B.E. FORMAN, A.C. MELISSINOS, P.F. SLATTERY, H. YUTA (UNIV. OF ROCHESTER, ROCHESTER, N. Y., USA)

ABSTRACT WE HAVE STUDIED THE FINAL STATES $P K^+(890)$, $K^0 N^{*++}(1236)$, AND $K^0(890) N^{*++}(1236)$ AS PRODUCED IN 12.7 GEV/C K^+ P INTERACTIONS. THE DENSITY MATRIX ELEMENTS AND T-DISTRIBUTIONS HAVE BEEN OBTAINED AND ARE COMPARED TO THEORETICAL MODELS.

CROSS REFERENCES

SEE ALSO CERN 66-17 (1966), NUOVO CIMENTO 51A, 401 (1967), AND NUOVO CIMENTO 49A, 9 (1967).
THIS ARTICLE SUPERSEDES PART OF CERN HADRON CONFERENCE 172 (1968).

ADDITIONAL FOOTNOTE REFERENCES

NUOVO CIMENTO 50A, 377 (1967), AND PHYS. REV. 134, 81099.

ARTICLE READ BY ODETTE BENARY IN 2/69, AND VERIFIED BY ROGER BLAND.

BEAM IS K^+ ON PROTON AT 12.7 GEV/C.

THIS EXPERIMENT USES THE B.N.L. 80 IN. (H) BUBBLE CHAMBER.

KEY WORDS = CROSS SECTION DALITZ PLOT MASS SPECTRUM ANGULAR DISTRIBUTION DENSITY MATRIX MODELS
DELTA(1238) $K^+(890)$ $K^+(1420)$

CROSS SECTION FOR K^+ PROTON $\rightarrow$ PROTON $K^+(890)^+$. (TABLE 1)
 $K^+(890)^+ \rightarrow K^0 \pi^+ [1]$

(THIS DATA REPLACES VALUES GIVEN EARLIER IN BERLINGHIERI, ET. AL., CERN HADRON CONFERENCE 172 (1968))

LABORATORY
BEAM MOMENTUM
GEV/C
12.7
MICRO-BARNS
51. $\pm$ 12.

[1] COUNTED ONLY EVENTS ABOVE BACKGROUND.

CROSS SECTION FOR K^+ PROTON $\rightarrow$ DELTA $^{++}$ K^0 . (TABLE 1)
DELTA $^{++} \rightarrow$ PROTON $\pi^+ [1]$

(THIS DATA REPLACES THE VALUE GIVEN EARLIER IN BERLINGHIERI, ET. AL., CERN HADRON CONFERENCE 172 (1968))

LABORATORY
BEAM MOMENTUM
GEV/C
12.7
MICRO-BARNS
44. $\pm$ 7.

[1] COUNTED ONLY EVENTS ABOVE BACKGROUND.

CROSS SECTION FOR K^+ PROTON $\rightarrow$ DELTA $^{++}$ $K^+(890)^0$. (TABLE 1)
DELTA $^{++} \rightarrow$ PROTON $\pi^+ [1]$
 $K^+(890)^0 \rightarrow K^+ \pi^- [2]$

LABORATORY
BEAM MOMENTUM
GEV/C
12.7
MICRO-BARNS
100. $\pm$ 30.

[1] COUNTED ONLY EVENTS ABOVE BACKGROUND.
[2] COUNTED ONLY EVENTS ABOVE BACKGROUND.

CROSS SECTION FOR K^+ PROTON $\rightarrow$ $K^+(1420)^+$ PROTON. (PAGE 333)
 $K^+(1420)^+ \rightarrow K^0 \pi^+ [1]$

LABORATORY
BEAM MOMENTUM
GEV/C
12.7
MICRO-BARNS
20. $\pm$ 5.

[1] COUNTED ONLY EVENTS ABOVE BACKGROUND.

CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow \Delta^{++} K^{*}(1420)0$. (PAGE 3331)
 $\Delta^{++} \rightarrow \text{PROTON } \pi^{+}$ (11)
 $K^{*}(1420)0 \rightarrow K^+ \pi^{-}$ (12)

LABORATORY
 BEAM MOMENTUM
 GEV/C
 12.7

MICRO-BARNS (3)
 25.

- (1) COUNTED ONLY EVENTS ABOVE BACKGROUND.
 (2) COUNTED ONLY EVENTS ABOVE BACKGROUND.
 (3) VALUE IS APPROXIMATE ONLY.

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^{*}(890)^+ \text{ PROTON}$. (FIGURE 3A1)
 $K^{*}(890)^+ \rightarrow K^0 \pi^{+}$ (11)

LABORATORY BEAM MOMENTUM = 12.7 GEV/C.

 . THIS DATA WAS READ FROM A GRAPH .

-T (GEV/C)**2	NO. EVENTS
.12 +- .05	12.0 +- 4.0
.22 .05	17.0 3.0
.32 .05	9.0 3.0
.42 .05	8.0 3.0
.67 .20	2.5 .7

T IS THE MOMENTUM TRANSFER BETWEEN THE [INCOMING K^+] AND THE [$K^{*}(890)^+$].

(1) USED SIMPLE MASS CUT.

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow \Delta^{++} K^0$. (FIGURE 3B)

LABORATORY BEAM MOMENTUM = 12.7 GEV/C.

 . THIS DATA WAS READ FROM A GRAPH .

-T (GEV/C)**2	NO. EVENTS
.065 +- .050	60.0 +- 12.0
.165 .050	23.0 9.0
.315 .100	11.0 2.5
.565 .150	4.0 2.0

T IS THE MOMENTUM TRANSFER BETWEEN THE [INCOMING K^+] AND THE [K^0].

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^{*}(890)0 \Delta^{++}$. (FIGURE 3C)
 $K^{*}(890)0 \rightarrow K^+ \pi^{-}$ (11)

LABORATORY BEAM MOMENTUM = 12.7 GEV/C.

 . THIS DATA WAS READ FROM A GRAPH .

-T (GEV/C)**2	NO. EVENTS
.07 +- .01	30.00 +- 5.00
.09 .01	15.00 4.00
.12 .02	6.00 2.00
.19 .05	3.20 1.00
.30 .06	1.60 .50
.81 .25	.11 .09

T IS THE MOMENTUM TRANSFER BETWEEN THE [INCOMING K^+] AND THE [$K^{*}(890)0$].

(1) USED SIMPLE MASS CUT.

89 SINGLE-PION PRODUCTION BY 1.96-BEV/C K^+ MESONS. [PHYS. REV. 142, 913 (1966)]

S. GOLDHABER, W. CHINDENSKY, G. GOLDHABER, T. A. O'HALLORAN [U.C. LAWRENCE RAD. LAB., BERKELEY, CALIF., USA, AND UNIV. OF CALIFORNIA, BERKELEY, CALIF., USA]

ABSTRACT SINGLE-PION PRODUCTION BY K^+ MESONS AT A MOMENTUM OF 1.96 BEV/C IS SHOWN TO LEAD PREDOMINANTLY TO K^+P AND N^+K FINAL STATES. STUDY OF THE DYNAMICS OF THE REACTION SHOWS THAT THE N^+K FINAL STATE IS PRODUCED ALMOST ENTIRELY VIA VECTOR EXCHANGE, WHEREAS THE K^+P FINAL STATE SEEMS TO BE PRODUCED PARTLY VIA PION EXCHANGE AND PARTLY VIA VECTOR EXCHANGE. NUMERICAL VALUES IN TERMS OF GOTTFRIED AND JACKSON'S PARAMETERIZATION OF THE SPIN-DENSITY MATRIX ARE GIVEN FOR THE K^+P FINAL STATE. COMPARISON WITH SIMILAR ANALYSES AT ENERGIES BELOW AND ABOVE THIS EXPERIMENT INDICATES THAT A STRONG ENERGY DEPENDENCE OF THE VECTOR-MESON-EXCHANGE CONTRIBUTION TO THESE PROCESSES AT 3 BEV/C DO NOT FIT THE DATA AT THIS ENERGY.

CROSS REFERENCES
CONTINUATION OF PREVIOUS EXPERIMENT IN PHYS. REV. 139, 1141 (1963).
SOME OF THIS DATA RENORMALIZED IN UCRL 18131 (1968).

ADDITIONAL FOOTNOTE REFERENCES
PHYS. REV. LETTERS 11, 90 (1963), PHYS. REV. 134, B1099 (1964), BULL. AM. PHYS. SOC. 8, 20 (1963), ATHENS CONFERENCE 92 (1963), PHYS. REV. LETTERS 11, 93 (1963), PHYS. LETTERS 6, 309 (1963), PHYS. REV. 133, B220 (1964), NUOVO CIMENTO 33, 309 (1964), PHYS. LETTERS 9, 144 (1964), NUCLEAR INSTRUMENTS AND METHODS 20, 37 (1963), NUCLEAR INSTRUMENTS AND METHODS 20, 42 (1963), PHYS. REV. 139, B1411 (1965), UCRL 9099, PHYS. REV. LETTERS 10, 351 (1963), NUOVO CIMENTO 33, 906 (1964), NUOVO CIMENTO 34, 1841 (1964), NUOVO CIMENTO 26, 186 (1962), PHYS. REV. LETTERS 12, 82 (1964), PHYS. REV. LETTERS 12, 399 (1964), PHYS. REV. LETTERS 12, 627 (1964), NUOVO CIMENTO 34, 735 (1964), AND PHYS. LETTERS 6, 62 (1963).

ARTICLE READ BY NAOMI BARASH-SCHMIDT IN 4/68, AND VERIFIED BY LEROY PRICE.

BEAM IS K^+ ON PROTON AT 1.96 GEV/C.

THIS EXPERIMENT USES THE B.N.L. 20 IN. (H) BUBBLE CHAMBER. A TOTAL OF 20000 PICTURES ARE REPORTED ON.

GENERAL COMMENTS ON THIS ARTICLE

1 CROSS SECTIONS ARE NORMALIZED TO TAU DECAYS.

KEY WORDS - CROSS SECTION DALITZ PLOT $K^+(890)$ DELTA(1238) ANGULAR DISTRIBUTION

CROSS SECTION FOR K^+ PROTON $\rightarrow K^+(890)^+ \text{ PROTON}$
 $K^+(890)^+ \rightarrow K^0 \pi^+ [1]$

ORIGINAL DATA		RENORMALIZED VALUE [2]
LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	MILLI-BARNS
1.96 $\pm$.02	1.3 $\pm$.2	1.2 $\pm$.2

[1] COUNTED ONLY EVENTS ABOVE BACKGROUND (MASS = .890 GEV; WIDTH = .100 GEV).
[2] R.W.BLAND, UCRL 18131 (1968).

CROSS SECTION FOR K^+ PROTON $\rightarrow K^0 \Delta^{++}$
 $\Delta^{++} \rightarrow \text{PROTON } \pi^+ [1]$

ORIGINAL DATA		RENORMALIZED VALUE [2]
LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	MILLI-BARNS
1.96 $\pm$.02	2.3 $\pm$.3	2.1 $\pm$.3

[1] COUNTED ONLY EVENTS ABOVE BACKGROUND (MASS = 1.220 GEV; WIDTH = .200 GEV).
[2] R.W.BLAND, UCRL 18131 (1968).

CROSS SECTION FOR K^+ PROTON $\rightarrow K^0 \text{ PROTON } \pi^+ [1]$
CROSS SECTION IS ONLY FOR THE NON-RESONANT PRODUCTION OF THESE PARTICLES.

ORIGINAL DATA		RENORMALIZED VALUE [1]
LABORATORY BEAM MOMENTUM GEV/C	MILLI-BARNS	MILLI-BARNS
1.96 $\pm$.02	1.0 $\pm$.2	.9 $\pm$.2

[1] R.W.BLAND, UCRL 18131 (1968).

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^0 \Delta^{++}$, [FIGURE 4]
 $\Delta^{++} \rightarrow \text{PROTON } \pi^+$ [1]

LABORATORY BEAM MOMENTUM = $1.96 \pm .02$ GEV/C.

 . THIS DATA WAS READ FROM A GRAPH .

ORIGINAL DATA		RENORMALIZED VALUES [2]	
COS(THETA)	D-SIGMA/D-OMEGA MB/SR		MILLI-BARNS
.95 $\pm$.05	.460 $\pm$.070	.	.425 $\pm$.055
.85 .05	.490 .070	.	.453 .055
.75 .05	.400 .070	.	.370 .055
.65 .05	.620 .070	.	.573 .055
.55 .05	.370 .070	.	.342 .055
.40 .10	.150 .020	.	.139 .018
.15 .15	.090 .015	.	.083 .014
-.25 .25	.030 .009	.	.028 .008
-.75 .25	.030 .009	.	.028 .008

THETA IS THE ANGLE THAT THE K^0 MAKES WITH THE BEAM IN THE GRAND C.M.

[1] USED SIMPLE MASS CUT [MASS = 1.120 GEV; WIDTH = .200 GEV].

[2] ORIGINAL VALUES HAVE BEEN MULTIPLIED BY .925 - THE SAME FACTOR THAT THE INTEGRATED CROSS SECTION WAS MULTIPLIED BY TO GET THE RE-NORMALIZED VALUES OF R.W.BLAND, UCRL 18131 (1968).

DIFFERENTIAL CROSS SECTION FOR $K^+ \text{ PROTON} \rightarrow K^*(890)^+ \text{ PROTON}$, [FIGURE 6]
 $K^*(890)^+ \rightarrow K^0 \pi^+$ [1]

LABORATORY BEAM MOMENTUM = $1.96 \pm .02$ GEV/C.

 . THIS DATA WAS READ FROM A GRAPH .

ORIGINAL DATA		RENORMALIZED VALUES [2]	
COS(THETA)	D-SIGMA/D-OMEGA MB/SR		MILLI-BARNS
.95 $\pm$.05	.620 $\pm$.100	.	.567 $\pm$.091
.85 .05	.500 .080	.	.457 .073
.75 .05	.210 .050	.	.192 .046
.65 .05	.190 .050	.	.174 .046
.50 .10	.130 .030	.	.119 .027
.20 .20	.062 .030	.	.057 .027
-.25 .25	.023 .020	.	.021 .018
-.75 .25	.034 .010	.	.031 .009

THETA IS THE ANGLE THAT THE $K^*(890)^+$ MAKES WITH THE BEAM IN THE GRAND C.M.

[1] USED SIMPLE MASS CUT [MASS = .890 GEV; WIDTH = .100 GEV].

[2] ORIGINAL VALUES HAVE BEEN MULTIPLIED BY .915 - THE SAME FACTOR THAT THE INTEGRATED CROSS SECTION WAS MULTIPLIED BY TO GET THE RE-NORMALIZED VALUES OF R.W.BLAND, UCRL 18131 (1968).

90 MEASUREMENT OF POLARIZATION IN $K^+ \text{ P}$ ELASTIC SCATTERING AT 1.37, 1.45, 1.71, AND 1.89 GEV/C AND PHASE SHIFT ANALYSIS. [PHYS. REV. LETTERS 23, 194 (1969)]

J.C. ASBURY, J.D. DOWELL, S. KATO, D. LUNDQUIST, T.B. NOVY, A. YOKOSAWA [ARGONNE NAT. LAB., ARGONNE, ILL., USA]
 B. BARNETT, P.F.M. KOEHLER, P. STEINBERG [UNIV. OF MARYLAND, COLLEGE PARK, MD., USA]

ABSTRACT MEASUREMENTS ARE REPORTED OF POLARIZATION IN $K^+ \text{ P}$ ELASTIC SCATTERING AT 1.37, 1.45, 1.71 AND 1.89 GEV/C. A PHASE SHIFT ANALYSIS INCLUDING THESE NEW DATA HAS BEEN PERFORMED. RESONANCE LINE BEHAVIOR IS OBSERVED IN THE $\pi^0 \pi^0$ PARTIAL WAVE AMPLITUDE.

ADDITIONAL FOOTNOTE REFERENCES
 PHYS. REV. 165, 1770 (1968), PHYS. REV. LETTERS 21, 1286 (1968), PHYS. REV. LETTERS 17, 102 (1966), PHYS. REV. 168, 1466 (1968), PHYS. REV. LETTERS 18, 1077 (1967), UCRL 18323 (1968), PHYS. LETTERS 288, 611 (1969), PHYS. REV. 174, 1859 (1968), BNL BNL 50031 (1965), CROVGENICS 6 207 (1966), ANL ANLHEP6938, PHYS. REV. LETTERS 19, 259 (1967), UCRL 17388 (1967), PHYS. LETTERS 16, 83 (1965), PHYS. REV. 129, 2743 (1953), PHYS. REV. 139B, 1411 (1965), PHYS. REV. LETTERS 21, 1282 (1968), UNIV. OF ROCHESTER UR-875-254, PHYS. REV. 177, 2318 (1969), ANL ANLHEP6819, AND NUC. PHYS. B9, 549 (1969).

ARTICLE READ BY ODETTE BENARY IN 7/69, AND VERIFIED BY ROGER BLAND.

BEAM IS K^+ ON HYDROGEN COMPOUND FROM 1.37 TO 1.89 GEV/C. TARGET IS POLARIZED 60 PER CENT (NORMAL TO THE BEAM DIRECTION).

THIS EXPERIMENT USES COUNTERS.

KEY WORDS = POLARIZATION PHASE SHIFT

ELASTIC POLARIZATION FOR K⁺ PROTON. (FIGURE 1A)LABORATORY BEAM MOMENTUM = 1.37 GEV/C $\pm$ 2.5(PER CENT).

(WRITTEN COMMUNICATION FROM T. B. NOVEY, JULY 1969)

COS(THETA)	POLARIZATION
.870	.548 $\pm$.090
.830	.698 .168
.790	.673 .060
.745	.592 .095
.695	.672 .070
.650	.590 .100
.580	.717 .075
.490	.765 .075
.430	.751 .000
.350	.727 .059
.230	.663 .090
.115	.763 .090
.020	.776 .089
-.080	.768 .157
-.125	.627 .102
-.225	.669 .140
-.305	.792 .151
-.360	-.007 .304
-.415	.796 .250
-.485	.724 .173
-.600	.805 .280
-.690	-.040 .300
-.760	.210 .274
-.820	.464 .346
-.895	-.279 .268

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.
 THE POLARIZATION IS OF THE PROTON ALONG THE NORMAL TO THE PRODUCTION PLANE IN THE GRAND C.M.

ELASTIC POLARIZATION FOR K⁺ PROTON. (FIGURE 1B)LABORATORY BEAM MOMENTUM = 1.45 GEV/C $\pm$ 2.5(PER CENT).

(WRITTEN COMMUNICATION FROM T. B. NOVEY, JULY 1969)

COS(THETA)	POLARIZATION
.8800	.1720 $\pm$.1640
.8600	.5490 .0990
.8150	.5520 .1020
.7775	.6290 .0500
.7350	.5360 .0710
.6735	.6280 .0620
.5750	.7520 .0950
.4750	.6600 .1000
.4200	.7100 .0800
.3265	.6940 .0710
.1975	.6600 .0740
.1050	.3740 .0850
.0180	.5980 .1210
-.0800	.5650 .1340
-.1700	.5700 .1700
-.2475	.6495 .1450
-.3300	.7505 .1610
-.5100	.5420 .2010
-.5900	.4290 .2030
-.7100	.3920 .1660
-.8150	.3190 .3650
-.8950	.1920 .2560

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.
 THE POLARIZATION IS OF THE PROTON ALONG THE NORMAL TO THE PRODUCTION PLANE IN THE GRAND C.M.

ELASTIC POLARIZATION FOR K⁺ PROTON. (FIGURE 1C)LABORATORY BEAM MOMENTUM = 1.71 GEV/C $\pm$ 2.5(PER CENT).

(WRITTEN COMMUNICATION FROM T. B. NOVEY, JULY 1969)

COS(THETA)	POLARIZATION
.910	.555 $\pm$.112
.880	.570 .070
.840	.576 .030
.800	.684 .057
.760	.660 .034
.720	.645 .058
.660	.652 .036
.615	.686 .073
.550	.674 .045
.510	.686 .081
.390	.625 .059
.320	.564 .120
.180	.648 .111
.150	.541 .118
.050	.596 .080
.070	.658 .080
-.040	.626 .080
-.140	.499 .075
-.240	.570 .115
-.380	.764 .206
-.560	.589 .270
-.690	.362 .190
-.830	-.131 .179
-.900	-.164 .800

THETA IS THE ANGLE THAT THE K⁺ MAKES WITH THE BEAM IN THE GRAND C.M.
 THE POLARIZATION IS OF THE PROTON ALONG THE NORMAL TO THE PRODUCTION PLANE IN THE GRAND C.M.

ELASTIC POLARIZATION FOR K^+ PROTON.

[FIGURE 10]

LABORATORY BEAM MOMENTUM = 1.89 GEV/C $\pm$ 2.5(PER CENT).

(WRITTEN COMMUNICATION FROM T. B. NOVEY, JULY 1969)

COS(THETA)	POLARIZATION	
.900	.410	-.120
.875	.505	.056
.830	.570	.031
.790	.561	.044
.742	.563	.033
.695	.520	.066
.637	.593	.033
.600	.585	.045
.520	.552	.044
.400	.512	.086
.340	.387	.087
.220	.396	.076
.125	.472	.099
.015	.394	.092
-.085	.326	.135
-.180	.425	.186
-.305	.910	.500
-.400	.350	.400
-.560	.450	.520
-.675	.560	.500
-.730	-.360	.385
-.779	-.040	.400
-.915	-.080	.650

THETA IS THE ANGLE THAT THE K^+ MAKES WITH THE BEAM IN THE GRAND C.M.
 THE POLARIZATION IS OF THE PROTON ALONG THE NORMAL TO THE PRODUCTION PLANE IN THE GRAND C.M.

Section IV.

INDICES

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

With all of the information for each article stored in a computer-searchable fashion, one could generate numerous types of indices; two types that we have found most useful are included in this section.

1. MOMENTUM INDICES—Here we list all of our K^+N articles classified by input channel (K^+p or K^+d) and then ordered according to increasing beam momentum. If a particular paper reports results at more than one energy, that paper is listed once for each momentum value reported. The reference number in the last column is the article number in Section III.

2. KEY WORDS CLASSIFICATION—As stated in Section III, each article is assigned certain KEY WORDS. These words (or phrases) are intended to indicate the contents of the article. As our list of KEY WORDS has grown, we generally have not yet gone back to older articles and inserted the appropriate new words. Thus references may be missing from some of the categories. We hope to have this remedied by our next edition of K^+N .

If you have any suggestions for other useful indices, please let us know. We believe that a good set of indices will make this report much more valuable.

Momentum Index ($K^+ p$)

BEAM MOMENTUM	1ST AUTHOR	JOURNAL	VOLUME, PAGE	INSTITUTIONS	DETECTOR	YEAR PUBLISHED	REF. NR.
.140	GOLDHABER	PRL	9 135	LRL BERK	HBC 15	62	8
.175	GOLDHABER	PRL	9 135	LRL BERK	HBC 15	62	8
.205	GOLDHABER	PRL	9 135	LRL BERK	HBC 15	62	8
.235	GOLDHABER	PRL	9 135	LRL BERK	HBC 15	62	8
.265	GOLDHABER	PRL	9 135	LRL BERK	HBC 15	62	8
.355	GOLDHABER	PRL	9 135	LRL BERK	HBC 15	62	8
.451	KYCIA	PR	118 553	LRL BERK	CNTR	60	27
.520	GOLDHABER	PRL	9 135	LRL BERK	HBC 15	62	8
.522	KYCIA	PR	118 553	LRL BERK	CNTR	60	27
.589	KYCIA	PR	118 553	LRL BERK	CNTR	60	27
.592	BUGG	PR	168 1466	RHEL BIRM	CNTR	68	30
.620	BUGG	PR	168 1466	RHEL BIRM	CNTR	68	30
.642	GOLDHABER	PRL	9 135	LRL BERK	HBC 15	62	8
.642	FISK	CERN CONF	358	UCLA LRL	HBC 15	62	6
.644	BUGG	PR	168 1466	RHEL BIRM	CNTR	68	30
.713	BUGG	PR	168 1466	RHEL BIRM	CNTR	68	30
.735	BLAND	UCRL	18758	LRL	HBC 25	68	72
.735	FILIPPAS	NC	51A 1053	CERN BGNA MODE	HBC 81	67	60
.755	FILIPPAS	NC	51A 1053	CERN BGNA MODE	HBC 81	67	60
.768	BUGG	PR	168 1466	RHEL BIRM	CNTR	68	30
.770	COOK	PRL	7 182	LRL BERK	CNTR	61	32
.770	FIORINI	NC	35 319	MILA EPOL	HLBC	65	10
.778	FOCARDI	PL	248 314	BGNA MODE CERN	HBC 81	67	76
.780	FEMIND	NC	50A 372	MSNA BGNA	HBC 81	67	71
.785	BLAND	UCRL	18758	LRL	HBC 25	68	72
.785	FILIPPAS	NC	51A 1053	CERN BGNA MODE	HBC 81	67	60
.810	FISK	CERN CONF	358	UCLA LRL	HBC 15	62	6
.810	STUBBS	PRL	7 188	LRL UCLA	HBC 15	61	7
.823	BUGG	PR	168 1466	RHEL BIRM	CNTR	68	30
.860	BLAND	UCRL	18131	LRL	HBC 25	68	68
.860	BLAND	PRL	18 1077	LRL BERK	HBC 25	67	19
.864	BLAND	UCRL	18758	LRL	HBC 25	68	72
.864	BLAND	UCRL	18889	LRL BERK	HBC 25	69	53
.864	BUGG	PR	168 1466	RHEL BIRM	CNTR	68	30
.865	ANDERSSON	CERN		CERN	CNTR	69	54
.891	COOL	PRL	17 102	BNL	CNTR	66	49
.900	COOL	PRL	17 102	BNL	CNTR	66	49
.904	BUGG	PR	168 1466	RHEL BIRM	CNTR	68	30
.910	HIRSHCH	PR	175 B171	LRL BERK	HLBC 32	64	73
.910	KEHDE	PRL	11 93	UNC	HLBC 32	63	21
.938	BUGG	PR	168 1466	RHEL BIRM	CNTR	68	30
.942	COOL	PRL	17 102	BNL	CNTR	66	49
.950	COOL	PRL	17 102	BNL	CNTR	66	49
.950	FIORINI	NC	35 319	MILA EPOL	HLBC	65	10
.960	BLAND	UCRL	18131	LRL	HBC 25	68	68
.962	BURROWES	PRL	2 117	MIT	CNTR	59	24
.969	BLAND	UCRL	18758	LRL	HBC 25	68	72
.969	BLAND	UCRL	18889	LRL BERK	HBC 25	69	53
.969	BUGG	PR	168 1466	RHEL BIRM	CNTR	68	30
.970	COOK	PR	129 2743	LRL BERK	SPRK	63	64
.970	COOK	PRL	7 182	LRL BERK	CNTR	61	32
.971	ANDERSSON	CERN		CERN	CNTR	69	54
.990	CARROLL	PRL	21 1282	BNL ROCH	SPRK	68	61
.992	COOL	PRL	17 102	BNL	CNTR	66	49
.1.000	COOL	PRL	17 102	BNL	CNTR	66	49
.1.029	BUGG	PR	168 1466	RHEL BIRM	CNTR	68	30
.1.040	CARROLL	PRL	21 1282	BNL ROCH	SPRK	68	61
.1.043	COOL	PRL	17 102	BNL	CNTR	66	49
.1.050	COOL	PRL	17 102	BNL	CNTR	66	49
.1.063	BURROWES	PRL	2 117	MIT	CNTR	59	24
.1.084	BUGG	PR	168 1466	RHEL BIRM	CNTR	68	30
.1.087	ANDERSSON	CERN		CERN	CNTR	69	54
.1.094	COOL	PRL	17 102	BNL	CNTR	66	49
.1.100	LUUL	PRL	17 102	BNL	CNTR	66	49
.1.110	CARROLL	PRL	21 1282	BNL ROCH	SPRK	68	61
.1.140	BOLDT	PR	133 8220	COEF SACL	HBC 35	64	62
.1.140	BUGG	PR	168 1466	RHEL BIRM	CNTR	68	30
.1.144	COOL	PRL	17 102	BNL	CNTR	66	49
.1.150	COOL	PRL	17 102	BNL	CNTR	66	49
.1.170	COOK	PR	129 2743	LRL BERK	SPRK	63	64
.1.170	CARROLL	PRL	21 1282	BNL ROCH	SPRK	68	61
.1.170	COOK	PRL	7 182	LRL BERK	CNTR	61	32
.1.189	BUGG	PR	168 1466	RHEL BIRM	CNTR	68	30
.1.194	COOL	PRL	17 102	BNL	CNTR	66	49
.1.200	BLAND	UCRL	18131	LRL	HBC 25	68	68
.1.200	CARROLL	PRL	21 1282	BNL ROCH	SPRK	68	61
.1.200	COOL	PRL	17 102	BNL	CNTR	66	49
.1.200	BLAND	PRL	17 939	LRL BERK	HBC 25	66	35
.1.207	BLAND	UCRL	18758	LRL	HBC 25	68	72
.1.207	BLAND	UCRL	18889	LRL BERK	HBC 25	69	53
.1.215	ANDERSSON	CERN		CERN	CNTR	69	54
.1.220	ANDERSSON	PL	288 611	CERN	CNTR	69	77
.1.238	BUGG	PR	168 1466	RHEL BIRM	CNTR	68	30
.1.245	COOL	PRL	17 102	BNL	CNTR	66	49
.1.250	COOL	PRL	17 102	BNL	CNTR	66	49
.1.245	BUGG	PR	168 1466	RHEL BIRM	CNTR	68	30
.1.295	COOL	PRL	17 102	BNL	CNTR	66	49
.1.300	CARROLL	PRL	21 1282	BNL ROCH	SPRK	68	61
.1.300	COOL	PRL	17 102	BNL	CNTR	66	49
.1.300	COOK	PRL	7 182	LRL BERK	CNTR	61	32
.1.345	COOL	PRL	17 102	BNL	CNTR	66	49
.1.347	BUGG	PR	168 1466	RHEL BIRM	CNTR	68	30
.1.350	COOL	PRL	17 102	BNL	CNTR	66	49
.1.360	BLAND	UCRL	18131	LRL	HBC 25	68	68
.1.367	BLAND	UCRL	18758	LRL	HBC 25	68	72
.1.370	ASBURY	PRL	23 194	ANL UND	CNTR	69	90
.1.370	BURROWES	PR	2 117	MIT	CNTR	59	24
.1.372	ANDERSSON	CERN		CERN	CNTR	69	54
.1.395	LUUL	PRL	17 102	BNL	CNTR	66	49
.1.400	CARROLL	PRL	21 1282	BNL ROCH	SPRK	68	61
.1.400	COOL	PRL	17 102	BNL	CNTR	66	49
.1.408	BUGG	PR	168 1466	RHEL BIRM	CNTR	68	30
.1.440	COOK	PRL	7 182	LRL BERK	CNTR	61	32
.1.445	COOL	PRL	17 102	BNL	CNTR	66	49
.1.450	ASBURY	PRL	23 194	ANL UND	CNTR	69	90
.1.450	BLAND	UCRL	18758	LRL	HBC 25	68	72
.1.450	COOL	PRL	17 102	BNL	CNTR	66	49
.1.453	ANDERSSON	CERN		CERN	CNTR	69	54
.1.455	BETTINI	PL	16 83	PADO OXF	HBC 23	65	63
.1.455	CHADWICK	PR	6 309	OXF	HBC 23	63	18
.1.468	BUGG	PR	168 1466	RHEL BIRM	CNTR	68	30
.1.490	CARROLL	PRL	21 1282	BNL ROCH	SPRK	68	61
.1.495	COOL	PRL	17 102	BNL	CNTR	66	49
.1.500	COOL	PRL	17 102	BNL	CNTR	66	49
.1.512	BURROWES	PRL	2 117	MIT	CNTR	59	24
.1.550	ABRAMS	PRL	19 259	BNL	CNTR	67	40
.1.563	BUGG	PR	168 1466	RHEL BIRM	CNTR	68	30
.1.580	BLAND	PRL	18 1077	LRL BERK	HBC 25	67	19
.1.585	BLAND	UCRL	18758	LRL	HBC 25	68	72

1.596	COOL	PRL	17	102	BNL	CNTR	66	49
1.600	COOL	PRL	17	102	BNL	CNTR	66	49
1.605	ABRAMS	PRL	19	259	BNL	CNTR	67	40
1.610	CARROLL	PRL	21	1282	BNL ROCH	SPRK	68	61
1.646	COOL	PRL	17	102	BNL	CNTR	66	49
1.650	COOL	PRL	17	102	BNL	CNTR	66	49
1.654	BURROWES	PRL	2	117	MIT	CNTR	59	24
1.670	CARROLL	PRL	21	1282	BNL ROCH	SPRK	68	61
1.680	COOK	PRL	7	182	LRL BERK	CNTR	61	32
1.696	COOL	PRL	17	102	BNL	CNTR	66	49
1.700	COOL	PRL	17	102	BNL	CNTR	66	49
1.700	ABRAMS	PRL	19	259	BNL	CNTR	67	40
1.710	ASBURY	PRL	23	194	ANL UMD	CNTR	69	90
1.710	CARROLL	PRL	21	1282	BNL ROCH	SPRK	68	61
1.746	COOL	PRL	17	102	BNL	CNTR	66	49
1.750	COOL	PRL	17	102	BNL	CNTR	66	49
1.750	ABRAMS	PRL	19	259	BNL	CNTR	67	40
1.790	CARROLL	PRL	21	1282	BNL ROCH	SPRK	68	61
1.796	COOL	PRL	17	102	BNL	CNTR	66	49
1.800	COOL	PRL	17	102	BNL	CNTR	66	49
1.800	ABRAMS	PRL	19	259	BNL	CNTR	67	40
1.820	CARROLL	PRL	21	1282	BNL ROCH	SPRK	68	61
1.850	ABRAMS	PRL	19	259	BNL	CNTR	67	40
1.877	BURROWES	PRL	2	117	MIT	CNTR	59	24
1.880	CARROLL	PRL	21	1282	BNL ROCH	SPRK	68	61
1.890	ASBURY	PRL	23	194	ANL UMD	CNTR	69	90
1.896	COOL	PRL	17	102	BNL	CNTR	66	49
1.900	COOL	PRL	17	102	BNL	CNTR	66	49
1.900	ABRAMS	PRL	19	259	BNL	CNTR	67	40
1.945	BUGG	PR	168	1466	RHEL BIRM	CNTR	68	30
1.950	ABRAMS	PRL	19	259	BNL	CNTR	67	40
1.960	GOLDMAN	PR	142	913	LRL BERK	HBC 20	66	89
1.960	BLAND	UCRL	18758	1411	LRL	HBC 25	68	72
1.960	CHINOWSKY	PR	13981	1411	LRL BERK	HBC 20	65	57
1.960	GOLDMAN	PL	18	76	LRL BERK	HBC 20	65	12
1.970	COOK	PR	129	2743	LRL BERK	SPRK	63	64
1.970	CUUK	PRL	7	182	LRL BERK	CNTR	61	32
1.996	COOL	PRL	17	102	BNL	CNTR	66	49
2.000	CARROLL	PRL	21	1282	BNL ROCH	SPRK	68	61
2.000	COOL	PRL	17	102	BNL	CNTR	66	49
2.000	ABRAMS	PRL	19	259	BNL	CNTR	67	40
2.050	ABRAMS	PRL	19	259	BNL	CNTR	67	40
2.096	COOL	PRL	17	102	BNL	CNTR	66	49
2.100	COOL	PRL	17	102	BNL	CNTR	66	49
2.100	ABRAMS	PRL	19	259	BNL	CNTR	67	40
2.147	BURROWES	PRL	2	117	MIT	CNTR	59	24
2.150	ABRAMS	PRL	19	259	BNL	CNTR	67	40
2.196	COOL	PRL	17	102	BNL	CNTR	66	49
2.200	CARROLL	PRL	21	1282	BNL ROCH	SPRK	68	61
2.200	COOL	PRL	17	102	BNL	CNTR	66	49
2.200	ABRAMS	PRL	19	259	BNL	CNTR	67	40
2.260	BOMSE	PR	158	1298	JHOP	HBC 20	67	65
2.260	COOK	PRL	7	182	LRL BERK	CNTR	61	32
2.260	BOMSE	PR	158	1281	JHOP	HBC 20	67	28
2.300	CNN	PRL	17	102	BNL	CNTR	66	49
2.300	ABRAMS	PRL	19	259	BNL	CNTR	67	40
2.330	CARROLL	PRL	21	1282	BNL ROCH	SPRK	68	61
2.350	ABRAMS	PRL	19	259	BNL	CNTR	67	40
2.387	BURROWES	PRL	2	117	MIT	CNTR	59	24
2.396	COOL	PRL	17	102	BNL	CNTR	66	49
2.400	COOL	PRL	17	102	BNL	CNTR	66	49
2.400	ABRAMS	PRL	19	259	BNL	CNTR	67	40
2.450	CARROLL	PRL	21	1282	BNL ROCH	SPRK	68	61
2.450	ABRAMS	PRL	19	259	BNL	CNTR	67	40
2.473	BUGG	PR	168	1466	RHEL BIRM	CNTR	68	30
2.480	ANDERSSON	PL	288	611	CERN	CNTR	69	77
2.500	ABRAMS	PRL	19	259	BNL	CNTR	67	40
2.550	ABRAMS	PRL	10	257	BNL	CNTR	67	40
2.550	COOK	PRL	7	182	LRL BERK	CNTR	61	32
2.600	ABRAMS	PRL	19	259	BNL	CNTR	67	40
2.650	NEWMAN	PR	158	1310	LRL BERK SLAC	HBC 20	67	75
2.650	BLAND	UCRL	18758	1411	LRL	HBC 25	68	72
2.650	ABRAMS	PRL	19	259	BNL	CNTR	67	40
2.700	ABRAMS	PRL	19	259	BNL	CNTR	67	40
2.750	ABRAMS	PRL	19	259	BNL	CNTR	67	40
2.760	ABRAMS	PRL	21	1407	ILL	HBC 33	68	66
2.800	ABRAMS	PRL	19	259	BNL	CNTR	67	40
2.830	COOK	PRL	7	182	LRL BERK	CNTR	61	32
2.850	ABRAMS	PRL	19	259	BNL	CNTR	67	40
2.900	ABRAMS	PRL	19	259	BNL	CNTR	67	40
2.950	ABRAMS	PRL	19	259	BNL	CNTR	67	40
2.965	FERRO-LUZZ	NC	36	1101	CERN	HBC 81	65	70
2.965	LYNCH	PL	9	359	CERN	HBC 81	64	20
2.970	BLAND	UCRL	18758	1411	LRL	HBC 25	68	72
2.970	FERRO-LUZZ	NC	39	417	CERN	HBC 81	65	56
2.970	DEBAISIEUX	NC	43A	142	LHC	HBC 81	68	20
2.970	FERRO-LUZZ	PL	17	133	ILL	HBC 81	65	15
2.972	SALLSTROM	AF	37	468	STOH	HBC 81	68	80
2.972	SALLSTROM	NC	49A	348	STOH	HBC 81	67	69
3.000	BASSOMPIER	PL	278	468	CERN	HBCS	68	74
3.000	ABRAMS	PRL	19	259	BNL	CNTR	67	40
3.000	JOLDERSMA	PRL	17	716	BNL	HBC 80	66	9
3.000	VON DARDDEL	PRL	5	333	CERN	CNTR	60	4
3.000	FERRO-LUZZ	PL	12	255	CERN	HBC 81	64	2
3.050	ABRAMS	PRL	19	259	BNL	CNTR	67	40
3.100	ABRAMS	PRL	19	259	BNL	CNTR	67	40
3.150	ABRAMS	PRL	19	259	BNL	CNTR	67	40
3.200	ABRAMS	PRL	19	259	BNL	CNTR	67	40
3.250	ABRAMS	PRL	19	259	BNL	CNTR	67	40
3.250	BAKER	PR	129	2285	BNL	CNTR	63	33
3.300	ABRAMS	PRL	19	259	BNL	CNTR	67	40
3.460	BLAND	UCRL	18758	1411	LRL	HBC 25	68	72
3.460	DEBAERE	NC	45A	885	BELG	HBC 81	66	51
3.460	GEORGE	NC	49A	9	CERN	HBC 81	67	50
3.460	FERRO-LUZZ	PL	17	155	CERN	HBC 81	65	15
3.500	BASSOMPIER	PL	278	468	CERN	HBCS	68	74
3.500	VON DARDDEL	PRL	5	333	CERN	CNTR	60	4
3.530	CLINE	PRL	19	675	WISC	HBC 80	67	41
3.533	GDSHAW	PL	22	347	WISC	HBC 80	66	59
3.533	THESIS	PL	16	1063	WISC	HBC 80	67	48
3.533	ERWIN	PRL	89	249	CERN	SACL	69	67
3.550	BAKER	NP	80	640	SACL CERN	SPRK	69	52
3.550	BANAIGS	PL	248	317	SACL CERN	SPRK	67	31
4.000	BAKER	PR	129	2285	BNL	CNTR	63	33
4.600	FU	UCRL	18417	2092	LRL BERK	HBC 80	68	81
4.600	ALEXANDER	PRL	17	726	LRL BERK	HBC 80	66	5
4.600	SHEN	PRL	45A	885	BELG	HBC 81	66	51
4.970	GEORGE	NC	49A	9	CERN	HBC 81	67	50
4.970	BASSOMPIER	NC	48	589	CERN	HBC 200	67	23
4.970	BASSOMPIER	PL	268	30	CERN	HBC 81	65	22
4.970	FERRO-LUZZ	PL	17	155	CERN	HBCS	68	15
5.000	BASSOMPIER	NP	89	295	CERN	HBCS	69	84

BEAM MOMENTUM	1ST AUTHOR	JOURNAL	VOLUME,PAGE	INSTITUTIONS	DETECTOR	YEAR PUBLISHED	REF.NO.	
.233	SLATER	PRL	7 378	UCLA LRL BERK	HBC	15	61	85
.330	SLATER	PRL	7 378	UCLA LRL BERK	HBC	15	61	85
.376	SLATER	PRL	7 378	UCLA LRL BERK	HBC	15	61	85
.377	STENGER	PR	13481111	UCLA LRL BERK	HBC	15	64	44
.529	SLATER	PRL	13481111	UCLA LRL BERK	HBC	15	64	44
.530	STENGER	PR	13481111	UCLA LRL BERK	HBC	15	64	44
.592	BUGG	PR	168 1466	RHEL BIRM	CNTR	30	68	30
.600	RAY	BNL	13616	CARN BNL	HBC	30	69	87
.620	BUGG	PR	168 1466	RHEL BIRM	CNTR	30	68	30
.641	SLATER	PRL	7 378	UCLA LRL BERK	HBC	15	61	85
.642	FISK	CERN CONF	7 378	UCLA LRL BERK	HBC	15	62	6
.644	BUGG	PR	168 1466	RHEL BIRM	CNTR	30	68	30
.713	BUGG	PR	168 1466	RHEL BIRM	CNTR	30	68	30
.768	BUGG	PR	168 1466	RHEL BIRM	CNTR	30	68	30
.770	COOK	PRL	7 182	LRL BERK	CNTR	61	32	
.810	FISK	CERN CONF	7 378	UCLA LRL BERK	HBC	15	62	6
.811	SLATER	PRL	7 378	UCLA LRL BERK	HBC	15	61	85
.823	BUGG	PR	168 1466	RHEL BIRM	CNTR	30	68	30
.863	HIRATA	PR	21 1485	LRL BERK	HBC	25	68	34
.864	BUGG	PR	168 1466	RHEL BIRM	CNTR	30	68	30
.891	COOL	PRL	17 102	BNL	CNTR	66	49	
.900	COOL	PRL	17 102	BNL	CNTR	66	49	
.904	BUGG	PR	168 1466	RHEL BIRM	CNTR	30	68	30
.938	BUGG	PR	168 1466	RHEL BIRM	CNTR	30	68	30
.942	COOL	PRL	17 102	BNL	CNTR	66	49	
.950	COOL	PRL	17 102	BNL	CNTR	66	49	
.962	BURROWS	PRL	2 117	MIT	CNTR	59	24	
.968	HIRATA	PRL	21 1485	LRL BERK	HBC	25	68	34
.969	BUGG	PR	168 1466	RHEL BIRM	CNTR	30	68	30
.970	COOK	PRL	7 182	LRL BERK	CNTR	61	32	
.992	COOL	PRL	17 102	BNL	CNTR	66	49	
1.000	COOL	PRL	17 102	BNL	CNTR	66	49	
1.029	BUGG	PR	168 1466	RHEL BIRM	CNTR	30	68	30
1.043	COOL	PRL	17 102	BNL	CNTR	66	49	
1.050	COOL	PRL	17 102	BNL	CNTR	66	49	
1.063	BURROWS	PRL	2 117	MIT	CNTR	59	24	
1.084	BUGG	PR	168 1466	RHEL BIRM	CNTR	30	68	30
1.094	COOL	PRL	17 102	BNL	CNTR	66	49	
1.100	COOL	PRL	17 102	BNL	CNTR	66	49	
1.140	BUGG	PR	168 1466	RHEL BIRM	CNTR	30	68	30
1.144	COOL	PRL	17 102	BNL	CNTR	66	49	
1.150	COOL	PRL	17 102	BNL	CNTR	66	49	
1.170	COOK	PRL	7 182	LRL BERK	CNTR	61	32	
1.109	BUGG	PR	168 1466	RHEL BIRM	CNTR	30	68	30
1.194	COOL	PRL	17 102	BNL	CNTR	66	49	
1.200	COOL	PRL	17 102	BNL	CNTR	66	49	
1.211	HIRATA	PRL	21 1485	LRL BERK	HBC	25	68	34
1.238	BUGG	PR	168 1466	RHEL BIRM	CNTR	30	68	30
1.245	COOL	PRL	17 102	BNL	CNTR	66	49	
1.250	COOL	PRL	17 102	BNL	CNTR	66	49	
1.293	BUGG	PR	168 1466	RHEL BIRM	CNTR	30	68	30
1.295	COOL	PRL	17 102	BNL	CNTR	66	49	
1.300	COOL	PRL	17 102	BNL	CNTR	66	49	
1.300	COOK	PRL	7 182	LRL BERK	CNTR	61	32	
1.345	COOL	PRL	17 102	BNL	CNTR	66	49	

1.364	HIRATA	PRL	21 1485	LRL	BERK	HBC	25	68	34
1.370	BURROWES	PRL	2 117	MIT		CNTR		59	24
1.395	COOL	PRL	17 102	BNL		CNTR		66	49
1.400	COOL	PRL	17 102	BNL		CNTR		66	49
1.408	BUGG	PR	168 1466	RHEL	BIRM	CNTR		68	30
1.440	COOK	PRL	7 182	LRL	BERK	CNTR		61	32
1.445	COOL	PRL	17 102	BNL		CNTR		66	49
1.450	COOL	PRL	17 102	BNL		CNTR		66	49
1.448	BUGG	PR	168 1466	RHEL	BIRM	CNTR		66	49
1.495	COOL	PRL	17 102	BNL		CNTR		68	30
1.500	COOL	PRL	17 102	BNL		CNTR		66	49
1.512	BURROWES	PRL	2 117	MIT		CNTR		66	49
1.550	ABRAMS	PRL	19 259	BNL		CNTR		59	24
1.563	BUGG	PR	168 1466	RHEL	BIRM	CNTR		67	40
1.596	COOL	PRL	17 102	BNL		CNTR		68	30
1.600	COOL	PRL	17 102	BNL		CNTR		66	49
1.600	ABRAMS	PRL	19 259	BNL		CNTR		66	49
1.646	COOL	PRL	17 102	BNL		CNTR		67	40
1.650	COOL	PRL	17 102	BNL		CNTR		66	49
1.654	BURROWES	PRL	2 117	MIT		CNTR		66	49
1.690	COOK	PRL	7 182	LRL	BERK	CNTR		59	24
1.696	COOL	PRL	17 102	BNL		CNTR		61	32
1.700	COOL	PRL	17 102	BNL		CNTR		66	49
1.700	ABRAMS	PRL	19 259	BNL		CNTR		66	49
1.746	COOL	PRL	17 102	BNL		CNTR		67	40
1.750	COOL	PRL	17 102	BNL		CNTR		66	49
1.750	ABRAMS	PRL	19 259	BNL		CNTR		66	49
1.796	COOL	PRL	17 102	BNL		CNTR		67	40
1.800	COOL	PRL	17 102	BNL		CNTR		66	49
1.800	ABRAMS	PRL	19 259	BNL		CNTR		67	40
1.850	ABRAMS	PRL	19 259	BNL		CNTR		67	40
1.877	BURROWES	PRL	2 117	MIT		CNTR		59	24
1.896	COOL	PRL	17 102	BNL		CNTR		66	49
1.900	COOL	PRL	17 102	BNL		CNTR		66	49
1.900	ABRAMS	PRL	19 259	BNL		CNTR		67	40
1.945	BUGG	PR	168 1466	RHEL	BIRM	CNTR		68	30
1.950	ABRAMS	PRL	19 259	BNL		CNTR		67	40
1.970	COOK	PRL	7 182	LRL	BERK	CNTR		61	32
1.996	COOL	PRL	17 102	BNL		CNTR		66	49
2.000	COOL	PRL	17 102	BNL		CNTR		66	49
2.000	ABRAMS	PRL	19 259	BNL		CNTR		67	40
2.050	ABRAMS	PRL	19 259	BNL		CNTR		67	40
2.096	COOL	PRL	17 102	BNL		CNTR		66	49
2.100	COOL	PRL	17 102	BNL		CNTR		66	49
2.100	ABRAMS	PRL	19 259	BNL		CNTR		67	40
2.147	BURROWES	PRL	2 117	MIT		CNTR		59	24
2.150	ABRAMS	PRL	19 259	BNL		CNTR		67	40
2.196	COOL	PRL	17 102	BNL		CNTR		66	49
2.200	COOL	PRL	17 102	BNL		CNTR		66	49
2.200	ABRAMS	PRL	19 259	BNL		CNTR		67	40
2.260	COOK	PRL	7 182	LRL	BERK	CNTR		61	32
2.300	BUTTERWORTH	PRL	15 734	LRL	BERK	DBC	20	65	43
2.300	COOL	PRL	17 102	BNL		CNTR		66	49
2.300	GOLDHABER	PRL	15 737	LRL	BERK	DBC	20	65	43
2.300	BUTTERWORTH	PRL	15 500	LRL	BERK	HBC	20	65	43
2.300	ABRAMS	PRL	19 259	BNL		CNTR		67	40
2.300	GOLDHABER	PL	18 76	LRL	BERK	HBC	20	65	12
2.350	ABRAMS	PRL	19 259	BNL		CNTR		67	40
2.387	BURROWES	PRL	2 117	MIT		CNTR		59	24
2.396	COOL	PRL	17 102	BNL		CNTR		66	49
2.400	COOL	PRL	17 102	BNL		CNTR		66	49
2.400	ABRAMS	PRL	19 259	BNL		CNTR		67	40
2.450	ABRAMS	PRL	19 259	BNL		CNTR		67	40
2.473	BUGG	PR	168 1466	RHEL	BIRM	CNTR		68	30
2.500	ABRAMS	PRL	19 259	BNL		CNTR		67	40
2.550	ABRAMS	PRL	19 259	BNL		CNTR		67	40
2.550	COOK	PRL	7 182	LRL	BERK	CNTR		61	32
2.600	ABRAMS	PRL	19 259	BNL		CNTR		67	40
2.650	ABRAMS	PRL	19 259	BNL		CNTR		67	40
2.700	ABRAMS	PRL	19 259	BNL		CNTR		67	40
2.750	ABRAMS	PRL	19 259	BNL		CNTR		67	40
2.800	ABRAMS	PRL	19 259	BNL		CNTR		67	40
2.830	COOK	PRL	7 182	LRL	BERK	CNTR		61	32
2.850	ABRAMS	PRL	19 259	BNL		CNTR		67	40
2.900	ABRAMS	PRL	19 259	BNL		CNTR		67	40
2.950	ABRAMS	PRL	19 259	BNL		CNTR		67	40
2.970	BUCHNER	NP	89 286	MPIM		HBC	81	69	58
2.970	GOLDSCHMID	PI	278 602	CERN		DBC	81	68	55
3.000	ABRAMS	PRL	19 259	BNL		CNTR		67	40
3.050	ABRAMS	PRL	19 259	BNL		CNTR		67	40
3.100	ABRAMS	PRL	19 259	BNL		CNTR		67	40
3.150	ABRAMS	PRL	19 259	BNL		CNTR		67	40
3.200	ABRAMS	PRL	19 259	BNL		CNTR		67	40
3.250	ABRAMS	PRL	19 259	BNL		CNTR		67	40
3.300	ABRAMS	PRL	19 259	BNL		CNTR		67	40
6.000	GALBRAITH	PR	138 8913	BNL	CORN	CNTR		65	39
8.000	GALBRAITH	PR	138 8913	BNL	CORN	CNTR		65	39
10.000	GALBRAITH	PR	138 8913	BNL	CORN	CNTR		65	39
12.000	GALBRAITH	PR	138 8913	BNL	CORN	CNTR		65	39
14.000	GALBRAITH	PR	138 8913	BNL	CORN	CNTR		65	39
16.000	GALBRAITH	PR	138 8913	BNL	CORN	CNTR		65	39
18.000	GALBRAITH	PR	138 8913	BNL	CORN	CNTR		65	39
20.000	GALBRAITH	PR	138 8913	BNL	CORN	CNTR		65	39

Key Words Classification

ANGULAR DISTRIBUTION

- [63] GOLDBABER, ET AL., PHYS. REV. LETTERS 19, 972 (1967)
 [64] FISK, ET AL., CERN CONFERENCE 358 (1962)
 [65] STUBBS, ET AL., PHYS. REV. LETTERS 7, 188 (1961)
 [66] GOLDBABER, ET AL., PHYS. REV. LETTERS 9, 135 (1962)
 [67] FIORINI, ET AL., NUOVO CIMENTO 35, 319 (1965)
 [68] ALEXANDER, ET AL., PHYS. REV. LETTERS 20, 755 (1968)
 [69] GOLDBABER, ET AL., PHYS. LETTERS 18, 76 (1965)
 [70] ALMEIDA, ET AL., PHYS. LETTERS 16, 184 (1965)
 [71] CHADWICK, ET AL., PHYS. LETTERS 6, 309 (1963)
 [72] BLAND, ET AL., PHYS. REV. LETTERS 18, 1077 (1967)
 [73] LYNCH, ET AL., PHYS. LETTERS 9, 359 (1964)
 [74] B.KEHO, PHYS. REV. LETTERS 11, 93 (1963)
 [75] BASSOMPIERRE, ET AL., PHYS. LETTERS 268, 30 (1967)
 [76] BOMSE, ET AL., PHYS. REV. LETTERS 20, 1519 (1968)
 [77] BERLINGHIERI, ET AL., CERN HADRON CONFERENCE 172 (1968)
 [78] KYCIA, ET AL., PHYS. REV. 118, 553 (1960)
 [79] BOMSE, ET AL., PHYS. REV. 158, 1298 (1967)
 [80] DEBAISIEUX, ET AL., NUOVO CIMENTO 43A, 142 (1966)
 [81] BANAIGS, ET AL., PHYS. LETTERS 248, 317 (1967)
 [82] ALEXANDER, ET AL., PHYS. REV. 177, 2092 (1969)
 [83] CLINE, ET AL., PHYS. REV. LETTERS 19, 675 (1967)
 [84] BERLINGHIERI, ET AL., PHYS. REV. LETTERS 18, 1087 (1967)
 [85] BUTTERWORTH, ET AL., PHYS. REV. LETTERS 15, 500 (1965)
 [86] STENGER, ET AL., PHYS. REV. 134, 8111 (1964)
 [87] GOLDBABER, ET AL., PHYS. REV. LETTERS 15, 737 (1965)
 [88] J.M.BISHOP, THESIS (1967)
 [89] GEORGE, ET AL., NUOVO CIMENTO 49A, 9 (1967)
 [90] DEBAERE, ET AL., NUOVO CIMENTO 45A, 885 (1966)
 [91] FERRO-LUZZI, ET AL., NUOVO CIMENTO 39, 417 (1965)
 [92] CHINDOWSKY, ET AL., NUC. PHYS. 89, 286 (1969)
 [93] BUCHNER, ET AL., NUC. PHYS. 89, 286 (1969)
 [94] FILIPPAS, ET AL., NUOVO CIMENTO 51A, 1053 (1967)
 [95] BOLDT, ET AL., PHYS. REV. 133, 8220 (1964)
 [96] BETTINI, ET AL., PHYS. LETTERS 16, 83 (1965)
 [97] COOK, ET AL., PHYS. REV. 129, 2743 (1963)
 [98] BOMSE, ET AL., PHYS. REV. 158, 1298 (1967)
 [99] R.W.BLAND, UCRL 18131 (1968)
 [100] SALLSTROM, ET AL., NUOVO CIMENTO 49A, 348 (1967)
 [101] FERRO-LUZZI, ET AL., NUOVO CIMENTO 36, 1101 (1965)
 [102] HIRSCH, ET AL., PHYS. REV. 135, 8191 (1964)
 [103] NEWMAN, ET AL., PHYS. REV. 158, 1310 (1967)
 [104] ODDO, ET AL., PHYS. REV. 177, 1991 (1969)
 [105] SALLSTROM, ET AL., ARKIV FOR FYSIK 37, 468 (1968)
 [106] DEBAERE, ET AL., NUOVO CIMENTO 61A, 397 (1969)
 [107] ANTICH, ET AL., PHYS. REV. LETTERS 21, 1842 (1968)
 [108] BUTTERWORTH, ET AL., PHYS. REV. LETTERS 15, 734 (1965)
 [109] BERLINGHIERI, ET AL., NUC. PHYS. 88, 333 (1968)
 [110] GOLDBABER, ET AL., PHYS. REV. 142, 913 (1966)
- ANTI-LAMBDA
 [111] ALEXANDER, ET AL., PHYS. REV. LETTERS 20, 755 (1968)
- ANTI-SIGMA
 [112] ALEXANDER, ET AL., PHYS. REV. LETTERS 20, 755 (1968)
- BARYON RESONANCE WITH $\gamma=2$ AT 2000 MEV
 [61] CARROLL, ET AL., PHYS. REV. LETTERS 21, 1282 (1968)
- BARYON RESONANCE WITH $\gamma=2$ AT 1480 MEV
 [61] FISK, ET AL., CERN CONFERENCE 358 (1962)
- BRANCHING RATIO
 [48] J.M.BISHOP, THESIS (1967)
- C MESON
 [3] GOLDBABER, ET AL., PHYS. REV. LETTERS 19, 972 (1967)
- CHARGE EXCHANGE
 [55] GOLDSCHMIDT-CLERMONT, ET AL., PHYS. LETTERS 278, 602 (1968)
 [85] SLATER, ET AL., PHYS. REV. LETTERS 7, 378 (1961)
 [87] RAY, ET AL., BNL 13616 (1969)
- CROSS SECTION
 [1] BERLINGHIERI, ET AL., PHYS. LETTERS 278, 665 (1968)
 [2] FERRO-LUZZI, ET AL., PHYS. LETTERS 12, 255 (1964)
 [4] VONHARDT, ET AL., PHYS. REV. LETTERS 5, 733 (1960)
 [6] FISK, ET AL., CERN CONFERENCE 358 (1962)
 [7] STUBBS, ET AL., PHYS. REV. LETTERS 7, 188 (1961)
 [8] GOLDBABER, ET AL., PHYS. REV. LETTERS 9, 135 (1962)
 [9] JOLDERSMA, ET AL., PHYS. REV. LETTERS 17, 716 (1966)
 [10] FIORINI, ET AL., NUOVO CIMENTO 35, 319 (1965)
 [11] ALEXANDER, ET AL., PHYS. REV. LETTERS 20, 755 (1968)
 [15] FERRO-LUZZI, ET AL., PHYS. LETTERS 17, 155 (1965)
 [21] B.KEHO, PHYS. REV. LETTERS 11, 93 (1963)
 [22] BASSOMPIERRE, ET AL., PHYS. LETTERS 268, 30 (1967)
 [23] BASSOMPIERRE, ET AL., NUOVO CIMENTO 48, 589 (1967)
 [24] BURROWS, ET AL., PHYS. REV. LETTERS 2, 117 (1959)
 [25] BOMSE, ET AL., PHYS. REV. LETTERS 20, 1519 (1968)
 [26] BERLINGHIERI, ET AL., CERN HADRON CONFERENCE 172 (1968)
 [27] KYCIA, ET AL., PHYS. REV. 118, 553 (1960)
 [28] BOMSE, ET AL., PHYS. REV. 158, 1281 (1967)
 [29] DEBAISIEUX, ET AL., NUOVO CIMENTO 43A, 142 (1966)
 [30] BUGG, ET AL., PHYS. REV. 168, 1466 (1968)
 [31] BANAIGS, ET AL., PHYS. LETTERS 248, 317 (1967)
 [32] COOK, ET AL., PHYS. REV. LETTERS 7, 182 (1961)
 [33] BAKER, ET AL., PHYS. REV. 124, 2285 (1963)
 [34] MIRATA, ET AL., PHYS. REV. LETTERS 21, 1485 (1968)
 [35] BLAND, ET AL., PHYS. REV. LETTERS 17, 939 (1969)
 [36] FOLEY, ET AL., PHYS. REV. LETTERS 11, 503 (1963)
 [39] CALBRAITH, ET AL., PHYS. REV. 138, 8913 (1965)
 [40] AARMS, ET AL., PHYS. REV. LETTERS 19, 259 (1967)
 [41] CLINE, ET AL., PHYS. REV. LETTERS 19, 675 (1967)
 [42] BUTTERWORTH, ET AL., PHYS. REV. LETTERS 15, 500 (1965)
 [44] STENGER, ET AL., PHYS. REV. 134, 8111 (1964)
 [45] GOLDBABER, ET AL., PHYS. REV. LETTERS 15, 737 (1965)
 [46] ERWIN, ET AL., PHYS. REV. LETTERS 16, 1063 (1966)
 [48] J.M.BISHOP, THESIS (1967)
 [49] COOL, ET AL., PHYS. REV. LETTERS 17, 102 (1966)
 [50] GEORGE, ET AL., NUOVO CIMENTO 49A, 9 (1967)
 [51] DEBAERE, ET AL., NUOVO CIMENTO 45A, 885 (1966)
 [55] GOLDSCHMIDT-CLERMONT, ET AL., PHYS. LETTERS 278, 602 (1968)
 [56] FERRO-LUZZI, ET AL., NUOVO CIMENTO 39, 417 (1965)
 [57] CHINDOWSKY, ET AL., NUC. PHYS. 89, 286 (1969)
 [58] BUCHNER, ET AL., NUC. PHYS. 89, 286 (1969)
 [60] FILIPPAS, ET AL., NUOVO CIMENTO 51A, 1053 (1967)
 [61] CARROLL, ET AL., PHYS. REV. LETTERS 21, 1282 (1968)

- [62] BOLOT, ET AL., PHYS. REV. 133, 8220 (1964)
 [63] BETTINI, ET AL., PHYS. LETTERS 16, 83 (1965)
 [64] COOK, ET AL., PHYS. REV. 129, 2743 (1963)
 [65] R.W.BLAND, UCRL 18131 (1968)
 [69] SALLSTROM, ET AL., NUOVO CIMENTO 49A, 348 (1967)
 [70] FERRO-LUZZI, ET AL., NUOVO CIMENTO 36, 1101 (1965)
 [72] BLAND, ET AL., UCRL 18758 (1968)
 [74] BASSOMPIERRE, ET AL., PHYS. LETTERS 278, 468 (1968)
 [75] NEWMAN, ET AL., PHYS. REV. 158, 1310 (1967)
 [76] FOCARDI, ET AL., PHYS. LETTERS 248, 314 (1967)
 [78] ODDO, ET AL., PHYS. REV. 177, 1991 (1969)
 [79] CHIEN, ET AL., PHYS. LETTERS 288, 615 (1969)
 [80] SALLSTROM, ET AL., ARKIV FOR FYSIK 37, 468 (1968)
 [81] C.FU, UCRL 18417 (1968)
 [82] DEBAERE, ET AL., NUOVO CIMENTO 61A, 397 (1969)
 [83] SLATER, ET AL., PHYS. REV. LETTERS 7, 378 (1961)
 [86] BUTTERWORTH, ET AL., PHYS. REV. LETTERS 15, 734 (1965)
 [88] BERLINGHIERI, ET AL., NUC. PHYS. 88, 333 (1968)
 [89] GOLDBABER, ET AL., PHYS. REV. 142, 913 (1966)
- DALITZ PLOT
 [9] JOLDERSMA, ET AL., PHYS. REV. LETTERS 17, 716 (1966)
 [17] ALMEIDA, ET AL., PHYS. LETTERS 16, 184 (1965)
 [22] BASSOMPIERRE, ET AL., NUOVO CIMENTO 48, 589 (1967)
 [26] BERLINGHIERI, ET AL., CERN HADRON CONFERENCE 172 (1968)
 [35] BLAND, ET AL., PHYS. REV. LETTERS 17, 939 (1966)
 [38] ALEXANDER, ET AL., PHYS. REV. 177, 2092 (1969)
 [43] BUTTERWORTH, ET AL., PHYS. REV. LETTERS 15, 500 (1965)
 [45] GOLDBABER, ET AL., PHYS. REV. LETTERS 15, 737 (1965)
 [48] J.M.BISHOP, THESIS (1967)
 [58] BUCHNER, ET AL., NUC. PHYS. 89, 286 (1969)
 [60] FILIPPAS, ET AL., NUOVO CIMENTO 51A, 1053 (1967)
 [62] BOLOT, ET AL., PHYS. REV. 133, 8220 (1964)
 [65] BOMSE, ET AL., PHYS. REV. 158, 1298 (1967)
 [69] SALLSTROM, ET AL., NUOVO CIMENTO 49A, 348 (1967)
 [70] FERRO-LUZZI, ET AL., NUOVO CIMENTO 36, 1101 (1965)
 [72] BLAND, ET AL., UCRL 18758 (1968)
 [75] NEWMAN, ET AL., PHYS. REV. 158, 1310 (1967)
 [78] ODDO, ET AL., PHYS. REV. 177, 1991 (1969)
 [80] SALLSTROM, ET AL., ARKIV FOR FYSIK 37, 468 (1968)
 [81] C.FU, UCRL 18417 (1968)
 [88] BERLINGHIERI, ET AL., NUC. PHYS. 88, 333 (1968)
 [89] GOLDBABER, ET AL., PHYS. REV. 142, 913 (1966)
- DELTA++
 [26] BERLINGHIERI, ET AL., CERN HADRON CONFERENCE 172 (1968)
 [28] BOMSE, ET AL., PHYS. REV. 158, 1281 (1967)
 [65] BOMSE, ET AL., PHYS. REV. 158, 1298 (1967)
 [69] SALLSTROM, ET AL., NUOVO CIMENTO 49A, 348 (1967)
- DELTA(1232)
 [12] FERRO-LUZZI, ET AL., PHYS. LETTERS 12, 255 (1964)
 [16] GOLDBABER, ET AL., PHYS. LETTERS 18, 76 (1965)
 [17] ALMEIDA, ET AL., PHYS. LETTERS 16, 184 (1965)
 [19] BLAND, ET AL., PHYS. REV. LETTERS 18, 1077 (1967)
 [20] LYNCH, ET AL., PHYS. LETTERS 9, 359 (1964)
 [25] BLAND, ET AL., PHYS. REV. LETTERS 17, 939 (1966)
 [43] BUTTERWORTH, ET AL., PHYS. REV. LETTERS 15, 500 (1965)
 [45] GOLDBABER, ET AL., PHYS. REV. LETTERS 15, 737 (1965)
 [48] J.M.BISHOP, THESIS (1967)
 [50] GEORGE, ET AL., NUOVO CIMENTO 49A, 9 (1967)
 [56] FERRO-LUZZI, ET AL., NUOVO CIMENTO 39, 417 (1965)
 [60] FILIPPAS, ET AL., NUOVO CIMENTO 51A, 1053 (1967)
 [62] BOLDT, ET AL., PHYS. REV. 133, 8220 (1964)
 [63] OCTYINI, ET AL., PHYS. LETTERS 16, 83 (1965)
 [68] R.W.BLAND, UCRL 18131 (1968)
 [70] FERRO-LUZZI, ET AL., NUOVO CIMENTO 36, 1101 (1965)
 [72] BLAND, ET AL., UCRL 18758 (1968)
 [75] NEWMAN, ET AL., PHYS. REV. 158, 1310 (1967)
 [80] SALLSTROM, ET AL., ARKIV FOR FYSIK 37, 468 (1968)
 [88] BERLINGHIERI, ET AL., NUC. PHYS. 88, 333 (1968)
 [89] GOLDBABER, ET AL., PHYS. REV. 142, 913 (1966)
- DENSITY MATRIX
 [20] LYNCH, ET AL., PHYS. LETTERS 9, 359 (1964)
 [25] BOMSE, ET AL., PHYS. REV. LETTERS 20, 1519 (1968)
 [26] BERLINGHIERI, ET AL., CERN HADRON CONFERENCE 172 (1968)
 [28] BOMSE, ET AL., PHYS. REV. 158, 1281 (1967)
 [45] GOLDBABER, ET AL., PHYS. REV. LETTERS 15, 737 (1965)
 [48] J.M.BISHOP, THESIS (1967)
 [50] GEORGE, ET AL., NUOVO CIMENTO 49A, 9 (1967)
 [56] FERRO-LUZZI, ET AL., NUOVO CIMENTO 39, 417 (1965)
 [58] BUCHNER, ET AL., NUC. PHYS. 89, 286 (1969)
 [62] BOLDT, ET AL., PHYS. REV. 133, 8220 (1964)
 [65] BOMSE, ET AL., PHYS. REV. 158, 1298 (1967)
 [68] R.W.BLAND, UCRL 18131 (1968)
 [69] SALLSTROM, ET AL., NUOVO CIMENTO 49A, 348 (1967)
 [70] FERRO-LUZZI, ET AL., NUOVO CIMENTO 36, 1101 (1965)
 [75] NEWMAN, ET AL., PHYS. REV. 158, 1310 (1967)
 [80] SALLSTROM, ET AL., ARKIV FOR FYSIK 37, 468 (1968)
 [81] C.FU, UCRL 18417 (1968)
 [82] DEBAERE, ET AL., NUOVO CIMENTO 61A, 397 (1969)
 [83] ANTICH, ET AL., PHYS. REV. LETTERS 21, 1842 (1968)
 [88] BERLINGHIERI, ET AL., NUC. PHYS. 88, 333 (1968)
- DIFFERENTIAL CROSS SECTION
 [47] BAKER, ET AL., PHYS. LETTERS 288, 291 (1968)
 [50] GEORGE, ET AL., NUOVO CIMENTO 49A, 9 (1967)
 [52] BANAIGS, ET AL., NUC. PHYS. 89, 640 (1969)
 [53] BLAND, ET AL., UCRL 18889 (1969)
 [54] ANDERSSON, ET AL., CERN (1969)
 [55] GOLDSCHMIDT-CLERMONT, ET AL., PHYS. LETTERS 278, 602 (1968)
 [66] AARMS, ET AL., PHYS. REV. LETTERS 21, 1407 (1968)
 [67] BAKER, ET AL., NUC. PHYS. 89, 249 (1969)
 [79] CHIEN, ET AL., PHYS. LETTERS 288, 615 (1969)
 [81] C.FU, UCRL 18417 (1968)
 [82] DEBAERE, ET AL., NUOVO CIMENTO 61A, 397 (1969)
 [85] SLATER, ET AL., PHYS. REV. LETTERS 7, 378 (1961)
 [87] RAY, ET AL., BNL 13616 (1969)
- ETA MESON(548)
 [48] J.M.BISHOP, THESIS (1967)

FITS

- [52] BANAIGS, ET AL., NUC. PHYS. 89, 640 (1969)
 [53] BLAND, ET AL., UCRL 18889 (1969)
 [58] BUCHNER, ET AL., NUC. PHYS. 89, 286 (1969)
 [67] BAKER, ET AL., NUC. PHYS. 89, 249 (1969)
 [72] BLAND, ET AL., UCRL 18758 (1968)
 [78] DODD, ET AL., PHYS. REV. 177, 1991 (1969)
 [79] CHIEN, ET AL., PHYS. LETTERS 288, 615 (1969)
 [81] C.FU, UCRL 18417 (1968)
 [82] DEBAERE, ET AL., NUOVO CIMENTO 61A, 397 (1969)
 [87] RAY, ET AL., BNL 13616 (1969)

KAPPA MESON

- [2] FERRO-LUZZI, ET AL., PHYS. LETTERS 12, 255 (1964)
 [10] FIORINI, ET AL., NUOVO CIMENTO 35, 319 (1965)
 [59] GOSHAU, ET AL., PHYS. LETTERS 22, 347 (1966)
 [60] FILIPPAS, ET AL., NUOVO CIMENTO 51A, 1053 (1967)
 [62] BOLDT, ET AL., PHYS. REV. 133, 8220 (1964)
 [68] R.W.BLAND, UCRL 18131 (1968)

K*(890)

- [2] FERRO-LUZZI, ET AL., PHYS. LETTERS 12, 255 (1964)
 [12] GOLDBERGER, ET AL., PHYS. LETTERS 18, 76 (1965)
 [17] ALMEIDA, ET AL., PHYS. LETTERS 16, 184 (1965)
 [19] BLAND, ET AL., PHYS. REV. LETTERS 18, 1077 (1967)
 [20] LYNCH, ET AL., PHYS. LETTERS 9, 359 (1964)
 [26] BERLINGHIERI, ET AL., CERN HADRON CONFERENCE 172 (1968)
 [28] BOMSE, ET AL., PHYS. REV. 158, 1281 (1967)
 [35] BLAND, ET AL., PHYS. REV. LETTERS 17, 939 (1966)
 [43] BUTTERWORTH, ET AL., PHYS. REV. LETTERS 15, 500 (1965)
 [45] GOLDBERGER, ET AL., PHYS. REV. LETTERS 15, 737 (1965)
 [48] J.M.BISHOP, THESIS (1967)
 [50] GEORGE, ET AL., NUOVO CIMENTO 49A, 9 (1967)
 [56] FERRO-LUZZI, ET AL., NUOVO CIMENTO 39, 417 (1965)
 [58] BUCHNER, ET AL., NUC. PHYS. 89, 286 (1969)
 [62] BOLDT, ET AL., PHYS. REV. 133, 8220 (1964)
 [63] BETTINI, ET AL., PHYS. LETTERS 16, 83 (1965)
 [65] BOMSE, ET AL., PHYS. REV. 158, 1298 (1967)
 [68] R.W.BLAND, UCRL 18131 (1968)
 [69] SALLSTROM, ET AL., NUOVO CIMENTO 49A, 348 (1967)
 [70] FERRO-LUZZI, ET AL., NUOVO CIMENTO 36, 1101 (1965)
 [72] BLAND, ET AL., UCRL 18758 (1968)
 [75] NEWMAN, ET AL., PHYS. REV. 158, 1310 (1967)
 [80] SALLSTROM, ET AL., ARKIV FOR FYSIK 37, 468 (1968)
 [88] BERLINGHIERI, ET AL., NUC. PHYS. 88, 333 (1968)
 [89] GOLDBERGER, ET AL., PHYS. REV. 142, 913 (1966)

K*(1320)

- [3] GOLDBERGER, ET AL., PHYS. REV. LETTERS 19, 972 (1967)
 [5] SHEN, ET AL., PHYS. REV. LETTERS 17, 726 (1966)
 [17] ALMEIDA, ET AL., PHYS. LETTERS 16, 184 (1965)
 [42] BERLINGHIERI, ET AL., PHYS. REV. LETTERS 18, 1087 (1967)
 [48] J.M.BISHOP, THESIS (1967)

K*(1420)

- [3] GOLDBERGER, ET AL., PHYS. REV. LETTERS 19, 972 (1967)
 [5] SHEN, ET AL., PHYS. REV. LETTERS 17, 726 (1966)
 [26] BERLINGHIERI, ET AL., CERN HADRON CONFERENCE 172 (1968)
 [48] J.M.BISHOP, THESIS (1967)
 [50] GEORGE, ET AL., NUOVO CIMENTO 49A, 9 (1967)
 [83] ANTICH, ET AL., PHYS. REV. LETTERS 21, 1842 (1968)
 [88] BERLINGHIERI, ET AL., NUC. PHYS. 88, 333 (1968)

L MESON

- [42] BERLINGHIERI, ET AL., PHYS. REV. LETTERS 18, 1087 (1967)

LIFETIME

- [62] BOLDT, ET AL., PHYS. REV. 133, 8220 (1964)
 [70] FERRO-LUZZI, ET AL., NUOVO CIMENTO 36, 1101 (1965)

MASS

- [62] BOLDT, ET AL., PHYS. REV. 133, 8220 (1964)
 [70] FERRO-LUZZI, ET AL., NUOVO CIMENTO 36, 1101 (1965)

MASS SPECTRUM

- [1] BERLINGHIERI, ET AL., PHYS. LETTERS 278, 665 (1968)
 [2] FERRO-LUZZI, ET AL., PHYS. LETTERS 12, 255 (1964)
 [3] GOLDBERGER, ET AL., PHYS. REV. LETTERS 19, 972 (1967)
 [5] SHEN, ET AL., PHYS. REV. LETTERS 17, 726 (1966)
 [6] FISK, ET AL., CERN CONFERENCE 358 (1962)
 [9] JOLDERSMA, ET AL., PHYS. REV. LETTERS 17, 716 (1966)
 [11] ALEXANDER, ET AL., PHYS. REV. LETTERS 20, 755 (1968)
 [12] GOLDBERGER, ET AL., PHYS. LETTERS 18, 76 (1965)
 [15] FERRO-LUZZI, ET AL., PHYS. LETTERS 17, 155 (1965)
 [17] ALMEIDA, ET AL., PHYS. LETTERS 16, 184 (1965)
 [22] BASSOMPIERRE, ET AL., PHYS. LETTERS 268, 30 (1967)
 [25] BOMSE, ET AL., PHYS. REV. LETTERS 20, 1519 (1968)
 [26] BERLINGHIERI, ET AL., CERN HADRON CONFERENCE 172 (1968)
 [28] BOMSE, ET AL., PHYS. REV. 158, 1281 (1967)
 [38] ALEXANDER, ET AL., PHYS. REV. 177, 2092 (1969)
 [42] BERLINGHIERI, ET AL., PHYS. REV. LETTERS 18, 1087 (1967)
 [43] BUTTERWORTH, ET AL., PHYS. REV. LETTERS 15, 500 (1965)
 [46] ERWIN, ET AL., PHYS. REV. LETTERS 16, 1063 (1966)
 [48] J.M.BISHOP, THESIS (1967)
 [50] GEORGE, ET AL., NUOVO CIMENTO 49A, 9 (1967)
 [56] FERRO-LUZZI, ET AL., NUOVO CIMENTO 39, 417 (1965)
 [59] GOSHAU, ET AL., PHYS. LETTERS 22, 347 (1966)
 [60] FILIPPAS, ET AL., NUOVO CIMENTO 51A, 1053 (1967)
 [62] BOLDT, ET AL., PHYS. REV. 133, 8220 (1964)
 [69] SALLSTROM, ET AL., NUOVO CIMENTO 49A, 348 (1967)
 [72] BLAND, ET AL., UCRL 18758 (1968)
 [74] BASSOMPIERRE, ET AL., PHYS. LETTERS 278, 468 (1968)
 [75] NEWMAN, ET AL., PHYS. REV. 158, 1310 (1967)
 [78] DODD, ET AL., PHYS. REV. 177, 1991 (1969)
 [80] SALLSTROM, ET AL., ARKIV FOR FYSIK 37, 468 (1968)
 [81] C.FU, UCRL 18417 (1968)
 [82] DEBAERE, ET AL., NUOVO CIMENTO 61A, 397 (1969)
 [83] ANTICH, ET AL., PHYS. REV. LETTERS 21, 1842 (1968)
 [88] BERLINGHIERI, ET AL., NUC. PHYS. 88, 333 (1968)

MESON WITH Y=1 AT 1260 MEV

- [78] DODD, ET AL., PHYS. REV. 177, 1991 (1969)

MESON WITH Y=1 AT 720 MEV

- [61] FISK, ET AL., CERN CONFERENCE 358 (1962)

MODELS

- [8] GOLDBERGER, ET AL., PHYS. REV. LETTERS 9, 135 (1962)
 [21] B.KEHO, PHYS. REV. LETTERS 11, 93 (1963)
 [25] BOMSE, ET AL., PHYS. REV. LETTERS 20, 1519 (1968)
 [26] BERLINGHIERI, ET AL., CERN HADRON CONFERENCE 172 (1968)
 [36] FOLEY, ET AL., PHYS. REV. LETTERS 11, 503 (1963)
 [38] ALEXANDER, ET AL., PHYS. REV. 177, 2092 (1969)
 [45] GOLDBERGER, ET AL., PHYS. REV. LETTERS 15, 737 (1965)
 [47] BAKER, ET AL., PHYS. LETTERS 288, 291 (1968)
 [48] J.M.BISHOP, THESIS (1967)
 [50] GEORGE, ET AL., NUOVO CIMENTO 49A, 9 (1967)
 [51] DEBAERE, ET AL., NUOVO CIMENTO 45A, 885 (1966)
 [52] BANAIGS, ET AL., NUC. PHYS. 89, 640 (1969)
 [55] GOLDSCHMIDT-CLERMONT, ET AL., PHYS. LETTERS 278, 602 (1968)
 [56] FERRO-LUZZI, ET AL., NUOVO CIMENTO 39, 417 (1965)
 [64] COOK, ET AL., PHYS. REV. 129, 2743 (1963)
 [66] ABRAMS, ET AL., PHYS. REV. LETTERS 21, 1407 (1968)
 [67] BAKER, ET AL., NUC. PHYS. 89, 249 (1969)
 [68] R.W.BLAND, UCRL 18131 (1968)
 [69] SALLSTROM, ET AL., NUOVO CIMENTO 49A, 348 (1967)
 [70] FERRO-LUZZI, ET AL., NUOVO CIMENTO 36, 1101 (1965)
 [77] ANDERSSON, ET AL., PHYS. LETTERS 288, 611 (1969)
 [79] CHIEN, ET AL., PHYS. LETTERS 288, 615 (1969)
 [80] SALLSTROM, ET AL., ARKIV FOR FYSIK 37, 468 (1968)
 [81] C.FU, UCRL 18417 (1968)
 [82] DEBAERE, ET AL., NUOVO CIMENTO 61A, 397 (1969)
 [83] ANTICH, ET AL., PHYS. REV. LETTERS 21, 1842 (1968)
 [84] BASSOMPIERRE, ET AL., NUC. PHYS. 89, 295 (1969)
 [88] BERLINGHIERI, ET AL., NUC. PHYS. 88, 333 (1968)

MOMENTUM TRANSFER

- [5] SHEN, ET AL., PHYS. REV. LETTERS 17, 726 (1966)

N*(1400)

- [48] J.M.BISHOP, THESIS (1967)
 [78] DODD, ET AL., PHYS. REV. 177, 1991 (1969)

N*(1512)

- [45] GOLDBERGER, ET AL., PHYS. REV. LETTERS 15, 737 (1965)
 [48] J.M.BISHOP, THESIS (1967)

N*(1680)

- [45] GOLDBERGER, ET AL., PHYS. REV. LETTERS 15, 737 (1965)
 [48] J.M.BISHOP, THESIS (1967)

OMEGA MESON

- [2] FERRO-LUZZI, ET AL., PHYS. LETTERS 12, 255 (1964)
 [48] J.M.BISHOP, THESIS (1967)

ONE PARTICLE EXCHANGE

- [17] ALMEIDA, ET AL., PHYS. LETTERS 16, 184 (1965)
 [18] CHADWICK, ET AL., PHYS. LETTERS 6, 309 (1963)
 [20] LYNCH, ET AL., PHYS. LETTERS 9, 359 (1964)
 [21] B.KEHO, PHYS. REV. LETTERS 11, 93 (1963)
 [28] BOMSE, ET AL., PHYS. REV. 158, 1281 (1967)
 [56] FERRO-LUZZI, ET AL., NUOVO CIMENTO 39, 417 (1965)
 [63] BETTINI, ET AL., PHYS. LETTERS 16, 83 (1965)
 [65] BOMSE, ET AL., PHYS. REV. 158, 1298 (1967)
 [70] FERRO-LUZZI, ET AL., NUOVO CIMENTO 36, 1101 (1965)

PHASE SHIFT

- [7] STUBBS, ET AL., PHYS. REV. LETTERS 7, 188 (1961)
 [8] GOLDBERGER, ET AL., PHYS. REV. LETTERS 9, 135 (1962)
 [13] LEA, ET AL., PHYS. REV. 165, 1770 (1968)
 [14] B.R.MARTIN, PHYS. REV. LETTERS 21, 1286 (1968)
 [16] LEA, ET AL., PHYS. LETTERS 23, 380 (1968)
 [27] KYCIA, ET AL., PHYS. REV. 118, 553 (1960)
 [37] A.A.CARTER, PHYS. REV. LETTERS 18, 801 (1967)
 [44] STENGER, ET AL., PHYS. REV. 134, 8111 (1964)
 [54] ANDERSSON, ET AL., CERN (1969)
 [64] COOK, ET AL., PHYS. REV. 129, 2743 (1963)
 [68] R.W.BLAND, UCRL 18131 (1968)
 [73] HIRSCH, ET AL., PHYS. REV. 135, 8191 (1964)
 [76] FOCARDI, ET AL., PHYS. LETTERS 248, 314 (1967)
 [85] SLATER, ET AL., PHYS. REV. LETTERS 7, 378 (1961)
 [87] RAY, ET AL., BNL 13616 (1969)
 [90] ASBURY, ET AL., PHYS. REV. LETTERS 23, 194 (1969)

POLARIZATION

- [54] ANDERSSON, ET AL., CERN (1969)
 [71] FEMINO, ET AL., NUOVO CIMENTO 50A, 372 (1967)
 [73] HIRSCH, ET AL., PHYS. REV. 135, 8191 (1964)
 [77] ANDERSSON, ET AL., PHYS. LETTERS 288, 611 (1969)
 [87] RAY, ET AL., BNL 13616 (1969)
 [90] ASBURY, ET AL., PHYS. REV. LETTERS 23, 194 (1969)

Q PEAK

- [22] BASSOMPIERRE, ET AL., PHYS. LETTERS 268, 30 (1967)
 [25] BOMSE, ET AL., PHYS. REV. LETTERS 20, 1519 (1968)
 [26] BERLINGHIERI, ET AL., CERN HADRON CONFERENCE 172 (1968)
 [42] BERLINGHIERI, ET AL., PHYS. REV. LETTERS 18, 1087 (1967)
 [48] J.M.BISHOP, THESIS (1967)

RHO MESON

- [12] GOLDBERGER, ET AL., PHYS. LETTERS 18, 76 (1965)

SPIN AND PARITY

- [25] BOMSE, ET AL., PHYS. REV. LETTERS 20, 1519 (1968)

STRANGE PARTICLES

- [48] J.M.BISHOP, THESIS (1967)

Z*

- [37] A.A.CARTER, PHYS. REV. LETTERS 18, 801 (1967)
 [74] BASSOMPIERRE, ET AL., PHYS. LETTERS 278, 468 (1968)

In Alphabetical Order

In Numerical Order

	Reference No.		Reference No.
ABRAMS, ET AL., PHYS. REV. LETTERS 19, 259 (1967)	[40]	[1]	BERLINGHIERI, ET AL., PHYS. LETTERS 278, 665 (1968)
ABRAMS, ET AL., PHYS. REV. LETTERS 21, 1407 (1968)	[66]	[2]	FERRO-LUZZI, ET AL., PHYS. LETTERS 12, 255 (1964)
ALEXANDER, ET AL., PHYS. REV. LETTERS 20, 755 (1968)	[11]	[3]	GOLDHABER, ET AL., PHYS. REV. LETTERS 19, 972 (1967)
ALEXANDER, ET AL., PHYS. REV. 177, 2092 (1969)	[38]	[4]	VONARDEL, ET AL., PHYS. REV. LETTERS 5, 333 (1960)
ALMEIDA, ET AL., PHYS. LETTERS 16, 184 (1965)	[17]	[5]	SHEN, ET AL., PHYS. REV. LETTERS 17, 726 (1966)
ANDERSSON, ET AL., CERN (1969)	[54]	[6]	FISK, ET AL., CERN CONFERENCE 358 (1962)
ANDERSSON, ET AL., PHYS. LETTERS 288, 611 (1969)	[77]	[7]	STUBBS, ET AL., PHYS. REV. LETTERS 7, 188 (1961)
ANTICH, ET AL., PHYS. REV. LETTERS 21, 1842 (1968)	[83]	[8]	GOLDHABER, ET AL., PHYS. REV. LETTERS 9, 135 (1962)
ASBURY, ET AL., PHYS. REV. LETTERS 23, 194 (1969)	[90]	[9]	JOLDERSMA, ET AL., PHYS. REV. LETTERS 17, 716 (1966)
BAKER, ET AL., PHYS. REV. 129, 2285 (1963)	[33]	[10]	FIORINI, ET AL., NUOVO CIMENTO 35, 319 (1965)
BAKER, ET AL., PHYS. LETTERS 288, 291 (1968)	[47]	[11]	ALEXANDER, ET AL., PHYS. REV. LETTERS 20, 755 (1968)
BAKER, ET AL., NUC. PHYS. 89, 249 (1969)	[67]	[12]	GOLDHABER, ET AL., PHYS. LETTERS 18, 76 (1965)
BANAIGS, ET AL., PHYS. LETTERS 248, 317 (1967)	[31]	[13]	LEA, ET AL., PHYS. REV. 165, 1770 (1968)
BANAIGS, ET AL., NUC. PHYS. 89, 640 (1969)	[52]	[14]	MARTIN, PHYS. REV. LETTERS 21, 1286 (1968)
BASSOMPIERRE, ET AL., PHYS. LETTERS 268, 30 (1967)	[22]	[15]	FERRO-LUZZI, ET AL., PHYS. LETTERS 17, 155 (1965)
BASSOMPIERRE, ET AL., NUOVO CIMENTO 48, 589 (1967)	[23]	[16]	LEA, ET AL., PHYS. LETTERS 23, 380 (1968)
BASSOMPIERRE, ET AL., PHYS. LETTERS 278, 468 (1968)	[74]	[17]	ALMEIDA, ET AL., PHYS. LETTERS 16, 184 (1965)
BASSOMPIERRE, ET AL., NUC. PHYS. 89, 295 (1969)	[84]	[18]	CHADWICK, ET AL., PHYS. LETTERS 6, 309 (1963)
BERLINGHIERI, ET AL., PHYS. LETTERS 278, 665 (1968)	[11]	[19]	BLAND, ET AL., PHYS. REV. LETTERS 18, 1077 (1967)
BERLINGHIERI, ET AL., CERN HADRON CONFERENCE 172 (1968)	[26]	[20]	LYNCH, ET AL., PHYS. LETTERS 9, 359 (1964)
BERLINGHIERI, ET AL., PHYS. REV. LETTERS 18, 1087 (1967)	[42]	[21]	KEHO, PHYS. REV. LETTERS 11, 93 (1963)
BERLINGHIERI, ET AL., NUC. PHYS. 88, 333 (1968)	[88]	[22]	BASSOMPIERRE, ET AL., PHYS. LETTERS 268, 30 (1967)
BETTINI, ET AL., PHYS. LETTERS 16, 83 (1965)	[63]	[23]	BASSOMPIERRE, ET AL., NUOVO CIMENTO 48, 589 (1967)
BISHOP, THESIS (1967)	[48]	[24]	BURROWS, ET AL., PHYS. REV. LETTERS 2, 117 (1959)
BLAND, ET AL., PHYS. REV. LETTERS 18, 1077 (1967)	[19]	[25]	BOMSE, ET AL., PHYS. REV. LETTERS 20, 1519 (1968)
BLAND, ET AL., PHYS. REV. LETTERS 17, 939 (1966)	[35]	[26]	BERLINGHIERI, ET AL., CERN HADRON CONFERENCE 172 (1968)
BLAND, ET AL., UCRL 18889 (1969)	[53]	[27]	KYCIA, ET AL., PHYS. REV. 118, 553 (1960)
BLAND, UCRL 18131 (1968)	[68]	[28]	BOMSE, ET AL., PHYS. REV. 158, 1281 (1967)
BLAND, ET AL., UCRL 18758 (1968)	[72]	[29]	DEBAISIEUX, ET AL., NUOVO CIMENTO 43A, 142 (1966)
BOLDT, ET AL., PHYS. REV. 133, 8220 (1964)	[62]	[30]	BUGG, ET AL., PHYS. REV. 168, 1466 (1968)
BOMSE, ET AL., PHYS. REV. LETTERS 20, 1519 (1968)	[25]	[31]	BANAIGS, ET AL., PHYS. LETTERS 248, 317 (1967)
BOMSE, ET AL., PHYS. REV. 158, 1281 (1967)	[28]	[32]	COOK, ET AL., PHYS. REV. LETTERS 7, 182 (1961)
BOMSE, ET AL., PHYS. REV. 158, 1298 (1967)	[65]	[33]	BAKER, ET AL., PHYS. REV. 129, 2285 (1963)
BUCHNER, ET AL., NUC. PHYS. 89, 286 (1969)	[58]	[34]	HIRATA, ET AL., PHYS. REV. LETTERS 21, 1485 (1968)
BUGG, ET AL., PHYS. REV. 168, 1466 (1968)	[30]	[35]	BLAND, ET AL., PHYS. REV. LETTERS 17, 939 (1966)
BURROWS, ET AL., PHYS. REV. LETTERS 2, 117 (1959)	[24]	[36]	FOLEY, ET AL., PHYS. REV. LETTERS 11, 503 (1963)
BUTTERWORTH, ET AL., PHYS. REV. LETTERS 15, 500 (1965)	[43]	[37]	CARTER, PHYS. REV. LETTERS 18, 801 (1967)
BUTTERWORTH, ET AL., PHYS. REV. LETTERS 15, 734 (1965)	[86]	[38]	ALEXANDER, ET AL., PHYS. REV. 177, 2092 (1969)
CARROLL, ET AL., PHYS. REV. LETTERS 21, 1282 (1968)	[61]	[39]	GALBRAITH, ET AL., PHYS. REV. 138, 8913 (1965)
CARTER, PHYS. REV. LETTERS 18, 801 (1967)	[37]	[40]	ABRAMS, ET AL., PHYS. REV. LETTERS 19, 259 (1967)
CHADWICK, ET AL., PHYS. LETTERS 6, 309 (1963)	[18]	[41]	CLINE, ET AL., PHYS. REV. LETTERS 19, 675 (1967)
CHIEN, ET AL., PHYS. LETTERS 288, 615 (1969)	[79]	[42]	BERLINGHIERI, ET AL., PHYS. REV. LETTERS 18, 1087 (1967)
CHINOWSKY, ET AL., PHYS. REV. 139, 81411 (1965)	[57]	[43]	BUTTERWORTH, ET AL., PHYS. REV. LETTERS 15, 500 (1965)
CLINE, ET AL., PHYS. REV. LETTERS 19, 675 (1967)	[41]	[44]	STENGER, ET AL., PHYS. REV. 134, 81111 (1964)
COOK, ET AL., PHYS. REV. LETTERS 7, 182 (1961)	[32]	[45]	GOLDHABER, ET AL., PHYS. REV. LETTERS 15, 737 (1965)
COOK, ET AL., PHYS. REV. 129, 2743 (1963)	[64]	[46]	ERWIN, ET AL., PHYS. REV. LETTERS 16, 1063 (1966)
COOL, ET AL., PHYS. REV. LETTERS 17, 102 (1966)	[49]	[47]	BAKER, ET AL., PHYS. LETTERS 288, 291 (1968)
DEBAERE, ET AL., NUOVO CIMENTO 45A, 885 (1966)	[51]	[48]	BISHOP, THESIS (1967)
DEBAERE, ET AL., NUOVO CIMENTO 61A, 397 (1969)	[82]	[49]	COOL, ET AL., PHYS. REV. LETTERS 17, 102 (1966)
DEBAISIEUX, ET AL., NUOVO CIMENTO 43A, 142 (1966)	[29]	[50]	GEORGE, ET AL., NUOVO CIMENTO 49A, 9 (1967)
DODD, ET AL., PHYS. REV. 177, 1991 (1967)	[78]	[51]	DEBAERE, ET AL., NUOVO CIMENTO 45A, 885 (1966)
ERWIN, ET AL., PHYS. REV. LETTERS 16, 1063 (1966)	[46]	[52]	BANAIGS, ET AL., NUC. PHYS. 89, 640 (1969)
FEMINO, ET AL., NUOVO CIMENTO 50A, 372 (1967)	[71]	[53]	BLAND, ET AL., UCRL 18889 (1969)
FERRO-LUZZI, ET AL., PHYS. LETTERS 12, 255 (1964)	[12]	[54]	ANDERSSON, ET AL., CERN (1969)
FERRO-LUZZI, ET AL., PHYS. LETTERS 17, 155 (1965)	[15]	[55]	GOLDSCHMIDT-CLERMONT, ET AL., PHYS. LETTERS 278, 602 (1968)
FERRO-LUZZI, ET AL., NUOVO CIMENTO 39, 417 (1965)	[56]	[56]	FERRO-LUZZI, ET AL., NUOVO CIMENTO 39, 417 (1965)
FERRO-LUZZI, ET AL., NUOVO CIMENTO 36, 1101 (1965)	[70]	[57]	CHINOWSKY, ET AL., PHYS. REV. 139, 81411 (1965)
FILIPPAS, ET AL., NUOVO CIMENTO 51A, 1053 (1967)	[60]	[58]	BUCHNER, ET AL., NUC. PHYS. 89, 286 (1969)
FIORINI, ET AL., NUOVO CIMENTO 35, 319 (1965)	[10]	[59]	GOSHAU, ET AL., PHYS. LETTERS 22, 347 (1966)
FISK, ET AL., CERN CONFERENCE 358 (1962)	[6]	[60]	FILIPPAS, ET AL., NUOVO CIMENTO 51A, 1053 (1967)
FOCARDI, ET AL., PHYS. LETTERS 248, 314 (1967)	[76]	[61]	CARROLL, ET AL., PHYS. REV. LETTERS 21, 1282 (1968)
FOLEY, ET AL., PHYS. REV. LETTERS 11, 503 (1963)	[36]	[62]	BOLDT, ET AL., PHYS. REV. 133, 8220 (1964)
FU, UCRL 18417 (1968)	[81]	[63]	BETTINI, ET AL., PHYS. LETTERS 16, 83 (1965)
GALBRAITH, ET AL., PHYS. REV. 138, 8913 (1965)	[39]	[64]	COOK, ET AL., PHYS. REV. 129, 2743 (1963)
GEORGE, ET AL., NUOVO CIMENTO 49A, 9 (1967)	[50]	[65]	BOMSE, ET AL., PHYS. REV. 158, 1298 (1967)
GOLDHABER, ET AL., PHYS. REV. LETTERS 19, 972 (1967)	[3]	[66]	ABRAMS, ET AL., PHYS. REV. LETTERS 21, 1407 (1968)
GOLDHABER, ET AL., PHYS. REV. LETTERS 9, 135 (1962)	[8]	[67]	BAKER, ET AL., NUC. PHYS. 89, 249 (1969)
GOLDHABER, ET AL., PHYS. LETTERS 18, 76 (1965)	[12]	[68]	BLAND, UCRL 18131 (1968)
GOLDHABER, ET AL., PHYS. REV. LETTERS 15, 737 (1965)	[45]	[69]	SALLSTROM, ET AL., NUOVO CIMENTO 49A, 348 (1967)
GOLDHABER, ET AL., PHYS. REV. 142, 913 (1966)	[89]	[70]	FERRO-LUZZI, ET AL., NUOVO CIMENTO 36, 1101 (1965)
GOLDSCHMIDT-CLERMONT, ET AL., PHYS. LETTERS 278, 602 (1968)	[55]	[71]	FEMINO, ET AL., NUOVO CIMENTO 50A, 372 (1967)
GOSHAU, ET AL., PHYS. LETTERS 22, 347 (1966)	[59]	[72]	BLAND, ET AL., UCRL 18758 (1968)
HIRATA, ET AL., PHYS. REV. LETTERS 21, 1485 (1968)	[34]	[73]	HIRSCH, ET AL., PHYS. REV. 135, 8191 (1964)
HIRSCH, ET AL., PHYS. REV. 135, 8191 (1964)	[73]	[74]	BASSOMPIERRE, ET AL., PHYS. LETTERS 278, 468 (1968)
JOLDERSMA, ET AL., PHYS. REV. LETTERS 17, 716 (1966)	[9]	[75]	NEWMAN, ET AL., PHYS. REV. 158, 1310 (1967)
KEHO, PHYS. REV. LETTERS 11, 93 (1963)	[21]	[76]	FOCARDI, ET AL., PHYS. LETTERS 248, 314 (1967)
KYCIA, ET AL., PHYS. REV. 118, 553 (1960)	[27]	[77]	ANDERSSON, ET AL., PHYS. LETTERS 288, 611 (1969)
LEA, ET AL., PHYS. REV. 165, 1770 (1968)	[13]	[78]	DODD, ET AL., PHYS. REV. 177, 1991 (1967)
LEA, ET AL., PHYS. LETTERS 23, 380 (1968)	[16]	[79]	CHIEN, ET AL., PHYS. LETTERS 288, 615 (1969)
LYNCH, ET AL., PHYS. LETTERS 9, 359 (1964)	[20]	[80]	SALLSTROM, ET AL., ARKIV FOR FYSIK 37, 468 (1968)
MARTIN, PHYS. REV. LETTERS 21, 1286 (1968)	[14]	[81]	FU, UCRL 18417 (1968)
NEWMAN, ET AL., PHYS. REV. 158, 1310 (1967)	[75]	[82]	DEBAERE, ET AL., NUOVO CIMENTO 61A, 397 (1969)
RAY, ET AL., BNL 13616 (1969)	[87]	[83]	ANTICH, ET AL., PHYS. REV. LETTERS 21, 1842 (1968)
SALLSTROM, ET AL., NUOVO CIMENTO 49A, 348 (1967)	[69]	[84]	BASSOMPIERRE, ET AL., NUC. PHYS. 89, 295 (1969)
SALLSTROM, ET AL., ARKIV FOR FYSIK 37, 468 (1968)	[80]	[85]	SLATER, ET AL., PHYS. REV. LETTERS 7, 378 (1961)
SHEN, ET AL., PHYS. REV. LETTERS 17, 726 (1966)	[5]	[86]	BUTTERWORTH, ET AL., PHYS. REV. LETTERS 15, 734 (1965)
SLATER, ET AL., PHYS. REV. LETTERS 7, 378 (1961)	[85]	[87]	RAY, ET AL., BNL 13616 (1969)
STENGER, ET AL., PHYS. REV. 134, 81111 (1964)	[44]	[88]	BERLINGHIERI, ET AL., NUC. PHYS. 88, 333 (1968)
STUBBS, ET AL., PHYS. REV. LETTERS 7, 188 (1961)	[7]	[89]	GOLDHABER, ET AL., PHYS. REV. 142, 913 (1966)
VONARDEL, ET AL., PHYS. REV. LETTERS 5, 333 (1960)	[14]	[90]	ASBURY, ET AL., PHYS. REV. LETTERS 23, 194 (1969)

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.