15

JUN 22 1965

MASTER

MASSACHUSETTS INSTITUTE OF TECHNOLOGY

Laboratory for Nuclear Science

PROGRESS REPORT

November 1, 1964

M. I. T. - 2098 - No. 142

PATENT CLEARANCE OBTAINED. RELEASE TO
THE PUBLIC IS APPROVED. PROCEDURES
ARE ON FILE IN THE RECEIVING SECTION.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

MASSACHUSETTS INSTITUTE OF TECHNOLOGY

LABORATORY FOR NUCLEAR SCIENCE

PROGRESS REPORT

November 1, 1964

M. I. T. -2098-No. 142

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

Reproduction in whole or in part is permitted for any purpose by the U. S. Government. The data and the results that are presented in this report should not otherwise be published without prior consultation with the Laboratory for Nuclear Science.

Researches herein reported by the Physics Groups have been supported by and large by the Atomic Energy Commission through AEC contract AT(30-1)-2098 and in part by separate contracts and grants with the Air Force Office of Aerospace Research (contract AF19(628)-2389), the National Aeronautics and Space Administration (contracts NASw-37, NASA-2053, NAS5-3205, grant number NsG-386) and the National Science Foundation (grant number G-19728). The work of the Chemistry Groups has been supported by the Atomic Energy Commission under contract AT(30-1)-905.

This is the sixty-fifth progress report of the Laboratory for Nuclear Science at the Massachusetts Institute of Technology. Progress reported covers the period for Physics and Chemistry through November 1, 1964.

TABLE OF CONTENTS

Chemistry of the Fission Elements Group	1
Nuclear Chemistry (Inorganic) Group	11
Nuclear Chemistry (Organic) Group	52
Cosmic Ray Group	62
Accelerator Physics Collaboration	74
Linear Accelerator Group	77
High Energy Accelerator Physics Group	82
ONR Generator Group	92
Radioactivity Group	101
Cyclotron Group	119
Theoretical Group	134
Personnel Listing	144

Reprints of published papers listed herein are available in limited numbers upon request from the Laboratory for Nuclear Science Document Room.

CHEMISTRY OF THE FISSION ELEMENTS

I. Complex Ions

A. Spectrophotometric Studies

It has been determined that TiH_2 dissolved in HBr free of atmospheric oxygen yields the Ti(III) aquo ion with the least amounts of contaminants. This ion exhibits a weak band in the visible region at $510m\mu$, which is suitable for study and avoids possible interference from an impurity in the HBr that absorbs in the same region as the UV band of the Ti(III) aquo ion. KSCN and o-phenanthroline have been found to yield complexes with Ti(III) having absorption maxima at 540 and 490 $m\mu$, respectively. These complexes and those of EDTA and fluoride are to be studied further as a function of both acid and ligand concentrations. (A. S. Gordon)

B. Mixed Ligand Complexes

In the study of the system Ag(I)-pyridine- γ -picoline it has been found that the gas chromatographic method of measuring only the ligand concentration is not accurate enough. The metal ion concentration also had to be determined, and a potentiometric method was decided upon.

After calibration of both methods, the two single ligands as well as the mixed ligand system will be studied. (J. N. Little)

II. Plasma Spectroscopy

Studies into specific inter-element effects in the excitation of metal spectra by plasma torch spectroscopy have been pursued. Using a pneumatic atomizer, the usual inter-element effects, e.g., phosphate depression of calcium emission, were noted. An attempt was made to eliminate these effects, since this atomizer system at present is more stable than the ultrasonic atomizer system employed. When sodium ion was added to the calcium solution containing phosphate, the calcium emission was increased above the level of emission without phosphate present. This effect was also noted for strontium. Both the atomic and ion lines are affected. The effect of the other alkalis was also determined. For a given concentration of alkali ion, the degree of emission enhancement produced was potassium > sodium $\gg$ lithium.

Previous studies had indicated no emission enhancement or depression due to inter-element effects when ultrasonic atomization of the sample was used. However, it was found that the presence of alkali metal ions in the sample solution caused marked enhancement in the emission of the alkaline earths. This problem is still to be resolved. The atomic ratios involved are approximately 5 alkali/alkaline earth atom. This type of system could readily be expected in samples, and so the interaction should be thoroughly studied. (A. L. Malenfant and T. W. Gilbert)

III. Thermometric Titrations

A preliminary investigation has been made of the thermometric titration of organic bases with perchloric acid in an isopropyl alcohol solvent system. The volumetric apparatus used is essentially according to the design of B. J. Duffield¹, while the thermistor bridge used is that of Keily². After an approximate calibration of the apparatus, aniline (pK_b in water 9.4) and n-butylamine in the concentration range 0.1 to 5×10^{-3} M were successfully titrated with perchloric acid, while m-nitroaniline (pK_b 11.5) could not be titrated.

The exact calibration of the buret and recorder was done by the method of Duffield. The syringe proved to be quite constant over most of its length, with the 95% confidence limits of the average flow rate being 0.1%. From the coefficient of cubic expansion of isopropyl alcohol it was calculated that a 4°C variation in the ambient temperature would introduce an additional error of 0.4% in the flow rate. (R. C. Berlo)

IV. Alternating Current Polarography

As mentioned in the last report, construction was begun on an improved version of an a.c. polarograph employing phase sensitive detection. This instrument has since been completed and partially calibrated.

With the major instrumental problems nearly over, attention is now being directed toward the chemical objectives of this work. As it was proposed to study the effect of high concentrations of supporting electrolytes on the reversibility of a.c. polarographic waves, it was decided to start with a classical reversible and irreversible electroactive substance to see what effect, if any, various concentrated supporting electrolytes had on them regarding their reversibility.

As a start, Cd is used for the reversible system and Ni and Co for the irreversible system. Supporting electrolytes to be studied include KCNS, K citrate and K tartrate. Preliminary results with Ni and Co in KCNS supporting electrolyte show no significant increase in reversibility, as measured by the height of the a.c. wave, when the concentration of the KCNS supporting electrolyte is increased from 0.1 to 9 M. Similar results have been indicated with potassium citrate as the supporting electrolyte. (T.F. Retajczyk, Jr.)

V. Determination of Amides

Work has been done in developing a method for quantitatively determining nitriles. The nitrile is first converted to the amide by reacting with a concentrated (~90%) basic solution of hydrogen peroxide. After conversion, the amide is saponified to the corresponding sodium salt of the acid by boiling the original solution to concentrate the base. After conversion of the amide,

the resulting solution is passed through a heated ($\sim 80^{\circ}\text{C}$) cation exchange resin in the hydrogen form. The cation of the acid exchanges for H^{+} , and the resulting acid solution is titrated potentiometrically with standard alkali solution.

The concentrations and amounts of the reacting components were optimized in order to effect the most complete conversion in the most reasonable amount of time. Also, the blank titration value needed to be minimized and stabilized. (D. M. Desiderio)

VI. Polarography of Cyclooctatetraene and Its Silver Adducts

Silver ion reduction at a DME has a pronounced maximum which obscures changes in the silver wave due to cyclooctatetraene (COT). The maximum depends upon Ag^{+} and supporting electrolyte concentrations. Thus, with less than $1 \text{ mM } \text{Ag}^{+}$ and $0.1 \text{ M } \text{KNO}_3$ the maximum was eliminated. The addition of COT to this solution results in a slight suppression of the current between $+0.35$ and 0 V vs SCE , which may be due to adduct formation. A sample of the prepared adduct AgCOTNO_3 will be used to check this behavior. (P. Vouros)

VII. Solubility of Metallic Mercury

It has been reported in the literature³ that dissolved Hg atoms in water have an absorption band at $2537 \text{ \AA}$. The absorbance at this wavelength has been followed as a function of the applied voltage in order to study the solubility of Hg as a function of voltage, the absorption cell itself serving as the electrolytic cell. It was found that the absorbance increases as the potential is increased in the cathodic direction. Further studies are necessary to elucidate the exact nature of this relationship. (A. S. Gordon)

VIII. The Galvanostatic Method for Fast Electrode Reactions

During the period covered by this report, the theory of the galvanostatic method was thoroughly reviewed and written up for publication in two parts under the title: "The Galvanostatic Method for Fast Reactions". The first paper covers the single-pulse method. A general equation, for the case where the double-layer is not considered, was derived and gives a quantitative guide to the kinetically useful region where the curve is linear. In addition, a more careful consideration of the approximations used to derive the equation where the double layer is considered allows the method to be extended to the measurement of rate constants one order of magnitude larger, although an additional time dependent term must be added to the equation. The measurement is achieved by a somewhat tedious iteration process.

In the second paper, the modified experimental technique of precharging the double layer was discussed. A theoretical derivation shows that the additional term in the previously mentioned equation is reduced to negligibility by using precharging, and the method is applicable without increased mathematical complexity. A comparison between derived equations and experimental results was made in both papers for the $\text{Hg}_2^{++}/\text{Hg}$ reaction and its kinetic rate constant was measured. (R. L. Birke)

IX. Potentiostatic Chromatography-Amalgam Electrodes

In the comparison of different forms of amalgamated nickel electrodes in anodic stripping voltammetry, it was found that different behavior resulted from edge effects. Exposed nickel was oxidized and passivated and the current-voltage curves of metal ion reduction and oxidation were distorted. A new electrode design avoids these problems, and it is now possible to observe the effect of nickel dissolved in the mercury upon the electrode reactions.

Column operation of amalgamated nickel granules, as previously described, is being attempted. Spherical granules have been obtained and a procedure for satisfactory amalgamation has been developed. (J. E. A. Toni)

X. Single Crystal Electrodes

During the past period, the 111 -- and to a lesser extent -- 100 faces of Ag single crystals were studied in an attempt to obtain reproducible potentials of these faces as a silver reference electrode. The faces were anodized at current densities from $1 \mu\text{a}/\text{cm}^2$ to $10 \mu\text{a}/\text{cm}^2$, the potential-time behavior being followed during the process. In general, the potential-time behavior was specific for a given 111 and 100 face on a crystal, and was different but reproducible for each of the four 111 faces studied on each crystal. As a rule, the final rest potential of the 100 face, after several anodizations, was 1 mV more negative than the corresponding value for the 111 faces. The results were complicated by variations in the potential of the reference electrode, and the much smaller area of the 100 face as compared with the 111 face.

Results among 111 faces indicate that the condition of the surface has a great affect on its behavior as regards the potential time characteristics during anodization and its rest potential afterwards. 111 faces that were rough under low magnification yielded rest potentials that were considerably less noble than those on a macroscopically undisturbed 111 face.

Recently equipment has been set up to determine the current potential curves for the 111 and 100 faces of silver. As yet, no results have been obtained. (W. M. Krebs)

XI. Behavior of HgSe as an Indicator Electrode

Further investigations of the surface oxidation processes of HgSe electrodes has led to the tentative assignment of half-reactions corresponding to three anodic peaks and one cathodic peak. These include the following couples and measured potentials vs SCE in 1 M NaNO₃: Hg, SeO₂/HgSe, 0.54 V; HgSeO₃/Hg, SeO₂, 0.59 V; Hg²⁺/Hg, 0.61 V; and the cathodic process of HgSeO₃/Hg²⁺, Se, 0.15 V. Since most of these processes involve H⁺, a series of measurements with buffer solutions will be made to permit a comparison of these potentials with tabulated free energies. (P. J. Sherwood)

XII. Experimental Evaluation of Transition Times

As a deviation from the preceding experimental approach to make a direct measurement of transition times in chronopotentiometry, an equation has been derived for the rapidly rising final portion of the potential-time curve. This equation is based on a solution of Fick's second law of diffusion with a series approximation of the Sand equation as an initial condition. The results give a good approximation to the difference between the applied current and the (decreasing) faradaic current beyond the transition time; it is this current that is available for double layer capacity charging. A series of measurements has been started, which will be used to test this equation and also to obtain a direct measurement of transition times. A novel mercury pool electrode with an amalgamated platinum circumferential ring was developed for these measurements so that the electrode surface would be planar. (D. K. Roe)

XIII. Voltammetry of Naphthalene and Naphthalenide Ion

Current-voltage curves of the oxidation of naphthalenide ion in tetrahydrofuran have been measured with a gold indicator electrode. Two oxidation waves resulted, starting at -1.7 and -0.7 vs AgCl (sat. LiCl)/Ag. The ratio of the limiting currents of the first to second wave was 1:2. This ratio was maintained as the concentration of radical anion decreased because of a slow influx of water. (P. D. Anderson)

XIV. I¹²⁹ as a Non-Radioactive Tracer

The detection limit of I¹²⁹ after neutron activation to I¹³⁰ (0.53 MeV γ , $t_{1/2}$ = 12.5 hr) was obscured by the high background due to Compton scattering of a peak at 1.29 MeV. This impurity was found to be Ar⁴¹ produced by activation of atmospheric Ar⁴⁰. By simply heating the open

vials for a short time before counting, this interference was eliminated. Without chemical pretreatment or separation, it was possible to measure 10^{-9} gm of I^{130} , which might be useful in certain cases of biological interest. A description of this method has been submitted to the Journal of Nuclear Medicine. (J. T. Watson)

XV. Electrochemically Generated Chemiluminescence

The initial program of research will involve the application of triangular wave cyclic voltammetry to systems of polynuclear aromatic hydrocarbons in various solvents and supporting electrolytes.

The apparatus for obtaining the cyclic voltammetric traces has been assembled, and preliminary runs have been carried out using the specially designed electrochemical cell.

It is also hoped to observe the chemiluminescence of these systems, and to record the electroluminescence intensity as functions of the applied voltage and frequency.

In addition, some thin layer chromatography has been attempted in order to investigate its applicability for estimating the purity of the various hydrocarbons to be used for the investigation. (R. C. Lansbury)

XVI. Electroluminescence of Organic Compounds

Various fluorescent organic compounds have been tested for electroluminescent activity by subjecting their layers to rapidly changing high voltage electric fields. The emitted light was examined spectroscopically using a Bausch and Lomb medium quartz spectrograph. It was found that bands were observed in every spectrum in addition to the ultraviolet fluorescence spectrum of the compound in the cell. These bands have been identified as belonging to the second positive system of nitrogen. It was realized that the light emission could be explained as the result of fluorescence caused by an internal emission of ultraviolet radiation from a nitrogen glow discharge.

In support of this mechanism it was found that no light emission was observed from cells run in an atmosphere of helium, although a commercial electroluminescent fluorophore continued to glow. Furthermore, it has been shown that all the observed data can be explained on the basis of a glow discharge-excitation mechanism. Previous work in the field of organic electroluminescence has been critically reviewed in the light of this work. This work is to be submitted for publication in the very near future. (G. D. Short)

XVII. Chemiluminescence of Luminol

Production of light from the oxidation of luminol (8-aminophthalhydrazide) by molecular oxygen in basic solution has been extensively studied. No single mechanism in either aqueous or non-aqueous systems has as yet gained general acceptance, and much remains to be done before the situation can be clarified. It can be foreseen that conclusive proof of a mechanism must involve several different, though mutually supporting, approaches. A combination of electrochemical, spectroscopic, and kinetic methods appears to be the best way to approach this problem. It is hoped that these, coupled with electron spin resonance, may yield enough data to propose a more definitive mechanism than has yet been put forward. (G. D. Short)

XVIII. Photochemical Analysis

A study has been undertaken to apply differential kinetic analysis to species which photolyze at different rates. A theoretical treatment has shown that it is possible, in principle, to set up simultaneous equations that may be solved for the individual concentrations of the species being photolyzed, even if there is an inter-filter effect and the reactions take place in an absorbing medium. At present the photochemical apparatus is being constructed and a system of methyl, methoxy, and hydroxy substituted methyl benzophenones is being purified in order to develop the method. It is hoped that the method will be useful for determining species that have similar structures and similar absorption spectra. (R. L. Birke)

XIX. Electrochemistry of Triplet States

Construction of flash equipment was completed and the flash duration was determined to be $\sim 70 \mu\text{sec}$. Although this duration was long compared to the desired $5 \mu\text{sec}$, it was decided to delay modifications of the circuits until it proves necessary to do so. The wave generator constructed during the last period was calibrated and operates satisfactorily with the Jaissle potentiostat.

The problems of shielding the electronics from the voltage surge caused by firing the flash lamp were largely eliminated by shielding the leads and grounding the working electrode. The spike was reduced from $\sim 40 \text{ V}$ to $\sim 250 \text{ mV}$ in this way. It now appears as if further shielding would be necessary around the spark gap and/or trigger transformer to further reduce this voltage surge.

The technique of preparing the deoxygenated THF has been greatly improved during this period. The time necessary for the solvent purification has been cut down from ~ 2 weeks to ~ 3 -4 days by this improved technique. (D. N. Bailey)

XX. Fluorescence Spectra of the Naphthoic Acids

The fluorescence spectra of 1- and 2-naphthoic acids are being studied under conditions of constant ionic strength in the pH range from 1 to 11. The initial measurements on the 2-naphthoic acid in this range have indicated three fluorescent species. In the same range 1-naphthoic acid gives evidence of only one fluorescent species. However, increase of the acidity to $\text{pH} < 1$ results in a second peak at longer wavelength, due to the protonated species in the strongly acid solution.

In the region of rapid change of the relative peak intensities of the fluorescent species in 2-naphthoic acid, spectra were run every 0.5 pH units in order to determine the excited state $\text{pK}_a(s)$. It is planned to investigate 1-naphthoic acid in the same manner. (A. L. Malenfant)

XXI. Fluorescence of 3-Aminophthalic Acid

The fluorescence of 3-aminophthalic acid (3APA) is being studied because 3APA is known to be the final product and the emitting species in the chemiluminescent reaction of luminol. The emission spectrum of the luminol reaction varies with solvent and does not always match the fluorescence of 3APA in the same medium. Also, energy transfer is observed during the luminol reaction when a suitable acceptor is present. It is desired to determine if energy transfer occurs from an excited state of 3APA, or if some intermediate species is responsible for the transfer.

To date, serious difficulties have been encountered in purification of 3APA. A green fluorescence impurity is present, which is virtually impossible to remove completely by recrystallization. However, it now appears that chromatography of 3APA on alumina from $\sim 3\text{N HCl}$ will enact the desired separation. (D. M. Hercules)

XXII. Fluorescence of Transition Metal Ion Chelates

Metal chelate compounds of transition metal ions usually do not fluoresce. Of the few that do show this property, in only one case has the phenomenon been explained in terms of the combining electronic energy levels. Several selected systems will be examined with a view to identification of the combining states and the critical parameters which lead to a fluorescent transition metal ion chelate. It is anticipated on the basis of preliminary results that several analytical procedures will be developed. (W. E. Ohnesorge, R. C. Berlo)

XXIII. Fluorescence of Hydroxy-Naphtoic Acids

The fluorescence of 1-hydroxy-2-naphthoic acid, 2-hydroxy-1-naphthoic acid and 2-hydroxy-3-naphthoic acid, their methyl esters, their methyl ethers and the corresponding methoxy-methyl esters has been studied as a function of pH. From these studies the emitting species for each compound has been determined, as well as equilibrium constants for excited-state ionization. The reasons for vast differences observed between the 2,3 derivatives and their corresponding 1,2 and 2,1 derivatives currently are being investigated. (R. M. King)

REFERENCES

1. B. J. Duffield, Ph. D. Thesis, M.I.T., 1960.
2. H. J. Keily, Ph. D. Thesis, M.I.T., 1956.
3. H. Reichardt and K. F. Bonhoeffer, Z. Physik 67, 780 (1932).

ADDRESSES

- D. M. Hercules, "Instrumental Techniques for Chemical Analysis", National Science Foundation Summer Institute for High School Teachers, Juniata College, Huntingdon, Pa., August 5, 6, 7, 1964.
- D. M. Hercules, "Analytical Luminescence Spectroscopy", Gordon Research Conference on Analytical Chemistry, New Hampton, N.H., August 25, 1964.
- D. M. Hercules, "Fundamental Luminescence Processes", American Chemical Society National Meeting, Chicago, Illinois, Sept. 1, 1964.
- D. M. Hercules, "Recent Advances in Fluorescence Analysis", Delaware Valley Section, American Chemical Society, Analytical Subsection, Wilmington, Delaware, October 14, 1964.
- D. N. Hume, "Radiofrequency Plasma Emission Spectroscopy", Olin Colloquium, New Haven, Conn., July 16, 1964.
- W. E. Ohnesorge, "Fluorescence of Metal Chelates", American Chemical Society National Meeting, Chicago, Ill., Sept. 1, 1964.
- D. K. Roe, "Electrode Reaction Kinetics: Measurements with the Galvanostatic Method", Tyco Laboratories, Waltham, Mass., October 8, 1964.

PUBLICATIONS

- N. R. Andersen and D. M. Hercules, "Fluorometric Determination of Uranium with Rhodamine B", Anal. Chem. 36, 2138 (1964).
- E. M. Barrall and L. B. Rogers, "Improved Precision of Calorimetric Measurements by Differential Thermal Analysis", Anal. Chem. 36, 1405 (1964).
- R. A. Durst, J. W. Ross, and D. N. Hume, "A Simple Electronic-Scan, Controlled-Potential Polarograph", J. Electroanal. Chem. 7, 245 (1964).
- D. W. Ellis and L. B. Rogers, "The Measurement of Ionization Constants of Electronically Excited Species from Aminonaphthols", Spectrochim. Acta 20, 1709 (1964).
- D. M. Hercules, "Chemiluminescence Resulting from Electrochemically Generated Species", Science 145, 808 (1964).

Chemistry of the Fission Elements Group

L. W. Marple and D. N. Hume, "Solubility Behavior of Lead Oxide in Perchloric Acid", J. Chem. Eng. Data 9, 432 (1964).

D. K. Roe, "Electrochemical Fractionation: Potentiostatic Chromatography and Elution Voltammetry", Anal. Chem. 36, 2371 (1964).

THESES

W. R. Matson, "The Mechanism of Galvanostatic Voltage Oscillations in Formic Acid", B. S. (Chemistry), June 1964.

P. J. Sherwood, "The Behavior of HgSe as an Indicator Electrode", B. S. (Chemistry), June 1964.

D. N. Hume
D. M. Hercules
W. E. Ohnesorge
D. K. Roe

NUCLEAR CHEMISTRY (INORGANIC) GROUP

I. Physicochemical Studies, Part I

A. The Uptake of Electrolytes by an Anion-Exchange Resin

The uptake by a strong-base polystyrene resin (Dowex 1-X8, 20-50 mesh, Baker lot 28358) in the chloride form of HCl, NaCl, BaCl₂, and LaCl₃ was measured by determining the appropriate weights, and the concentration of the equilibrium solution either by conductance or by chloride analysis by methods previously reported.

The concentration of fixed groups is about 3.8 moles per kilogram of water and it changes but little with electrolyte uptake. Since conversion to the hydroxyl form appeared to increase the few parts per thousand of weak base groups, which are apparently tertiary amines (D. A. Freeman, J. Phys. Chem., 64, 1048 (1960)), the resin was merely washed first with 2M HCl and then with conductivity water until the pH of the solution was close to the asymptotic value of 4.30.

The determinations could not be carried to very dilute solutions because the uptake appeared negative. It usually corresponded to less than $2 \times 10^{-4}m$ in the beads, but was very occasionally ten times this much. At only slightly greater concentrations, the uptake became greater than would be calculated for ideal solutions.

These systems were studied because Glueckauf (E. Glueckauf, Proc. Roy. Soc., A268, 350 (1962)) says that the slope of the graph curve of $\log \bar{m}_M$ vs $\log m_M$, should be constant and equal to $(2-Z)$, independent of the valence of the co-ion, in which $\bar{m}_M$ is the concentration of co-ion in the resin and m_M that in the equilibrium solution, and Z is the measure of physical uniformity of the resin. For ideal solutions, however, the slope would be 2 for HCl and NaCl, 3 for BaCl₂ and 4 for LaCl₃.

These slopes for our measurements are far from constant. For m between 0.1 and 1.0 they were fairly constant at 2 for HCl and NaCl, 1.8 for BaCl₂ and 2.6 for LaCl₃, corresponding to $Z=0$, 0.2 and -0.6, but at smaller concentrations the slopes are very much smaller. Any tendency to negative uptake would increase the slope in dilute solutions. We conclude that this resin is not significantly non-uniform in concentration.

A good way to express the deviations from ideality is to consider $\log \bar{\gamma}$ as a function of $\bar{m}_M$ or of the activity of the electrolyte. The latter is a steadier measure in the dilute solutions. At the higher activities $\log \gamma$ is nearly constant, at low activities it is nearly proportional to $\log a$. (W. T. Yap and G. Scatchard)

B. The Osmotic Pressure, Light Scattering, and Ultracentrifuge Equilibrium of Polyelectrolyte Solutions

The address with this title will appear in the collected addresses at the Electrochemical Society Meeting in Toronto, May 4-6 (1964).

The differential equations for the distribution of diffusible components are derived for constant pressure and composition on the side of the membrane with no diffusible component, in terms of the partial derivatives $b_{ik} = (\partial \mu_i / \partial m_k)_{T, P, m} / RT$, in which μ_i is the chemical potential of component i and m_k is the quantity of component k per unit quantity of component 1, called the solvent. In dilute solutions there is no practical advantage over the previous derivation with constant pressure and concentrations of diffusible components on the side of the membrane with diffusible component (G. Scatchard, J. Am. Chem. Soc., 68, 2315, 1946), but the derivation is simpler and esthetically more pleasing.

The expression for the osmotic pressure is compared with those in the same terms for light scattering (W. H. Stockmayer, J. Chem. Phys. 18, 58, 1950) and for ultracentrifuge equilibrium (G. Scatchard, "Ultracentrifugal Analysis in Theory and Experiment". J. W. Williams, Ed., Academic Press, New York, N.Y. 1963). The equations in terms of the b_{ik} 's are simple though exact. For a solution with n solute components there are needed n equations, each containing n b_{ik} 's, but the total number of b_{ik} 's is less than n^2 because $b_{ki} = b_{ik}$.

Because of errors in the recent literature there is detailed discussion of some matters already known to many workers. The first is the appropriate diluent in order to extrapolate to zero concentration of polyelectrolyte. For osmotic pressure it is that solution which keeps constant the chemical potential of each diffusible component as the concentration of the non-diffusible component and the pressure are changed. In light scattering the pressure is kept constant, and the optimum solution is that which keeps constant the chemical potential of each solute but one, usually the polyelectrolyte, but permits that of the solvent to vary. In the ultracentrifuge the optimum average composition is the same as for light scattering, but the concentration at each point is determined by the centrifugation.

With osmotic pressure or light scattering, the determined molecular weight is for the composition chosen by the experimenter, but the centrifuge selects its own. In a solute system with a polyelectrolyte $ax_{(z'_a/z'_x)}$ and a simple electrolyte $bx_{(z'_b/z'_x)}$, the molecular weight is given by $w_a^* + (w_x^* - w_b^*)z'_a / (z'_b + z'_x)$ in which $w_i^* = w_i - RT\rho(\partial P / \partial m_i)_{T, m}$, in which w_i is the molecular weight of species i , z'_i is the absolute value of its valence, w_i is the molecular weight, ρ is the density, and P is the osmotic pressure.

The effect of the ionization of the solvent is discussed. Except for "membrane hydrolysis" with a semipermeable membrane and hydrolysis of a salt of a weak acid or base, this effect becomes appreciable only for equivalent concentrations less than ten times the concentration of ions in the pure solvent. The proper, and usual, treatment of two component systems is to extrapolate to zero concentration in an ideal solvent which differs from the real one only in being un-ionized. In the light scattering of two component systems there is an additional effect of the independent fluctuations of the ionic concentrations, which becomes appreciable when the ionic strength is less than about 4×10^{-4} times the number of ions per mole (for the blue light and water at room temperature).

The treatment of two component systems is illustrated by the light scattering and ultra-centrifuge equilibrium of 12-tungstosilicic acid, usually called silicotungstic acid, (J. S. Johnson, K. A. Kraus, and G. Scatchard, *J. Phys. Chem.*, 64, 1967 (1960); M. J. Kronman and S. N. Timasheff, *J. Phys. Chem.*, 63, 629 (1959); M. Kerker, J. P. Kratochvily, R. H. Ottewill, and E. Matijevic, *J. Phys. Chem.*, 67, 1097 (1963)). This is a strong quadrivalent acid with molecular weight 2878, which behaves in dilute solutions (for JKS), like a Debye-Hückel electrolyte with $\kappa a = 2.5 \sqrt{I}$. (G. Scatchard)

C. The Computation of the Activity Coefficients of Ions from the Loeb, Overbeek, and Wiersema Tables

In their treatment of electrolyte solutions Debye and Hückel expanded $[\exp(z_i e \psi_k / kt) - 1]$ in a power series but dropped all but the first term, in which z_i is the valence of the i 'th ion and ψ_k is the electrical potential due to the k 'th ion and its ion atmosphere. Almost immediately there were three attempts to extend the treatment to higher terms. Bjerrum used a physico-chemical method, calculating an apparent association constant from the probability that an ion i of opposite sign of valence to the central ion, k , is less than a certain distance from the central ion in a very dilute solution. Gronwall, LaMer, and their collaborators expanded to two more terms and computed analytically. Müller computed for the complete exponential by numerical integration. Recently the availability of electronic computers has led to a revival and extension of Müller's method. (E. A. Guggenheim, *Trans. Faraday Soc.*, 56, 1152 (1960); 58, 86 (1962); A. L. Loeb, Th. G. Overbeek, and P. H. Wiersema, "The Electrical Double Layer around a Spherical Colloid Particle", the M.I.T. Press, Cambridge, Massachusetts 1961 (LOW)). Guggenheim made calculations for 2-2 electrolytes in water with the distance of nearest approach, a , about $4 \text{ \AA}$, and for 2-1 (or 1-2) electrolytes with a about $4 \text{ \AA}$ and about $3.5 \text{ \AA}$, for dilute solutions ($\kappa a \leq 0.4$). His program was not powerful enough to compute for the corresponding 3-3 electrolytes. The LOW computations are obviously for another purpose; they extend from $\kappa a = 0.1$ to very large values and over a very large range of the ratio of the square of the charge to the dielectric constant. There are over three hundred pages of tables. Scatchard (G. Scatchard, "Annual Review of Physical Chemistry", 14, 161, 1963) noted that they may be used for small ions and may be extended to more dilute solutions by combination with the Kirkwood expression for the limiting value at extremely low concentrations.

This paper shows how the LOW tables may be used and gives examples of their application. The comparison with the computations of Guggenheim shows practically perfect agreement, which is especially gratifying for the empirical interpolation from the Kirkwood limit to the LOW tables.

A recent combination and extension of the methods of Bjerrum and of Gronwall - La Mer by Fuoss and Onsager (R. M. Fuoss and L. Onsager, *Proc. Nat. Acad. Sciences, U.S.A.*, 47, 818 (1961)) is discussed and shown to contain numerical errors.

The uncertainty due to the failure to fulfill the conditions of integrability is discussed briefly and the opinion expressed that the most probable values are those of Mayer (J. E. Mayer, *J. Chem. Phys.* 18, 1426 (1950)). (G. Scatchard)

II. Physicochemical Studies, Part II

A. Relative Affinity of Simple Anions for Amine Salts

The order of preference of anions in an anionic exchange process between an aqueous phase and organic solutions of tertiary ammonium salt has been determined. The values of the equilibrium constant for the exchange reaction

(where the bar indicates the organic phase) have been obtained by distribution measurements with tracers. The equilibrium constant for the equilibrium (1) is

$$K = \frac{\overline{a}_{R_3NHY} \cdot a_X}{\overline{a}_{R_3NHX} \cdot a_Y} \quad (2)$$

Introducing the activity coefficients $\bar{\gamma}$ for the organic phase and the mean activity coefficient $\gamma_{MX}^{\pm}$ for the aqueous phase we get

$$K = \frac{(R_3NHY)(X^-)}{(R_3NHX)(Y^-)} \cdot \frac{\bar{\gamma}_{RY}}{\bar{\gamma}_{RX}} \cdot \frac{\gamma_{MX}^{\pm 2}}{\gamma_{MY}^{\pm 2}} \quad (3)$$

Working at constant ionic strength we can include the aqueous activity coefficients in the equilibrium constant, $K' = K(\gamma_{MX}^{\pm 2}/\gamma_{MY}^{\pm 2})$, and compare the K' for different anions between them. In order to study the influence of the organic activity coefficient ratio, Γ_{org} , on the equilibrium constant, the values of K' are plotted against the organic concentration of the amine salt in Figures 2.1, 2.2, and 2.3. For the exchange of anions of approximately the same size, like the pairs $ReO_4^- - ClO_4^-$, $-Br^-$, $-Cl^-$, K is independent of the amine salt concentrations. The value of Γ_{org} is in this case very close to unity (within the experimental errors in K). The values of K' give the following trend: $ReO_4^- > ClO_4^- > Br^- > NO_3^- > Cl^-$. Of course the values of K' for other pairs of different anion size are those obtained by extrapolation at low amine concentration where Γ_{org} is expected to be close to unity.

Figure 2.1

K' vs organic concentration for 3/12 ammonium- NO_3^- -aqueous ReO_4^- . Temperature=298° K. Aqueous (ReO_4^-) = 10^{-3} M.

Figure 2.2

K' vs organic concentration for 3/12 ammonium Cl^- - aqueous ReO_4^- . Temperature = 298°K . Aqueous $(\text{ReO}_4^-) = 10^{-3} \text{ M}$.

Figure 2.3

K' vs organic concentration for 3/12 ammonium Br^- - aqueous ReO_4^- . Temperature = 298°K . Aqueous $(\text{ReO}_4^-) = 10^{-3} \text{ M}$.

(R. A. Nathan, A. S. Kertes, G. Scibona, and J. W. Irvine, Jr.)

B. Infrared and NMR Studies of Amine Hydrochloride in Organic Solvents

The HCl and HBr salts of triheptyl-, trioctyl-, tridecyl-, and tridodecylamine and the HI salt of tridodecylamine were prepared and their infrared spectra recorded as mulls in Nujol or in the conjugate amine. Hydrogen bonding in these salts was shown to exist by the large shift of the N-H stretching frequency from its "normal" position near 3450 cm^{-1} . The IR spectra of the four HCl salts were similar to each other. The IR spectra of the HBr salts were found to be similar to each other but different from the spectra of the HCl salts. The $\text{N}^+ - \text{H} \cdots \text{X}^-$ stretching frequencies in these long chain tertiary amine salts were assigned as: chloride - 2425 cm^{-1} , bromide - 2544 cm^{-1} , iodide - 2662 cm^{-1} , resulting in a linear relationship between the $\text{N}^+ - \text{H} \cdots \text{X}^-$ stretching frequencies and either the ionic radii or the proton affinity of the halide ions. The chain length of these amine salts was shown not to have an effect on the hydrogen bond strength of the amine salt, contrary to the effect reported for the low molecular weight tertiary amine salts.

From studies of the effect of concentration of the anhydrous tridodecylamine (TLA) salts in CCl_4 on the $\text{N}^+ - \text{H} \cdots \text{X}^-$ stretching frequency, it was proposed that a change in aggregation becomes important at approximately 0.06 M TLAHBr, probably between a monomer and a relatively large aggregate. TLAHCl and TLAHI in CCl_4 were shown to probably have a continuous change in aggregation over the concentration range 10^{-3} to 10^{-1} M , with a monomer-dimer equilibrium, with neither species predominating. From absorbance measurements of TLAHCl and TLAHBr in CCl_4 , equilibrated with the corresponding aqueous acid, the acid affinity constants were calculated. Values of 1.5×10^4 and 2.1×10^3 were found for 0.1 M and 0.01 M TLAHCl, with values of 1.5×10^5 and 2.1×10^4 for 0.1 M and 0.01 M TLAHBr.

Spectral evidence was found for the formation of the hydrogen dichloride ion, HCl_2^- , when TLAHCl was equilibrated with high concentrations of aqueous HCl. Absorption bands characteristic of HCl_2^- and of the $\text{N}^+ - \text{H} \cdots \text{HCl}_2^-$ stretching frequency were observed, and the corresponding decrease in the absorbance of the $\text{N}^+ - \text{H} \cdots \text{Cl}^-$ stretching frequency was used to calculate the equilibrium constant for the reaction: $\text{TLAHCl}_0 + \text{H}^+ + \text{Cl}^- \rightleftharpoons \text{TLA} \cdot 2\text{HCl}_0$. Values of 2.4×10^{-4} and 4.5×10^{-4} for 0.1 M and 0.01 M TLAHCl were obtained.

No spectral evidence was found for the formation of HCl_2^- when TLAHCl in CCl_4 was equilibrated with 0.1 M HCl plus LiCl up to 10 M .

No spectral evidence was found for HBr_2^- formation with TLAHBr in CCl_4 either at high aqueous HBr concentrations of 0.1 M HBr were high LiBr concentrations.

Proton chemical shifts of tridodecyl amine hydrochloride salts in carbon tetrachloride solutions have been measured with a Varian A-60 Analytical NMR Spectrometer at a field sweep width of 1000 cps. The spectra of the dry solutions prepared by dissolving the solid anhydrous salt in the solvent are in general agreement with similar systems published in the literature (Keder, Wilson, and Burger, Amine System in Solvent Extraction, in Symposium on Aqueous Reprocessing Chemistry of Irradiated Fuels, Brussels, Belgium, April 1963). From the spectra of wet organic solutions prepared by equilibrating various concentrations of amine salts with

10 M LiCl, 0.01 M HCl solutions, the following observations have been made. The position of the N-H peak shifts toward high field with increasing amine salt concentration; its intensity varies with concentration rising from 0.09 M to a maximum at 0.29 M and falling to background at 0.40 M. The α -methyl peak of the amine salt solutions shifts down field with respect to that of the free amine solution. At 0.25 M concentration a sharp peak appears between the α -methyl and methylene peaks ($\sim 3-4$ ppm) and its intensity increases with increasing amine salt concentration. The low-field shoulder of the methylene peak changes shape with increasing concentration of amine salt. These preliminary results do not allow detailed interpretation but seem consistent with the picture of aggregation equilibria in the organic phase as discussed in Section C. (C. D. Strehlow, J. E. Canty, A. S. Kertes, G. Scibona, and J. W. Irvine, Jr.)

C. Extraction of Gallium from Chloride Solutions by Quaternary Ammonium Salts in Organic Solvent

In a previous report (Nathan, Kertes, and Irvine, LNS Progress Report, November 1963, p. 29) the results of the extraction of Ga(III) at tracer concentrations from aqueous chloride solutions by tetraheptyl ammonium chloride were described. A non-integral slope over at least two orders of magnitude of amine concentration for the plots $\log E_n^0 - \log (\text{amine salt molarity})$ was obtained. The hypothesis of an aggregation of the organic phase was advanced to account for the deviation from the integral slope. More experiments over a wide range of amine concentration from 10^{-8} to 10^{-2} M have been carried out at different LiCl concentrations. The reproducibility of the data was very poor at concentrations lower than 10^{-6} M. In Figure 2.4 are shown the experimental results. A non-integral slope ranging from 1 towards the value of 2 is obtained and it seems to be a function of the aqueous LiCl concentration. In order to attempt a quantitative evaluation of the data a model, in terms of monomer and dimer equilibrium of the amine salt, has been set up, and the experimental results for 4, 5, 6, and 7 M LiCl fit this hypothesis. In the light of the results we have concluded that physical chemistry investigations on the structure of the organic phase in equilibrium with the aqueous phase must be available before an adequate interpretation of the extraction data can be made.

Figure 2.4

(S. Basol, A. S. Kertes, G. Scibona, and J. W. Irvine, Jr.)

D. Aggregation of Tertiary Ammonium Salts in Organic Solvents

The determination of the number-average molecular weight, $\overline{M}_n$, of the benzene solutions of tri-n-dodecyl amine chloride and perchlorate has been undertaken with the aim of calculating the average aggregation number of the solutions, $\bar{n} = \overline{M}_n / M_n$, where M_n is the molecular weight of the monomer. An osmometer (Microlab Model 301) has been used and the obtained values of $\bar{n}$ vs the concentrations of the ammonium salts are shown in Figure 2.5. A comparison of our values of $\bar{n}$ is made in Figure 2.5 with those obtained for the chloride salt by Erik Høgfældt (Trans. Royal Inst. Technol., No. 228, 1964) by using an e.m.f. biphasic titration method. The differences obtained in these two sets of measurements may be due to the effect of the aqueous phase on the aggregation equilibrium in the organic phase, to the difference in the solvent, or to the lack of correspondence between the experimental observation and interpretation in either or both methods of measurement. For example, presence in our solutions of surface active agents may affect the measurements. For the aqueous phase it has been shown that the effect of the presence of surface films is negligible (Brady, Huff, and McBain, J. Phys. and Colloid Chem. 55, 304, 1951). In addition the measurements above 0.1 M amine salt may be affected by the lack of precision since 0.1 M amine salt is the upper limit for the optimum precision of our instrument. Furthermore, to interpret the deviation of the electrical resistance measurements, ΔR , from the calibration curves in terms of aggregation, two assumptions must be made: a) the substances used in the calibration system are quite ideal; b) the deviation from the calibration is due only to

the solute-solute interaction and no solute-solvent interactions are present. These latter assumptions are involved in the E.M.F. measurements, too.

Figure 2.5

Average aggregation number vs concentration of the amine salt. Dashed line is calculated from e.m.f. measurements of E. Högfeldt for $Z=1$. Solid line is from the measurement of vapor-pressure lowering with matched thermistors.

(S. Basol, G. Scibona, and J. W. Irvine, Jr.)

E. Dielectric Constant Studies of Amine Salt Solutions

In the effort to substantiate the idea of the aggregation of the amine salt in the organic phase, reported in Section C, the dependence of the dielectric constant (at 1 Mc) on the tri-*n*-dodecyl amine hydrochloride molar fraction was measured. The measurements were made on benzene and carbon tetrachloride solutions of the salt. The molar polarizations of the solution calculated by the method of P. Debye (Polar Molecules, Dover, New York, 1945) and G. P. Smith (Dielectric Behavior and Structure, McGraw-Hill, New York, 1955) are plotted in Figure 2.6. The decrease of the polarization as a consequence of anti-parallel dipole interaction may be expected. The rather complicated dependence of the molar polarization on the concentration of the amine salt in benzene is not easy to explain. Difficulties are of course included in the evaluation of end-on and anti-parallel fractions of solute that contribute to the total molar polarization of the solutions. At this moment it is only possible to state that the change of dipole moment of the solutions as a function of the amine salt molarity supports the idea of an aggregation process in the phase.

Figure 2.6

Molar polarization vs molar fraction of the amine salt.

(J. Chang, H. Nakahara, A. S. Kertes, G. Scibona, and J. W. Irvine, Jr.)

F. Effect of Temperature on Extraction by Amines

In a previous report (Kimura, Kertes, and Irvine, LNS Progress Report, May 1964, p. 16) the apparent enthalpy changes for the over-all process of extraction of tracer amounts of various anionic species by tri-n-dodecyl amine hydrochloride in carbon tetrachloride have been reported. A dependence of ΔH_{E_a} for the Cl^- - Br^- exchange on the aqueous composition was observed. As expected the change of ΔH_{E_a} at low HCl molarity (below 0.01 M) is due to the hydrolysis effect of the amine salt. When the chloride and hydrogen ion concentration in the aqueous LiCl -HCl solutions is such as to make negligible the hydrolysis of the amine salt in the range of temperature studied, the chloride concentration does not affect the ΔH_{E_a} values. Also, the effect of the concentration of the amine salt on ΔH_{E_a} for the Cl^- - Br^- exchange at fixed aqueous composition has been investigated. The amine concentration in the range 10^{-4} - 10^{-2} M does not affect the ΔH_{E_a} values of the Cl - Br exchange either. (A. Toth, A. S. Kertes, K. Kimura, G. Scibona, and J. W. Irvine, Jr.)

G. Extraction Studies of Tritiated Amine Salts

By the reduction of the double bond in di-dodecyl mono-dodecenyl amine hydrochloride with tritium, a high specific activity, tri-dodecyl amine hydrochloride was prepared (preparation made by New England Nuclear Corporation). Efforts to study the distribution of this salt between carbon tetrachloride or toluene and various aqueous systems as a function of R_3NHCl concentration have not been satisfactory. Adsorption of the amine salt on the equipment has resulted in very poor reproducibility of data. Moreover, as each labelled molecule contains two H^3 atoms, the decay of one breaks C-C bonds and leaves a labelled impurity which further obscures results. It is probable that this route to the study of the characteristics of amine salt systems is not feasible. (J. F. Byrum and J. W. Irvine, Jr.)

H. Solvent Extraction of Salts of $B_{10}X_{10}^-$ Acids

Cesium at trace concentration has a finite extraction coefficient into nitrobenzene when the anion is from one of the boron cage acids, $B_{10}X_{10}^-$. The extraction coefficients increase in the order of increasing molecular weight of the anion, i.e.:

At ~ 0.1 M acid concentration the extraction coefficients for the halogen derivatives are in the range of 1 to 100. The extraction is not sensitive to the pH of the solution.

Extraction coefficients for tracer cesium as a function of acid concentration for the bromo acids tends to flatten off at $\sim 10^{-4}$ M and then show a second rise at $\sim 10^{-2}$ M acid. Two possible explanations are the presence of a hydrolysis product such as $H_2B_{10}Br_9OH$ or an increased extraction of the free acid. I-R spectra of salts of the bromo acids show OH absorption peaks but assays of the acids have not given a conclusive answer as yet. Measurements of the extraction of the free acids are also under way. (C. Schlaikjer and J. W. Irvine, Jr.)

I. Solvent Extraction of Phosphoric Acid

Dodson, Forney, and Swift (J. Am. Chem. Soc., 58, 2573, 1963) observed that phosphate extracted into isopropyl ether in the presence of Fe(III) and HCl. This observation has been confirmed and an effort has been made to determine what happens in such systems. It has been established that the species extracted is not a mixed $H_2PO_4^- - Cl^-$ complex of iron as the visible absorption spectrum of the ether solution remains constant over a wide range of H_3PO_4 concentrations. The P/Fe ratio in the ether varies from 10^{-2} to 5 over a H_3PO_4 concentration range of 0.03 M to 10 M. At high concentrations of H_3PO_4 , this acid extracts appreciably even in the absence of HCl and Fe(III). The basicity of the solvent appears to play a role in this extraction as does the availability of "free" water in the aqueous phase of the concentrated electrolytes. (R. H. McCorkell and J. W. Irvine, Jr.)

J. Study of Divalent Metal Perchlorates in Concentrated Aqueous Solution by Means of the Ion Activity Function

Previous studies have shown the promise of the ion activity function for the evaluation of the behavior of electrolytes in concentrated aqueous solutions (L. Leifer and E. Høgfældt, LNS Progress Reports, Nov. 1, 1961, p. 14; Nov. 1, 1962, p. 9, and May 1, 1963, p. 26; Høgfældt and Leifer, Acta Chem. Scand. 17, 338, 1963). The experimental Hammett acidity function H_0 provides a measure of the activity a_{H^+} of unhydrated protons in concentrated solutions of acids, based on indicator studies of the conversion of neutral base B to acid BH^+ .

$$-H_0 = \log a_{H^+} + \log (y_B / y_{BH^+}) = \log \phi_0 a_{H^+} \quad (1)$$

where ϕ_0 is the activity coefficient ratio y_B / y_{BH^+} on a molar scale, an undetermined function of the concentration of the acid, but independent of the specific indicator. It is found that the H_0 values of various strong acids are the same at a given water activity a_{H_2O} , indicating that ϕ_0 as well as the hydration equilibria of the proton are the same at the same water activity. We postulate

that the analogous function for the anion A^- will have the same generality for solutions of acid or salt at given water activity, as has been confirmed with Cl^- for HCl and $LiCl$ and Br^- for HBr and $LiBr$ (*loc. cit.*). This ion activity function $\varphi_0^{-1}a_{A^-}$ is determined from the H_0 for the acid of molarity M_{HA} and the mean molar activity coefficients $y_{\pm}$ of the acid HA as

$$\log \varphi_0^{-1}a_{A^-} = H_0(M_{HA}) + \log M_{HA}^2 y_{\pm}^2 = H_0(M_{HA}) + \log(a_{H^+} a_{A^-}) \quad (2)$$

We assume that the $\varphi_0^{-1}a_{A^-}$ value is the same in a salt solution CA of the same water activity, thus establishing $\varphi_0^{-1}a_{A^-}$ over the range of CA concentration for which a_{H_2O} and $y_{\pm}$ are known. Then we can establish the ion activity function for the cation $\varphi_0 a_{C^+}$ as a function of a_{H_2O} , since at given a_{H_2O} for HA and CA :

$$\log \varphi_0 a_{C^+} = \log M_{CA}^2 y_{\pm}^2 - \log \varphi_0^{-1}a_{A^-} = \log M_{CA}^2 y_{\pm}^2 - H_0 - \log(a_{H^+} a_{A^-}) \quad (3)$$

Further, it appears that $\log \varphi_0 a_{C^+}$ is to a fair approximation a property of the cation and its solvent interactions, independent of the anion.

The ion activity function for a hydrated ion $D(H_2O)_n^{\pm z}$ is related to the function for the bare ion $D^{\pm z}$ simply as:

$$\log \varphi_0^{\pm z} a_{D(H_2O)_n^{\pm z}} = \log \varphi_0^{\pm z} a_{D^{\pm z}} + n \log a_{H_2O} \quad (4)$$

Such functions permit the exploration of hydration equilibria.

From ion activity functions there can easily be derived ionic activity coefficient functions $\varphi_0^{\pm} y_{D^{\pm z}}$, since

$$\log \varphi_0^{\pm z} y_{D^{\pm z}} = \log \varphi_0^{\pm z} a_{D^{\pm z}} - \log M_{D^{\pm z}} \quad (5)$$

where M is the molarity of the ion.

The ionic activity coefficient functions over a range of water activities can be used to approximate the average hydration number $\bar{n}''$ of the ion by the equation:

$$\bar{n}'' = \frac{-d \log \varphi_0^{\pm z} y_{\pm}}{d \log a_{H_2O}} \quad (6)$$

as shown by Høgfjeldt (*Act. Chem. Scand.* **14**, 1627, 1960).

These methods have been applied to the two-one salts MA_2 of the ions Mg^{2+} , Ca^{2+} , Sr^{2+} , Ba^{2+} , and Pb^{2+} as perchlorates, using H_0 from $HClO_4$, and data on $y_{\pm}$ and a_{H_2O} in all solutions considered, with obvious adjustments in equations 3-5 above for cation charge $z=2$. The values for the logarithm of the ion activity functions $\log \varphi_0^2 a_{M^{2+}}$ rise from values near zero as $-\log a_{H_2O}$ increases from zero, showing for $-\log a_{H_2O}=0.5$ the approximate values 6.0, 5.4, 4.8, and 4.0 for Mg^{2+} , Ca^{2+} , Sr^{2+} , and Pb^{2+} . The curve for Ba^{2+} rises more slowly than that for Pb^{2+} , but data are not available for water activities below $10^{-0.2}$. The plots of $\log \varphi_0^{-1} y_{ClO_4^-}$ vs $\log a_{H_2O}$ for these ions are all parallel, with slope 1.0, the values for $Pb(ClO_4)_2$ extending to $-\log a_{H_2O}=-1$. The slopes imply that on the average there is 1.0 water of hydration for ClO_4^- in all the salt solutions, as also is found for $HClO_4$ solutions (where H^+ binds four water molecules).

The ionic activity coefficient function for M^{2+} is derived as follows from H_0 , a_{H_2O} , and $y_{\pm}(HA)$ for perchloric acid solution and a_{H_2O} , mean molarity $M_{\pm}(MA_2)$, and $y_{\pm}(MA_2)$ for the metal perchlorate solutions, considering data at a given a_{H_2O} :

$$\log \phi^2 y_{M^{2+}} = \log M_{MA_2}^3 y_{MA_2}^3 - 2[H_0 + \log M_{HA}^2 y_{HA}^2] - \log M_{MA_2} \quad (7)$$

The available data for $\log \phi^2 y_{M^{2+}}$ are plotted vs $-\log a_{H_2O}$ in Fig. 2.7. It is seen then that the hydration for a given cation (slope of the curves) decreases with increasing concentration, decreasing to about 2 for Pb^{++} when $-\log a_{H_2O}$ is 1 [5.7 moles per liter or 4.6 moles of H_2O per mole of $Pb(ClO_4)_2$]. This solution is nearly the composition of liquid $Pb \cdot (H_2O)_2^{2+}(ClO_4 \cdot H_2O)^-$, with very little unbound water. There is also a pronounced fall in hydration at given water activity in the series Mg^{2+} , Ca^{2+} , Sr^{2+} , Pb^{2+} , and Ba^{2+} . Values of the average hydration number $\bar{n}_{M^{2+}}$ at $-\log a_{H_2O} = 0.3$ are estimated from Fig. 2.7 as 8.7, 7.9, 6.3, 5.3, and 4.2 respectively.

Figure 2.7

Plot of logarithm of the ionic activity coefficient function $\phi^2 y_{M^{2+}}$ vs. logarithm of water activity a_{H_2O} for the perchlorates of Mg^{2+} , Ca^{2+} , Sr^{2+} , Ba^{2+} , and Pb^{2+} .

(L. Leifer and E. Høgfældt)

K. Study of the HCl- H_2O System by Means of the Ion Activity Function

The relations among $\phi_0 a_{H^+}$ and $\phi_0^{-1} a_{Cl^-}$, the ion activity functions (See Section J above) for H^+ and Cl^- for concentrated HCl solutions have been examined in detail for determination of the hydration of various components. If $\log(y_{\pm}^2 a_{H_2O}^n)$ is independent of $-\log a_{H_2O}$ over a significant range in concentrated solution, the condition for determining hydration of an ion by equation (6) of Section J above is justified (E. Høgfældt, Acta Chem. Scand., **14**, 1627, 1960). Various values of n were tried for HCl, and the values 4 led to the results of curve A, Fig. 2.8, where the function, called $\log y_{H(H_2O)_4^+} y_{Cl^-}$ is independent of $-\log a_{H_2O}$ above the value 0.4. Previous studies (Høgfældt and Leifer, Acta Chem. Scand., **17**, 338, 1963) have shown that the hydration of the chloride ion is zero above ~ 10 M HCl ($-\log a_{H_2O} \sim 0.5$). We can then go to equation (6), Section J, for the hydration; but for convenience we treat the product of $\phi \cdot y_{H^+}$ and $\phi_0^{-1} y_{Cl^-}$. Attention must also be given to the fact that the ionization of HCl in concentrated aqueous solutions is incomplete, going to the extent α . This has been shown by Raman spectroscopy, proton nuclear magnetic resonance, and the departure of the Hammett acidity function at given a_{H_2O} .

from that for the strongest acids H_2SO_4 and HClO_4 . A comparison of the α values is given by Høgfeldt (J. Inorg. Nucl. Chem. 17, 302, 1961). The mean molar activity coefficient for the ionized H^+Cl^- is $y'_{\pm} = y_{\pm}/\alpha$. The average gross hydration $\bar{n}'$ of H^+Cl^- is given as:

$$\bar{n}' = \frac{-d \log y'_{\text{H}^+} y'_{\text{Cl}^-}}{d \log a_{\text{H}_2\text{O}}} \quad (1)$$

Curve B in Fig. 2.8 shows that $\bar{n}' = 4.0$ for the concentrated solutions, a value which we ascribe to the proton. The principal form is thus $\text{H}(\text{H}_2\text{O})_4^+$.

Figure 2.8

Analysis of hydration in the system $\text{HCl}-\text{H}_2\text{O}$. Curve A is the dependence of the molar activity coefficient product for $\text{H}(\text{H}_2\text{O})_4^+$ and Cl^- on water activity $a_{\text{H}_2\text{O}}$. Curve B is a similar analysis for H^+ and Cl^- considering incomplete dissociation. Curve C is the analysis for undissociated HCl .

For HCl of stoichiometric concentration C_{HCl} , with dissociation α , we can define the thermodynamic ionization constant K as

$$\frac{C_{\text{HCl}}^2 \alpha^2 y'_{\text{H}^+} y'_{\text{Cl}^-}}{C_{\text{HCl}}(1-\alpha)} = K y_{\text{HCl}} \quad (2)$$

where y_{HCl} is the molar activity coefficient of undissociated HCl . Thus y_{HCl} is proportional to the left side of equation (2), which is known experimentally.

It can be shown, analogous to equation (6) of Section J above, that $\bar{n}_{\text{HCl(und)}}$, the average hydration number of undissociated HCl is given by:

$$\bar{n}_{\text{HCl(und)}} = \frac{-d \log K y_{\text{HCl}}}{d \log a_{\text{H}_2\text{O}}} \quad (3)$$

This may be clarified by recognizing that the ion activity function $\phi_0^{\pm z} a_D^{\pm z}$ and the ionic activity coefficient function $\phi_0^{\pm z} y_D^{\pm z}$ reduce to a_D and y_D respectively for a species D of charge 0, since $\phi_0^0 = 1$.

Curve C of Fig. 2.8 shows the plot of $K_{y_{HCl}}$ vs $-\log a_{H_2O}$. Its slope is constant above $-\log a_{H_2O} = 0.3$, in the region where $\bar{n}_{H^+Cl^-}$ is 4.0 (or $\bar{n}_{H^+}$ is 4.0), and has the value of 3.0, corresponding to an average of 3.0 water molecules bonded to unionized HCl. The ionization equation in strong HCl solutions is thus seen to be:

(L. Leifer and E. Høgfældt)

L. Detailed Hydration Behavior of Li^+ and Cl^- in Concentrated Solutions from the Ion Activity Function

Following the principles of the ion activity function of ionic species $D^{\pm z}$ in concentrated aqueous media, defined as $\varphi_0^{\pm z} a_{D^{\pm z}}$ (see Section II.J above), the ionic activity coefficient function $\varphi_0^{\pm z} y_{D^{\pm z}}$, and the determination of approximate hydration numbers $\bar{n}_{D^{\pm z}}$, we address ourselves to the data for concentrated solutions of LiCl and LiBr.

In the case of the LiCl- H_2O system we began with the generalized expression for the anionic hydration:

We can then write the following equation in which brackets indicate concentration:

$$\frac{[Cl(H_2O)_m^-]}{[Cl^-] y_{Cl^-}} = K_m a_{H_2O}^m \quad (2)$$

In equation (2) the molar activity coefficients of the hydrated species are absorbed into the constant, K_m . Taking the logarithm of equation (2) we have

$$\log \frac{[Cl(H_2O)_m^-]}{\{Cl^-\}} = \log K_m + m \log a_{H_2O} \quad (3)$$

We show that $-d \log \Sigma ([Cl(H_2O)_m^-] / \{Cl^-\}) / d \log a_{H_2O} = \bar{m}_{Cl^-}$, the average degree of hydration of the chloride ion. We write the generalized hydration expression in the following form:

$$\frac{\sum_0^m [Cl(H_2O)_m^-]}{\{Cl^-\}} = \sum_0^m K_m a_{H_2O}^m \quad (4)$$

Differentiating equation (4) with respect to the water activity and then multiplying through by a_{H_2O} we obtain:

$$\frac{d \left(\frac{\sum_0^m [Cl(H_2O)_m^-]}{\{Cl^-\}} \right)}{d \log a_{H_2O}} = \sum_0^m m K_m a_{H_2O}^m \quad (5)$$

Dividing (5) by (4) yields the average value of the hydration number, $\bar{m}_{Cl^-}$

$$\frac{d \log \left(\sum_0^m [\text{Cl}(\text{H}_2\text{O})_m] / \{\text{Cl}^-\} \right)}{d \log a_{\text{H}_2\text{O}}} = \frac{\sum m K_m a_{\text{H}_2\text{O}}}{\sum K_m a_{\text{H}_2\text{O}}} \equiv \bar{m}_{\text{Cl}^-} \quad (6)$$

It has been shown previously (Høgfældt, Acta Chem. Scand. 14, 1627, 1960) that

$$\frac{-d \log \varphi_0^{-1} y_{\text{Cl}^-}}{d \log a_{\text{H}_2\text{O}}} = \bar{m}_{\text{Cl}^-} \quad (7)$$

Thus we identify $(\varphi_0^{-1} y_{\text{Cl}^-})^{-1}$ with the summation on the left side of Equation (4) or (6). If over a substantial region of $a_{\text{H}_2\text{O}}$, $\log \varphi_0^{-1} y_{\text{Cl}^-}$ is linear with $-\log a_{\text{H}_2\text{O}}$, with slope n , this is taken as evidence that only one term of the series, the n th one, is important, and we have the opportunity to use Equation (2) to evaluate K_n for the concentration region used. In the case of the LiCl-H₂O system it has been established (Høgfældt and Leifer, Acta Chem. Scand. 17, 338, 1963) that for the region $0.5 < -\log a_{\text{H}_2\text{O}} < 0.7$, $n=0$ and $\log \varphi_0^{-1} y_{\text{Cl}^-} \equiv 0.19$, so that

$$-\log \varphi_0^{-1} y_{\text{Cl}^-} = \log K_0 = -0.19 \quad (K_0=0.65) \quad (8)$$

But the ion activity function, $\varphi_0^{-1} a_{\text{Cl}^-} = \varphi_0^{-1} \{\text{Cl}^-\}$ is also known as a function of M_{LiCl} or $a_{\text{H}_2\text{O}}$, which permits us to determine the free chloride ion concentration $[\text{Cl}^-]_{\text{free}}$

$$\log \varphi^{-1} a_{\text{Cl}^-} - 0.19 = \log [\text{Cl}^-]_{\text{free}} \quad (9)$$

$$[\text{Cl}^-]_{\text{total}} - [\text{Cl}^-]_{\text{free}} = [\text{Cl}^-]_{\text{hydrated}} = \Delta \quad (10)$$

Assuming that Equation (8) has validity for the $n=0$ (anhydrous) form over a wider range, the ionic activity coefficient function for the remaining terms of the series, corrected for contribution of this constituent is:

$$\log \varphi_0^{-1} y_{\text{Cl}(\text{corr})} = \log \varphi_0^{-1} a_{\text{Cl}^-} - \log \Delta \quad (11)$$

Thus we can go through a series of "curve peelings" to evaluate successive important members of the summation on the left side of Equation (4). The next term is expected to be given for an adjacent region of $-\log a_{\text{H}_2\text{O}}$ where again a constant slope p is seen in $\log \varphi_0^{-1} y_{\text{Cl}(\text{corr})}$, or the function is treated as the sum of several. In the case of integral slope p :

$$-\log \varphi_0^{-1} y_{\text{Cl}(\text{corr})} = \log K_p + p \log a_{\text{H}_2\text{O}} \quad (12)$$

This will mark the p th term as the next representative one of the summation. For the LiCl data $p=4.0$ gave good fit over the region $0.05 < -\log a_{\text{H}_2\text{O}} < 0.4$, so that

$$-\log \varphi_0^{-1} y_{\text{Cl}(\text{H}_2\text{O})_4} - 4 \log a_{\text{H}_2\text{O}} = \log K_4 = 0.34 \quad (K_4=2.2) \quad (13)$$

and hydration numbers intermediate between 0 and 4 seem to be of small importance, corresponding to very small values of K_1 , K_2 , and K_3 . It does not seem justifiable to go to lower values of $-\log a_{\text{H}_2\text{O}}$ because the assumption of constancy of K_m values becomes poorer.

The fall in the function $\log \varphi_0^{-1} a_{\text{Cl}^-}$ with $-\log a_{\text{H}_2\text{O}}$ in very concentrated solutions is ascribed to the formation of a solvated ion-pair $\text{LiCl}(\text{H}_2\text{O})_q$. By procedures analogous to those above, it is found that the data are in good accord with $q = 1.0$, $\text{LiCl}(\text{H}_2\text{O})$. This is reflected in the equation:

$$\log [\text{LiCl}(\text{H}_2\text{O})] = \log \{\text{LiCl}\} + \log a_{\text{H}_2\text{O}}^{-4.84} \quad (14)$$

Figure 2.9 shows the balance for hydrates of species involving chloride ion in LiCl solutions. The ordinate is molarity M of the different species, the abscissa is the molality (moles per kg H_2O) of LiCl, increasing with $-\log a_{H_2O}$. It is seen that the molarity of $Cl(H_2O)_4^-$ computed by Equation (13) rises to a maximum of 2.3 at ~ 5 M LiCl and then falls off. The anhydrous Cl^- is the principal species after 5 M LiCl, and rises to 11.5, where it levels off. At very high abscissas, $LiCl(H_2O)$ begins to rise. These three individual curves sum to values close to the envelope for stoichiometric molarity, showing the general utility of this formal treatment.

Figure 2.9
Concentrations (molarity) of various chloride species, $Cl(H_2O)_4^-$, Cl^- (anhydrous), and $Li(H_2O)Cl$ (ion pair) in LiCl solutions as a function of molality m_{LiCl} .

The hydration of Li^+ in the same solutions can be obtained in analogous fashion using the experimental functions $\varphi_0 a_{Li^+}$ and $\varphi_0 y_{Li}$. After making correction for the $LiCl(H_2O)$ in very concentrated solutions, the limiting slope of the $\varphi_0 y_{Li}(\text{corr})$ vs $-\log a_{H_2O}$ is +3.0, corresponding to $\log K_3^{Li} = -1.02$. The next term found at lower values of $-\log a_{H_2O}$ had slope 5.0, corresponding to $\log K_5^{Li} = 0.34$. We thus consider the main cation hydrate above 2 M LiCl to be $Li(H_2O)_5^+$, later dehydrated to $Li(H_2O)_3^+$. Further extraction of water leads to formation of the solvated ion-pair $LiCl(H_2O)$. Its formation in concentrated solution is

Fig. 2.10 shows the balance for hydrates of species involving lithium ion in LiCl solutions. Here, analogous to the chloride data in Fig. 2.9, molarity of the species is plotted against the stoichiometric molality. The species $Li(H_2O)_5^+$ is the principal one up to ~ 7 M LiCl, after which $Li(H_2O)_3^+$ predominates. At 14 M LiCl (20 m), $LiCl(H_2O)$ represents about 18% of the total Li^+ .

Figure 2.10
Concentrations (molarity) of various lithium species $\text{Li}(\text{H}_2\text{O})_5^+$, $\text{Li}(\text{H}_2\text{O})_3^+$, and $\text{Li}(\text{H}_2\text{O})\text{Cl}$ (ion pairs) in LiCl solutions as a function of molality m_{LiCl} .

This picture of the solvation properties of ions is quite definite, and thus subject to check by measurements of the other properties of concentrated solutions. Efforts of this type are in progress. (L. Leifer and E. Høgfældt)

III. Equipment

Pneumatic Tube System and Fast-Unloading Station

The design, construction, and installation of equipment suitable for the rapid handling of irradiated solutions have been completed. The system minimizes the time between the production and subsequent radiochemical processing of material irradiated in the MIT reactor. An extension of the pneumatic-tube system of the reactor enables samples to be transmitted to the nuclear chemistry laboratory in the adjoining building. The extension tube is not connected directly to the reactor sample-transfer tube and the rabbit containing the irradiated solution must pass clear of the reactor-system tube by jumping an open gap of about 6 inches before entering the extension. A separate centrifugal pneumatic pump, located in the chemistry building draws the rabbit through a tube which penetrates the reactor containment shell and passes underground beneath the reactor building to the chemistry building. A micro-switch signals the approach of the rabbit to the end of the tube where a solenoid valve causes suction on the tube to be broken, allowing the rabbit to exit via a hinged door into the receiver station. The total elapsed time for ejection from the irradiation position in the reactor to arrival at the receiver station is approximately 5 seconds. The transfer of the irradiated solution from the receiver station into a chemical processing system is accomplished by piercing the irradiation container with a needle assembly and drawing the liquid into an evacuated system. The handling times required for this technique are as short as 2 seconds and are achieved with minimal manipulation. This system for rapid handling of irradiated liquids

has general applicability and has incorporated in the design numerous safety features, reducing likelihood of radiation accidents, yet permitting the solution transfer to be made easily and rapidly.

In order to provide for safe handling and containment of medium-level radioactivity in the laboratory containing the receiver station, a number of radiological-safety features have been added. A small hood, which exhausts into the room's fume-hood system has been placed around the receiver station. The source of vacuum for transfer of the irradiated solution from the rabbit and for subsequent handling of the sample is a 5 ft³ evacuated epoxy-lined tank. Short-lived, gaseous activities that are produced during the irradiation and released during sample transfer and chemical operations are stored for several hours in the tank before being pumped out and exhausted into the fume-hood system. A radiation monitor has been placed in a gas-sampling line attached to the main fume-hood exhaust system. A change room, containing a foot-operated sink, radiation monitors, and a rack for laboratory coats has been installed at the laboratory entrance.

Examples of experiments involving the use of the system are given in Sections IV.A and IV.B. A manuscript describing the system, "Design of Equipment for Rapid Handling of Irradiated Solutions", by C. E. Bemis, Jr., and J. W. Irvine, Jr., has been accepted for publication in Nuclear Instruments and Methods. (C. E. Bemis, Jr., J. W. Irvine, Jr., and D. A. Muga)

IV. Nuclear Decay Schemes

A. Search for Fission-Product Antimony Delayed-Neutron Precursors

Using the pneumatic-tube system described in Section III possible delayed-neutron emission following the decay of 2-sec Sb¹³⁵ has been investigated further (see C. E. Bemis, G. E. Gordon, and C. D. Coryell, J. Inorg. Nucl. Chem. 26, 213, 1964). Solutions of enriched U²³⁵ have been irradiated for short periods, and the solution transferred to separation apparatus for the chemical separation of antimony from the gross fission products. The antimony separation was performed by reduction of antimony to SbH₃ gas with zinc metal in HCl solution and passage of the SbH₃ through a trap, three 0.5 M NaOH traps, and a drying tube consisting of CaSO₄ and Ascarite before the AgNO₃ collection tube. In most of the experiments the latter was placed inside of the neutron counting system. Using this rapid antimony chemistry, it has been possible to start counting as soon as 12 seconds after irradiation. The counting system, having an efficiency of 7.2%, consisted of eight BF₃ tubes placed in circular array in a paraffin moderator. The rapid decay was followed by multiscaling with 256- and 400-ch. pulse-height analyzers.

A typical delayed-neutron decay curve from a fission-product antimony sample is shown in Figure 2.11. The resolution of the decay curves gave the following component half lives: 55 sec (Br⁸⁷), 23 sec (Br⁸⁸ and I¹³⁷), and 7 sec (Br⁸⁹ and I¹³⁸). These components arise from halogen contamination in the chemical separation and from the products remaining in the distilling

flask, whose neutrons are counted with a small, but finite, efficiency. The upper limit on the amount of a 2-sec delayed-neutron component that could be admitted in the decay curve of Fig. 2.11 has allowed us to set a conservative upper limit of 1% on the neutron branching in the decay of Sb^{135} .

Figure 2.11

Delayed-neutron decay curve observed from a fission-product antimony sample. Counting began 12 sec after a 10-sec irradiation of 5 mg U^{235} .

Because of the inefficiency of the bubble traps in scavenging out the halogen gases (iodine and bromine) in the chemical separation scheme, we are investigating a new procedure which eliminates these traps and consists of the following: Addition of iodine carrier and 5 ml HBr directly to the reaction flask, reduction of antimony to SbH_3 gas with zinc metal, drawing the gas stream through a 41-mm dia. 2-ft drying tube of CaSO_4 and Ascarite, and precipitation of antimony with AgNO_3 collection tube within the neutron counter. Continued efforts in the resolution of the gross fission decay curves of antimony samples using the new chemical operation scheme are being pursued at the present time. (C. E. Bemis, D. A. Muga, G. E. Gordon, and C. D. Coryell)

B. Decay Schemes of Sb^{130} and Sb^{131}

We have prepared fission-product Sb samples highly enriched in 37-min Sb^{130} and 23-min Sb^{131} by making a fast Sb separation from fission products immediately after irradiation of U^{235} with thermal neutrons (for chemical procedures, see Sects. IV.A and IV.D), to reduce the activities of Sb isotopes having long-lived Sn parents. The sample is allowed to stand for about 20 min before a second separation of Sb is made to remove the daughter products of very short-lived Sb products. The activities remaining are essentially those of Sb^{130} and Sb^{131} , with the 25-min Te^{131} daughter of the latter subsequently growing into the sample.

The γ -ray spectrum of an Sb^{130} - Sb^{131} - Te^{131} sample as obtained with a 3-x3-in. NaI detector is shown in Fig. 2.12. Contributions from Te^{131} have been removed as indicated by means of pure Te^{131} spectra obtained by Te^{130} (n, γ) (see Walters, Bemis, Gordon, and Ragaini, Sect. IV.C).

Individual γ -ray peaks were obtained by "stripping" the spectrum by the use of response functions constructed from the spectra of γ -ray standards taken under the same conditions. We attempted to assign the γ -rays to Sb^{130} or Sb^{131} by determining relative intensities of the γ -rays in spectra taken at successive time intervals, but uncertainties in the stripping process and the similarity of the half lives made it impossible. Most of our assignments were based on β - γ , γ - γ , and sum-coincidence experiments, aided by previous studies of Sb^{130} at times long compared with the Sb^{131} half life (B. J. Dropesky and C. J. Orth, J. Inorg. Nucl. Chem. 24, 130, 1962; G. Lange, Ph.D. thesis, Univ. of Mainz, 1963). We have also studied the γ rays by observation with a Li-drifted Ge detector (see Fig. 2.13). In general, the correspondence between spectra obtained with NaI and Li-drifted Ge is good for peaks of large magnitude.

Figure 2.12

Gamma-ray spectrum of 37-min Sb^{130g} and 23-min Sb^{131} analyzed for intensities of individual γ rays. Contribution from 25-min Te^{131g} (shown as the dashed curve) has been subtracted.

Figure 2.13

Gamma-ray spectrum of 37-min Sb^{130g} , 23-min Sb^{131} , and 25-min Te^{131g} obtained with a Li-Ge detector at high-gain setting.

The workers mentioned above have assigned γ rays of 186, 329, 830 (complex, i.e. more than a single line), and 930 keV to the decay of 37-min Sb^{130}g . In Figure 2.13, strong lines are seen at 807 and 849 keV, with a weaker line at 820 keV. In γ - γ and γ -sum-coincidence experiments, we have found very strong coincidences among members of the ~ 830 keV complex and between the 830 keV complex and 186- and 326-keV lines. Thus, the main feature of the Sb^{130} decay scheme indicated by our experiments is a 5-member cascade involving 186-, 326-, 807-, 820-, and 849-keV γ rays. On the basis of intensities we have placed the 807 $\rightarrow$ 849 members at the bottom of the cascade as shown in Fig. 2.14. The placement of the 849-keV line from the first excited state to the ground state is in good agreement with many inelastic-scattering and Coulomb-excitation measurements of the energy of the first excited state, 845 ± 7 keV. The order of the 186-, 326- and 820-keV members is somewhat arbitrary and has been made mainly on the basis of systematics. The long chain of coincident γ rays is probably the result of a very high spin (5 to 8 $\hbar$) of 37-min Sb^{130} . If so, however, this leads to a serious discrepancy with the (p,p') data for Te^{130} obtained by Cookson and Darcey (Univ. of Oxford, Ref. 144/64, 1964). They found second and third excited states at 1588 ± 7 and 1633 ± 7 keV, with apparent spins of 4^+ and 2^+ . The high-spin level at 1588 is far outside of the limits of error on our state at 1657 keV (about ± 10 keV), and the only other state that might be expected in the vicinity (as yet, unobserved) is the 0^+ member of the 0^+ , 2^+ , 4^+ two-phonon triplet. Clearly, more work is needed to resolve this discrepancy.

Figure 2.14

Proposed partial decay scheme of 37-min Sb^{130}g . Numbers in parentheses are approximate intensities of γ rays relative to the 849-keV line.

The decay of Sb^{131} is almost devoid of strong γ - γ coincidences, the major exceptions being a strong ~ 950 -950 keV coincidence and some indications for a 1050-950 one. The partial decay scheme shown in Fig. 2.15 includes these along with a strong 652-keV line (also seen by Cooper, Gatti, Rasmussen, and Thompson, UCRL-11255, Feb. 1964) and a number of weaker lines. Also,

in the case of Sb^{131} , much more work, preferably with the aid of an isotope separator, is needed to construct completely the decay scheme. Both partial decay schemes (Figs. 2.14 and 2.15) must be considered quite tentative.

Figure 2.15

Proposed partial decay scheme of 23-min Sb^{131} . Numbers in parentheses are approximate intensities of γ rays relative to the sum of the two ~ 950 -keV lines.

As the half life of Sb^{131} was quite important in the analysis of the isomer-yield ratio, $\text{Te}^{131\text{m}}/\text{Te}^{131}$ (see Sarantites, Coryell, and Gordon, Sec. V.B, this report), we made a careful determination of the half life. We followed the decay of 950-950-keV γ - γ coincidences from a fission-product Sb sample. The decay curve showed a single component of half-life 23 ± 1 min in good agreement with the 23-min value previously reported by Pappas (AECU-2806, 1953). No explanation is given for the somewhat lower value of 19.4 min reported by Strom (presented at Univ. of Oxford, Sept. 1962).

More detailed results of this work are given in the thesis of C. E. Bemis, Oct. 1964. An abstract covering the Sb^{130} results has been submitted for the New York meeting of the American Physical Society, January 1965. We would like to thank Dr. H. Kraner and Mr. J. Sovka for their generosity with the Li-Ge detectors. (C. E. Bemis and G. E. Gordon)

C. Decay-Scheme Studies of 25-min $\text{Te}^{131\text{g}}$

In the course of studies of the decay schemes of Sb^{130} and Sb^{131} (see Bemis and Gordon, Sect. IV.B), it became necessary to have an accurate knowledge of the decay scheme of the 25-min $\text{Te}^{131\text{g}}$ daughter of 23-min Sb^{131} . Gamma-ray spectra obtained with NaI(Tl) detectors and Te^{131} produced by enriched- $\text{Te}^{130}(n, \gamma)$ at the MIT reactor showed two γ rays that were not reported

in the recent study by Devare, Tandon, and Devare (Phys. Rev. 131, 1750, 1963), one of 727 keV and another of ~ 840 keV energy. Using a Li-drifted Ge detector, we were able to determine γ -ray energies to ± 2 keV in most cases and to confirm the suggestion by Devare et al. that there is more than one γ -ray in the 350 keV and 970 keV regions. A typical spectrum obtained with a Li-drifted Ge detector is shown in Fig.2.16. A list of the energies and relative intensities of the observed γ rays is given in Table 2.I. We have used the relative intensities of prominent lines as determined from NaI spectra to construct an efficiency-vs-energy curve for the Li-drifted Ge detector. The latter was used to determine the relative intensities of lines not clearly resolved in the spectra taken with NaI crystals.

Figure 2.16

Gamma-ray spectrum of 25-min Te^{131g} observed with a Li-drifted Ge detector. Numbers given on the peaks are energies of the lines in keV.

Table 2.I

Relative intensities of γ rays emitted in the decay of
25-min Te^{131g}

E_{γ} (keV)	Devare <u>et al.</u> ^b		This Work	
	NaI	NaI	NaI	Li-Ge
150	(100)	(100)	(100)	(100)
343	2.1	1.53		0.5
384				1.1
450	28.0	23.5		(23.5)
493	4.6	6.9		(6.9)
602	7.9	6.5		5.0
655	1.3	2.7		1.8
727	not seen	0.9		0.7
840	not seen	0.9		unclear
951	4.5	4.5		2.0
997	7.0	7.2		6.7
1008				1.0
1148	8.0	8.9		(8.9)
1296	1.2	1.1		0.6

^a Energies taken from the Li-Ge spectra

^b Devare, Tandon, and Devare, Phys. Rev. 131, 1750 (1963).

We have used γ - γ , sum-peak, and sum-coincidence techniques to establish coincidence relationships and construct the decay scheme. A typical sum-peak spectrum is shown in Fig. 2.17. Results of these experiments are listed in Table 2.II. In Figure 2.18 is shown our proposed partial level scheme for the resulting ^{131}I . The main features of our decay scheme are in good agreement with those of Devare and co-workers' proposed scheme. We have confirmed their suggestion that there are two γ rays of ~ 985 keV feeding levels at 150 and 493 keV. We find the energies of the γ rays to be 997 and 1008 keV, the latter showing up as the high point on the upper edge of the 997-keV peak. We have observed this indication repeatedly in similar spectra. Also, the intensity of the 1008-keV peak relative to that of the 997-keV one is greatly diminished in preliminary Li-drifted Ge spectra taken in coincidence with 150-keV events observed with a NaI(Tl) detector. We are now investigating the suggestion by Devare *et al.* that there are two γ rays in the vicinity of 920 to 935 keV which feed levels at 150 and 493 keV. We observe only one strong line at 951 keV (found to be coincident with 150- and 493-keV γ rays), but it is difficult to determine whether or not there are γ rays at energies slightly below 950 keV. This difficulty occurs because of the sizable Compton edge of the 1148-keV line in that area. Preliminary Li-Ge — NaI coincidence experiments suggest that there are too many 951-keV γ rays in coincidence with the 150-keV line to be explained by the single 951-keV line which we show as feeding the 493-keV level. This result indicates a 951-keV γ ray (from a level at 1101 keV) that feeds the 150-keV level directly.

Figure 2.17

Sum-peak spectrum of γ rays from 25-min Te^{131}g . Observed with two 3- x 3-in NaI(Tl) detectors and sum-coincidence electronics. Numbers assigned to the peaks indicate energies in keV.

Table 2. II
Gamma-ray coincidence relationships in decay of 25-min $\text{Te}^{131\text{g}}$

$E_{\gamma 1}$ (keV)	$E_{\gamma 2}$ (keV)													
	150	275	343	384	450	493	602	655	727	840	951	997	1148	1296
150			G, S		G, S	N	N	G	G, S	G	G, S	G, S	N	G, S
343	G, S			S		N		S			G, S		N	
384			S			G, S							N	
450	G, S												N	
493	N			G, S				G, S			G, S	G, S	N	
1148	N	G	N	N	N	N	N	N	N	N	N	N	N	

G: Coincidence observed in γ - γ coinc. expt.

S: Sum of the two γ rays observed in sum-peak spectrum

N: Coinc. not observed in γ - γ coinc. expt.

blank: Not investigated

Figure 2.18

Proposed tentative level scheme for I^{131} . Numbers in parentheses indicate intensities relative to that of the 150-keV line.

Our major disagreement with Devare *et al.* is in the placement of the 343- and 384-keV γ 's. The latter placed them much higher in the level scheme: a 350-keV line between levels at 1480 and 1130 keV; a 360-keV line between levels at 1425 and 1065. We have strong evidence for our placement of the lines: the 493- and 727-keV sum peaks, other coincidence data, and the fact that the sum of 343 and 384 is exactly 727. We have looked for coincidences with the 1148-keV γ ray, finding no indications for 350- to 360-keV γ rays. We observed only a small number of ~ 275 -keV γ rays coincident with the 1148-keV, perhaps from the decay of 1.2-day $\text{Te}^{131\text{m}}$.

We are continuing this investigation, particularly making use of the powerful technique of NaI-Li-Ge coincidence experiments. In addition to the open questions mentioned above, we will also look for confirmation of the ~ 840 -keV γ ray which is present in the NaI spectra, but not apparent in the Li-Ge spectra. We wish to thank Dr. Hobart Kraner and Mr. J. Sovka for their generous loan of Li-drifted Ge detectors. (W. Walters, C. E. Bemis, Jr., G. E. Gordon, and R. C. Ragaini)

D. Half Life and Gamma-Ray Spectrum of Te^{133g}

A method for the separation of Te daughter-products arising from the decay of Sb fission products was developed. Following short irradiations of $\text{UO}_2(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$, the Sb fission products were separated by generating SbH_3 gas (see Sect. IV. A). The SbH_3 was collected by passing the gas through a 10% NaOCl solution. The Sb was oxidized to Sb(V). Ten mg of Te carrier containing equal amounts of Te(IV) and Te(VI) was added. The solution was made 3 M in HCl by addition of conc. HCl. About 3 ml of a 15% solution of hydrazinium chloride ($\text{N}_2\text{H}_5\text{Cl} + \text{HCl}$) was added to the solution and Te reduced to the metal by passing in SO_2 gas. The metal was removed by filtration, washed twice with 3 M HCl, and mounted between layers of tape on a sample card for γ -ray counting. The Te daughter products could be counted about 3 min after irradiation, allowing a 1.5 min growth period after the initial Sb separations. The method of production of Te samples, has the advantage that Te daughters of medium-lived (a few minutes) Sb fission products are highly enriched relative to those of very short (a few sec) or long (~ 20 min or longer) Sb parents.

The γ -ray spectra of the Te daughter(s) separated from Sb are shown in Fig. 2.19. Only the first two of a series of successive spectra are shown. It was expected that the γ spectra would be essentially that of 2-min Te^{133g} (from 2.4-min Sb^{133}) as previously measured by Ferguson, Love, and Sam (J. Inorg. Nucl. Chem. 24, 1, 1962). We expected interference from other Te daughter products to be very small. No significant quantities of 78-hr Te^{132} nor 25-min Te^{131} are produced with the short Sb decay times used. Te^{134} and Te^{135} are not obtained because of the very short half lives of <1.5 -sec Sb^{134} and 2-sec Sb^{135} .

Figure 2.19

Successive γ -ray spectra of Te separated from short-lived Sb fission products. The top spectrum was recorded 1.5 min after separation, the lower one ~ 15 min later.

The γ -ray activity decayed with a 12 ± 3 -min half life. Prominent γ -rays were observed at 150, 310, 403, 715, 812, 912, 1015, 1330, 1710, and 1790 keV. As noted below, these are attributed to the decay of Te^{133g} . Gamma rays at 530 and 850 keV appeared to grow into the sample.

Recently, S. Prussin (Ph.D. thesis, Univ. of Michigan, 1964) reported the existence of a 12.4-min isomer of Te^{133} . The 12.4-min species was obtained by separating samples of Te(IV) from a stock solution of Te(VI) previously separated from fission products. The parent half-life of the 12.4-min activity was established as ~ 50 min (Te^{133m}). The assignment of the activity to Te^{133g} was based on the above information and the apparent growth of 21-hr I^{133} in the Te samples.

We conclude that the 12 ± 3 -min Te activity observed in this work is the same as that reported by Prussin, Te^{133g} . Gamma-ray spectra are similar and the 530- and ~ 850 -keV γ rays growing into the sample are characteristic of 21-hr I^{133} . The half life of the Sb parent (2.4-min for Sb^{133}) would have to be in the range of a few minutes in order to produce large amounts of 12-min Te^{133g} within a growth period of a few minutes. (C. E. Bemis, Jr.)

E. Decay Scheme of the Cd^{117} Isomers

This work is a continuation of the study of the decay of ~ 2.7 -hr $\text{Cd}^{117}(1/2+)$ and ~ 3.2 -hr $\text{Cd}^{117m}(11/2-)$ reported previously (Tang, Gordon, and Coryell, LNS Progress Report, Nov. 1963, p. 36).

1. Energies of Gamma Rays

Gamma-gamma coincidence studies have suggested the presence of a number of γ rays that were not resolved in the stripping of singles spectra taken with a 3- x 3-in NaI(Tl) scintillator. With the use of Li-drifted Ge detectors, many of these γ rays have now been resolved and definitely established. However, many more peaks suggesting the presence of other previously unknown γ 's have been observed and await further verification from coincidence work. The following 27 γ rays (in keV) attributed to the Cd^{117} isomers have been definitely established by this work: 88, 221, 274, 346, 434, 461, 567, 630, 750, 834, 862, 882, 980, 1033, 1052, 1070, 1140, 1230, 1300, 1390, 1430, 1570, 1720, 1990, 2310, 2370, and 2440. Indium-117 was milked and cleaned from Cd^{117} and its spectrum on the Li-drifted Ge detector shows only the 158-, 316-, and 552-keV peaks. Methods for obtaining relative γ ray intensities from Li-drifted Ge detector spectra are being studied. Figure 2.20 shows a γ -ray spectrum of a Cd^{117} - In^{117} source taken with the Li-drifted Ge detector.

Figure 2.20

Li-drifted Ge spectrum of Cd^{117} - In^{117} Gamma Rays

2. Half-Life Measurements

Decay curves of Cd^{117} were obtained by using the RIDL 400-channel pulse-height analyzer in the time mode and using coincidence sorting for selective counting to reduce Compton-shoulder interference from other radiations of different half lives. A 1.5-in Pilot-B plastic scintillator was used as the β -ray detector and a 3- x 3-in NaI(Tl) crystal as the γ -ray detector. The following results were obtained:

Gated Radiation	Counted Coincident Radiations	Observed Half Life
274-keV γ	1.300-MeV γ	2.5 hr
434 } 461 } 439-keV γ	$\beta > 1.0$ -MeV	2.5 hr
434 } 461 } 439-keV γ (repeat)	$\beta > 1.0$ -MeV	2.5 hr
1.033 } 1.052 } 1.054-MeV γ	$\gamma > 274$ keV	3.2 hr
1.070 }		
1390 } 1430 } 1.428-MeV γ	$\gamma > 274$ keV	3.4 hr
834 } 862 } 867-keV γ	$\gamma > 550$ keV	3.1 hr
882 }		

The 2.5-hr value obtained for Cd^{117g} is more accurate than the ~ 3.2 -hr one (Cd^{117m}) because the gated radiations are better resolved, and the coincident radiations counted were more intense.

3. Q_{β^-} of Cd^{117}

It is now possible to determine Q_{β^-} for Cd^{117} based on new mass spectrometric data for Cd^{116} and Sn^{117} , having mass excesses of -90.405 and -88.713 MeV, respectively (Damerow, Rice, and Johnson, Phys. Rev. 132, 1673, 1963), the measurement of the Q_{β^-} value for the $\text{Cd}^{116}(\text{d}, \text{p})\text{Cd}^{117}$ reaction, 3.550 ± 0.020 MeV (Silva and Gordon, Phys. Rev. 136, B618, 1964), and the previous information on Q_{β^-} for In^{117} . The resulting Q_{β^-} for Cd^{117} is 2.54 ± 0.02 MeV.

4. Coincidence Work

Fig. 2.21 shows a coincidence spectrum of Cd^{117} taken on the Li-drifted Ge detector with the analyzer gated by 1300 ± 50 -keV γ rays detected in a 3-x 3-in NaI crystal. This technique holds much promise for future coincidence work. Standard γ - γ coincidence studies using two NaI(Tl) scintillators as detectors have also been performed. Sum-coincidence techniques are also being applied in the present work.

Figure 2.21

Spectrum observed in a Li-Ge detector in coincidence with 1300 ± 50 -keV γ rays of Cd^{117} detected by a 3-x 3-in NaI(Tl) crystal.

5. Assignment of Gamma Rays

Most of the principal γ rays can now be assigned to one of the two Cd^{117} isomers by analyzing the spectra of Cd^{117} produced from the $\text{Cd}^{116}(\text{n}, \gamma)\text{Cd}^{117}$ reaction and $\text{Cd}^{116}(\text{d}, \text{p})\text{Cd}^{117}$ reaction (giving about 3-fold higher yield of $\text{Cd}^{117\text{m}}$ than in the former reaction) and by comparing spectra taken immediately after bombardment with those taken 23 hours later.

We wish to thank Dr. H. Kraner of the Radioactivity Center and Mr. J. Sovka of the Nuclear Engineering Department for their help in obtaining the Li-Ge spectra. (C. -W. Tang and C. D. Coryell)

V. Nuclear Fission

A. Kinetic Energy Release in Photofission of U^{238} and Th^{232}

Data from the semiconductor measurements of the energies of coincident fragments from photofission of U^{238} and Th^{232} have been analyzed (see earlier reports, Brenner and Cooper, LNS Progress Report, May 1964, p. 35; Nov. 1963, p. 46). The energy scales for the two detectors were determined by observation of fragments from a Cf^{252} spontaneous fission source and the use of the calibration scheme described by Kiker (Ph.D. thesis, Univ. of Tennessee; also issued as ORNL-3692, 1964). Transformation of the data obtained in the energy-energy coordinates to other coordinate systems of interest were made with the aid of a program, Kerif, a modification of the interpolation scheme developed by Thomas and Gibson (Princeton-Penn. Accelerator Report No. PPAD 473E, Oct. 1962).

In Fig. 2.22 we show the single fragment kinetic energy spectra for the Cf^{252} spontaneous-fission standard and the photofission of U^{238} and Th^{232} . In Figs. 2.23 and 2.24 we show average total kinetic energy release and yield-mass as a function of fragment mass number for photofission of U^{238} and Th^{232} . In Figs. 2.25 and 2.26 we show the fission fragment frequency divisions as a function of total kinetic energy release for various 4-mass-unit intervals.

Figure 2.22

Single fragment kinetic energy spectra for Cf^{252} spontaneous fission and photofission of U^{238} and Th^{232} .

Figure 2.23
Average total kinetic energy release and fragment yield
as a function of mass number for photofission of U^{238} .

Figure 2.24
Average total kinetic energy release and fragment yield
as a function of mass number for photofission of Th^{232} .

Figure 2.25

Fragment frequency distributions for U^{238} photofission as a function of total kinetic energy release summed over intervals of four mass units.

Figure 2.26

Fragment frequency distributions for Th^{232} photofission as a function of total kinetic energy release summed over intervals of four mass units.

Kinetic energies in photofission of U^{238} and Th^{232} were previously studied by Bochagov, Komar, and Solyakin (Soviet Phys. -JETP 11, 990, 1960) who used ionization chambers to measure the fragment energies. Their data, in the representation of Figs. 2.25 and 2.26 of this report, showed strong indications of two fragment energy groups for mass ratios ~ 1.0 to 1.3 , especially in the case of U^{238} . Those results, if true, lent strong support to the explanation of the dip in kinetic energy release at symmetric fission on the basis of the two-mode fission hypothesis (see Gordon and Brenner, LNS Progress Report, May 1961, p.66). Because of the difficulties of measuring fragment kinetic energies with ionization chambers and the lack of details about corrections for energy loss in the target, etc., in the paper by Bochagov, et al., we felt it necessary to repeat the experiments with the use of semiconductor detectors. As can be seen in Figs. 2.25 and 2.26, there is no indication of more than one kinetic energy group at any given mass division. Instead, we find a gradual, continuous shift of the energy distributions toward lower kinetic energies as symmetric fission is approached in both systems studied. Thus, our results rule out the two-mode hypothesis as an explanation for the kinetic energy dip at symmetric fission. We find the magnitudes of the dips to be approximately 5 and 14 MeV for U^{238} and Th^{232} , respectively.

A more detailed account of this work is given in the thesis of D. S. Brenner. (D. S. Brenner, R. D. Cooper (Dept. of Nucl. Eng., MIT, on leave from U.S. Army Natick Lab., Natick, Mass.), and G. E. Gordon).

B. Ratios of Independent Yields of Te^{131} and Te^{133} Isomers in Fission

Measurements of the ratios of the yields of tellurium isomers of spin $11/2^-$ and $3/2^+$ formed independently in fission permit inference of the average spin of the primary fission fragments of mass ~ 133 . Such measurements have been made for the isomeric pairs of masses 131 and 133 (LNS Progress Report, May 1, 1963), based on rapid separation of radiochemically pure tellurium and analysis of the quantities of 8.05-da I^{131} and 20.8-hr I^{133} found at successive time intervals after the separation. The growth-decay curve of each iodine isotope can be treated as a three-component curve with positive amount $A_I^{eff}(0)$ decaying with the I^{131} or I^{133} half life, and negative amounts $-A_m^{eff}(0)$ and $-A_g^{eff}(0)$ decaying with the half-lives of the metastable and ground-state isomers. The curves were analyzed by the Frantic computer program (P. C. Rogers, LNS Technical Report No. 76 (NYO-3303), June 1961). The initial yield ratio is derived from the A^{eff} values. The fission modes explored were $U^{235}(n_{th}, F)$, $Th^{232}(\alpha_{33}, F)$, $Th^{232}(d_{18}, F)$, $U^{238}(\alpha_{33}, F)$, and $U^{238}(d_{18}, F)$, the subscripts designating energy in MeV or "th" for thermal.

For the Te^{131} isomers, correction was made for contribution by β decay to the amounts of 30-hr Te^{131m} and 25-min Te^{131g} from independently formed 3.4-min Sb^{131} and 23-min Sb^{131} (decaying $6.8 \pm 1.0\%$ to Te^{131m}), using conventional treatment of charge distribution in fission, with the aid of a computer program (called Growth-Decay) written for the Bateman equations in general form including chain branching. The independent isomer yield ratios (reported previously, loc. cit.) are somewhat sensitive to the half-life of Sb^{131} , and would be higher if the half-life of the latter should be shorter, with the only large correction being 30% for 19-min Sb^{131} in thermal neutron fission.

At the time of the previous report it was thought that Te^{133g} had a half-life of about 3 min. Dr. S. G. Prussin (private communication) has established that the ground state has a half-life of 12.5 min, as corroborated by Dr. C. E. Bemis of our group (see Section IV. D). We have re-evaluated our data for the I^{133} growth and decay (8 points) from 53-min Te^{133m} and 12.5-min Te^{133g} , using the Frantic values for $A_I^{\text{eff}}(0)$ and $-A_m^{\text{eff}}(0)$, taking $A_g^{\text{eff}}(0)$ as the difference between these, assigning a rather large uncertainty to the value. Corrections were then made for growth of the Te^{133} isomers during bombardment and up to time of tellurium separation considering the independent fission yields of 30-sec Sn^{133} and 2.64-min Sb^{133} (decaying 72% to Te^{133m}). The independent isomer-yield ratios are lower for $\text{U}^{235}(\text{n}_{\text{th}}, \text{F})$ and higher for charged particle fission than reported previously with the assumed half lives of 53 min (Te^{133m}) and 3 min (Te^{133g}). The final results are given in Table 2.III. The fractional yield of mass 131 high-spin isomer in thermal neutron fission of U^{235} , $\text{Te}^{131m} / (\text{Te}^{131m} + \text{Te}^{131g})$, of 0.64 ± 0.05 is in excellent agreement with that of M. Talat-Erben, L. E. Glendenin, H. C. Griffin, and E. P. Steinberg (private communication) of 0.63 ± 0.03 . Our value for mass 133 is similar, $0.61^{+0.06}_{-0.10}$.

Table 2.III

Ratios of Independent Yields of Te^{131} and Te^{133} Isomers
in Fission 30 hr Te^{131m} and 25-min Te^{131g} ; 53-min Te^{133m}
and 12.5-min Te^{133g}

Type of fission	$\text{U}^{235}(\text{n}_{\text{th}}, \text{F})$	$\text{Th}^{232}(\alpha_{33}, \text{F})$	$\text{Th}^{232}(\text{d}_{18}, \text{F})$	$\text{U}^{238}(\alpha_{33}, \text{F})$	$\text{U}^{238}(\text{d}_{18}, \text{F})$
$\text{Te}^{131m}/\text{Te}^{131g}$	1.8 ± 0.4	3.3 ± 0.5	2.7 ± 0.5	5.0 ± 1.0	3.3 ± 0.5
$\text{Te}^{133m}/\text{Te}^{133g}$	1.5 ± 0.5	2.8 ± 0.7	1.7 ± 0.9	3.1 ± 2.0	1.8 ± 1.1

We have carried out much more detailed investigation of the dependence of fractional yield of high-spin isomer upon assumed spin distribution of the primary fission fragments. First, exhaustive exploration of the yields and neutron emission from primary Te isotopes of mass numbers A, A+1, A+2, and A+3 that give final Te^A and the residual excitation of Te^A was made by the Monte Carlo method (G. E. Gordon, N. K. Aras, and L. A. Wilson, LNS Progress Report, May 1, 1964). This analysis gives good predictions of the most-probable charge Z_p of the secondary fission products, of the average number of neutrons ν_A as function of A, of the gross kinetic energy spectrum, and of the average prompt γ -ray energy release. We next assume that the angular momentum distribution of the primary fission products is given by the statistical formula:

$$P(J_c) = (2J_c + 1) \exp[-J_c(J_c + 1)/B^2] \quad (1)$$

following H. Warhanek and R. Vandenbosch (J. Inorg. Nucl. Chem. 26, 669, 1964), where $P(J_c)$ is the probability for primary fragments of spin J_c and B is a spin-cutoff parameter. Then the effect of neutron boil-off in changing the angular-momentum distribution was carried out using the statistical treatment of R. Vandenbosch and J. R. Huizenga (Phys. Rev. 120, 1313, 1960) with spin-cutoff parameter σ_n taken as 4 and barrier coefficients $T_l(E_n)$ from optical model calculations. A similar treatment was applied to γ emission, assuming two γ rays, dipole radiation only. Of

the distribution resulting from the emission of these two γ rays, we assume that all nuclei having $J < 7/2$ decay to the lower isomer ($3/2^+$), all with $J > 7/2$ decay to the upper isomer ($11/2^-$) and that those with $J = 7/2$ divide equally between the isomers upon emission of the third γ ray. The average of the results for spin-cutoff parameter σ_γ of 3 and of 4 was taken.

It was found that changes in average spin $\bar{J}$ from neutron and γ -ray emission were not large for $\bar{J}_c$ near 8 ($B=10$), and that they were not very sensitive to the σ_n and σ_γ values chosen. The results were also not very sensitive to neutron kinetic energy or the number of γ rays emitted. Fig. 2.27 shows the high-spin isomer fraction $\sigma_m/(\sigma_m + \sigma_g)$ predicted for Te^{133} as a function of B , the spin-cutoff parameter in Equation (1), using our best estimates of the parameters. The predictions for Te^{131} are virtually identical. We estimate that for a given observed high-spin fraction, the uncertainty in determining B is about $\pm 1.5 \hbar$ units at most. The average initial spin $\bar{J}_c$ varies more slowly than B , so we expect an uncertainty of about $\pm 1.2 \hbar$ in the range of interest.

The measured fractional chain yield values of 0.64 and 0.61 for Te^{131} and Te^{133} in thermal neutron fission correspond to B values of 6.0 and 5.9, or $\bar{J}_c$ values of $(5 \pm 1) \hbar$ for both. M. Hoffman (Phys. Rev. 133, B174, 1964) reports the value $(7 \pm 2) \hbar$ for angular correlations between γ rays and gross fission fragments in thermal neutron fission; Warhanek and Vandenbosch (loc. cit.) infer the value of about $6.5 \hbar$ from the Cs^{134} isomer ratio formed in photofission [2.9-hr $\text{Cs}^{134m}(8^+)$, 2.1-yr $\text{Cs}^{134g}(4^+)$]. The agreement is gratifying.

The spins of the fissioning U^{236} nuclei are either 3 or 4 $\hbar$ for s-wave neutron capture, by no means enough to explain the $\bar{J}_c$ value. The relatively high value, with a corresponding one of nearly opposite direction for the complementary light fragment can be explained as the result of torque arising if the main axes of the fragments form initially at an angle somewhat less than 180° . This would also be in accord with Hoffman's finding (loc. cit.) that the fragment spin vectors are directed preferentially to the direction of fragment motion.

Figure 2.27

Calculated high-spin fraction, $\sigma_m/(\sigma_m + \sigma_g)$, for Te^{133} as a function of B .

In the cases of charged-particle fission, the primary fragments have somewhat higher $\bar{J}_c$ values. The case of $\text{Th}^{232}(\alpha_{33}, \text{F})\text{Te}^{131}$ indicates a value of about $7 \hbar$ assuming the emission of about 2 prompt neutrons to yield Te^{131} (in contrast to 0.78 for thermal neutron fission). However, the compound nucleus is expected to have an average angular momentum J_f of about $13 \hbar$. The fission anisotropy implies that the projection K along the symmetry axis is $\sim 8 \hbar$. The projection perpendicular to the symmetry axis, $\sim 10 \hbar$, becomes orbital angular momentum of the fission fragments. If the light fragment has $\bar{J}_c$ about the same as the heavy ($7 \hbar$), it is apparent that there is some cancellation of the two $\bar{J}_c$ values, again implying off-axis fission. (D. G. Sarantites, C. D. Coryell, and G. E. Gordon)

C. Ranges of Shielded Nuclides from Thermal-Neutron Fission of U^{233} and U^{235}

As the shielded nuclides of interest, Rb^{86} , Sb^{126} , and Cs^{136} have very low independent yields, it has been necessary for us to develop chemical-separation methods capable of producing samples of those elements of very high radiochemical purity with reasonably high chemical yields. In addition, we want to be able to produce samples of several high-yield (Ba^{140} , Mo^{99} , Sr^{89}) and near-symmetric fission products (Ag^{111} , Cd^{115} , and isotopes of Sn and Sb). Considering the fission yields of each nuclide, a certain value was taken as the required value of the decontamination factor for various fission products from the element of interest. For example, Sb^{126} should be free from Mo^{99} with the decontamination factor of $\sim 10^6$; from Sn with $\sim 10^3$; from Cd with $\sim 10^2$; etc.

The chemical techniques developed are principally based on the usual qualitative analysis scheme and are as follows: the Al catcher foil is dissolved in 6 N HCl containing 30 to 50 mg each of Sb, Sn, Cd, Mo, Ba, Sr, Cs, and Rb carriers. After dissolution, ~ 50 mg of Ag^+ carrier is added, and the precipitated AgCl separated by centrifuging. The supernatant solution is adjusted to 2N in HCl and H_2S gas is passed through the solution repeatedly. The sulfides of Sb, Sn, and Mo are separated by centrifuging and treated with conc. HCl in order to separate Mo from the Sb + Sn fraction. The supernatant from (Mo + Sb + Sn)-sulfide-precipitation step is adjusted to 0.2 N in HCl and treated with H_2S gas. The CdS precipitate is removed from the solution and the supernatant solution is scavenged by $\text{Fe}(\text{OH})_3$ precipitation. Barium and strontium are precipitated as oxalates by adding $(\text{NH}_4)_2\text{C}_2\text{O}_4$ to the remaining solution. After centrifuging the (Ba + Sr) oxalate, the remaining solution is passed through an anion exchange column in the OH^- form, the eluent is evaporated under an infrared lamp in order to remove NH_4OH , and the resulting Rb and Cs fraction is obtained as the hydroxide salt. After several purification steps, Ag, Sb, and Sn are mounted on Pt discs by using self-electrodeposition techniques. Cadmium is also mounted as an electrodeposit on a Pt disc. Barium is purified and weighed as $\text{BaCl}_2 \cdot 2\text{H}_2\text{O}$, Sr as $\text{SrC}_2\text{O}_4 \cdot \text{H}_2\text{O}$, and Mo as oxalate. Each precipitate is mounted on a Pt disc by means of sedimentation and fixed by using some organic binder. We are investigating the separation and purification of Rb and Cs by means of cation-exchange resin, eluted with 0.5 N HCl in 10% ethyl alcohol solution. (H. Amano and H. Nakahara)

D. X-Rays and γ Rays from the Fission Products of U^{235}

The measurement of the radiations in the x-ray region associated with thermal-neutron induced fission of U^{235} was continued using a 2-meter bent quartz crystal spectrograph of the Cauchois type and Ilford G-5 100 micron plates (LNS Progress Report, p. 41, May 1963). The composite results of the exposures spanning the 25-110 keV region are tabulated in Table 2.IV. The most prominent feature of the spectrum is the pattern of K_{α} and K_{β} x-ray doublets of the elements. The intensity of the K_{α} doublet rises from Sb to a maximum at I and falls to background for Pm with appreciable odd-even alternation, odd Z intensity being stronger.

The K_{α} doublet of Xe exhibits a pronounced deficiency of intensity while the K_{β} doublet is just detectable. The $K_{\alpha 2}$ line of Cs appears to be of higher intensity than the $K_{\alpha 1}$ line, in disagreement with the normally expected intensity proportion $K_{\alpha 1}:K_{\alpha 2}::2:1$. (These two features had been incorrectly ascribed to I and Xe respectively, in LNS Progress Report, P. 41, May 1963). The range of the intensities of the K_{α} doublets exceeds the normal mass yield vs A distribution. This may be explained by the moderately high x-ray intensities expected from fission products of low Z from the Mayer-Jensen $h_{11/2}-d_{3/2}$ isomeric decay and for high Z from the Bohr-Mottleson low-lying rotational levels. Thirty-two γ rays were observed at energies above 57 keV while only seven were detected below that energy. Several of the gamma rays exhibited full width at half maximum corresponding closer to those of x-rays rather than gamma rays.

Three additional experiments have been completed and their results are being processed. The energy region between 40-60 keV has been re-examined to confirm the apparent lack of radiations in this region. A nuclear emulsion spanning the 25-35 keV region was exposed to a fission source immediately following a radiation period of one week to observe whether any change in the intensity ratio of Cs K_{α} doublet occurred. A nuclear emulsion was exposed to a fission source after a week of cooling in order to detect the contribution of the x-ray intensities expected for known Sn, Sb, Te, and Xe^{131m} fission products with the Mayer-Jensen $h_{11/2}-d_{3/2}$ isomeric decay.

Table 2.IV

X-ray and γ ray lines observed in the 25-110 keV region

Energy ^a (keV)	Relative ^b Intensity (D)	Probable Source	Energy ^a (keV)	Relative ^b Intensity (D)	Probable Source
26.111	0.025	Sb K	57.449	0.022	Ce ¹⁴³
26.319	0.010	— α_2	57.788	0.012	
26.359	0.044	Sb K	58.237	0.014	
27.203	0.031	Te K α_1	59.767	0.005	
27.473	0.062	Te K α_2	62.712	0.012	
28.317	0.125	I K α_1	64.466	0.08	
28.553	0.052	— α_2	64.787	<0.01	
28.611	0.216	I K	65.158	0.04	
29.454	0.027	Xe K α_1	65.445	0.04	
29.677	0.015	Sb K α_2	66.023	0.02	
29.723	0.010	Sb K β_3	67.233	0.010	
29.773	0.066	Xe K β_1	68.318	0.008	
29.956	0.020	— α_1	68.783	0.018	
30.631	0.255	Cs K	69.076	0.004	
30.968	0.199	{Cs K α_2 Te K α_1	69.194	0.005	
		{Ba K $\beta_1, 3$ Ba K α_2	69.772	0.008	Sm ¹⁵³
31.813	0.046	Ba K α_1	70.142	0.008	
32.188	0.071	I K β_3	70.429	0.010	
32.287	0.008	I K β_1	76.669	0.012	
32.325	0.020	La K α_2	77.661	0.010	
33.027	0.050	— α_1	77.904	0.005	
33.435	0.120	Ce K	78.764	0.010	
34.181	0.010	Ce K α_2	79.540	0.022	
34.267	0.035	Cs K α_1	81.052	0.020	Xe ¹³³
34.715	0.064	Cs K β_3	81.833	0.016	
34.918	0.028	Pr K β_1	85.599	0.004	
34.976	0.040	Pr K α_2	87.321	0.014	
35.542	0.094	Ba K β_3	93.628	0.030	
36.019	0.140	Ba K β_1	94.921	0.060	U K α_2
36.375	0.018	Nd K α_2	97.810	0.24	
36.448	0.028	La K α_1	98.542	0.072	U K α_1
36.839	0.030	La K β_3	108.677	0.018	
37.358	0.056	Pm K β_1			
37.627	0.026	Pm K α_2			
37.707	0.040	Ce K β_3			
38.129	0.024	Ce K α_1			
38.688	0.040	Pr K β_3			
39.073	0.012	Pr K β_1			
39.161	0.020	Pr K α_2			
40.553	0.009	Ce K β_3			
40.647	0.018	Pr K β_1			

^a Precision of the energy value is ± 0.01 keV^b The relative intensities (D scale) have been corrected for background but not for attenuation or geometry. The precision of the measurements is $\sim 20\%$.

(J. E. Canty and C. D. Coryell)

ADDRESSES

- L. Leifer, "The Ion Activity Function - A Quantitative Formulation of Species in Aqueous LiCl", Int. Symposium on Electrolyte Solutions, Toronto, Ontario, May 6, 1964.
- L. Leifer, "Low-Energy Electromagnetic Radiation from Thermal-Neutron Fission of U^{235} ", Dept. of Physics, Boston College, Chestnut Hill, Mass., Oct. 7, 1964.
- G. E. Gordon, "Monte Carlo Calculation of Prompt-Neutron Emission", Dept. of Chemistry, Clark University, Worcester, Mass., Nov. 4, 1964.
- G. Scatchard, "The Osmotic Pressure, Light Scattering, and Ultracentrifuge Equilibrium of Polyelectrolyte Solutions", American Electrochemical Society, Theoretical Section, Symposium, Toronto, Canada, May 5, 1964.
- G. Scatchard, "Lars Onsager, the Recipient. Introduction of T. W. Richards Medalist." Northeastern Section, American Chemical Society, May 14, 1964.

PUBLICATIONS

- E. L. Bahn, B. D. Pate, R. D. Fink, and C. D. Coryell, "Decay of 20-minute Ag^{115} ", Phys. Rev. 136, B603 (1964.)
- R. J. Silva, and G. E. Gordon, "Nuclear Levels of Cd^{115} and Cd^{117} from (d,p) Reaction Studies", Phys. Rev. 136, B618 (1964).
- C. D. Coryell, review of "Radionuklide" by K. Schmeiser, Springer-Verlag, Berlin, 1963, J. Am. Chem. Soc. 86, 2754 (1964).
- G. Scibona and B. Scuppa, "Potentiometric Determination of Nitrates in Benzene", J. Phys. Chem. 68, 2003 (1964).
- A. S. Kertes and Y. E. Habousha, "Miscibility Gap in Extraction Systems involving Alkylamines", Third Int. Conf. on the Peaceful Uses of Atomic Energy, Geneva, 1964, Paper P/755.
- A. S. Kertes, "Solubility Parameters of Long-Chain Alkylamines and their Activities in Organic Solvents", J. Inorg. Nucl. Chem. 26, 1764 (1964).
- G. Scatchard, G. M. Wilson, and F. G. Satkiewicz, "Vapor-Liquid Equilibrium X. An Apparatus for Static Equilibrium Measurements", J. Am. Chem. Soc. 86, 125 (1964).
- G. M. Wilson, "XI. A New Expression for the Excess Free Energy of Mixing", ibid., 127 (1964).
- G. Scatchard and F. G. Satkiewicz, "XII. The System Ethanol-Cyclohexane from 5 to 65°", ibid., 130 (1964).
- G. Scatchard and W. T. Yap, "The Physical Chemistry of Protein Solutions. XII. The Effects of Temperature and Hydroxide Ion on the Binding of Small Anions to Human Serum Albumin", J. Am. Chem. Soc. 86, 3434 (1964).
- G. Scatchard, "Lars Onsager, the Recipient. Introduction of T. W. Richards Medalist", Nucleus 41, 270 (1964).
- L. Leifer and E. Høgfældt, "An Approach to the Study of Electrolyte Behavior in Concentrated Solutions", Proc. First Austral. Conf. Electrochem., A. Friend and F. Gutmann, Eds., Pergamon Press, Oxford, 1964, p. 107.

- L. Leifer, "Report on the Melbourne Symposium on Tunnelling Effects", Proc. First Austral. Conf. Electrochem., A. Friend and F. Gutmann, Eds., Pergamon Press, Oxford, 1964, p. 949.

THESES

- C. E. Bemis, Jr., "Decay Studies of Antimony Fission Products", Ph.D. Oct. 1964.
- R. P. Black, "Heavy Ion Nuclear Reactions in the Medium Mass Region", Ph.D., May 1964.
- D. S. Brenner, "Kinetic Energy Release in Photofission of Uranium and Thorium", Ph.D. Sept., 1964.
- R. A. Duce, "Determination of Iodine, Bromine, and Chlorine in the Marine Atmosphere by Neutron Activation Analysis", Ph.D., May 1964.
- R. L. Yoest, "Isotonic Solutions of Aqueous Electrolyte Mixtures", Ph.D., June 1964.
- C. D. Strehlow, "An Infra-red Study of Hydrogen Bonding in Tertiary Amine Salts", S. M., Aug. 1964.

NUCLEAR CHEMISTRY (ORGANIC) GROUP

Summary

Radical anions of a number of aliphatic and aromatic systems have been investigated by electron spin resonance spectroscopy. 1-Deuterocyclopropane affords a paramagnetic species in dimethoxyethane-tetrahydrofuran, sodium-potassium alloy at -150° . The spectrum shows an even number of lines, consistent with expectations for the 1-deuterocyclopropane radical anion. Radical anions of a number of alkyl aryl ethers have been generated and shown to undergo a variety of reactions including dimerization and loss of methoxyl. The reactions observed are markedly dependent on orientation of substituents in the aryl moiety.

Reduction of either *cis*- or *trans*-4-*t*-butyl-1-methylcyclohexyl chloride with organotin hydrides affords the same mixture of *cis*- and *trans*-4-*t*-butyl-1-methylcyclohexane (1:12 at 30°) via a free radical chain reaction of chain length greater than 1000. Hydrogen atom transfer to the 4-*t*-butyl-1-methylcyclohexyl radical occurs preferentially *cis* to the 4-*t*-butyl group. Efforts to examine by electron spin resonance spectroscopy the 4-*t*-butyl-1-methylcyclohexyl radical and other tertiary carbon radicals of varying angular restraint are in progress.
(F. D. Greene)

The extent of transfer of the α -hydrogen from carbon to oxygen in the transition state for enolization of a simple ketone by hydroxide ion in water at 25° is 26-30% as measured by the secondary isotope effect (k_{DO}/k_{HO}), in satisfactory agreement with the 25% value deduced from the 0.25 slope of the Bronsted catalysis law plot of enolization by strong bases. This extent of transfer (order of the O---H bond) increases when weaker bases are employed, as shown by the increasing Bronsted slope with 8 weaker bases (up to 0.8 for trimethylacetate) and by the increasing percentage decrease in rate caused by addition of a fixed concentration of sodium perchlorate when weaker bases are used.

Substantial inverse entering group oxygen isotope effects exclude an acylium ion mechanism for the hydrolysis of benzoyl chloride in 75% dioxane-25% water or 50% dioxane-50% water solutions at 25° . Relative rates with H_2O^{16} and H_2O^{18} are 0.980 ± 0.005 and 0.986 ± 0.004 respectively.

The rate of decomposition of benzenediazonium salts is practically independent of solvent and anion over a wide range of structure (water to methylene chloride, chloride to hexafluoroantimonate). In the presence of bromine or iodine, low yields of halobenzenes are formed, but the rates depend on the halogen concentration. (C. G. Swain)

I. Radical Ions

Recent work in the electron-spin resonance area has been concentrated on adamantanes, cyclopropanes and aryl ethers. Mono- and dideuteriocyclopropane have been prepared. The former affords a paramagnetic species under our usual conditions (sodium-potassium alloy, tetrahydrofuran-dimethoxyethane, -150°). Although the signal is weak, it is an even-line spectrum of line separation and g-value approximately the same as those previously observed here for cyclopropane. The intensity is not adequate to state the total number of lines; the central portion consists of four lines, in accord with expectations for the middle four of the six lines predicted for the monodeuterocyclopropane radical ion.

A substantial quantity of adamantane, the principal precursor to many derivatives of interest in this work, has been synthesized. Part of the material has been converted to 1-bromo-, 1-nitro-, 1-methoxy- and 1-deuterioadamantane; 2-deuterioadamantane is being prepared via "Meerwein's Ester".

In conjunction with the studies on the alkoxyadamantanes, an examination has been undertaken of alkoxybenzenes in the sodium-potassium, tetrahydrofuran-dimethoxyethane system. The anisole radical anion is observed at low temperatures (-100°); at higher temperatures it disappears and is replaced by the radical anion of biphenyl. o-Dimethoxybenzene and the meta-isomer have afforded only the anisole radical anion; from o- and meta-methoxytoluene only the toluene radical anion has been observed. p-Methoxytoluene and p-dimethoxybenzene have afforded the corresponding radical ions with no evidence for loss of methoxyl or for dimerization at elevated temperatures. Possible relationships between loss of methoxy and orbital degeneracy in the aromatic compounds are under examination in other systems.

The new electron-spin resonance spectrometer (obtained by funds provided by the Office of the Provost, M.I.T.) has arrived and is in operation. A new high vacuum line has been constructed which increases several-fold our facilities for sample purification and preparation. (K.W. Bowers, G.J. Nolfi, Jr. J. Pazos)

II. Selectivity in Free Radical Reactions

Selectivity in atom transfer to the 4-t-butyl-1-methylcyclohexyl radical has been examined by use of the alkyl halide-tin hydride reaction. The isomeric cis- and trans-4-t-butyl-1-methylcyclohexyl chlorides have been prepared, structures have been established, and the chlorides have been reduced to mixtures of cis- and trans-4-t-butyl-1-methylcyclohexane by various tin hydrides. The reaction proceeds by a free radical chain reaction of chain length greater than 1000.

The principal findings regarding selectivity are summarized in Table 3.1.

Table 3.1

Reduction of cis- and trans-4-t-butyl-1-methylcyclohexyl chloride^a with tin hydrides to cis- and trans-4-t-butyl-1-methylcyclohexane in benzene solution, azobisisobutyronitrile-hv initiation

Reactants ^a trans-RCI/cis-RCI	Temperature	Tin hydride	Products trans-RH/cis-RH
1/0 ^b	30°	(n-Butyl) ₃ SnH	11.4 ± 1.2
0/1 ^b	30°	"	12.0 ± 0.7
4.7/1	0°	"	18.5 ± 0.3
4.7/1	80°	"	8.4 ± 0.2
4.7/1	0°	(C ₆ H ₅) ₃ SnH	12.4 ± 1.2
4.7/1	0°	(n-Butyl) ₂ SnH ₂	14.3 ± 0.7

^aCis and trans refer to orientation of alkyl groups. ^bRecovery of chlorides after partial reaction showed no isomerization of reactants.

The data indicate that: 1) both the cis- and the trans-reactants proceed through a common intermediate, the 4-t-butyl-1-methylcyclohexyl radical, 2) the hydrogen atom transfer to this radical is largely from the "axial" direction, 3) selectivity in the atom transfer step decreases with increasing temperature and with increasing reactivity of tin hydride. The relation of selectivity to geometry of the intermediate radical is under further investigation. (Añña Fang)

A number of approaches have been made for the direct observation by electron-spin resonance spectroscopy of the 9-decalyl radical (see previous reports). Efforts to prepare di-9-decalyl mercury have been unsuccessful. Irradiation of 9-bromo and cis and trans 9-chlorodecalins in hydrocarbon glasses at -182° afforded paramagnetic species. The spectra obtained to date lack adequate resolution but it is hoped that use of other glasses may resolve this difficulty and permit ultimate examination of the radicals labelled with carbon-13. (N. N. Lowry)

III. Enolization of Ketones by Bases

The structure of the transition state in the base-catalyzed enolization of ketones is being studied by observing effects on rate and equilibrium in the enolization of sodium p-(α -isobutylphenacyl) benzenesulfonate, an optically active ketone with only one α -hydrogen, caused by variations in base, ketone, solvent, inert salts and temperature. With different alkoxide, phenoxide and carboxylate anions in water at 25° , the Brønsted catalysis law plot is a smooth curve with slope varying from 0.25 for hydroxide to 0.8 for trimethylacetate, indicating that as the basicity of the oxygen anion increases there is a progressive variation in the structure of the transition state from one close to reactants to one close to products with respect to bond order of the O---H bond which is forming.

The secondary isotope effect from comparison of DO^- and HO^- as bases (in D_2O and H_2O respectively) affords another criterion of the degree of formation of the O---H bond at the transition state. We have now found that this ratio is 1.23. Since the maximum effect is the equilibrium constant for the reaction $\text{DO}^- + 1/2 \text{H}_2\text{O} \rightleftharpoons \text{HO}^- + 1/2 \text{D}_2\text{O}$, for which recent values range from 2.0 to 2.2, this corresponds to $\log 1.23 / \log K$ or 26-30%, in satisfactory agreement with the value of 25% derived

from the slope of the Brønsted catalysis law plot in the region of catalysis by strong bases.

Salt effects provide further evidence for this steady shift of transition state structure with decreasing basicity of the anion. Added sodium perchlorate decreases the rate attributable to each anion, but this salt effect increases in the sequence $\text{HO}^- < \text{CF}_3\text{CH}_2\text{O}^- < \text{C}_6\text{H}_5\text{O}^-$ showing again that the transition state decreasingly resembles reactants or increasingly resembles products as weaker bases are employed.

The reacting bond rule (Swain and Thornton, J. Am. Chem. Soc., 84, 817 (1962)) predicted all of the above results qualitatively before this work was begun. However, unambiguously curved Brønsted plots had not been previously demonstrated, and the curvature found and variation in transition state structure which must be responsible for it are much larger quantitatively than we had expected. (M. F. Grundon)

For the next phase of this problem, we have calculated the primary kinetic isotope effect for the α -hydrogen of the ketone from a linear three-center model ($\text{O} \cdots \text{H}^* \cdots \text{C}$) of the activated complex, as a function of transition state parameters (bond orders and coupling force constant). The calculated values for various $\text{O} \cdots \text{H}$ bond orders (in parentheses), neglecting bends and assuming that the total bond order to this hydrogen remains unity and that the coupling force constant (f_{12}) is the minimum one ($\sqrt{f_{11}f_{12}}$), are 1.000 (0.0); 2.698 (0.1); 4.363 (0.2); 6.179 (0.3); 6.917 (0.4); 5.820 (0.5); 4.269 (0.6); 3.003 (0.7); 2.058 (0.8); 1.338 (0.9); 0.597 (1.0). We have also calculated primary kinetic isotope effects for entering group, leaving group and central atom for a number of other displacement reactions ($\text{O} \cdots \text{H}^* \cdots \text{O}$, $\text{O} \cdots \text{C}^* \cdots \text{Cl}$, $\text{O} \cdots \text{C}^* \cdots \text{Cl}$, $\text{O} \cdots \text{C}^* \cdots \text{Br}$, $\text{O} \cdots \text{C}^* \cdots \text{I}$ and $\text{O} \cdots \text{C}^* \cdots \text{Cl}^*$). Experimental values of the α -hydrogen isotope effects in this ketone enolization for various $\text{O} \cdots \text{H}$ bond orders (equal to the Brønsted slopes) will be obtained and compared with the calculated values to test the assumptions and theory underlying these calculations. The special importance of this work derives from the fact that although most current theories predict a maximum for about half transfer, in fact no system is known in which an experimental primary isotope effect first rises and then unambiguously falls with a systematic monotonic change of structure. The present system must give this result if these theories are correct because the transfer ranges from 0.25

complete for hydroxide ion to 0.8 complete for trimethylacetate ion, with seven bases in between. The maximum primary isotope effect is predicted for 0.40 transfer. (R. L. Schowen)

IV. Solvolysis of Benzoyl Chloride

The transition state for solvolysis of benzoyl chloride is being examined through isotope effects to attempt to distinguish between one resembling an acylium ion (S_N1 mechanism), the one corresponding to a concerted displacement (S_N2), or the one expected for an addition-elimination mechanism with the second step (elimination step) rate determining. The principal difference is in the bond order of the O---C bond to the entering solvent molecule at the transition state, which should be approximately zero, 0.2-0.8, and unity for the three mechanisms respectively.

The results of two determinations of the oxygen isotope effect in the hydrolysis of benzoyl chloride in aqueous dioxane are given in Table 3.2. Errors

Table 3.2
Oxygen isotope effects in the hydrolysis of benzoyl chloride

Solvent	50% water $\rightleftharpoons$ 50% dioxane	25% water $\rightleftharpoons$ 75% dioxane
Isotope ratios, O^{18}/O^{16}		
Water	0.001962 ± 0.000005	0.001965 ± 0.000007
Benzoyl chloride	0.001973 ± 0.000004	0.001943 ± 0.000005
Benzoic acid	0.001981 ± 0.000003	0.001974 ± 0.000004
Isotope effect k_{16}/k_{18}	0.986 ± 0.004	0.980 ± 0.005

noted are standard deviations from the mean. In 50% dioxane $\rightleftharpoons$ 50% water (by volume) at 25° the isotope effect (k_{16}/k_{18}) is 0.986 ± 0.004 . In 75% dioxane $\rightleftharpoons$ 25% water at 25° the isotope effect is 0.980 ± 0.005 . Since the two isotope effects are the same within experimental error, it is quite probable that the hydrolysis mechanism is the same in both cases. It has been suggested previously that the hydrolysis mechanism changes as the amount of water in the solvent changes, but apparently such a change does not occur in the range between 25% water and 50% water. The magnitude of the observed isotope effect is not what would be expected for a mechanism involving ionization of

benzoyl chloride to an acylium ion followed by addition of water. More determinations of the isotope effect are needed in order to increase the precision of the results. Isotope effects measured in other water-dioxane mixtures will also be of interest.

In order to determine the isotope effect in the hydrolysis it is necessary to analyze the oxygen-18 content of the benzoyl chloride and water which are used in the reaction as well as the benzoic acid which is formed. All oxygen is converted to carbon dioxide, which is analyzed with a Consolidated-Nier isotope ratio mass spectrometer, model 21-201.

Water is analyzed by exchange with carbon dioxide. At least a tenfold excess of water is used. The equilibrium constant for the partitioning of oxygen-18 between water and carbon dioxide at 25° is known.

Oxygen analysis in benzoyl chloride is done by conversion of a sample of benzoyl chloride to benzamide by use of ammonia in monoglyme. The amide is recrystallized from xylene, dried, and pyrolyzed to water and benzonitrile over glass wool at 700°. The water formed is exchanged with carbon dioxide as before.

The benzoic acid isolated after completion of the hydrolysis of benzoyl chloride is recrystallized from petroleum ether, sublimed, and pyrolyzed to carbon dioxide and benzene over glass wool at 775°. The pyrolysis is quantitative.

(M. H. O'Leary)

V. Decomposition of Benzenediazonium Salts

The mechanism by which benzenediazonium fluoborate yields phenol and chlorobenzene when heated in aqueous hydrochloric acid may involve rate-determining cleavage to a phenyl cation, (possibly a diradical or triplet), concerted displacement by water or chloride, an addition-elimination process, or intermediate formation of a covalent azo compound and its decomposition to phenyl radical.

In order to determine whether free phenyl radical intermediates are present in solution, an attempt was made to use bromine as a radical scavenger. Decomposition in 0.05 M. sulfuric acid saturated in bromine produced p-dibromobenzene in yields up to 38%, a mixture of oxidation and bromination products of phenol, and traces of bromobenzene. This solvent has serious disadvantages

because of the limited solubility of bromine (0.3 M.) and the insolubility of the products. The precipitated solids absorb a considerable amount of bromine. Therefore 75% acetic acid-25% water (by weight) was used in further work, because it is miscible with bromine and dissolves all of the products but tetrabromobenzoquinone.

When 0.1 M. benzenediazonium fluoborate was decomposed at 25° for 48 hours in solutions 0.5-4.0 M. in bromine in 75% acetic acid, the results in Table 3.3 were obtained by gas chromatography.

Table 3.3
Bromobenzenes formed in the presence of bromine

<u>Br₂ M.</u>	<u>Bromobenzene, %</u>	<u>p-Dibromobenzene, %</u>	<u>Total, %</u>
0.44	7.7		7.7
1.1	12.2		12.2
2.2	12.1	5.9	18.0
3.0	.8	21.5	22.3
4.0		28.0	28.0

The reaction causing formation of bromobenzene in approximately first order in bromine up to 4 M., whereas trapping of a free radical in the presence of such high bromine concentration would be zero order. The reaction converting bromobenzene to p-dibromobenzene is higher than first order in bromine and certainly does not correspond to bromination of a diradical intermediate. Bromobenzene in 75% acetic acid 4 M. in bromine was 98% converted to p-dibromobenzene in 10 hours at 25°. It is evident that any free radicals present are not being scavenged by bromine. Iodine also failed as a scavenger. Decomposition in 75% aqueous acetic acid saturated with iodine produced only 13% iodobenzene and no detectable p-diiodobenzene.

Table 3.4 gives rate constants for decomposition in a variety of solvents (without bromine added).

Table 3.4

Decomposition of benzenediazonium fluoborate at $25.0 \pm .1^\circ$

Solvent	$k \times 10^5, \text{sec.}^{-1}$
25.0% CH_3COOH	$4.41 \pm .027$
50.0% CH_3COOH	$4.13 \pm .025$
75.0% CH_3COOH	$3.86 \pm .023$
99.9% CH_3COOH	$2.26 \pm .013$
85% H_3PO_4	2.332
75% H_2SO_4	2.14 ± 0.02
100% H_2SO_4	$2.065 \pm .011$
108% H_2SO_4	$2.078 \pm .011$
Dioxane	$1.15 \pm .020$
Sulfolane	$1.24 \pm .010$
Methylene chloride	$2.20 \pm .048$
Methylsulfolane	$1.22 \pm .030$
Methanol	$9.6 \pm .2$

There seems to be little correlation between rate of reaction and nucleophilicity of solvent. Dioxane gives the slowest rate yet obtained, half of that of methylene chloride, which is neither very basic nor very nucleophilic. The rate of decomposition is virtually constant in sulfuric acid solutions in range 75%–108%, although the nucleophilicity of the medium decreases greatly. Variation of dielectric constant has little effect, since the rate is the same rate in dioxane as in sulfolane. Bond formation to the solvent at the transition state must therefore be very slight or nonexistent.

Since the reaction in methylene chloride, dioxane, sulfolane, and methyl sulfolane is primarily with the anion, a variety of anions were tried. The results in Table 3.5 were obtained. Bonding to the anion is evidently also very slight.

Table 3.5

Decomposition of benzene diazonium salts at $25.0 \pm .1^\circ$

Anion	Solvent	$k \times 10^5, \text{sec.}^{-1}$
BF_4^-	Methylene chloride	2.20 ± 0.048
PF_6^-	" "	$2.61 \pm .036$
SbF_6^-	" "	$3.2 \pm .5$
SbCl_6^-	" "	$6.35 \pm .065$
TsO^-	" "	$3.7 \pm .3$
BF_4^-	Methylsulfolane	$1.22 \pm .05$
TsO^-	"	$1.20 \pm .1$

We plan to distinguish between the two most likely remaining mechanisms, a concerted displacement with the transition state coming very early (close to reactants in structure) and one proceeding through a phenyl cation (which would have the transition state then close to this very unstable intermediate), by measuring the magnitude of the nitrogen isotope effect: this should be unity for the concerted displacement and a maximum for phenyl cation. (J. E. Sheats)

ADDRESSES

- C. G. Swain, "The Effect of Structure on Solvation Bonds in Transition States", Seventh Gordon Research Conference on Chemistry and Physics of Isotopes, Tilton, N.H., July 13, 1964.
- C. G. Swain, "Isotope Effects for Study of Organic Reaction Mechanisms", Eighth Organic Chemistry Symposium, Quartermaster Corps, Natick, Mass., October 14, 1964.

PUBLICATIONS

- F. D. Greene, C-C. Chu, and J. Walia, "Decomposition of Alkyl Hypochlorites. The 4-*t*-Butylcyclohexyl Radical", J. Org. Chem., 29, 1285 (1964).
- F. D. Greene, H. P. Stein, C-C. Chu and F. M. Vane, "Mechanisms of Decomposition of Diacyl Peroxides", J. Am. Chem. Soc., 86, 2080 (1964).
- C. G. Swain and C. R. Morgan, "A Convenient Spectrophotometric Method for Following the Reactions of Arenesulfonates", J. Org. Chem., 29, 2097 (1964).

THESES

- J. C. Stowell, "Stereospecificity in Radical Termination Reactions", Ph.D. Thesis in Chemistry, M.I.T., August, 1964.
- Barbara Streicher Magid, "Substituent Effects on the Chlorine Isotope Effect in Methanolyses of Triarylmethyl Chlorides", Ph.D. Thesis in Chemistry, M. I.T., June, 1964.
- D. W. Henderson, "Substituent Effects on the Nitrogen Isotope Effect in the Thermal Decomposition of Areneazotriphenylmethanes", Ph.D. Thesis in Chemistry, M.I.T., June, 1964.

COSMIC RAY GROUP

I. X-ray Astronomy

A. Techniques have been developed for the determination of angular sizes of celestial x-ray sources. Of the techniques considered one type of collimator has been specifically studied and developed. The resolving angle of this device is typically 10 arc minutes at present and it is expected that a resolving angle of one arc minute will be achieved soon.

B. Collimators of the above type were mounted in front of x-ray detectors and flown on an Aerobee-rocket in collaboration with Giacconi, Gursky, and Waters (American Science and Engineering, Inc., Cambridge, Mass.). The angular size of the Scorpius x-ray source was found to be smaller than 10 arc minutes. This result makes any of the physical explanations of the source which have been recently advanced very doubtful. On the other hand, the Sagittarius x-ray source which was resolved from the Scorpius source seems to be a complex of localized sources each smaller than 30 arc minutes. Further analysis of the data is in progress.

C. Physical explanations of the x-ray sources have been sought. Thus far, none of the known physical processes seems adequate to explain the results concerning these newly discovered astronomical objects. (G. Clark, G. Garmire, M. Oda and M. Wada)

II. Time Variations of Cosmic Rays

Continuous observations of cosmic ray meson intensities are still being made in Building 26. The data are punched on IBM cards, tabulated, and plotted by automatic processing.

The analysis of the daily variations has been carried out for six months. The data obtained at Deep River, Canada, were used together with MIT data. The hourly data in each day are fitted to the Fourier components up to the fourth order. The curve built up from the four components for each day shows a complicated shape which varies day by day. There is, however, a tendency that the same shape appears on the next day with slight change. It means that the shape is genuine and reflects a complicated field configuration outside the earth.

In order to get some idea about the shape of daily variation, the times of maximum and of minimum intensity in a day are picked up. The time is converted to the direction of arrival of cosmic rays outside the terrestrial magnetic field by using the knowledge of the deflection of charged particles in the field (see MIT, LNS Technical Report No. 77, August 1962, K. G. McCracken, U. R. Rao and M. A. Shea). The histogram of arrival direction for the maximum shows a peak at 90° east of the sun, while that of the minimum shows broader distributions with peak at 45° west of the sun. The fact that the difference in directions of maximum and minimum is not 180° is another expression of the complicated shape of the daily variation.

As we know from the direct measurement of the interplanetary magnetic field by IMP satellite (see N. F. Ness and J. M. Wilcox, Phys. Rev. Letters 13, 461 (1964)), the direction of the line of force is 45° west of the sun, which is very reasonable with the velocity of the plasma as about 400 km/s (H. Bridge, A. Egidi, A. Lazarus, E. Lyon, and L. Jacobson in this volume). The similarity of the directions of the line of force and least directional intensity of cosmic rays gives us the following speculation. Cosmic rays coming outside the solar system along the interplanetary magnetic lines of force are generally reflected back at respective mirror points between the earth and the sun. In such case, cosmic rays distribute isotropically at any region in the solar system. If, on the other hand, there are irregularities of line of force between the earth and the sun along the line of force to which the earth crosses, cosmic rays going toward the sun are scattered out and no longer resume their original line. As there is a lack of returning particles the intensity is less than normal value. As there is a possibility that the scattering-out effect is almost cancelled out by scattering-in effect, it should be examined by solution of strict diffusion equation which matches with the situation mentioned above. The analysis is in progress with the help of S. Olbert. (V. Sarabhai, G. L. Pai, and M. Wada)

III. Primary Gamma Rays

A. Explorer XI

The data from Explorer XI have now been completely analyzed and the results will appear in the April 1, 1965, issue of the Astrophysical Journal. The final results are in no basic way different than those previously reported. We observe an apparent isotropic intensity of gamma ray in the 70 MeV and greater region of about $3 \times 10^{-4} \text{ cm}^{-2} \text{ sec}^{-1} \text{ sterad}^{-1}$. The observed lack of dependence of this intensity upon the geomagnetic latitude of the satellite would be strong evidence of a cosmic origin were the statistical significance of the results better. Unfortunately, we have used all the available dates and the device, while still in orbit, has long been inoperative. (W. Kraushaar, G. Clark, G. Garmire, and P. Higbie)

B. OAO Experiment

We have been invited to include the duplicate to the Explorer XI instrument as part of the first Orbiting Astronomical Observatory. The duplicate instrument was taken out of storage and proves to be working well. The Laboratory for Space Experiments (part of the Center for Space Research) is collaborating in this effort and has taken on the responsibility for adapting the data to digital form suitable for transmission over the OAO telemetry system. (W. Kraushaar, G. Clark, and G. Garmire)

C. OSO; S-57 Gamma Ray Experiments

The gamma ray detector for the Orbiting Solar Observatory is similar to the Explorer XI instrument but includes provision for rough energy measurement and more definitive study of

possible sources of background. Many technical difficulties have appeared in recent months, including corona in the photomultiplier wiring and electronic circuit failure. These difficulties seem presently to be in hand. One instrument is installed in the spacecraft for comprehensive environmental tests. Another, the spare, is due for completion and delivery next week. A third instrument is being prepared for calibration in the tagged gamma ray beam at the California Institute of Technology synchrotron. (G. Clark, G. Garmire, and W. Kraushaar)

D. Low Energy Gamma Rays

This balloon-borne experiment is designed to detect with a triggered, thin-plate spark chamber, gamma rays in the 10-70 MeV region. The importance lies in the possibility of distinguishing, through the shape of the low energy spectrum, the production mechanism. (P. Higbie, W. Kraushaar)

IV. Bolivian Air Shower Joint Experiment (BASJE)

During this period the low-mu run mentioned in the last report was completed and two major modifications to the apparatus were installed: (1) the automatic fast timing apparatus, and (2) the five density detectors in an outer ring at 150 m. At the time of this writing (Dec. 1, 1964), Dr. Suga is visiting in La Paz to assist Dr. Toyoda with the final testing of these new systems. Preliminary testing at MIT in La Paz indicates that both systems are operating satisfactorily.

A meeting at MIT of the principle participants of BASJE was held in November. At this time, we decided to run the apparatus essentially continuously during the next year to accumulate large showers ($> 5 \times 10^6$ particles) for use in the study of the longitudinal development of extensive air showers and the μ/e content of these showers. At present, the rather poor statistics indicate no separate group of low-mu air showers in the size range. With the enlarged detector array it now becomes feasible to improve the statistics. Concurrently with this run, we shall accumulate low-mu showers of smaller sizes ($\leq 5 \times 10^6$). Finally, the Michigan group intends to make concurrent runs to study the electromagnetic core structure with their neon bulb hodoscope and also to select cloud chamber events characterized by one incident nuclear active particle and no associated extensive air shower (i.e. study of surviving primaries).

Mr. A. Krieger is preparing his equipment for the study of the Cerenkov light associated with EAS. He plans to leave MIT for La Paz in the spring of 1965.

Two papers were presented at the Buenos Aires Conference in August 1964 by the MIT group. They report a nuclear core pattern which fluctuates dramatically from shower to shower and a primary energy spectrum with statistics greatly improved over that reported at Jaipur in December 1963. (G. Clark, M. Oda, H. Bradt, M. La Pointe, A. Krieger, and S. Rappaport)

V. Solar Neutrons and Gamma Rays

It remains our intention to monitor the high energy (1-100 MeV) neutral radiation from the sun by means of a balloon-borne detector during a period of high solar activity. Presently we are designing a detector which consists of a large ($\sim 1000 \text{ cm}^3$) CsI crystal and a large plastic scintillator of similar dimensions, both completely surrounded by an anti-coincidence shield. As stated in previous reports, we shall search for an increase in the neutral counting rate in near coincidence with the time of a visible flare. The relative increase of counting rates in the plastic and crystal will indicate the nature of the radiation (gamma rays or neutrons). This information together with the energy spectrum of the neutral particles should cast important light upon the mechanism of the acceleration of charged particles in solar flares. (H. Bradt, C. Forline, P. Oddone, and A. Krieger)

VI. Balloon X-Ray Astronomy

We searched for high energy x-rays from the Crab Nebula in a balloon experiment at an altitude above 130,000 feet carried out on July 21, 1964. We observed x-rays with energies between 15 keV and 60 keV emanating from a concentrated source lying in a region that included the Crab Nebula. Since the Crab Nebula is known to be a source of 4 keV x-rays and was the only one in the region from which the higher energy x-rays were observed to come, we concluded that we had observed the high energy end of the electromagnetic spectrum of the Crab Nebula. It now appears that this spectrum extends from radio frequencies to over 10^{19} c/s .

If one attempts to explain the observed x-ray emission of the Crab Nebula above 15 keV as blackbody emission from the surface of a neutron star as in the early speculations regarding the nature of the x-ray sources, one must assume a surface temperature of about $8 \times 10^6 \text{ K}^\circ$ which is far above the theoretical limits set by considering the cooling effects of neutrino emission from the core. Our observation of high energy x-rays is therefore new evidence against the neutron star hypothesis.

Figure 4.1 is a schematic diagram of the detector used in the experiment. Figure 4.2 is a block diagram of the electronic and recording system. The counting rates observed at various azimuths relative to the Crab Nebula are shown in Figure 4.3 where one can see the evident existence of a peak in the three middle channels near zero relative azimuth. We interpret these count rate peaks as being produced by high energy x-rays from the Crab Nebula. Figure 4.4 shows the predicted counting rates in the various channels relative to channel 3 for various assumed spectra. Figure 4.5 is a summary of data on the electromagnetic spectrum of the Crab Nebula including recent results of the NRL group near 4 keV and the present experiment.

SCHEMATIC DIAGRAM OF X-RAY DETECTOR.

Figure 4.1

SCHEMATIC
DIAGRAM OF ELECTRONIC
AND DATA RECORDING
APPARATUS

Figure 4.2

COUNTING RATES VERSUS AZIMUTH RELATIVE TO THE CRAB NEBULA

Figure 4.3

COMPARISON BETWEEN OBSERVED COUNTING RATES
AND THOSE EXPECTED FOR POWER LAW SPECTRA
WITH VARIOUS SPECTRAL INDICES α

Figure 4.4

SUMMARY OF DATA ON THE ELECTROMAGNETIC
SPECTRUM OF THE CRAB NEBULA

Figure 4.5

This experiment has been described more completely in a paper submitted for publication.
(G. Clark, W. Smith, A. Wan)

VII. Digitized Spark Chamber for Gamma Ray Astronomy

A triggering system for a spark chamber was built using an EFP 60 secondary emission tube to fire an Amperex 6279 hydrogen thyratron which in turn fired a nitrogen-filled spark gap. The coincidence pulse fed into the grid of the EFP 60 was supplied by 2 RCA 6810 photomultipliers covered with scintillating plastic and placed above and below the spark chamber.

Threshold-sparking measurements were made on a 6-inch square, 1/2 cm-gap first prototype spark chamber. It was found that 90% sparking efficiency could be attained with applied voltages in the range from 1000 to 2000 volts by flowing a mixture of 90% neon and 10% helium through the chamber. A set of 16 magnetic cores was wired and attached to one plate of the chamber which consisted of 64 closely spaced copper strips coupled in groups of four.

An electronic system for automatic readout of a spark chamber core memory has been built and tested. Using this system, we have confirmed that under proper conditions the spark current through the chamber was suitable for writing into these magnetic cores, and a series of tests showed that every spark through the chamber did indeed write in a core. Some sparks however wrote in two cores connected to adjacent groups of copper strips.

Three later prototype spark chambers approximately 18 inches square have recently been completed. These chambers are to be employed in testing the angular determination of particles through the chambers. Also the accompanying amplified set of memory cores is in the process of being wired. The three chambers consist of an aluminum plate and a printed circuit plate containing 256 separate conducting strips separated by clear ground glass spacers. (G. Clark, K. Dzugin, K. Richards, J. Stein)

VIII. Theoretical Studies of Cosmic Rays as a Relativistic Gas

The investigations outlined in the previous Progress Report (see Progress Report, May 1, 1964, Cosmic Ray Group, Section XII) are continued. As mentioned in that report, a detailed study of the behavior of cosmic rays in our galaxy requires the knowledge of the spatial distribution of the interstellar gas as well as the distribution of its velocity field. For this purpose we have compiled the astronomical and radio-astronomical data (especially those obtained from the 21-cm radiation) that exhibit (a) the density variation of the interstellar gas as a function of the galactic coordinates (r, θ, ϕ) , (b) the spatial dependence of the azimuthal component of the bulk velocity of the gas. Combining these two aspects of the experimental information and applying the continuity equation under steady-state conditions we were able to construct, by numerical computations, the radial component of the bulk velocity of the gas as a function of the galactic coordinates. The calculations show that the galactic gas ought to stream outward from the galactic center with a finite but small velocity (of the order of 10 km/sec). There are, however, regions where the radial velocity is substantially higher (particularly near the galactic center and at the edge of the galaxy).

Using this obtained velocity field, we then proceeded to construct the mean galactic magnetic field with the help of the mhd equations (essentially, $\frac{\partial \vec{B}}{\partial t} = \text{curl}(\vec{V} \times \vec{B})$). The results reflect the spiral nature of the large-scale magnetic field and show the appropriate strengthening and weakening of this field as one crosses the galactic arms radially.

If we accept such magnetic field pattern as a working hypothesis we may study explicitly their effect on the large scale transport of cosmic rays in our galaxy via the equations quoted in the previous Progress Report. Although the resulting transport equations appear to be rather complicated we believe that we shall be able to arrive at approximate solutions for several plausible boundary conditions. (A. Klimas, S. Olbert)

IX. Interplanetary Plasma

A. Satellite Instrumentation

Work on plasma instrumentation for EOGO, Pioneer, IMP and AIMP is continuing. The design of these instruments is supported under separate NASA contracts and is not discussed here.

in detail. A considerable effort is however being devoted to ensure the current designs, utilize fully recent theoretical and experimental results, so as to ensure the maximum scientific return from the experiments. (H. Bridge, A. Lazarus, E. Lyon, and F. Scherb)

B. Analysis of Plasma Measurements from Explorer XVIII

The analysis of data from IMP -I has yielded significant results during the past half year.

The most important of these are as follows:

1. The physical parameters which characterize the properties of the plasma in the interplanetary region have been computed for the first four months of satellite operation. This has been done for the shortest time scale possible for the IMP data, five minute averages, and also on an hourly basis. Further analysis is in progress. (E. Lyon)
2. In the case of the IMP satellite the MIT experiment measures the angular distribution of the plasma flux in the equatorial plane of the satellite with an angular resolution of about 20° . It has proved possible to superimpose the data for many satellite revolutions and to improve this resolution by a factor between five and ten. Approximately 20% of the data suitable for such a treatment have been analyzed to date and two important conclusions drawn. In the first place, the plasma flow in the interplanetary region is radially outward from the sun for 90% of the time covered by the observations. The accuracy of the result is about 5° ; i.e., the flow lies within a cone of 5° half angle centered on the earth-sun line. In obtaining this result, the velocity of the probe relative to the plasma (orbital velocity of the earth) has been taken into account.

The second result of this analysis concerns the plasma flow within the transition region. The anisotropy of the plasma flux within the transition region can be examined by measuring the angular distribution in exactly the manner just described. When this is done, it is found that the flux distribution in the "subsolar region" is essentially isotropic. However, the degree of anisotropy becomes larger and larger as the sun-earth-satellite angle increases. If it is assumed that the velocity vector which specifies the plasma flow in the transition region lies in a plane which contains the earth-sun line, the flow vector can be determined from the experimental results. The calculation leads to the result that the flow is roughly parallel to the boundaries of the magnetosphere and the shock transition. (A. Egidi)

3. The results obtained by the MIT plasma probe aboard IMP -I have been analyzed in an interpretive manner. Thus far the data referring to the subsolar region, both in front of the bow shock and in the transition region between the shock front and the magnetopause, have been subjected to a detailed, quantitative analysis. In particular, we have endeavored to derive the following quantities from the direct flux measurements of individual energy channels in our plasma probe: the "thermal" speed of protons (i.e., the random speed of protons in the frame of reference Σ_0 moving with the wind) and the proton densities on both sides of the shock. The above quantities may be computed unambiguously only after a sufficient amount of information is obtained about the form of

the distribution function in Σ_0 . For this purpose we have utilized the results from the first few inbound orbits of IMP -I in the transition region where the fluxes are observed to be nearly isotropic in 90 % of the cases. The mathematical task consisted in "unfolding" the five integrals (one for each proton channel) which relate the measured fluxes to the (unknown) distribution function. In constructing these integrals a particular care has been exercised to take into account accurately the laboratory tested response of the cup. The analysis has shown that the distribution of protons in the transition region is far broader than the maxwellian, it could be well approximated in all cases studied (about 70) by a form $f \propto (1 - \frac{w^2}{2w_0^2})^{-3}$ where w_0 represents the most probable speed of protons. The computed values of w_0 ranged between 120 and 180 km/sec. Similar unfolding procedures allowed us also to estimate w_0 in the interplanetary regime (the region ahead of the shock); the typical values obtained there ranged between 30 and 50 km/sec.

The computed proton densities exhibit persistently a surprisingly high ratio across the shock: while ahead of the shock the densities run typically between 4 and 7 protons per cc, behind the shock their values fluctuate between 30 and 100 particles per cc; the typical average ratios across the shock amount to about 10, i.e., significantly higher than the theoretically anticipated value of 4.

Using the above mentioned empirical distribution function we were also able to compute the kinetic energy densities of protons on both sides of the shock. Again, we have found that the energy density in the transition region is typically 10 times higher than that in front of the shock. The momentum conservation across the shock can be roughly accounted for, provided the electron and the magnetic energy densities are small compared with the derived energy densities of positive ions. (There seems to be a tentative experimental support of this assumption.)

Accepting the above observational findings as unbiased by some unknown effects of plasma on our Faraday cup, we are now confronted with a new challenge of justifying them on theoretical grounds. (G. Moreno, S. Olbert)

X. Composition of Primary Cosmic Rays*

A series of experiments have been carried out at the Yale Heavy Ion accelerator using O^{16} , B^{11} , and C^{12} nuclei in order to analyze in detail the limitations in identifying these particles by using the decay time of fluorescence from CsI(Tl). Decay time measurements have also been made using deuterium from the MIT cyclotron. We have now explored the low energy (approximately 3-10 MeV per nucleon) response of CsI(Tl) for particles from protons to O^{16} . The results will be a Master's thesis by A. Womack. Senior theses are being done on 1) the dependence of the decay time on the thallium content, and 2) measurement of the decay time from thin scintillators in which particles traverse at almost constant dE/dx . (A. Lazarus, A. Womack, M. Joseph, A. Russell, D. Grimes, and D. Slevin)

* See Progress Report, Nov. 1, 1963.

ADDRESSES AND INVITED PAPERS

- B. Rossi, Memorial Galileo Lecture on "X-Ray and Gamma Ray Astronomy" on the occasion of the COSPAR meeting in Florence, May 11, 1964.
- B. Rossi, Seminar on "X-Ray Astronomy"
Department of Physics, University of Bologna, May 18, 1964.
Department of Physics, University of Naples, May 20, 1964.
CERN, Geneva, Switzerland, June 25, 1964.
- B. Rossi, Seminar on "Interplanetary Plasma", at the Imperial College, London, June 16, 1964.
- B. Rossi, "Remarks on X-Ray Astronomy", given at the Solvay Conference in Brussels, Belgium, Sept. 1964.
- B. Rossi, "Recent Results concerning Interplanetary Plasma", Invited paper at the Nuclear Scientific Symposium of the Institute of Electrical and Electronic Engineers, Philadelphia, Pa., Octo. 1964.
- W. Kraushaar, "Observational Results on Gamma Rays", presented at the IAU Symposium No. 23 in Liege, Belgium on Aug. 17-20, 1964.
- A. Lazarus, "Measurements of Plasma in Space", Northeast Electronics Research and Engineering Meeting, Boston, November 1964.
- E. Lyon, "Results of the MIT Plasma Experiment on IMP -I", Space Science Seminar, Rice University, Houston, Texas, Oct. 1964.
- S. Olbert, "Interplanetary Magnetic Fields", invited paper presented at the Rome Session of the Quadricentennial Celebration of Galileo's Birth, Sept. 14-16, 1964.
- V. Sarabhai, G. L. Pai, D. Patel, and P. N. Pathak, "Geomagnetic Disturbance, Plasma Wind and Solar Activity", presented at the Second Bendum Earth Magnetic Symposium, Pittsburgh, October 1964.

PUBLICATIONS

- H. Bradt, G. Clark, M. LaPointe, M. Oda, V. Domingo, I. Escobar, K. Murakami, K. Suga, and Y. Toyoda, "Nuclear Core Structure of small EAS at 5200 m Altitude", IQSY Conference, Buenos Aires, 1964, to be published.
- H. Bradt, G. Clark, I. Escobar, K. Murakami, M. La Pointe, V. Domingo, K. Suga, Y. Toyoda, and J. Hersil, "Notes on the Primary Energy Spectrum from 6×10^{14} eV to 4×10^{17} eV", IQSY Conference, Buenos Aires (1964), to be published.
- W. L. Kraushaar, "Cosmic Gamma Rays - Experimental", I.A.U. Symposium on Astronomy from Space Vehicles, Liege, August 1964.
- U. Rao and V. Sarabhai, "Probing Interplanetary Plasma and Magnetic Fields with Galactic Cosmic Rays", Planetary and Space Sciences, October 1964.
- E. A. Womack, A. J. Lazarus, and M. Joseph, "The Fluorescent Decay of $\text{ScI}(\text{Tl})$ ", IEEE Transactions on Nuclear Science NS-11, 24 (1964). Proceedings of the 9th Scintillation and Semiconductor Counter Symposium.
- M. Oda, "The Neutron Star Hypothesis of Celestial X-ray Sources", Nature 202, June 1964.

THESES

- F. W. Floyd, Jr., "A Solar Wind Deuterium Detection and Charge-Spectrum Analyzer", M.S. in the Department of Electrical Engineering.
- J. W. Overbeck, "Experimental Test of the Inverse Compton Effect Hypothesis for the X-Ray Source in Scorpius", June 1964, Ph.D. in the Department of Physics.
- M. Oliven, "The Infrared Albedo of the Moon", B.S., June 1964.
- H. Hinteregger, "Measurement of Stellar Infrared Intensities", B.S., Physics, June 1964.
- W. Watson, "Studies in Optical Photon Correlation", B.S., June 1964.

ACCELERATOR PHYSICS COLLABORATION

Last year we projected a five point program. This program was:

1. Analysis of the Spark Chamber Film Taken in 1963

This film has been analyzed and two papers were presented at the Dubna Conference on High Energy Physics and two papers were published in the Physical Review Letters. These papers are:

"Total Cross sections and Angular Distributions for $\pi^- + p \rightarrow \eta^0 + n$ from Threshold to 1151 MeV".

"Total Cross Sections and Angular Distributions for $\pi^- p$ Charge Exchange in the 2nd and 3rd Resonance Regions".

" $\pi^- p$ Charge Exchange Scattering between 500 and 1150 MeV".

"The Production of Eta Zero Mesons in $\pi^- p$ Collisions".

2. Extension of the Spark Chamber work to the Energy Region 1.3 BeV to 3.0 BeV at Brookhaven National Laboratory.

Our equipment at the Cosmotron was dismantled, rebuilt and assembled on the floor of the AGS at Brookhaven National Laboratory. Preliminary test runs have been completed and data in this energy region should be acquired in early 1965.

3. The Taking of 500,000 Pictures in the 12" H₂ Chamber at CEA for the Study of $\gamma + p$ Interactions.

Approximately 800,000 good pictures (out of 1,700,000 pictures taken) were acquired at CEA with the 12" H₂ chamber. About 10% of the film was analyzed and a paper on this 10% was presented at the Dubna High Energy Physics Conference, and two papers were published in the Physical Review Letters. These papers are:

"Gamma Ray Proton Interactions between 0.5 and 4.8 BeV".

"N*₃₃ and ρ^0 Production by High Energy Photons".

"Multiple Pion and Strange Particle Production in γ -ray-proton Reactions between 0.5-4.8 BeV".

4. Pressing for Completion of the 500-liter H₂ Chamber.

The chamber is not yet complete. However, the LNS technicians have successfully rebuilt the helium refrigerator and have completed all plumbing on the chamber. The glass window has been installed. The first attempt at cooling down to liquid hydrogen temperature should occur in early 1965.

5. An Attempt to Study $\bar{p} + p$ Interactions with PEPR using Film from the 20" Brookhaven National Laboratory H_2 Chamber.

This program has been changed. There are no current plans to use 20" Brookhaven National Laboratory film. The prototype 5" system has undergone a stringent series of tests. The results of these tests show that PEPR, as a measuring instrument, has an absolute accuracy of ± 1 micron over a square 4.25" x 4.25". The 1 micron is an RMS deviation. The system has a 12 hour stability. Detailed plans now exist to use PEPR to analyze the CEA experiment. Construction on the 35 mm system (which was to be used on the 20" Brookhaven National Laboratory film) was stopped. All efforts are being made to convert the prototype PEPR into a production PEPR for our CEA film.

We have plans to continue our present work at Brookhaven next year with our steel plate spark chamber.

Our major effort next year is to finish the 500 liter chamber, take and analyze production pictures. This effort should produce an order of magnitude more data over what is known now. As part of this program we shall continue the design and prototype construction of a 7" three-view PEPR system.

PUBLICATIONS

- "Total Cross Sections and Angular Distributions for $\pi^- p \rightarrow \eta^0 + n$ from Threshold to 1151 MeV"
F. Bulos, R. Lanou, A. Pifer, A. Shapiro, M. Widgoff, R. Panvini, A. Brenner, C. Bordner, M. Law, E. Ronat, K. Strauch, J. Szymanski, P. Bastien, B. Brabson, Y. Eisenberg, B. Feld, V. Fischer, I. Pless, L. Rosenson, R. Yamamoto, G. Calvelli, L. Guerriero, G. Salandin, A. Tomasin, L. Ventura, C. Voci, F. Waldner,
Phys. Rev. Letters 13, No. 15, 486 (1964).
- "Total Cross Sections and Angular Distributions for $\pi^- p$ Charge Exchange in the 2nd and 3rd Resonance Regions".
F. Bulos, R. Lanou, A. Pifer, A. Shapiro, M. Widgoff, R. Panvini, A. Brenner, C. Bordner, M. Law, E. Ronat, K. Strauch, J. Szymanski, P. Bastien, B. Brabson, Y. Eisenberg, B. Feld, V. Fischer, I. Pless, L. Rosenson, R. Yamamoto, G. Calvelli, L. Guerriero, G. Salandin, A. Tomasin, L. Ventura, C. Voci, F. Waldner
Phys. Rev. Letters 13, No. 18, 558 (1964).
- " $\pi^- p$ Charge Exchange Scattering between 500 and 1150 MeV"
C. A. Bordner, A. E. Brenner, M. E. Law, E. E. Ronat, K. Strauch, J. J. Szymanski, P. Bastien, B. Brabson, Y. Eisenberg, B. Feld, V. Fischer, I. Pless, L. Rosenson, R. Yamamoto, F. Bulos, R. Lanou, A. Pifer, A. Shapiro, M. Widgoff, R. Panvini, G. Calvelli, L. Guerriero, G. A. Salandin, A. Tomasin, L. Ventura, C. Voci, F. Waldner,
Presented at Dubna Conference on High Energy Physics, August 1964.
- "The Production of Eta Zero Mesons in $\pi^- p$ Collisions"
F. Bulos, R. Lanou, A. Pifer, A. Shapiro, M. Widgoff, R. Panvini, L. Guerriero, G. Calvelli, G. Salandin, A. Tomasin, L. Ventura, C. Voci, F. Waldner, C. Bordner, A. Brenner, M. Law, E. Ronat, K. Strauch, J. Szymanski, P. Bastien, B. Brabson, Y. Eisenberg, B. Feld, V. Fischer, I. Pless, L. Rosenson, R. Yamamoto.
Presented at the Dubna Conference on High Energy Physics, August 1964.

"Gamma Ray Proton Interactions between 0.5 and 4.8 BeV"

H. Crouch, R. Hargraves, B. Kendall, R. Lanou, A. Shapiro, M. Widgoff, G. E. Fischer, A. Brenner, M. Law, E. Ronat, K. Strauch, J. Street, J. Szymanski, J. Teal, V. Fischer, I. Pless, A. Rogers, C. Rogers, P. Bastien, Y. Eisenberg, B. Feld, L. Rosenson, T. Watts, R. Yamamoto, L. Guerriero, G. A. Salandin.

Phys. Rev. Letters 13, No. 21, 636 (1964).

"N^{*}₃₃ and ρ^0 Production by High Energy Photons"

H. Crouch, R. Hargraves, B. Kendall, R. Lanou, A. Shapiro, M. Widgoff, G. Fischer, A. Brenner, M. Law, E. Ronat, K. Strauch, J. Street, J. Szymanski, J. Teal, P. Bastien, Y. Eisenberg, B. Feld, V. Fischer, I. Pless, A. Rogers, C. Rogers, L. Rosenson, T. Watts, R. Yamamoto, L. Guerriero, G. A. Salandin

Phys. Rev. Letters 13, No. 21, 640 (Nov. 23, 1964).

"Multiple Pion and Strange Particle Production in γ -ray-proton Reactions between 0.5 and 4.8 BeV".

Y. Eisenberg, P. Bastien, B. Feld, V. Fischer, I. Pless, A. Rogers, C. Rogers, L. Rosenson, T. Watts, R. Yamamoto, M. Widgoff, A. Shapiro, R. Lanou, F. Bulos, A. Brenner, M. Law, E. Ronat, K. Strauch, J. Street, J. Szymanski, J. Teal, L. Guerriero, G. Salandin, G. Fischer,

Dubna Conference, August 1964.

LINEAR ACCELERATOR

I. Photoneutron Spectrometry

A. $\text{Be}^9(\gamma, n)$

During this period, data was collected on the angular distribution of photo-induced neutron emission from beryllium. Neutron energies were measured by time-of-flight. Neutron energy spectra were measured at 25° , 80° , and 155° . Runs were made at bremsstrahlung end point energies from 5 MeV to 12 MeV in steps of 1.5 MeV. A preliminary analysis of the data indicates some forward to backward asymmetry in the angular distribution of neutrons of approximately 1.2 MeV energy. At present further analysis is being done to test this preliminary result. We are also hoping to obtain more low energy data. If these preliminary results remain valid, we shall have evidence for the existence of an odd parity state mixed with the known even parity state of Be^9 at 2.94 MeV. (W. Bertozzi, S. Kowalski, T. Phillips, C. P. Sargent, and W. Turchinets)

B. Photoneutron Systematics

We have analyzed our photoneutron data, which consist of neutron energy spectra taken at 24° , 75° , and 156° , with bremsstrahlung energies of 15.0 MeV and $13\frac{1}{2}$ MeV for several elements from Bi to Sn. The data have been reduced to the form $W(\theta) = a_0(P_0 + a_1P_1 + a_2P_2)$.

In general, we found that a_1 was small, and since the background and counting rate in the 24° counter were appreciably worse than in the other two, a_1 was taken to be zero. (Bismuth was an exception, where $a_1 \approx 0.05$ at higher neutron energies). Using this assumption, the second Legendre coefficient ($-\frac{a_2}{a_0}$) exhibited the following behavior as a function of energy and mass: $-\frac{a_2}{a_0}$ is small (~ 0) from 0.4 to 2 or 3 MeV. At this energy it begins to increase sharply. Near the closed neutron shells it continues to grow with increasing neutron energy; Pb, Bi, Tl, Hg and Au reach 0.6 - 0.7, and Pr and La reach 0.8. In the region of deformed nuclei, W, Ta, Er, Ho and Sm, it reaches a maximum of about 0.25 and levels off, and perhaps even declines. Below the closed shell at $N=82$, I, Sn, and In increase to 0.3. Then I levels off, Sn increases to 0.4 and In decreases. However, the cross sections are small and the statistics are bad, therefore the behavior above about 4-5 MeV is not really determined.

Some elements, Tl, Pb^{206} , Sm and In, show a backward peaking below 2 MeV. This appears to be a real effect, which increases with decreasing γ -ray energy.

The preliminary experimental results for A_2/A_0 for various elements are shown in Fig. 6.1 as a function of mass number A. Similar measurements by other groups are indicated along with the corresponding references.

Figure 6.1

At present we are analyzing the data to determine the nuclear temperature using runs made at 15.0 MeV and 13.5 MeV to simulate neutron spectra from an equivalent γ -ray beam with an energy of 14.0 MeV and FWHM of 1.75 MeV. (W. Bertozzi, G. Mutchler, C. P. Sargent, and W. Turchinetz)

C. Polarization Measurements

The He^4 neutron polarimeter, described in earlier reports, was used for some polarization measurements the first two weeks in July. As in the past the experimental work was done at the Linear Accelerator Facility of the Rensselaer Polytechnic Institute, Troy, New York. These measurements were primarily extensions of work on oxygen-16 photoneutrons in October, 1963, and were presented at the April meeting of the American Physical Society in Washington, D. C.

For the measurements under discussion the five-inch-diameter, ten-inch-high, stainless steel chamber was used. By matching the thickness of the wavelength-shifter coating on the glass plates of the counter and by very careful attention to the purity of the He-Xe mixture the resolution of a 5.3 MeV alpha source was reduced to about 7%. This is for three axial sources spread over a five inch distance. Tests on the chamber showed that pulse height and resolution of the 5.3 MeV alphas was constant to about 1% over the entire sensitive volume of the counter, radially as well as axially. When the angular resolution of the helium-scattered neutron detectors is combined with the gas-counter resolution an overall value of about 10% for neutron energy resolution is obtained. This is more than two times better than the October runs with resolutions of 20-25%.

On O^{16} runs were made at 45° , at 24, 29.5, and 38 MeV bremsstrahlung end points. The results, with no polarimeter-scattering or target-scattering corrections, are given in Figure 6.2. Some data were also taken on Pb^{208} and carbon, at 45° and 90° to the bremsstrahlung beam.

Figure 6.2

The data have been analyzed for preliminary values of the polarizations and neutron spectra. At present work is being done on scattering corrections to these early results. (W. Bertozzi, F. Hanser, S. Kowalski, C. P. Sargent, and W. Turchinets)

II. Collaboration with University of Saskatchewan

A. $Li^6 (\gamma, np)$

During the past period work continued on the instrumentation for the $Li^6 (\gamma, np)$ experiments to be done at the University of Saskatchewan Linear Accelerator. One experiment involves measurement of the energies of coincident neutron-proton pairs in the photodisintegration (140 MeV bremsstrahlung) of Li^6 , using an array of 16 proton counters and 4 neutron counters.

The electronic circuitry for all the counters has been assembled and tested.

The proton counters (2 1/2" dia. x 5" long plastic scintillators) were tested at the MIT Cyclotron with 25-30 MeV protons obtained from the $Al(\alpha, p)$ reaction, and at the Harvard Cyclotron with elastically scattered protons (50-130 MeV). The energy resolution was around 3.3% for the 30 MeV protons and apparently somewhat worse at the higher energies, although time did not permit a thorough study in the latter case.

Calculations indicated that at forward angles ($\lesssim 45^\circ$) high energy electrons would be detected at a rate comparable with that of the protons. Eight "dE/dx counters" (2" dia., 1/2" thick plastic scintillators mounted on lucite light pipes to be viewed from the side by a second phototube) were constructed. Their resolutions, tested at the MIT cyclotron, was around 3.5%.

An earlier idea -- that of using a thin anthracene crystal optically coupled to the front end of the 5" plastic, with the same phototube viewing both -- was tried at the Harvard Cyclotron and proved to be unsuccessful. The hope was to distinguish protons from electrons by observing the longer decay component from the anthracene, but the large light pulse from the plastic turned out to be too wide.

The fourth neutron detector (three already exist and are in use in (γ, n) experiments at the MIT Linac) was put together and tested. A test of the timing circuitry was made by looking at the annihilation radiation from Na^{22} with this counter and one proton counter. The flight time resolution turned out to be around 4.5 nsec. The causes of this width are not entirely known; we may be able to improve it.

The data will be recorded in a Nuclear Data 4096 channel analyzer, through use of an adapter unit which routes the digital information associated with each event -- proton counter number (1-16), neutron counter number (1-4), proton pulse height (64 channels), neutron flight time (64 channels) -- to the 18 bit memory and advances the address. The data are stored sequentially, one event per channel. A computer program has been written to decode the analyzer output and sort the data in various ways. (J. Bergstron, W. Bertozzi, S. Kowalski, J. Matthews, E. Reinhard, C. P. Sargent, and W. Turchinetz)

B. Photoneutron Spectrometry

The five neutron detectors and their associated circuitry have been assembled. Three of them were available from the photoneutron work at the MIT Linac. Two others have been constructed and are being tested. (W. Bertozzi, S. Kowalski, G. Hurford, J. Matthews, T. Phillips, C. P. Sargent, W. Turchinetz and M. Zombek)

III. Fast Clock

Construction of the fast clock system has been completed. The present version of the system is indicated schematically in Figure 6.3. A brief description of the system operation follows.

Figure 6.3

The 300 Mc strip-line oscillator is operated c.w. The START signals, which initiate the time interval to be measured, phase the oscillator and after a short delay interval (~ 100 ns) open the gate allowing the first FLIP-FLOP to scale the oscillator pulses. FLIP-FLOP No. 1 is driven alternately positive and negative by means of charge storage diodes and the output is a 300 Mc train whose length is a measure of the time interval of interest. The arrival of a STOP pulse closes the gate and determines the end of the measured time interval. The uncertainty in the measured time interval is less than 1.7×10^{-9} sec.

The output of each FLIP-FLOP is amplified, shaped and applied to each successive FLIP-FLOP in the series. The oscillator and first three binary stages were constructed using high speed tunnel diodes (D5066C, in 3858) and strip-line technique. The output of the third binary stage is less than 100 Mc and from this point on the scaling is done directly using available commercial modules. Having set up the various bits of information any sort of core or magnetic tape memory may be used for storage. In the tests made so far, we used a Nuclear Data 4096 channel pulse-height analyzer.

The high frequency parts of the system -- oscillator, gate, three binary stages -- are enclosed in a well-shielded environment relatively stable and free from outside interference. Measurements of frequency stability relative to a crystal controlled oscillator, indicates a stability of $(2-4) \times 10^{-4}$ (over a 12 hour period) at 300 Mc. It is believed that some of the drift is due to temperature variations, so we plan to include a temperature controlling device in the system.

Operational measurements on the complete system to date have included stability measurements on time intervals determined by a cable and "white spectrum" measurements. In the latter case the oscillator is phased at several kc rates using an external trigger source and the STOP signal is provided by a phototube viewing a radioactive source. Correcting for dead time losses, the differential nonlinearity has been found to be less than 1% for 1.7 nsec channel widths and a maximum time interval of 6 μ sec.

We plan to make use of the above system shortly in the measurement of high resolution neutron energy spectra using a 50 meter flight path at the University of Saskatchewan. (I. Hayashi, S. Kowalski)

ADDRESSES

- W. Turchinets, "Some recent studies of photonuetron reactions", McGill University, November, 1964.
- W. Bertozzi, P. T. Demos, S. Kowalski, C. P. Sargent, W. Turchinets, R. Fullwood and J. Russell, "Polarization of Photoneutrons from D, O¹⁶ and C¹²", Paris Conference, Series B:4d, July 1964.

HIGH ENERGY ACCELERATOR PHYSICS GROUP

I. Photoproduction of Mesons at CEA

A. General

The data taken in the fall of 1963 were published in the Physical Review letters^{1,2} in June. In summary, the cross section for $\gamma + p \rightarrow \pi^0 + p$ as measured at 90 degrees in the center of mass system shows a fast drop as a function of energy; this was also true at $\theta_\pi = 60^\circ$ although the drop was not as sharp. Superimposed on this behavior are two bumps in the differential cross section, one at a bombarding energy of roughly 3.6 BeV corresponding to a center of mass total energy, M^* , of 2.7 BeV. There was also a bump on the 60° data corresponding to $M^*=2.5$ BeV. We also found at 3.6 BeV a yield of coincidences between low energy gamma rays and protons, which were interpreted as $\eta + p$ production.

We ran again this summer with the 48-counter shower detector; thus we were able to detect the π^0 sometimes when both decay γ rays struck the counter, and also when one γ ray struck since its energy is correlated with the angle it makes to the π^0 line of flight. We have now finished the preliminary analysis of the 90° data. The single and double gamma ray data were treated separately. They were analyzed by computer to take into account the efficiency of the gamma ray counter. The resulting preliminary cross sections are shown in Figure 7.1. It is gratifying to note that both the one- and two-gamma ray events give roughly the same cross section. Points from our published data are also shown; there is a clear disagreement, for which we have not ascertained the explanation.

Figure 7.1

π^0 Photoproduction - 90° c. of m.

We are proceeding with data analysis to see if the remaining $\gamma\gamma$ coincidences, which are not consistent with π^0 production, can be analyzed in terms of a simple mechanism such as η production. (R. Alvarez, R. Fessel, V. Elings, D. Garelick, S. Homma, R. Lewis, W. Lobar, D. Luckey, L. Osborne, J. Uglum, Z. Bar-Yam, W. Kern)

B. On-Line Computer

The data were taken during the summer with the Linc computer attached to the equipment. We had enough software ready so that the computer could present histograms giving momentum plots for the π^+ , protons, and protons with gamma rays in coincidence. It could give counting rate profiles of the hodoscope counters, which served as a good running check on their behavior. We have extended and written programs so that the Linc can bring in information on machine energy, magnet currents, etc.

With the advent of wire chambers and other hodoscopes we envisage the need for greater information processing ability. We have initiated a program to connect the LNS IBM 7044 to our experiment through telephone lines. The computer would operate on a simple time sharing program. The computer systems program has been examined and the project appears entirely feasible. (R. Fessel, D. Garelick, L. Osborne)

C. Wire Chambers

This summer we designed and did preliminary tests on some simple wire chambers. This led to the setting of specifications for core stacks and reading electronics; these have now been ordered and have just been delivered. We plan to read the cores by the coincident current technique which will make the reading electronics considerably cheaper.

The parts for the wire chambers have arrived and we are now assembling a first chamber. (D. Bellenger, D. Garelick, S. Homma)

D. Magnetic Spectrometer

We have installed an arm on the rear end of the spectrometer platform, which will allow bending in a vertical plane by raising the final detecting hodoscopes. The equipment operates satisfactorily. We hope to be back in operation in January. (J. Davin)

E. π^0 and η^0 Production at Forward Angles

The lead glass counters to be used in this experiment are the ones used in the π^0 experiment mentioned above. They work satisfactorily. A mount to hold these counters in a double stack to measure $\eta^0 \rightarrow 2\gamma$ and $\pi^0 \rightarrow 2\gamma$ is now in the process of design. (D. Luckey, R. Lewis)

F. Meson Production at Back Angles

An experiment has been designed to measure π^+ photoproduction cross sections at very large angles. Basically, a bending magnet and two spark chambers, together with scintillation

trigger counters, are used to detect and momentum-analyze the pion. We hope to measure the excitation function from about 1100 to 3000 MeV at about 165° c. of m. and enough of the backward hemisphere c. of m. angular distribution at energies corresponding to the formation of intermediate state pion-nucleon resonances to gain some information about their spins. It appears to be possible with essentially the same apparatus to distinguish π^+ and K^+ mesons at momenta corresponding to a c. of m. production angle of about 165° and photon energies from roughly 2 to 3 BeV. One of the spark chamber optical systems has been constructed and work is currently being done on the design and construction of the rest of the detection apparatus. (R. Alvarez, G. Cooperstein)

II. Elastic π^-p Charge Exchange

In a recent spark chamber experiment on pion charge exchange at BNL-AGS from 2.8 to 18 GeV/c³, we measured the angular distributions of the π^0 's near 0° to see both the shape of the "diffraction" peak and to extrapolate to $\frac{d\sigma}{d\Omega}(0^\circ)_{\text{ch.exch.}}$.

At two momenta, namely 6 and 10 GeV/c, we had the spark chamber close enough to our liquid hydrogen target to have good efficiency for comparatively large scattering angles ($\theta^* \gtrsim 40^\circ$). Angular distributions at these two momenta are shown in Figures 7.2 and 7.3. In these distributions, the π^0 direction is taken to be the bisector (θ_h^*) of the two gammas from the $\pi^0 \rightarrow 2\gamma$ decay. The error bars on the histogram give the statistical counting errors only, after taking into account the target empty subtraction.

Figure 7.2

Angular Distribution in C.M. of $\pi^- + p \rightarrow \pi^0 + n$ at 6 GeV/c.

Figure 7.3

Angular Distribution in C.M. of $\pi^- + p \rightarrow \pi^0 + n$ at
10 GeV/c.

Since the statistics are much better for the 10 GeV/c data, we will look at it in more detail. Figure 7.4 is identical with Figure 7.3; except that the scale has been expanded by a factor of 50 to show the structure more clearly. The angular distribution shows a minimum around $\cos \theta_b^* = 0.93$ and then a second, broad maximum. Due to the large statistical errors, however, we can fit the data almost as well by a diffraction peak plus a flat distribution between $\cos \theta_b^* = 0.94$ and 0.82. A similar strong discontinuity of slope, also giving the appearance of a minimum, is evident in the 6 GeV/c data around $\cos \theta_b^* = 0.88$.

Figure 7.4

Angular Distribution in C.M. of $\pi^- + p \rightarrow \pi^0 + n$ at
10 GeV/c with expanded scale.

Figure 7.5 shows the same 10 GeV/c data plotted as a function of t , the 4-momentum transfer squared. The shape of this curve, especially near $t=0$ where it flattens off considerably, is quite different from the nearly straight line observed for $\pi^\pm p$ scattering. The 6.0 GeV/c data, with the same flattening of slope near $t=0$ and a minimum near the same value $t=-0.6$.

Figure 7.5

Angular Distribution in C.M. of $\pi^- + p \rightarrow \pi^0 + n$ at 10 GeV/c but with square of 4-momentum transfer instead of angle as abscissa.

The flat differential cross section at the larger t values might be due to an inelastic contamination of our elastic events if such inelastics are present in our data and have the required flat shape. The inelastic π^0 s must have larger $\gamma=\gamma$ opening angles than the elastics, so if we accept only events with opening angles between 1.00 and 1.05 times the minimum allowed for elastic π^0 s, we will eliminate most inelastics. As we increase our acceptance criteria to 1.10, and 1.15, and 1.20 times the minimum angle, we accept a larger and larger fraction of inelastic events. However, the differential cross section, within statistics, is the same for all four of these opening angle cuts, and this indicates that the inelastic events are not the cause of the flat distribution. We are using the data with the cut at 1.15.

Figure 7.6 gives the "mass spectrum" of the $\gamma\text{-}\gamma$ system at 10 GeV/c. This is not a true mass spectrum; it is obtained from the $\gamma\text{-}\gamma$ decay opening angle in the $X^0 \rightarrow 2\gamma$ decay by assuming that each measured $\gamma\text{-}\gamma$ angle is at its minimum allowed value. Using $\theta_{\gamma\gamma\min} = 2M_{\gamma\gamma}/P_{X^0}$, where $M_{\gamma\gamma}$ is the mass of the $\gamma\text{-}\gamma$ system, and P_{X^0} is the X^0 momentum in the $\pi^- p$ C.M. system, we can

change from a $\theta_{\gamma\gamma}$ to a $M_{\gamma\gamma}$ distribution. Thus at a $M_{\gamma\gamma}$ corresponding to a particle or resonance which decays into 2 γ s, we should see a peak which rises sharply on the low mass side, and falls more slowly on the high mass side.

Figure 7.6

Mass Spectrum of $\gamma\text{-}\gamma$ system at
10 GeV / c

The only statistically significant masses observed are the π^0 and the η^0 mesons. In particular, we see no evidence that the 960 MeV meson decays strongly into 2 γ s. We can observe $\gamma\text{-}\gamma$ opening angles corresponding to $\gamma\text{-}\gamma$ masses up to ~ 1700 MeV, at which point our detection efficiency falls off rapidly, but we see no resonant behavior in the 1000 to 1700 MeV region.

(I. Mannelli (Pisa), M. Wahlig, and L. Sodickson)

III. Dipion Mass Distribution in $\pi^- + p \rightarrow (2\pi^0) + n$

Figure 7.7 ($p_\pi=10$ GeV) shows the f^0 peak⁴ and also shows an enhancement in the low mass region. We are studying this distribution as a function of opening angle cut to verify that the enhancement is real.

Figure 7.7

Approximate $2\pi^0$ Mass Distribution in $\pi^- + p \rightarrow (2\pi^0) + n$.

(M. Wahlig, E. Shibata, and D. Gordon)

IV. MASCOT

This large-solid-angle and high-Z sparkchamber and scintillator array is assembled and is now being tested at BNL. (A. Buffington, O. Fackler, S. Smith, M. Strovink, and C. Ward)

V. Λ^0 Gyromagnetic Ratio

All of the data film has been scanned and measured. We are repeating measurements on several samples to verify our procedure. Corrections are being calculated for some small systematic errors. The preliminary result is $\mu_{\Lambda} = -0.6 \pm 0.25$ nuclear magnetons (statistical error only) based on about half the data. This value is compatible with the SU(3) prediction, $\mu_{\Lambda} = -0.95$ n.m., but disagrees with several alternative theories. (D.A. Hill, K. K. Li, C. Stassis; E. Jenkins, T. Kycia, H. Ruderman (BNL))

VI. Search for Point Sources of High-Energy Cosmic Gamma Rays

Progress has been slow because of bad observing conditions and frequent mirror fogging. Probably the winter months will be more productive. We have made further tests of the detector

and found an arrangement which gives a shower counting rate of 100 cpm with good angular resolution. (G. Fazio and H. Helmkin (Smithsonian Astrophysical Observatory), and D. A. Hill)

VII. The Photoproduction of Wide Angle Muon Pairs

The analysis of the data from the measurement of the photoproduction of μ -pairs from carbon carried out in collaboration with physicists from Northeastern University was finished in the spring of 1964. The results were published in Phys. Rev. Letters 26, 739 (1964). Measurements were made of the differential cross sections for the following kinematic intervals:

$$1.8 \text{ BeV} \leq E_1, E_2 \leq 2.4 \text{ BeV}$$

$$4.5^\circ \leq \theta_1, \theta_2 \leq 11.5^\circ$$

$$-15^\circ \leq \phi_1 \leq 15^\circ \quad 165^\circ \leq \phi_2 \leq 195^\circ$$

where E_1, E_2 ; θ_1, θ_2 ; and ϕ_1 and ϕ_2 are the total energies, polar angles, and azimuthal angles for the members of the pair. The cross sections for $\theta_1 \approx \theta_2$ were used to test the validity of the quantum electrodynamic description of the muon propagator at four momentum transfers q to the virtual muon up to about $3f^{-1}$. A least squares analysis of the ratio of the experimental cross section to the theoretical cross section as a function of q^2 gives

$$\frac{\sigma_{\text{exp}}}{\sigma_{\text{theor}}} = (1.18 \pm 0.15) [1 - (0.011 \pm 0.021) |q^2|]$$

An analysis of the results in terms of a breakdown model⁵ for the muon propagator sets a limit of $\frac{1}{\Lambda} < 0.16 f$ with 95% confidence for a possible cut-off distance.

These measurements have recently been repeated with improved apparatus and greater beam intensities. The amount of data collected was about a factor of 40 times greater than that of our first set of measurements. We thus expect to be able to improve considerably the limits placed by our first experiment on a possible breakdown in the theory. The analysis of the data from the second set of measurements is now being carried out. (J. I. Friedman, G. Glass)

A. Elastic Electron-Deuteron Scattering

All of the equipment which is to be used in the experiment has been constructed and tested. We have had a number of preliminary runs and have taken data at momentum transfers q in the range of 4.0 - $4.2 f^{-1}$. This data is now being analyzed and some consequent improvements in the equipment have been made and tested. We are now continuing our measurements. We expect to reach values of q up to nearly $5 f^{-1}$. (J. I. Friedman, H. W. Kendall, L. Van Speybroeck, G. Hartman, J. Elias)

B. The Two Nucleon Correlation Function

All of the primary data for this experiment have been analyzed with the use of our radiative correction computer programs. The normalizations have been rechecked experimentally at the linear accelerator at Orsay this past fall and these data, combined with our earlier data, should result in a completed program early this winter. The experiment will then be written up for publication. (Experiment in collaboration with Dr. Didier Isabelle, Orsay) (J. I. Friedman, H. W. Kendall)

C. Phase Shift Analysis of Inelastic High Energy Electron Scattering

At this writing an initial version of the entire program has been tested and debugged. The source deck contains about 7500 cards. Square well and delta function nuclear charge densities were employed. A set of subroutines which will calculate the required potentials for arbitrary point specified densities has been written and nearly debugged. When these are introduced into the main program we can begin comparisons with experimental scattering. (J. I. Friedman, H. W. Kendall)

REFERENCES

1. R. Alvarez, Z. Bar-Yam, R. Fessel, W. Kern, D. Luckey, L. Osborne and S. Tazzari, "Photoproduction of π Mesons between 0.9 and 4.0 BeV", Phys. Rev. Letters 12, no. 25, (June 22, 1964).
2. R. Alvarez, Z. Bar-Yam, R. Fessel, W. Kern, D. Luckey, L. Osborne and S. Tazzari, "Evidence for Two New Resonances in the π -Nucleon System", Phys. Rev. Letters, 12, no. 25, (June 22, 1964).
3. M. A. Wahlig, I. Mannelli, L. Sodickson, O. Fackler, C. Ward, T. Kan, and E. Shibata, Phys. Rev. Letters 13, 103 (1964), and May 1964 Progress Report.
4. L. Sodickson, I. Mannelli, D. Frisch, O. Fackler, and M. Wahlig, "Observation of $f^0 \rightarrow 2\pi^0$ ", Phys. Rev. Letters 133 (Feb. 10, 1964).
5. S. D. Drell, Ann. Phys. 4, 75 (1958).

PUBLICATIONS

- R. Alvarez, A. Bar-Yam, R. Fessel (CEA), W. Kern, D. Luckey, L. Osborne, and S. Tazzari, "Photoproduction of π Mesons between 0.9 and 4.0 BeV", Phys. Rev. Letters 12, No. 25, (June 22, 1964).
- R. Alvarez, Z. Bar-Yam, R. Fessel (CEA), W. Kern, D. Luckey, L. Osborne, and S. Tazzari "Evidence for Two New Resonances in the π -Nucleon System", Phys. Rev. Letters, 12, No. 25 (June 22, 1964).
- O. Fackler, D. Frisch, T. Kan, I. Mannelli, E. Shibata, L. Sodickson, M. Wahlig, and C. Ward, "Charge Exchange Scattering $\pi^- + p \rightarrow \pi^0 + n$ in the Forward Direction at 6.0 and 10 GeV/c " presented by L. Osborne at the Dubna Conference, USSR, August 1964.

- O. Fackler, T. Kan, I. Mannelli, E. Shibata, L. Sodickson, M. Wahlig, and C. Ward, "Structure in Forward Direction π^- -p Charge Exchange Scattering in the 3 GeV/c Region", Phys. Rev. Letters 13, No. 3 (July 20, 1964).
- G. Cooperstein and D. Garelick, "Photoproduction of Charged Mesons from Free Nucleons for Bombarding Gamma-Ray Energies near 275 MeV", Phys. Rev. 136, No. 1B, B201, (October 13, 1964).

ONR GENERATOR GROUP

I. Introduction

In the past six-month period, a number of exposures have been made with the multiple-gap spectrograph, notably in the $1f_{7/2}, 2p_{3/2}$ shells. Some of the data are being analyzed at MIT; other parts at the Institutes with which we collaborate, such as Yale University, University of Mexico, University of Chile, University of Bergen (Norway), and Max-Planck Institut in Heidelberg, Germany.

Unusually long delays in scheduled experiments have been encountered during this period, mostly because of breakdowns of terminal equipment in the ONR generator.

II. Studies of Energy Levels in the $f_{7/2}$ Shell

The following seven exposures have been completed and are presently being scanned and analyzed as part of the continuing study of nuclei in the $(f_{7/2})$ -shell region.

A. The $\text{Ca}^{42}(\text{d}, \text{p})\text{Ca}^{43}$ Reaction

An exposure of 2500 μC at an incident deuteron energy of 7.00 MeV was made, using a thin 93 percent enriched Ca^{42} target. Some of the plates from this experiment were scanned automatically by Cyclops.

B. The $\text{Ca}^{42}(\text{d}, \text{d}')\text{Ca}^{42}$ Reaction

An exposure of 2500 μC at an incident deuteron energy of 7.50 MeV was made. Thin 1.5-mil aluminum foils were used as absorbers in front of the 50-micron NTA plates and were found to help considerably in the discrimination of deuteron tracks from proton tracks. Elastic-scattering data were taken at this energy with the use of the new Rutherford slit system discussed elsewhere.

C. The $\text{Ca}^{44}(\text{d}, \text{d}')\text{Ca}^{44}$ Reaction

An exposure of 2500 μC at an incident deuteron energy of 7.50 MeV was made. The target was enriched to 98 percent in Ca^{44} . Elastic-scattering data at 7.50 MeV and target-thickness measurements at 3.0 MeV were also made.

D. The $\text{Ca}^{42}(\text{d}, \alpha)\text{K}^{40}$ Reaction

An exposure of about 3500 μC at an incident deuteron energy of 7.50 MeV was made. Ilford K0 nuclear-track plates were used to detect the alpha particles.

E. The $Ti^{48}(p, p')Ti^{48}$ Reaction

An exposure of 2000 μC at an incident deuteron energy of 7.00 MeV has been carried out. The preliminary results of this exposure are to be presented at the American Physical Society meeting in New York in January 1965.

F. The $Ti^{48, 50}(d, d')Ti^{48, 50}$ Reactions

Exposures of 3000 μC at an incident energy of 6.00 MeV were made. This work is being done in conjunction with Professor C. K. Böckelman of Yale's Accelerator Laboratory.

Preliminary experiments on the $Sn^{118}(d, p)Sn^{119}$ reaction have also been carried out at 7.5-MeV bombarding energy. (T. A. Belote, W. E. Dorenbusch, and O. Hansen)

III. Mercury-vapor Target

A mercury-vapor target has been developed for use in nuclear reaction studies. The principle of operation of the target consists of passing a dense vapor stream from a heated crucible with mercury across the path of the bombarding particles. Excess vapor is then condensed on a water-cooled hood and circulated back into the boiler. Preliminary tests indicate that an effective target thickness of 2 $\mu g/cm^2$ of Hg can be achieved with a recirculation efficiency of better than 90 percent. A series of water-cooled and liquid nitrogen-cooled baffles collect most of the mercury that escapes recirculation. The advantages of such a target over conventional thin-film and gas-cell targets are (a) the contaminant spectrum from materials in backings is eliminated; (b) the energy width of spectrum peaks from the presence of backings and windows is removed; and (c) unlimited beam currents can be used without fear of damaging the target. (E. Cosman and H. A. Enge)

IV. Combined Magnetic Quadrupole Lens and Deflector Magnet System

The multiple-gap spectrograph has one large (1 1/2") gap at $\theta = 45^\circ$ intended to be used in conjunction with a quadrupole lens as a spectrometer¹ for work where a large solid angle is required. Preliminary calculations indicate that an increase by a factor of about 50 in solid angle over the existing spectrograph gaps may be obtained with such an arrangement without appreciable loss of resolving power.

For a given transition, a slight variation of energy versus reaction angle occurs ("kinematic effect", also called Doppler shift). It has been found that this effect can be corrected for (and resolution restored) by a small rotation of the quadrupole lens about its axis. The required angle of rotation is a function of the masses and kinetic energies of the incident and emitted particles, the mass of the residual nucleus, and the reaction angle. For a set of typical

lens-deflector parameters and the reaction

$\text{Cr}^{52}(\text{d}, \text{p})\text{Cr}^{53}$ ground state

$Q = 5.725 \text{ MeV}$

$E_{\text{deuteron}} = 7.500 \text{ MeV},$

a rotation angle of approximately 3 degrees is required.

Plans are now being made for construction of a magnetic quadrupole lens to be used in conjunction with the 45-degree gap and of a scintillation spectrometer for gamma rays, to be mounted close to the target. This system will be used for measuring gamma-ray branching from specific levels fed by a charged-particle reaction, and possibly later may be used for correlation measurements between charged particles and gamma rays. (D. L. Smith and H. A. Enge)

V. He^3 Elastic Scattering

In the immediate future it is planned to run a series of He^3 elastic-scattering experiments. Mg^{24} , S^{32} , and Ca^{40} targets will be exposed to He^{3++} beams at 10, 12, and 14 MeV; or if it is found impossible to operate with a doubly ionized beam, a singly charged beam with the highest energies feasible will be utilized. The angular distributions will be analyzed with the ABACUS code to obtain "best fit" optical-model parameters. If possible, the angular distributions from C^{12} and O^{16} in the target backings and from heavier contaminants in the targets will also be analyzed.

In connection with these experiments, a "Rutherford slit assembly" has been designed and constructed. This assembly fits in the forward quadrant of the spectrograph in front of the existing " α slits". It masks the α slits with additional slits whose openings vary as $\sin^4(\theta/2)$ from 0.75" at 90° (89 gap) to 0.0135" at 30° . From 30° to 7° , the slit opening is kept constant, since at slit openings less than 10 mils excessive multiple scattering from the slits would be expected. The masking slits can be moved out of position from the exterior of the spectrograph so that the scattered beam is limited by the normal " α slits". The Rutherford slits have been calibrated using the results from 5-MeV elastic scattering of deuterons on Sn^{118} . The angular distribution from this experiment is assumed to be pure Rutherford, and the solid angle for each gap has been calculated from the observed angular distributions. (Helen J. Young)

VI. Reaction Studies from the Chromium Isotopes

A summary of the work in progress on the isotopes of chromium was given in the May 1964 Progress Report (p. 94). Since then the following progress has been made.

A. The $\text{Cr}^{50}(\text{d}, \text{p})\text{Cr}^{51}$ Reaction

The majority of the nuclear emulsions exposed in the multiple-gap spectrograph are in the process of being scanned at the Max-Planck Institut in Heidelberg, Germany. No data are yet available for analysis.

B. The $\text{Cr}^{53}(\text{d}, \text{d}')\text{Cr}^{53}$ Reaction

The proposed experiment to study the angular distribution of inelastic deuterons from 7.5-MeV deuteron bombardment of a Cr^{53} target was performed. Eastman NTB plates were used, and, since no absorber was placed in front of the emulsions, there was greater difficulty in distinguishing protons from deuterons in this energy range than has been the experience with NTA emulsions. However, in the region of low proton level density from the $\text{Cr}^{53}(\text{d}, \text{p})\text{Cr}^{54}$ reaction, it was possible to identify inelastic groups associated with the $\text{Cr}^{53}(\text{d}, \text{d}')\text{Cr}^{53}$ reaction. Data on the angular distribution of these levels in Cr^{53} up to about 2.0 MeV are being assimilated.

C. The $\text{Cr}^{54}(\text{d}, \text{p})\text{Cr}^{55}$ Reaction

A twenty-four-angle exposure was made of the proton spectra from bombardment of enriched Cr^{54} with 7.5-MeV deuterons. The scanning of the majority of these plates has been carried out in Heidelberg and is nearly completed. Angular-distribution data have been assimilated for a number of levels up to 3.0 MeV in Cr^{55} , and a DWBA analysis is under way.

With the completion of an analysis of an earlier 90-degree single-gap proton spectrum obtained from bombardment of Cr^{54} with 6.5-MeV deuterons, precise measurements of a number of levels in Cr^{55} have been made. Excitation energies of the most prominent levels are shown in Table 8.I. Evidence for weaker proton groups above about 1.5 MeV is seen from the 90-degree single-gap exposure, as well as at other angles in the multiple-gap run. A more complete listing will be made on completion of the angular-distribution study. The values in Table 8.I are in good agreement with those of the corresponding levels from other recent measurements.²

Table 8.I
Single-gap Measurements on $\text{Cr}^{54}(\text{d}, \text{p})\text{Cr}^{55}$ Reaction;
 $Q=4.027 \pm 0.008$ MeV.

Excitation Energies in MeV						
0.242	1.212	2.275	3.154	3.835	4.172	4.662
0.517	1.475	2.607	3.300	4.035	4.390	4.739
0.564	2.020	2.895	3.354	4.046	4.423	
0.882	2.091	3.023	3.527	4.151	4.467	

(A. Sperduto, C. A. Wiedner, N. M. Rao, and G. Nair)

VII. The Automatic Plate Scanner

A new illumination system has been installed, and this has resulted in improved performance of the scanner, particularly for tracks with low-grain density. The system consists of two high-pressure mercury arc lamps with reflectors and heat filters.

A spiral-disk scanning unit, mentioned briefly in the last Progress Report, is now being installed to replace the scanning tape. The new unit will allow us to scan a zone of 10 mm instead of 4 mm, and we also hope that the resolution will be improved, thereby reducing the counting losses when the track density is high. (P. Hanssen and H. A. Enge)

VIII. Ion Optics Calculations

During this period the 7090 ray-tracing program³ was converted into a subroutine capable of tracing a ray through a system of "n" magnets with fields symmetric about the median plane within the magnet and with realistic fringing fields at the entrance and exit of each magnet. A further improvement, included as an option, is the ability to extract all first- and second-order focusing coefficients for such an array.

Another computer program is being developed to obtain the approximate numerical solution of elliptic partial differential equations in two dimensions for determining the field in a magnet with arbitrary boundary and coil configurations. It will be used to ascertain the proper physical coil arrangements for the proposed 185-degree analyzing magnet that will give the desired analytic field expressions in the median plane. The analytic expressions for the field in the median plane will then be used to determine the field off the median plane, using Taylor expansions in the ray-tracing program. In this way all focusing coefficients through second order will be obtained so that a more thorough study of the design can be made. (J. E. Spencer and H. A. Enge)

IX. DWBA Calculations and Optical-Model Analysis of Elastic-Scattering Data

DWBA calculations were made using the computer program "Julie" on a number of nuclear reactions, namely: $\text{Ne}^{20}(\text{d}, \text{p})\text{Ne}^{21}$; $\text{Ne}^{22}(\text{d}, \text{p})\text{Ne}^{23}$; $\text{Ca}^{46}(\text{d}, \text{p})\text{Ca}^{47}$; $\text{Ti}^{46}(\text{d}, \text{p})\text{Ti}^{47}$; $\text{V}^{51}(\text{d}, \text{p})\text{V}^{52}$; $\text{Cr}^{52}(\text{d}, \text{p})\text{Cr}^{53}$; $\text{Ca}^{40}(\text{He}^3, \text{d})\text{Sc}^{41}$; and $\text{Ca}^{40}(\text{He}^3, \alpha)\text{Ca}^{39}$. When the elastic-scattering data were available, an optical-model analysis was made in order to obtain the optical-model parameters. The computer program "ABACUS" was used in this analysis to fit the elastically scattered deuterons on Ca^{46} at 7 MeV, on Cr^{52} at 7.5 MeV, and to fit the He^3 ions scattered at 12 MeV on Ca^{40} . (H. Y. Chen)

X. The $\text{Ne}^{20}(\text{d}, \text{p})\text{Ne}^{21}$ and $\text{Ne}^{22}(\text{d}, \text{p})\text{Ne}^{23}$ Reactions

Since our last report the distorted-wave analyses of the proton angular distributions from these reactions has been completed. This has enabled spin-parity assignments to be made for eleven states in Ne^{21} and seven in Ne^{23} , and these are listed in Tables 8. II and 8. III.

Table 8. II
Level Spins and Parities in Ne^{21}

Level Number	E_x (MeV) (a)	ℓ_n (b)	J^π (b)	Previous Work
0	0			$3/2^+$
1	0.349	2	$3/2^+, 5/2^+$	$5/2^+$ (c, d)
2	1.750			$7/2^+$ (d)
3	2.800	0	$1/2^+$	$1/2^+$ (c, d)
4	2.870			$\geq 5/2^+$ (d)
5	3.666			$3/2$ (d)
6	3.737			$5/2^+$ (d)
7	3.889			
8	4.435			
9	4.528	2	$3/2^+, 5/2^+$	$3/2^+, 5/2^+$ (c)
10	4.685	2	$3/2^+, 5/2^+$	
11	4.729	(1)	$(1/2^-, 3/2^-)$	$3/2^-$ (d)
12	5.388	2	$3/2^+, 5/2^+$	
13	5.434			$(3/2^+, 5/2^+)$ (c)
14	5.550	2	$3/2^+, 5/2^+$	
15	5.632			$(3/2^+, 5/2^+)$ (c)
16	5.694	(0)	$(1/2^+)$	
17	5.777	(1)	$(1/2^-, 3/2^-)$	$(1/2^-, 3/2^-)$ (c)
18	5.820	(2)	$(3/2^+, 5/2^+)$	
19	5.997	(2)	$(3/2^+, 5/2^+)$	

Table 8. III
Level Spins and Parities in Ne²³

0	0	2	$3/2^+, 5/2^+$	$3/2^+, 5/2^+(c)$
1	1.018	0	$1/2^+$	$1/2^+(c)$
2	1.703			
3	1.826			
4	2.314	2	$3/2^+, 5/2^+$	
5	2.520			
6	3.218	0	$1/2^+$	
7	3.433	2	$3/2^+, 5/2^+$	
8	3.836	1	$1/2^-, 3/2^-$	
9	3.988	2	$3/2^+, 5/2^+$	

^a J. M. Freeman, Phys. Rev. 120, 1436 (1960).

^b Present work.

^c H. B. Burrows, T. S. Green, S. Hinds, and R. Middleton, Proc. Phys. Soc., A69, 310 (1956).

^d A. J. Howard, D. A. Bromley, and E. K. Warburton, Phys. Rev. 137, 32 (1965).

Although the distributions for the 4.73-, 5.69-, and 5.78-MeV states in Ne²¹ are strongly forward peaked, there remains some doubt concerning the ℓ_n values in the transitions, even though the indicated values are considered most probable. Because of the negative Q-values for these states, the distinction between $\ell_n=0$ and $\ell_n=1$ is not clear at this incident deuteron energy of 7.0 MeV. These states are now being studied at a lower bombarding energy of 4.0 MeV where the distinction between $\ell_n=0$ and 1 should be much clearer.

In both Ne²¹ and Ne²³, attempts have been made to describe the observed level schemes in terms of the strong-coupling collective model of Nilsson. Thus, in Ne²¹ rotational bands may be expected based on Nilsson orbit 6 with $K=3/2^+$, orbit 5 with $K=5/2^+$, orbits 9 and 11 with $K=1/2^+$, and orbit 14 with $K=1/2^-$. The head of the $K=3/2^+$ band can here be identified with the ground state of Ne²¹. Since the heads of the two $K=1/2^+$ bands are the only two states that may be formed by $\ell_n=0$ stripping, then the lowest of these, corresponding to orbit 9, can be identified with the 2.80-MeV level and the other $K=1/2^+$ head, probably to the 5.69-MeV level.

The state at 3.73 MeV is known from the work of Howard, Bromley and Warburton to be $5/2^+$ which may therefore qualify as a candidate for the $K=5/2^+$ band head. However, we observe no sign of stripping to this state, and it may be that mixing of the $K=1/2^+$, $3/2^+$, and $5/2^+$ bands in this case dilutes the stripping reduced width to negligible proportions. We have not yet carried out any calculations of the stripping reduced widths with band mixing included. It does not appear necessary to invoke band mixing to explain the nonstripping behavior of the ground state of Ne²¹. Even without this, the Nilsson model predicts the $\ell_n=2$ transition strength to the ground

state to be only about 4 percent of that to the first excited state, and this is not strongly dependent upon the deformation.

The first negative-parity state that we observe is probably that at 4.73 MeV, and this is consistent with the $3/2^-$ assignment of Howard et al. to this state. The only other $\ell_n=1$ transition is probably that observed to the 5.77-MeV level, which could therefore correspond to the $K=1/2^-$ band head.

In Ne^{23} , the 1.70-, 1.83-, and 2.52-MeV states were only weakly excited, the remaining states being well described by a stripping mechanism. The only states whose properties were already well known were the ground and first excited states, and within the framework of the Nilsson model these have usually been assumed to be the $K=5/2^+$ and $K=1/2^+$ band heads, respectively, corresponding to orbits 5 and 9.

The higher intrinsic configurations are expected to have $K=1/2^+$ and $K=1/2^-$, corresponding to orbits 11 and 14, respectively. The $K=1/2^+$ band head can here be identified with the 3.218-MeV level. The first state that can be assigned negative parity with certainty is that of 3.836 MeV. This state can therefore be $1/2^-$ or $3/2^-$, and if it follows the general pattern observed in Ne^{21} , Mg^{25} , and Al^{25} , the $3/2^-$ level lies lower than the $1/2^-$ band head. (D. J. Pullen)

XI. The $\text{N}^{15}(\text{d}, \text{p})\text{N}^{16}$ and $\text{O}^{18}(\text{d}, \alpha)\text{N}^{16}$ Reactions

These reactions have recently been studied at an incident energy of 7.5 MeV using a rotating gas cell. The nitrogen gas was enriched to 97.8 percent in N^{15} and the oxygen to 97.4 percent in O^{18} . Angular distributions were measured from 7.5 to 172.5 degrees using the multiple-gap magnetic spectrograph. At present, the scanning in the region of the unbound states is still incomplete.

The $\text{N}^{15}(\text{d}, \text{p})\text{N}^{16}$ reaction has been previously studied in this Laboratory, and the proton angular distributions leading to the ground and first three excited states in N^{16} have been measured. In the present work, angular distributions are being measured for the 3.34-, 3.50-, and 3.96-MeV states. Of these, only the 3.96-MeV state appears to show any marked forward peaking.

In the $\text{O}^{18}(\text{d}, \alpha)\text{N}^{16}$ reaction, angular distributions of alpha particles leading only to the ground and first three excited states are complete, and these exhibit strong forward and backward peaking. All the known states of N^{16} up to 5.53 MeV appear to be equally strongly excited by this reaction. This may indicate the principal mode of formation of these states to be via compound nucleus at this incident energy. (D. J. Pullen and E. Chalbaud)

REFERENCES

1. H. A. Enge, "Combined Magnetic Spectrograph and Spectrometer", Rev. Sci. Instr. 29, 885 (1958).
2. J. H. Bjerregaard, P. F. Dahl, O. Hansen, and G. Sidenius, Nuclear Phys. 51, 641 (1964); and H. H. Duham, private communication.
3. Progress Report, Laboratory for Nuclear Science, MIT, November 1961, p. 130.

PUBLICATIONS

- E. Kashy, A. Sperduto, H. A. Enge, and W. W. Buechner, "Shell-Model States in Ca^{49} ", Phys. Rev. 135, B865 (1964).
- H. A. Enge, "Magnetic Spectrographs and Beam Analyzers," Proc. of Third International Accelerator Conference, Nucl. Instr. and Methods 28, 119 (1964).
- P. D. Barnes and C. K. Bockelman (Yale), O. Hansen (Copenhagen), and A. Sperduto, "Inelastic Deuteron Scattering and (d, p) Reactions from Isotopes of Ti(I): $\text{Ti}^{50}(\text{d}, \text{p})\text{Ti}^{51}$ ", Phys. Rev. 136, B438 (1964).

THESES

- D. M. Sheppard, "An Investigation of He^3 -Induced Reactions on Ca^{40} at 12 MeV", Ph.D. September 1964.
- M. M. Castro, "A Study of the $\text{Cl}^{35}(\text{He}^3, \text{p})\text{Ar}^{37}$ Reaction", M.S.
- K. A-A. Hamza: "Analysis of Experimental Data from the $\text{V}^{51}(\text{d}, \text{p})\text{V}^{52}$ Reactions," M.S.
- K. M. I. Abdo, "A Study of the Reaction $\text{Cl}^{35}(\text{He}^3, \alpha)\text{Cl}^{34}$ ", M.S.

RADIOACTIVITY GROUP

I. Platinum Experiments

A number of experiments on the platinum isotopes have been completed, the principal aim having been the determination of the magnetic moments of the vibrational states in even-even platinum isotopes, $Pt^{192, 194, 196}$. The relevant lifetimes of the states are of the order of 50 psec, too short a time to measure Larmor frequencies reliably with standard external fields. Therefore, internal fields have been used, and these have been investigated using the Mössbauer effect in Pt^{195} . The summaries below describe this work.

A. Mössbauer Effect in Pt^{195}

Two levels in Pt^{195} , at 99 and 129 keV, are in principle observable by the Mössbauer effect; the decay scheme is shown in Fig. 9.1. A number of experiments have been carried out on the 99 keV, M1, transition using various source and absorber matrices. The most important are summarized in Table 9.1. Several conclusions can be drawn from this data:

1. The mean-life, deduced from the full-width-at-half-maximum is 190 psec in good agreement with Coulomb excitation measurements.
2. No evidence for either isomer shifts or quadrupole interactions is observed. None was expected.
3. The Debye temperature of Pt, Pt in Cu, and Pt in Ir, are 230 ± 15 , 230 ± 20 , and 320 ± 20 ° K respectively. The value for Pt was confirmed by a temperature dependence study from 10° to 80° of the resonance scattering and is in good agreement with specific heat data.

Figure 9.1

Table 9.I
Mössbauer Effect for 99 keV Pt¹⁹⁵

Source	Absorber	T(°K)	$\Delta(\%)$	$\Gamma(\text{cm/sec})$
Au ¹⁹⁵ in Cu	1.5 mil Pt	4.2	2.4	1.98
	3.0 mil Pt	4.2	4.6	2.00
	6.0 mil Pt	4.2	6.3	2.56
	6.0 mil Pt	77	0.9	---
	K ₂ Pt Cl ₆	4.2	2.1	2.00
	(1.8 mil Pt eqv.)			
	PtO ₂	4.2	4.5	2.00
Au ¹⁹⁵ in Ir	6.0 mil Pt	4.2	12.1	2.56
	11.2 mil Pt	4.2	11.9	2.85
Pt ^{195m} in Pt	4.5 mil Pt	4.2	~4.5	---
	6.0 mil Pt	4.2	~4.6	---

Many experiments on the 129 keV transition have been run, yet no unambiguous effect has been observed. Early indications of an effect have disappeared as the apparatus was improved. For example, using a Pt in Ir source and an 11.2 mil Pt absorber, two 50 point spectra were run each with more than 200,000 counts per point. In the first run the velocity range was ± 1.5 cm/sec, and in the second, ± 3.67 cm/sec. The expected full-width is 0.35 ± 0.05 from Coulomb excitation and lifetime measurements. No effect was observed and we can place a conservative estimate, taking into account background, that the effect is less than 0.6%. Assuming the Debye temperatures above we expect, again conservatively, 3%.

We have examined several possibilities which could explain this puzzle, although none looks promising:

1. The lifetime of the state is incorrect. This possibility seems to be ruled out both by the range of velocities over which we searched, i.e., from 1/2 to 10 times the expected width.
2. Broadening by internal fields. The lack of such perturbations for the 99 keV transition and the cubic nature of both source and absorber makes this possibility remote.
3. The Debye temperature is energy dependent. There is no doubt that to some degree this is the case but the data would require a reduction of from 320°K to ~200°K, a reduction difficult to understand for only a 30% change in transition energy (only a 70% change in recoil energy).
4. The decay scheme is incorrect and the 129 keV transition is either not to the ground state or there are two unresolved transitions, the ground state being less than 25% of the other. There is no evidence for either suggestion in the extensive

decay scheme work on these transitions. No 30 keV Mössbauer effect has been observed. No 00-129 keV coincidences have been observed. The 129 keV is observed in Coulomb excitation, from the decay of the 13/2 isomer in both singles and in coincidence with a 130 keV converted transition, and from Au^{195} both in singles and in coincidence with x-rays. Moreover, internal conversion studies show that the transition energies are correct to better than 1 keV and no other transitions than those in Fig. 9.1 are observed.

(A. Buyrn and L. Grodzins)

II. Internal Fields in Pt-Fe Alloys

The Mössbauer spectra of some Pt-Fe alloys taken at 4.2°K is shown in Fig. 9.2; the atomic percent of Pt is indicated. Over the range from 3 to 30% there is no observable change within about 15% in the splitting, hence in the value of H_{eff} . Neither a quadrupole interaction nor an isomer shift is observed for any composition except FePt. The ground state gyromagnetic moment is known, $g_0 = +1.02$. This large value dominates the 6-line hyperfine pattern hence the deduced value of H_{eff} is much more accurate than is g_{ex} . These values, obtained from a computer fit to the data, are as follows:

$$g_{\text{ex}} = -0.33 \pm 0.04$$

$$H_{\text{int}}(\text{Pt}_{03}\text{Fe}_{97}) = -1.380 \pm 0.085 \text{ moe.}$$

Figure 9.2

An intermetallic compound, FePt, displayed only a broadened line which could be fit by the same internal field. Line-broadening due to absorber thickness and quadrupole interaction smeared the pattern.

All samples were prepared from Pt foil and reagent-grade iron wire. These were melted in an argon arc furnace. A number of remelts -- up to 50 for Pt-Fe -- were carried out until no gross segregation was observable. Further analysis using a "microbeam probe" indicated homogeneous samples. The annealing and quenching assured that all samples were in the α phase at 4.2°K.

The value of H_{eff} in $\text{Pt}_{0.3}\text{Fe}_{0.7}$ is almost identical with that of dilute Au in Fe, -1.350 ± 50 MG, and within error the same as in Ir in Fe, $|1.300| \pm 0.300$ MG. Angular correlation measurements (see below), indicate that the sign of the internal field for Pt in Fe is negative, the same as for Au in Fe.

The internal field at the Pt nucleus is almost certainly mainly due to s-d exchange coupling which leads to a polarization of the conduction electrons. The s-electron density at the nucleus of Pt is high so that large internal fields result from little electron polarization and the value of 1.38 MG may be understood from current theories.¹ It is, however, surprising to find that the average internal field is essentially independent of the number of nearest neighbor Fe atoms since studies of H_{eff} on the iron site in various Fe alloys (not with Pt) show strong dependences. (A. Buyrn and L. Grodzins; N. Blum - National Magnet Lab; J. Wulff - Metallurgy)

III. Magnetic Moments of 2+ States in Pt^{192} , 194 , and 196

Using the internal fields in PtFe alloys discussed above, the g values of the first 2+ states in Pt^{192} , 194 , and 196 as well as the second 2+ state in Pt^{194} were measured. The pertinent levels used in the various angular correlations are shown in Fig. 9.3. For Pt^{192} and Pt^{194} , and $\text{Ir}_{0.3}\text{Pt}_{0.7}$ wire was activated in the reactor giving sources of Ir^{192} and 194 . Two sources of Au^{196} (which decays to Pt^{196}) in Fe were prepared. Source I was made by diffusing essentially carrier free Au^{196} into Fe. Source II was prepared by deuteron irradiation of $\text{Pt}_{30}\text{Fe}_{70}$; the source was annealed following bombardment.

In all cases, pulse height spectra were obtained both for the angular correlation and field flip measurements in order to subtract the backgrounds properly. Relative intensities of close lying levels were checked using a lithium drifted germanium detector. In the case of the Pt^{194} 622 keV level and the 356 keV level of Pt^{196} , this procedure allowed the separation of the near degenerate energies used for the correlation.

Figure 9.3

The results are given in Table 9.II. A_2 and A_4 are the angular correlation parameters, Δ is the field reversal effect, ωt is the precession angle and g is the deduced gyromagnetic ratio assuming an internal field of 1.38 moe for states in Pt^{192} and Pt^{194} and 650 koe for Pt^{196} . In this last case, the field was obtained directly from the Mössbauer effect in Pt^{195} measured at 4.2° K after the Au^{196} had diminished in intensity leaving Au^{195} dominant. There is, therefore, the further assumption that the internal field diminished little from 4.2 degrees to room temperature; this is a good assumption for source I and for $\text{Ir}_{03}\text{Fe}_{97}$ but a poor one for source II. The underlined value for ωt for the 328 keV state of Pt^{194} is felt to be more reliable since the 1151-328 γ - γ cascade is freer of unresolved and disturbing angular correlations.

Table 9.II
Data for Magnetic Moments of 2^+ States in Pt

Nucleus	Level (keV)	t_{mean} (psec)	Cascade	A_2	A_4	(%)	ωt	g
Pt^{192}	316	51	468-316 (4-2-0)	$.094 \pm .003$	$.008 \pm .004$	$2.38 \pm .16$	$.092 \pm .008$	$.27 \pm .03$
Pt^{194}	328	65	939-328 (0-2-0)	$.25 \pm .02$	$.78 \pm .03$	23.4 ± 1.0	$.097 \pm .04$	$.22 \pm .01$
			1151-328 (0-2-0)	$.12 \pm .02$	$.69 \pm .03$	26 ± 1.0	$.115 \pm .05$	$.26 \pm .02$
	622	89	645-622 (0-2-0)	$.26 \pm .04$	$.80 \pm .04$	26.7 ± 3.6	$.086 \pm .01$	$.14 \pm .02$
Pt^{196}	356	52	333-355 (2-2-0)	$.069 \pm .006$	$.325 \pm .008$	$4.28 \pm .58$	$.041 \pm .006$	$.23 \pm .06$

Several conclusions may be drawn from this work.

- 1) The magnetic moments are nearly constant and considerably lower than the value of $Z/A = 0.4$ which simple models might give.
- 2) The g values for the first and second 2^+ states of Pt^{194} are not equal as would be predicted by the harmonic approximation in which the magnetic moments of all members of the same band are proportional to the spin, I .
- 3) The magnetic moments of the $3/2^-$ state of Pt^{195} and the 2^+ states of Pt^{194} are not in agreement with a simple core-particle coupling picture for the $3/2^-$ state of Pt^{195} . A similar remark may be made about the $1/2^+$ first excited state of Au^{197} . A simple core-particle picture requires that the g value of the core, i.e., the first 2^+ state of Pt^{196} , be ~ 0.5 which is far outside the present errors.
- 4) The sign of the precession requires that $\mu_{2+} H_{\text{eff}}$ be negative. A recent external field measurement of $\mu(2+)\text{Pt}^{194}$ shows that its value is positive, hence H_{eff} is negative.

(A. Buyrn and L. Grodzins)

IV. Mössbauer Resonance Scattering Calculations

The Mössbauer resonance scattering yield, $N^M(\gamma)$ in the solid angle $d\Omega$ at Θ , at zero relative velocity, using a non-resonantly absorbing source, is given by

$$N^M(\gamma, \Theta) = \int_{-\infty}^{+\infty} dE \int_0^t dt' \int d\Omega \left\{ f_s N_s(E) e^{-\sigma_T(E) \eta t' / a} \frac{d\sigma^M}{d\Omega}(\Theta, E) \eta \left[(1 - f_{sc}) e^{-\sigma_e \eta t' / a} + f_{sc} e^{-\sigma_T \eta t' / b} \right] \right\} \quad (1)$$

where f_s and f_{sc} are the "recoilless" fractions for source and absorber respectively,

$$N_s(E) = \frac{\Gamma}{2\pi} \frac{N_s}{(E - E_0)^2 + (\frac{\Gamma}{2})^2} = \frac{2\Gamma N_s}{\pi [4(E - E_0)^2 + \Gamma^2]} \quad (2)$$

$$\sigma_T(E) = \sigma^M(E) + \sigma_{\text{Compton}}(E) + \sigma_{\text{Rayleigh}}(E)$$

$$\sigma_e = \sigma_T(E) - \sigma^M(E)$$

$$\frac{d\sigma^M}{d\Omega}(E, \Theta) = \frac{\sigma_0 f_{sc} \Gamma^2}{4(E - E_0)^2 + \Gamma^2} W(\Theta)$$

$$\sigma_0 = 2\pi \lambda^2 g$$

η is the density and a and b are the direction cosines for incident and exiting photons in the scatterer of thickness t .

Equation 9.1 can be integrated immediately over t . The result may be integrated over E in special cases, the most important being the case where

$$\sigma_T(E) \eta \frac{t}{a} \ll 1 \quad (3)$$

In that case we can integrate the leading term exactly obtaining

$$N^M(\gamma, \Theta) = \frac{N_s f_s f_{sc}}{2(1+\alpha)^2} t \frac{d\sigma_0^M}{d\Omega}(\Theta) \quad (4)$$

To estimate the yield for Mössbauer scattering, calculations have been carried out for

$$\frac{d\sigma^M}{d\Omega}(\text{effective}, \gamma) = \frac{f_s f_{sc}}{2(1+\alpha)^2} \frac{d\sigma_0^M(\Theta)}{d\Omega} \quad (5)$$

as a function of Z and E_0 . These are given on the left graphs of Figures 9.4 through 9.10 (Differential cross sections in $\text{cm}^2/\text{steradian}$ for equations 9.5 and 9.6 and for Rayleigh, Compton, and photo-effect processes at a scattering angle of $\Theta=150^\circ$.) The Debye temperature of source and scatterer are assumed to be the same and equal to 270°K . The scattering distribution has been assumed isotropic. The solid line is for $\alpha=2$. The dashed curves are for M1 and E2 transitions respectively; values of α were taken from Rose's tables. The Rayleigh and Compton cross sections are also given for a scattering angle of $\Theta=150^\circ$.

Figure 9.4

Figure 9.5

Figure 9.6

Figure 9.7

Figure 9.8

Figure 9.9

Figure 9.10

In principle the Mössbauer effect for a thin scatterer is given by $\frac{d\sigma^M}{d\Omega} / \frac{d\sigma^R}{d\Omega}$ since the Compton gamma rays may be discriminated against. This theoretical ratio is given in Fig. 9.11 for $Z = 30$ and 90 .

Figure 9.11

Theoretically observable Mössbauer effect at $\Theta=150^\circ$,
for $Z=30$ and 90 .

If the de-excitation internal conversion electrons are detected the above equations are nearly unchanged, one only replaces Γ_γ^2 in Equation 9.2 by $\Gamma_\gamma \Gamma_e$. For K shell electrons Equation 9.5 becomes

$$\frac{d\sigma^M}{d\Omega}(\text{effective, } e_K) = \frac{f_s f_{sc} \alpha_K}{2(1+\alpha)^2} \frac{d\sigma_0^M}{d\Omega}(\Theta) \quad (6)$$

This effective cross section as a function of Z and E_0 is given on the right graphs of Figures 9.4 through 9.10. The value of $\alpha_K = 1$ for the solid curve.

The competitive process is the photo-effect which is strongly peaked forward. However, over most of the range of Z and E considered, a scatterer which may be thin both as regards the Mössbauer effect and the electron energy lost, may be thick insofar as multiple scattering is concerned. We have, therefore, included both the thin ($\Theta=150^\circ$) and the thick (isotropic) scattering cases. Fig. 9.12 is the K-electron analogue of Fig. 9.11.

Figure 9.12
Theoretically observable Mössbauer effect at $\Theta=150^\circ$,
for $Z=30$ and 90 .

Calculations of intensity and line widths for scatterers of finite thickness will be presented in a subsequent report. (L. Grodzins and M. Slade)

V. Liquid Helium Cryostat for Mössbauer Scattering Experiments

A simply constructed cryostat designed for Mössbauer gamma-ray scattering has been constructed and used for more than a year; see Fig. 9.13 and 9.14. The sample volume is part of the transfer line from a liquid helium reservoir to a mechanical pump. The helium vapor exiting from the sample volume cools the transfer line and radiation shield. The temperature of the sample is controlled by the helium flow rate determined by a needle valve at the pump. The operating range is from 9° to $\sim 300^\circ\text{K}$. The liquid helium consumption in liters/hour is 0.3 at 10°K , 0.2 at 16°K , and 0.1 at 34°K . Source and scatterer temperatures differ by less than $1/2^\circ$ at 10°K and less than 7° at 80°K . (L. Grodzins and P. H. Barrett)

Figure 9.13

Schematic drawing of the helium flow cryostat and source-scatterer-detector geometry. Each portion of apparatus is drawn approximately to scale. The inner diameter of the cryostat is $3/4$ ", the source to scatterer distance is 3" and the overall length, exclusive of the detachable tip, is about 40".

Figure 9.14

Schematic drawing of the Mössbauer scattering insert. The lower portion is to scale; the outer diameter of C is $23/32$ ".

V. Magnetic Moments of 1st 2+ States by the Precession of the Mössbauer Scattering Distribution

This work, preliminary accounts of which were reported in the May, 1964, Progress Report, is essentially complete. The results are summarized below, values in parentheses are still not firm.

The electric quadrupole interaction of the 2+ state in Os^{186} in osmium metal is

$$\omega_E \tau = 0.043 \pm 0.004$$

where $\omega_E = eqQ/4I(2I-1)\hbar$.

Gyromagnetic Ratios of 2+ States

Isotope	Energy (keV)	$t_{1/2}(\text{n sec})$	g
W^{182}	100	1.40	$(.27 \pm 0.27)$
W^{186}	122	0.98	$.34 \pm 0.03$
Os^{186}	137	0.84	$.32 \pm 0.02$
Os^{188}	155	0.73	$(.30 \pm 0.03)$

(Y.-W. Chow and L. Grodzins)

VI. Internal Magnetic Field in W-Fe Alloys

The internal magnetic field at the tungsten nucleus in alloys containing 5, 10, and 15% by weight of tungsten have been determined from the Mössbauer absorption spectra of the 100 keV state of W^{182} . Fig. 9.15 shows one of the spectra. The computer fit splitting yield values of 668 ± 100 , 678 ± 80 , and 684 ± 75 koe for H_{eff} for three alloys, assuming that $g(2+) = 0.27$. The experiments do not yield a value for the sign of the field.

Fig. 9.15

(Y.-W. Chow and L. Grodzins - LNS; J. Wulff - Metallurgy; N. Blum - National Magnet Lab.)

VII. Magnetic Moments of W^{182} and W^{184} , II

The determination of the magnetic moment of the 100 keV state of W^{182} by the precession of the Mossbauer scattering distribution -- see above -- has been supplemented by a direct measurement of the hyperfine splitting using an external magnetic field. A Ta^{182} source, outside the field, was run against a 1 mil natural tungsten absorber placed in a longitudinal field of 105 koe. The radiation was detected by a 4 mm thick lithium-drifted germanium detector. The Mossbauer spectrum is shown in Fig. 9.16; the width at zero field is indicated. The result is $g(2+) = 0.34 \pm 0.10$ assuming that $H_{eff} = 105$ koe. The error includes the uncertainty in the width of the hyperfine lines.

Ta^{182} SOURCE - 1 MIL W ABSORBER UNDER
EXTERNAL MAGNETIC FIELD 105 KOE
USING SOLID STATE DETECTOR AT LIQ.
HE. TEMPERATURE.

Figure 9.16

The ratio of the magnetic moments of the first $2+$ states of W^{182} and W^{184} was measured by comparing the hyperfine interactions in a W-Fe alloy. Only the source differed in the two measurements, being in one case, Ta^{182} from neutron capture in Ta^{182} and in the other Re^{184} from the α, n reaction on Ta^{181} . The result is $g(184)/g(182) = 1.06 \pm 0.15$. (Y. -W. Chow and L. Grodzins - LNS; N. Blum - NML)

VIII. Lead-Loaded Plastic Scintillators

Plastic scintillator detectors containing up to 10% by weight of lead (Pb plastic) are now being manufactured commercially.² We have been using such detectors for more than a year in studies of magnetic moments of excited states in heavy nuclei by the spin precession of α - γ

correlations in an external magnetic field. Some of the pragmatic test results we have obtained using 1"-diameter x 1"-high cylindrical samples of this material are presented here in order to indicate the usefulness for various problems.

The loss of light, over that of unloaded plastic, is approximately 30 %, 50% and 65% for 2%, 5%, and 10% Pb plastic. The photopeak efficiency, relative to that of NaI, was measured for the 73 keV Pb x-ray from a source of Bi²⁰⁷, giving values of 15%, 35%, and 52% for 2%, 5%, and 10% Pb loaded, respectively. For the 122 keV gamma ray of Co⁵⁷ the values were 10%, 23%, and 35%. The pulse height spectrum from Co⁵⁷ in a 5% Pb plastic is shown in Fig. 9.17. The energy resolution is roughly 54%.

PULSE-HEIGHT SPECTRUM OF Co⁵⁷
5% PB PLASTIC

Figure 9.17

Time resolution results obtained using 56 AVP photomultipliers and Chronetics circuitry are as follows: the slopes (drop-off by 1/2) are for the Pb plastic side:

	<u>F W H M</u> <u>(psec)</u>	<u>Slope</u> <u>(psec)</u>
Bi ²⁰⁷ source: 73 keV x-ray versus 976 keV electrons detected in a thin Naton crystal		
2 and 5% Pb plastic	800	150
10% Pb plastic	1000	200
Co ⁶⁰ , γ - γ , 10% Pb plastic versus thin Naton plastic	500	75

The detection efficiency, rapid response, photo-peak yield, relatively large size, and low cost make Pb plastic advantageous for a number of applications, among them being:

1. Fast coincidence experiments involving gamma rays, especially of low energy. The Compton peak, which is usually used for gamma detection in plastic, has an energy

$$E_c = \frac{E}{1 + \frac{mc^2}{2E}}$$

so that E_c decreases, and hence time resolutions worsen, rapidly as the photon energy decreases below about 200 keV. For example, the energy lost in the plastic detecting the Compton edge electrons is 90 and 8 keV for photon energies of 200 and 50 keV respectively.

2. Experiments in which the high detection efficiency is important, such as the measurement of positron decay lifetimes and angular distributions or as an anti-Compton shield surrounding NaI(Tl) or solid state detectors.
3. The detection of a high flux of gamma rays where pile-up can be minimized by the use of short pulses.

(W. Schick, D. Murnick, L. Grodzins)

IX. Nuclear Excited State Lifetimes

The lifetimes of the first and second excited rotational states of Ra^{224} , Th^{228} and Th^{230} together with the first vibrational state of Rn^{220} were measured and will appear presently in the Physical Review. The final results are tabulated in Table 9.III along with some of Bell's ³ values.

Table 9.III

Nucl	$I\pi$	E_γ (keV)	$T_{1/2}$ (psec)	$T_{1/2}$ (Bell) (psec)	Q_0 (10^{-24} e-cm ²)	β	$\frac{B(E2; 4+ \rightarrow 2+)}{B(E2; 2+ \rightarrow 0+)}$
Ra^{224}	2+	84.5	744 ± 19	760 ± 30	6.27 ± 0.09	0.166	1.37 ± 0.07
	4+	168.5	181 ± 9				
Th^{228}	2+	57.5	402 ± 10	400 ± 30	8.35 ± 0.12	0.210	1.46 ± 0.05
	4+	128.6	161 ± 5				
Th^{230}	2+	52.4	354 ± 9	370 ± 20	8.95 ± 0.13	0.222	1.40 ± 0.05
	4+	117.6	166 ± 5				
Ra^{220}	2+	241.	145 ± 5	150 ± 10			

(W. R. Neal and H. W. Kraner)

X. Lithium-drifted Germanium Detectors

Development of lithium-drifted germanium detectors for gamma ray spectroscopy has continued with several small ($\sim 0.5 \text{ cm}^3$) successful counters having been produced. These detectors have been used for general spectroscopy as well as the unique application to a Mossbauer experiment with relatively high energy transitions in a strong magnetic field. Further work on the method of fabrication will, we hope, yield useful detectors having up to $\sim 5 \text{ cm}^3$ active volumes. A very useful and simple dewar has been designed and built by Arthur D. Little Inc. which allows convenient access to the detector and a long LN_2 hold time of up to three days for a 1.5 liter charge. (W. R. Neal and H. W. Kraner)

REFERENCES

1. E. Daneil and J. Friedel, J. Phys. Chem. Solids 24, 1601 (1963).
2. Pilot Chemicals, Inc., Watertown, Massachusetts .
3. R. E. Bell, S. Bjørnholm and J. C. Severiens, Kgl. Danske Videnskab. Selskab, Mat.-Fys. Medd. 32, No. 12 (1960).

PUBLICATIONS

- N. Blum and L. Grodzins, "Sign of the Magnetic Hyperfine Field in Dilute Iron Alloys using Mössbauer Effect", Phys. Rev. 136, A133 (Oct. 5, 1964).

THESIS

- D. Yeboah-Amankwah, M.S. in the Dept. of Physics, June 1964.

CYCLOTRON GROUP

I. Cyclotron Operation and Equipment

During the past summer we installed a new bending magnet and quadrupole pair right outside the cyclotron. The present setup is shown in Fig. 10.1 (the new magnets are A, Q_1 and Q_2). The purpose of this improvement was to increase beam intensity without sacrificing resolution. A study of the beam characteristics was made during the fall and as an example of the improvements made we shall compare two (α , α') experiments. During the spring we did Ca^{40} and obtained a beam $\approx 0.030 \mu\text{a}$ with a resolution of ≈ 130 keV. This fall we did $\text{Ca}^{44}(\alpha, \alpha')$ with a resolution of ≈ 110 keV and a beam intensity of $\approx 0.3 \mu\text{a}$. This beam is approximately $1/8''$ wide by $1/2''$ high. To facilitate work on the beam optics two viewing ports were added before and after the 30° bending magnet (B). At the present time we are still making some beam measurements, and we are calculating the beam optics in order to further understand and improve the system.

Figure 10.1
Cyclotron and External Beam
Layout

This fall we have also added a cold trap and baffles to the scattering chamber to cut down the oil vapor in the system. This should help to eliminate carbon deposits on our targets. This may also increase the longevity of solid state counters as well as enabling us to cool them. A system to cool detectors was also installed and it has improved our resolution and also apparently improved the rise time of thicker detectors. (N. S. Wall, G. Heymann, F. Fay, E. White, A. Bernstein)

II. $\text{Ca}^{40}(\alpha, 2\alpha)\text{Ar}^{36}$ Reaction

The correlations in energy and angle of the alpha particles produced by the reaction $\text{Ca}^{40}(\alpha, 2\alpha)\text{Ar}^{36}$ have been measured for incident alpha energies of 28.7 and 30.5 MeV. Data were taken with a two-dimensional pulse-height analyzer for an angular range from 20° to 135° with solid-state detectors (angular aperture $\pm 2^\circ$) confined to the reaction plane. It was found that the first excited state (1.97 MeV) in Ar^{36} is much more strongly populated than the ground state. Pronounced structure was observed in practically all correlation spectra taken. Fig. 10.2 shows the data for one pair of angles, the area of a circle in proportion to the number of counts in the corresponding channel. The triangular portion near the origin has been arbitrarily left blank. The number of counts per channel increases as one approaches the origin. At the higher incident energy and at detector angles of 35° - 35° and 55° - 55° with respect to the beam, the alpha particles were found to be emitted preferentially with equal energies. When the incident energy was lowered one alpha particle energy remained constant. This implies a two-step process via an excited state in Ca^{40} near 18.8 MeV. Measurements with unequal detector angles were carried out to determine alpha correlations with respect to the recoil axis. This experiment is being continued.

Figure 10.2

A two-dimensional spectrum from the $\text{Ca}^{40}(\alpha, 2\alpha)$ reaction. The areas of the circles are proportional to the number of counts. The axes represent the energies of coincident particles in the counters. The line labeled "1st" represents the kinematic line for leaving the residual nucleus (Ar^{36}) in its first excited state. The area near the origin has been left blank but there were many counts in that region.

(R. W. Bauer, C. R. Gruhn, G. Heymann, W. Kossler, N.S. Wall)

III. $\text{Ti}^{50}(\alpha, \alpha')$

The elastic and inelastic cross sections for scattering of 30.5 MeV alpha particles on Ti^{50} were measured with an overall resolution of 130 keV. The target was a thin self-supporting foil of separated isotope Ti^{50} but unfortunately was only 70 percent pure. For this reason only angular distributions for five excited states could be observed. These include known levels at 1.55 (2^+) and 2.67 (4^+) MeV and states at 4.42 and 7.1 MeV which are assigned as 3^- states, and a state at 4.32 which is assigned as 2^+ . The 4.42 MeV assignment agrees with previous (p,p') assignments^{1,2}. The 7.1 MeV state is approximately 40 percent as strong as the 4.42 MeV state. These data are presented in Figs. 10.3 through 10.6. An optical model fit was made to the elastic scattering data (Fig. 10.3) using a Woods Saxon potential with parameters: $V=47.6$ MeV, $W=14.3$ MeV, $R=5.88$ f, and $a=0.55$ f. In Fig. 10.4 the results for the 2^+ states at 1.55 and 4.32 MeV are presented. The solid lines represent DWBA fits, as calculated by JULIE, with $\beta_2=0.13$ and 0.08 respectively. Coulomb excitation has been included. Fig. 10.5 shows the results for the 2.67 MeV 4^+ state, the solid line represents the DWBA fit for $\beta_4=0.08$. Coulomb excitation has not been included here. Fig. 10.6 shows the results for the 3^- states. For the lowest one at 4.42 MeV we have $\beta_3=0.12$, while for the state at 7.1 MeV we have $\beta_3=0.09$. The fits to all of this data are quite good. These fits have been made with collective form factors. We hope to calculate the results using shell model wave functions.

Figure 10.3

Elastic Scattering of 30.5 MeV alpha particles from Ti^{50} . The solid line is an optical model fit.

Figure 10.4

Differential cross section for inelastic scattering of 30.5 MeV alpha particles to the 1.55 MeV and 4.32 MeV states in Ti^{50} . The solid curves are DWBA fits.

Figure 10.5

Differential cross section for inelastic scattering of 30.5 MeV alpha particles to the 2.67 MeV state of Ti^{50} . The solid curves are DWBA fits.

Figure 10.6

Differential cross section for inelastic scattering of 30.5 MeV alpha particles to the 4.42 and 7.1 MeV states of Ti^{50} . The solid lines are DWBA fits.

(E. P. Lippincott and A. Bernstein)

IV. $\text{Zr}^{90}(\alpha, \alpha')$

We have obtained preliminary data on the differential cross section for the scattering of 30.5 MeV alpha particles from Zr^{90} . The elastic scattering data is shown in Fig. 10.7 as the ratio to Rutherford scattering. We have obtained an optical model fit using a Wood Saxon potential with $V=45.5$ MeV, $W=12.6$ MeV, $R=6.97$, and $a=0.60$ f. For inelastic scattering we have data for states at 2.18 (2^+), 2.33 (5^-), 2.745 (3^-), 3.07 (4^+), 3.31, 3.45 (6^+), 3.89, 4.37, and 5.71 MeV. All of our values have an uncertainty of ± 40 keV except the 2.745 state upon which the energy calibration was based. The 0^+ state at 1.75 MeV was not excited. The first four observed states exhibit strong oscillatory behavior which is somewhat different from the usual phase rules observed in inelastic alpha particle scattering. This had been previously observed for the 3^- state³ and is apparently a Coulomb effect as can be demonstrated from the fact that there are normal phase relations for the 3^- and 2^+ states at 42 MeV.⁴ Fig. 10.8 presents the scattering from the 3^- and 5^- states with DWBA fits as computed by JULIE. Coulomb excitation cannot be included because our code does not have enough partial waves for an accurate calculation of this value of Z . The calculation assumes a collective form factor. The values of β obtained are

$\beta_3=0.12$ and $\beta_5=0.054$. Fig. 10.9 shows the data and DWBA fit for the 2^+ state. Here $\beta_2=0.077$. Fig. 10.10 shows the data and DWBA for the 4^+ state, for which we obtain $\beta_4=0.031$.

Figure 10.7

Elastic scattering cross section for 30.5 MeV alpha particles on Zr^{90} plotted as the ratio to Rutherford scattering. The solid line is an optical model fit to the data.

Figure 10.8

Differential cross sections for the inelastic scattering of 30.5 MeV alpha particle to the 2.75 and 2.32 MeV states of Zr^{90} . The solid lines are DWBA fits.

Figure 10.9

Differential cross section for the inelastic scattering of 30.5 MeV alpha particle from the 2.18 MeV state of Zr^{90} . The solid line is a DWBA fit.

Figure 10.10

Differential cross section for the inelastic scattering of 30.5 MeV alpha particles from the 3.07 MeV state of Zr^{90} . The solid line is a DWBA fit.

We are presently attempting to fit the 40 MeV data and to extract parameters for the 3^- and 2^+ states. We are also working on the question of the Coulomb effects on the phase relationships. We shall attempt to use the shell model to calculate differential cross sections. We are also attempting to assign spins and parities to the other states that we have data for, but for which no assignments have been made. (E. Martens and A. Bernstein)

V. $\text{Ca}^{40}(\alpha, p)\text{Sc}^{43}$

This experiment represents the first in a series of multi-nucleon transfer reaction experiments in medium weight nuclei. A natural Calcium foil (approximately 1 mg/cm^2 thick) was bombarded with 30.5 MeV α particles. The resulting proton spectra were recorded at 5 degree intervals from 15 to 90 degrees. The detector consisted of a 500μ silicon surface barrier transmission counter (thick enough to stop the scattered α particles in front of a 4.6 mm silicon detector consisting of two 2.3 mm surface barrier detectors connected together. The two signals were added together and gated by a discriminator on the signal from the thick counter. Since the (α, p) Q value (-3.5 MeV) is at least 8 MeV more positive than the Q value for (α, d) , (α, t) or (α, He^3) reactions only protons were recorded at the high energy end of the spectra. A typical spectrum is shown in Fig. 10.11. The angular distributions (Fig. 10.12, 10.13, and 10.14) were extracted by visually fitting the experimental spectra with a background and gaussian peaks generated utilizing the LNS PDP -1 computer. Cross sections for states up to 3.10 MeV excitation have been extracted. The Q values obtained agree quite well with those previously found in this reaction using 20 MeV particles.⁵

Figure 10.11

Spectrum of protons from the $\text{Ca}^{40}(\alpha, p)\text{Sc}^{43}$ reaction at 45 degrees (lab.).

Figure 10.12

Differential cross sections for the $\text{Ca}^{40}(\alpha, p)\text{Sc}^{43}$ reaction leading to the ground state and 1.17 and the 1.38 MeV states of Sc^{43} . The lines have been drawn through the data for visual purposes only.

Figure 10.13

Differential cross sections for the $\text{Ca}^{40}(\alpha, p)\text{Sc}^{43}$ reaction leading to states at 0.46, 1.82, and 2.61 MeV in Sc^{43} . The lines have been drawn through the data for visual purposes only.

Figure 10.14

Differential cross sections for the $\text{Ca}^{40}(\alpha, p)\text{Sc}^{43}$ reaction leading to states 2.27, 2.96 and 3.10 MeV in Sc^{43} . The lines have been drawn through the data for visual purposes only.

Figure 10.15

Laboratory cross sections for the elastic scattering of 160 MeV protons from Be^9 , Ca^{40} , Sn^{120} , and Pb^{208} . The data points are not shown -- just lines drawn through the data.

Theoretical fits to the angular distributions are being developed using a DWBA calculation carried out by the JULIE program on the 7094 computer. The form factors for the reaction have been worked out by Dr. B. Bayman of Princeton and will be used in the calculation.

Of special note is the fact that the $d_{3/2}$ hole state is Sc^{43} at 150 keV⁶ and has been excited by less than approximately 10 percent of the ground state transition. One expects the ground state transition to go by a stripping of three particles from He^4 and putting them in the $f_{7/2}$ shell. The $d_{3/2}$ hole state, on the other hand, would be populated by knocking out a proton from the closed $d_{3/2}$ shell and then putting the four nucleons in the $f_{7/2}$ shell. This latter probability is at least a factor of 10 smaller than the stripping one. (R. Ginaven and A. Bernstein)

VI. Data Analysis on the PDP -1 Computer

A data analysis program has been written which has as its philosophy leaving the maximum amount of decision making to the experimenter and most of the repetitive operations to the computer.

Data from a nuclear reaction experiment consisting of counts versus channel number are put on punched paper tape by a multi-channel analyzer. This tape is read conveniently by the PDP -1 computer and is then displayed on a scope. Any 64 channel subset can be expanded for convenience. The areas under isolated peaks are extracted by two methods. First by adding up the counts between two channel numbers, designated by the experimenter, with a linear background subtracted. In the second method the parameters of a gaussian curve (mean position, height, and width) are adjusted by switches on the console until a good fit to the data is obtained. The quality of fit is judged by the experimenter comparing the data and the fit on a scope.

When the fit is considered satisfactory, the experimenter types 0 on the typewriter and the computer types out the parameters of the fit. Multiple peaks are fit as a superposition of gaussian peaks plus a linear background.

This method has already proved to be reliable and rapid, and has been used in the analysis of the $Ca^{40}(\alpha, p)Sc^{43}$ reaction. (T. Provost and A. Bernstein)

VII. Elastic and Inelastic Scattering of High Energy Protons from Nuclei

This experiment was performed in order to study the scattering of 160 MeV protons using the Harvard Cyclotron. The energy of the scattered protons was measured using a 3"x3" Na I crystal. The experiment can be divided into three parts.

A. Elastic Scattering

Angular distributions were obtained (approx. 5 degrees to approx. 50 degrees) for elastic scattering from Be^9 , Ca^{40} , Ni^{58} , Sn^{120} , and Pb^{208} and are presented in Fig. 10.15. The angular

distributions were then fit with optical model calculations using a Saxon-Woods shape for the potential well. In order to fit the experimental data, it was found necessary to make the imaginary well radius larger than the real well radius.⁷ The results for Ca^{40} are shown in Fig. 10.16. The results for the other nuclei are equally good.

Figure 10.16

Elastic scattering of 160 MeV protons from Ca^{40} in the C.M. System. The solid line represents an optical model fit to the data.

B. Inelastic Scattering

Angular distributions were obtained for the 2.43 MeV level in Be^9 (10° - 50°), the 3.73 MeV level (10° - 40°) and 4.5 MeV, level (20° - 40°) in Ca^{40} , and the 1.45 MeV level (14° - 24°) and 4.5 MeV, level (14° - 24°) in Ni^{58} . Calculations of these inelastic cross sections were made using the Distorted Wave Born Approximation (DWBA). The interaction used was that appropriate to a non-spherical well, and therefore describing collective states of the nucleus. In addition one calculation for Be^9 was made using the impulse approximation, and the nuclear matrix element derived from electron scattering and the results are shown in Fig. 10.17. The impulse approximation calculation fits the data quite well. The DWBA calculations, however, do not give good results.

Figure 10.17

Inelastic scattering of 160 MeV protons to the 2.43 MeV state in Be^9 . The solid lines are theoretical curves as described in text.

C. "Quasi-Elastic" Scattering

The proton continuum of the scattered protons has been measured for approximately 35 MeV to 160 MeV. This measurement has been made for Be^9 , C^{12} , Ca^{40} , Ni^{58} , Sn^{120} and Bi^{209} in 10° steps from 20° to 80° . The continua show broad peaks which at any particular angle are very similar for all nuclei. An example of this data is shown in Fig. 10.18. The peak energy position shifts to lower energy with an increase in angle, but not exactly according to free nucleon-nucleon kinematics. One would expect the proton continuum to reflect the momentum distribution of the nucleons in the nucleus if the process was "quasi-elastic" scattering; that is, if the scattering was the result of a single proton-target nucleon interaction. The results are being considered on the basis of "quasi-elastic" scattering. Copies of this thesis are available from Dr. Roos at the University of Maryland.

"QUASI-ELASTIC" SPECTRA AT 50°

Figure 10.18

"Quasi-elastic" spectra at 50°.

(P. G. Roos and N. S. Wall)

VIII. A Search for Two-Photon Decays in the Spectrum of In^{114}

Data on this process has been taken using the two-dimensional mode of the 4096 channel analyzer. The analysis of this data is in progress. (E. Romberg and N.S. Wall)

REFERENCES

1. H. O. Funsten, N. R. Roberson, and E. Rost, Phys. Rev. 134, B117 (1964).
2. W. S. Gray, R. A. Kenefick, and J. J. Kraushaar, to be published.
3. M. Crut, D. R. Sweetman and N. S. Wall, Nucl. Phys. 17, 655 (1960).
4. D. L. Hendrie, and G. W. Farwell, Phys. Letters 9, 321 (1964).
5. N. O. Lassen and O. Nathan, Nucl. Phys. 42, 183 (1963).
6. R. E. Holland, F.J. Lynch, and K.E. Nysten, Phys. Rev. Letters 13, 241 (1964). R.D. Lawson and M. H. McFarlane, to be published.
7. The same conclusion was reached by G. R. Satchler and R. M. Haybron, Phys. Letters 11, 313 (1964).

PUBLICATIONS

- A. M. Bernstein, R. A. Haberstroh, D. R. Hamilton, M. Posner, and J. L. Snider, "Spin and Magnetic Moment of N^{13} ", Phys. Rev. 136, B27 (1964).
- N. S. Wall, G. Heymann, R. W. Bauer, and A. M. Bernstein, "The Relative Size of Ca^{44} and Ca^{40} ", Comptes Rendus du Congres International de Physique Nucleaire, Vol. II, p. 866 (1964).
- R. A. Haberstroh, W. J. Kossler, and D. R. Hamilton, "Hyperfine Structure and Nuclear Moments of $20.4 \text{ min } C^{11}$ ", Phys. Rev. 136, B932 (1964).

ADDRESSES

- N. S. Wall, "Elastic and Inelastic Scattering of He^3 and He^4 ", Gordon Conference, August 1964.
- N. S. Wall, "Elastic Scattering of Alpha Particles", Paris Conference, July 1964.
- A. Bernstein, "A Search for Particle Stable State in H^4 and H^5 ", Paris Conference, July 1964.
Yugoslave Summer School, July 1964.
Northeastern University, November 1964.
- W. J. Kossler, "Study of the $Ca^{40}(\alpha, 2\alpha)$ Reaction", Gatlinburg Conference, October 1964.

THESES

- R. Burn, "The Decay of Mn^{57} ", B. S., May 1964.
- P. Roos, "Elastic and Inelastic Scattering of High Energy Protons from Nuclei", Ph.D., September 1964.

THEORETICAL GROUP

As can be seen from the enclosed abstracts, the theoretical group has been interested in the major problems of nuclear physics and elementary particle physics. Field theory, unitary symmetry group description of elementary particles and phenomenological analysis of data are being actively studied in elementary particle physics. In nuclear physics reaction theory, at both high and low energies, gamma ray and inelastic scattering phenomena, nuclear matter and the bound states of nuclei are under consideration.

In addition to the visiting research associates, we also have a number of guests who are supported either by fellowships or by their governments. We have two such people from France, two from Germany and one from Australia. We also have a number who are here on sabbatical from institutions whose support we only contribute in part.

I. Field Theory

A. Boot-strap solutions

Work on boot-strap solutions of static models has been completed and submitted for publication. Final results, for the models examined, are: no boot-strap solutions exist for un-subtracted dispersion relations; for subtracted dispersion relations, boot-strap solutions exist for all the models examined, but do not uniquely determine the expected number of parameters. (F. Low and K. Huang)

B. Topology and Feynman Diagrams

The geometric method of studying Feynman diagrams initiated by Fotiadi, Froissart, Lascoux, and Pham has been applied to a Feynman diagram containing two internal loops. A bound on the number of invariant functions associated with the diagram was obtained and some non-Landau singularities studied. The increase in difficulty over the single loop cases is great. (P. Federbush)

C. A Note on Non-Landau Singularities

The calculation of some homology groups relevant to sixth order Feynman diagrams, has been submitted for publication to the Journal of Mathematical Physics. (P. Federbush)

D. Analysis of Q.E.D. Energy-Momentum Tensor

Work has been in progress to find a finite expression for the energy-momentum tensor in quantum electrodynamics. Invariance under coordinate transformations is used to construct a non-local expression which in a limit is the desired tensor. An analogous approach was used by K. A. Johnson (Brandeis Summer Lecture 1964) to construct a non-local current vector from a gauge invariance principle. (E. J. Irwin, Jr.)

E. Three-Body Problems in Fundamental Particle Physics

Relativistic (Bethe-Salpeter) equations have been found which are direct analogues to the Fadeev equations for the scattering of three non-relativistic particles with two-body interaction only. Some approximation schemes for their solution have been considered. (R. F. Sawyer)

F. Solution of the Bethe-Salpeter Equation

The solution of the Bethe-Salpeter equation for non-zero energies is being studied. The ladder approximation is made and finite mass particles are considered. (R. P. Zia, A. P. Stamp, R. Saenger, H. Feshbach)

G. Three-Body Forces in Nuclear Physics

We are attempting to ascertain whether or not there exists a unique long-range tail to the nuclear three-body force analogous to the one-pion exchange potential for the nuclear two-body force. On the assumption that only the long-range part of this force will be important, we are preparing to calculate the configuration splitting in nuclei with three particles outside of closed shells. (H. K. Quang and A. K. Kerman)

II. Unitary Symmetry

A. Mass Formula and Symmetry Breaking in SU(6)

A study of the mass formula and symmetry breaking in the SU(6) symmetry scheme is continuing as well as an attempt to understand the electromagnetic mass splitting.¹ (T. Yao)

B. The Electromagnetic Structure of Vector Mesons and SU(3) Symmetry

In an investigation of the electromagnetic structure of vector mesons in SU(3) symmetry a table of the moments of vector mesons has been given. It was found that in the model of pole dominance for the electromagnetic form factors the anomalous magnetic moment of all members of the vector-meson octet is zero.

An investigation of the force in the baryon-vector-meson system is in progress.
(D. Flamm)

III. Phenomenological Analyses

A. ϕ -Meson Decay and Amplitude Parity

The amplitude parity quantum number² predicts that the ϕ -meson does not decay into ρ and π mesons. However, it seems likely that the quantum number is violated by the presence of baryon-antibaryon pairs. Accordingly, if amplitude parity alone is what suppresses the $\rho\pi$ decay of the ϕ , there must be a small decay amplitude due to the process $\phi \rightarrow B + \bar{B} \rightarrow \rho + \pi$. It is conjectured that the high mass of the $B + \bar{B}$ system relative to the ϕ mass, and the limitations imposed by unitarity on the weight function of the decay amplitude dispersion relation will cause the decay rate to be small in this mode. Alternatively, the decay rate could be small if the ϕ couples very strongly to the $B + \bar{B}$ system so that pair suppression is operative.

In our work we are studying the decay by dispersion methods. It is our aim to determine quantitatively how much suppression of $\phi \rightarrow \rho + \pi$ can be provided by the mechanisms cited above. Since we are primarily concerned with an order of magnitude estimate, only the NN intermediate baryon state is included. NN scattering is parametrized by a scattering length approximation, and $NN \rightarrow \rho\pi$ is assumed to proceed via an intermediate ω -meson. (J. B. Bronzan)

B. The Pion-Nucleon "Second Resonance"

Coupling to the π -N* channel alone was shown not to lead to the $D_{1/2} \ 3/2$ "second resonance". This confirmed the theoretical implication that inter-channel coupling would cause resonances only below the threshold of the coupled channel. Previously coupling to the π -N channel had been shown to describe adequately the complex $D_{1/2} \ 3/2$ amplitude up through the resonance.³ We have now shown that a moderate additional coupling to the π -N* channel will cause considerable coupling. We will compare the three-body final states predicted by this three-channel model with experiment.

The details of the appearance of the resonance and its movement down from threshold with increased inter-channel coupling have been studied. (E. Lomon and A. Miller)

C. The Nucleon-Nucleon Boundary Condition Model

In order to improve the charge-independent fit to the neutron-proton data, two alterations of the previous form of the model are being tested. One is the use of separate boundary-condition radii for $T=0$ and $T=1$. The other is the addition of the meson potential (with no force parameters) to the two pion exchange potential. In the first case, our present proton-proton fit is automatically maintained. In the second we have successfully refitted to the proton-proton data. This new fit allows the equality of the coupling constant in the one- and two-pion exchange contributions, while still maintaining physically meaningful values of the potential parameters and the boundary radius. (H. Feshbach and E. Lomon)

D. Mandelstam Relation for the Boundary Condition Model

A proof of sufficient conditions for the existence of Mandelstam-type relations of the kind that has been demonstrated to exist for the hard core potential by Nussenzweig is being developed for the boundary condition model. (F. Newman)

E. Two-Body Forces with Finite Repulsive Cores

Some work has been done on the problem of finite repulsive potentials to replace the hard cores in nucleon-nucleon scattering. We have found that an equally good fit to the data can be obtained with finite cores whose radius is considerably larger than the infinite core conventionally used. Work is continuing on this subject. (A. K. Kerman)

IV. Nucleon Reactions ("High" Energy) and Electron Scattering

A. Quasi-Elastic Contribution to Electron Nucleus Scattering

A calculation of electron scattering on Bi^{209} is being carried through using a shell model for the ground state of Bi^{209} and various assumptions about the outgoing nucleon. This will enable us to obtain a better picture of shell model effects on the shift and width of the quasi-elastic peak in electron nucleus scattering. In addition it is of great interest to know the shell model contributions to the electron scattering with large energy loss because in this region one hopes to see the two-particle correlation function. To obtain this from the data it will be necessary to know the shell model contribution. (S. Klawansky and A. K. Kerman)

B. Angular Distribution of Protons Stripped from High Energy Deuterons

We are investigating the angular distribution of stripped protons obtained by scattering high energy deuterons from nuclei. Previous results for high energy deuteron stripping by nuclei^{4, 5, 6} and for scattering of protons by carbon⁷ are being applied. (V. Franco and J. Reading)

C. Deuteron-deuteron collisions at High Energies

The Glauber-Moliere high-energy approximation is used to investigate deuteron-deuteron collisions at high (~500 MeV) energies. Elastic scattering, stripping, break-up, absorption, and total cross sections are calculated. The expressions include double, triple, and quadruple scattering corrections. (V. Franco)

D. Two Solutions of High Energy Multiple Scattering Problem

The R. Glauber approximation and the optical potential (A. K. Kerman, H. McManus, R. M. Thaler) are compared as methods of solving the particle-nucleus scattering problem. (J. F. Reading)

E. Deuteron-Carbon Elastic Scattering at 650 MeV

A calculation of the deuteron-carbon elastic differential cross section using the Glauber approximation has been performed and compared with experiment. (J. F. Reading)

F. Effect of Repulsive-Core Correlation

The effect of repulsive-core correlation in nuclei on optical-potential is being considered and calculated in a naive approximation. (J. F. Reading)

G. Production of Composite Particles in High-Energy Nucleon-Nucleus and γ -Nucleus Collisions

A rather detailed literature search for experimental and theoretical papers on the reactions (n, d), (p, d), and (γ , d) has essentially been completed. It has been found generally, in many experiments over the last 15 years, that in such reactions an unusually large number of deuterons are observed at all energies from a few MeV up to several BeV. Although many of the experiments contain questionable data because of the experimental difficulties involved, there are certain fairly well-established conclusions:

1. A statistical model of nuclear reactions cannot explain the large number or the angular distribution of the deuterons at any energy.
2. An important contribution to the reactions appears to come from one-stage or multi-stage pickup mechanisms.
3. There are significant variations in the cross sections for deuteron emission among neighboring nuclei, suggesting nuclear structure effects.
4. The A dependence of the cross section at constant energy is less than linear ($A^{2/3}$ or $A^{1/3}$ perhaps) suggesting a surface effect may be important.
5. The large number of deuterons even at hundreds of MeV incident energies suggest strong correlations among nuclear particles.

Initially we are concentrating on the (γ , d) reaction. Since a photon has a low momentum in relation to its energy, it has difficulty giving up its energy to a nuclear particle unless a second nucleon is available to satisfy momentum conservation. Therefore, results at high energy should be sensitive to the effects of correlations among nuclear particles. Current work will be oriented to an investigation of the effects of pair correlations on the (γ , d) reaction. (L. R. Peterson and R. H. Lemmer)

V. Nuclear Reactions ("Low"Energy)

A. Intermediate Resonances and Doorway States

A formulation which connects intermediate resonances and doorway states has been developed. The assumption is made that there exist states, called "doorway states", which connect directly to the open channels and to more complicated structures; the latter are not directly connected to the open channels.

Such a circumstance leads to intermediate resonances if the width of the doorway state is not too large. That width is shown to be composed of two parts, which are: the width for decay into the open channel; and the width for decay into the more complicated excitation.

The appropriate definitions of these various widths have been obtained, and some qualitative applications to isobar analogues, dipole giant resonances, etc., are being investigated.

(R. Lemmer, A. Kerman, H. Feshbach)

B. On the Damping of Isobar Analogue States

We are making an attempt to compute the isotopic spin mixing into the excited isobaric analogue state of Yt^{89} . This involves calculating matrix elements of Coulomb force between the analogue state and the large number of two-particle one-hole proton excitations of different isotopic spin. Most of these matrix elements are in the range of less than 10 keV and this implies that the damping will be of this order of magnitude. There is one large matrix element to the state made of the same configuration as the analogue state. However, this state is always far away in energy and makes only a small contribution to the damping. (A. F. R. deToledo and A. K. Kerman)

C. Role of Collective Excitations in the Neutron Strength Function

The theoretical study of the role played by collective nuclear states in the resonance scattering of neutrons, as described in the previous progress report, continues. Definite conclusions and results have not yet been obtained.

To recapitulate briefly, we are studying in some detail the scattering of neutrons from spherical even-even nuclei in the vibrational region, $A=58$ to $A=80$. It is hoped thereby to obtain a clearer understanding of the deviations of the s-wave neutron strength function from predictions of the standard optical model in that mass region.

Recent work on the construction of the optical potential, has suggested that the strength function fluctuations might be fitted by a formula appropriate to resonance scattering through a single intermediate state consisting of a particle plus vibration. (It has been found that only one such state occurs in the right energy range for the nuclei of interest.) The essential point in making this argument is the fact that such an intermediate state, in addition to an escape width for neutron emission, has a width for damping into more complex states. A single-level Breit-Wigner form for the strength function can be written in which the damping width plays the role of an absorption width. As the other quantities occurring in this formula, namely the resonance energy and escape width, are known roughly for the nuclei in question (and can be calculated to greater accuracy), we can try to fit the strength function by regarding the damping width as an adjustable parameter.

It was pointed out in the last progress report that intermediate states consisting of particle plus vibration are strongly coupled to the incident channel. For this reason the proper treatment of the elastic scattering involves the simultaneous solution of the coupled equations. This has been done for a simple schematic form of coupling analogous to that used for scattering through particle-hole states.⁸ (H. Picker and R. Lemmer)

D. T=2 Excitation of O

Current configuration mixing calculations for the lowest even-parity excited states in O^{16} have been specialized to $J=0^+$, $T=2$. These states are of interest because a narrow resonance with these quantum numbers has been observed by Garvey, Cerny and Pehl in the $O^{18}(p, t)O^{16}$ reaction.⁹ If the $T=2$ assignment is correct, particle decays can occur only by violation of isospin conservation, and this would account for the narrow width. It is interesting to study the shell model description of this excitation, which must be a mixture of two-particle, two-hole configurations. Estimates of the energies of the possible $J=0^+$, $T=2$ configurations indicate that the lowest state will be primarily a mixture of $(1p_{1/2})^{-2}(2s_{1/2})^2$ and $(1p_{1/2})^{-2}(1d_{5/2})^2$. The mixing of these two configurations has been calculated with the zero-range interaction of Brown, *et al.*¹⁰ The results show that these two configurations are strongly mixed, and the lower of the resulting excitation energies is very near the 22.9 MeV reported by Garvey *et al.* If this description of the $T=2$ state is correct, radiative decay will be almost entirely by E1 radiation to the lowest 1^- , $T=1$ particle-hole excitation. (H. P. Jolly)

E. Alpha Inelastic Scattering on Even Isotopes

In collaboration with N. K. Glendenning (UCLRL, Berkeley) calculations of inelastic $\alpha-\alpha'$ scattering on even single closed shell nuclei are in progress. Ingredients are the distorted wave Born approximation, a finite range interaction between the projectile and the nucleons of the target, and two quasi-particles configuration mixing for all the states of the nucleus. Comparison is made with the usual analysis in terms of the vibration description of the nuclear states. (M. Veneroni)

F. Calculation of Damping Width for 2p-1h States in Spherical Nuclei

The damping width down into the compound nucleus is being calculated under the assumption that it is small compared to the escape width. The results for a δ -function force have been expressed in terms of the appropriate 3-j and 9-j symbols and are now being coded. A preliminary calculation for a Pb^{208} target has indicated that there is no damping from the 2p-1h states to the 3p-2h states. This is due to the large energy gap between any 2p-1h state and the 3p-2h states which connect to it. (V. Franco and R. H. Lemmer)

G. Elastic Neutron Scattering from a Pb^{208} Target

Feshbach's unified theory of nuclear reactions is being applied to elastic neutron scattering from Pb^{208} . The elastic scattering total cross section and angular distributions are being calculated as a function of the incident neutron energy. We are using the two-body-function force $V_0(\pi_t + p\pi_s) \delta(\vec{\gamma}_1 - \vec{\gamma}_2)$ with

$$\pi_t = \frac{3 + \vec{\sigma}_1 \cdot \vec{\sigma}_2}{4}$$

and

$$\pi_s = \frac{1 - \vec{\sigma}_1 \cdot \vec{\sigma}_2}{4}$$

where p is a constant. (V. Franco and R. H. Lemmer)

H. A Theoretical Investigation of the Triple Scattering

A theoretical investigation of the triple scattering parameters P, D, R and A for neutrons scattered from non-zero spin nuclei is being made using the optical model. Spin-spin and spin-orbit forces are included. (A. P. Stamp)

I. Nuclear and Photonuclear Reactions in Light Nuclei

A completely antisymmetrized formalism for nuclear reactions has been developed in which we have started from Villars scattering amplitude $|A+1, 0, k_0^+ \rangle$ in configuration space, and evaluated what G. M. Vagnadov calls the channel function $f_n(q)$ where

$$f_n(q) = \langle A, n | \psi(q) | A+1, 0, k_0^{(+)} \rangle \quad (1)$$

$\langle A, n |$ is taken as the residual nucleus in Hartree-Fock (H.F.) basis, while $\psi(q)$ is the field operator for a nucleon.

We then find that $f_0(q)$, the elastic channel function satisfies an integro-differential equation in which the open channel has been subtracted as an intermediate state. Solving the integro-differential equation for $f_0(q)$, and taking the limit as $q \rightarrow \infty$ gives the usual form for the scattering amplitude.

$$T = T_p + \sum_{\gamma} \frac{R_{\gamma}}{E - \text{Re}(E_{\gamma}) + i\Gamma_{\gamma}/2} \quad (2)$$

where T_p is the amplitude for potential scattering. Apart from a phase factor, R_{γ} is the width Γ_{γ} , and the sum is over all resonances in the vicinity of E.

The width Γ_{γ} is given by

$$\Gamma_{\gamma} = 2\pi \left| \int dg \langle \Psi_{\gamma} | P_c J(q) | A, 0 \rangle \chi^{(+)}(q) \right|^2 \quad (3)$$

where Ψ_{γ} is an eigenstate of the complex operator $= H + \Delta$ (H is the total Hamiltonian), i.e.,

$$\Psi_{\gamma} = E_{\gamma} \Psi_{\gamma} \quad (4)$$

where

$$\Delta = \int dq' P_c J(q') | A, 0 \rangle \frac{1}{\epsilon_{k_0}^{(+)} - H_0(q')} \langle A, 0 | J^+(q') P_c$$

P_c is a projection operator that excludes the continuous spectrum of the (A+1) nucleon states, $J(q)$ is the current defined by Villars while $\chi^{(+)}$ is the out-going solution that gives asymptotically the amplitude for potential scattering T_p . We are now extending the above formalism to include (n,n') and (γ ,n) reactions, for application to nuclear and photonuclear reactions in light nuclei. (I. Afnan and R. Lemmer)

J. Optical Model Calculations

Some calculations begun at Brookhaven have been continued. Among these are re-calculation of the 80 MeV π -C scattering experiments of Rainwater, et al, using a Kisslinger-type potential which takes into account the p-wave nature of the π -nucleon scattering process. A

more extensive calculational survey of scattering of heavy ions by medium and heavy nuclei, following the work reported by Auerbach and Porter at the Conference on Reactions between Complex Nuclei has been completed.

An abstract was submitted for the New York American Physical Society Meeting covering continuing work in collaboration with L. Rosen and others at Los Alamos on the polarization resulting from elastic scattering of 10.5 and 14.5 MeV protons from 50 nuclei. The average potentials thus derived describe scattering in the 4-23 MeV range quite well. (E. H. Auerbach)

K. The Effects of Pauli Principle on the Stripping and Pick-up Process

The formulation of the (d,p) process, including the effects of the identity of the particles in the deuteron and those in the target nucleus is being considered. (L Dohnert)

VI. Finite Nuclei

A. Hartree-Fock Calculations in Spherical Nuclei

We have completed and are now testing a computer program for doing a complete Hartree-Fock calculation for closed-shell spherical nuclei using an arbitrary but finite local two-body force. No firm results have been obtained yet. This work is being done in collaboration with Andrew M. Lockett of the Los Alamos Scientific Lab. (A. K. Kerman)

B. Hartree-Fock Calculation with a Non-local Nuclear Force

Calculation of the Hartree-Fock self-consistent single-particle well is proceeding (see Progress Report, May 1964). Computer programs are being written and tested to calculate the matrix elements of the two-body potential with respect to harmonic oscillator states, and to perform the iterations and diagonalizations necessary to achieve a self-consistent solution. Preliminary results indicate that the method may have some validity.

In the present form, the programs are specific to the separable potential of Tabakin¹¹ and to closed shell nuclei. More general cases will also be considered. (J. P. Svenne)

VII. Nuclear Matter

A. Energy Gap at the Fermi Surface of Nuclear Matter

On the basis of the Brueckner-Goldstone-Bethe Theory of nuclear matter, where the single particle potential is defined to cancel third-order self-energy insertions, it was found that there is an energy gap of order 13 MeV (paper in press). A mistake in the original calculation has been corrected leading to the above result. The calculation has been reconsidered in the light of Bethe's new suggestion, that $U(k)$ should compensate for all three body clusters. It turns out that the gap will be reduced to about 3 MeV, the latter a rough estimate based on (i) a factor of

1/3 from suppression of three body clusters and (ii) central forces are about 1/3 less effective than tensor forces in generating an off energy shell contribution to the energy denominator.

(D. Sprung)

B. Properties of Nuclear Matter

A calculation of the properties of nuclear matter according to the boundary condition model of nuclear forces is in progress. The reference spectrum approximation as modified by Razavy and Sprung¹² is used. (D. Sprung, in collaboration with P. C. Bhangava, McMaster University).

REFERENCES

1. T. K. Kuo and S. Yao, Phys. Rev. Letters 13, 415 (1964).
2. J. B. Bronzan and F. E. Low, Phys. Rev. Letters 12, 522 (1964).
3. H. Goldberg and E. Lomon, Phys. Rev. 134, 659 (1964).
4. V. Franco, Ph.D. Thesis (Harvard University), unpublished.
5. V. Franco and R. J. Glauber, to be published
6. R. J. Glauber, private communication
7. J. Reading, Ph.D. Thesis (University of Birmingham), unpublished
8. R. H. Lemmer, Phys. Letters 4, 205 (1963).
9. G. T. Garvey, J. Cerny and R. Pehl, Phys. Rev. Letters 13, 548 (1964).
10. G. E. Brown, L. Castillejo and J. A. Evans, Nucl. Phys. 22, 1 (1961).
11. F. Tabakin, Ann. Phys. 30, 51 (1964).
12. Razavy and Sprung, Phys. Rev. 133, B300 (1964).

PUBLICATIONS

H. Feshbach and E. Lomon, "The Boundary Condition Model of Strong Interactions", Ann. Phys. 29, 19 (1964).

THESIS

R. P. Perrin, Elimination of the Infrared Divergence from Quantum Electrodynamical Dispersion Relations, Ph.D. Thesis, September 1964.

APPENDIX

PERSONNEL LISTING BY PROJECTS

(as of November 1, 1964)

Visiting Committee

Prof. S. D. Drell
Department of Physics
Stanford University
Stanford, California

Prof. W. A. Fowler
Physics Department
California Institute of Technology
Pasadena 4, California

Prof. H. E. Gove
Department of Physics
University of Rochester
Rochester, New York

Professor Kenneth Greisen
Laboratory of Nuclear Studies
Cornell University
Ithaca, New York

Dr. Edward J. Lofgren
University of California
Lawrence Radiation Laboratory
Berkeley, California

Prof. D. H. Wilkinson
Department of Nuclear Physics
Nuclear Physics Laboratory
University of Oxford
21 Banbury Road
Oxford, England

MIT Interdepartmental Laboratory Committee for the Laboratory for Nuclear Science

P. T. Demos
Professor of Physics
Director of the Laboratory

M. Benedict
Professor of Nuclear Engineering
Head of the Department

W. W. Buechner
Professor of Physics
Head of the Department

A. C. Cope
Professor of Chemistry
Head of the Department

J. B. Wiesner
Institute Professor
Dean of the School of Science

Laboratory Committees

IBM 7044 Computer

Prof. E. Lomon, Chairman
Prof. B. T. Feld
Prof. H. Feshbach
Prof. L. Osborne
Prof. I. Pless

Inner Space

Prof. D. Frisch, Chairman
Prof. W. Buechner
Prof. M. Deutsch
Prof. B. T. Feld
Prof. H. Feshbach

Reading Room

Prof. B. T. Feld, Chairman
Prof. A. Bernstein
Prof. M. Deutsch
Prof. J. I. Friedman
Prof. A. Kerman
Prof. W. Kraushaar

Personnel Listing

Headquarters

Prof. P. T. Demos
Director

Dr. F. J. Eppling
Associate Director

R. L. Loria
Headquarters

B. Bailey
Engineering

R. W. Calileo
General Services

E. DeAgazio
Electronics Shop

R. Gustavson
Machine Shop

J. L. Sargeant
Purchasing

A. J. Scully
Accounting

S. A. Tardivo
Property

C. W. Tourtellotte
Drafting

RESEARCH GROUPS

Chemistry of the Fission Elements Group

Research Staff

Prof. D. N. Hume (on leave)
Prof. D. M. Hercules
Prof. W. E. Ohnesorge
Prof. D. K. Roe
R. C. Lansbury
G. D. Short
J. E. A. Toni

Guest

T. W. Gilbert

Graduate Students

N. R. Andersen
P. D. Anderson
D. N. Bailey
R. C. Berlo
R. L. Birke
D. M. Desiderio
Adrienne Gordon
*J. M. Hayes
*R. M. King
W. M. Krebs
J. N. Little
A. L. Malenfant
T. F. Retajczyk
P. Vouros
J. T. Watson

Undergraduate Students

*W. R. Matson
*P. J. Sherwood
Mrs. Barbara T. Vickers

*Terminated

Personnel Listing

Nuclear Chemistry (Inorganic) Group

<u>Research Staff</u>	<u>Guests</u>	<u>Graduate Students</u>
Prof. C. D. Coryell	Dr. H. Amano	S. Basol (Ford Founda-
Prof. G. E. Gordon	Division of Radio Isotope Prod.	tion Fellow)
Prof. J. W. Irvine, Jr.	Takai Research Establishment	*C. E. Bemis
Prof. G. Scatchard	Japan Atomic Energy Research Inst.	*R. P. Black (NIH
*Dr. A. S. Kertes	Takai-Mura, Japan	fellow)
*Dr. K. Kimura		*D. S. Brenner (NIH
Dr. R. H. McCorkell	Prof. L. Leifer	Fellow)
Dr. R. A. Rousseau	Department of Chemistry	J. F. Byrum
Dr. G. Scibona	Boston College	J. E. Canty
Dr. W. B. Walters	Boston, Massachusetts	*R. A. Duce
*Dr. W. T. Yap		A. M. Ehrlich
		J. L. Fasching
		H. F. Gibbard, Jr.
		D. J. Hnatowich
		H. Nakahara
		D. A. Muga
		B. Parsa
		R. C. Ragaini
		C. Schlaikjer
		*C. D. Strehlow
		C. -W. Tang
		J. F. Wild
		R. L. Yoest

Nuclear Chemistry (Organic) Group

<u>Research Staff</u>	<u>Graduate Students</u>	<u>Undergraduate Student</u>
Prof. F. D. Greene, II	R. V. Cartwright	F. J. Weigert
Prof. C. G. Swain	R. D. Cramer	
Dr. A. L. Powell	J. A. Eckert	
Dr. K. W. Bowers	Anna Fang	
Dr. C. R. Morgan	K. G. Harbison	
*Dr. M. F. Grondon	Mrs. Judith M. Harbison	
Dr. A. Ohno	*Dr. W. Henderson	
	Mrs. Nancy N. Lowry	
	T. J. Lynch	
	*Mrs. Barbara S. Magid	
	G. J. Nolfi, Jr.	
	M. H. O'Leary	
	J. Pazos	
	*Mrs. Barbara J. Schowen	
	J. E. Sheats	
	H. Stein	
	J. T. Viola	
	Joanne Worosz	

*Terminated

Cosmic Ray Group

<u>Research Staff</u>	<u>Graduate Students</u>	<u>Undergraduate Students</u>
Prof. B. B. Rossi	J. Binsack	R. Chang
Dr. H. S. Bridge	K. Brecher	T. Dally
Prof. H. Bradt	F. Floyd	C. Forline
Prof. G. W. Clark	I. Glass	H. Houranian
Dr. J. Davis	D. Grimes	C. Leong
Dr. A. Egidi	P. Higbie	P. Oddone
Dr. G. Garmire	H. Hinteregger	P. Scheffler
Dr. G. Gordon	*M. Joseph	A. Tanenbaum
Prof. W. L. Kraushaar	A. Klimas	K. Yip
Prof. A. J. Lazarus	A. Krieger	
Dr. M. La Pointe	E. F. Lyon, III	
Dr. J. D. Linsley	W. Meyer	
LNS Cosmic Ray Station	S. Rappaport	
Albuquerque, N.Mex.	*A. Russell	
Dr. G. Moreno	D. Slevin	
Dr. M. Oda	J. Stein	
Prof. S. Olbert	R. Thorne	
Dr. L. Pai	V. Vasyliunas	
Prof. F. Scherb	A. Womack	
W. B. Smith		
*D. Stowell		
Dr. M. Wada		

Accelerator Physics Collaboration (formerly Bubble Chamber Group)

<u>Research Staff</u>	<u>Graduate Students</u>	<u>Undergraduate Students</u>
Dr. P. Bastien	B. Brabson	S. Bronstein
S. Brudno	*J. Campbell	L. Cox
*M. Donahue	L. Kirkpatrick	W. Podolsky
L. Dunn	B. Nelson	C. Reeve
*Dr. Y. Eisenberg	R. Panvini	
Prof. B. T. Feld	M. Partovi	
Dr. V. K. Fischer	A. Rogers	
Prof. I. A. Pless	C. Rogers	
Prof. A. H. Rosenfeld	K. Schneider	
Prof. L. Rosenson	J. Wolfson	
B. Wadsworth		
Dr. T. Watts		
*C. Laywine		

Linear Accelerator Group

<u>Research Staff</u>	<u>Graduate Students</u>	<u>Undergraduate Students</u>
Prof. P. T. Demos	F. Hanser	R. Rudziaski
Prof. W. Bertozzi	J. Matthews	D. Trevvett
Dr. C. P. Sargent	G. Mutchler	A. Woreck
Dr. W. Turchinetz	T. Phillips	
Dr. S. Kowalski	E. Reinhardt	
*Dr. I. Hayashi	J. Bergstrom	
	M. Zombek	
	G. Hurford	

*Terminated

Personnel Listing

High Energy Accelerator Physics Group

Research Staff

Dr. R. Alvarez
Prof. B. T. Feld
Dr. R. Fessel (CEA)
Prof. J. I. Friedman
Prof. D. H. Frisch
Dr. D. Garelick
Dr. G. Glass
Dr. S. Homma
Prof. W. H. Kendall
W. Lobar
Dr. D. Luckey
Prof. L. Osborne
*Dr. L. Sodickson
C. J. Strumski
Dr. M. Wahlig

Visitors

Dr. Z. Bar-Yam (SMTI)
Dr. W. Kern (SMTI)

Graduate Students

A. Broadfoot
A. Buffington
K. J. Cohen
G. Cooperstein
J. Elias
V. B. Elings
O. Fackler
D. Gordon (Brandeis)
G. Hartman
P. Kirk
R. A. Lewis
R. Logan
P. Mockett
E. I. Shibata
S. Smith
M. Sogard
L. M. Sompayrac
R. W. Stimets
J. R. Uglum
L. Van Speybroeck
C. Ward

Undergraduate Students

W. R. Alves
S. P. Apsell
D. Bellenger
M. Breidenbach
D. Cannell
S. B. Deutsch
D. J. Ernst
R. Fisher
B. Goodman
S. W. Gray
E. Groves
R. J. Long
J. R. Nagy
M. P. Ryan
S. Schutz
A. R. Stottlemeyer
M. W. Strovink
M. Teague

ONR Generator Group

Research Staff

Prof. W. W. Buechner
Prof. H. A. Enge
Prof. T. A. Belote
H. Y. Chen
Dr. W. E. Dorenbusch
Dr. F. J. Eppling
O. Hansen
*P. Hanssen
Dr. W. H. Moore
Dr. D. J. Pullen
M. K. Salomaa
A. Sperduto

Visitors

*A. Graue
University of Bergen
Bergen, Norway
*Dr. C. A. Wiedner
University of Heidelberg
Heidelberg, Germany

Graduate Students

*K. M. I. Abdo
*M. M. Castro
E. Chalbaud
E. Cosman
*K. A. -A. Hamza
G. Nair
M. N. Rao
*D. M. Sheppard
J. E. Spencer
Helen J. Young

Undergraduate Students

M. Keehner
M. Skerry
D. Schramm
A. Spearer
J. Ottesen
S. Scott
Louise Wilson

*Terminated

Radioactivity Group

<u>Research Staff</u>	<u>Graduate Students</u>	<u>Undergraduate Students</u>
Prof. R. D. Evans	J. Alonso	M. Slade
Prof. L. Grodzins (on leave)	A. Buyrn	M. Schuster
Universitets Institut for	Y. W. Chow	
Teoretisk Fysik	D. Murnick	
Copenhagen, Denmark	Gatti Pramilla	
Prof. R. Bauer	W. Schick	
Dr. S. G. Cohen	D. Yeboah-Amankwah	
Dr. H. W. Kraner		

Cyclotron Group

<u>Research Staff</u>	<u>Graduate Students</u>
*Dr. N. S. Wall	R. Ginaven
Prof. A. Bernstein	P. Lippincott
*Prof. R. Bauer	E. Martens
*Dr. G. Heymann	*E. Romberg
E. F. White	*P. Roos
W. J. Kossler	S. Smith
T. Provost	*R. Brown
	H. Tannenbaum
	M. Slade
	*S. Dostrovsky

Theoretical Group

<u>Research Staff</u>	<u>Graduate Students</u>	
Prof. V. F. Weisskopf (on leave)	I. Afnan	R. M. Saenger
CERN	A. K. Bhattacharyya	E. Salz
Geneva, Switzerland	M. L. Blackmon	C. Schmid
Prof. F. Villars (on leave)	C. N. Bressel	J. P. Svenne
Nuclear Research Center	J. D. Cornwell	N. M. Tomljanovich
Saclay, France	A. deToledo	R. P. Zia
Prof. H. Feshbach	L. Dohnert	
Dr. E. Auerbach	C. Dover	
Dr. J. Bronzan	J. P. Dowd	
Prof. P. Federbush	M. Feinroth	
Dr. D. Flamm	T. Fitzgerald	
Dr. V. Franco	W. A. Friedman	
Prof. K. Huang	J. Harte	
Dr. J. Irwin	A. Huffman	
Prof. K. Johnson (on leave)	P. N. Kirk	
Institute for Theoretical Physics	S. Klawansky	
Copenhagen, Denmark	R. Logan	
Dr. P. Jolly	A. I. Miller	
Prof. A. Kerman	A. H. Mueller	
Prof. R. Lemmer	F. M. Newman	
Prof. E. Lomon	V. J. Newton	
Prof. F. Low	R. Perrin	
Dr. J. Reading	L. R. Peterson	
Prof. R. Sawyer	H. Picker	
Dr. D. Sprung	H. Quang	
Dr. A. Stamp	E. F. Redish	
Dr. M. Veneroni	N. Rogerson	
Dr. T. Yao	A. I. Rubenstein	

*Terminated