Radiative Transition Probabilities
for the $A^1\Sigma^+ - X^1\Sigma^+$ Bands of $^7\text{LiH}^*$

Warren T. Zemke
Department of Chemistry
Warburg College
Maverly, IA 50607

NOTICE
This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

and

MASTER

William C. Stwalley
Departments of Chemistry and Physics
University of Iowa
Iowa City, IA 52242

*Supported by the United States Energy Research and Development Administration [Contract No. EY-76-S-02-2326.*001].

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

ABSTRACT

The recently reported (J. Chem. Phys. 66, 5412 (1977)) hybrid potential curves for the $X^1\Sigma^+$ and $A^1\Sigma^+$ states of LiH are combined with the ab initio transition moments of Docken and Hinze (J. Chem. Phys. 57, 4936 (1972)) to calculate radiative transition probabilities between all upper ($v' \leq 26$) and all lower ($v'' \leq 23$) vibrational levels of ^7LiH . By far the strongest single band found is the 23-26 band between the highest "long-range" levels in each potential. A comparison with experimental intensity observations is made; it appears that the experiments are globally in accord with our calculations, but with some detailed differences discussed herein.

I. INTRODUCTION

The simplest neutral heteronuclear molecule, LiH, has been the object of intense theoretical and spectroscopic interest¹ since the 1930s. We have embarked on a long-term project to fully understand the spectroscopy of the three lowest singlet states ($X^1\Sigma^+$, $A^1\Sigma^+$ and $B^1\Pi$) of LiH for a variety of reasons discussed elsewhere.²

As an initial step in this project, we have constructed full potential curves (Figure 1) for the $X^1\Sigma^+$ and $A^1\Sigma^+$ states of ^7LiH and from these curves calculated the vibrational (and vibrotational) energy levels of the isotopic lithium hydrides.² In this paper we summarize our calculations for the radiative transition probabilities between all vibrational levels of the $X^1\Sigma^+$ ($0 \leq v'' \leq 23$) and the $A^1\Sigma^+$ ($0 \leq v' \leq 26$) states of ^7LiH and compare with available experimental data. A more complete compilation of the results of these calculations is also available.³ Subsequent publications will report ^7LiH calculations of the lifetimes of all vibrational levels in the $A^1\Sigma^+$ state,⁴ the transition probabilities of the $B^1\Pi - X^1\Sigma^+$ and $B^1\Pi - A^1\Sigma^+$ bands⁵, the lifetimes of all vibrational levels in the $B^1\Pi$ state,⁶ the IR transition probabilities for the $X^1\Sigma^+$ state,⁷ and a detailed examination of the "long-range molecules"⁸ corresponding to $X^1\Sigma^+$ ($v=23$), $A^1\Sigma^+$ ($v=26$) and $B^1\Pi$ ($v=2$).⁹

In the present manuscript, we first examine the appropriate formulae for absorption and emission transition probabilities in the A-X bands of LiH in Section II. In Section III we discuss construction of the dipole strength function (electronic transition dipole moment as a function of internuclear distance). Results of the transition probability calculations are summarized in Section IV. In Section V we discuss the physical inter-

Figure 1. Potential energy curves from Ref. 2 for the $X^1\Sigma^+$ and $A^1\Sigma^+$ state of ^7LiH , including calculated vibrational levels $v'' = 0 - 23$ and $v' = 0 - 26$. The Franck Condon regions for $v'' = 0$ (— — —) and $v' = 0$ (— — —) are indicated. The curves are parallel at $\sim 7.3a_0$, implying a classical satellite and enhanced emission e.g. for $A(v'=5) \rightarrow X(v''=18)$. Note also the large outer turning points for $v' = 26$ and $v'' = 23$. See Figure 3 for comparison.

C-674-387-1

pretation of these results and compare with other (more limited) theoretical and especially experimental results.

II. RADIATIVE TRANSITION PROBABILITIES

There are four basic kinds of spectroscopic intensity measurement, depending on whether absorption or emission is involved, and whether photon or energy flux is detected. The absolute intensities in each case involve a power of the transition frequency $\nu_{v',J'; v'',J''} = h^{-1}(E'(v', J') - E''(v'', J''))$ times the square of an electronic transition dipole matrix element

$$D_{v',J'; v'',J''} \equiv \int_0^{\infty} \psi_{v',J'}(R) D(R) \psi_{v'',J''}(R) dR$$

(where the upper and lower state wavefunctions $\psi_{v',J'}$ and $\psi_{v'',J''}$ are normalized and where $D(R)$ is the dipole strength function (electronic transition dipole moment as a function of internuclear distance), discussed below in Section III). For the case of absorption measured by a decrease in photon flux, the transition probability is proportional to $|D_{v',J'; v'',J''}|^2$ (no frequency dependence) and commonly expressed as line strength S or Einstein B coefficient. For absorption measured by a decrease in energy flux, the transition probability is proportional to $|D|^2 \nu$; the absorption oscillator strength f or absorption coefficient α is commonly used in that case. For emission, the transition probabilities go as $|D|^2 \nu^3$ and $|D|^2 \nu^4$ for photon and energy detectors, respectively; the Einstein A coefficient is commonly used in the former case. An excellent discussion of the different quantities with conversion factors is given by Wiese, Smith and Glennon.¹⁰ A final quantity of interest is the Franck-Condon factor

$$q_{v',J'; v'',J''} = \left| \int_0^{\infty} \psi_{v',J'}(R) \psi_{v'',J''}(R) dR \right|^2$$

which is proportional to $|D|^2$ when $D(R)$ does not change significantly with R in the regions of appreciable integrand.

We shall restrict our attention here to $J' = J'' = 0$ only (although we have some limited $J > 0$ results); i.e. we shall ignore rotation, and report vibrational transition probabilities. We shall here report only Franck-Condon factors, $q_{v',v''}$; squares of transition dipole matrix elements, $|D_{v',v''}|^2$ (in units of $(e\text{\AA})^2$); and spontaneous photon emission intensities, $I(v',v'')$ (in units of $(e\text{\AA})^2 (\text{cm}^{-1})^3$). A more complete listing³ and conversion factors^{3,10} (e.g. $A_{v',v''}(\text{sec}^{-1}) = 7.2356 \times 10^{-6} I(v',v'') ((e\text{\AA})^2 (\text{cm}^{-1})^3)$) are given elsewhere. The transition frequencies $\nu_{v',v''}$ are available using the energies listed in Ref. 2.

III. THE DIPOLE STRENGTH FUNCTION

The dipole strength function $D(R)$ (i.e. the electronic transition dipole moment as a function of internuclear distance) has been constructed in a manner similar to that used previously for potential curves (e.g. in Ref. 2). The $D(R)$ function consists of three regions, given in Table I.

Region I assumes a constant value of 0.421 a.u. for $D(R)$ for $R < 1.0$, which is well into the classically forbidden region for all v' and v'' and hence of little physical significance. The value of 0.421 for the transition dipole moment was obtained by considering the He atom as the model for $(\text{LiH})^{++}$ with two valence electrons. This "semi-united atom" model seems more reasonable than considering Be at $R=0a_0$ and resembles the work of Mulliken et al.¹¹⁻¹³, where N_2 and CO take on a "semi-united" Mg atom character at $\sim 0.5 - 0.8a_0$ before going to the true Si united atom as $R \rightarrow 0$. Considering the $A - X$ LiH transition at $1a_0$ to be modelled by the sigma component of the $1s^2 \rightarrow 1s2p$ He atomic transition ($f = 0.2762$, $\Delta E = 171135 \text{ cm}^{-1}$, from Ref. 10), one calculates a transition moment of 0.421 a.u. This value matches well with the innermost Region II points, as seen in Figure 2.

Region II is the region of predominant significance in our calculations. It consists of a cubic spline fit to the points given in Table II, which (except at $1.0a_0$) are the highly regarded ab initio results of Docken and Hinze.¹⁴ As discussed elsewhere⁴, the excellent agreement (within mutual uncertainties of $\sim 10\%$) between experimental and theoretical $A^1\Sigma^+$ lifetimes lends additional support for this ab initio dipole strength function. It is our opinion that the $D(R)$ in the physically important region from $2 - 12a_0$ is unlikely to be anywhere in error by as much as 20%. Finally it should be noted that the change in $D(R)$ with internuclear distance is

Table I. The dipole strength function $D(R)$ in atomic units for the $A\ ^1\Sigma^+ - X\ ^1\Sigma^+$ transition in ^7LiH .

Region	Range of R	Function $D(R)$	Reference
I	$R \leq 1.0$	0.421	a
II	$1.0 \leq R \leq 12.0$	Tensioned spline fit	Table II
III	$12.0 \leq R$	2.3425	b

^aFrom He $(1s^2 \rightarrow 1s2p)$ oscillator strength value; see text.

^bThis value is essentially the separated atom limit; see text.

Figure 2. Dipole strength function $D(R)$ (in atomic units) for the $A^1\Sigma^+ - X^1\Sigma^+$ transition of ^7LiH . See Tables I and II for details.

A-G77-785

Table II. The dipole strength function $D(R)$ (in a.u.) for the LiH A-X transition in Region II (see Table I) consists of a cubic spline fit to the points listed below. Except for the point at $1.0a_0$ discussed in the text, all points are from the work of Docken and Hinze.¹⁴

$R(a_0)$	$D(R) (e \cdot a_0)$
1.0	0.421
2.0	0.6247
2.25	0.7034
2.5	0.7855
2.75	0.8714
3.0	0.9599
3.25	1.0530
3.5	1.1516
4.0	1.3739
4.5	1.6323
5.0	1.9448
5.5	2.3100
6.0	2.6956
6.5	3.0202
7.5	3.1453
8.5	2.7686
10.0	2.3613
12.0	2.3425

far from negligible and thus simple approximations involving Franck-Condon factors or R -centroids can not be expected to be quantitative.

In the asymptotic Region III, we have used 2.3425 (the $12a_0$ value of Docken and Hinze¹⁴). This value differs negligibly from the best experimental value¹⁰ of 2.35 a.u. Since the value is so large, we feel justified in neglecting the small long-range change in the dipole strength function. Again, this region is not physically significant except for the highest levels.⁸

IV. RADIATIVE TRANSITION PROBABILITY RESULTS

The Franck-Condon factors ($q_{v',v''}$, unitless); dipole strength function matrix element squares ($|D_{v',v''}|^2$, in $(e\text{\AA})^2$ units); and spontaneous photon emission intensities ($I(v',v'')$, in $(e\text{\AA})^2 (\text{cm}^{-1})^3$ units) as defined in Section II are given in Tables III, IV and V for all vibrational levels v' of the $A\ ^1\Sigma^+$ state and v'' of the $X\ ^1\Sigma^+$ state of ^7LiH . The spontaneous emission intensities $I(v',v'')$ are also displayed in Figure 3. A more complete listing is given elsewhere.³

These calculations have been carried out using an extensively modified version of the Franck-Condon factor program of R. N. Zare and coworkers,¹⁷ based on the Numerov-Cooley numerical solution of the radial Schrödinger equation. Except for the higher levels (discussed elsewhere⁹), the calculations are typically carried out with a $0.005\text{\AA}$ grid from $0.5 - 10.5\text{\AA}$, and have been extensively checked for convergence with respect to grid spacing and inner and outer limits. Our values agree very well with the more limited results of other workers. For example, Halmann and Laulicht¹⁸ evaluated $q_{v',0}$ for $A-X\ ^7\text{LiH}$ for $v' = 0 - 13$ with slightly different potentials and with a narrow grid ($1.2 - 4.7\text{\AA}$). Their $q_{v',0}$ values agree with ours to <0.001 , except for higher v' levels (≥ 7) where our potentials differ and their outermost grid points are perhaps too small. Likewise, our transition moment matrix elements $D_{v',0}$ values differ from those reported by Docken and Hinze¹⁴ (who used their ab initio potentials) for various $v' \leq 12$ by $\lesssim 0.008\text{ ea}_0$. More recent $D_{v',v''}$ values for $v' = 5-10$ (also in ea_0 units) from Oppenheimer and Docken¹⁹ (using the same ab initio potentials) showed less good agreement, but this is quite reasonable considering the different potentials; e.g. $D_{5,0} = -0.3046$ vs. our -0.3095 ; $D_{10,19} = -0.3751$ vs. our -0.4055 .

Table III. Franck-Condon factors $q_{v',v''}$ (unitless) for the $A^1\Sigma^+ - X^1\Sigma^+$ bands of ^7LiH .

	v''					
	0	1	2	3	4	5
0	.0007	.0043	.0140	.0327	.0607	.0933
1	.0044	.0220	.0545	.0917	.1119	.0978
2	.0145	.0546	.0961	.0999	.0568	.0084
3	.0326	.0883	.0970	.0429	.0005	.0263
4	.0570	.1030	.0552	.0009	.0305	.0674
5	.0833	.0900	.0115	.0188	.0644	.0276
6	.1052	.0567	.0012	.0550	.0384	.0006
7	.1173	.0219	.0239	.0570	.0028	.0323
8	.1182	.0021	.0516	.0271	.0098	.0499
9	.1093	.0028	.0603	.0023	.0387	.0236
10	.0937	.0187	.0464	.0050	.0462	.0005
11	.0754	.0398	.0230	.0259	.0264	.0109
12	.0576	.0574	.0051	.0434	.0047	.0329
13	.0421	.0673	.0001	.0452	.0011	.0375
14	.0296	.0686	.0070	.0329	.0141	.0225
15	.0201	.0634	.0199	.0162	.0297	.0054
16	.0135	.0548	.0331	.0041	.0370	.0001
17	.0088	.0447	.0424	.0000	.0332	.0072
18	.0057	.0350	.0468	.0030	.0228	.0186
19	.0037	.0266	.0464	.0100	.0116	.0266
20	.0024	.0197	.0427	.0174	.0037	.0278
21	.0015	.0143	.0372	.0232	.0003	.0235
22	.0010	.0103	.0309	.0262	.0004	.0167
23	.0006	.0072	.0244	.0259	.0024	.0100
24	.0004	.0048	.0177	.0222	.0042	.0049
25	.0002	.0026	.0101	.0142	.0038	.0019
26	.0000	.0001	.0006	.0010	.0004	.0001
$\sum_{v'} q_{v',v''}$	.999	.981	.899	.744	.657	.674

Table III (cont.)

v'	v''					
	6	7	8	9	10	11
0	.1244	.1452	.1481	.1321	.1034	.0704
1	.0566	.0140	.0007	.0287	.0799	.1207
2	.0063	.0491	.0826	.0639	.0161	.0026
3	.0700	.0559	.0082	.0105	.0613	.0730
4	.0323	.0002	.0390	.0636	.0196	.0050
5	.0022	.0477	.0462	.0014	.0296	.0595
6	.0446	.0389	.0000	.0396	.0423	.0001
7	.0429	.0001	.0374	.0346	.0007	.0448
8	.0047	.0270	.0369	.0002	.0415	.0188
9	.0094	.0432	.0012	.0330	.0225	.0080
10	.0378	.0134	.0174	.0322	.0015	.0394
11	.0356	.0011	.0383	.0018	.0308	.0119
12	.0105	.0224	.0191	.0119	.0272	.0040
13	.0002	.0350	.0002	.0326	.0018	.0288
14	.0132	.0210	.0108	.0219	.0086	.0224
15	.0290	.0031	.0276	.0023	.0270	.0017
16	.0306	.0018	.0259	.0037	.0226	.0061
17	.0191	.0139	.0107	.0182	.0056	.0217
18	.0061	.0244	.0006	.0245	.0004	.0218
19	.0001	.0248	.0027	.0171	.0091	.0090
20	.0023	.0170	.0114	.0059	.0179	.0004
21	.0084	.0077	.0181	.0002	.0183	.0022
22	.0141	.0017	.0188	.0014	.0121	.0086
23	.0164	.0000	.0147	.0056	.0051	.0123
24	.0148	.0008	.0090	.0081	.0011	.0111
25	.0094	.0014	.0041	.0064	.0001	.0065
26	.0011	.0003	.0005	.0011	.0000	.0012
$\sum_{v'=v''} q_{v'v''}$	.642	.611	.630	.603	.606	.612

Table III (cont.)

v'	v''					
	12	13	14	15	16	17
0	.0409	.0197	.0075	.0021	.0003	.0000
1	.1274	.1001	.0584	.0241	.0061	.0007
2	.0496	.1147	.1356	.0979	.0421	.0087
3	.0232	.0032	.0704	.1456	.1301	.0540
4	.0589	.0647	.0064	.0353	.1500	.1566
5	.0124	.0160	.0716	.0272	.0190	.1669
6	.0400	.0455	.0000	.0556	.0412	.0183
7	.0249	.0071	.0545	.0076	.0382	.0417
8	.0112	.0460	.0010	.0421	.0190	.0281
9	.0418	.0006	.0383	.0130	.0266	.0244
10	.0030	.0310	.0153	.0195	.0246	.0169
11	.0186	.0234	.0092	.0298	.0064	.0290
12	.0322	.0006	.0335	.0001	.0331	.0012
13	.0045	.0260	.0058	.0259	.0032	.0289
14	.0070	.0222	.0078	.0189	.0130	.0091
15	.0263	.0008	.0266	.0000	.0248	.0044
16	.0185	.0095	.0128	.0157	.0041	.0224
17	.0016	.0233	.0000	.0216	.0043	.0107
18	.0038	.0153	.0106	.0060	.0186	.0001
19	.0161	.0019	.0197	.0006	.0145	.0101
20	.0194	.0017	.0131	.0100	.0023	.0160
21	.0120	.0101	.0027	.0158	.0010	.0086
22	.0034	.0149	.0002	.0115	.0076	.0010
23	.0000	.0125	.0039	.0042	.0112	.0007
24	.0009	.0070	.0070	.0004	.0089	.0036
25	.0018	.0026	.0057	.0000	.0043	.0039
26	.0005	.0003	.0012	.0000	.0007	.0009
$\sum_{v'=v''} q_{v'v''}$	.600	.621	.619	.631	.655	.667

Table III. (cont.)

v'	v''						$\Sigma q_{v''v'} v''$
	18	19	20	21	22	23	
0	.0000	.0000	.0000	.0000	.0000	.0000	1.000
1	.0000	.0000	.0000	.0000	.0000	.0000	1.000
2	.0004	.0000	.0000	.0000	.0000	.0000	1.000
3	.0069	.0000	.0001	.0000	.0000	.0000	1.000
4	.0523	.0015	.0008	.0000	.0000	.0000	1.000
5	.1722	.0300	.0012	.0005	.0003	.0000	1.000
6	.2106	.1559	.0018	.0065	.0011	.0000	.999
7	.0385	.2772	.0733	.0158	.0001	.0000	.995
8	.0272	.1111	.2851	.0003	.0113	.0007	.971
9	.0241	.0032	.2705	.0893	.0590	.0021	.947
10	.0229	.0190	.0273	.3160	.0345	.0003	.883
11	.0134	.0180	.0027	.1928	.0140	.0026	.681
12	.0273	.0161	.0180	.0264	.1100	.0065	.601
13	.0002	.0207	.0311	.0441	.0649	.0022	.549
14	.0234	.0002	.0077	.0911	.0624	.0012	.537
15	.0121	.0194	.0004	.0046	.1685	.0026	.536
16	.0010	.0113	.0180	.0004	.1082	.0001	.455
17	.0168	.0003	.0068	.0272	.0474	.0008	.386
18	.0140	.0121	.0003	.0002	.1036	.0006	.395
19	.0010	.0134	.0093	.0001	.0459	.0050	.325
20	.0035	.0028	.0104	.0097	.0187	.0103	.287
21	.0116	.0007	.0031	.0055	.0433	.0117	.281
22	.0111	.0063	.0001	.0022	.0034	.0271	.231
23	.0049	.0089	.0028	.0000	.0095	.0345	.218
24	.0008	.0063	.0049	.0011	.0031	.0648	.208
25	.0000	.0026	.0036	.0016	.0000	.1473	.234
26	.0000	.0004	.0007	.0004	.0001	.5640	.570
$\Sigma q_{v'v''} v''$	.696	.737	.780	.836	.909	.884	

Table IV. Dipole strength function matrix element squares $|D_{v',v''}|^2$ (in $(e\text{\AA})^2$ units) for the $A^1\Sigma^+ - X^1\Sigma^+$ bands of ^7LiH .

v'	v''					
	0	1	2	3	4	5
0	2.717(-4)	1.935(-3)	7.150(-3)	1.901(-2)	4.003(-2)	6.971(-2)
1	1.667(-3)	9.509(-3)	2.696(-2)	5.156(-2)	7.133(-2)	7.044(-2)
2	5.243(-3)	2.285(-2)	4.593(-2)	5.424(-2)	3.478(-2)	5.667(-3)
3	1.136(-2)	3.562(-2)	4.464(-2)	2.234(-2)	2.745(-4)	1.806(-2)
4	1.910(-2)	3.995(-2)	2.441(-2)	4.232(-4)	1.773(-2)	4.385(-2)
5	2.680(-2)	3.355(-2)	4.823(-3)	9.308(-3)	3.565(-2)	1.699(-2)
6	3.241(-2)	2.028(-2)	5.357(-4)	2.593(-2)	2.024(-2)	3.867(-4)
7	3.458(-2)	7.490(-3)	9.554(-3)	2.568(-2)	1.315(-3)	1.890(-2)
8	3.327(-2)	6.881(-4)	1.969(-2)	1.162(-2)	4.993(-3)	2.766(-2)
9	2.934(-2)	8.757(-4)	2.197(-2)	9.238(-4)	1.848(-2)	1.237(-2)
10	2.397(-2)	5.609(-3)	1.617(-2)	2.037(-3)	2.099(-2)	2.211(-4)
11	1.834(-2)	1.140(-2)	7.651(-3)	9.996(-3)	1.142(-2)	5.533(-3)
12	1.330(-2)	1.571(-2)	1.616(-3)	1.600(-2)	1.927(-3)	1.577(-2)
13	9.224(-3)	1.755(-2)	3.527(-5)	1.590(-2)	4.747(-4)	1.710(-2)
14	6.146(-3)	1.706(-2)	2.043(-3)	1.107(-2)	5.497(-3)	9.784(-3)
15	3.959(-3)	1.502(-2)	5.562(-3)	5.230(-3)	1.106(-2)	2.223(-3)
16	2.502(-3)	1.238(-2)	8.841(-3)	1.259(-3)	1.317(-2)	5.758(-4)
17	1.541(-3)	9.600(-3)	1.083(-2)	3.088(-8)	1.131(-2)	2.847(-3)
18	9.410(-4)	7.172(-3)	1.142(-2)	8.799(-4)	7.439(-3)	7.033(-3)
19	5.694(-4)	5.184(-3)	1.085(-2)	2.754(-3)	3.620(-3)	9.622(-3)
20	3.478(-4)	3.697(-3)	9.651(-3)	4.653(-3)	1.109(-3)	9.755(-3)
21	2.115(-4)	2.573(-3)	8.063(-3)	5.946(-3)	7.468(-5)	7.950(-3)
22	1.299(-4)	1.772(-3)	6.459(-3)	6.472(-3)	1.331(-4)	5.462(-3)
23	7.952(-5)	1.192(-3)	4.916(-3)	6.200(-3)	6.803(-4)	3.165(-3)
24	4.712(-5)	7.596(-4)	3.451(-3)	5.160(-3)	1.142(-3)	1.522(-3)
25	2.349(-5)	3.972(-4)	1.929(-3)	3.228(-3)	1.025(-3)	5.637(-4)
26	1.091(-5)	1.872(-4)	9.292(-4)	1.613(-3)	5.700(-4)	2.292(-4)

Table IV (cont.)

v'	v''					
	6	7	8	9	10	11
0	1.054(-1)	1.396(-1)	1.619(-1)	1.646(-1)	1.468(-1)	1.140(-1)
1	4.592(-2)	1.263(-2)	8.681(-4)	3.532(-2)	1.107(-1)	1.896(-1)
2	5.241(-3)	4.464(-2)	8.426(-2)	7.318(-2)	2.036(-2)	4.227(-3)
3	5.358(-2)	4.774(-2)	7.564(-3)	1.226(-2)	7.806(-2)	1.039(-1)
4	2.326(-2)	2.271(-4)	3.731(-2)	6.729(-2)	2.266(-2)	7.414(-3)
5	1.696(-3)	3.848(-2)	4.101(-2)	1.218(-3)	3.493(-2)	7.697(-2)
6	3.058(-2)	2.933(-2)	2.351(-5)	3.918(-2)	4.558(-2)	7.090(-5)
7	2.771(-2)	5.113(-5)	3.144(-2)	3.144(-2)	9.677(-4)	5.331(-2)
8	2.754(-3)	1.942(-2)	2.878(-2)	3.184(-4)	4.181(-2)	2.021(-2)
9	5.876(-3)	2.901(-2)	8.021(-4)	2.861(-2)	2.074(-2)	9.154(-3)
10	2.197(-2)	8.368(-3)	1.300(-2)	2.585(-2)	1.656(-3)	4.016(-2)
11	1.958(-2)	7.644(-4)	2.664(-2)	1.229(-3)	2.730(-2)	1.086(-2)
12	5.413(-3)	1.362(-2)	1.241(-2)	9.284(-3)	2.223(-2)	4.168(-3)
13	1.378(-4)	2.009(-2)	8.355(-5)	2.338(-2)	1.279(-3)	2.600(-2)
14	6.683(-3)	1.142(-2)	6.956(-3)	1.464(-2)	6.933(-3)	1.867(-2)
15	1.386(-2)	1.523(-3)	1.656(-2)	1.362(-3)	1.994(-2)	1.187(-3)
16	1.399(-2)	9.984(-4)	1.470(-2)	2.557(-3)	1.562(-2)	5.073(-3)
17	8.340(-3)	7.003(-3)	5.742(-3)	1.142(-2)	3.485(-3)	1.648(-2)
18	2.528(-3)	1.171(-2)	2.560(-4)	1.450(-2)	3.299(-4)	1.553(-2)
19	4.621(-5)	1.136(-2)	1.487(-3)	9.632(-3)	5.999(-3)	5.974(-3)
20	9.319(-4)	7.520(-3)	5.819(-3)	3.170(-3)	1.115(-2)	2.136(-4)
21	3.294(-3)	3.258(-3)	8.789(-3)	1.036(-4)	1.085(-2)	1.630(-3)
22	5.303(-3)	6.835(-4)	8.810(-3)	8.116(-4)	6.863(-3)	5.687(-3)
23	6.004(-3)	2.022(-7)	6.675(-3)	2.898(-3)	2.785(-3)	7.771(-3)
24	5.271(-3)	3.440(-4)	3.970(-3)	4.053(-3)	5.852(-4)	6.771(-3)
25	3.286(-3)	5.855(-4)	1.747(-3)	3.160(-3)	2.527(-5)	3.878(-3)
26	1.632(-3)	3.871(-4)	7.655(-4)	1.690(-3)	4.254(-7)	6.870(-4)

Table IV (cont.)

v'	v''					
	12	13	14	15	16	17
0	7.562(-2)	4.158(-2)	1.804(-2)	5.649(-3)	1.054(-3)	6.740(-5)
1	2.273(-1)	2.028(-1)	1.344(-1)	6.269(-2)	1.798(-2)	2.199(-3)
2	8.627(-2)	2.239(-1)	2.985(-1)	2.429(-1)	1.171(-1)	2.690(-2)
3	3.660(-2)	6.562(-3)	1.492(-1)	3.439(-1)	3.431(-1)	1.574(-1)
4	9.205(-2)	1.123(-1)	1.179(-2)	8.043(-2)	3.748(-1)	4.313(-1)
5	1.717(-2)	2.798(-2)	1.349(-1)	5.606(-2)	4.568(-2)	4.342(-1)
6	5.692(-2)	7.042(-2)	9.432(-5)	1.124(-1)	9.112(-2)	4.522(-2)
7	3.188(-2)	1.149(-2)	9.151(-2)	1.317(-2)	8.209(-2)	9.690(-2)
8	1.497(-2)	6.463(-2)	1.259(-3)	7.607(-2)	3.631(-2)	6.318(-2)
9	4.969(-2)	6.080(-4)	5.826(-2)	2.048(-2)	5.130(-2)	4.916(-2)
10	2.938(-3)	4.020(-2)	2.054(-2)	3.223(-2)	4.187(-2)	3.422(-2)
11	2.102(-2)	2.730(-2)	1.335(-2)	4.361(-2)	1.161(-2)	5.213(-2)
12	3.290(-2)	1.001(-3)	4.264(-2)	2.711(-4)	5.173(-2)	2.654(-3)
13	3.969(-3)	2.907(-2)	6.303(-3)	3.547(-2)	4.129(-3)	4.775(-2)
14	7.195(-3)	2.262(-2)	9.762(-3)	2.322(-2)	1.931(-2)	1.309(-2)
15	2.403(-2)	6.028(-4)	2.943(-2)	6.862(-5)	3.269(-2)	7.245(-3)
16	1.558(-2)	9.668(-3)	1.283(-2)	1.896(-2)	4.633(-3)	3.132(-2)
17	1.172(-3)	2.151(-2)	4.646(-5)	2.365(-2)	5.808(-3)	1.336(-2)
18	3.335(-3)	1.308(-2)	5.830(-3)	2.188(-2)	1.997(-4)	5.591(-5)
19	1.263(-2)	1.406(-3)	1.844(-2)	7.703(-4)	1.565(-2)	1.278(-2)
20	1.432(-2)	1.601(-3)	1.137(-2)	1.022(-2)	2.169(-3)	1.847(-2)
21	8.324(-3)	8.206(-3)	2.108(-3)	1.503(-2)	1.295(-3)	9.081(-3)
22	2.189(-3)	1.143(-2)	2.416(-4)	1.025(-2)	7.949(-3)	8.299(-4)
23	1.243(-5)	9.153(-3)	3.416(-3)	3.454(-3)	1.092(-2)	8.996(-4)
24	7.046(-4)	4.879(-3)	5.732(-3)	2.927(-4)	8.246(-3)	3.912(-3)
25	1.287(-3)	1.721(-3)	4.505(-3)	5.894(-5)	3.856(-3)	4.006(-3)
26	3.368(-4)	2.255(-4)	9.055(-4)	5.620(-5)	5.969(-4)	8.740(-4)

Table IV (cont.)

v'	v''					
	18	19	20	21	22	23
0	3.288(-7)	7.067(-7)	5.316(-8)	4.495(-10)	2.501(-9)	1.074(-10)
1	1.126(-5)	1.948(-5)	3.574(-8)	3.354(-7)	1.525(-7)	7.212(-9)
2	1.167(-3)	1.341(-4)	1.216(-5)	9.253(-6)	1.630(-6)	7.425(-8)
3	2.157(-2)	5.408(-5)	4.038(-4)	5.947(-5)	1.499(-6)	4.691(-8)
4	1.566(-1)	4.507(-3)	3.037(-3)	1.668(-5)	4.973(-5)	3.619(-6)
5	4.878(-1)	9.016(-2)	4.482(-3)	1.580(-3)	1.144(-3)	4.775(-5)
6	5.618(-1)	4.458(-1)	4.838(-3)	2.121(-2)	4.074(-3)	1.038(-4)
7	9.611(-2)	7.470(-1)	2.056(-1)	4.996(-2)	3.723(-4)	1.788(-5)
8	6.625(-2)	2.789(-1)	7.614(-1)	1.597(-3)	3.201(-2)	2.107(-3)
9	5.648(-2)	9.254(-3)	6.752(-1)	2.288(-1)	1.624(-1)	6.004(-3)
10	4.752(-2)	4.672(-2)	5.828(-2)	7.720(-1)	9.075(-2)	8.045(-4)
11	2.813(-2)	3.722(-2)	8.997(-3)	4.294(-1)	3.309(-2)	6.201(-3)
12	5.097(-2)	3.376(-2)	3.457(-2)	4.593(-2)	2.448(-1)	1.451(-2)
13	5.009(-4)	3.959(-2)	6.264(-2)	7.835(-2)	1.257(-1)	4.163(-3)
14	4.022(-2)	4.723(-4)	1.520(-2)	1.706(-1)	1.074(-1)	2.059(-3)
15	1.841(-2)	3.403(-2)	1.132(-3)	7.057(-3)	2.972(-1)	4.680(-3)
16	2.000(-3)	1.767(-2)	3.164(-2)	3.152(-4)	1.806(-1)	1.592(-4)
17	2.475(-2)	6.228(-4)	1.077(-2)	4.531(-2)	7.108(-2)	1.439(-3)
18	1.846(-2)	1.849(-2)	6.164(-4)	2.970(-4)	1.621(-1)	8.865(-4)
19	1.025(-3)	1.836(-2)	1.445(-2)	2.682(-4)	7.120(-2)	7.512(-3)
20	4.870(-3)	3.271(-3)	1.462(-2)	1.487(-2)	2.774(-2)	1.596(-2)
21	1.425(-2)	1.120(-3)	3.773(-3)	7.569(-3)	6.599(-2)	1.798(-2)
22	1.266(-2)	8.253(-3)	1.393(-4)	2.711(-3)	5.776(-3)	4.131(-2)
23	5.166(-3)	1.076(-2)	3.889(-3)	5.967(-5)	1.381(-2)	5.328(-2)
24	6.715(-4)	7.156(-3)	6.249(-3)	1.562(-3)	4.942(-3)	9.917(-2)
25	1.263(-5)	2.820(-3)	4.402(-3)	2.145(-3)	5.093(-5)	2.261(-1)
26	3.959(-5)	3.860(-4)	8.108(-4)	4.932(-4)	1.550(-4)	8.670(-1)

Table V. Spontaneous photon emission intensities $I(v', v'')$ (in units of $(e\text{\AA})^2 (\text{cm}^{-1})^3$) for the $A \ ^1\Sigma^+ \rightarrow X \ ^1\Sigma^+$ bands of ^7LiH .

v'	v''					
	0	1	2	3	4	5
0	4.745(9)	2.873(10)	9.011(10)	2.023(11)	3.584(11)	5.226(11)
1	3.007(10)	1.461(11)	3.523(11)	5.702(11)	6.651(11)	5.512(11)
2	9.796(10)	3.643(11)	6.241(11)	6.252(11)	3.388(11)	4.645(10)
3	2.203(11)	5.908(11)	6.324(11)	2.692(11)	2.802(9)	1.555(11)
4	3.853(11)	6.906(11)	3.611(11)	5.337(9)	1.899(11)	3.974(11)
5	5.627(11)	6.048(11)	7.460(10)	1.230(11)	4.012(11)	1.622(11)
6	7.085(11)	3.815(11)	8.665(9)	3.591(11)	2.394(11)	3.891(9)
7	7.874(11)	1.471(11)	1.617(11)	3.731(11)	1.635(10)	2.004(11)
8	7.890(11)	1.410(10)	3.485(11)	1.769(11)	6.524(10)	3.092(11)
9	7.246(11)	1.872(10)	4.066(11)	1.474(10)	2.538(11)	1.456(11)
10	6.162(11)	1.251(11)	3.127(11)	3.405(10)	3.026(11)	2.740(9)
11	4.905(11)	2.650(11)	1.545(11)	1.749(11)	1.727(11)	7.211(10)
12	3.698(11)	3.803(11)	3.405(10)	2.927(11)	3.053(10)	2.158(11)
13	2.664(11)	4.420(11)	7.750(8)	3.038(11)	7.874(9)	2.456(11)
14	1.842(11)	4.469(11)	4.676(10)	2.207(11)	9.534(10)	1.473(11)
15	1.230(11)	4.085(11)	1.324(11)	1.087(11)	2.002(11)	3.501(10)
16	8.051(10)	3.492(11)	2.187(11)	2.723(10)	2.488(11)	9.467(8)
17	5.131(10)	2.806(11)	2.781(11)	6.944(5)	2.224(11)	4.889(10)
18	3.237(10)	2.169(11)	3.038(11)	2.053(10)	1.521(11)	1.258(11)
19	2.020(10)	1.620(11)	2.986(11)	6.658(10)	7.681(10)	1.788(11)
20	1.271(10)	1.191(11)	2.742(11)	1.163(11)	2.437(10)	1.880(11)
21	7.945(9)	8.532(10)	2.360(11)	1.533(11)	1.695(9)	1.586(11)
22	5.005(9)	6.034(10)	1.944(11)	1.718(11)	3.113(9)	1.124(11)
23	3.134(9)	4.157(10)	1.516(11)	1.688(11)	1.634(10)	6.698(10)
24	1.894(9)	2.702(10)	1.087(11)	1.436(11)	2.807(10)	3.298(10)
25	9.573(8)	1.434(10)	6.170(10)	9.128(10)	2.560(10)	1.243(10)
26	4.470(8)	6.798(9)	2.990(10)	4.591(10)	1.435(10)	5.088(9)

Table V (cont.)

v'	v''					
	6	7	8	9	10	11
0	6.583(11)	7.225(11)	6.913(11)	5.769(11)	4.203(11)	2.653(11)
1	3.001(11)	6.864(10)	3.904(9)	1.308(11)	3.363(11)	4.702(11)
2	3.598(10)	2.556(11)	4.007(11)	2.878(11)	6.592(10)	1.122(10)
3	3.876(11)	2.890(11)	3.816(10)	5.133(10)	2.702(11)	2.963(11)
4	1.776(11)	1.456(9)	2.001(11)	3.006(11)	8.407(10)	2.275(10)
5	1.369(10)	2.616(11)	2.340(11)	5.810(9)	1.390(11)	2.545(11)
6	2.609(11)	2.114(11)	1.428(8)	1.997(11)	1.945(11)	2.526(8)
7	2.499(11)	3.909(8)	2.031(11)	1.712(11)	4.429(9)	2.045(11)
8	2.625(10)	1.574(11)	1.978(11)	1.851(9)	2.050(11)	8.345(10)
9	5.916(10)	2.491(11)	5.860(9)	1.774(11)	1.089(11)	4.062(10)
10	2.334(11)	7.606(10)	1.008(11)	1.707(11)	9.297(9)	1.913(11)
11	2.194(11)	7.345(9)	2.191(11)	8.638(9)	1.636(11)	5.544(10)
12	6.386(10)	1.382(11)	1.081(11)	6.929(10)	1.420(11)	2.274(10)
13	1.711(9)	2.150(11)	7.695(8)	1.851(11)	8.689(9)	1.514(11)
14	8.711(10)	1.287(11)	6.764(10)	1.227(11)	5.000(10)	1.158(11)
15	1.895(11)	1.804(10)	1.697(11)	1.206(10)	1.524(11)	7.824(9)
16	2.003(11)	1.241(10)	1.585(11)	2.388(10)	1.263(11)	3.546(10)
17	1.248(11)	9.120(10)	6.500(10)	1.122(11)	2.971(10)	1.218(11)
18	3.947(10)	1.594(11)	3.035(9)	1.496(11)	2.960(9)	1.212(11)
19	7.514(8)	1.614(11)	1.843(10)	1.041(11)	5.652(10)	4.904(10)
20	1.574(10)	1.112(11)	7.520(10)	3.579(10)	1.099(11)	1.840(9)
21	5.765(10)	4.998(10)	1.181(11)	1.218(9)	1.116(11)	1.467(10)
22	9.588(10)	1.085(10)	1.227(11)	9.906(9)	7.341(10)	5.332(10)
23	1.118(11)	3.310(6)	9.594(10)	3.657(10)	3.085(10)	7.555(10)
24	1.006(11)	5.777(9)	5.862(10)	5.261(10)	6.675(9)	6.788(10)
25	6.385(10)	1.002(10)	2.631(10)	4.185(10)	2.944(8)	3.976(10)
26	3.194(10)	6.674(9)	1.162(10)	2.257(10)	5.006(6)	7.116(9)

Table V (cont.)

v'	v''					
	12	13	14	15	16	17
0	1.425(11)	6.311(10)	2.199(10)	5.512(9)	8.230(8)	4.219(7)
1	4.582(11)	3.312(11)	1.771(11)	6.654(10)	1.537(10)	1.517(9)
2	1.870(11)	3.952(11)	4.276(11)	2.820(11)	1.102(11)	2.057(10)
3	8.567(10)	1.257(10)	2.334(11)	4.386(11)	3.571(11)	1.341(11)
4	2.331(11)	2.340(11)	2.017(10)	1.129(11)	4.323(11)	4.100(11)
5	4.706(10)	6.342(10)	2.526(11)	8.666(10)	5.837(10)	4.605(11)
6	1.689(11)	1.737(11)	1.932(8)	1.913(11)	1.289(11)	5.345(10)
7	1.024(11)	3.082(10)	2.049(11)	2.463(10)	1.284(11)	1.274(11)
8	5.196(10)	1.883(11)	3.078(9)	1.561(11)	6.268(10)	9.219(10)
9	1.861(11)	1.920(9)	1.551(11)	4.601(10)	9.749(10)	7.940(10)
10	1.186(10)	1.374(11)	5.948(10)	7.911(10)	8.737(10)	6.099(10)
11	9.124(10)	1.008(11)	4.191(10)	1.166(11)	2.653(10)	1.022(11)
12	1.533(11)	3.980(9)	1.448(11)	7.878(8)	1.290(11)	5.703(9)
13	1.980(10)	1.243(11)	2.310(10)	1.116(11)	1.120(10)	1.121(11)
14	3.837(10)	1.037(11)	3.851(10)	7.896(10)	5.678(10)	3.346(10)
15	1.366(11)	2.956(9)	1.246(11)	2.513(8)	1.039(11)	2.009(10)
16	9.415(10)	5.057(10)	5.812(10)	7.457(10)	1.587(10)	9.390(10)
17	7.519(9)	1.197(11)	2.246(8)	9.956(10)	2.136(10)	4.316(10)
18	2.262(10)	7.722(10)	5.560(10)	2.618(10)	8.609(10)	6.920(8)
19	9.036(10)	8.776(9)	1.006(11)	3.677(9)	6.564(10)	4.733(10)
20	1.077(11)	1.053(10)	6.548(10)	5.163(10)	9.656(9)	7.281(10)
21	6.559(10)	5.665(10)	1.277(10)	8.008(10)	6.094(9)	3.791(10)
22	1.800(10)	8.250(10)	1.533(9)	5.734(10)	3.933(10)	3.652(9)
23	1.061(8)	6.873(10)	2.260(10)	2.017(10)	5.654(10)	4.147(9)
24	6.213(9)	3.781(10)	3.925(10)	1.773(9)	4.432(10)	1.876(10)
25	1.161(10)	1.369(10)	3.164(10)	3.665(8)	2.130(10)	1.976(10)
26	3.072(9)	1.814(9)	6.435(9)	3.547(8)	3.339(9)	4.369(9)

Table V (cont.)

v'	v''					
	18	19	20	21	22	23
0	1.660(5)	2.912(5)	1.827(4)	1.326(2)	6.698(2)	2.759(1)
1	6.308(6)	8.976(6)	1.383(4)	1.126(5)	4.644(4)	2.111(3)
2	7.306(8)	6.954(7)	5.330(6)	3.539(6)	5.680(5)	2.492(4)
3	1.515(10)	3.169(7)	2.014(8)	2.604(7)	6.007(5)	1.813(4)
4	1.236(11)	2.989(9)	1.726(9)	8.367(6)	2.292(7)	1.618(6)
5	4.324(11)	6.762(10)	2.899(9)	9.072(8)	6.058(8)	2.456(7)
6	5.587(11)	3.775(11)	3.554(9)	1.390(10)	2.471(9)	6.125(7)
7	1.070(11)	7.124(11)	1.710(11)	3.726(10)	2.578(8)	1.205(7)
8	8.233(10)	2.987(11)	7.151(11)	1.350(9)	2.521(10)	1.617(9)
9	7.811(10)	1.109(10)	7.130(11)	2.184(11)	1.448(11)	5.224(9)
10	7.289(10)	6.240(10)	6.892(10)	8.283(11)	9.120(10)	7.897(8)
11	4.770(10)	5.519(10)	1.186(10)	5.156(11)	3.730(10)	6.834(9)
12	9.516(10)	5.537(10)	5.062(10)	6.143(10)	3.080(11)	1.787(10)
13	1.026(9)	7.152(10)	1.014(11)	1.162(11)	1.758(11)	5.701(9)
14	9.004(10)	9.361(8)	2.708(10)	2.794(11)	1.661(11)	3.121(9)
15	4.489(10)	7.370(10)	2.210(9)	1.270(10)	5.062(11)	7.814(9)
16	5.289(9)	4.163(10)	6.746(10)	6.204(8)	3.370(11)	2.914(8)
17	7.073(10)	1.591(9)	2.495(10)	9.715(10)	1.447(11)	2.876(9)
18	5.677(10)	5.097(10)	1.545(9)	6.904(8)	3.582(11)	1.924(9)
19	3.378(9)	5.438(10)	3.900(10)	6.767(8)	1.700(11)	1.762(10)
20	1.712(10)	1.036(10)	4.228(10)	4.003(10)	7.114(10)	4.025(10)
21	5.318(10)	3.771(9)	1.163(10)	2.174(10)	1.809(11)	4.845(10)
22	4.990(10)	2.941(10)	4.549(8)	8.263(9)	1.682(10)	1.183(11)
23	2.136(10)	4.031(10)	1.337(10)	1.917(8)	4.242(10)	1.610(11)
24	2.893(9)	2.796(10)	2.244(10)	5.246(9)	1.588(10)	3.135(11)
25	5.604(7)	1.136(10)	1.631(10)	7.437(9)	1.690(8)	7.385(11)
26	1.780(8)	1.577(9)	3.048(9)	1.736(9)	5.220(8)	2.875(12)

Figure 3. Matrix of intensities for $A^1\Sigma^+ (v' = 0 - 26) \rightarrow X^1\Sigma^+ (v'' = 0 - 23)$ emission (photon detection) of ^7LiH . As also shown in Fig. 1, strong transitions connect $v'' = 0$ to $v' = 2-15$ and $v' = 0$ to $v'' = 3 - 12$. Intense transitions corresponding to $v' = 5$ to $v'' = 18$ and its immediate vicinity in frequency also occur (a "head of heads" or classical "satellite"¹⁶). The most intense transition is the long-range molecular one: $v' = 26$ to $v'' = 23$ (the wavefunctions overlap 75%¹⁵). The observed bands (primarily from currently unpublished and still incomplete work in this laboratory by K. C. Li and F. B. Orth (see discussion below), using plates measured from ~ 3400 - $8700 \text{ \AA}$) are also indicated. The heavy lines ending on the large number (e.g. 3500) delimit the spectroscopic wavelength region in $\text{\AA}$ in which ν_{00} for the various bands occurs. Note that a rich spectrum extending from ~ 3000 - $10000 \text{ \AA}$ is predicted, consistent with observations.

B-677-258-1

■ VERY STRONG ($I \geq 10^{12}$)

▨ WEAK ($I \geq 10^{10}$)

▩ STRONG ($I \geq 10^{11}$)

□ VERY WEAK ($I \geq 10^9$)

• OBSERVED

In summary (except for transitions involving $v'' = 23$ or $v' = 26$) we are confident of the numerical reliability of our q values to $< 10^{-4}$, $|D|^2$ values to $< 10^{-4} \text{ ea}_0$, and our I values to $\sim 10^8 (\text{eA})^2 (\text{cm}^{-1})^3$. Further discussion of the highest levels which present particular numerical difficulties will be provided elsewhere.⁹

V. DISCUSSION

The matrix of spontaneous photon emission intensities $I(v', v'')$ (Figure 3) is worth examining in greater detail in terms of the potential energy curves involved (figure 1) (matrices of $q_{v', v''}$ or $|D_{v', v''}|^2$ could equally well be given). As shown in figure 1, emission for $v' = 0$ should occur to a variety of lower levels; a rough criterion for a reasonably intense transition is that the outer turning points of the lower levels ($R_{v'',+}$) be between the inner and outer turning points of the $v' = 0$ level ($R_{0,+}$) (see Figure 1). The levels v' satisfying this relationship ($R_{0,-} \leq R_{v'',+} \leq R_{0,+}$) are termed "allowed" in Figure 4a. The emission from $v' = 1$ is similar (Figure 4b) except for a central dip in intensity near $v'' = 8$. The slightly expanded "allowed" region corresponds here to $R_{1,-} \leq R_{v'',+} \leq R_{1,+}$. The central dip is associated with the node in the $v'=1$ wavefunction being at a distance close to $R_{8,+}$. The emission from $v' = 2$ is similar (Figure 4c) but with an additional slight expansion of the "allowed" region ($R_{2,-} \leq R_{v'',+} \leq R_{2,+}$) and with two wavefunction nodes at distances close to $R_{6,+}$ and $R_{11,+}$, respectively. Note that except for the nodal dips, the intensities of all "allowed" transitions are comparable ($I \approx 1.5 \times 10^{11} \text{ (eA)}^2 \text{ (cm}^{-1})^3$ or $A \approx 1 \times 10^5 \text{ sec}^{-1}$). The large weak (mostly blank) triangle in upper right corner of Figure 3 represents emission which is "forbidden" because $R_{v'',+} > R_{v',+}$ for low v' . The much smaller weak triangle in the upper left corner represents emission which is "forbidden" because $R_{v'',+} < R_{v',-}$ for low v' . For higher v' the emission again becomes "allowed" because of predominantly inner turning point transitions, and becomes quite complex because in general neither wavefunction has simple nodal structure. For the highest v' of 26 (a "long-range molecule"), however, a new simplification occurs (discussed further elsewhere⁹) as most of the wavefunction amplitude is outside the

Figure 4. Absolute spontaneous photon emission intensities $I(v',v'')$ (in $(\text{e}\text{\AA})^2 (\text{cm}^{-1})^3$) for the ${}^7\text{LiH } A \ ^1\Sigma^+ (v') \rightarrow X \ ^1\Sigma^+ (v'')$ bands for (a) $v' = 0$; (b) $v' = 1$; and (c) $v' = 2$, as a function of v'' . Note the smooth behavior of these curves, with strong intensities only in the "allowed" regions (where $R_{v',-} \leq R_{v'',+} \leq R_{v',+}$). Note also the nodal dips (one in $v' = 1$ and two in $v' = 2$) where the internuclear distance of the $\psi_{v'}$ wavefunction approximately equals $R_{v'',+}$.

outermost node and hence transitions to similar levels (e.g. $v'' = 23$) become the strongest molecular transitions in the entire band system (almost atomic in strength).

The satellite band (or head-of-heads) behavior¹⁶ which occurs when potential curves are parallel (at $\sim 7.3a_0$ in Figure 1) is also apparent in Figure 3. A large number of strong bands (and also bound to free emission) all occur in the same spectral region near the 5-18 band at $\sim 10400\text{\AA}$.

The experimental observations of emission²⁰⁻²² do not yet include precise measurement of even relative intensities. Nevertheless, it is clear from Figure 3 that the strong bands expected are indeed often observed. The inability to observe the 0-0 band and adjacent bands in the upper left hand corner, for example, is consistent with the present calculations. There does appear to be a discrepancy in the position of the nodal dip for $v' = 1$ emission bands in that the $v' = 8$ transition is predicted to be weakest, whereas the 1-8 band is in fact observed while the 1-9 and 1-10 bands seem too weak to observe²². Further study of the relative emission intensities for low v' has begun.

Similar results are obtained in the case of absorption. A matrix of $|D|^2$ or $|D|^2_v$ values would differ from the I matrix of Figure 3 only in detail and not in overall structure. For example, the (energy) absorption coefficient from $v'' = 0$ given in Table VI and shown in Figure 5 has a roughly gaussian shape similar to $v' = 0$ emission in Figure 4a. The "allowed" transitions in this case would be absorption into v' levels where $R_{0''-} \leq R_{v'-} \leq R_{0''+}$, i.e. $v' = 4 - 13$, which are clearly the strongest. Again $v' = 1$ and 2 absorption show nodal dips (at $v' \sim 8$, and at $v' \sim 6$ and 13, respectively). The extreme lower left corner of an absorption coefficient matrix similar to Figure 3, for example, shows a weak region resulting from $R_{v''-} > R_{v'-}$.

Table VI. Relative (energy) absorption coefficients for the ${}^7\text{LiH}$ $A\ ^1\Sigma^+ (v') \leftarrow X\ ^1\Sigma^+ (v'' = 0)$ bands are given normalized to a value of 47.5 at $v'=8$. The accurate $v|D|^2$ calculation ($\times$) is similar in shape to the simpler Franck-Condon approximation (vq ; $-\rightarrow-$), but significantly different in detail (e.g. a maximum at $v' = 7$ rather than 8). Note that this is for a $v'' = 0$ which only vibrates (classically) $\sim 0.33\text{\AA}$, and for which the change of $D(R)$ might easily have been neglected. Also given are experimental values²³, which presumably are uncertain by roughly their deviation ($\sim \pm 30\%$) from a smooth Gaussian shape which must result in any case.

v'	Absorption coefficient (arbitrary units)		
	Expt. ^a	$v_{v',0} D_{v',0} ^2{}^b$	$v_{v',0}q_{v',0}{}^b$
0	----	0.4	0.3
1	----	2.2	1.6
2	3.8	6.9	5.4
3	10.2	15.2	12.3
4	16.1	25.8	21.7
5	18.7	36.7	32.1
6	24.0	45.0	41.1
7	42.1	48.7	46.5
8	47.5	47.5 ^b	47.5 ^b
9	32.5	42.5	44.5
10	34.3	35.2	38.7
11	39.7	27.3	31.5
12	41.8	20.0	24.4
13	32.6	14.1	18.1
14	----	9.5	12.9
15	----	6.2	8.8
16	----	4.0	6.0
17	----	2.5	4.0
18	----	1.5	2.6

^aEstimated from Figure 8 of Ref. 23.

^bNormalized to 47.5 at $v' = 8$.

Figure 5. Relative (energy) absorption coefficients from Table VI for the ${}^7\text{LiH A } {}^1\Sigma^+ (v') + \chi {}^1\Sigma^+ (v'' = 0)$ bands are given normalized to 47.5 at $v' = 8$.

Again absorption from higher v'' becomes more complex because of the interference of the highly oscillatory wavefunction, simplifying again only for the long-range molecule⁹ $v'' = 23$, whose absorption spectrum is dominated by $v' = 26$.

In summary, we have used hybrid potentials² and dipole strength function to accurately calculate absolute radiative transition probabilities for absorption and emission involving all vibrational levels of the $A^1\Sigma^+$ and $X^1\Sigma^+$ electronic states of ^7LiH and found these calculations to be in general in reasonable accord with our expectations (in terms of the Franck-Condon Principle) and with experimental observations.

REFERENCES

1. W. C. Stwalley and W. T. Zemke, "Selected Bibliography on Ground and Excited Potential Curves of H_2 , LiH and Li_2 ", AEC Report C00-2326-6 (1974). This and other C00-2326-xx reports are available on request from WCS.
2. W. C. Stwalley, W. T. Zemke, K. R. Way, K. C. Li and T. R. Proctor, J. Chem. Phys. 66, 5412 (1977); erratum 67, xxxx (1977).
3. W. T. Zemke and W. C. Stwalley, "Tabulated Radiative Transition Probabilities for the $A\ ^1\Sigma^+ - X\ ^1\Sigma^+$, $B\ ^1\Pi - X\ ^1\Sigma^+$, and $B\ ^1\Pi - A\ ^1\Sigma^+$ Bands of 7LiH ", ERDA Report C00-2326-24(1977).
4. W. T. Zemke, J. B. Crooks and W. C. Stwalley, "Radiative and Nonradiative Lifetimes for Vibrational Levels of the $A\ ^1\Sigma^+$ state of 7LiH ", ERDA Report C00-2326-23(1977); following paper.
5. W. T. Zemke, W. C. Stwalley, and K. R. Way, "Radiative Transition Probabilities for the $B\ ^1\Pi - X\ ^1\Sigma^+$ and $B\ ^1\Pi - A\ ^1\Sigma^+$ Bands of 7LiH ", ERDA Report C00-2326-29(1977); to be submitted for publication.
6. W. T. Zemke and W. C. Stwalley, "Radiative Lifetimes for Vibrational Levels of the $B\ ^1\Pi$ State of 7LiH ", ERDA Report C00-2326-32(1977); to be submitted for publication.
7. W. T. Zemke and W. C. Stwalley, "Radiative Transition Probabilities and Lifetimes for Vibrational Levels of Ground State 7LiH ", to be submitted for publication.
8. W. C. Stwalley, Contemp. Phys. 18, xxxx (1977).
9. W. T. Zemke and W. C. Stwalley "Radiative Transition Probabilities Involving Long-Range Molecular Levels of 7LiH ", to be submitted for publication.
10. W. L. Wiese, M. W. Smith, and B. M. Glennon, Atomic Transition Probabilities, Volume I Hydrogen Through Neon (National Bureau of Standards, Washington, D. C. 1966).
11. R. S. Mulliken, Intl. J. Quantum Chem. 1, 103(1967);
12. R. S. Mulliken, Intl. J. Quantum Chem. 8, 817(1974);
13. W. C. Ermler, R. S. Mulliken and A. C. Wahl, J. Chem. Phys. 66, 3031 (1977).
14. K. K. Docken and J. Hinze, J. Chem. Phys. 57, 4936 (1972).
15. W. C. Stwalley and W. T. Zemke, Intl. J. Quantum Chem. S10, 223 (1976).
16. L. K. Lam, A. Gallagher and M. M. Hessel, J. Chem. Phys. 66, 3550 (1977).
17. R. N. Zare, J. Chem. Phys. 40, 1934 (1964).
18. M. Halmann and I. Lauicht, J. Chem. Phys. 46, 2684 (1967); also supplementary material discussed therein.

19. M. Oppenheimer and K. K. Docken, Chem. Phys. Letters 29, 349(1974); signs of matrix elements from K. K. Docken, private communication.
20. F. H. Crawford and T. Jorgensen, Phys. Rev. 47, 358 and 932 (1935); T. Jorgensen, Ph.D. Thesis, Harvard Univ. (1935).
21. K. C. Li and W. C. Stwalley, "The A $1\Sigma^+ \rightarrow X 1\Sigma^+$ Bands of the Isotopic Lithium Hydrides", ERDA Report C00-2326-28(1977), submitted to J. Mol. Spectrosc.; K. C. Li and W. C. Stwalley, "Assigned Lines in the A $1\Sigma^+ - X 1\Sigma^+$ Bands of ^6LiH , ^6LiD , ^7LiH and ^7LiD ", ERDA Report C00-2326-20(1977).
22. F. B. Orth and W. C. Stwalley, unpublished.
23. I. Fernandez-Florez and R. Velasco, Optica Puray Aplicada 2, 153(1969).