AN EVALUATION OF THE ECOLOGICAL CONSEQUENCES OF PARTIAL POWER OPERATION OF K REACTORS, SRS (U)

by

J. B. Gladden, H. E. Mackey, M. M. Paller, W. L. Specht, L. D. Wike, and E. W. Wilde

Westinghouse Savannah River Company
Savannah River Site
Aiken, South Carolina 29808

DISCLAIMER

A technical being deposited at OSTI

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

This report was prepared in connection with work done under Contract No. DE-AC09-89SR18035 with the U.S. Department of Energy. By acceptance of this report, the publisher and/or recipient acknowledges the U.S. Government's right to retain a nonexclusive, royalty-free license in and to any copyright covering this report, along with the right to reproduce and to authorize others to reproduce all or part of the copyrighted report.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

D. B. Moore, Manager

WSRC-RP-91-688

D. B. Moore, Manager
Authorized Derivative Classifier

**An Evaluation of the Ecological Consequences of
Partial-Power Operation of the K Reactor, SRS**

By: J. B. Gladden
H. E. Mackey
M. H. Paller
W. L. Specht
L. D. Wike
E. W. Wilde

Environmental Sciences Section
Savannah River Laboratory

Approved by: D. B. Moore, Section Manager
D. B. Moore, Section Manager
Environmental Sciences Section

Publication Date: JUNE, 1991

WESTINGHOUSE SAVANNAH RIVER COMPANY
SAVANNAH RIVER SITE
AIKEN, SC 29808

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by the trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Summary

The K Reactor at the Savannah River Site (SRS) shut-down in spring 1988 for maintenance and safety upgrades. Since that time the receiving stream for thermal effluent, Indian Grave Branch and Pen Branch, have undergone a pattern of post-thermal recovery that is typical of other SRS streams following removal of thermal stress. Diversity of fish and aquatic macroinvertebrate communities has increased and available habitats have been colonized by numerous species of herbaceous and woody plants.

K Reactor is scheduled to resume operation in 1991 and operate through 1992 without a cooling tower to cool the discharge. It is likely that the reactor will operate at approximately one-third to one-half of full power (800-1200 MW thermal) during this period and effluent temperatures will be substantially lower than earlier operation at full power. Monthly average discharge temperatures at half-power operation will range from approximately 42°C in winter to 49°C in summer. The volume of water discharged will not be affected by altered power levels and will average approximately 10-11 m³/s. The ecological consequences of this mode of operation on the Indian Grave/Pen Branch stream system have been evaluated.

Thermal modeling studies predicted that with half-power operation under average monthly meteorological conditions, temperatures exceeding 35°C will occur throughout the year during reactor operation from the reactor outfall through the upper and mid-reaches of the receiving stream. Under summer meteorological conditions, water temperatures exceeding 40°C will occur as far downstream as the Pen Branch delta. When the reactor operates at approximately one-third power, effluent discharge temperatures will range from approximately 31°C in winter to 42°C in summer. Summer stream temperatures will remain as high as 38°C at the lower end of the stream where it enters the Pen Branch delta.

The combination of high temperatures and high stream discharge will preclude the continued development and maintenance of normal stream biotic communities from the K-Reactor discharge to the Pen Branch delta. Much of the aquatic and wetland biotic communities that have become established during the current reactor outage will be lost from most of this stream reach because of high water temperatures, physical displacement of aquatic organisms and erosional undercutting of vegetation.

Phytoplankton, periphyton and zooplankton are not major contributors to the structure or function of small stream ecosystems, so adverse effects on these communities in the

upper reaches of the stream will be of minimal consequence. Water temperatures will exceed the thermal tolerances of most fish and invertebrates throughout most of the year in the upper reaches of Indian Grave/Pen Branch, and during the summer in the lower reaches resulting in exclusion of most macroinvertebrate and fish species from the upper and middle reaches of the stream. Those portions of the stream that do not exceed 33°C should maintain diverse and productive macroinvertebrate communities, while temperatures of 33-35°C will result in reductions in species richness. Similarly, stream reaches that do not exceed 33°C, decrease to at least 20°C for spring spawning, and do not undergo rapid temperature drops of more than 15°C during winter shut-down will provide acceptable thermal regimes for fish. Summer temperatures will exceed the thermal tolerance of most stream biota for both 30% and 50% power operation.

In the Pen Branch delta, water velocities are reduced and sediment eroded from upstream will be deposited. Water levels in the delta will be elevated, which may eliminate some of the less flood-tolerant plant species that have recolonized the delta during the reactor outage. However, water temperatures in the Pen Branch delta will vary seasonally with winter temperatures being low enough to permit occupancy by many species of fish and invertebrates. Summer temperatures will likely preclude the presence of most aquatic organisms in much of the delta. Areas on the outer periphery of the delta where warm reactor effluent water mixes with ambient temperature swamp water will likely be inhabited by common swamp organisms throughout the year. Above ambient water temperatures and increased nutrients from the Savannah River water may enhance production of some aquatic species in regions where temperatures remain below lethal limits.

High summer temperatures may result in the resumption of tree kill on the edge of the delta, but the rate of tree kill, if it occurs, should be substantially lower than observed during full power operations. The emergent and submergent herbaceous wetland vegetation community is expected to revert to a community dominated by thermally tolerant species because the growing season for these species is concurrent with periods when water temperatures are maximum.

The Pen Branch delta has historically provided marginal foraging habitat for the endangered wood stork. The elevated water levels and probable thermal exclusion of fish from this area during the summer months will likely preclude the utilization of the area by wood storks. However, the Department of Energy funded construction of Kathwood Lake in 1985 to provide supplemental foraging habitat for wood storks as mitigation for lost foraging habitat associated with L Reactor operation.

The American alligator will be excluded from most thermally affected areas of the Pen Branch corridor, but populations on or near the Pen Branch delta will not likely be affected adversely and alligators may be attracted to the area in winter.

At the time of the expected K-Reactor restart, the Indian Grave/Pen Branch stream ecosystem will have undergone four years of post-thermal recovery. Adverse impacts to the stream and delta ecosystems are expected to be similar to those observed during earlier operations, with the exception that the impacts should be somewhat reduced in magnitude because of the lower discharge temperatures. Consequently, the biota of the affected stream and delta areas should recover to their current status within a period of three-to-four years following mitigation of thermal effluents with the cooling tower. The exception to this predicted recovery pattern arises from the possibility that additional mature cypress and tupelo trees would be killed in the Pen Branch swamp. Should additional mature trees be killed, recovery time for this ecosystem component would be measured in decades.

Introduction

The K Reactor at the Savannah River Site (SRS) discharges water into Indian Grave Branch, a tributary of Pen Branch (Figure 1). During previous K-Reactor operation at full power, cooling water effluent temperatures to Indian Grave Branch were $\geq 70^{\circ}\text{C}$, which resulted in substantial degradation of biological communities in affected areas of Indian Grave Branch, Pen Branch and portions of the Savannah River swamp. Upstream communities were largely restricted to thermophilic blue-green algae, while communities in downstream areas contained thermo-tolerant algae and a low diversity of macroinvertebrates (Specht, 1987). Small numbers of thermally tolerant fish occupied cooler coves and backwaters adjacent to the creek and thermally tolerant plants occupied stumps and sandbar/mudflat areas that were above the water level. From 1954 to 1985, thermal impacts resulted in the loss of wetland tree canopy from approximately 93 hectares (ha) in the stream corridor upstream from the swamp and approximately 152 ha in the swamp; the rate of swamp canopy loss during full power operation was 5-10 ha per year (Mackey, 1987). High discharges during earlier operation resulted in substantial erosion and channel enlargement in upstream areas and sediment deposition in the downstream swamp areas producing a sedimentary delta where Pen Branch enters the Savannah River floodplain swamp.

Following the cessation of thermal discharges in 1988, biota began to recolonize the effluent streams. Previous research has shown that within days of cessation fish begin to reinvade the lower portions of Pen Branch (Specht, 1987). Some species of aquatic macroinvertebrates and wetland/aquatic vegetation also rapidly became established in the habitats when ambient temperatures were reestablished, although the rates of reinvasion were likely not as rapid as for fish (Wike et al., 1989).

K Reactor is scheduled to resume operation in 1991, prior to the completion of the K-Reactor cooling tower. From start-up in 1991 through much of 1992, K Reactor is scheduled to operate without the cooling tower. During this period the reactor will operate at a maximum of approximately one-third to one-half power [800-1200 megawatts (MW) thermal]. The volume of cooling water discharge will be approximately 10-11 m^3/s during reactor operation because the volume of cooling water discharged is independent of reactor power levels. Cooling water effluent temperatures will vary depending on reactor power levels and intake water temperatures. The relationship between cooling water intake temperature and effluent temperature is depicted in Figure 2. Although K-Reactor power levels are predicted to decline somewhat in summer, cooling water effluent temperatures are higher than

in winter because of the higher temperature of Savannah River water that is used to cool the reactor. With half-power operation average monthly discharge temperatures will range from approximately 40°C in winter to 51°C in summer. At one-third power, cooling water effluent temperatures will range from approximately 31°C in winter to approximately 42°C in summer.

The reduction in effluent temperature coupled with the instream cooling will result in temperature decreases as water flows down Pen Branch. Upon exiting the reactor, water cooling in the open channels of Indian Grave Branch and then Pen Branch is strongly influenced by seasonal meteorological conditions such as air temperature, relative humidity and wind conditions. Lower air temperatures result in greater heat transfer from the water to the air, and therefore the rate of water temperature decline from the reactor outfall to the swamp is greater in winter than in summer.

Stream Temperatures

During the early phase of K-Reactor operation, discharge temperatures were comparable to those predicted for half-power operation under current conditions. Representative values for temperatures in Pen Branch are presented in Table 1. Although the data are not complete for all stations, and measurements are not available from the swamp, the results indicate that in the range of predicted discharge temperatures for half-power operation, temperatures higher than 35-37°C occurred in the lower reaches of Pen Branch. Several potentially important factors differ between operations in 1955 and projected operations for 1991-92. Discharge volumes were substantially lower in 1955 than now, which would affect the water transport rates, cross-sectional area, and surface area of the stream. Additionally, little tree canopy loss had occurred in the stream and swamp in 1955, which would reduce both air circulation near the heated water surface and solar heating of the water. The relative effects of these factors on overall cooling performance of the stream is not known quantitatively.

Pen Branch water temperatures were also simulated for current stream conditions using predicted monthly discharge temperatures and average monthly meteorological conditions (Table 2). These simulations indicate that summer temperatures would exceed 35°C as far downstream as the entrance to the swamp for both the 30% and 50% reactor power scenarios. Predicted winter temperatures are substantially lower, but nearly reach 30°C at Hwy. 125 and 25°C at the swamp for 30% and exceed those values for 50% K-Reactor power operation.

Phytoplankton

Phytoplankton are a minor component of small stream ecosystems and contribute little to the food base of the Indian Grave Branch/Pen Branch stream community. Consequently, impacts on phytoplankton caused by the restart of K Reactor without a cooling tower will be of little environmental significance and would be reversible once the cooling tower is on line.

Periphyton

The periphyton community of the Pen Branch system was studied between 1983 and 1985 when K Reactor was operating at full power (Specht 1987). The stream periphyton community was dominated by thermally tolerant blue-green algae in the thermally impacted reaches of Indian Grave Branch and Pen Branch. However, periphyton are not considered a dominant contributor to the food chain base in this lotic stream system. Restarting the reactor without a cooling tower would be expected to change the species composition of periphyton in the discharge area toward a community dominated by thermophilic blue-green algae, as was the case when the reactor was operating prior to 1988. However, this change would not be ecologically significant and the communities would revert to normal once the cooling tower becomes operational.

Zooplankton

Zooplankton are generally considered to be a minor component of small stream ecosystems and contribute little to the food base of higher organisms in such ecosystems. Although the zooplankton community in the Pen Branch system has not been studied since August 1985, prior to the April 1988 outage, the greatly reduced flows and ambient temperature water in the stream between the reactor outfall and the delta has provided an environment more suitable for zooplankton development than was the case when the reactor was operating. A restart of the reactor without a cooling tower would disrupt this community. The increased flows and elevated temperatures would make the corridor region mostly uninhabitable for zooplankton when the reactor is operating. Conversely, restarting the reactor without a cooling tower would result in increased aquatic habitat in the Pen Branch delta and swamp, and increased nutrient loading from the discharge of Savannah River water. With operation at half-power or less, the increased temperatures may be below the upper thermal tolerance limits for most zooplankton in much of the Pen Branch delta and adjacent swamp. In these downstream areas zooplankton abundance could be enhanced.

Wetlands

Swamp forests associated with the Savannah River floodplain on the SRS are dominated by bald cypress (Taxodium distichum) and water tupelo (Nyssa aquatica). Thermal discharges from the nuclear reactors and a coal-fired power plant are or have been released into tributaries which flow into this riverine swamp. Thermal discharges have increased water volume, flow rates, and water temperatures, subjecting the original swamp forest and replacement communities to substrate instability, increased flooding, and thermal stress (Donovan and McLeod, 1985). By 1985, approximately 93 hectares (ha) of bottomland forest had been lost in the stream corridor from K-Reactor discharge canal to the swamp and 152 ha of forest had been lost in the swamp (Figure 3). During full power operations of K Reactor, approximately 5-10 ha of swamp forest canopy was lost annually during full power operation.

The combinations of increased flow, temperature, and sediment erosion/deposition has resulted in the formation of zones of wetland vegetation which reflect past and continuing perturbations from thermal discharges. Table 3 summarizes the typical dominant wetland species and/or wetland community types found under thermal conditions on or near the deltas of the SRS Savannah River swamp when the reactors operated at nominal full power and with discharge rates of 10-11 m³/s (Christy 1976; Christy and Sharitz 1980; Jensen et al. 1983; Jensen et al. 1986; Jensen et al. 1987;). In addition to Ludwigia spp. and Ammannia coccinea, Christy and Sharitz (1980) also list Boehmeria cylindrica and several graminoids as common on the Pen Branch delta.

Under thermal conditions the herbaceous wetland communities of the thermal deltas of the SRS are dominated primarily by a few thermally tolerant species. Irwin (1975) reported Eupatorium spp., Ludwigia leptocarpa, Fimbristylis spadicea, Paspalum dilatatum, Lycopus sp., and Panicum anceps as common on stump and log communities of Pen Branch at thermal sites (50°C, 45°C, and 40°C). Some old field species such as Lespedeza cuneata and Andropogon virginicus occurred on stumps and sand/mud islands but are not typically regarded as wetland species. Christy and Sharitz (1980), Sharitz et al. (1974) McCaffrey (1982), Christy (1976), Jensen et al (1987) and Mackey (1990) also reported Cyperus strigosus and Arundinaria gigantea as common on the thermal Pen Branch and Four Mile Creek deltas. Lemna spp. (duckweed) was also commonly seen in the Pen Branch delta (Mackey, 1990). A band of Typha spp. (cattails) extended southeast from the Pen Branch delta along the Savannah River floodplain terrace edge (Mackey, 1990). This band of cattails has expanded by several hundred meters onto the delta since 1988 when thermal

discharges stopped. When high flows and temperatures are maintained, the mud and sand islands of thermal deltas will be dominated by Ludwigia spp., which although influenced by flooding and siltation effects can germinate and survive under these conditions (Christy, 1976; McCaffrey, 1982; Dolan and Sharitz, 1984; Christy and Sharitz, 1980; Mackey, 1990; McLeod and Sherrod, 1981). In 1987, K Reactor operated at nominal half-power and there was an expansion of the cattail communities (Table 4). Ludwigia occupied the shallow-water, mudflat areas of the central delta and the cattails expanded primarily in southeastern section of the delta.

Once thermal effluents were removed from these creeks the herbaceous wetland species composition changed rapidly. Ludwigia leptocarpa, Ammania coccinea and Boehmeria cylindrica began to decline as dominant species (Table 3). The cattail beds on the Pen Branch delta expanded and Sagittaria spp., Myriophyllum spp. and Polygonum spp. developed in areas of slightly deeper water. These initial changes are indications of changes to a less flooded and cooler delta. On elevated sites within the delta the 'old field' species noted above invaded. In addition seedlings of red maple, buttonbush, willow and cypress are evident in certain locations. Under reduced power operations, these 'old field' species would be expected to decline in abundance, Ludwigia would remain as a dominant, and the persistent emergent species would probably remain but not expand in coverage. Other annual species such as Lemna, Myriophyllum and Polygonum would be expected to remain as well, especially in the outer margins of the delta where water will be cooler and deeper.

The distribution of wetland plant species is influenced by hydrologic conditions in addition to temperature. Following the shutdown of K Reactor in 1988, stream discharge in Pen Branch declined sharply (Figure 4). The tolerance of common SRS wetland woody tree and shrub species to flooding and heat was evaluated by McLeod et al. (1987). Seedlings of bald cypress, water tupelo, black willow and buttonbush were grown from near ambient soil temperatures up to 10°C above ambient and from unsaturated (drained) to flooded soil (6 cm of standing water above the soil surface) for one year. The four study species acclimated and survived continuous flooding at 30°C and 35°C water temperatures, but flood water at 40°C reduced morphological acclimation to varying degrees (McLeod et al., 1987; Donovan et al., 1988). Adverse effects were observed in the 40°C flooding treatment in which only 20 percent of water tupelo survived. Survival of black willow (47 percent), bald cypress (60 percent), and button-bush (87 percent) was also reduced by flooding at 40°C. Flooding cypress and tupelo with 40°C water resulted in reduced root and stem biomass production, lower root starch content, impaired leaf function, and reduced ethylene production which may have resulted in the impaired morphological response to

flooding. Black willow exhibited less severe response to the high temperature flooding and buttonbush exhibited the greatest tolerance for these conditions.

Seedling and mature tree survival and growth can also be influenced by timing of flooding from thermal effluents as well as river flooding events (Scott et al., 1985; Schneider and Sharitz, 1988; Schneider et al., 1989; Sharitz and Lee, 1985). Extensive winter time flooding as well as long periods of flooding in the growing season limit woody species colonization and growth (Birch and Cooley, 1983). For example, seedlings of bald cypress 20 - 30 cm in height were all killed within two weeks by complete submergence by flood waters (Demaree, 1932). Many woody wetland species such as cypress, tupelo, willow and buttonbush tolerate inundation during the dormant season and brief flooding during the growing season (Witlow and Harris, 1979). Most seedlings, however, are killed if foliage is submerged for more than two weeks after they have leafed out (Broadfoot and Williston, 1973).

Extensive recruitment of young cypress (0.5 to 4 ft high) has occurred in recent years in the Pen Branch 'tail' portion of the delta. These seedling and young cypress are not normally flooded by flows from Pen Branch unless river flooding occurs also. They have survived frequent flooding in the winter of 1990-91 with depths greater than 1 meter on occasion. Given the lower temperatures of reactor restart at reduced power the low frequency of river flooding, these young cypress are likely to survive water temperatures resulting from low power operations during 1991-92.

With a restart of K Reactor at approximately half-power in 1991, much of the herbaceous revegetation of sand islands and mud flats in Indian Grave Branch and the middle and lower reaches of Pen Branch would be expected to be lost due to high temperatures and flows. The losses would consist primarily of grasses, and some early invading wetland species (e. g. cattails and Sagittaria spp.) along with scrub-shrub species such as willows and young bottomland hardwood species [e.g. red maple (Acer rubrum)]. In the lower Pen Branch corridor and the delta area, depending on river flooding patterns, Ludwigia spp. would be expected to dominate the Pen Branch delta. If temperatures remain low enough (i.e. 30-35°C) then species such as cattails, scrub-shrub seedlings and other wetland species such as young cypress may be able to survive flows from restart at half-power. The results of McLeod et al (1987) suggest that most of these woody species should survive at the expected water temperatures. Expansion of the tree kill zone on the Pen Branch delta, if it occurs, would slow with lowered temperatures as has been observed with Steel Creek and Beaver Dam Creek deltas (Christensen et al. 1988), but the level of change is difficult to predict.

Operation of K Reactor at 30% power levels would result in substantially lower winter and spring water temperatures, but relatively slight decreases in summer water temperatures relative to operation at half-power. These water temperatures should result in early emergence of some thermally tolerant macrophyte species, such as cattails, and earlier occurrence of such species as Polygonum spp. and Myriophyllum spp. in the lower stream corridor and delta regions of Pen Branch. Generally, herbaceous plant communities would shift to species more tolerant of higher temperature and increased flooding conditions. Early colonizing woody species, such as willow and buttonbush, would likely persist through the moderately thermal and increased flooding conditions. Young cypress that have become established in the Pen Branch delta 'tail' region would likely persist, unless the thermal conditions are accompanied by flooding of the swamp by Savannah River water. However, further recruitment of bottomland woody plant species is unlikely to occur until low flow and ambient water temperature conditions are restored.

However, revegetation of the delta would not be expected to continue at these flow levels (Sharitz et al. 1974; Sharitz and Lee 1985; Dunn and Sharitz 1987) because the slight increases in water depth would result in less area that is suitable for cypress seedling establishment. Long term, natural revegetation of the elevated sediment portions of the delta, as with Steel Creek and Four Mile Creek deltas, to cypress/tupelo forest is unlikely.

Macroinvertebrates

During reactor operation at full power, the thermally perturbed portions of Pen Branch were dominated by nematode worms and chironomids and tanytarsine midges (Wike et al. 1989). These organisms were probably able to survive in the stream either by burrowing deep into the sediments, or in the case of chironomids, by having sufficiently short generation times to be able to colonize the stream during brief reactor outages.

Macroinvertebrate studies conducted in Pen Branch subsequent to the initiation of the extended outage of K Reactor in April 1988 (during the fall of 1988 and the winter and spring of 1989) indicate that the macroinvertebrate community of Pen Branch recovered substantially during the first year of the reactor outage. During three months of sampling, at least 86 macroinvertebrate taxa were collected in the portions of the stream that had been thermally perturbed (Wike et al. 1989). Dominant taxa during recolonization included several species of orthoclad midges (Corynoneura nr. tarsis, Cricotopus spp., Orthocladus spp., and Tvetenia discoloripes gr.),

chironomini midges (Microtendipes pedellus, and Polypedilum spp.), tanytarsine midges (Rheotanytarsus distinctissimus and Tanytarsus spp.), and blackflies (Simulium tuberosum and S. vittatum). Other taxa that were abundant at one or more stations included the mayflies, Caenis sp. and Stenonema modestum/smithae; the stonefly, Perlesta placida; the caddisflies Cheumatopsyche spp. and Hydropsyche spp.; the beetle Macronychus glabratus, and danceflies (Empididae; Wike et al. 1989).

At 50% power, discharge temperatures at the reactor outfall will average approximately 40°C in the winter and 51°C in the summer. Based on data collected at Road A-13.2 during 1987 and 1988, it is anticipated that stream temperatures at Road A-13.2 near the delta will average about 35°C in the winter and about 43°C in the summer. Most species of macroinvertebrates are not able to survive in temperatures in excess of 33 to 35°C. Therefore, it is expected that Indian Grave Branch and Pen Branch, at least down to Road A-13.2 will be too warm to support all but the most thermally tolerant species of macroinvertebrates and that the organisms that presently inhabit these portions of the stream will be killed by the warm water or displaced downstream by the increased water velocities. The Pen Branch delta will probably be cool enough during the winter months to support a somewhat diverse community, but temperatures during the summer will eliminate the more sensitive species from the community, or restrict their colonization to the cooler side channels and margins of the delta. The elevated temperatures will also accelerate the development of many species of aquatic insects, which may result in early emergence of those species whose rate of development is temperature dependent. High mortality rates and/or reproductive failure may result if emergence occurs during periods of cold weather.

At 30% power, discharge temperatures at the reactor outfall will average approximately 31°C in the winter and 42°C in the summer (Table 2). Resulting temperatures in the stream corridor and delta will be cool enough to support a reasonably diverse community during the winter months (25°C at the delta), but stream temperatures during the summer months (37°C at the delta) will be too warm for all but the most thermally tolerant species. However, the slightly lower stream temperatures that would result from reactor operation at 30% power would extend the time period during which stream temperatures would be low enough to support most species, possibly by one to two months. Early emergence, with the risk of increased mortality and reproductive failure will also occur at 30% power.

Subsequent to start-up of the cooling tower, macroinvertebrates are expected to rapidly recolonize the stream. Based on data collected from Pen Branch in the year

following reactor shut-down in April 1988, and more extended recovery data from Four Mile Creek following the shut-down of C Reactor in 1985, it is anticipated that a reasonably stable and diverse macroinvertebrate community will develop in Pen Branch and Indian Grave Branch within two years of the time that the cooling tower is brought on line. Further changes in the community will occur in response to the long term changes in the vegetation and stream channel morphometry brought about by the decreases in discharge, water velocity, and temperature associated with cooling tower operation.

Fish

The average upper tolerance levels of SRS fish range from approximately 31 - 37°C. Centrarchids (e.g. bass, bream) and mosquitofish generally exhibited slightly higher tolerances than other taxa. Paller and Saul (1987) found no relationship between electrofishing catch per unit effort (CPUE) and temperatures below 30°C in SRS streams (Figure 5). CPUEs declined somewhat at temperatures between 30 and 35°C; however, the percentage of zero catches (60) was considerably higher than at temperatures under 30°C (10.5%). At temperatures above 35°C, catch per unit effort was generally zero.

Paller and Saul (1987) found that the most abundant taxa at temperatures approaching 35°C were sunfishes, largemouth bass, gar, and gizzard shad (Figure 5). They found little obvious difference among largemouth bass, spotted sunfish, warmouth, redbreast sunfish, redear sunfish, and bluespotted sunfish in terms of temperature tolerance; all except bluegill were collected at temperatures between 30 and 35°C. However, mosquitofish are abundant in SRS thermally impacted streams and occur at temperatures slightly above 40°C.

Maximum temperatures for the survival of fish embryos range from 20°C for the black crappie and yellow perch to 34°C for the bluegill and threadfin shad (Brungs and Jones, 1977). Field data on the spawning temperature requirements of SRS fishes were gathered by Paller et al (1987). Larval densities were generally highest (up to 290/1000 m³) at temperatures ranging from 10 - 26°C. Lower densities ($\leq$ 100/1000 m³) were observed between 27 and 35°C.

Expected stream water temperatures associated with K Reactor operation exceed the lethal level for most fish, which will result in elimination of fish from Indian Grave Branch and the upper reaches of Pen Branch. However, instream cooling may reduce water temperatures to acceptable levels further downstream in Pen Branch. The impacts of half-power operation will be acceptable in portions of Pen Branch that do not exceed approximately 33°C, decrease to at least 20°C during the spring (to ensure successful spawning), and do not

undergo rapid drops of more than about 15°C when the reactor goes down during winter. Consequently, when the reactor is operating fish will be excluded from Indian Grave Branch and much of Pen Branch throughout the year, but will likely inhabit portions of the delta during late fall through early spring.

Stream water temperatures with K Reactor operating at 30% power will permit some thermally tolerant species of fish to occur in Indian Grave Branch, Pen Branch and the Pen Branch delta during reactor operation. Water temperatures will be sufficiently high to preclude successful spawning by many species, however. Overall, operation of K Reactor at 30% power will allow a greater variety of fish to occupy the affected stream areas for a greater period of time during the year, but adverse effects of the thermal effluents will be evident in the fish community.

The response of fish to current velocity is species and life stage specific. Some fishes found in SRS streams, such as the northern hogsucker, prefer runs with relatively high current velocities while others, such as largemouth bass and sunfishes prefer the slow flowing or still waters found in pools (Meffe and Sheldon 1988). The latter group of fishes are seldom found in association with current velocities exceeding approximately 10 cm/s (Table 5). Because of their comparatively weak swimming ability, the early life stages of most species require lower current velocities than the adults.

Flow related impacts will not be as severe as temperature related impacts but will reduce habitat quality for fish in portions of Pen Branch areas where current velocities exceed species specific levels of tolerance. Current velocities will be highest in the relatively narrow, channelized upper and mid reaches of the Indian Grave/Pen Branch system. With the exception of stream edges, slack waters behind obstructions, and side channels, these reaches will have current velocities that exceed the levels of tolerance of most species. The lower reaches of Pen Branch, which are comparatively wide and braided or swampy, will undergo structural changes (i.e., scouring, erosion, and channel widening) in some areas as a result of increased flow but are likely to contain extensive areas where current velocities are acceptable to a range of SRS fishes.

Recovery of the fish community will begin as soon as half-power operation ceases. Fish will return to areas from which they were formerly excluded when temperatures and current velocities return to acceptable levels. Previous research on Four Mile Creek and Pen Branch indicated that recolonization by many species, especially larger and more mobile ones, began within 12 hours after reactor operations ceased (Aho et al. 1986). Community structure in Indian Grave Branch and

Pen Branch is likely to reach equilibrium with the existing habitat within several years after the cessation of half-power operations. At this time, the fish community will be very similar to that which existed in Indian Grave Branch and Pen Branch prior to the resumption of reactor operations.

American Alligator

The status of the American alligator has recently been changed from Endangered to Threatened by Similarity of Appearance (DOI 1987). Surveys conducted on the SRS during 1981-87 indicated that the alligator populations in the Pen Branch drainage were small and the population was not self-sustaining (Seigel 1989). These studies also indicated that under full power operating conditions, there was little threat of mortality to alligators in Pen Branch because the alligators responded rapidly to increasing temperatures by seeking refuge on land where they are capable of extended overland movements. Alligators actively avoid areas when water temperatures exceed 31°C, which is well below the lethal body temperature of 38°C.

The expected temperature regime in the Pen Branch drainage from the K-Reactor outfall to the delta will likely preclude the presence of alligators in the stream. However, no mortality should occur as a result of reactor operations, and alligators may be attracted to the Pen Branch delta during winter months because of the warmer, but sublethal, temperatures.

Wood Stork

The decline in wood stork populations in North America prompted the United States Fish and Wildlife Service of the Department of Interior to list the wood stork as an endangered species in 1984 (DOI 1984). The Birdsville Colony, the most northern and inland wood stork colony, is located at "Big Dukes Pond" near Millen, Georgia. This wood stork colony was believed to be the source of storks observed at the Steel Creek Delta in the SRS Savannah River swamp during 1980-1982. The SRS Savannah River swamp is 45 km from the Birdsville Colony.

Studies of wood storks at the Birdsville Colony indicated that the storks most frequently foraged in cypress and tupelo swamps with little understory vegetation (Coulter 1986, 1987, 1988, 1989; Hodgson et al. 1988). The median water depth at the location where storks landed at the foraging sites was 18 to 23 cm during the six years of the survey. Sunfish comprise the major component of the diet of birds from the Birdsville Colony.

Wood storks are most commonly observed in the SRS Savannah River swamp during July and August (Table 6). In 1983, storks were observed most frequently in July when storks were present during approximately 55% of the surveys, with an average of 30 storks per survey. The Steel Creek and Beaver Dam Creek delta areas had the greatest frequency (15%) of stork observations during the surveys and the greatest amount of wood stork use. The Pen Branch and the Four Mile Creek deltas had little stork use during the same period (Coulter, 1986).

The water depth in the SRS Savannah River swamp is dependent on the water flow from both the Savannah River and onsite streams. The river flow is primarily influenced by the amount of water discharged from the Clarks Hill Dam. The water level in the swamp also depends on flows from Beaver Dam Creek, Four Mile Creek, Pen Branch, and Steel Creek, and other smaller streams on the SRS.

The numbers of wood storks seen during aerial surveys of the SRS Savannah River swamp are greater in those areas of the swamp receiving lower flows, especially in the flow class of 51 to 100 cubic feet per second (cfs) (Table 7).

Wood stork use of the Pen Branch delta areas has been the lowest of all the delta areas on the SRS (Table 7). This has probably been the result of high flow and thermal conditions (Jensen et al., 1988) in this delta with flows in excess of 100 cfs most of the time (greater than 90% frequency) during the summer survey months from 1983 through 1987. No storks were observed during the summer months of 1988 under low flow conditions when the reactor was not operating. Low stork use was observed during the summer 1988 for the entire SRS Savannah River swamp.

Data from the aerial wood stork surveys of the SRS Savannah River swamp and the studies at the Birdsville Colony suggest that the SRS Savannah River swamp is probably not used extensively during the breeding nor pre-fledging phase of the Birdsville Colony. Most of the observations of storks on the SRS occur during late fledging or the post-fledging period. High flows (greater than 100 cfs) to the SRS Savannah River creeks probably limit the usefulness of the swamp as a foraging site. High flows (350 to 400 cfs) would be expected during the summer months from K-Reactor; therefore, the potential for stork foraging would be limited in the Pen Branch delta area. Low flows (less than 50 cfs) may likewise limit the usefulness of the SRS swamp for stork foraging. During periods of low flow (less than 50 cfs) the streams are more confined to their channels, and habitats are not suitable for stork foraging. During periods of high flow, water depths may be limiting. Flows and flooding patterns also can influence vegetation patterns, thus influencing available foraging sites. Typical wood stork foraging sites

have reduced quantities of both submerged and emergent macrophytes (Coulter, 1986). Moderate flows (greater than 50 cfs, but less than 150 cfs) to the SRS deltas during the summer months would probably benefit and/or assist in maintaining wood stork foraging sites in the SRS Savannah River swamp.

In 1985, Kathwood Lake was constructed near the SRS and began operation to mitigate the loss of foraging habitat in Steel Creek. Wood storks have used the foraging ponds extensively each year since summer 1986 and the project has compensated for the loss of foraging habitat in Steel Creek (Wike et al. 1989).

REFERENCES:

- Aho, J.M., C.S. Anderson, K.B. Floyd and M.T. Negus. Patterns in fish assemblage structure and dynamics in waters of the Savannah River Plant. SREL-27, UC-66e. Savannah River Ecology Laboratory, Aiken, SC. (1986)
- Aho, J. M., J. W. Terrell. Habitat Suitability Index Models and Instream Flow Suitability Curves: Redbreast Sunfish. Biological Report 82(10.119). U. S. Department of the Interior, Fish and Wildlife Service. (1986)
- Birch, J. B., and J. L. Cooley. The Effect of Hydroperiod on Floodplain Forest Production. Report ERC 08-83. Institute of Ecology, University of Georgia, Athens, Georgia 30602 (1983)
- Broadfoot, W. M. and H. L. Williston. "Flooding effects on southern forests," J. For. 71(9): 584-587. (1973)
- Brungs, W.A. and B.R. Jones. Temperature criteria for freshwater fish; protocol and procedures. EPA-600/3-77-061, U.S. Environmental Protection Agency, Environ. Res. Lab. Ecol. Res. Ser., 130 pp. (1977)
- Christensen, E. J. , J. R. Jensen, E. W. Ramsey, H. E. Mackey, Jr. "Aircraft MSS Data Registration and Vegetation Classification for Wetland Change Detection." Int. J. Remote Sensing. 9(1):23-38 (1988)
- Christy, E. J. Population Dynamics of Two Semi-aquatic Macrophytes Growing Under Different Temperature Regimes. Masters Thesis, University of Georgia, Athens, GA (1976)
- Christy, E. J. and R. R. Sharitz. "Characteristics of three populations of a swamp annual under different temperature regimes." Ecology 61(3): 454-460 (1980)
- Coulter, M. C. Wood Storks of the Savannah River Plant: Foraging and Breeding Ecology. Comprehensive Cooling Water Study Final Report. SREL-24, Savannah River Ecology Laboratory, Aiken, SC (January 1986).
- Coulter, M. C. Wood Storks of the Birdsville Colony and Swamps of the Savannah River Plant, 1986 Annual Report. SREL-31, Savannah River Ecology Laboratory, Aiken, SC (April 1987).

- Coulter, M. C. Wood Storks of the Birdsville Colony and Swamps of the Savannah River, 1987 Annual Report. SREL-33, Savannah River Ecology Laboratory, Aiken, SC (May 1988).
- Coulter, M. C. Wood Storks of the Birdsville Colony and Swamps of the Savannah River Plant, 1988 Annual Report. SREL-37, Savannah River Ecology Laboratory, Aiken, SC (1989).
- Demaree, D. "Submerging experiments with Taxodium. Ecology. 12 (3):258-262. (1932)
- Department of Interior (DOI). U.S. Fish and Wildlife Service. "Endangered and Threatened Wildlife and Plants: U. S. Breeding Population of the Wood Stork Determined to be Endangered." Federal Register. 49(40):7332-7335 (1984).
- Department of Interior (DOI). U. S. Fish and Wildlife Service. "Endangered and Threatened Wildlife and Plants; Reclassification of the American Alligator to Threatened Due to Similarity of Appearance Throughout the Remainder of It's Range." Federal Register. 52(107):21059-21064 (1987)
- Dolan, R. W. and R. R. Sharitz. "Population dynamics of Ludwigia leptocarpa (Onagracea) and some factors affecting size hierarchies in a natural population." J. Ecology. 72:1031-1041. (1984)
- Donovan, L. A. and K. W. McLeod. "Morphological and root carbohydrate responses of bald cypress to water level temperature regimes." J. Therm. Biol. 10 (4):227-232 (1985)
- Donovan, L. A., K. W. McLeod, K. C. Sherrod, Jr.k, and J. J. Stumpff. "Response of Wood swamp seedlings to flooding and increased water temperatures. I. Growth, biomass, and survivorship. Amer. J. Bot. 75(8):1181-1190 (1988)
- Falk, L. L. Savannah River Plant Temperature Study, Study 25845. E. I. Du Pont de Nemours & Co, Engineering Department, Wilmington, Delaware. (1956)
- Hodgson, M. E., J. R. Jensen, and H. E. Mackey. "Monitoring Wood Stork Foraging Habitat with Remote Sensing and Geographic Information Systems." Photogrammetric Engineering and Remote Sensing. 55(11):1601-11607 (1988).

- Irwin, J. E. Structure of Stump Communities in a Stream affected by Thermal Effluent. Masters Thesis, University of North Carolina Chapel Hill, NC. (1975)
- Jensen, J. r., E. J. Christensen, and R. Sharitz. "Mapping of thermally altered wetlands using high resolution multispectral scanner data." In: Renewable Resources Management - Applications of Remote Sensing, May 22-27, 1983. Proceedings, American Society Photogrammetry, Falls Church VA pp. 318-336. (1983)
- Jensen, J. R. , M. E. , E. Christensen, He. E. Mackey, Jr., L. R. Tinney, and R. Sharitz. "Remote sensing inland wetlands: a multispectral approach. "Photogrammetric Engineering and Remote Sensing. 52(1): 87-100 (1986)
- Jensen, J. R., E. W. Ramsey, H. E. Mackey, Jr., E. J. Christensen, R. R. Sharitz. "Inland wetland change detection using aircraft MSS data." "Photogrammetric Engineering and Remote Sensing. 53(5):521-529. (1987)
- Jensen, J. R., E. Ramsey, II. H. E. Mackey, Jr., and M. E. Hodgson. 1988. "Thermal modeling of heat dissipation in the Pen Branch delta using thermal infrared imagery. "Geocarto International. 1988 (4): 17-28. (1988)
- Mackey, H. E. Comprehensive Cooling Water Study, Final Report, Vol IV: Wetlands. DP-1739-4, E.I. du Pont de Nemours and Company, Savannah River Laboratory, Aiken, SC (1987)
- Mackey, H. E. "Monitoring seasonal and annual wetland changes in a freshwater marsh with SPOT HRV data. Proceedings, 1990 ACSM-ASPRS Annual Convention, Vol. 4, Image Processing/Remote Sensing, March, 1990, Denver CO. pp. 283-292. (1990)
- McCaffrey, C. A. Effects of Flooding and Sedimentation on Germination and Survival of Ludwigia leptocarpa (Nutt.) Hara Masters Thesis, University of Georgia, Athens, GA (1982)
- McLeod, K. W. and C. Sherrod, Jr. "Baldcypress seedling growth in thermally altered habitats." "Amer. J. Bot. 68(7): 918-923. (1981)
- McLeod, K. W., L. A. Donovan, and N. J. Stumpff. "Responses of woody seedlings to elevated flood water temperatures." In: The Ecology and Management of Wetlands, Timber Press, Portland, Oregon pp. 441-451. (1987)

- Meffe, G.K. and A.L. Sheldon. The influence of habitat structure on fish assemblage composition in southeastern blackwater streams. *The American Midland Naturalist*, 120(2): 225-240. (1988)
- Mertz, S. M. Cooling Water Effluent Temperatures (U). RRD-RTS-910188. (1991)
- NUS. Thermal Modeling of K-Reactor Cooling Water Discharge. Fax from A. Toblin to M. Paller, J. Gladden. June 17, (1991).
- Paller, M.H. and B.M Saul. Effects of thermal discharges on the distribution and abundance of adult fishes in the Savannah River and selected tributaries, annual report, November 1984 - August 1985, ECS-SR-28. (1987)
- Paller, M.H., B.M Saul, and D.W. Hughes. The distribution of ichthyoplankton in thermal and nonthermal creeks and swamps on the Savannah River Plant, February - July 1985, DPST-86-796. (1987)
- Schneider, R. R. and R. R. Sharitz. "Hydrochoroy and regeneration in a bald cypress-water tupelo swamp forest." Ecology. 69 (4): 1055 - 1063 (1988)
- Schneider, R. L., N. E. Martin, and R. R. Sharitz. "Impact of dam operations on hydrology and associated floodplain forests of southeastern rivers." In: Freshwater Wetlands and Wildlife CONF- 8603101, DOE Symposium Series No. 61, R. R. Sharitz and J. W. Gibbons (Eds). USDOE Office of Scientific and Technical Information, Oak Ridge, TN pp. 1113 -1122. (1989)
- Scott, M. L., R. R. Sharitz, and L. C. Lee. "Disturbance in a cypress-tupelo wetland: an interaction between thermal loading and hydrology." Wetlands 5:53-68. (1985)
- Seigel, R. A. Population Status and Ecology of the American Alligators on the Savannah River Plant. Savannah River Ecology Laboratory, Aiken, SC (1989)
- Sharitz, R. R. , L. C. Lee. "Limits on Regeneration Processes in Southeastern Riverine Wetlands." In: *Riparian Ecosystems and Their Management: Reconciling Uses*. First North American Conference, April 16-18, 1985, Tucson, AZ, pp. 139-160 (1985)
- Sharitz, R. R., C. P. Dunn. Revegetation of a Taxodium-Nyssa Forested Wetland Following Complete Vegetation Destruction. *Vegetation*, 72:151-157. (1987)

- Sharitz, R. R., J. E. Irwin, and E. J. Christy. "Vegetation of swamps receiving reactor effluents." Oikos. 25:7-13. (1974)
- Specht, W.L. Comprehensive Cooling Water Study, Final Report, Vol V: Aquatic Ecology. DP-1739-5, E.I. du Pont de Nemours and Company, Savannah River Laboratory, Aiken, SC (1987)
- Stuber, R. J., G. Gebhart, O. E. Maughan. Habitat Suitability Index Models: Bluegill. FWS/OBS-82/10.8. U. S. Department of the Interior, Fish and Wildlife Service. (1982a)
- Stuber, R. J., G. Gebhart, O. E. Maughan. Habitat Suitability Index Models: Largemouth Bass. FWS/OBS-82/10.16. U. S. Department of the Interior, Fish and Wildlife Service. (1982b)
- Whitlow, T. H. and R. W. Harris. Flood Tolerance in Plants: A State-of-the-Art Review. Technical Report E-79-2, US Army Engineer Waterways Experiment Station, Environmental Laboratory, Vicksburg, Miss. (1979)
- Wike, L. D., W. L. Specht, H. E. Mackey, M. H. Paller, E. W. Wilde, A. S. Dicks. Reactor Operation Environmental Information Document, Volume II, Ecology. WSRC-RP-89-816. Westinghouse Savannah River Company, Aiken, SC. (1989)

Table 1 Pen Branch Water Temperatures (°C) During Operation in 1955*

DATE	MEASUREMENT LOCATION			
	<u>DISCHARGE</u>	<u>RD.B</u>	<u>RD.A</u>	<u>RD.A-13.2</u>
4/18**	41.6		34.5	
4/26**	40.4		34.4	
5/3**	40.0		35.0	
5/10**	42.0		37.0	
5/23**	41.2	42.1	37.5	33.9
5/31**	45.1		39.3	
6/7**	45.3		38.0	
6/13**	45.2		41.3	
6/21**	45.3		40.3	
6/23	48.4	47.0	43.6	38.9
6/28**	47.2		38.1	
7/5**	47.4		36.8	
7/12**	48.4		39.1	
10/6	49.8	48.2	42.8	37.1
10/21**	47.8			34.3
10/27**	47.1			31.5

* Source: Falk, 1956, Table IX, Only data for discharge temperatures of 40-50°C are presented.

** Discharge temperature estimated from production data.

Table 2 Predicted February and August Water Temperatures
(°C)^a in the Pen Branch Drainage Under 50% and 30%
Power Operation Conditions at the K Reactor.
(Modified from NUS, 1991)

Month	LOCATION				
	Ambient	Discharge Pipe	End of Canal	Hwy. 125 (SRS Rd.A)	Entrance to Swamp
<u>50% Power</u>					
February	12	40	48.7	36.3	29.9
August	26	51	49.6	48.3	41.7
<u>30% Power</u>					
February	12	31	30.2	28.7	24.8
August	26	42	41.2	40.4	36.8

a. Simulation results based on average meteorological conditions at
Bush Field, Augusta, GA.

Table 3. Typical Wetlands Classes and Species Found On or Near the SRS Savannah River Swamp Thermal Deltas

CLASS	COMMON NAME	SCIENTIFIC NAME
Open Water		
Algal Mat		
Nonpersistent Emergent Marsh	Water primrose Ammannia	<u>Ludwigia</u> spp. <u>Ammannia coccinea</u>
Persistent Emergent Marsh	Knotweed Cutgrass False nettle	<u>Scripus cyperinus</u> <u>Leersia</u> spp <u>Boehmeria cylindrica</u>
Scrub/Shrub	Willow Buttonbush	<u>Salix</u> spp. <u>Cephalanthus occidentalis</u>
Mixed Deciduous Swamp Forest	Bald cypress Tupelo gum	<u>Taxodium distichum</u> <u>Nyssa aquatica</u>
Mixed Deciduous Upland Forest	Oak Loblolly pine Sweetgum Red maple Hickory	<u>Quercus</u> spp. <u>Pinus taeda</u> <u>Liquidambar styraciflua</u> <u>Acer rubrum</u> <u>Carya</u> spp.

Sources: Christy, 1976
Christy and Sharitz, 1980
Jensen et al., 1983
Jensen et al., 1986
Jensen et al., 1987

Table 4. Areas (hectares) of Major Wetland Cover Types Based on Classification of Satellite Imaging Data for the Pen Branch Delta, 1987 Through Spring 1989 (Mackey, 1990).

Wetland Cover Type	Date and Year of HRV Data Used											
	1987				1988				1989			
	Apr 09	Apr 24	May 24	Aug 22	Oct 22	Apr 17	May 02	May 08	Oct 25	Jan 28	May 17	
1. Deep/Open Water	10.4	13.1	13.6	1.5	11.8	4.8	4.0	1.2	0.6	0.0	0.0	
2. Non-Persistent Emergent Marsh (NPE)												
2a. Shallow Water/ Mud Flats	82.1	65.5	33.4	12.7	75.0	44.4	29.0	0.2	56.2	0.1	0.4	
2b. Ludwigia spp.	13.6	34.0	43.5	55.4	-	30.9	56.8	88.2	53.9	155.9 ^a	97.4 ^a	
2c. Duckweed	-	3.2	-	-	0.3	7.5	10.1	0.3	-	-	3.6	
3. Persistent Emergent Marsh (PE)												
3a. Cattails	10.7	31.2	29.5	65.7	21.0	23.0	38.5	*	*	*	*	
4. Total	116.8	147.0	119.0	135.3	108.1	110.6	138.4	89.9	110.7	156.0	101.4	

^a After mid-May 1988, it was not possible to distinguish between the non-persistent beds of Ludwigia spp. and stands of cattails with the SPOT HRV data.

Table 5. Preferred and Maximum Current Velocities (cm/sec)
for Several Species of Fish Found in SRS Streams.

Species	Preferred	Maximum Tolerable	Reference
Bluegill			
adult	<10	45	Stuber et al. (1982a)
juvenile	<5	15	Stuber et al. (1982a)
fry	<5	12	Stuber et al. (1982a)
embryo	<8	30	Stuber et al. (1982a)
Largemouth bass			
adult	<10	20	Stuber et al. (1982b)
juvenile	<10	20	Stuber et al. (1982b)
fry	<1	2.6	Stuber et al. (1982b)
embryo	<6	10	Stuber et al. (1982b)
Redbreast sunfish			
adult	<35	-a	Aho and Terrell (1986)
juvenile	<35	-a	Aho and Terrell (1986)
fry	<20	-a	Aho and Terrell (1986)
embryo	<20	-a	Aho and Terrell (1986)

^a Data not available

Table 6. Total Numbers of Wood Storks Observed in the SRS
Savannah River Swamp, June through September, 1983-
1988.

Period	Number of Storks	Percentage
Before June	3	0.2
June 1 -15	88	6.1
June 16 - 30	66	4.5
July 1 - 15	539	37.1
July 16 - 31	315	21.7
August 1 - 15	139	9.6
August 16 - 31	156	10.7
September 1 - 15	37	2.5
September 16 - 30	94	6.5
After September	15	1.0
Total	1452	99.9

Table 7. Total Numbers of Wood Storks Observed in the SRS
Savannah River Swamp, June Through September, 1983-
1988.

Flow Classes in the Creeks (cfs)	<u>Number of Wood Storks</u>					Total	Percentage	Cumulative Percentage
	BDC	FM	FMS	PB	SC			
0-50	0	97	3	0	92	192	12.43	12.43
51-100	291	291	235	27	249	1093	70.74	83.17
101-150	1	0	0	0	67	68	4.40	87.57
151-200	-	0	8	0	0	8	0.52	88.09
201-250	-	0	16	1	0	17	1.10	89.19
251-300	-	0	0	10	0	10	0.65	89.84
301-350	-	0	0	0	0	0	0.00	89.84
351-400	-	0	0	6	0	6	0.39	90.23
401-450	-	36	101	4	0	141	9.13	99.36
> 450	-	0	0	10	-	10	0.65	100.01
Total	292	424	363	58	408	1545	100.01	

cfs = cubic feet per second

BDC = Beaver Dam Creek Delta area

FM = Four Mile Creek Delta area

FMS = Four Mile Creek sloughs between Four Mile Creek and Pen Branch
deltas

PB = Pen Branch Delta area

SC = Steel Creek Delta area

Figure 1. Map of Pen Branch Drainage.

Figure 2. Predicted Reactor Power Levels and Effluent Temperatures from K Reactor (Mertz, 1991)

Figure 3. Historic Patterns of Water Discharge and Temperature from K Reactor, and Wetland Tree Canopy Loss in the Pen Branch Corridor and Swamp.

Figure 4. Flow in Pen Branch from January, 1985 through June, 1989 (Mackey, 1990).

Figure 5. Catches (Number of Fish/100 meters) of Fish Species in SRS Streams in Relation to Water Temperature.

END

**DATE
FILMED**

8 / 10 / 92