EGG--2656

DE92 016826

Public Acceptability of the use of Gamma Rays from Spent Nuclear Fuel as a Hazardous Waste Treatment Process

**B. J. Mincher
R. P. Wells
H. J. Reilly**

Published January 1992

**Prepared by the
Idaho National Engineering Laboratory
EG&G Idaho, Inc.
for the U.S. Department of Energy
Office of Environmental Restoration and
Technology Development
DOE Idaho Field Office
Under Contract No. DE-AC07-76ID01570**

MASTER

EP

ABSTRACT

Three methods were used to estimate public reaction to the use of gamma irradiation of hazardous wastes as a hazardous waste treatment process. The gamma source of interest is spent nuclear fuel. The first method is Benefit-Risk Decision Making, where the benefits of the proposed technology are compared to its risks. The second analysis compares the proposed technology to the other, currently used nuclear technologies and estimates public reaction based on that comparison. The third analysis is called Analysis of Public Consent, and is based on the professional methods of the Institute for Participatory Management and Planning. The conclusion of all three methods is that the proposed technology should not result in negative public reaction sufficient to prevent implementation.

CONTENTS

ABSTRACT	iii
INTRODUCTION	1
BENEFIT-RISK DECISION MAKING	2
COMPARISON TO EXISTING IRRADIATION TECHNOLOGIES	3
Waste Water Irradiation	3
Medical Irradiations	4
Miscellaneous Irradiation Technology	5
Food Irradiation	6
Comparison of Current Irradiation Technologies	8
ANALYSIS OF POTENTIAL FOR PUBLIC CONSENT	9
Legitimacy Questions	9
Potential Issues	9
Proper Conduct	10
CONCLUSIONS AND RECOMMENDATIONS	12
REFERENCES	13

INTRODUCTION

Organochlorine pollutants are widely disseminated in the natural environment. Of special concern are those pollutants that persist because of chemical resistance to natural modes of decomposition, namely pesticides and polychlorinated biphenyls (PCBs). Some compounds have no acceptable treatment methods and even incineration does not guarantee zero release of PCBs. Hazards linked to the above compounds range from toxicity to potential carcinogenicity. This environmental situation has spawned much research in the past several decades, including using nuclear technology as a source of energy to promote the decomposition of hazardous organochlorine compounds.

Research is currently underway at the Idaho National Engineering Laboratory (INEL), [in collaboration with Lawrence Livermore National Laboratory (LLNL)] examining decomposition rates and products of PCBs exposed to gamma irradiation.¹ This work is based on studies of pesticide and PCB decomposition found in the literature of research pioneers Sherman et al.^{2,3} They postulated mechanisms based on free-radical induced chain dechlorinations. Numerous authors have performed more recent similar studies, including Schweitzer et al.⁴ and especially Sawai et al.⁵ and Singh et al.^{6,7}

Research at the INEL is concentrated on expanding these studies. Using higher gamma ray doses available at the spent fuel pool of the Advanced Test Reactor, research indicates that PCBs decompose beyond simple dechlorination, with resultant production of volatile organic degradation products.⁸ PCBs are totally destroyed by sufficient gamma ray exposure. Further research proves that other organochlorines are susceptible as well.

The use of spent fuel sources of gamma rays for hazardous waste decomposition is particularly efficient because one waste problem is used to solve another. Because fuel must be stored anyway, it is a source of free gamma rays, and thus a potentially cost effective source for treating hazardous waste. Spent fuel is available from operating reactors throughout the U.S.

The INEL's goal is design and demonstration of a hazardous waste treatment process, employing spent reactor fuel as a source of gamma rays, for decomposition of organochlorine pollutants. Selected PCBs are the initial compounds being tested in laboratory experiments. A treatment facility is envisioned where wastes (such as PCB contaminated transformer oils) flow at a predetermined rate; either around or located adjacent to spent fuel. These gamma irradiations could occur in a preexisting spent fuel storage facility at the reactor site. Because gamma radiation is highly penetrating, multiple layers of containment would prevent radioactive contamination of the treated solution. Treatment would continue until enough dose was absorbed thereby rendering the wastes nonhazardous. The technology exhibits an ability to clean up many waste types, from contaminated groundwater to stockpiled drums of hazardous waste.

This paper addresses the probable public acceptance of such technology. Three methods are used to assess public acceptance. The first approach is Benefit-Risk Decision-Making.⁹ This approach compares economic and health benefits of the proposed technology to its estimated risk. A second compares proposed technology to similar, and in some cases, accepted nuclear technology. The third, called Analysis of Public Consent, investigates the elements believed to contribute to public consent for new technologies.

BENEFIT-RISK DECISION-MAKING

The rate at which any technology becomes exploited is largely controlled by the public acceptability of that technology. Public acceptance can be based on several things: reduced consumer costs, minimization of public risk, and enhancement of life and the environment.⁹ Public acceptance of a technology can also be widely influenced by both accuracy of public information and media coverage. Benefit-Risk Decision-Making is a method of estimating public acceptance where economic or other benefits are compared to human risks.⁹

Spent fuel pools are operating in government and commercial facilities throughout the U.S. Circulation systems or dry tubes (for irradiation of hazardous wastes) do not require significant design changes for existing facilities that currently contain circulation systems for cooling and dry tubes for insertion of experiments. The addition of hazardous waste irradiation facilities does not impose further risk.

The potential for radioactive contamination of the hazardous waste by the fuel needs to be addressed. An escape of fission product activity from the fuel requires a breach of fuel cladding; a rare event even when the fuel is exposed to the high temperatures of reactor operation. The fuel fabrication industry has invested substantial time and resources into the integrity of reactor fuel cladding. However, in the unlikely event of leaking fuel, the highly penetrating nature of gamma rays allows for the design of multiple containment layers between the fuel and the hazardous waste being treated. Thus, a double or triple failure must occur to contaminate the hazardous waste container. These waste systems would not require elevated pressure or temperature. Fuel pool water is already maintained to minimize corrosion and is ion exchanged, which limits contamination spread. Radiological contamination of hazardous waste is highly unlikely. Standard operating procedures ensure that contamination is easily detected before material is released to the environment. Risks associated with transport of highly radioactive materials can be avoided by treating hazardous

wastes onsite (where spent fuel is produced). For some wastes, such as PCBs, the same owner is often involved—the commercial electric power company.

The risks of using spent fuel irradiators are minimal or already accepted; the benefits are many. The nation's most vexing hazardous waste problems might be solved by the application of this technology. Some very stable hazardous compounds have been demonstrated susceptible to gamma-ray induced decomposition.¹⁻⁸ For many of these wastes, few acceptable mitigation techniques exist. Polychlorinated biphenyls have been incinerated; however, this process is often inefficient, hard to monitor, and therefore may release hazardous compounds to the environment.

Hazardous organics in aqueous solution, such as trichloroethylene or carbon tetrachloride, are usually collected on activated carbon. This material then becomes the hazardous waste because the organochlorines have merely found a new place to reside. Gamma irradiation offers the possibility of total destruction of these and other hazardous compounds to nonhazardous decomposition products.

Mixed hazardous wastes that contain both radioactive and hazardous components are an especially difficult problem, because limited methods exist at sites for treating or storing such materials. If the hazardous component of mixed wastes is eliminated it would significantly simplify disposition of waste.

A great benefit of gamma-ray irradiation would be a cleaner environment. Economic benefits may be envisioned as well. As industry solves some of its more difficult environmental compliance issues, operating costs may decline and this might affect the costs of many consumer goods. The cost of modifications to existing spent fuel pools would probably be small. Thus, a simple benefit-risk comparison predicts that the proposed technology will be acceptable to the public.

COMPARISON TO EXISTING IRRADIATION TECHNOLOGIES

Waste Water Irradiation

A process similar to hazardous waste irradiation is waste water and sewage irradiation. Both technologies use radiation sources to treat potentially health threatening waste material before safely returning it to the environment. Epidemics of typhoid fever, cholera, parasitic infections, and viral diseases have resulted from the use of untreated sewage in agriculture.¹⁰ Offshore sewage disposal has produced hazardous swimming conditions in New York and a current cholera epidemic in Peru is related to polluted seafoods.

Thermal pasteurization of sewage sludge is the only conventional method available for disinfection, before it returns to the environment for agricultural use. Objections against the use of sewage sludge are based on the presence of pathogens and parasites. Sterilization by irradiation offers numerous advantages over thermal pasteurization, including destruction of viruses not killed by thermal treatment, destruction of weed seeds, no generation of obnoxious odors, and decreased use of fossil fuels. In fact, no conventional treatment offers an effluent free of pathogens.¹⁰ An irradiation treatment facility is also simpler to operate, maintain, and control as sterilization occurs at low, noncorrosive temperatures and irradiated sludge does not require flocculation or decantation facilities.¹¹

Irradiators demonstrate high sterilization efficiency of waste waters as compared to conventional thermal treatment. Studies at the Massachusetts Institute of Technology show a 99.99% kill of bacteria and viruses using electron beams.¹² Gamma rays should provide similar efficiency with higher penetrating ability. Herrenberger¹³ compared spent fuel to ⁶⁰Co sources for sewage sludge irradiation and found fuel to be economically competitive. A review of waste water irradiation studies was given by Gerrard¹⁴ and similar high kill efficiencies were demonstrated for various bacteria, amoeba, and bacteriophages in numerous studies in four countries. Other benefits reported in the review include increased sedimentation of particulates,

detergent foam elimination, and decomposition of cyanide.

Probably the most famous sewage sludge irradiator is at Geiselbullach, near Munich, Germany.¹¹ The facility started operation in 1973 and uses a 120 KCi ⁶⁰Co source to process sludge at 30 m³/day to 300 KR. The source is triply encapsulated ensuring radioactive contamination of sludge does not occur. The sterilized sludge, free of salmonella and parasite eggs, is used as fertilizer.

A commercial sewage irradiator was started up at Palmdale, Florida, March 10, 1970.¹⁵ The plant has a nominal capacity of 10⁴ gal/day and uses a 16 KCi ⁶⁰Co source. It produces irrigation quality water at the effluent, including destruction of viruses. The authors of a paper about Palmdale suggest that adding irradiators to conventional sewage treatment plants would eliminate sewage bypass around treatment during operating difficulties.¹⁵ Irradiators have no moving parts and extremely low downtime.

The benefits of waste water irradiation appear to be numerous; the only conceivable disadvantages are those common to all irradiator technology. Transportation of radioactive materials has been receiving increased media attention and appears to be experiencing some public resistance. Isotope sources require periodic replenishment as they decay that necessitates routine shipments of radioactive materials. The negative public attitude toward shipping of radioactive materials is unwarranted. In Canada, where ⁶⁰Co food irradiation is used, 190 million curies of ⁶⁰Co were shipped between 1955 and 1988¹⁶ and involved 870 separate shipments. Not one shipment caused the release of radioactive materials to the environment or presented any danger to the public. In the U.S., there were 51 accidents associated with approximately 40,000 radioactive shipments since 1971.¹⁷ No shipping cask has ever failed.

The escape of radioactivity is a potential problem associated with isotope sources. However, isotope sources rely on penetrating gamma rays and allow multiple layers of containment between the source

and the environment. Sources are contained in cells with no moving parts and are maintenance free. The contamination hazard presented is virtually nonexistent except for rather routine operations involved in replacing spent source material. The infamous instances of public exposure to isotope source radioactivity in Latin America have resulted from irresponsible abandonment of small, highly radioactive sources where engineering and administrative controls were purposefully defeated.¹⁸ Such acts are akin to sabotage and should not be used as an argument against safe and valuable technology. In fact, after more than 20 years of operation, the public has neither been endangered nor exposed to radiation from any irradiator operated by the Atomic Energy of Canada Limited (AECL), where the technology is finding wide use.¹⁹ In summation, the transportation of isotope sources has an excellent safety record, and the penetrating nature of gamma rays allows for multiple layers of containment.

The literature does not document public resistance to the concept of waste water or sewage sludge irradiation. Economic competition with more conventional heat treatment and chlorination are probably responsible for the low visibility irradiation has gotten in the waste water treatment industry. Cost effectiveness of the process could easily change should effluent standards become more restrictive, fossil fuels more expensive, and water increasingly scarce.

In conclusion, waste water irradiation attracts little negative public comment; possibly due to the obvious public health and environmental benefits. It is interesting to note that this has occurred despite the fact that the irradiated waste water is returned to the environment.

Medical Irradiations

The most well established industrial use of gamma-ray irradiations is the sterilization of medical products. The popularity of disposable plastics in medicine has resulted in many products, such as disposable syringes, which can only be sterilized by irradiation.²⁰

Radiosterilization of medical products was introduced to the U.S. in 1956 by Ethicon.^{20,21} Since 1964, all sterilization of sutures has been accomplished by gamma irradiation with ⁶⁰Co sources.²¹ Before 1964, accelerated electrons were used. In 1967, a Code of Practice for Radiosterilization of Medical Products was adopted by the International Atomic Energy Agency (IAEA).²⁰ The Code recommends hygiene, packaging, and biological calibrations of the radiation source; and is meant to be used as a guideline for IAEA member nations to establish indigenous legislation.

Great Britain has employed radiation sterilization of medical products since the mid-60's, and designed the first gamma irradiation plants. By 1966, four ⁶⁰Co plants and one electron beam plant were in operation.²⁰ Early microbiological studies show a 2.5 MRad dose was necessary to ensure product sterility.

In addition to disposable plastic, sutures, blood transfusion equipment, and tissue grafts are sterilized by irradiation. There are many benefits to radiation as opposed to more conventional sterilization methods. The usual conventional methods of steam or dry heat sterilization are too rigorous for disposable plastics and tissue grafts, and sutures require lengthy drying and heating, which makes them stiff and brittle.¹⁸ For these reasons, radiation sterilization of sutures and most plastics is preferred. In fact, the rapid development and use of disposable sterile plastics would have been impossible without radiosterilization.²⁰

Chemical sterilization (usually wet alcohol or ethylene oxide) may be too rigorous on some materials, and undesirable residues can result. Blood transfusion equipment sterilized with toxic gases has caused adverse patient reactions.²⁰ Toxic chemical traces are obviously undesirable on tissue grafts, and tissue grafts cannot be freeze dried. A properly balanced dose to tissue grafts changes antigenic properties that can help prevent tissue rejection.²⁰ Past tissue irradiation testing included nerves and dura matter treated with electrons; and bone, heart valves, and fascia later treated with gamma rays from spent fuel.²⁰ Based on experience with more than 15,000 tissue transplants, Poland's Central Tissue Bank offers justification

for widespread use of radiosterilization.²¹ General advantages of tissue irradiation include long durability of the irradiated tissues, suppression of antigenicity, high degree of sterility, and easy transport and storage of the sterile product.

General advantages of medical product radiation treatment include the following: increased bactericidal sterility, ability to sterilize thermolabile materials, possibility to treat materials of different shapes, sizes, and container types (including porous containers), and the possibility of a continuous process sterilization.^{22,23} In addition to disposables and tissue, products commonly irradiated are intravenous feeding tubing, catheters, pacemakers, saline water, eye drops, vaccines, and vitamins.^{20,24}

Few drawbacks are foreseen for radiation treatment of medical products. The previous comments concerning radioactive sources can be made and were addressed in this paper. A primary concern of radiation sterilization of medical products is product degradation. Extensive testing for effects of irradiation was performed on plastic and rubber materials used in medical supplies. In general, plastics may be classified as predominantly degraded by radiation or predominantly crosslinked. Materials having a high crosslink ratio have high radiation resistance.²³ For example, most rubbers are stable at sterilizing doses of radiation, thus the process is applicable to known specific materials.

In conclusion, isotope radiosterilization of supplies, equipment, some tissues, and pharmaceuticals remains the state-of-the-art process for producing safe medical products. The method has a long successful history and is credited with making the use of disposable medical supplies possible. Available literature contains no references to adverse public reaction to this technology. It is possible that confidence in the medical profession concerning medical matters has led to public trust and acceptance of this technology.

Miscellaneous Irradiation Technology

In addition to waste water and medical products irradiations, the use of isotope and accelerator radiation sources is applied to other less known processes. Radiation is used as a free radical generating mechanism for the catalysis of certain chemical reactions. Radiation polymer crosslinking has been performed commercially in the U.S. since the early 1960's.²⁵ Cross-linking of polymers produces a plastic possessing higher thermal stability. Radiation processing results in a product without residual chemical catalysts. Graft copolymers are also produced in this fashion by a free radical mechanism. These products are used for battery separators, ion exchange membranes, and liquid viscosity modifiers.

Ethyl bromide synthesis, conducted by Dow Chemical, was the first industrial process using isotope irradiation.²⁵ Three thousand curies of ⁶⁰Co is used to produce one-million pounds of product per-year at a plant running continuously. The plant requires minimal supervision or maintenance.

Wood, plastic, and concrete are pressure impregnated with monomers for irradiation. Upon irradiation, the monomer polymerizes and a composite is produced possessing higher physical and mechanical properties.^{25,26} Suggestions for the use of improved concretes include improved highway pavements and underwater applications requiring low water permeability. A major product of impregnated irradiated wood is parquet flooring.

Radiation vulcanization of rubber is being researched in Japan and other countries with advantages being the production of rubber without nitrosamines and low cytotoxicity.²⁷ One application is production of highly crosslinked nondermatite rubber gloves.

Finally, composites of waste materials and polymers were experimented with. This includes a composite of methyl methacrylate and sewage waste solids and methyl methacrylate and crushed glass for building materials.²⁸

These technologies produce new materials or significantly improved materials using what has become standard, safe radiation technology. Many of these processes are unknown to the public who have come to accept and rely upon their products.

Food Irradiation

Although some might offer it as a counterexample (of the hypothesis of this study) it is impossible to ignore food irradiations when discussing irradiation applications. It can be argued that food irradiation, which has not been widely accepted in the U.S., possesses an important difference from those irradiation technologies in use.

Perhaps the most controversial of all irradiation applications is the irradiation of foods. Since a U.S. patent was issued in 1921 for elimination of *Trichinella* in foods,²⁹ at least 50 countries have pursued the use of ionizing radiation as a means for improving safety or product "shelf life;" and as of 1988, more than 260 clearances were given in over 30 countries for food treatment.^{29,30} Of these countries, 19 are applying this technology commercially. Since the early 1970's, Canada, Israel, the Netherlands, and the USSR have irradiated foods for public consumption.³¹ In general, the three main benefits are spoilage elimination, elimination of microorganisms, pest eradication, and retardation of ripening or sprouting ("shelf life" extension).

Any technology that would disinfest stored foodstuffs and extend shelf life would benefit developing as well as industrialized nations. Warm climates in Third World countries often result in rapid food spoilage. India estimates stored wheat losses (due to insects) at 20 to 50%, and sprouting and rotting losses of onions at 25 to 30%.³² This is roughly three times the amount of onions exported, robbing India of much needed revenue. Indonesia estimates more than 10% stored foodstuff is destroyed by insects.³³ Indonesian and Indian waters are fish abundant, yet as a valuable protein resource it is barely exploited, because of inadequate distribution, transportation, and storage facilities. Irradiation of fish could extend shelf life without refrigeration, thus alleviating these difficulties. The United Nations Food and Agriculture

Organization (FAO) estimates that one-fifth of the world's produced food supply is destroyed by insects and microorganisms.³⁴ Estimates for Africa are even higher.

Ionizing radiation acts as a cytotoxic agent to prevent contamination and spoilage of food. Studies in Hawaii demonstrated disinfestation of tropical fruits from fruit flies and weevils using radiation doses of 25 KRad.³⁵ Mango seed weevils are not destroyed by commercial fumigation procedures. Shelf life extension of tropical fruits has also been reported. The Hawaiian Development Irradiator (HDI) Project operated between 1967-1973 and used a 220,000 Ci ⁶⁰Co source. Delayed ripening of mangoes and bananas, due to irradiation, was also reported in India as was inhibition of sprouting in onions, potatoes, and disinfestation of wheat.³² Canadian studies of preservation of foods using ⁶⁰Co irradiation have demonstrated sprout inhibition, disinfestation, shelf life extension, and pathogen elimination.³⁶ There have been numerous similar studies worldwide.

Despite the obvious benefits of food irradiation, it has not gained widespread acceptance in the U.S. Several government agencies, like the Occupational Safety and Health Administration (OSHA), the Nuclear Regulatory Commission (NRC), and the Department of Transportation (DOT) have long standing procedures for the control and monitoring of radiation sources and processes. Major international organizations such as FAO, the World Health Organization (WHO), and the IAEA have recognized and endorsed food irradiation.^{32,17} National health organizations such as the American Medical Association (AMA) and the American Council on Science and Health have followed suit.¹⁷ Irradiation of spices, produce, pork, and wheat has been approved by the U.S. Food and Drug Administration (USFDA).³⁷ Food irradiation in the U.S., apart from spices, is primarily limited to demonstration projects such as the HDI and the Cs-137 Agricultural Commodities Irradiator (CACI).³⁸

A forum for the transfer of knowledge called the International Consultative Group on Food Irradiation (ICGFI) was established by 15 governments, including the U.S., on May 9, 1984. In April 1988,

experts from 24 nations convened to share recent global developments on food irradiation; some of this information is referenced in this paper.³⁹

There are three major problems related to food irradiation that hamper future domestic development: public perception of induced radioactivity in irradiated products, chemical changes caused by irradiation, and the transport of radioactive sources (previously discussed). The first two are addressed as follows:

The problem of induced radioactivity in irradiated foods is one of perception rather than technology. Food irradiation is performed with pure gamma emitters, such as ⁶⁰Co, or machine generated electrons. If spent fuel were used as a source, estimates suggest natural neutron activation levels would be higher than neutron activation due to irradiation with fuel of normal burnup.³⁹ Isotope gamma rays do not induce radioactivity. Machine electron sources are limited to energies less than 10 MeV and machine x-ray sources to 5 MeV for irradiation of foods by the Codex General Standard for Irradiated Foods to prevent induced radioactivity.³⁹ Others suggest the limit is conservative.⁴⁰ The Codex Standard is the official approval and adoption standard for food irradiation by the IAEA, the FAO, and the WHO. Food irradiation does not induce product radioactivity. Containment in multiple layers can prevent foodstuff contamination that may result from a leaking gamma source.

The potential for chemical changes to foodstuffs because of irradiation have caused general concern. Currently, no radiolytic product has been discovered unique to irradiation. The chemical changes produced by doses high enough for microbial disinfection are the result of free radical reactions; comparable to those produced by high cooking temperatures or fermentation.⁴¹ At low doses (i.e., those used for insect disinfection or sprout inhibition), chemical changes are difficult to detect. Numerous animal feeding studies do not detect toxic or carcinogenic effects associated with a pure irradiated food diet.³⁹ In the U.S., comprehensive animal feeding studies and toxicological tests conducted on chicken meat irradiated to 5.8 MRad showed no adverse effects.³⁹ This suggests that the

Codex General Standard limit of 1 MRad offers a large margin of safety. It has been demonstrated that vitamins are no more degraded by radiation than by cooking and that proteins are little affected even at high doses.³⁹ Despite the overwhelming evidence of the safety of irradiated foods, health authorities in many nations are insisting upon classifying and testing the physical process of food irradiation as a chemical additive.⁴¹ This has delayed acceptance of irradiated foods despite improvements in food quality and WHO endorsement; and has led to slow legislation, that allows free trade in irradiated foods.³²

Public acceptance of nuclear technologies in the U.S. is difficult to obtain despite the proven, positive technological solutions they provide for potential health problems. Therefore, it is interesting to investigate the public acceptance of irradiated foods in those countries where it has been done. The world's first commercial food irradiation plant opened in the Netherlands in 1968.⁴² Both ⁶⁰Co and machine sources were used. Marketing of irradiated foods was accompanied by a government public relations campaign that led to the products being publicly well received. Dutch consumers reported the major factor influencing their purchase decision was product freshness. The Psychology Department of the Hebrew University (Israel) conducted similar consumer acceptance studies of irradiated potatoes and onions in 1968–1970.⁴³ The results of the study showed a 2:1 to 4:1 preference for irradiated products because of their product quality. In 1970, large quantities of irradiated products were marketed through retail outlets (clearly labeled as radiation preserved) and were met with practically no resistance.

In the U.S., numerous market tests were performed. In a one-day trial at two markets in Southern California, consumers were offered a choice between both irradiated papayas and papayas treated by the conventional thermal double-dip method for disinfection.⁴⁴ Over 60% at one location preferred the irradiated papayas. At the second location, the preference was split between the two methods. This is not to say that consumers would not buy irradiated papayas if they were the only ones offered. At the first location, 81% said they would buy irradiated papayas, and 66% at the

second location agreed. In this one-day test, irradiated papayas outsold the double-dipped papayas by more than 10:1. In a similar March 1987 study, consumers preferred irradiated Hawaiian papayas to unirradiated papaya 11:1. A market test in Florida in October 1986 also showed a preference for irradiated Puerto Rican mangoes.^a

In a recent American study, consumers were asked about concerns over alternative food treatment methods. A larger percentage of those questioned were more concerned about pesticide residues (85%) and preservatives (80%) than about irradiation (46%).³⁷ Consumers also showed a preference for irradiation treatment of spices (58%) over more conventional ethylene oxide (35%). Other opinion studies show similar percentages of consumers are more concerned about pesticides than radiation.^{45,46}

Other surveys on consumer attitudes toward irradiated foods show the majority of consumers (55–65%) are uncertain rather than opposed (5–10%).^{17,47} Twenty-five to 30% of those surveyed favored the technology over the conventional methods. Public education could reduce concerns and foster more positive attitudes from the undecided consumer majority. When the facts are presented about perceived induced radioactivity, and the absence of novel chemical changes in irradiated foods, irradiated products could be as well received in the U.S. as they were in the Netherlands and Israel.

In conclusion, irradiated foods are well received when tried. Acceptance is increased when official attempts are undertaken to dispel misconceptions associated with the technology. The deciding factor for consumers is increased freshness of irradiated products. The benefits are tangible whereas perceived risks are abstract.

a. Personal communication with J. Moy, Food Engineering Department, University of Hawaii.

Comparison of Current Irradiation Technologies

The current uses of isotope irradiator technology were reviewed. Research indicates that gamma-ray irradiation (of appropriate target materials) results in numerous beneficial changes. Among them are the following: disinfection, disinfestation, and extended shelf life of foods; sterilization of waste, pharmaceuticals, medical supplies, suppression of antigenicity in tissue grafts, and polymerization of monomers.

The said technologies, with the exception of food irradiation, are becoming publicly accepted. The controversial process of food irradiation is still met with mixed approval from an informed public; especially when alternative methods of food preservation are compared. Irradiator technology is improving the safety of foods, and medicines, the use of waste water for irrigation, and structural integrity of common building materials. The public expects direct benefits, regardless of the source.

Proposed technology could be compared to the above technologies in an attempt to estimate public opinion. Hazardous waste irradiation is similar to waste water irradiation because both technologies seek to mitigate effects of a waste form with public health consequences. Despite public health consequences however, sewage is largely ignored by the public, while hazardous waste is a more controversial topic. Likewise, building materials and new polymers are obscured from the public eye; their irradiation is ignored. Of more direct concern are foods and medical supplies. Past research proves that medical supply irradiation is acceptable to the public and that food irradiation could become acceptable if the circumstances were right.

Therefore, it is the position of this paper that widespread acceptability of hazardous waste irradiations will meet public acceptance much the same as waste water or industrial irradiations.

ANALYSIS OF POTENTIAL FOR PUBLIC CONSENT

The following analysis of the elements that are believed to contribute to public consent is based principally on techniques presented in a handbook developed by the Institute for Participatory Management and Planning.⁴⁸

The techniques presented in the handbook seem less rigorous than the analyses in the preceding two sections of this report, because the techniques come directly from the social sciences rather than from the physical sciences. The techniques are based on former case studies, including some projects implemented successfully and some projects that failed because of little public consent. In this context, the techniques provide a useful checklist of factors that are considered if public consent is to be obtained for projects based primarily on this technology.

The handbook's principal use is for proposed projects requiring public consent for implementation. The proposed irradiation technology is not yet at this stage. Projects requiring public consent are some distance in the future; public consent would presumably not be required for pilot program irradiations on the INEL site. However, now is the ideal time to begin consideration of the potential for problems with public acceptability and to develop programs that will end future problems. This report provides an initial opportunity to do so.

The following three courses of action should be considered:

- Questions of legitimacy of future projects based on this technology
- Potential issues and potentially-affected interests (PAIs)
- Proper conduct of the public relations program.

Legitimacy Questions

Legitimacy questions include the following:

- Is a real problem being addressed by the project?

- Is the project within the mission of the Department of Energy (DOE) and the INEL?
- Have possible disadvantages of the project been identified and are they being dealt with in an open and responsible manner?

A serious health problem concerning accumulated chlorinated hydrocarbons in the environment had to be addressed. This was discussed in the Introduction section of this report. This technology presents a cost-effective solution to the problem. The federal government clearly has the responsibility to deal with hazardous wastes created by federal operations. Whether DOE is the appropriate agency to perform research leading to methods of addressing waste problems is a matter for Congress to decide. It does not *appear* to be outside the DOE charter. The DOE is the appropriate agency to be dealing with DOE wastes.

Earlier sections discussed the possible disadvantages of the technology and ways to minimize the disadvantages. The planned research will determine whether disadvantages are outweighing the advantages of the technology. There is no intent to pursue further projects based on this technology, if disadvantages outweigh the advantages (or have unknown or unevaluated disadvantages).

The existence of this report demonstrates DOE's willingness and intent to conduct this program in an open and responsible manner. Therefore, it is reasonable to conclude that projects based on this technology will be legitimate projects pursued with public funds.

Potential Issues

To address the second area, potential issues, an issues matrix was prepared. It shows potential issues, hypothetical issue-related newspaper-headlines, and PAIs that have been identified; others may be added as the program progresses.

The first two potential issues should not arise if the program is conducted properly. Potential risks

to personnel should be routine and acceptable if proper precautions are taken. Transportation of hazardous wastes should not be an issue because other materials equally as hazardous are transported without problems.

If irradiations are conducted at commercial reactor facility spent fuel pools, the Nuclear Regulatory Commission (NRC) would become interested to a degree until specific facilities and designs are selected. Some communications between each facility's owner and NRC would be necessary to establish the acceptability of the operations to NRC; and to determine whether there are any issues that must be resolved before NRC's permission can be obtained. Early informal discussions with NRC are advisable to assess the extent of this potential problem.

Certification of irradiated hazardous wastes as safe for disposal, at specific locations, should not be issues if research is completed as planned, thereby demonstrating technological reliability.

The possible issue of competitiveness with private industry is a difficult issue. If the technology is developed and it becomes the preferred method of disposal of some hazardous wastes, it could be divested to private industry. The actual development of specific projects could be performed by government agencies if it is not practical for private industry to participate or if private industry does not demonstrate an interest. This potential issue appears to involve matters of law that are beyond our purview, but does not appear to obviate development of the technology by DOE.

It is concluded that none of these potential issues or PAIs should stall future projects. The hypothetical newspaper headlines seem credible enough that some measures be taken to obviate the development of such issues. Compliance with transportation requirements, adequate safety analysis, and thorough technical work, should minimize problems.

Proper Conduct

The third area for consideration is the proper conduct of the program. This requires DOE to comply with DOE's internal procedures and that public reporting of the program remains adequate. The DOE and its contractors will follow its own procedures and all applicable regulations in performance of the program. Also, because this program is at an early stage, it can be assumed that DOE will conduct the program in an open and responsible manner; with appropriate consideration given to public concerns. Press releases and other communications to the public will be designed to keep the public informed about the program as it develops and should include candid and balanced presentations about the technology. The public officials responsible for the program should take full advantage of the considerable experience gained throughout the U.S. for project implementation requiring public consent.

The analysis of public consent indicates that, given successful completion of the technical work as planned, no overriding issues prevent the implementation of projects based on this technology.

Potential Issues & Headlines

Hazards to personnel handling materials and operating equipment at sites - "Workers at Any site Reactor to be Exposed to Hazardous Materials."

"Transportation of Hazardous Materials through or past communities - "DOE Plans to Route Hazardous Waste Shipments through Publietown."

Potentially-Affected Interests

Watchdog agencies

Local citizenry
Transportation regulatory agencies

Potential Issues & Headlines

Potential interactions of hazardous materials with spent fuel in pools; e.g., participation in pool accidents, seismic, tornadoes or sabotage; accidents of project that threaten spent fuel – "Hazardous Materials add to Hazards at Anysite Reactor."

Certification that irradiated material is safe to dispose of – "Radioactive PCBs to be Dumped at Publictown."

Ultimate disposal of irradiated material – "Irradiated PCBs to be dumped here."

"Proposed project competes unfairly with private sector."

Potentially-Affected Interests

Watchdog agencies
Nuclear Regulatory Commission

Watchdog agencies
Citizenry near disposal sites

Citizenry near disposal sites

Private companies
Watchdog agencies

CONCLUSIONS AND RECOMMENDATIONS

Three methods of assessing public reaction to the proposed technology were presented. No methods identified a public concern serious enough to prevent implementation of hazardous waste irradiations.

The benefits of the technology are obvious. No other technology can decompose numerous hazardous compounds to nonhazardous products before release to the environment. Public concern is high regarding the chlorinated hydrocarbon accumulation in the environment and it expects a solution from the scientific community. A technology that produces an environmentally acceptable effluent (a process treating materials such as polychlorinated biphenyls and pesticides) would be greatly appreciated by a public that believes cancer rates and other illnesses are on the rise due to environmental contamination. Reservations will arise in some due to the use of spent fuel isotope sources, as opposed to ^{60}Co or machine generated sources. Spent fuel has its own reputation as a hazardous waste and negative connotations are associated with it in public perception. It is believed, however, that many will appreciate the efficiency inherent in using one waste to treat another. Spent fuel already exists at storage locations throughout the U.S., using it for this technology will not increase its danger to the public.

The proposed technology should be accepted by the public with only minor reservations and the following recommendations could help passage:

- Hazardous waste treatment irradiators should be integral or adjacent to facilities where spent fuel is generated. Although the history of radioactive materials shipping has an outstanding safety record, accidents involving radiation are of public concern. Shipping waste to the source rather than the source to the waste is preferred.
- The penetrating ability of gamma rays should be implemented by using a system of multiple containment barriers against radioactive contamination of the effluent. This is easily achieved using the fuel cladding itself as one barrier.
- The real benefits of the technology should be given favorable attention, including press releases. Press releases must be balanced and explain the minimal risks associated with irradiation technology. Treating hazardous waste should be conducted in a manner to allow for good public relations.

By employing recommendations suggested within the context of this paper, an acceptable technology should receive little adverse publicity. Therefore, gamma-ray irradiation can be used in one more way to benefit humankind.

REFERENCES

1. S. M. Matthews, A. J. Boegel, S. F. Eccles, D. W. Rice, J. A. Loftis, M. C. Jovanovich, R. A. Caufield, B. J. Mincher, D. H. Meikrantz, R. J. Murphy, and G. L. Gresham, "High Energy Irradiation of Chlorinated Hydrocarbons," *Presented at the American Nuclear Society International Topical Conference, Kona, HI, April 1991.*
2. W. V. Sherman, R. Evans, E. Nesyto, and C. Radlowski, "Dechlorination of DDT in Solution by Ionizing Radiation," *Nature*, 232, 118, July 1971.
3. R. Evans, E. Nesyto, C. Radlowski, and W. V. Sherman, "Mechanism of Radiation-Induced Dechlorination of 1,1,1-Trichloro-2,2-bis (p-chlorophenyl) ethane in Alcoholic Solution," *J. Phys. Chem.*, 75 (18) 2762, 1971.
4. J. F. Schweitzer, G. S. Born, J. E. Etzel, and W. V. Kessler, "Evaluation of Gamma Radiation for Degradation of a Polychlorinated Biphenyl in Solution and on Activated Carbon," *J. Rad. Nuc. Chem. Let.*, 118, (5), 323, 1987.
5. T. Sawai, T. Shimokawa, and Y. Shinozake, "Radiolytic Dechlorination of Polychlorinated Biphenyls (PCB) in Organic Solutions," *Chem. Let.*, 865, 1972.
6. A. Singh, N. H. Sagert, J. Bursa, H. Singh, and G. D. Bennett, "The Use of High Energy Radiation for the Treatment of Wastewater," *A Review, Environment Canada, Proceedings of the 8th Wastewater Symposium, Montreal, 1985.*
7. A. Singh, W. Kremers, P. Smalley, and G. S. Bennett, "Radiolytic Dechlorination of Polychlorinated Biphenyls," *Rad. Phys. Chem.* 25, 11, 1985.
8. B. J. Mincher, D. H. Meinrantz, R. J. Murphy, G. L. Gresham, and M. J. Connolly, "Gamma Ray Induced Degradation of PCB's and Pesticides Using Spent Reactor Fuel," *Appl Radiat. Isot.*, 42, 1061, 1991.
9. G. P. Dix, "Considerations in the Public Acceptance of Sewage Sludge Irradiation Systems," *Proceedings of the International Symposium on the Use of High-Level Radiation in Waste Treatment, IAEA, Munich, 399, 1975.*
10. E. Lund, "Public Health Aspects of Waste-Water Treatment," *Proceedings of a Symposium, Radiation for a Clean Environment, Munich, 45, 1975.*
11. A. Suess, H. Moetsch, E. Bosshard, G. Schurmann, O. Luescher, *Kerntechnik* 16, (2), Jahrgang, 1974.
12. A. J. Sinskey, D. Shah, K. A. Wright, E. W. Merrill, S. Summer, and J. G. Trump, "Biological Effects of High-Energy Electron Irradiation of Municipal Sludge," *Proceedings of a Symposium, Radiation for a Clean Environment, Munich, 139, 1975.*
13. V. R. D. Gernberger, "Fission Product Gamma-Ray Sources as an Alternative to Cobalt-60 Sources for Sewage Sludge Irradiation," *Radiation for a Clean Environment, IAEA, Munich, 367, 1975.*
14. M. Gerrard, "Sewage and Waste-Water Processing with Isotopic Radiation: Survey of the Literature," *Isot. Radiat. Tech.*, 8, (4), 429, 1971.

15. L. A. Mann, "Biological-Gamma-Radiation System for Sewage Processing," *Isot. Radiat. Tech.*, 8, (4), 439, 1971.
16. P. Kunradt, "Transport and Disposal of Cobalt-60 Industrial Radiation Sources," *Report of a Task Force Meeting on Public Information of Food Irradiation*, ICGFI, Cadarache, France, April 1988.
17. R. E. Engel, and D. D. Derr, "Regulatory Control of Food Irradiation Processes," *Report of a Task Force Meeting on Public Information of Food Irradiation*, ICGFI, Cadarache, France, April 1988.
18. "The Goiania Radiation Accident," *Health Physics* 60, (1) Special Issue, January 1991.
19. J. Taylor, "What are Consumer Demands in Accepting Irradiated Food," *Proceedings of an Advisory Group Meeting, Factors Affecting Practical Application of Food Irradiation*, Vienna, 1988.
20. International Atomic Energy Agency, *Manual on Radiation Sterilization of Medical and Biological Materials*, Technical Report Series No. 149, Vienna, 1973.
21. K. Ostrowski, and S. Moskalewski, *Organization of Tissue Banks based on the Radiosterilization Technique, Manual on Radiation Sterilization of Medical and Biological Materials*, Technical Report Series No. 149, Vienna, 283, 1973.
22. A. E. Kiselev, and G. I. Kozinets, *Report from the Union of Soviet Socialist Republics, Reports of a Panel on the Radiosterilization of Medical Products, Pharmaceuticals and Bioproducts*, Vienna, 77, 1967.
23. International Atomic Energy Agency, *Guidelines for Industrial Sterilization of Disposable Medical Products, (Cobalt-70 Gamma Irradiation)*, Vienna, 1990.
24. B. L. Gupta, "Chemical and Biological Effects of Radiation Sterilization of Medical Products," *Proceedings of the Symposium on Ionizing Radiation for Sterilization of Medical Products and Biological Tissues*, IAEA, Bombay, December, 1974.
25. D. E. Harmer, and D. S. Ballantine, "Applying Radiation to Chemical Processing," *Chemical Engineering* 91, May 3, 1971.
26. J. T. Dikeow, W. C. Cowan, G. W. DePuy, W. G. Smoak, G. B. Wallace, M. Steinberg, L. E. Kukacka, A. Auskern, P. Columbo, J. M. Hendrie, and B. Manowitz, *Concrete-Polymer Materials, Third Topical Report BNL-50275*, January 1971.
27. *Proceedings of the International Symposium on Radiation Vulcanization of Natural Rubber Latex, Gunma, Japan, January 1990.*
28. E. E. Fowler, "Recent Advances in the Application of Isotopes and Radiation in the United States," *Proceedings of the Fourth International Conference*, IAEA, Geneva, Vol. 14, 29, 1971.
29. T. Sadate, "High Energy Units in Food and Pharmaceutical Processing," *Proceedings of a Seminar on Industrial Applications of Electron Beams, Nuclear Energy Board, Dublin, Ireland, April 1987.*
30. P. A. Wills, "Status of Current Research Activities in Food Irradiation, Australian Institute of Nuclear Science and Engineering," *11th AINSE Radiation Biology Conference, Lucas Heights, N.S.W., Australia, August 1987.*

31. "Peaceful Uses of Atomic Energy," *Proceedings of the Fourth International Conference*, IAEA, Geneva, Vol. 12, 1971.
32. A. Sreenivasan, A. S. Aiyar, D. R. Bongirwar, U. S. Kumta, G. W. Rahalkar, P. Sundarsan, P. Thomas, and U. K. Vakil, "Nuclear Methods of Reducing Food Losses," *Proceedings of the Fourth International Conference*, IAEA, Geneva, Vol. 12, 265, 1971.
33. N. Abdullah, E. G. Siagian, N. Isnaeni, and M. Ismachin, "Laboratory Work on Food Irradiation in Indonesia," *Proceedings of the Fourth International Conference*, IAEA, Geneva, Vol. 12, 337, 1971.
34. E. E. Fowler, "Overview of USAEC Interest in Food Irradiation," *Proceedings of a Panel on the Use of Irradiation to Solve Quarantine Problems in the International Fruit Trade*, FAO/IAEA, Honolulu, 1, 1970.
35. K. K. Otagaki, "Hawaiian Development Irradiator – A Tool in Tropical Fruit Processing," *Proceedings of a Panel on the Use of Irradiation to Solve Quarantine Problems in the International Fruit Trade*, FAO/IAEA, Honolulu, 1, 1970.
36. F. K. MacQueen, J. W. Ketcheson, K. O. Lapins, M. D. Proverbs, L. H. Lyall, V. W. Nutall, and D. A. Rennie, "Canadian Studies on Applications of Isotopes and Radiation in Agriculture and Food Preservation," *Proceedings of the Fourth International Conference*, IAEA, Geneva, Vol. 12, 67, 1971.
37. H. G. Schutz, C. M. Bruhn, K. V. Diaz-Knauf, *Food Technology* 43, (10), 80, 1989.
38. G. Subbaraman, H. Farrar IV, and S. B. Ahlstrom, "The Cesium-137 Agricultural Commodities Irradiator (CACI)," *Food Irradiation Processing, Proceedings of a Symposium*, IAEA/FAO, Washington, D. C., 1985.
39. J. F. Diehl, "Toxicological Safety and Nutritional Adequacy of Irradiated Foods," *Report of a Task Force Meeting on Public Information of Food Irradiation*, ICGFI, Cadarache, France, April 1988.
40. M. C. Lagunas-Solar, S. M. Matthews, and D. R. Slaughter, "Scientific Considerations for the Use of 10 MeV X-Radiation in Food Processing," *Proceeding of a Symposium, Food Irradiation Processing*, IAEA/FAO, Washington, D.C., 337, 1985.
41. G. S. Lawrence, "Radiation Chemistry's Contribution to the Food Irradiation Debate," *Australian Institute of Nuclear Science and Engineering 11th AINSE Radiation Biology Conference, Lucas Heights, N.S.W., Australia, August 1987*.
42. R. M. Ulmann, "Introducing Irradiated Foods to the Producer and Consumer," *Proceedings of the Fourth International Conference*, IAEA, Geneva, Switzerland, Vol. 12, 299, 1971.
43. M. Lapidot, "The Israel Food Irradiation Program 1965–1970: Application of a Conceptual Approach to the Commercialization of Irradiated Agricultural Products," *Proceedings of the Fourth International Conference*, IAEA, Geneva, Switzerland, Vol. 12, 279, 1971.
44. C. M. Bruhn, and J. W. Noell, "Consumer In-Store Response to Irradiated Papayas," *Food Technology* 41, (9), 83, 1987.
45. K. W. McNutt, *Consumer Acceptance of Irradiated Foods*, Presented to R&DA, Irradiated Food Products Committee, Boston, MA, 1985.

46. Opinion Research Corporation, *Trends: Consumer Attitudes and the Supermarket*, Food Marketing Institute, Washington, D.C., 1988.
47. *Brand Group, Irradiated Seafood Product*, a position paper for the Seafood Industry, Final Report, Chicago, IL, 1986.
48. H. Bleiker, and A. Bleiker, *Citizen Participation Handbook for Public Officials and Other Professionals Serving the Government, Institute for Participatory Management and Planning*, Sixth Edition, 1990.
49. IAEA/FAO, "Food Irradiation Processing," *Proceedings of Food Irradiation Processing Symposium*, Washington, D.C., 1985.

END

**DATE
FILMED**

8 / 28 / 92