ANNUAL REPORT

to

UNITED STATES DEPARTMENT OF ENERGY

ARCTIC HYDROLOGY AND METEOROLOGY

DOE GRANT NO. 60247

1988

from

Douglas L. Kane, Director
Water Research Center
University of Alaska Fairbanks
460 Duckering Building
Fairbanks, AK 99775-1760

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

ds
DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

INTRODUCTION

The behavior of arctic ecosystems is directly related to the ongoing physical processes of heat and mass transfer. Furthermore, this system undergoes very large fluctuations in the surface energy balance; this can be appreciated by looking at the swings in air temperature (Figure 1). The buffering effect of both snow and the surface organic soils can be seen in Figure 1 by looking at the surface and 40 cm soil temperatures. The active layer, that surface zone above the permafrost table, is either continually freezing or thawing. A large percentage of energy into and out of a watershed must pass through this thin veneer that we call the active layer. Likewise, most water entering and leaving the watershed does so through the active layer. These are a few reasons why it is so critical to understand the hydrology and meteorology of this region.

To date, we have been very successful at monitoring the hydrology of Imnavait Creek with special emphasis on the active layer processes. The major contribution of this study is that year-round hydrologic data are being collected. Although runoff only occurs from May through September, there are a number of ongoing processes taking place throughout the winter such as freezing of the active layer, snow accumulation, snow redistribution by the wind, snowmelt, etc. These processes set the stage for many of the summer ecosystem processes.

The original overall objective of our R4D Phase I study was to define how the thermal and moisture regimes within the active layer change during an annual cycle under natural conditions, and then to define how the regime will be impacted by some imposed terrain alteration. It is proposed that the last part of this objective will be addressed in Phase II. During the course of Phase I, certain adjustments were made to various projects such that all relevant processes were being studied. We inherited the task of collecting meteorological data at Imnavait Creek; two other remote sites were added at Franklin Bluffs and Sagwon. Carl Benson, Kay Everett and Donald Walker are also collecting other types of data at these sites. Some tasks related to snowmelt hydrology

AVERAGE DAILY AIR AND SOIL TEMPERATURES

Figure 1. Measured temperatures of air, ground surface, and soil at 40 cm depth, 1986.

and summer hydrology on both the plot and watershed scale were initiated in cooperation with Kay Everett and Carl Benson so that all aspects of the hydrologic cycle were being studied.

Our major analysis of the hydrologic data sets for Imnavait Creek have been water balance evaluations for plots during snowmelt, water balance for the watershed during both rainfall and snowmelt, and the application of a hydrologic model to predict the Imnavait Creek runoff events generated by both snowmelt and rainfall. Pie charts that illustrate the partitioning of snowmelt into soil storage, runoff and evaporation are shown in Figure 2. Runoff is greatest from those plots that have the greatest snowpack water content. Evaporation becomes more important for those plots with less snow as runoff constitutes a smaller percentage. Water content of the snowpack varies over the watershed because of redistribution by the wind. During the 1986 winter which had no major wind events, the plot average was similar for that of the entire watershed. However, both in 1985 and 1987, major wind events increased the water content of the snowpack in water-tracks, along the valley bottom and on the leeward side of the slopes. This results in a higher percentage of runoff than the average from the plots would indicate.

Water balance calculations for the watershed in 1986 and 1987 are shown in Table 1. Both years had similar snowpacks, but summer rainfall was substantially more in 1987. Consequently, total runoff was much greater. Evapotranspiration varied only 15% between the two summers.

Table 1. Annual water balance for Imnavait Creek watershed.

Year	Snowpack Water Content (cm)	Summer Precip (cm)	Total Precip (cm)	Snowmelt Runoff (cm)	Summer Runoff (cm)	Total Runoff (cm)	Evapo- trans- piration (cm)	Pan Evap (cm)
1986	10.9	16.3	27.2	5.7	6.2	11.9	15.3	31.0
1987	10.8	27.2	38.0	7.1	17.9	25.0	13.0	32.0

1985

1986

1987

Figure 2. Partitioning of snowmelt into runoff, evaporation and soil storage

HBV-3 hydrologic model of streamflow has been used effectively in Sweden, Norway and in Alaska. But, to our knowledge, HBV-3 has never before been used to model a basin completely underlain by permafrost. The ultimate goal of our application is for hydrologic forecasting incorporating physical changes in the basin, but that has not yet been attempted. Such predictions are quite difficult because this not only requires an accurate model, but also one whose components can, with certainty, be related to the characteristics of the catchment.

In this part of the world, there are relatively few short-term data available, and no long-term data. From the analysis of four spring melt events and two full summer seasons, we feel that HBV-3 models soil moisture, evapotranspiration, snow ablation and runoff well. It does a fair job of modeling summer rain events, but more data are needed to better calibrate the model to accurately predict spring snowmelt runoff. Since the calibration data are so limited, prediction of the hydrologic response to changes in watershed characteristics will indeed be difficult.

Hydrology in an arctic region conforms to the accepted hydrologic principles developed in more temperate regions, but the extreme environmental differences cause marked differences in response to various events. The average annual air temperature in the watershed is -8°C and the radiation balance actually totals to a net loss, keeping the ground perpetually frozen. The presence of the permafrost will effectively limit the role of the subsurface system in the hydrologic cycle. The ice-rich mineral soils act as a barrier to prevent infiltration from snowmelt or summer rains, so the contribution to baseflow from the subsurface system is an insignificant portion of the streamflow. Since none of the annual water balance is lost as groundwater recharge, all water leaving the basin is either through near-surface runoff or evapotranspiration.

The two years with modeling results (Figures 3 and 4) were quite different, hydrologically demonstrating the flexibility of the model. A late spring thaw marked 1986. Temperatures during snowmelt were quite

Figure 3. Results from hydrologic simulation with HBV-3, 1986

Figure 4. Results for hydrologic simulation with HBV-3, 1987

high, and incoming solar radiation was very near the season maxima. The entire summer was quite dry and spring snowmelt accounted for about half the total volume of runoff. In 1987, spring snowmelt was much earlier, spring temperatures were much lower, and the summer precipitation was much greater. It was difficult to model the 1987 summer precipitation events which overlapped, with some of the summer precipitation falling as snow. HBV-3 does have an advantage over other similar models in that it does consider the state of summer precipitation (i.e., snow or rain), and it determines runoff appropriately.

**DATE
FILMED**
8/11/92