The submitted manuscript has been authored by a contractor of the U. S. Government under contract No. W-31-109-ENG-38. Accordingly, the U. S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or allow others to do so, for U. S. Government purposes.

ANL-HEP-TR--91-84

DE92 001904

SEP 28 1991

SDC

SOLENOIDAL DETECTOR NOTES

Luminosity Evolution at the SSC

Thomas B. W. Kirk

25 September 1991

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Luminosity Evolution at the SSC

Thomas B. W. Kirk
High Energy Physics Division
Argonne National Laboratory

Abstract

A "realistic" model for the time evolution of the SSC luminosity is given. The SDC instrumentation and physics strategy should be informed by this profile.

Introduction

For many reasons, the evolution of SSC luminosity in time is important to SDC. Although we cannot predict this function with great precision, it will certainly *not* be a step-function up to the design value of 10^{33} . Also, even though the start-time for collisions in the SSC may be quite uncertain, the evolution of luminosity after that time is probably predictable. Taking these statements seriously, and also noting that SSCL has not yet produced an official expectation for the luminosity growth, the present document introduces an informed model for the interim use of the Collaboration.

The relevant time scale for this exercise is years. The key variable for physics is the integrated luminosity over time. From the point of view of trigger requirements, radiation damage and data analysis plans, both the integrated and the instantaneous luminosity are relevant. We provide a model of the instantaneous luminosity (averaged over a day or so) and integrate this function to arrive at the integral values.

Model and Results

We take as our model the turn-on behavior as anticipated by SSCL accelerator scientists at the present time¹. This model predicts:

SSC Luminosity Growth

<i><u>Time</u></i> <i><u>(years)</u></i>	<i><u>Luminosity</u></i> <i><u>(cm⁻²sec⁻¹)</u></i>	<i><u>Luminosity</u></i> <i><u>(SSC years/year)</u></i>
0	0	0
0.5	1.0 x 10 ³¹	0.01
1.0	1.0 x 10 ³²	0.1
1.5	5.0 x 10 ³²	0.5
2.0	1.0 x 10 ³³	1.0
2.5	1.2 x 10 ³³	1.2
3.0	1.5 x 10 ³³	1.5
3.5	1.7 x 10 ³³	1.7
4.0	2.0 x 10 ³³	2.0
5.0	2.8 x 10 ³³	2.8
6.0	3.2 x 10 ³³	3.2
7.0	4.0 x 10 ³³	4.0
8.0	5.0 x 10 ³³	5.0
9.0	6.0 x 10 ³³	6.0
10.0	8.0 x 10 ³³	8.0
11.0	9.0 x 10 ³³	9.0
12.0	10.0 x 10 ³³	10.0

* the value taken here is assumed to be the average over a 'fill', taken as a 24-hour period.

This model for luminosity growth is shown in graphical form in Fig. 1. From this plot we can deduce several useful conclusions:

- the first 'SSC year' of integrated luminosity occurs only after 2.4 calendar years
- the fifth 'SSC year' will be achieved in less than five years of collider operation
- 100 'SSC years' will be reached after about 17 years of operation.

These conclusions assume that the luminosity of the machine will 'top out' at about 10³⁴ cm⁻² sec⁻¹ and that the SSC Laboratory will pursue an aggressive strategy of increasing luminosity as fast as accelerator operating capabilities permit.

If, on the other hand, conditions conspire to limit luminosity at a lower value, the limits shown here could prove optimistic and the achievement of '100 years of SSC luminosity' never realized. Obviously, we should anticipate and plan for the more optimistic curve but be ready to present good physics even if a lower delivery is achieved.

Work supported by the U.S. Department of Energy, Division of High Energy Physics, under Contract W-31-109-ENG-38.

References

- ¹ SSC, Accelerator Physics Division (private communication), July 1991.

SSC Luminosity vs Time

Version: Sept 1991

END

**DATE
FILMED**

11/15/91