LA--11253-MS

DE88 010483

*SESAME Equation of State
Number 7530, Basalt*

J. F. Barnes

S. P. Lyon

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

SESAME EQUATION OF STATE NUMBER 7530, BASALT

by

J. F. Barnes and S. P. Lyon

ABSTRACT

A full-range equation of state (EOS) for dry, nonporous basalt with a grain density of 2.868 g/cm^3 has been constructed and placed on the SESAME library as material number 7530.

As with most rocks, basalt is found with varying composition. It is an igneous rock, composed largely of complex minerals of the plagioclase feldspar and pyroxene groups; typically it contains very little identifiable quartz. In oxide abundance it is fairly similar to the diabbases and gabbros, other igneous rocks of about the same density but different texture. The particular molecular composition represented by equation-of-state (EOS) 7530 is given in Table I, where data are taken from Clark¹; corresponding atomic abundance is shown in Table II. For this mixture the average atom, as used in the GRIZZLY² code, is calculated to have an atomic number $\bar{Z} = 10.665$ and an atomic weight $\bar{A} = 21.574$. The EOS is constructed to have a reference density $\rho_0 = 2.868 \text{ g/cm}^3$, the assumed dry grain density of basalt. The EOS can be mixed with a suitable SESAME EOS for water if desired, and through the hydro-code setup the user can allow for the desired degree of porosity.

High pressure shock Hugoniot data on basalt are somewhat sparse, and since in fact the composition of the samples used in the existing

experiments was not as given in Table I anyway, we chose to use the more complete set of shock data on diabase given by McQueen, Marsh, and Fritz³. Ahrens and Gregson⁴ showed that at the lower pressures where data exist the Hugoniot for gabbro, diabase, and basalt are in good agreement when account is taken of their differing initial density. We would expect the Hugoniot to agree also at higher pressures where the mineral makeup is "forgotten." Because the diabase samples had a density of 2.99 g/cm³, we constructed the EOS at that density and then scaled it to the basalt density given above. We assumed that the breaks in the experimental shock data reported in Reference 3 are due to a single phase change and approximated the U_s-U_p Hugoniot by three straight lines.

A new version of the "CHART JD" ion thermal (nuclear) model⁵ that incorporates an improved treatment of the melting transition was used, along with the CHART-D⁶ prescription for the variation of the Grüneisen parameter, γ , with compression. Reference γ , Debye temperature, cohesive energy, and melt temperature were simply computed as the number-weighted average of the major constituents, as shown in Table III. Although this is a crude procedure, better treatment for these quantities isn't warranted when one considers 1) the other approximations used in constructing equations of state for mixtures (such as the average atom approximation) and 2) the uncertainties of some of the entries in Table III. Note that, for example, since sound speeds and other data for calculation of γ_0 and Θ_D seem to be unavailable for ferrous oxide we used the calculated values for Fe₂O₃. It should also be noted that, as used here, cohesive energy is the energy required per mole of average atoms to separate a substance into its separate atoms at

0 K. The last line of Table III gives the calculated number-weighted averages of the quantities that we used as input to GRIZZLY. In construction of the cold curve, the Lennard-Jones factor was taken to be 0.185, which resulted in a critical temperature of about 6675 K.

Results are shown in Figures 1 through 4. Figure 1 shows selected pressure isotherms over the entire temperature range of the SESAME library table. Below the calculated critical isotherm van der Waal loops are obtained, and the 301 table in SESAME contains these loops. Figure 2 shows the corresponding isotherms with Maxwell equal-area construction. Data for effecting this replacement are stored in the 401 table of the library. Figure 3 shows the energy isotherms and Figure 4 the principal Hugoniot.

TABLE I

Chemical Composition of Basalt as Used in EOS 7530 (Ref. 1)

	<u>By Weight</u>	<u>By Number</u>
SiO_2	50.83%	53.04%
CaO	10.42	11.65
MgO	6.34	9.86
Al_2O_3	14.07	8.65
FeO	9.00	7.86
H_2O	0.91	3.17
Na_2O	2.23	2.26
TiO_2	2.03	1.59
Fe_2O_3	2.88	1.13
K_2O	0.82	0.55
MnO	0.18	0.16
P_2O_5	0.23	0.10

TABLE II

Atomic Abundance as Used in EOS 7530

<u>Element</u>	<u>Abundance (By Number)</u>
O	60.11 %
Si	18.26
Al	5.96
Ca	4.01
Fe	3.48
Mg	3.39
H	2.18
Na	1.56
Ti	0.55
K	0.38
P	0.07
Mn	0.06

TABLE 111

Estimates of the Grüneisen Parameter, Debye Temperature, Cohesive Energy, and Melt Temperature for the Major Constituents of Basalt and Their Resulting Weighted Averages

	<u>γ_o</u>	<u>θ_D</u>	<u>E_{coh}</u>	<u>T_m</u>
SiO_2	0.653	950 K	146 kcal/mol	1883 K
CaO	2.40	640	127	2853
MgO	1.72	630	118	3125
Al_2O_3	1.595	1034	145	2288
FeO	(1.64)	(610)	110	1642
Fe_2O_3	<u>1.64</u>	<u>610</u>	<u>114</u>	<u>1838</u>
Basalt	1.17	851	137	2158

REFERENCES

1. K. F. Clark, "Mineral Composition of Rocks," in Handbook of Physical Properties of Rocks, Volume I, R. S. Carmichael, Ed. (CRC Press, Inc., Boca Raton, Florida, 1984), p.70.
2. J. Abdallah, Jr., "User's Manual for GRIZZLY," Los Alamos National Laboratory report LA-10244-M (September 1984).
3. R. G. McQueen, S. P. Marsh, and J. N. Fritz, J. Geophys. Res. 72, 4999 (1967).
4. T. J. Ahrens and V. G. Gregson, Jr., J. Geophys. Res. 69, 4839 (1964).
5. J. D. Johnson, Los Alamos National Laboratory, private communication, September 1986.
6. S. L. Thompson and H. S. Lawson, "Improvements in the Chart-D Radiation-Hydrodynamic Code III: Revised Analytic Equations of State," Sandia National Laboratories report SC-RR-710714 (March 1972).

7530-1ST
TEMP (K)
A-0.00+00
B-2.98+02
C-1.16+03
D-5.80+03
E-1.16+04
F-2.32+04
G-3.48+04
H-5.80+04
I-1.16+05
J-2.32+05
K-3.48+05
L-5.80+05

changes ?

TUESDAY 2FEB 88

TABLE 301

T4 CRAY SES2D 1R

Fig. 1.

Selected pressure isotherms, containing van der Waal loops, for EOS 7530.

7530-1ST
TEMP (K)
A-0.00+00
B-1.45+02
C-2.98+02
D-5.80+02
E-1.16+03
F-2.90+03
G-5.80+03
H-8.70+03
I-1.16+04
J-1.74+04
K-2.32+04
L-2.90+04

changes ?

TUESDAY 2FEB 88

TABLE 301

M-3.48+04
N-4.64+04
O-5.80+04
P-9.28+04

T4 CRAY SE52D 1R

Fig. 2.

Selected pressure isotherms for EOS 7530 with van der Waal loops replaced by calculated equilibrium vapor pressures.

7530-1ST
TEMP (K)
A-0.00+00
B-2.98+02
C-1.16+03
D-5.80+03
E-1.16+04
F-2.32+04
G-3.48+04
H-5.80+04
I-1.16+05
J-2.32+05
K-3.48+05
L-5.80+05

changes ?

TUESDAY 2FEB 88

TABLE 301

M-1.16+06
N-2.98+06
O-6.96+06
P-1.74+07
Q-6.96+07
R-2.32+08
S-1.16+09

Fig. 3.

Selected energy isotherms for EOS 7530.

T4 CRAY SES2D 1R

7530-HJ1
EO. HJ/KG
0.00+00

changes ?

TUESDAY 2FEB 88

TABLE 301

T4 CRAY SES2D 1R

Fig. 4.

Shock velocity versus particle velocity on principal Hugoniot for EOS 7530.